TITLE: Wood capacitance is related to water content, wood density, and anatomy across 30

temperate tree species

AUTHORS: Kasia Ziemińska[1]*, Emily Rosa[2], Sean Gleason[3], and N. Michele Holbrook[4]

[1]Arnold Arboretum of Harvard University, Boston, MA 02131, USA; [2]Sonoma State

University, Rohnert Park, CA 94928, USA; [3]United States Department of Agriculture –

Agricultural Research Service, Water Management and Systems Research Unit, Fort Collins,

CO 80526, USA; [4]Department of Organismic and Evolutionary Biology, Harvard University,

Cambridge, MA 02138, USA

*kasia.s.zieminska@gmail.com

SUMMARY

____

Water released from wood tissue during transpiration (capacitance) can meaningfully affect

daily water use and drought response. To provide context for better understanding of

capacitance mechanisms, we investigated links between capacitance and wood anatomy.

On twig wood of 30 temperate angiosperm tree species, we measured capacitance, water

content, wood density, and anatomical traits, i.e., vessel dimensions, tissue fractions, and

vessel-tissue contact fractions (fraction of vessel circumference in contact with other tissues).

Across all species, the strongest predictors of capacitance were wood density (WD) and

predawn lumen volumetric water content (VWC$_{L-pd}$, r^2_{adj}=0.44, P<0.0001). Vessel-tissue

contact fractions explained an additional ~10% of the variation in capacitance. Regression

models were not improved by including predawn relative water content (RWC$_{pd}$) or tissue

lumen fractions. Among diffuse-porous species, VWC$_{L-pd}$ and vessel-ray contact fraction

were the best predictors of capacitance, whereas among ring/semi-ring-porous species,

VWC$_{L-pd}$, WD and vessel-fibre contact fraction were the best predictors. Mean RWC$_{pd}$ was

0.65±0.13 and uncorrelated with WD. VWC$_{L-pd}$ was weakly negatively correlated with WD.

Our findings imply that capacitance depends on the amount of stored water, tissue

connectivity and the bulk wood properties arising from WD (e.g., elasticity), rather than the

fraction of any particular tissue.

Keywords: angiosperm trees, capacitance, fibres, parenchyma, vessels, water storage, wood

anatomy, wood density

INTRODUCTION

Water stored in wood can buffer excessive water demand on diurnal (Goldstein *et al.* 1998; Meinzer, James & Goldstein 2004; Meinzer *et al.* 2008; Meinzer, Johnson, Lachenbruch, McCulloh & Woodruff 2009; Scholz *et al.* 2007; Köcher, Horna & Leuschner 2013; Lachenbruch & McCulloh 2014; Carrasco *et al.* 2015) and seasonal time scales (Hao, Wheeler, Holbrook & Goldstein 2013; Pratt & Jacobsen 2017; Salomón, Limousin, Ourcival, Rodríguez-Calcerrada & Steppe 2017). Estimates of the contribution of stored water to a tree's daily water budget range from 5 to 50% (Goldstein *et al.* 1998; Kobayashi & Tanaka 2001; Phillips *et al.* 2003; Meinzer *et al.* 2004; Scholz *et al.* 2007; Köcher *et al.* 2013; Carrasco *et al.* 2015). Thus, water storage can be an important component of whole-plant water balance (Gleason, Blackman, Cook, Laws & Westoby 2014; Blackman *et al.* 2016; Christoffersen *et al.* 2016). However, the mechanisms of water storage and release, and their structural underpinnings remain unclear. Consequently, our understanding of cost vs. benefits of water storage and how storage is coordinated with other tree functions is limited. Here, our objective was to quantify water storage (amount of stored water) and diurnal capacitance (water released per wood volume per change in stem water potential, kg m^{-3} MPa^{-1}) across a diverse suite of 30 temperate angiosperm trees.

Capacitance is most often measured using psychrometers and estimated as the initial slope extracted from a water release-potential curve (Meinzer, James, Goldstein & Woodruff 2003; Meinzer *et al.* 2008; Scholz *et al.* 2007; McCulloh *et al.* 2012; Trifilò *et al.* 2015; Jupa, Plavcová, Gloser & Jansen 2016; Santiago *et al.* 2018; Siddiq, Zhang, Zhu & Cao 2019). However, if tissues *in natura* never become fully saturated, or if the operating water potential of xylem falls outside the initial water release curve, then this way of estimating capacitance may be functionally irrelevant. Oversaturating wood prior to measuring capacitance may also skew results. Moreover, using psychrometers on excised material is likely burdened with an artefact due to water being released from open vessel ends (Tyree & Yang 1990; Jupa *et al.* 2016). Few studies have estimated diurnal capacitance across the range of water potentials experienced by field plants, with only two studies using pressure chamber measurements of non-transpiring leaves (Zhang, Meinzer, Qi, Goldstein & Cao 2013; Wolfe & Kursar 2015) and one study using psychrometers on excised tissues (Richards, Wright, Lenz & Zanne 2014). We addressed this shortcoming by measuring capacitance between predawn and midday during peak summer conditions. To avoid open vessels and oversaturation artefacts, we estimated capacitance from the difference in wood water content between predawn and midday with the corresponding change in stem water potential measured using bagged, non-transpiring (equilibrated) leaves (Begg & Turner 1970; Clearwater & Meinzer 2001). We refer to this measure as 'day capacitance', following Zhang *et al.* (2013).

Anatomical structure should determine wood capacitance (Zimmermann 1983; Tyree
& Yang 1990; Holbrook 1995). Tyree & Yang (1990) proposed that water release consists of
three phases linked to anatomy: 1) an initial phase (0 to *ca.* -0.6 MPa), when water is
released from capillary storage (already embolized fibres, vessels and tracheids, and
intercellular spaces), 2) a second phase (< -0.6 MPa but prior to vessel embolization), when
water is released from elastic storage (cells with elastic cell walls), and 3) a final phase
(below the embolization threshold), when water is released from embolizing vessels and
fibres, likely resulting in permeant damage to the xylem tissue, at least in large trees. Despite
this well-developed and popular theory, few studies have quantified wood anatomy in relation
to capacitance, and none of them used methods, which would avoid open vessels and
oversaturation artefacts. As a result, we lack quantitative understanding of the anatomical
drivers of capacitance, especially for angiosperm trees, which exhibit substantial variation in
stem cell sizes, types, their proportions and geometry.

Given these knowledge gaps, it is surprising that parenchyma, the main living tissue
in wood commonly believed to have elastic cell walls, is often assumed the primary source of
capacitance in stems (e.g, Meinzer *et al.* 2003; Steppe & Lemeur 2007; Plavcová & Jansen
2015; Vergeynst, Dierick, Bogaerts, Cnudde & Steppe 2015; Morris *et al.* 2016; Li *et al.*
2018; Rungwattana & Hietz 2018; Santiago *et al.* 2018). Several studies have extended this
idea to suggest that a higher parenchyma fraction may therefor confer higher capacitance
(Borchert & Pockman 2005; Scholz, Phillips, Bucci, Meinzer & Goldstein 2011; Pratt &
Jacobsen 2017; Secchi, Pagliarani & Zwieniecki 2017; Nardini, Savi, Trifilò & Lo Gullo 2018).
While this may indeed be the case in highly parenchymatous stems like palms or baobabs
(Holbrook & Sinclair 1992; Chapotin, Razanameharizaka & Holbrook 2006a b), it may be
less likely for the majority of tree species, which tend to have lower wood parenchyma
fractions (Morris *et al.* 2016). In these species, parenchyma are connected with other cells
(fibres, vessels) via highly lignified middle lamella impeding parenchyma independent
volumetric changes. Moreover, recent microCT studies, suggest that water released from
cavitating fibres and vessels may contribute meaningfully to capacitance at moderate water
potentials (Knipfer *et al.* 2017, 2019).

Here, we measured detailed anatomical traits, water storage and capacitance to
address the following questions: 1) how large is day capacitance and how does it vary
among diverse temperate tree species?, 2) how much water is stored in wood and does this
water volume limit day capacitance during peak summer?, 3) which anatomical traits are
related to capacitance?

MATERIALS AND METHODS

Site, species and individuals

All measurements were taken from trees growing at the Arnold Arboretum of Harvard

University in Boston (Massachusetts, USA). Mean annual temperature in 2017, the year

when capacitance measurements were taken, was 10.9°C and annual precipitation was 787

mm. Growing season (March to October) mean temperature was 15.3°C and precipitation

was 562 mm. Mean temperature during sampling (August) was 21.2°C and precipitation was

22 mm. The soil texture is variable across the Arboretum and sampled trees grew on: sandy

loam, slit loam, loamy sand, and an outcrop complex. The studied trees were excluded from

the Arboretum's supplemental watering regime during the summer 2017.

Thirty species of deciduous, angiosperm trees were selected, spanning 30 genera

and 26 families (Table 1; tree accession numbers are listed in Table S1). InsideWood

database (InsideWood 2004; Wheeler 2011) was used as a guide to select species with the

most diverse anatomies, e.g., parenchyma abundance, porosity (diffuse-, ring- or semi-ring-

porous) and vessel size.

Due to the limited number of individuals per species, trees were sampled across the

existing variation in topography and soil texture. Trees growing near streams and ponds

were avoided and only healthy trees were sampled. When possible, similar height trees per

given species were studied. We measured branches exposed to the midday sun. However,

variation in forest density near the measured trees contributed to variation in the amount of

sun branches received throughout the course of a day and across measurements and

species.

Capacitance and water storage: sampling and measurements

Water storage and capacitance measurements were performed during the second half of

August 2017. Ten individuals per day were sampled, resulting in a total of nine sampling

days. In the case of rain, sampling was delayed for one or two days. For stem water

potential, four leaves per tree were enclosed in opaque, silver, 4 mil (0.1 mm) thick Mylar zip-

lock bags the evening before sampling. In principle, the water potential of a bagged, non-

transpiring leaf will come into equilibrium with the stem water potential. As such, bagged leaf

water potential (i.e., after equilibration) provides an estimate of stem water potential (Begg &

Turner 1970; Clearwater & Meinzer 2001). Two leaves at predawn and two leaves at midday

were collected. The bagged leaf was cut with a razor blade and the bag was immediately

closed and placed in an insulated box at ambient temperature. Six terminal, 0.5-0.7 m long

twigs per tree, four of which also included the leaves bagged the previous evening, were

sampled. Three of these twigs (per tree) were collected at predawn and three twigs at

midday. Twigs for each predawn-midday pair were located near each other, as close as tree

architecture allowed (usually within 0.5 m) to limit the variation in water content that might be

expected along the length of a branch. After collecting the predawn twig, the cut surface on

the branch still attached to a tree was covered with Vaseline to reduce drying from the exposed surface. Twigs were quickly de-leafed (apical meristem was removed together with most distal leaves), double-bagged in zip-lock bags, and placed in an additional opaque bag at ambient temperature. Each sampling event took approximately two hours. After collecting all leaves and twigs, material was transported to the onsite laboratory within 5 min.

Immediately after arrival to the laboratory, leaf water potential was measured on one leaf per tree using a pressure chamber (Model 1000, PMS Instrument Company, USA). Next, twig segments *ca.* 60 mm long and 5 mm diameter (excluding bark and pith) were cut at a minimum 50 mm distance from the initial, field cut. The segments were quickly wrapped in Parafilm, placed in a 4 mil zip-lock bag, and then into a cooled box with ice. After preparing all samples, the following steps were carried out in a humidity-controlled room (at relative humidity of 75-80%) to minimize water loss. Bark and pith were removed, the ends were trimmed by a few mm, and fresh mass was measured on an analytical balance (Sartorius, 0.00001g), after which the samples were placed in distilled water.

These wood samples were then stored at 4°C for two weeks and, afterwards, saturated mass and volume was measured using Archimedes principle as described in (Ziemińska, Westoby & Wright 2015). In several cases, twigs did not sink even after two weeks of soaking, thus indicating that gas was still present in those samples. These samples were held under water at *ca.* 60°C for several hours. This treatment resulted, in all cases, in samples that sank, and, next, volume and saturated mass was recorded. For logistic reasons, we were unable to measure volume within 48 hours after sampling, as is commonly done in other studies. However, our preliminary tests confirmed that across all species studied, volume measured on fresh vs. saturated samples (after two weeks of soaking at 4°C) differed on average by 2% ± 1.1% SD, and as such, should not meaningfully influence our volume estimates. As comparison, soaking for 48 hours resulted in 1.6% ± 0.9% SD volume change – similar to the change after two weeks. After saturated mass and volume measurements, the samples were dried at 102°C for three days and dry mass was recorded.

Broadly, water storage is defined here as the amount of water contained in a wood sample. It can be expressed as: relative water content or volumetric water content. Relative water content (RWC) is the proportion of water in a sample relative to the maximum amount of water that could be stored in that sample and was calculated as follows:

$$RWC = M_F - M_D \big/ M_S - M_D$$

where M_F is sample fresh mass, M_D is sample dry mass, and M_S is sample saturated mass. Volumetric water content (VWC) indicates total water volume per sample volume (Gartner, Moore & Gardiner 2004) and was calculated as follows:

$$VWC = (M_F - M_D) \big/ V \times \rho$$

where V is sample volume and ρ is water density, assumed to equal 1 g cm^{-3}. We also

partitioned VWC to lumen (VWC_L) and wall (VWC_W) water content. For that, we assumed

that the fibre saturated point (FSP), defined as the point where only water bound in cell wall

is present, is at 30% moisture content (MC) (Ross 2010; Dlouhá, Alméras, Beauchêne, Clair

& Fournier 2018). The standard equation for moisture content is:

$$MC = (M_F - M_D)/M_D \times 100\%$$

(Ross 2010). From this equation, we calculated sample mass at FSP (M_{FSP}) and estimated

VWC_L as:

$$VWC_L = (M_F - M_{FSP})/(V \times \rho)$$

and VWC_W as:

$$VWC_W = (M_{FSP} - M_D)/(V \times \rho)$$

Next, we estimated lumen relative water content (RWC_L, Longuetaud $et\ al.$ 2016):

$$RWC_L = (M_F - M_{FSP})/(M_S - M_{FSP})$$

All water content indices were estimated at predawn and midday, indicated by subscripts ($_{pd}$

and $_{md}$).

Following a simplified version of the equation from Meinzer $et\ al.$ (2003) and Richards

$et\ al.$ (2014), cumulative water released (CWR, kg m^{-3}) was calculated, separately for

predawn and midday, as follows:

$$CWR = (M_S - M_F)/V$$

and then multiplied by 1000 to convert units from g cm^{-3} to kg m^{-3}. Wood day capacitance (kg

m^{-3} MPa^{-1}) was estimated as:

$$Capacitance = (CWR_{md} - CWR_{pd})/(\Psi_{pd} - \Psi_{md})$$

where Ψ is stem water potential. This calculation of capacitance takes saturation as a

reference point, to minimize the effect of intrinsic differences between predawn and midday

samples (e.g., in their wood density). However, capacitance could also be estimated more

directly from fresh and dry samples only:

$$Capacitance_{F-D} = ([(M_F - M_D)/V]_{pd} - [(M_F - M_D)/V]_{md})/(\Psi_{pd} - \Psi_{md})$$

In principle, if predawn and midday water content was measured on exactly the same

samples, the two measures of capacitance would lead to exactly same results. However, this

was not the case in our study, because predawn and midday water content was assessed on

separate samples. Nevertheless, the two measures were well correlated with each other

(r^2=0.76, P<0.001) and the intercept and slope were not significantly different from 0 and 1,

respectively (P=0.80 and P=0.59; Fig. S1 in supporting information; $Paulownia\ tomentosa$

was excluded because, as an extreme outlier (see Results), it would have a
disproportionately strong effect on this analysis). This strong correlation suggests that
findings based on 'capacitance' as presented in the current study would be very similar to
findings based on 'capacitance$_{F-D}$'.

All measured traits, their abbreviations used in the text and units are listed in Table 2.
219

Anatomy: sampling and measurements
For anatomical measurements, one, midday sun-exposed twig per tree was collected in
August 2016. Twigs were transported to the lab in semi-opaque, plastic bags. Next, several
pieces with wood diameter of 4-6mm (excluding bark and pith) were cut and stored in 70%
ethanol for about three months until further processing. In about a third of cases, samples
collected in 2016 and 2017 came from different individuals (Table S1) due to either
deteriorated foliage health state between the two years or considerable storm damage. Twig
size (diameter and length) and location on the tree (e.g., aspect) were consistent between
2016 and 2017 sampling, although in some cases slightly lower branches were measured in
2017 to allow for leaf bagging.

Cross-sections ~10-20 µm thick were made using Reichert sledge microtome, blade
holder (Accu-Edge M7321-43) and low-profile blades (Accu-Edge, 4689). To facilitate
flattening, sections were stored between two glass slides held together by paper clips in 50%
ethanol for several days. Next, sections were stained in a mixture of safranin O and Alcian
Blue (0.35g safranin O in 35ml 50% ethanol + 0.65g of Alcian Blue in 65ml of distilled water)
for three minutes, and then rinsed and mounted on a glass slide in glycerol. Longitudinal
radial and tangential sections of one sample per species were also taken to assist in
anatomical interpretations. A pie-shaped region of one cross-section per tree (three trees per
species) was photographed using a Zeiss Axiophot microscope, AxioCam 512 camera and
Zen Blue 2.3 software (ZEISS, Germany). Magnification was provided by a Plan-Neofluar
20x objective. Several photos were taken to cover the entire pie region, stretching from pith
to cambium, which then were stitched together in freeware Image Composite Editor
(Microsoft, 2015).

Anatomical measurements were done on three individuals per species (Table S1),
across all growth rings (4.4 ± 1.5 SD) with the exception of vessel-tissue contact fractions
(see below), which were measured on all except the innermost ring due to the substantial
time/labour cost of this measurement. The potential error resulting from this approach is
likely minimal because tissue fractions measured on all growth rings exhibited a strong
correlation with tissue fractions measured on all except innermost ring (r^2>0.90 for most
tissues, r^2= 0.79 for fibre fraction and r^2= 0.76 for living fibre fraction), and the innermost ring
usually contributes little to the whole-twig cross-section.

Lumen and wall were measured separately for all tissue fractions: fibre, living fibre,

axial parenchyma, ray parenchyma, vessels, and conduits with maximum diameter <15 µm

(for these, lumen and wall were counted together). The latter category (denoted hereafter

conduits$_{15}$) likely encompassed small vessels, the tapered ends of vessels (tails), and

tracheids. Living fibres were identified by the presence of starch, nuclei or septa, and their

wall thickness was similar to fibres and/or thicker than parenchyma walls. From these

measurements, we calculated the fraction of lumen per given tissue. The vessel

characteristics measured included: vessel mean area and diameter (mean of minimum and

maximum diameters of a given vessel), hydraulically weighed diameter ($D_H = \Sigma$diameter^5 /

Σdiameter^4; Sperry, Nichols, Sullivan & Eastlack 1994), vessel number per cross-sectional

area, and the vessel size-to-number ratio (denoted S in Zanne *et al.* 2010), calculated as

mean vessel area divided by vessel number per cross-sectional area). We also measured

the proportion of vessel circumference in contact with other tissues, referred to collectively as

'vessel-tissue contact fraction'. For example, 'vessel-axial parenchyma contact fraction' is the

proportion of vessel circumference in contact with axial parenchyma.

For vessel traits, we first colored vessels in black using the "magic wand" tool in

freeware GIMP (GNU Image Manipulation Program, www.gimp.org) excluding protoxylem

vessels. Next, vessels and the total pie-shaped sample area were automatically measured in

freeware ImageJ (ImageJ; Schneider, Rasband & Eliceiri 2012). Image manipulation and

analysis were processed using Wacom Cintiq 22HD pen display (Wacom Technology

Corporation, Portland, USA).

For tissue fractions, a grid method was applied (Ziemińska *et al.* 2015). Briefly, each

grid point overlaid over the pie region in ImageJ was classified ('Cell counter' plugin,

https://imagej.net/Cell_Counter) depending on which tissue it fell into (Fig. 1 in Ziemińska *et*

al. 2015). The distance between grid points was 55 µm. On average, we analysed 484

(±128) points per pie region and the total number of points depended on the wood sample

and the pie region sizes. For ray wall fraction we counted only walls perpendicular to the

cross-section, and as such our measurements are likely underestimates. For vessel-tissue

contact fractions, all vessels in a pie region were analysed except for the innermost growth

ring (see above). For this measurement, we modified the grid method. In Photoshop CS4

(Adobe Systems Incorporated, USA), dots were placed at equal distances to each other at

70 pixels (22.5 µm, Fig. 1) along the vessel lumen circumference, with the exception of the

first and last dot, whose distance to each other was determined by vessel circumference. We

counted the last dot only if its distance from the first dot was larger than 35 pixels (half-

distance between all other dots). Next, using the 'Cell counter' plugin as for tissue fractions,

we classified each dot depending on the neighbouring tissue. For instance, if the dot fell on

the border with ray, it was classified as vessel-ray contact dot, and so forth. We then took the

ratio of all vessel-ray contact dots to total analysed dots in a pie region, and did the same for

the other tissues. This 'dot-ratio' is vessel-tissue contact fraction for a given, abutting tissue.

On average, we analysed 943 dots (±605) per each pie region. The number of dots

depended on the total vessel circumference (e.g., it was the highest for species with large

vessel fraction composed of many small vessels).

Wood density

Wood density (WD) was measured on the same samples as water storage and capacitance

measurements, as sample dry mass divided by saturated volume (g cm^{-3}).

Data analysis

The studied species encompassed 14 diffuse-porous species, 10 ring-porous species, and 6

semi-ring-porous. Anatomically, diffuse-porous species were very different from the two other

groups, which in turn were very similar to each other (see Results). Consequently, we ran

analyses across all species, as well as within the two porosity groups: diffuse-porous and

ring/semi-ring-porous.

All analysis were performed in R (R Core Team 2018). We used linear multiple

regression models to assess predictors of capacitance using the 'lm' function. The

distribution of residuals was evaluated using the 'residualPlot' and 'qqPlot' functions in the

"car" package (Fox & Weisberg 2011). Colinearity of predictors was assessed using the

variance inflation factor (VIF, 'vif' function). When VIF>4, we removed the variables from the

model. *Paulownia tomentosa* was an outlier in many relationships, so we excluded it from

several models and indicate that accordingly in the Results. Correlation matrices were

obtained using the 'corrplot' function in "corrplot" package (Wei & Simko 2017), and

scatterplot matrices were obtained using the base function 'pairs' . Selected figures (e.g.,

correlation matrices) were additionally edited in Illustrator (Adobe Systems Incorporated,

USA).

We note that tissue fractions and vessel-tissue contact fractions are not independent, i.e.,

two fractions taken from the same whole should be expected to vary inversely with one

another. However, this expectation is weakened when more than two tissues create the

whole, as is the case for our study. Nevertheless, r^2 and *P* values calculated from plotting

one fraction against another should be interpreted with caution and in the context of all other

tissue fractions.

To assess the relationship between total lumen fraction and saturated VWC_L (VWC_{L-sat}), we

fitted major axis models and tested for differences in slope and elevation using the 'ma'

function in the "smatr" package (Warton, Duursma, Falster & Taskinen 2011).

RESULTS

Anatomical variation

Anatomical variation in tissue fractions, vessel properties and vessel-tissue contact fractions was considerable (Table S2, Fig.2a). Fibres represented the highest fraction (0.45 ± 0.08, here and thereafter we report mean ± one standard deviation), followed by axial+ray parenchyma (0.28 ± 0.08), and then vessels (0.24 ± 0.11). Average axial+ray parenchyma fraction was higher than that found by global analysis of ~400 temperate species (0.21 ± 7.9) and ranged from 0.16 to 0.43, spanning more than half of the reported in that global analysis (Morris *et al.* 2016). Vessel fraction encompassed almost the entire spectrum of global variation (Zanne *et al.* 2010). Conduits$_{15}$ were the least abundant (0.016 ± 0.023, absent in *P. tomentosa*). Living fibres were observed in eight species, but comprised only a small fraction of total wood area (0.026 ± 0.037) with *Acer saccharum* as an outlier (0.11). Axial and ray parenchyma had similar fractions (axial: 0.13 ± 0.09, ray: 0.15 ± 0.04). The fraction of tissue lumen per total area of a given tissue, called here 'lumen fraction$_{tissue}$' (Table 2) varied considerably across tissue types, with vessels exhibiting the largest lumen fraction$_{vessel}$ (0.82 ± 0.04), followed by rays (0.64 ± 0.05), axial parenchyma (0.52 ± 0.10), and then fibres (0.30 ± 0.15). Although fibres had the lowest lumen fraction$_{fibre}$, it also varied the most (8-fold) across species. As mentioned in Materials and Methods, ray lumen fractions (per entire wood cross-section as well as per ray tissue) were likely overestimated (wall was likely underestimated).

Vessel-tissue contact fractions differed significantly between tissues and across species, more so than tissue fractions (Table S2, Fig.2b). The largest contact fraction was between vessels and axial parenchyma (0.36 ± 0.28) and between vessels and fibres (0.27 ± 0.22 SD). Vessel-vessel contact fraction was on average 0.18 ± 0.09, followed by vessel-ray contact (0.13 ± 0.07).

Anatomical characteristics differed considerably between diffuse-porous, semi-ring-porous and ring-porous species (Fig. 3), and these differences strongly influenced the investigated trait-trait relationships. Semi-ring-porous and ring-porous species tended to group with each other and many of these species occur in both forms (InsideWood 2004), hence we grouped them into one category denoted 'ring/semi-ring-porous' (Fig. 3).

Within diffuse-porous species, fibre fraction was inversely related to vessel fraction (r^2=0.68, P<0.001 P<0.001, Fig.4a), but not to axial+ray parenchyma fraction (r^2=0.00, P=0.87, Fig. 4b). In contrast, within ring/semi-ring-porous species, fibre fraction was negatively correlated with axial+ray parenchyma fraction (r^2=0.50, P<0.01, Fig. 4b) but not with vessel fraction (r^2=0.14, P=0.15, Fig. 4a). Across all species, vessel fraction was negatively related to axial+ray parenchyma fraction (r^2=0.43, P<0.001, Fig. 4c) and this was driven by porosity type, i.e., diffuse-porous species tended to have higher vessel fraction and lower axial+ray parenchyma as oppose to ring/semi-ring-porous species. Axial

parenchyma fraction was negatively correlated (weakly) with ray fraction, with diffuse-porous

species tending towards higher ray fraction and ring/semi-ring-porous toward higher axial

parenchyma fraction (r^2=0.15, P<0.05, Fig. S2).

Species with more abundant axial and ray parenchyma tended to have more contact

between these tissues and vessels. However, having abundant fibre or vessel fractions did

not translate to higher vessel-fibre or vessel-vessel contact (Fig. S2). Vessel-fibre contact

fraction was inversely correlated with vessel-axial parenchyma contact fraction (r^2=0.75,

P<0.001, slope: -1.1, Fig. 5), Vessel-ray contact fraction was positively related to vessel-fibre

contact fraction (r^2=0.34, P<0.001, slope: 1.9) and negatively to vessel-axial parenchyma

contact fraction (r^2=0.40, P<0.001, slope: -2.6). The relatively steep slopes of these

relationships indicated that per one unit change in vessel-ray contact fraction, the two other

tissues changed by approximately two units. All these trade-offs were driven by strong

differences between porosity groups (Fig. 3).

Vessel diameter was 33 ± 7 µm (mean vessel area: 1077 ± 531 µm^2) and D_H was 61

± 25 µm averaged across all species. Vessel properties differed markedly between diffuse-

porous and ring/semi-ring-porous species (Fig. 3). Mean vessel area was negatively

correlated with vessel number per mm^2 (i.e., species with small vessels had many of them,

log10 r^2=0.84, P<0.001, Fig. S2). Species with smaller vessels, measured as mean vessel

area or D_H, also tended to have higher total vessel lumen fraction (log10 mean vessel area:

r^2=0.49, P<0.001, log10 D_H: r^2=0.51, P<0.001, Fig. S2). Vessel size was positively correlated

with axial parenchyma fraction (log10 mean vessel area: r^2=0.33, P<0.001, log10 D_H:

r^2=0.56, P<0.001), and negatively (albeit weakly) with ray fraction (mean vessel area:

r^2=0.25, P<0.01, D_H: r^2=0.10, P<0.10). All of these relationships were strongly affected by

porosity, with diffuse-porous species at the small vessel area end of spectrum and ring/semi-

ring-porous species at the large vessel area end.

Anatomy and wood density

Total lumen and total wall fraction were the strongest drivers of WD (negative with lumen:

r^2=0.66, P<0.001, positive with wall: r^2=0.60, P<0.001, Fig. S2). Among the specific tissues,

fibre lumen and fibre wall fractions were strongest drivers of WD (negative with fibre lumen:

r^2=0.48, P<0.001, positive with fibre wall: r^2=0.36, P<0.001). While axial+ray parenchyma

fraction was positively correlated with WD (all species: r^2=0.17, P<0.05, excluding *P.*

tomentosa: r^2= 0.44, P<0.001). Vessel lumen fraction was negatively correlated with density

(all species: r^2=0.13, P<0.1, excluding *P. tomentosa*: r^2=0.23, P<0.01,). Mean vessel area

and D_H were not related to WD across all species (mean vessel area: r^2=0.003, P=0.77, D_H:

r^2=0.06, P=0.19).

Water storage

Relative water content (RWC) was 0.65 ± 0.13 at predawn and 0.59 ± 0.12 at midday. Lumen relative water content (RWC_L) was 0.52 ± 0.17 at predawn and 0.45 ± 0.16 at midday (Fig. 6). Lumen volumetric water content (VWC_L) was on average 0.25 ± 0.09 at predawn, 0.21 ± 0.08 at midday and 0.47 ± 0.09 at saturation (Table S2). The relationship between VWC_{L-sat} and total lumen fraction was strong, as expected ($r^2=0.63$, $P<0.001$, Fig. S3). The slope of that relationship was not significantly different from 1 (major axis fit, $P=0.53$) and the intercept was not significantly different from 0 ($P=0.29$) suggesting that the two measurements are directly related to each other. Wall VWC (VWC_W) averaged 0.16 ± 0.03 and was positively correlated with total wall fraction ($r^2=0.62$, $P<0.001$).

Predawn water content was strongly associated with midday water content for all water content indices ($r^2>0.97$, $P<0.001$). Therefore, the following relationships are reported for predawn values only unless stated otherwise. VWC_{L-pd} was positively correlated with RWC_{L-pd} ($r^2=0.70$, $P<0.001$) and RWC_{pd} ($r^2=0.49$, $P<0.001$). RWC_{pd} and RWC_{L-pd} were strongly correlated with each other ($r^2=0.91$, $P<0.001$), and, we use RWC_{L-pd} in subsequent analyses.

Relationships between anatomical traits and water content differed depending on water content indices and porosity type. Across all species, vessel lumen fraction was positively correlated with VWC_L (all species: $r^2=0.19$, $P<0.05$, excluding *P. tomentosa*: $r^2=0.39$, $P<0.001$, Fig. 7a). Other tissue lumen fractions were weakly or not correlated with VWC_{L-pd},(Fig. S2). VWC_{L-pd} tended to be higher in species with smaller vessels (log10 mean vessel area: all species: $r^2=0.20$, $P<0.05$, excluding *P. tomentosa*, $r^2=0.44$, $P<0.001$; log10 D_H: all species: $r^2=0.32$, $P<0.01$, excluding *P. tomentosa*, $r^2=0.48$, $P<0.001$). These were the strongest pairwise relationships. VWC_{L-pd} was negatively correlated with WD ($r^2=0.19$, $P<0.05$, Fig. S1) across all species. Within the two porosity groups, tissue lumen fractions or WD were not correlated with VWC_{L-pd} (Fig.S2).

Relationships between RWC_{L-pd} and tissue fractions were stronger than for VWC_{L-pd}, and with different tissues. Across all species, the ones with higher fibre lumen fraction tended to have lower RWC_{L-pd} (all spp: $r^2=0.19$, $P<0.05$, excluding *P. tomentosa*: $r^2=0.37$, $P<0.001$, Fig. 7b), while other tissue lumen fractions were weakly or not related (Fig. S2). Within porosity type, fibre lumen was also the strongest, inverse correlate of RWC_{L-pd} (diffuse-porous: $r^2=0.65$, $P<0.001$, all ring/semi-ring-porous: $r^2=0.07$, $P>0.1$, excluding *P. tomentosa*: $r^2=0.58$, $P<0.01$). Higher RWC_{L-pd} was associated with higher ray lumen fraction in diffuse-porous ($r^2=0.33$, $P<0.05$) and higher axial lumen fraction in ring/semi-ring-porous (excluding *P. tomentosa*: $r^2=0.50$, $P<0.01$). In neither of the porosity groups was vessel lumen fraction related to RWC_{L-pd}.

RWC$_{L-pd}$ was not related to WD across all species (r^2=0.00, P=0.79), nor within

ring/semi-ring-porous species (r^2=0.00, P=0.95). But it was positively correlated within

diffuse-porous species (r^2=0.45, P=0.01).

Day cumulative water released

The amount of water released between predawn and midday (ΔCWR$_{pd-md}$) was on average

37 ± 14 kg per m^3 of wood, and was positively correlated with VWC$_{L-pd}$ (r^2=0.31, P<0.01) and

more weakly with RWC$_{L-pd}$ (r^2=0.15, P<0.05). ΔCWR$_{pd-md}$ was also weakly negatively related

to WD across all species (r^2=15, P<0.05). Tissue lumen fraction or vessel-tissue contact

fractions were not related to ΔCWR$_{pd-md}$, except for a weak positive correlation with vessel

lumen fraction (all species: r^2=0.12, P<0.1, excluding *P. tomentosa*: r^2=0.21, P<0.05) driven

by porosity type.

Across diffuse-porous species, ΔCWR$_{pd-md}$ was negatively correlated with vessel size

(mean vessel area: r^2=0.36, P<0.05, D$_H$; r^2=0.36, P<0.05) and not correlated with tissue

fractions or contact fractions (on pairwise basis). Across ring/semi-ring-porous species,

higher ΔCWR$_{pd-md}$ was associated with higher VWC$_{L-pd}$ (r^2=0.55, P<0.01, excl. *P. tomentosa*:

r^2=0.39, P<0.05), and lower WD (r^2=0.30, P<0.05).

Stem water potential

Predawn stem water potential (Ψ_{pd}) ranged from -0.21 to -0.93 MPa and midday stem water

potential (Ψ_{md}) from -0.37 to -2.1 MPa (Table S2). The largest change between predawn and

midday water potential ($\Delta\Psi_{pd-md}$) was 1.6 MPa and the smallest was 0.13 MPa. WD was

higher in more negative Ψ_{md} (r^2=0.53, P<0.001) and larger $\Delta\Psi_{pd-md}$ (r^2=0.51, P<0.001)

species, but correlated weakly with Ψ_{pd} (r^2=0.11, P<0.1, Fig. S2).

Capacitance

Day capacitance was on average 53 ± 26 kg m^{-3} MPa^{-1}, excluding *P. tomentosa*, and ranged

from 8 to 99 kg m^{-3} MPa^{-1} (Table S2). The outlier, *P. tomentosa*, had a capacitance of 505 kg

m^{-3} MPa^{-1}, and was excluded from further analysis. Across all species, those with higher

capacitance tended to have lower WD (r^2=0.35, P<0.001, Fig.8a) and higher VWC$_{L-pd}$

(r^2=0.29, P<0.01, Fig. 8b, Table 3). Together, WD and VWC$_{L-pd}$ explained 44% of the

variation in capacitance (r^2_{adj}=0.44, P<0.001), meaning that for a given WD, species with

higher VWC$_{L-pd}$ tended to have higher capacitance and for a given VWC$_{L-pd,}$ species with

lower WD had higher capacitance. Adding tissue lumen fractions to that model did not

improve its strength but adding contact fraction did (Table 3). For a given VWC$_{L-pd}$ and WD,

capacitance was higher in species with higher vessel-axial parenchyma contact fraction

(r^2_{adj}=0.56, P<0.001), lower vessel-fibre contact fraction (r^2_{adj}=0.54, P<0.001) and lower

vessel-ray contact fraction (r^2_{adj}=0.54, P<0.001, Table 3). These were the strongest models

describing correlates of day capacitance across all species.

Relationships between capacitance and other traits differed between porosity types

and across all species. Among diffuse-porous species, WD or VWC_{L-pd} were not or were only

marginally correlated with capacitance (Table3). The trait combination that was most strongly

related to capacitance, was VWC_{L-pd} minus vessel-ray contact fraction (r^2=0.52, P<0.01) and

plus vessel-vessel contact fraction (r^2=0.41, P<0.05). In ring/semi-ring-porous species,

VWC_{L-pd} together with WD explained the largest proportion of capacitance variation

(r^2_{adj}=0.59, P<0.01, or after excluding *Sassafras albidum*, r^2_{adj}=0.73, P<0.001; *S. albidum* was

the only species with the majority, if not all, fibres being gelatinous and an outlier in vessel-

fibre contact fraction, Fig. 5). In addition, per given VWC_{L-pd}, species with higher vessel-fibre

contact fraction also had higher capacitance (r^2_{adj}=0.63, P<0.01) but only after removing *S.*

albidum (not significant, when *S. albidum* included).

When we substituted VWC_{L-pd} with RWC_{pd} in the models listed in Table 3, their

strength was either comparable ('RWC_{pd} + WD' model across diffuse-porous species),

weaker by 2-7% or not statistically significant (for other models). We also ran a set of models

substituting VWC_{L-pd} with WD. Across all species, no models surpassed the ones including

VWC_{L-pd} (0.28<r^2_{adj}<0.37, P<0.01). Among diffuse-porous species, none of the models were

significant (P>0.15), whereas across ring/semi-ring-porous species only WD was statistically

significant.

Apart from the multiple regression models, species with higher capacitance tended to

have less negative Ψ_{pd} (log10, r^2=0.44, P<0.001) and narrower $\Delta\Psi_{pd-md}$ (log10, r^2=0.50,

P<0.001).

DISCUSSION

Day capacitance and cumulative water release

Across all species, WD and VWC_{L-pd} were the strongest correlates of capacitance (Table 3,

Fig. 8a, b). An inverse relationship between WD and capacitance was also found in other

studies, using the bagged leaf/shoot method (Wolfe & Kursar 2015; Li *et al.* 2018), as well as

in studies having used psychrometers , across a wide range of water potentials (Richards *et*

al. 2014, Meinzer *et al.* 2003, 2008; Scholz *et al.* 2007; Trifilò *et al.* 2015; Jupa *et al.* 2016;

Santiago *et al.* 2018; Siddiq *et al.* 2019). Taken together, these studies offer strong evidence

linking capacitance to the water content and elasticity of wood tissues.

The finding that higher absolute water content (here VWC_{L-pd}) conferred higher

capacitance is new, yet, it makes intuitive sense, i.e., the more water there is, the more can

potentially be released. Perhaps less intuitive is the result that WD was a stronger driver of

capacitance than the proportions of any individual tissues. This implies that capacitance is an
emergent property of the whole-wood, rather than being linked strongly to any one
anatomical component. All cells in wood are tightly packed (Fig. 1), in contrast to leaves, and
joined via highly lignified lamella. Therefore, any change in cell volume, resulting from water
release, would require a coordinated change across neighbouring cells (Holbrook 1995).
This, presumably, would be more feasible in lower WD species that have a greater whole-
organ ability to shrink (Irvine & Grace 1997). This shrinkage would result from cell volume
change due to water release and/or cell wall shrinkage due to adhesive forces between
water and the surfaces of the water transporting conduits (Rosner, Karlsson, Konnerth &
Hansmann 2009). Indeed, daily sapwood shrinkage has been observed in several
angiosperm trees (Scholz *et al.* 2008; Sevanto, Hölttä & Holbrook 2011; Lintunen, Lindfors,
Nikinmaa & Hölttä 2017; Hölttä *et al.* 2018), although Scholz *et al.* (2008) found no
correlation between WD and diurnal sapwood shrinkage across six species. The latter study,
however, encompassed a narrow WD range (~0.42-0.62 g cm^{-3}) and when lower density
tissues from bark were included, strong covariation with capacitance was observed.
Enlarging the number of species and broadening the wood density range in future studies
would likely help to clarify this issue.
Parenchyma lumen fraction, as well as all other tissue lumen fractions, either were not
correlated with capacitance or were less strongly correlated than WD. Furthermore, the
tissue fractions that were significant in multiple regression models were also correlated with
WD (fibre lumen and axial+ray lumen parenchyma fractions, but not vessel lumen fraction,
see below). This finding is in concordance with previous studies, which also found no
relationship between parenchyma fraction and capacitance in four (Jupa *et al.* 2016) and
nine angiosperm species (Pratt, Jacobsen, Ewers & Davis 2007). Wood parenchyma is often
assumed to be a reservoir of capacitance water (Meinzer *et al.* 2003; Steppe & Lemeur
2007; Plavcová & Jansen 2015; Vergeynst *et al.* 2015; Morris *et al.* 2016; Li *et al.* 2018;
Rungwattana & Hietz 2018; Santiago *et al.* 2018) with some studies suggesting that more
abundant parenchyma confers higher capacitance (Borchert & Pockman 2005; Scholz *et al.*
2011; Pratt & Jacobsen 2017; Secchi *et al.* 2017; Nardini *et al.* 2018). Here, we show that
parenchyma lumen fraction does not limit capacitance (Fig. 8c). If anything, vessel lumen
fraction had the strongest, albeit still weaker than WD, link to capacitance (r^2=0.14, *P*<0.05,
Fig. 8d) and ΔCWR_{pd-md} (r^2=0.21, *P*<0.05) across all species, as well as within ring/semi-ring-
porous species (multiple regression, r^2_{adj}=0.55, *P*<0.01, Table 3). This result is aligned with
the idea that cavitating vessels might contribute to capacitance, even at xylem water
potentials well above critical thresholds (Hölttä, Cochard, Nikinmaa & Mencuccini 2009;
Vergeynst *et al.* 2015; Knipfer *et al.* 2019). Moreover, $\Delta VWC_{L-pd-md}$ averaged at 0.04 ± 0.02,
which is small in comparison with tissue lumen fractions (Fig. 2, Table S2), suggesting that

lumen fraction might not limit water release or that water could be released from multiple
tissues. The interpretation of these results is further complicated by the possibility of variable
water distribution across growth rings as well as within growth rings (Umebayashi *et al.* 2008,
2010). More sophisticated methods, for example, microCT or MRI (De Schepper, van
Dusschoten, Copini, Jahnke & Steppe 2012; Knipfer *et al.* 2019), would be necessary to
resolve these questions.

Instead of tissue lumen fractions, vessel-tissue contact fractions tended to be more
strongly linked to capacitance (as independent variables in multiple regression models)
across all species, as well as within porosity groups. Within ring/semi-ring-porous species,
per given $VWC_{L\text{-pd}}$, capacitance increased with higher vessel-fibre contact fraction (Table 3),
suggesting that water may be released from fibres in these species (as well as vessels, see
above). This is in agreement with microCT evidence of emptying fibres (and vessels) as
water potential decreases (Knipfer *et al.* 2017, 2019). An interesting path forward would be to
measure the size of fibre lumina and taper, because these characteristics could have an
additional effect on capillary water release as capillary tension and lumen diameter are
negatively correlated (Tyree & Yang 1990; Hölttä *et al.* 2009). Across diffuse-porous species,
per given $VWC_{L\text{-pd},}$ capacitance decreased with higher vessel-ray contact fraction (Table 3).
Furthermore, across all species, vessel-axial (positive), vessel-fibre (negative) and vessel-
ray (negative) contact fractions were all linked to capacitance with similar strength (Table 3).
Because the three contact fractions are correlated with each other (Fig. 5), it is not possible
to decipher which one of these models may represent a mechanistic link. Nevertheless, the
vessel-ray contact influence is consistent across all species as well as within the diffuse-
porous species (Table 3). How could the inverse influence of vessel-ray contact fraction on
capacitance be explained? One possibility is that, in species limited by a small volume of
water released from wood (low capacitance), additional water could be supplied from bark or
pith (Goldstein, Meinzer & Monasterio 1984; Cochard, Forestier & Améglio 2001;
Pfautsch, Renard, Tjoelker & Salih 2015b; Pfautsch, Hölttä & Mencuccini 2015a;
Mason Earles *et al.* 2016) and higher vessel-ray contact would potentially facilitate the
release of this radially transported water into vessels. Partitioning capacitance between bark,
wood and pith could possibly clarify the influence of contact fractions and ray parenchyma on
whole-stem capacitance. We are aware of only one study (Martínez-Cabrera, Jones, Espino
& Schenk 2009) that quantified contact fractions, but not in relation to any physiological trait.
Our results suggest that tissue connectivity may be an important functional trait.
579
Water storage

None of the studied species was fully saturated – far from it, about half of the cell lumen
was empty (average RWC_{L-pd}: 0.52 ± 0.17). Similar values were reported in the trunk wood of
three temperate angiosperms (Longuetaud *et al.* 2016, 2017). Lack of saturation may explain
why our capacitance values are lower, sometimes by an order of magnitude, than
capacitance estimated as the initial slope of the water release curve (on saturated samples)
in several studies of tropical (Meinzer *et al.* 2003, 2008; Carrasco *et al.* 2015; Santiago *et al.*
2018) and temperate species (Jupa *et al.* 2016), but in a similar range to Cerrado species
(Scholz *et al.* 2007). Given that most of these studies measured capacitance on short xylem
segments, it is possible that water may have been released from open conduits, thus
resulting in an overestimation of capacitance (Tyree & Yang 1990; Jupa *et al.* 2016). Our
values overlap more so with studies which used the bagged leaf method on tropical species
(Zhang *et al.* 2013; Wolfe & Kursar 2015). Note that (Zhang *et al.* 2013) measured
capacitance on entire stems, including bark and pith, which may explain the presence of
species with high capacitance in that study. Our results also overlap with capacitance values
of Australian angiosperms, estimated from excised material using psychrometers within the
native operating shoot water potential range of these species (Richards *et al.* 2014), as well
as with capacitance values estimated from the second, "flatter" phase of the water release
curve (Jupa *et al.* 2016). Overall, these findings highlight the importance of *in natura* water
content and capacitance measurements and the need for a better understanding of how
these measurements compare with ones in which the water potentials were measured on
small, excised samples.
WD, a direct outcome of total lumen and wall fractions, was weakly or not related to fresh
water content indices, similar to other studies on three temperate angiosperms (Longuetaud
et al. 2016, 2017) and across *ca.* 290, 180 and 100 species from Southeast Asia (Suzuki
1999; Kenzo, Tomoaki, Yuta, Joseph Jawa & Sophal 2016; Kenzo, Sano, Yoneda & Chann
2017). This lack of correlation is likely due to a considerable portion of lumen (most likely
fibre lumen, Fig.7) being empty as indicated by RWC_{L-pd}. These results caution against WD
being taken as a direct proxy of water content in the fresh state.
VWC_L was more strongly correlated with capacitance than total VWC or the more
commonly measured RWC, suggesting that absolute, lumen-based water indices are more
relevant for capacitance than relative ones. Moreover, we estimated that the amount of wall-
bound water was considerable (VWC_{W-pd} average: 0.16 ± 0.03) in comparison with lumen
water content (VWC_{L-pd} average: 0.25 ± 0.09). Although, it needs to be noted, that these
estimates may carry some error because of the assumption that fibre saturation point is
attained at 30% moisture content in any species, which is not always the case (Kellogg &
Wangaard 1969; Dlouhá *et al.* 2018). Nevertheless, these findings highlight the need to

better understand the function of lumen and wall water and their explanatory power in

comparison with RWC, which conflates lumen and wall water.

Anatomical landscape

Differences between diffuse-porous and ring/semi-ring-porous species were startling not only

in vessel properties, as has been well documented, but also in tissue fractions and vessel-

tissue contact fractions. To our knowledge, this has not been reported before on a broad

species set and/or with such detail (but see Fujiwara, Sameshima, Kuroda & Takamura

1991; Fujiwara 1992). The anatomical patterns described here agree with ones found across

six angiosperm species (Jupa *et al.* 2019). Other studies also found marked differences in

wood and leaf physiology between these two porosity groups (Bush *et al.* 2008; Klein 2014;

von Allmen, Sperry & Bush 2015). Although this topic is beyond the scope of this study, our

anatomical results raise intriguing questions regarding structure-function relationships in

diffuse-porous vs. ring/semi-ring-porous species.

We found that the wall fraction per total axial or ray parenchyma was relatively high

and surprisingly consistent across species (0.48 ± 0.10 in axial parenchyma and 0.36 ± 0.05

in rays, Fig.S4). Somewhat lower values for axial parenchyma were found in ten tropical

species (average: 0.29, range: 0.07 to 0.53) (Ogunwusi & Ibrahim 2017). Larger values were

found for rays in 50 temperate species (range 0.25-0.65; Fujiwara 1992) and similarly for ten

tropical species (0.25-0.68; Ogunwusi & Ibrahim, 2017). Parenchyma walls, especially axial

parenchyma, are often considered to be thin and of insignificant contribution to the overall

parenchyma proportion (e.g., Martínez-Cabrera *et al.* 2009; Zheng & Martínez-Cabrera 2013;

Ziemińska, Butler, Gleason, Wright & Westoby 2013; Ziemińska *et al.* 2015; Fortunel, Ruelle,

Beauchêne, Fine & Baraloto 2014), yet the results reported here and elsewhere suggest

otherwise. Moreover, parenchyma wall thickness overlaps with that of thin-wall fibres

(Fujiwara *et al.* 1991; Fujiwara 1992; Jupa *et al.* 2016). As such, parenchyma walls may

influence the mechanical properties of wood, across-wall water and solute transport, and

result in lower than expected parenchyma lumen fraction.

Conclusions

This work examined the anatomical correlates of wood water storage and day capacitance.

Contrary to our expectations, tissue lumen fractions did not constrain capacitance. Instead,

WD, VWC_{L-pd} and the connectivity between vessels and other tissues were more closely

related to capacitance than were tissue lumen fractions. Our findings challenge several

common assumptions: 1) that capacitance is positively related to parenchyma fraction, and

2) that wood density is negatively correlated with water content in fresh samples. Given that

fresh wood was never saturated (mean RWC_{pd}: 0.65 ± 0.13), we also question the functional

relevance of capacitance estimated on saturated samples., at least in temperate species.
Our study is limited by a lack of information on: 1) water distribution across and within growth
rings, and 2) the combined and independent effects of wood, bark and pith. Addressing these
issues would likely improve our understanding of capacitance and the anatomical traits that
are aligned with it. Notwithstanding these limitations, our findings offer new insights into
capacitance and its anatomical determinants, and suggest intriguing avenues for future
research.
661

ACKNOWLEDGEMENTS

KZ and ER were supported by the Arnold Arboretum of Harvard University through the Postdoctoral Putnam Fellowship and the DaRin Butz Foundation Research Internship Program, respectively. We are grateful to all the Arnold Arboretum staff, and especially: Faye Rosin (Director of Research Facilitation), Living Collections, Horticulture, and the admin staff for excellent help and support. We also would like to cordially thank the Holbrook Lab at the Department of Organismic and Evolutionary Biology of Harvard University, for discussions about capacitance and providing a pressure chamber.

AUTHOR CONTRIBUTION

KZ designed the study and SMG and NMH contributed to concept and method development. KZ carried out fieldwork and lab measurements. KZ and ER performed anatomical analysis. KZ analysed the data and wrote first draft of the manuscript. All authors contributed to subsequent draft revisions.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

von Allmen E.I., Sperry J.S. & Bush S.E. (2015) Contrasting whole-tree water use,
hydraulics, and growth in a co-dominant diffuse-porous vs. ring-porous species pair.
Trees **29**, 717–728.

Begg J.E. & Turner N.C. (1970) Water Potential Gradients in Field Tobacco. *Plant*
Physiology **46**, 343–346.

Blackman C.J., Pfautsch S., Choat B., Delzon S., Gleason S.M. & Duursma R.A. (2016)
Toward an index of desiccation time to tree mortality under drought. *Plant, Cell &*
Environment **39**, 2342–2345.

Borchert R. & Pockman W.T. (2005) Water storage capacitance and xylem tension in
isolated branches of temperate and tropical trees. *Tree Physiology* **25**, 457–466.

Bush S.E., Pataki D.E., Hultine K.R., West A.G., Sperry J.S. & Ehleringer J.R. (2008) Wood
anatomy constrains stomatal responses to atmospheric vapor pressure deficit in
irrigated, urban trees. *Oecologia* **156**, 13–20.

Carrasco L.O., Bucci S.J., Francescantonio D.D., Lezcano O.A., Campanello P.I., Scholz
F.G., … Goldstein G. (2015) Water storage dynamics in the main stem of subtropical
tree species differing in wood density, growth rate and life history traits. *Tree*
Physiology **35**, 354–365.

Chapotin S.M., Razanameharizaka J.H. & Holbrook N.M. (2006a) Baobab trees (Adansonia)
in Madagascar use stored water to flush new leaves but not to support stomatal
opening before the rainy season. *New Phytologist* **169**, 549–559.

Chapotin S.M., Razanameharizaka J.H. & Holbrook N.M. (2006b) Water relations of baobab
trees (Adansonia spp. L.) during the rainy season: does stem water buffer daily water
deficits? *Plant, Cell & Environment* **29**, 1021–1032.

Christoffersen B.O., Gloor M., Fauset S., Fyllas N.M., Galbraith D.R., Baker T.R., … Meir P.
(2016) Linking hydraulic traits to tropical forest function in a size-structured and trait-
driven model (TFS v.1-Hydro). *Geoscientific Model Development; Katlenburg-Lindau*
9, 4227–4255.

Clearwater M.J. & Meinzer F.C. (2001) Relationships between hydraulic architecture and leaf
photosynthetic capacity in nitrogen-fertilized Eucalyptus grandis trees. *Tree*
Physiology **21**, 683–690.

Cochard H., Forestier S. & Améglio T. (2001) A new validation of the Scholander pressure
chamber technique based on stem diameter variations. *Journal of Experimental*
Botany **52**, 1361–1365.

De Schepper V., van Dusschoten D., Copini P., Jahnke S. & Steppe K. (2012) MRI links
stem water content to stem diameter variations in transpiring trees. *Journal of*
Experimental Botany **63**, 2645–2653.

Dlouhá J., Alméras T., Beauchêne J., Clair B. & Fournier M. (2018) Biophysical
dependences among functional wood traits. *Functional Ecology* **32**, 2652–2665.

Fortunel C., Ruelle J., Beauchêne J., Fine P.V.A. & Baraloto C. (2014) Wood specific gravity
and anatomy of branches and roots in 113 Amazonian rainforest tree species across
environmental gradients. *New Phytologist* **202**, 79–94.

Fox J. & Weisberg S. (2011) *An {R} companion to applied regression.*, Second. Thousand
Oaks CA: Sage.

Fujiwara S. (1992) Anatomy and properties of Japanese hardwoods II. Variation of
dimensions of ray cells and their relation to basic density. *IAWA Bulletin n.s.* **13**, 397–
402.

Fujiwara S., Sameshima K., Kuroda K. & Takamura N. (1991) Anatomy and properties of
Japanese hardwoods. I. Variation of fibre dimensions and tissue proportions and their
relation to basic density. *IAWA Bulletin n.s.* **12**, 419–24.

Gartner B., Moore J. & Gardiner B. (2004) Gas in stems: Abundance and potential
consequences for tree biomechanics. *Tree physiology* **24**, 1239–50.

Gleason S.M., Blackman C.J., Cook A.M., Laws C.A. & Westoby M. (2014) Whole-plant
capacitance, embolism resistance and slow transpiration rates all contribute to longer
desiccation times in woody angiosperms from arid and wet habitats. *Tree Physiology*
34, 275–284.

Goldstein G., Andrade J.L., Meinzer F.C., Holbrook N.M., Cavelier J., Jackson P. & Celis A.
(1998) Stem water storage and diurnal patterns of water use in tropical forest canopy
trees. *Plant, Cell & Environment* **21**, 397–406.

Goldstein G., Meinzer F. & Monasterio M. (1984) The role of capacitance in the water
balance of Andean giant rosette species. *Plant, Cell & Environment* **7**, 179–186.

Hao G.-Y., Wheeler J.K., Holbrook N.M. & Goldstein G. (2013) Investigating xylem embolism
formation, refilling and water storage in tree trunks using frequency domain
reflectometry. *Journal of Experimental Botany*.

Holbrook N.M. (1995) Stem water storage. In *Plant stems: physiology and functional*
morphology. pp. 151–174. Academic Press.

Holbrook N.M. & Sinclair T.R. (1992) Water balance in the arborescent palm, Sabal palmetto.
II. Transpiration and stem water storage. *Plant, Cell & Environment* **15**, 401–409.

Hölttä T., Carrasco M.D.R.D., Salmon Y., Aalto J., Vanhatalo A., Bäck J. & Lintunen A.
(2018) Water relations in silver birch during springtime: How is sap pressurised?
Plant Biology **20**, 834–847.

Hölttä T., Cochard H., Nikinmaa E. & Mencuccini M. (2009) Capacitive effect of cavitation in
xylem conduits: results from a dynamic model. *Plant, Cell & Environment* **32**, 10–21.
ImageJ.
InsideWood (2004) Published on the Internet. http://insidewood.lib.ncsu.edu.
Irvine J. & Grace J. (1997) Continuous measurements of water tensions in the xylem of trees
based on the elastic properties of wood. *Planta* **202**, 455–461.
Jupa R., Doubková P. & Gloser V. (2019) Ion-mediated increases in xylem hydraulic
conductivity: seasonal differences between coexisting ring- and diffuse-porous
temperate tree species. *Tree Physiology*.
Jupa R., Plavcová L., Gloser V. & Jansen S. (2016) Linking xylem water storage with
anatomical parameters in five temperate tree species. *Tree Physiology* **36**, 756–769.
Kellogg R. & Wangaard F. (1969) Variation in the cell-wall density of wood. *Wood and Fiber*
Science **1**, 180–204.
Kenzo T., Sano M., Yoneda R. & Chann S. (2017) Comparison of Wood Density and Water
Content Between Dry Evergreen and Dry Deciduous Forest Trees in Central
Cambodia. *Japan Agricultural Research Quarterly: JARQ* **51**, 363–374.
Kenzo T., Tomoaki I., Yuta I., Joseph Jawa K. & Sophal C. (2016) Wood density and water
content in diverse species from lowland dipterocarp rainforest and dry dipterocarp
forest. *Proceedings of the symposium "Frontier in tropical forest research: progress in*
joint projects between the Forest Department Sarawak and the Japan Research
Consortium for Tropical Forests in Sarawak," 94–103.
Klein T. (2014) The variability of stomatal sensitivity to leaf water potential across tree
species indicates a continuum between isohydric and anisohydric behaviours.
Functional Ecology **28**, 1313–1320.
Knipfer T., Cuneo I.F., Earles J.M., Reyes C., Brodersen C.R. & McElrone A.J. (2017)
Storage Compartments for Capillary Water Rarely Refill in an Intact Woody Plant.
Plant Physiology **175**, 1649–1660.
Knipfer T., Reyes C., Earles J.M., Berry Z.C., Johnson D., Brodersen C.R. & McElrone A.J.
(2019) Spatiotemporal coupling of vessel cavitation and discharge of stored xylem
water in a tree sapling. *Plant Physiology*, pp.01303.2018.
Kobayashi Y. & Tanaka T. (2001) Water flow and hydraulic characteristics of Japanese red
pine and oak trees. *Hydrological Processes* **15**, 1731–1750.
Köcher P., Horna V. & Leuschner C. (2013) Stem water storage in five coexisting temperate
broad-leaved tree species: significance, temporal dynamics and dependence on tree
functional traits. *Tree Physiology* **33**, 817–832.

Lachenbruch B. & McCulloh K.A. (2014) Traits, properties, and performance: how woody
plants combine hydraulic and mechanical functions in a cell, tissue, or whole plant.
New Phytologist **204**, 747–764.

Li X., Blackman C.J., Choat B., Duursma R.A., Rymer P.D., Medlyn B.E. & Tissue D.T.
(2018) Tree hydraulic traits are coordinated and strongly linked to climate-of-origin
across a rainfall gradient. *Plant, Cell & Environment* **41**, 646–660.

Lintunen A., Lindfors L., Nikinmaa E. & Höltta T. (2017) Xylem diameter changes during
osmotic stress, desiccation and freezing in Pinus sylvestris and Populus tremula.
Tree Physiology **37**, 491–500.

Longuetaud F., Mothe F., Fournier M., Dlouha J., Santenoise P. & Deleuze C. (2016) Within-
stem maps of wood density and water content for characterization of species: a case
study on three hardwood and two softwood species. *Annals of Forest Science* **73**,
601–614.

Longuetaud F., Mothe F., Santenoise P., Diop N., Dlouha J., Fournier M. & Deleuze C.
(2017) Patterns of within-stem variations in wood specific gravity and water content
for five temperate tree species. *Annals of Forest Science* **74**, 64.

Martínez-Cabrera H.I., Jones C.S., Espino S. & Schenk H.J. (2009) Wood anatomy and
wood density in shrubs: responses to varying aridity along transcontinental transects.
American Journal of Botany **96**, 1388–1398.

Mason Earles J., Sperling O., Silva L.C.R., McElrone A.J., Brodersen C.R., North M.P. &
Zwieniecki M.A. (2016) Bark water uptake promotes localized hydraulic recovery in
coastal redwood crown. *Plant, Cell & Environment* **39**, 320–328.

McCulloh K.A., Johnson D.M., Meinzer F.C., Voelker S.L., Lachenbruch B. & Domec J.-C.
(2012) Hydraulic architecture of two species differing in wood density: opposing
strategies in co-occurring tropical pioneer trees. *Plant, Cell & Environment* **35**, 116–
125.

Meinzer F.C., Campanello P.I., Domec J.-C., Gatti M.G., Goldstein G., Villalobos-Vega R. &
Woodruff D.R. (2008) Constraints on physiological function associated with branch
architecture and wood density in tropical forest trees. *Tree Physiology* **28**, 1609–
1617.

Meinzer F.C., James S.A. & Goldstein G. (2004) Dynamics of transpiration, sap flow and use
of stored water in tropical forest canopy trees. *Tree Physiology* **24**, 901–909.

Meinzer F.C., James S.A., Goldstein G. & Woodruff D. (2003) Whole-tree water transport
scales with sapwood capacitance in tropical forest canopy trees. *Plant, Cell &*
Environment **26**, 1147–1155.

Meinzer F.C., Johnson D.M., Lachenbruch B., McCulloh K.A. & Woodruff D.R. (2009) Xylem
hydraulic safety margins in woody plants: coordination of stomatal control of xylem
tension with hydraulic capacitance. *Functional Ecology* **23**, 922–930.

Morris H., Plavcová L., Cvecko P., Fichtler E., Gillingham M.A.F., Martínez-Cabrera H.I., …
Jansen S. (2016) A global analysis of parenchyma tissue fractions in secondary
xylem of seed plants. *New Phytologist* **209**, 1553–1565.

Nardini A., Savi T., Trifilò P. & Lo Gullo M.A. (2018) Drought Stress and the Recovery from
Xylem Embolism in Woody Plants. In *Progress in Botany Vol. 79*. Progress in Botany,
(eds F.M. Cánovas, U. Lüttge & R. Matyssek), pp. 197–231. Springer International
Publishing, Cham.

Ogunwusi A. a & Ibrahim H.D. (2017) Variations in Axial and Ray Parenchyma Cells in Ten
Hardwood Species Growing in Nigeria. *Journal of Resources Development and*
Management **38**, 64-68–68.

Pfautsch S., Hölttä T. & Mencuccini M. (2015a) Hydraulic functioning of tree stems—fusing
ray anatomy, radial transfer and capacitance. *Tree Physiology* **35**, 706–722.

Pfautsch S., Renard J., Tjoelker M.G. & Salih A. (2015b) Phloem as Capacitor: Radial
Transfer of Water into Xylem of Tree Stems Occurs via Symplastic Transport in Ray
Parenchyma. *Plant Physiology* **167**, 963–971.

Phillips N.G., Ryan M.G., Bond B.J., McDowell N.G., Hinckley T.M. & Čermák J. (2003)
Reliance on stored water increases with tree size in three species in the Pacific
Northwest. *Tree Physiology* **23**, 237–245.

Plavcová L. & Jansen S. (2015) The Role of Xylem Parenchyma in the Storage and
Utilization of Nonstructural Carbohydrates. In *Functional and Ecological Xylem*
Anatomy. (ed U. Hacke), pp. 209–234. Springer International Publishing.

Pratt R.B. & Jacobsen A.L. (2017) Conflicting demands on angiosperm xylem: Tradeoffs
among storage, transport and biomechanics. *Plant, Cell & Environment* **40**, 897–913.

Pratt R.B., Jacobsen A.L., Ewers F.W. & Davis S.D. (2007) Relationships among xylem
transport, biomechanics and storage in stems and roots of nine Rhamnaceae species
of the California chaparral. *New Phytologist* **174**, 787–798.

R Core Team (2018) *R: A language and environment for statistical computing. R Foundation*
for Statistical Computing, Vienna, Austria.

Richards A.E., Wright I.J., Lenz T.I. & Zanne A.E. (2014) Sapwood capacitance is greater in
evergreen sclerophyll species growing in high compared to low-rainfall environments.
Functional Ecology **28**, 734–744.

Rosner S., Karlsson B., Konnerth J. & Hansmann C. (2009) Shrinkage processes in
standard-size Norway spruce wood specimens with different vulnerability to
cavitation. *Tree Physiology* **29**, 1419–1431.

Ross R.J. ed. (2010) *Wood handbook : wood as an engineering material*, Centennial. US
Department of Agriculture. USDA Forest Service. Forest Products Laboratory,
Madison, WI, USA.

Rungwattana K. & Hietz P. (2018) Radial variation of wood functional traits reflect size-
related adaptations of tree mechanics and hydraulics. *Functional Ecology* **32**, 260–
272.

Salomón R.L., Limousin J.-M., Ourcival J.-M., Rodríguez-Calcerrada J. & Steppe K. (2017)
Stem hydraulic capacitance decreases with drought stress: implications for modelling
tree hydraulics in the Mediterranean oak Quercus ilex. *Plant, Cell & Environment* **40**,
1379–1391.

Santiago L.S., De Guzman M.E., Baraloto C., Vogenberg J.E., Brodie M., Hérault B., …
Bonal D. (2018) Coordination and trade-offs among hydraulic safety, efficiency and
drought avoidance traits in Amazonian rainforest canopy tree species. *New*
Phytologist, n/a-n/a.

Schneider C.A., Rasband W.S. & Eliceiri K.W. (2012) NIH Image to ImageJ: 25 years of
image analysis. *Nature Methods* **9**, 671–675.

Scholz F.C., Bucci S.J., Goldstein G., Meinzer F.C., Franco A.C. & Miralles-Wilhelm F.
(2008) Temporal dynamics of stem expansion and contraction in savanna trees:
withdrawal and recharge of stored water. *Tree Physiology* **28**, 469–480.

Scholz F.G., Bucci S.J., Goldstein G., Meinzer F.C., Franco A.C. & Miralles-Wilhelm F.
(2007) Biophysical properties and functional significance of stem water storage
tissues in Neotropical savanna trees. *Plant, Cell & Environment* **30**, 236–248.

Scholz F.G., Phillips N.G., Bucci S.J., Meinzer F.C. & Goldstein G. (2011) Hydraulic
Capacitance: Biophysics and Functional Significance of Internal Water Sources in
Relation to Tree Size. In *Size- and Age-Related Changes in Tree Structure and*
Function. Tree Physiology, (eds F.C. Meinzer, B. Lachenbruch & T.E. Dawson), pp.
341–361. Springer Netherlands.

Secchi F., Pagliarani C. & Zwieniecki M.A. (2017) The functional role of xylem parenchyma
cells and aquaporins during recovery from severe water stress. *Plant, Cell &*
Environment **40**, 858–871.

Sevanto S., Hölttä T. & Holbrook N.M. (2011) Effects of the hydraulic coupling between
xylem and phloem on diurnal phloem diameter variation: Xylem-phloem hydraulic
coupling and phloem diameter variation. *Plant, Cell & Environment* **34**, 690–703.

Siddiq Z., Zhang Y.-J., Zhu S.-D. & Cao K.-F. (2019) Canopy water status and
photosynthesis of tropical trees are associated with trunk sapwood hydraulic
properties. *Plant Physiology and Biochemistry* **139**, 724–730.

Sperry J.S., Nichols K.L., Sullivan J.E.M. & Eastlack S.E. (1994) Xylem embolism in ring-
porous, diffuse-porous, and coniferous trees of Northern Utah and Interior Alaska.
Ecology **75**, 1736–1752.

Steppe K. & Lemeur R. (2007) Effects of ring-porous and diffuse-porous stem wood anatomy
on the hydraulic parameters used in a water flow and storage model. *Tree Physiology*
27, 43–52.

Suzuki E. (1999) Diversity in specific gravity and water content of wood among Bornean
tropical rainforest trees. *Ecological Research* **14**, 211–224.

Trifilò P., Nardini A., Gullo M.A.L., Barbera P.M., Savi T. & Raimondo F. (2015) Diurnal
changes in embolism rate in nine dry forest trees: relationships with species-specific
xylem vulnerability, hydraulic strategy and wood traits. *Tree Physiology* **35**, 694–705.

Tyree M.T. & Yang S. (1990) Water-storage capacity of *Thuja*, *Tsuga* and *Acer* stems
measured by dehydration isotherms. *Planta* **182**, 420–426.

Umebayashi T., Utsumi Y., Koga S., Inoue S., Fujikawa S., Arakawa K., … Oda K. (2008)
Conducting pathways in north temperate deciduous broadleaved trees. *IAWA Journal*
29, 247–263.

Umebayashi T., Utsumi Y., Koga S., Inoue S., Matsumura J., Oda K., … Otsuki K. (2010)
Xylem water-conducting patterns of 34 broadleaved evergreen trees in southern
Japan. *Trees* **24**, 571–583.

Vergeynst L.L., Dierick M., Bogaerts J.A.N., Cnudde V. & Steppe K. (2015) Cavitation: a
blessing in disguise? New method to establish vulnerability curves and assess
hydraulic capacitance of woody tissues. *Tree Physiology* **35**, 400–409.

Warton D.I., Duursma R.A., Falster D.S. & Taskinen S. (2011) smatr 3– an R package for
estimation and inference about allometric lines. *Methods in Ecology and Evolution*.

Wei T. & Simko V. (2017) *R package "corrplot": visualization of a corelation matrix (Version*
0.84).

Wheeler E.A. (2011) InsideWood - a web resource for hardwood anatomy. *IAWA Journal* **32**,
199–211.

Wolfe B.T. & Kursar T.A. (2015) Diverse patterns of stored water use among saplings in
seasonally dry tropical forests. *Oecologia* **179**, 925–936.

Zanne A.E., Westoby M., Falster D.S., Ackerly D.D., Loarie S.R., Arnold S.E.J. & Coomes
D.A. (2010) Angiosperm wood structure: global patterns in vessel anatomy and their
relation to wood density and potential conductivity. *American Journal of Botany* **97**,
207–215.

Zhang Y.-J., Meinzer F.C., Qi J.-H., Goldstein G. & Cao K.-F. (2013) Midday stomatal
conductance is more related to stem rather than leaf water status in subtropical
deciduous and evergreen broadleaf trees. *Plant, Cell & Environment* **36**, 149–158.

Zheng J. & Martínez-Cabrera H.I. (2013) Wood anatomical correlates with theoretical
conductivity and wood density across China: evolutionary evidence of the functional
differentiation of axial and radial parenchyma. *Annals of Botany* **112**, 927–935.

Ziemińska K., Butler D.W., Gleason S.M., Wright I.J. & Westoby M. (2013) Fibre wall and
lumen fractions drive wood density variation across 24 Australian angiosperms. *AoB*
PLANTS **5**.

Ziemińska K., Westoby M. & Wright I.J. (2015) Broad Anatomical Variation within a Narrow
Wood Density Range—A Study of Twig Wood across 69 Australian Angiosperms.
PLoS ONE **10**, e0124892.

Zimmermann M.H. (1983) *Xylem Structure and the Ascent of Sap.* Springer-Verlag.

TABLES

Table 1. Species list, their abbreviations and porosity type.

Family	Species	Abbr	Porosity
Sapindaceae	*Acer saccharum*	ace	Diffuse
Sapindaceae	*Aesculus turbinata*	aes	Diffuse
Betulaceae	*Betula dahurica*	bet	Diffuse
Cercidiphyllaceae	*Cercidiphyllum japonicum*	cer	Diffuse
Cornaceae	*Cornus kousa*	cor	Diffuse
Eucommiaceae	*Eucommia ulmoides*	euc	Diffuse
Fagaceae	*Fagus grandifolia*	fag	Diffuse
Altingiaceae	*Liquidambar styraciflua*	liq	Diffuse
Magnoliaceae	*Liriodendron tulipifera*	lir	Diffuse
Magnoliaceae	*Magnolia cylindrica*	mag	Diffuse
Ericaceae	*Oxydendrum arboreum*	oxy	Diffuse
Theaceae	*Stewartia pseudocamellia*	ste	Diffuse
Styracaceae	*Styrax obassia*	sty	Diffuse
Tiliaceae	*Tilia japonica*	til	Diffuse
Fabaceae	*Albizia julibrissin*	alb	Semi-ring
Bignoniaceae	*Catalpa speciosa*	cat	Semi-ring
Fabaceae	*Cladrastis kentukea*	cla	Semi-ring
Ebenaceae	*Diospyros virginiana*	dio	Semi-ring
Fabaceae	*Gleditsia triacanthos*	gle	Semi-ring
Scrophulariaceae	*Paulownia tomentosa* var. 'Coreana'	pau	Semi-ring
Juglandaceae	*Carya laciniosa*	car	Ring
Oleaceae	*Fraxinus angustifolia* ssp. *oxycarpa*	fra	Ring
Moraceae	*Maclura pomifera*	mac	Ring
Moraceae	*Morus alba*	mor	Ring
Rutaceae	*Phellodendron amurense*	phe	Ring
Simaroubaceae	*Picrasma quassioides*	pic	Ring
Fagaceae	*Quercus muehlenbergii*	que	Ring
Lauraceae	*Sassafras albidum*	sas	Ring
Rutaceae	*Tetradium daniellii*	tet	Ring
Ulmaceae	*Zelkova sinica*	zel	Ring

948 Table 2. Traits, abbreviations and units. Fractions of wall (w), lumen (l) and wall+lumen (w+l)

949 of a given tissue were measured. Subscripts denote: lumen (L), predawn (pd), midday (md)

950 and difference between predawn and midday (pd-md).

Trait	Abbr	Units
Day capacitance	capacitance	kg m^{-3} MPa^{-1}
Cumulative water released	CWR$_{pd}$, CWR$_{md}$, ΔCWR$_{md-pd}$	kg m^{-3}
Stem water potential	Ψ$_{pd}$, Ψ$_{md}$, ΔΨ$_{pd-md}$	MPa
Volumetric water content	VWC$_{pd}$, VWC$_{md}$, ΔVWC$_{pd-md}$	Unitless
Volumetric water content: saturated	VWC$_{sat}$	"
Lumen volumetric water content	VWC$_{L-pd}$, VWC$_{L-md}$, ΔVWC$_{L-pd-md}$	"
Lumen volumetric water content: saturated	VWC$_{L-sat}$	"
Wall volumetric water content	VWC$_W$	"
Relative water content	RWC$_{pd}$, RWC$_{md}$, ΔRWC$_{pd-md}$	"
Lumen relative water content	RWC$_{L-pd}$, RWC$_{L-md}$, ΔRWC$_{L-pd-md}$	"
Wood density	WD	g cm^{-3}
Fibre fraction: w, l, w+l		Unitless
Axial+ray parenchyma fraction: w, l, w+l		"
Axial parenchyma fraction: w, l, w+l		"
Ray fraction: w, l, w+l		"
Vessel fraction: w, l, w+l		"
Living fibre fraction: w, l, w+l		"
Conduits$_{15}$ fraction: w+l		"
Vessel-fibre contact fraction		"
Vessel-axial+ray parenchyma contact fraction		"
Vessel-axial parenchyma contact fraction		"
Vessel-ray contact fraction		"
Vessel-vessel contact fraction		"
Vessel-conduits$_{15}$ contact fraction		"
Vessel mean area		µm^2
Vessel mean diameter		µm
Hydraulically weighted vessel diameter	D$_H$	"
Vessel number per area		mm^{-2}
Vessel size to number ratio		mm^2 mm^{-2}
Lumen fraction per total fibre area	lumen fraction$_{fibre}$	Unitless
Lumen fraction per total axial+ray parenchyma area	lumen fraction$_{axial+ray}$	"
Lumen fraction per total axial parenchyma area	lumen fraction$_{axial}$	"
Lumen fraction per total rays area	lumen fraction$_{ray}$	"
Lumen fraction per total vessels area	lumen fraction$_{vessel}$	"
Lumen fraction per total living fibres area	lumen fraction$_{living fibre}$	"

952
953

Table 3. Best models predicting wood capacitance, based on r^2 (for bivariate) or r_{adj}^2 (for multiple

models). Analysed were lumen fractions of: fibre, vessel, axial parenchyma, ray parenchyma,

axial+ray parenchyma, and contact fractions: vessel-fibre, vessel-vessel, vessel-axial parenchyma,

vessel-ray parenchyma, vessel-axial+ray parenchyma. Three best models with $P<0.1$ are shown. For

all species (n=29, excluding *Paulownia tomentosa*), we allowed maximum three explanatory variables,

and for diffuse-porous (n=14) and ring/semi-ring-porous (n=15) species, we allowed maximum two

explanatory variables, to avoid model overfitting. For models including vessel-fibre contact fraction

among ring/semi-ring-porous species one outlier was removed (*Sassafras albidum*, see Fig. 5). VWC_{L-pd}:

lumen volumetric water content at predawn, WD: wood density, axial: axial parenchyma. Bold: r^2 or

$r_{adj}^2>0.5$, regular: $0.4<r^2$ or $r_{adj}^2<0$, grey: r^2 or $r_{adj}^2<0.4$. Significance levels: '***' $P<0.001$, '**' $P<0.01$, '*'

$P<0.05$, '.' $P<0.1$, 'ns' $P>0.1$.

Species group	Model rank	Model predicting capacitance	r^2/r_{adj}^2	P
		Bivariate		
All species	1	- WD***	0.35	0.001
	2	+ VWC$_{L-pd}$**	0.29	0.003
Diffuse	1	- average vessel area*	0.30	0.041
	2	- WD.	0.25	0.069
Ring/semi-ring	1	+ VWC$_{L-pd}$**	0.41	0.010
	2	- WD*	0.37	0.016
		VWC$_{L-pd}$ + WD		
All species	na	+ VWC$_{L-pd}$* - WD**	0.44	0.000
Diffuse	na	+ VWC$_{L-pd}$ns - WD.	0.23	0.095
Ring/semi-ring	na	+ VWC$_{L-pd}$** - WD*	**0.59**	0.002
		VWC$_{L-pd}$ + WD + tissue fractions		
All species	1	+ VWC$_{L-pd}$** - WD** + axial lumen fractionns	0.47	0.000
	2	+ VWC$_{L-pd}$** - WD* - ray lumen fractionns	0.46	0.000
	3	+ VWC$_{L-pd}$** - WD* - vessel lumen fractionns	0.43	0.001
		VWCL-pd + WD + contact fractions		
All species	1	+ VWC$_{L-pd}$*** - WD*** + vessel-axial contact fraction**	**0.56**	0.000
	2	+ VWC$_{L-pd}$*** - WD*** - vessel-fibre contact fraction*	**0.54**	0.000
	3	+ VWC$_{L-pd}$** - WD** - vessel-ray contact fraction*	**0.54**	0.000
		VWC$_{L-pd}$ + tissue fractions		
All species	1	+ VWC$_{L-pd}$*** + fibre lumen fraction. - ray lumen fraction.	0.42	0.001
	2	+ VWC$_{L-pd}$*** + fibre lumen fraction** + axial lumen fractionns	0.40	0.001
	3	+ VWC$_{L-pd}$*** + fibre lumen fraction*	0.38	0.001
Diffuse	1	+ VWC$_{L-pd}$. + fibre lumen fraction.	0.25	0.081
Ring/semi-ring	1	+ VWC$_{L-pd}$** + vessel lumen fraction*	**0.55**	0.003
	2	+ VWC$_{L-pd}$** - axial+ray lumen fraction*	**0.55**	0.003
	3	+ VWC$_{L-pd}$** + fibre lumen fraction.	**0.50**	0.006
		VWC$_{L-pd}$ + contact fractions		
All species	1	+ VWC$_{L-pd}$*** + vessel-vessel contact fraction* + vessel-axial contact fraction*	0.42	0.001
	2	+ VWC$_{L-pd}$*** - vessel-ray contact fraction*	0.41	0.000
	3	+ VWC$_{L-pd}$*** + vessel-vessel contact fraction.	0.32	0.003
Diffuse	1	+ VWC$_{L-pd}$** - vessel-ray contact fraction**	**0.52**	0.007
	2	+ VWC$_{L-pd}$* + vessel-vessel contact fraction*	0.41	0.022
Ring/semi-ring	1	+ VWC$_{L-pd}$** + vessel-fibre contact fraction*	**0.62**	0.002
		VWC$_{L-pd}$+ tissue fractions + contact fractions		
All species	1	+ VWC$_{L-pd}$*** + fibre lumen fraction* - vessel-ray contact fraction*	**0.50**	0.000
	2	+ VWC$_{L-pd}$*** + fibre lumen fraction** + vessel-axial contact fraction*	0.48	0.000
	3	+ VWC$_{L-pd}$*** + fibre lumen fraction** - vessel-fibre contact fraction*	0.47	0.000

FIGURES

Figure 1. Illustration of the method used to estimate vessel-tissuecontact fractions between

vessels and other tissues (fibres, axial parenchyma, rays, vessels, conduits$_{15}$). The image

shows fragment of a cross-section of *Fraxinus angustifolia* ssp. *oxycarpa*. Yellow dots

distributed on vessel (black) circumference are classified based on the tissue in contact with

a vessel (1 - axial parenchyma, 2 - rays, 3 - fibres, 4 - vessels; conduits$_{15}$ - not shown).

Figure 2. Tissue fractions sorted along axial+ray parenchyma fraction (a) and vessel-tissue
contact fractions sorted along vessel-axial+ray parenchyma contact fraction (b). Three-letter
codes denote species (Table 1).

Figure 3. Boxplots illustrating differences between diffuse-porous (D, n=14), semi-ring-porous (SR, n=6) and ring-porous species (R, n=10). Circles denote species averages. Traits shown: wood density (WD), tissue fractions (F), vessel-tissue contact fractions (CF), hydraulically weighted diameter (D_H), vessel size to number ratio (S), predawn lumen volumetric water content (VWC_{L-pd}), predawn relative water content (RWC_{pd}), midday to predawn difference in cumulative water released (ΔCWR_{md-pd}) and capacitance. An outlier is *Paulownia tomentosa* (pau).

Figure 4. Relationships between tissue fractions (F): fibre fraction vs. vessel fraction (a), fibre

fraction vs. axial+ray parenchyma fraction (b), and vessel fraction vs. axial+ray parenchyma

fraction (c). Legends illustrate trait used to scale bubble size and porosity type.

Figure 5. Relationship between vessel-axial parenchyma contact fraction and vessel-fibre

contact fraction. CF: contact fraction. Legend displays porosity type. R^2 is across all species.

Outliers are *Quercus muehlenbergii* (que) and *Sassafras albidum* (sas). Legend illustrates

porosity type.

1000

1001
1002

1003   Figure 6. Barplot showing predawn and midday relative water content (RWC) ± 1 SD bars in

1004   diffuse-porous (a) and ring/semi-ring-porous species (b). Three-letter codes denote species

1005   (Table 1).

1006

1007
1008

1009   Figure 7. Relationships between water content and tissue lumen fractions (F): predawn

1010   lumen volumetric water content (VWC$_{L-pd}$) vs. vessel lumen fraction (a) and predawn lumen

1011   relative water content (RWC$_{L-pd}$) vs. fibre lumen fraction (b). Legend displays porosity type. r^2

1012   is across all species excluding *Paulownia tomentosa* (pau).

1013

1014
1015

1016 Figure 8. Relationships between capacitance and: wood density (WD, (a)), predawn lumen

1017 volumetric water content (VWC$_{L-pd}$, (b)), axial+ray parenchyma lumen fraction (axial+ray

1018 lumen F, (c)), and vessel lumen fraction (vessel lumen F, (d)). Legend displays porosity type.

1019 R^2 values are across all species, excluding *Paulownia tomentosa* (pau).

1020