Systematic cysteine-crosslinking in native membranes establishes the transmembrane architecture in Ire1 clusters

Kristina Väth[1,2], Roberto Covino[3], John Reinhard[1,2],

Gerhard Hummer[3,4], and Robert Ernst[1,2]

[1]Medical Biochemistry and Molecular Biology, Medical Faculty, Saarland University, Kirrberger Str. 100, Building 61.4, 66421 Homburg, Germany

[2]PZMS, Center for Molecular Signaling (PZMS), Medical Faculty, Saarland University, 66421 Homburg

[3]Department of Theoretical Biophysics, Max-Planck-Institute of Biophysics, Max-von-Laue-Strasse 3, 60438 Frankfurt am Main, Germany.

[4]Institute of Biophysics, Goethe-University, 60438 Frankfurt am Main, Germany.

*To whom correspondence should be addressed

robert.ernst@uks.eu

Running title: The TMHs of signaling-active Ire1 are X-shaped

Abstract

The endoplasmic reticulum (ER) is a key organelle of membrane biogenesis and crucial for the folding of both membrane and secretory proteins. Stress sensors of the unfolded protein response (UPR) monitor the unfolded protein load in the ER and convey effector functions for the maintenance of ER homeostasis. More recently, it became clear that aberrant compositions of the ER membrane, referred to as lipid bilayer stress, are equally potent activators of the UPR with important implications in obesity and diabetes. How the distinct signals from lipid bilayer stress and proteotoxic stress are processed by the highly conserved UPR transducer Ire1 remains unknown. Here, we have generated a functional, cysteine-less variant of Ire1 and performed systematic cysteine crosslinking experiments to establish the transmembrane architecture of signaling-active clusters in native membranes. We show that the transmembrane helices of two neighboring Ire1 molecules adopt an X-shaped configuration and that this configuration is independent of the primary cause for ER stress. Based on these findings, we propose that different forms of stress converge in a single, signaling-active conformation of Ire1.

Keywords

UPR, IRE1, PERK, lipid bilayer stress, UPR-transducer, cysteine crosslinking, membrane stress, ER stress, membrane quality control

Introduction

The endoplasmic reticulum (ER) marks the entry-point to the secretory pathway for soluble and membrane proteins. Under adverse conditions, accumulation of unfolded proteins causes ER stress and initiates the unfolded protein response (UPR) mediated by the Inositol-requiring enzyme 1 (Ire1) in budding yeast, and by the troika of IRE1α, the PKR-like Endoplasmic Reticulum Kinase (PERK), and the activating transcription factor 6 (ATF6) in vertebrates (1). Once activated, the UPR attenuates the production of most proteins and initiates a wide transcriptional program to upregulate ER chaperones, ER-associated degradation (ERAD), and lipid biosynthesis (2). Through these mechanisms, the UPR is centrally involved in cell fate decisions between life, death, and differentiation (3). Insulin-producing β-cells, for example, rely on UPR signals for their differentiation into professional secretory cells, while chronic ER stress caused by an excess of saturated fatty acids kills them (4). The transcriptional program of the UPR controls to expression of more than five percent of all genes with tremendous impact on virtually all aspects of cell physiology (2). Consistent with these broad effector functions, the UPR is associated with numerous diseases including diabetes, cancer, and neurodegeneration (5).

Ire1 is highly conserved among eukaryotes and represents the only transducer of ER stress in baker's yeast (6, 7). It is a type I transmembrane protein equipped with an ER-luminal sensor domain and two cytosolic effector domains: a kinase and an endoribonuclease (RNase) (8–10). How exactly unfolded proteins activate the UPR via direct and indirect mechanisms is a matter of active debate (11–15). ER stress caused by the accumulation of unfolded proteins leads to the oligomerization of Ire1 (16), which activates the cytosolic effector kinase and RNase domains (17). The unconventional splicing of the *HAC1* precursor mRNA initiated by the RNase domain facilitates the production of an active transcription factor that controls a broad spectrum of UPR-target genes via unfolded protein response elements (UPRE) in the promotor region (2, 18).

Lipid bilayer stress due to aberrant compositions of the ER membrane is equally potent in activating the UPR (19–21). This membrane-based mechanism is conserved throughout evolution (20, 22, 23) and has been implicated in the lipotoxicity associated with obesity and type II diabetes (4, 24). We have shown that Ire1 from baker's yeast inserts an amphipathic helix (AH) into the luminal leaflet of the ER-membrane, thereby forcing the adjacent TMH in a tilted configuration that locally squeezes the bilayer (25).

Aberrant stiffening of the ER membrane during lipid bilayer stress would increase the
energetic penalty for membrane deformation and thus stabilizes an oligomeric state of
Ire1 via a membrane-based mechanism (25, 26).

The accumulation of unfolded proteins in the ER can be triggered with
dithiothreitol (DTT) and Tunicamycin (TM), which interfere with protein folding in the
ER by inhibiting disulfide bridge formation and N-linked glycosylation, respectively.
Lipid bilayer stress is caused, for example, by an accumulation of saturated lipids or
by inositol-depletion (19, 21). While both proteotoxic and membrane-based stress lead
to the formation of Ire1 clusters (16, 25, 27), it remains unexplored if there is only one
signaling-active configuration of Ire1 is or if distinct conformations trigger custom-
tailored responses.

Here, we report on a systematic dissection of Ire1's TMH region domain in
signaling-active clusters. We have engineered a cysteine-less variant for genomic
integration at the endogenous locus, generated a series of constructs featuring single
cysteines in the TMH region, and developed a new crosslinking approach to study the
transmembrane configuration of Ire1 in the natural environment of ER-derived
membrane vesicles featuring the native complexity of lipids and proteins. This
approach uncovers the transmembrane architecture of Ire1 in signaling-active clusters
and suggests – irrespective of the primary cause of ER-stress (i.e. unfolded proteins
or lipid bilayer stress) an X-shaped configuration of the TMHs from neighboring Ire1
molecules. Our findings underscore the crucial importance of Ire1's highly bent
configuration in the TMH region for stabilizing an oligomeric state via a membrane-
mediated mechanism.

Results

We employed systematic cysteine-crosslinking in the TMH region of Ire1 to gain insight into the structural organization of its signaling-active clusters during ER-stress. Recognizing that Ire1 is activated by aberrant physicochemical membrane properties (25, 26), which are exceedingly hard to mimic using bottom-up approaches, we performed these crosslinking experiments in microsomes exhibiting the natural complexity of proteins and lipids of the ER membrane.

Cysteine-less Ire1 is functional

We have generated a cysteine-less version of Ire1 that would allow us to introduce single cysteine residues in the TMH region for a subsequent crosslinking using copper sulfate ($CuSO_4$). The cysteine-less construct is based on a previously established knock-in construct of *IRE1* that provides a homogeneous, near-endogenous expression level (25) and encodes for a fully-functional, epitope-tagged variant of Ire1, with an 3xHA tag and monomeric, yeast-enhanced GFP (yeGFP) inserted in a flexible loop at the position H875 (25, 28) (Figure 1A). To generate a cysteine-less version, we have substituted each of the twelve cysteines in the luminal, transmembrane and cytosolic regions of Ire1 with serine. Two cysteines in the signal sequence, which are co-translationally removed, remained in the final construct to ensure correct ER-targeting and membrane insertion (Figure 1A). Cysteine 48 of yeGFP ($C48^{yeGFP}$) was mutated to serine, while $C70^{yeGFP}$ is present in the cysteine-less construct to ensure correct folding of the fluorescent protein (29). Notably, $C70^{yeGFP}$ is buried inside the fluorescent protein (30) and thus inaccessible for crosslinking agents under non-denaturing conditions.

The steady-state level of wildtype and cysteine-less Ire1 are comparable (Figure S1A) and both proteins are properly integrated into the membrane as shown by subcellular fractionation (Figure S1B) and extraction assays (Figure S1C). The functionality of cysteine-less Ire1 was analyzed using a sensitive assay scoring the growth of cells exposed to chronic ER stress (25). Liquid cultures in minimal (synthetic complete dextrose; SCD) and full (yeast peptone dextrose; YPD) media were exposed to different concentrations of the reducing agent DTT and cultivated for 18 h prior to determining of the optical density (OD) of the culture. Cells producing either the wildtype or cysteine-less Ire1 are phenotypically indistinguishable and substantially

more resistant to DTT than cells lacking *IRE1* (Figure 1B). This suggests that the
cysteine-less variant of Ire1 is functional and capable to mount an adaptive UPR.
The functionality of the cysteine-less variant during ER-stress was further validated by
an alternative assay based on the degree of *HAC1* splicing (Figure 1C) and the mRNA
level of the UPR-target gene *PDI1* (Figure S1D). Exponentially growing cells were
acutely stressed for 1 h with either DTT or Tunicamycin, characterized by RT-qPCR,
and compared to unstressed cells. Expectedly, the spliced *HAC1* mRNA was
upregulated several-fold in stressed cells compared to unstressed cells and this
upregulation was observed in both wildtype and cysteine-less Ire1-producing cells.
(Figure 1C). Consequently, we also observed an upregulation of the mRNA of PDI1 in
response to ER-stress albeit to slightly lower degree for the cysteine-less version
compared to the wildtype construct (Figure S1D). The number of signaling-active
clusters per cell during ER-stress was almost identical for cells expressing either
wildtype and cysteine-less Ire1 (Figure S1E). We conclude that the cysteine-less
variant of Ire1 is functional during chronic and acute ER stress.
175
Crosslinking of Ire1's TMH in ER-derived microsomes
We established a strategy to crosslink a series of single-cysteine variants of clustered
Ire1 via copper sulfate ($CuSO_4$) in microsomes derived from the ER of stressed cells
(Figure 2A-C). This approach has several advantages over previous attempts and
provides unprecedented insights into structure-function relationships of this UPR-
transducer. Ire1 is studied 1) in the context of the full-length protein, 2) in ER-derived
microsomes featuring the native lipid environment, and 3) at a physiological protein-
to-lipid ratio with the full range of transmembrane proteins being present in the same
membrane. Using this system, we have studied the configuration of Ire1's TMH in
UPR-signaling clusters. Notably, these long-lived clusters are stable for several
minutes (16, 31). Thus, $CuSO_4$-mediated crosslinking that occurs on the same time-
scale can provide useful structural information even though it leads to the irreversible
formation of disulfide bonds under our experimental conditions.
Cells expressing either a cysteine-less variant of Ire1 or a variant containing a single-
cysteine in the TMH region (F544C) were cultivated to the mid-exponential phase in
minimal medium. These cells were either left untreated or stressed for 1 h with either
DTT (2 mM) or TM (1.5 µg/ml) to cause ER-stress, which leads to the formation of Ire1-
clusters (16, 25). We then isolated crude microsomes from these cells and incubated

them on ice for 5 min either in the presence or absence of 10 mM CuSO$_4$ to catalyze
the formation of disulfide bonds by oxidizing nearby sulfhydryl groups (32). Given the
low copy number of ~260 for Ire1 (33) and the fragmentation of the ER during
microsome preparation, we expect to detect crosslinking of single-cysteine variants of
Ire1 only when it was pre-clustered prior to the preparation.

Immunoblotting of the resulting samples revealed a prominent, HA-positive signal
corresponding to monomeric Ire1 and for some samples, a less-pronounced HA-
positive signal from a band with lower electrophoretic mobility that was only observed
when 1) Ire1 contained a single-cysteine in the TMH region (F544C), 2) the
microsomes were prepared from stressed cells (either DTT or TM), and 3) when
crosslinking was facilitated by CuSO$_4$ (Figure 2C). This suggests a remarkably specific
formation of covalent, disulfide bonds between two Ire1 molecules in the TMH region,
despite the presence of numerous other, potentially competing membrane proteins
with exposed cysteines in the ER. A Co-IP analysis using Flag-tagged and HA-tagged
Ire1 variants co-produced in a single cell and crosslinked in microsomes via the native
cysteine in the TMH region (C552) verified that the additional band with low
electrophoretic mobility represented presumably disulfide-linked, SDS-resistant dimers
of Ire1 (Figure S2). We conclude that Ire1 variants with single cysteines in the TMH
region can be specifically cross-linked via CuSO$_4$, when present as pre-formed clusters
in microsomes isolated from stressed cells.

A crosslinking screen identifies an X-shaped configuration of Ire1's TMH
Next, we systematically introduced single cysteine-residues in the TMH-region of Ire1
in order to characterize the structural configuration of Ire1's TMH in signaling-active
clusters. We generated a total of thirteen variants each containing a single cysteine in
the TMH region starting with E540C at the transition between the AH and the TMH
(Figure 3A) and ending at the residue C552, a native TMH residue that is substituted
to serine in the cysteine-less Ire1. Consequently, this scanning approach covered more
than three helical turns and almost the entire short TMH of Ire1 (Figure 3A, B). All
variants with engineered cysteine residues (E540C to F551C) were subjected to a
sensitive, cell-based assay to ascertain the functionality of the UPR under condition of
prolonged ER stress (Figure S3A). Consistent with the functional role of the AH in the
ER-luminal membrane leaflet (25), we found that the substitution of an AH-residue to
cysteine (E540C, T541C, or G542C) impaired the stress response of Ire1 as evident

from the increased sensitivity of the respective cells (Figure S3A). Notably, this
increased sensitivity to DTT is indeed due to a functional defect because the mutations
do not alter the steady-state level of the Ire1 variants (Figure S2B). In contrast,
substitution of TMH residues (V543C-F551C) did not result in an impaired resistance
to DTT in the respective cells (Figure S3A). This suggests that these TMH residues
have no specific relevance for Ire1 activation (Figure S3A). We then subjected this set
of single cysteine variants to the established cysteine-crosslinking procedure (Figure
3C). While some variants (e.g. G542C or L546C) showed no detectable crosslinking,
a significant portion of other variants (e.g. T541C or L549C) was crosslinked under the
given experimental conditions (Figure 3C,D). The F544C variant consistently exhibited
the highest crosslinking efficiency (Figure 3C,D). The overall pattern of crosslinking
residues in the TMH residues was independent of the condition of ER stress
(Figure 3D). Proteotoxic stress induced by either DTT, TM or lipid bilayer stress caused
by inositol-depletion (19, 34) showed essentially the same crosslinking pattern
(Figure S3C, 3D). These data suggest that the structural organization of Ire1 in
signaling-active clusters is comparable for different types of stress, at least in the TMH
region.
Cysteine crosslinking can be used to infer structural models. The observed pattern of
crosslinking residues in the TMH of Ire1 is very distinct to those observed in the TMH
region of the growth hormone receptor (35) and the thrombopoietin receptor (36),
which form parallel dimers leading to a helical periodicity of crosslinking. Our data also
disfavor a model of the two TMHs - attached to the ER-luminal and cytosolic portions
of Ire1 via flexible loops – forming dynamic complexes in an entirely random
orientation. Instead, our crosslinking data suggest an X-shaped configuration of the
TMHs with the best-crosslinking residue F554 positioned at the crossing point.
Intriguingly, such an arrangement would be consistent with the previously reported,
highly tilted orientation of the monomeric TMH of Ire1, which is enforced by the
adjacent, ER-luminal AH (25).
In order to obtain a structural representation of these crosslinking data, we modeled a
dimer of the TMH region of Ire1 based on extensive molecular dynamics (MD)
simulations in lipid membranes that integrated the crosslinking data, and in particular
the contact between the two F544 (Figure 4, Supplementary Movie) and restrained the
two F544 residues to face each other. The resulting model of the TMH region
highlighted the overall, highly bent architecture of the TMH region of each protomer

leading to an X-shaped configuration of the dimer (Figure 4, Supplementary Movie).
Substantial membrane thinning and water penetration, which were previously
observed around the monomeric TMH region of Ire1 (25), become clearly apparent
and are most significant in the region of the AHs (Figure S4, Supplementary Movie).
266

Validating the structural model of the TMH region of Ire1
Our crosslinking approach has identified the TMH residue F544 as a candidate residue
that might stabilize the transmembrane configuration of Ire1 in signaling-active
clusters. Interestingly, an aromatic residue in the TMH region of the mammalian IRE1α
(W547) was reported to stabilize the oligomeric state of IRE1α (37). Hence, we
embarked on testing a similar role for F544 in Ire1 from baker's yeast. We generated
a F544A mutation, which contained the native C552 in the TMH. A cell-based assay
revealed that the F544A mutant was phenotypically indistinguishable from cysteine-
less Ire1 (Figure 5A) just as the F544C mutant (Figure S1A), thereby suggesting that
F544 is not functionally relevant. This finding was corroborated by Cu^{2+}-mediated
crosslinking of C552 in microsomes isolated from cells stressed either with DTT or TM.
The intensity of the band corresponding to crosslinked Ire1 was unaffected by the
F544A mutation (Figure 5B). Thus, F544 does not contribute to the stability of Ire1-
oligomers even though it is placed in close proximity to equivalent residues on the TMH
of neighboring Ire1 molecules.
Our structural model of the monomeric TMH (25) and the dimeric TMH
(Figure 4, Supplementary Movie) suggested an unusual, tilted architecture of the TMH
region, which is stabilized by the ER-luminal AH. We proposed that this configuration
of the TMH region facilitates Ire1 to sense aberrant membrane properties by
contributing to the stability of oligomeric Ire1 during lipid bilayer stress (25, 38). We put
this model to the test, by combining the single-cysteine F544C for crosslinking with the
AH-disruption F531R mutation. The resulting F531R/F544C double mutant exhibited
only a very mild functional defect as implied by the cellular resistance to prolonged
stress caused by DTT (Figure 5C). Strikingly, the disruption of the AH by the F531R
mutation greatly impaired the crosslinking propensity of the F544C thereby suggesting
an important contribution of the AH to the transmembrane architecture of Ire1
(Figure 5D). An analogous set of experiments was performed with a construct that
contained an AH-disrupting F531R mutation (25) and, as a single cysteine the native
TMH residue C552. This F531R/C552 construct was functionally impaired

(Figure S5A) and showed a significantly reduced crosslinking propensity compared to
the equivalent construct with an intact AH (Figure S5B), thereby highlighting structural
and functional importance of the ER-luminal AH.
299
Discussion
Here, we establish a structural model for the TMH region of Ire1 in signaling-active
clusters (Figure 4). Employing a systematic cysteine crosslinking approach in native
membranes and aided by molecular dynamics simulations, we show that neighboring
TMHs of clustered Ire1 are organized in an X-shaped configuration during ER-stress.
Based on the remarkable similarity of the crosslinking patterns observed in the context
of lipid bilayer stress caused by inositol-depletion and proteotoxic stress caused by
DTT or TM (Figure 3, Figure S3C), we propose that different forms of ER-stress
converge in a single, signaling-active configuration in the TMH region. Our structural
and functional analyses reveal a crucial role of the transmembrane architecture in
controlling the oligomeric state of Ire1. The finding that the F544A mutation of the best-
crosslinking residue at the intersection of neighboring TMHs causes no functional
defect (Figure 5A,B) reinforces our interpretation that Ire1 senses lipid bilayer stress
via its unusual architecture rather than by stereospecific, structural elements in the
TMH.
Our findings are consistent with a crucial role of the ER-luminal AH in forcing
the adjacent TMH in a highly tilted and bent configuration and deforming the membrane
(25) (Figure 4, Figure S4, Supplementary Video). We show that disrupting the
functionally important AH by incorporation of a charged residue into the hydrophobic
face substantially reduces the crosslinking efficiency, thereby implying a significant
structural change (Figure 5C,D and Figure S5A,B).
Modeling the dimeric transmembrane organization aided by the crosslinking
data provides intriguing insights into membrane-deforming potential of Ire1 (Figure, 4,
Figure S4, Supplementary Video). The ER-luminal AH and a cluster of charged
residues at the cytosolic end of the TMH locally distort the bilayer, which is best
evidenced by a decreased membrane thickness in the proximity of the monomeric (25)
and dimeric transmembrane domain of Ire1 (Supplementary Video). Such membrane
deformations also cause disordering of lipid acyl chains and come at an energetic cost,
which is controlled by the composition and physical properties of the surrounding
bilayer (38, 39). The higher this cost e.g. due to increased lipid saturation or inositol-

depletion, the higher is the energetic gain upon coalescing these regions and thus the
propensity of Ire1 to oligomerize.

This membrane-based activation of the UPR is not restricted to yeast, but has
also been reported for UPR transducers in worms (23, 40), fishes (41), and mammals
(20, 37, 42). In fact, all three transducers of the UPR from mammals can be activated
by aberrant membrane signals albeit via distinct mechanisms (20, 37, 43). The IRE1α-
and PERK-dependent branches of the UPR respond to increased lipid saturation even
when the ER-luminal domain is removed (20). While some studies suggested that
stress-sensing relies on generic features in the TMH of mammalian IRE1α, which are
independent of the exact amino acid sequence in the TMH region (20, 44), a more
recent study proposed a mechanism of sensing that is based on specific residues in
the TMH (37). Cho *et al*. scored the oligomerization propensity of IRE1α in cells via
bimolecular fluorescence complementation in palmitate-treated cells and suggested
that the increased lipid saturation might induce a conformational switch in the TMH
region, which relies on a tryptophan (W457) as putative sensing residue and a
conserved leucine zipper motif (SxxLxxx) involving serine 450 (37, 44). Intriguingly,
such a rotation-based mechanism of sensing would be reminiscent of the lipid
saturation sensor Mga2 from baker's yeast controlling the expression of the essential
fatty acid desaturase-encoding gene *OLE1* (45, 46).

We propose that Ire1 from baker's yeast uses a mechanism distinct from that
proposed for mammalian IRE1α based on the following evidence: 1) Mutagenesis in
the entire TMH of Ire1 from baker's yeast causes no functional defect unless the
integrity of the AH is disrupted (25). 2) Mutagenesis of individual TMH residues (V543-
F551) including three aromatic residues lining one side of the TMH – one of which
being the best-crosslinking residue F544C - causes no relevant functional defects
(Figure S3A). These findings are notable because aromatic residues have been
implicated in lipid/membrane sensing in other systems (37, 45, 46). 3) Our crosslinking
data provide no evidence for a rotational re-organization of the TMHs during lipid
bilayer stress (Figure 3C,D, Figure S3C). 4) Our cell-based assays and crosslinking
data underscore the central importance of an intact AH adjacent to the TMH for normal
UPR activity and crosslinking in the TMH region.

Based on similar physicochemical features of the TMH region including the
evolutionary conserved AH, we propose that the mammalian PERK might use a similar
mechanism as Ire1 from baker's yeast, while the mammalian IRE1α -as a variation to

a common theme- seems to employ a distinct mechanism (37, 44). Clearly, more
studies with an increased temporal and spatial resolution that do not interfere with the
dynamic properties of Ire1 will be required to identify the unique specializations of
individual UPR transducers and to unequivocally establish their structural and dynamic
features that govern their membrane sensitivity.

Our combined results lead to the following model. The activity of Ire1 requires
an oligomerization of its cytosolic effector domains, which mediate the splicing of the
HAC1 in yeast of *XBP1* in mammals for signal propagation (1). This oligomerization
involves the entire protein as a response to divergent activating signals, which are
sensed by different portions of the protein.
Unfolded proteins are sensed by the ER-luminal domain via a binding groove (12),
which is formed by two interacting ER-luminal domains and is further facilitated by a
subsequent formation of higher oligomers (47). Under these conditions, the
transmembrane domain and the cytosolic effector domains 'follow' the oligomerization
of the ER-luminal domains. A large diversity of ER-luminal and cytosolic interactors
including chaperones can tune and specify the activity of mammalian UPR transducers
(14, 48). This may reflect a way to custom-tailor the globally-acting UPR to different
cell types with distinct protein folding requirements during steady-state and particularly
during differentiation. Lipid bilayer stress activates the UPR in both yeast and
mammals via a membrane-based mechanism and does not require the binding of
unfolded proteins to the ER-luminal domain and/or associated chaperones (19, 20,
25).

We propose that prolonged and chronic forms of ER-stress, which can be
mimicked by sub-lethal doses of DTT or TM, activate the UPR predominantly via this
membrane-based mechanism. When Ire1 is rendered insensitive to proteotoxic stress
by mutations in the binding groove for unfolded proteins (19, 25) and even when the
entire ER-luminal domain of IRE1α was substituted by a zipper domain (49) the
resulting proteins are activated under these conditions. We suggest that all portions of
Ire1, the ER-luminal domain, the TMD and the cytosolic domains cooperate in
determining the stability of oligomeric configurations of Ire1. An intriguing implication
of this model is that different forms of ER-stress would be independent of each other,
but inter-dependent in a somewhat additive fashion. Lipid bilayer stress would sensitize
Ire1 to unfolded proteins and, conversely, an increased load of unfolded proteins in the
ER lumen would increase the sensitivity of Ire1 to aberrant lipid compositions (20, 38).

The intriguing hypothesis that chemically distinct signals from the lumen and the
membrane of the ER cooperate in UPR activation remains to be experimentally tested.
Clearly, more studies are required to better understand the molecular
mechanisms of signal integration by UPR transducers particularly in the context of
prolonged and chronic ER stress. The clinical relevance is obvious as ER-stress and
UPR activation have been implicated the inflammatory response (50), B-cell
differentiation into plasma cells (51), insulin resistance, and pancreatic β-cell failure
(4). It has been speculated that the sensing of lipid bilayer stress by IRE1α and PERK
might account for instances, where relatively modest levels of unfolded protein load
lead to substantial UPR signaling and has been termed anticipatory UPR (52). An
exciting alternative possibility is that the membrane-based UPR may be caused by an
overcrowding of the ER membrane with membrane proteins (38, 53).
We have established a cysteine-less, functional variant of Ire1 allowing for a
detailed structure-function analysis of this UPR transducer in its native membrane
context. This construct and the established protocol for cysteine-crosslinking can be
readily adapted to address other central questions regarding the structural organization
of signaling-active clusters of Ire1, which may be inaccessible by other approaches.
415

Materials and Methods

Reagents and Antibodies

All chemicals and reagents used in this study were purchased from Sigma Aldrich, Carl Roth or Millipore and are of analytical or higher grade. The following antibodies were used: mouse anti-Flag monoclonal (M2) (Santa Cruz), rat anti-HA monoclonal (3F19) (Roche), mouse anti-Dpm1 monoclonal (5C5A7) (Life Technologies), mouse anti-Pgk1 (22C5D8) (Life Technologies), mouse anti-MBP monoclonal (NEB), anti-mouse-HRP (Dianova), anti-rat-HRP (Dianova).

Strains and plasmids

All strains used in this study are listed in Table 1, plasmids are listed in Table 2.

Genetic manipulation of *S. cerevisiae*

For genetic manipulation of *S. cerevisiae*, we used a previously described knock-in strategy (25). This construct comprised the endogenous promoter of *IRE1* (−1 to −551 bp), the sequence of *IRE1* with a sequence coding for a 3xHA tag and a monomeric version of yeGFP (A206R ^{yeGFP}) inserted at the position of H875, and the endogenous 5′ terminator sequence on the plasmid pcDNA3.1-IRE13xHA-GFP (25). A cysteine-less variant of this construct was generated by substituting all 12 cysteines in *IRE1* with serine using a strategy based on the QuikChange method (Stratagene). Additionally, the cysteine 48 (C48 ^{yeGFP}) of the monomeric yeGFP was substituted to serine, while the inaccessible, functionally relevant cysteine 70 (C70 ^{yeGFP}) remained in the final construct (29, 30). Derivatives of this constructs were generated by substituting individual codons of TMH residues to cysteine. All mutants used in this study are listed in Table 2. These plasmids were linearized using Hind*III* and Xho*I* restriction enzymes and used for transforming a previously established cloning strain lacking both the gene of IRE1 and its promotor (25). Strains used in this study are listed in Table 1.

Additionally, a Flag-tagged cysteine-less Ire1 version based on the CEN-based Ire1 construct from the pPW1628/pEv200 plasmid (28) was generated. The 3xHA epitope tag in the knock-in construct was replaced by a 3xFlag epitope tag using the Q5 site-directed mutagenesis kit (NEB). The newly generated knock in sequence was amplified in a multi-step PCR reaction adding the terminator sequence from the

pEv200 plasmid and *BssH*I and *Hind*III restriction site. The transfer of the *IRE1$_{3xFlag-GFP}$*

sequence in the *CEN*-based vector pPW1628/pEv200 (28) was performed using

*BssH*I/*Hind*III restriction sites.

Cultivation of *S. cerevisiae*

The yeast strains were cultivated at 30°C on agar plates containing SCD

complete medium or selection medium. Liquid yeast cultures were inoculated with a

single colony and typically cultivated at 30°C for a minimum of 18 h to reach the

stationary phase. This overnight culture was used to inoculate a fresh culture, which

was inoculated to an OD_{600} = 0.2 and cultivated until the mid-exponential cultures.

For microsomal membrane preparation, stationary cells were used to inoculate

a fresh culture in SCD complete medium to an OD_{600} of 0.2. After cultivation at 30°C

to an OD_{600} of 0.7, the cells were either left untreated or stressed with either 2 mM DTT

or 1.5 µg/ml Tunicamycin for 1 h. For inositol depletion, exponentially growing cells

were washed with inositol-depleted media and used to inoculate the main culture to an

OD_{600} of 0.5 in SCD complete w/o inositol, which was further cultivated for 3 h.

Live cell confocal microscopy

A fresh culture in SCD medium was inoculated to an OD_{600} = 0.2 and cultivated

for 5.5 h. To induce ER-stress, DTT was added to a final concentration of 2 mM

followed by additional cultivation for 1 h. The cells were harvested placed on

microscopic slides coated with a thin layer of SCD containing 1.5% agarose for

immobilization. Microscopy was performed using a Zeiss LSM 780 confocal laser

scanning microscope (Carl Zeiss AG) with spectral detection and a Plan-Apochromat

63x 1.40 NA oil immersion objective. GFP fluorescence was excited at 488 nm and the

emission was detected between 493 and 598 nm. Transmission images were

simultaneously recorded using differential interference contrast (DIC) optics. Z-stacks

(450 nm step-size, 62.1 µm pinhole size) were recorded. Ire1-clusters were identified

by automated segmentation using CellProfiler (54). In brief, cellular areas of the field

of view were determined based on sum projections of recorded z-stacks. Clusters

within these cells were identified by adaptive thresholding of the respective maximum

intensity z-projections.

Preparation of cell lysates and immunoblotting

Lysates were prepared from exponentially growing cells, which were harvested by centrifugation (3.000xg, 5 min, 4°C) and then washed once with ddH$_2$O and once with PBS. During washing, the cells were transferred into 1.5 ml reaction tubes allowing for a more rapid centrifugation (8.000xg, 20 sec, 4°C). The tubes with the washed cell pellet were placed in a -80°C freezer and stored until further use.

For preparing a cell lysate, either 5 or 20 OD equivalents were resuspended in 400 µl or 1000 µl lysis buffer (PBS containing 30 µg/ml protease inhibitor cocktail), respectively. After addition of either 100 µl or 500 µl of zirconia beads, respectively, the cells were disrupted by bead beating for 5 min at 4°C. Four volumes units of the resulting lysate were mixed with one volume of 5x reducing sample buffer (8M urea, 0.1 M Tris-HCl pH 6.8, 5 mM EDTA, 3.2% (w/v) SDS, 0.15% (w/v) bromphenol blue, 4% (v/v) glycerol, 4% (v/v) β-mercaptoethanol) and then incubated at 95°C for 10 min for fully unfolding and solubilizing the proteins therein. 0.1 OD equivalents of the resulting sample was subjected to SDS-PAGE and the proteins were separated on 4-15% Mini-PROTEAN-TGX strain-free gels (BioRad). For subsequent immuno-blotting, proteins were transferred from the gel to methanol-activated PVDF membranes using semi-dry Western-Blotting. Specific proteins were detected using antigen-specific primary antibodies, HRP-coupled secondary antibodies, and chemiluminescence.

Assaying the resistance to ER-stress

Stationary overnight cultures were used to inoculate a fresh culture to an OD$_{600}$ of 0.2. After cultivation for 5 to 7 h at 30°C the cells were diluted with pre-warmed medium to an OD$_{600}$ of 0.05. 50 µl of these diluted cultures were mixed in a 96-well plate with 180 µl of medium and 20 µl of a DTT dilution series leading to a final concentration of DTT between 0 and 2 mM and 0 and 4 mM, respectively. After incubation at 30°C for 18 h, the cultures were thoroughly mixed and 200 µl of the cell suspension were transferred to a fresh 96-well plate for determining the density of the culture via a photospectrometer using the OD$_{600}$/OD$_{620}$.

RNA preparation, cDNA synthesis and quantitative real-time (qPCR) PCR analysis

Exponentially growing cells were used to inoculate a 20 ml culture to an OD_{600} of 0.2 in YPD. After cultivation for 3 h, the cells were either left untreated or stressed with either 4 mM DTT or 1.0 µg/ml Tunicamycin. After an additional cultivation for 1 h, 5 OD equivalents were harvested by centrifugation (3.000xg, 5 min, RT). The supernatant was discarded while the pellet was snap frozen in liquid N_2 and stored at -80°C. The RNA was prepared from these cells using the RNeasy Plus RNA Isolation Kit (Qiagen). 500 ng RNA of the total isolated RNA were used as a template for the synthesis of cDNA using Oligo(dT) primers and the Superscript^{TM} II RT protocol (Invitrogen). qPCR was performed using ORA qPCR Green ROX L Mix (HighQu) in a Piko Real PCR system (Thermo Scientific). The following primers were used to amplify DNA sequences of interest:

HAC1s forward primer:	5' – CTTTGTCGCCCAAGAGTATGCG – 3'
HAC1s reverse primer:	5' – ACTGCGCTTCTGGATTACGC – 3'
ACT1 forward primer:	5' – TGTCACCAACTGGGACGATA – 3'
ACT1 reverse primer:	5' – AACCAGCGTAAATTGGAACG – 3'

All reactions were performed in technical duplicates. Non-template controls (RNA) and non-reaction controls (H_2O) were additionally performed. The qPCR program included the following steps: 1) 95°C, 15 min; 2) 95°C, 20 sec; 3) 58°C, 20 sec; 4) 72°C, 30 sec; 5) 72°C, 5 min; steps 2-4 were repeated 40 times. The relative quantification of *HAC1* splicing levels was based on the comparative $\Delta\Delta$CT method using normalization to *ACT1* levels (StepOnePlus^{TM} user Manual, Applied Biosystems).

Microsomal membrane preparation

80 OD_{600} equivalents were harvested from a mid-exponential culture by centrifugation (3.000xg, 5 min, 4°C), washed with PBS, and stored at -80°C. All steps of membrane fractionation were performed on ice or at 4°C with pre-chilled buffers. The cell pellets were resuspended in 1.5 ml lysis buffer (50 mM HEPES pH 7.0, 150 mM NaCl, 1 mM EDTA, 10 µg/ml chymostatin, 10 µg/ml antipain, 10 µg/ml pepstatin). For cysteine crosslinking experiments, a lysis buffer without EDTA was used. Cells were disrupted by using zirconia beads (Roth) and a bead beater (2 x 5 min). Cell debris was removed from the lysate by centrifugation (800x g, 5 min, 4°C) and further cleared from major mitochondrial contaminations (5,000 x g, 10 min, 4°C). The pellet

551　was discarded, and the supernatant was spun again (100.000x g, 45 min, 4°C) to
552　obtain a crude microsomal fraction in the pellet. The microsomes were resuspended
553　in 1.4 ml lysis buffer, sonicated for homogenization (50%, 5x1sec, MS72 tip on a
554　sonifier cell disrupter from Branson Ultrasonic) snap frozen in liquid N_2 in small aliquots
555　to avoid freeze-thaw cycles, and stored at -80°C for further use.
556

Test of membrane association

The cleared supernatant of a 5.000xg step was divided into equal parts, which
were then mixed with an equal volume of lysis buffer supplemented with 0.2 M Na_2CO_3
resulting in a final pH of 11, 5 M urea, 2% Triton X-100 or without additional additives.
After incubation for 1 h on a rotator, these samples were centrifuged (100,000x g, 45
min, 4°C) to separate soluble from insoluble material. The supernatant and pellets from
these fractions corresponding to 0.2 OD equivalents were further analyzed by SDS-
PAGE and immunoblotting.
565

CuSO$_4$-induced cysteine crosslinking

For cysteine crosslinking with CuSO$_4$, the microsomes were thawed on ice. 8 µl
of microsomes with a protein concentration of 1 ± 0.2 mg/ml were then incubated with
2 µl of 50 mM CuSO$_4$ in ddH$_2$O resulting in a final concentration of 10 mM CuSO$_4$ and
incubated for 5 min. A sample supplemented with 2 µl ddH$_2$O served as a non-
crosslinking control. The reaction was stopped with 8 µl of membrane sample buffer
(4 M urea, 50 mM Tris-HCl pH 6.8, 1.6 % (w/v) SDS, 0.01% (w/v) bromophenol blue,
2% (v/v) glycerol) including 125 mM EDTA and 250 mM NEM. Samples were heated
for 5 min at 95°C and further analyzed by SDS-PAGE and immunoblotting. The
percentage of dimer formation was determined using the densiometric signals of the
bands corresponding to the monomeric and covalently dimeric protein, which were
determined using ImageJ.
578

Immunoprecipitation from microsomes after CuSO$_4$-induced cysteine
crosslinking

300 µl of microsomes with a typical protein concentration of 1 mg/ml were
incubated with 12.5 µl 250 mM CuSO$_4$ (final concentration of 10 mM) for 5 min on ice.
The reaction was stopped by adjusting the sample to a final concentration of 50 mM
EDTA and 111 mM NEM by adding 30 µl of 0.5 M EDTA stock solution and 44 µl of 1

M NEM stock solution, respectively. The final volume was adjusted to 1.3 ml with lysis
buffer with a final concentration of 5 mM EDTA. The $CuSO_4$ concentration was thus
reduced to 2.4 mM and the NEM concentration to 33.6 mM, respectively.

After crosslinking, the microsomes were solubilized using 2% Triton X-100 and
incubated for 1 h at 4°C under constant agitation. Insoluble material was removed by
centrifugation (20.000x g, 10 min, 4°C). The resulting supernatant was incubated with
8 µl Flag beads (Sigma Aldrich), equilibrated with IP wash buffer (lysis buffer + 5 mM
EDTA + 0.2 % Triton X-100), for 3 h under constant shaking. Flag beads were washed
five times with IP wash buffer by centrifugation (8.000xg, 30 sec, 4°C). For elution, the
Flag beads were incubated with 10 µl IP-Wash and 10 µl 5x reducing sample buffer
for 5 min at 95°C, which did not disrupt the disulfide bond formed between to protomers
of Ire1. These samples were analyzed by SDS-PAGE and immunoblotting.

Modelling of the dimer

We extracted an equilibrated conformation of a monomeric 56 amino-acid long
sequence 516-SRELD EKNQNSLLLK FGSLVYRIIE TGVFLLLFLI FCAILQRFKI
LPPLYVLLSK I-571 from a previously performed 10 µs long equilibrium MD simulation.
We duplicated the configuration in order to create a new system containing two
identical protomers. We then rotated and translated one of the two protomers to form
a dimer structure, such that the two F544 faced each other with the distance between
their Cβ atoms at around 0.7 nm. A short energy minimization in solution resolved all
steric clashes between side-chains. The structure of the model dimer was prepared by
using gromacs/2019.3 tools (55) and VMD (56). We used Charmm-GUI (57, 58) to
reconstitute the dimer in a bilayer containing 248 POPC and 62 cholesterol molecules
modelled in the Charmm36m force-field (59–61). We solvated the system with 24813
TIP3P water molecules, 72 chloride and 66 sodium ions, corresponding to a salt
concentration of 150 mM.

Equilibrium and restrained simulations of the dimer model

After an initial energy minimization and quick relaxation, we carefully
equilibrated the dimer model in the bilayer. We first ran a 50 ns long simulation
restraining the position of protein atoms by using harmonic potentials with force-
constants (in units of kJ mol^{-1} nm^2) of 500 for backbone atoms and 200 for side-chain
atoms. We then ran further 50 ns lowering the force-constants to 200 and 50,

respectively. After this equilibration, we relieved all restraints and ran two independent
500 ns long MD simulations. In the first simulation, the system evolved according to its
unbiased dynamics; in the second, we restrained the distance between the two Cβ
atoms of F544 on each protomer, gradually driving them closer in time. We first
centered the restraining harmonic potential around 0.5 nm with a force-constants (in
units of kJ mol^{-1} nm^2) of 1000. We then lowered the equilibrium distance of the potential
and decreased the force-constants (in units of kJ mol^{-1} nm^2): 0.4 nm and 750 after
10 ns from the beginning of the run; 0.3 nm and 500 after 20 ns; and 0.2 nm and 500
after 30 ns and for the remaining duration of the simulation.

We ran both the restrained equilibration and unbiased production simulation in
gromacs/2019.3, and the restrained production simulation in gromacs/2018.7 patched
with the open-source, community-developed PLUMED library (62) version 2.5 (63),
using a time step of 2 fs. Electrostatic interactions were evaluated with the Particle-
Mesh-Ewald method (64). We maintained a constant temperature of 303 K (65),
applying separate thermostats on the protein, membrane, and solvent with a
characteristic time of 1 ps. We applied the semi-isotropic Berendsden barostat (66) for
the restrained equilibration, and the Parrinello-Rahman barostat (67) for the production
runs, acting separately on the x-y plane and z direction to maintain a constant pressure
of 1 atm, and with a characteristic time of 5 ps. We constrained all hydrogen bonds
with the LINCS algorithm (68). Molecular visualizations were obtained with VMD and
rendered with Tachyon.
640

Data analysis, representation, and statistical tests
The amount of crosslinked dimer was determined using the densiometric signals of the
bands corresponding to the monomeric and covalently dimeric protein. The percentage
was calculated from the ratio of dimer signal to whole Ire1 signal. The software Image
J was used for the determination if the densiometric signal.
Throughout the manuscript, the data are represented as the average ± SEM. All
experiments were performed at least in triplicates. The significance was tested by
unpaired Student's t tests. Further information regarding statistical analysis can be
found in the figure legends. In the following, the exact value of n is provided for every
dataset of this paper.

Fig. 1B: left panel: Δ*IRE1*, WT: n=21 (technical replicates from four individual
colonies); cysteine-less: n=12 (technical replicates from four individual
colonies)
right panel: Δ*IRE1*: n=15 (technical replicates from three individual
colonies); cysteine-less: n=12 (technical triplicates from four individual
colonies); WT: n=9 (technical triplicates from three individual colonies)
Fig. 1C: left panel: WT -DTT: n= 4; WT +DTT: n=6; cysteine-less -DTT: n=6;
cysteine-less +DTT: n=5
Right panel: WT -TM: n=4; WT +TM: n=5; cysteine-less -TM: n=6; cysteine-
less +TM: n=4
Fig. 3D: DTT: E540C, T541C: n=8 (technical duplicates from four individual
colonies); G542C, V543C, L545C, L546C, L547C, F548C, L549C, I550C,
F551C: n=4 (technical duplicates from two individual colonies); F544C:
n=14 (technical duplicates from seven individual colonies); C552: n=11
(technical duplicates from six individual colonies)
TM: E540C, T541C: n=8 (technical duplicates from four individual colonies);
G542C, V543C, L545C, L546C, L547C, F548C, L549C, I550C, F551C: n=4
(technical duplicates from two individual colonies); F544C: n=10 (technical
duplicates from five individual colonies); C552: n=8 (technical duplicates
from four individual colonies)
Inositol depletion: n=6 (technical duplicates from three individual colonies)
Fig. 5A: Δ*IRE1*, cysteine-less: n=12 (technical replicates from two individual
colonies), F544A C552: n=12 (technical triplicates from four individual
colonies)
Fig. 5B: DTT: C552: n=11 (technical duplicates from six individual colonies), F544A
C552: n=4 (technical duplicates from two individual colonies)
TM: C552: n=8 (technical duplicates from four individual colonies), F544A
C552: n=4 (technical duplicates from two individual colonies)
Fig. 5C: Δ*IRE1*, cysteine-less: n=12 (technical replicates from two individual
colonies), F531R F544C: n=12 (technical triplicates from four individual
colonies)
Fig. 5D: DTT: F544C: n=14 (technical duplicates from seven individual colonies),
F531R F544C: n=4 (technical duplicates from two individual colonies)

684		TM: F544C n=10 (technical duplicates from five individual colonies), F531R
685		F544C: n=4 (technical duplicates from two individual colonies)
686	**Fig. S1B**:	left panel: WT -DTT: n= 5; WT +DTT: n=5; cysteine-less -DTT: n=6; cysteine-
687		less +DTT: n=6
688		Right panel: WT -TM: n=5; WT +TM: n=5; cysteine-less -TM: n=6; cysteine-
689		less +TM: n=3
690	**Fig. S3A**:	OD_{620}: $\Delta IRE1$: n=9 (technical triplicates from three individual colonies):
691		cysteine-less: n=12 (technical triplicates from four individual colonies);
692		E540C – L546C: n=6 (technical triplicates from two individual colonies)
693		OD_{600}: $\Delta IRE1$: n=10 (technical replicates from three individual colonies):
694		cysteine-less: n=22 (technical triplicates from four individual colonies);
695		L547C – F551: n=6 (technical triplicates from two individual colonies)
696	**Fig. S4A**:	$\Delta IRE1$, cysteine-less: n=12 (technical replicates from two individual
697		colonies), F544R C552: n=9 (technical triplicates from three individual
698		colonies)
699	**Fig. S5B**:	DTT: C552: n=11 (technical duplicates from six individual colonies), F531R
700		C552: n=4 (technical duplicates from two individual colonies)
701		TM: C552: n=8 (technical duplicates from four individual colonies), F531R
702		C552: n=4 (technical duplicates from two individual colonies
703		

References

1. Walter P, Ron D (2011) The unfolded protein response: from stress pathway to

homeostatic regulation. *Science* 334(6059):1081–1086.

2. Travers KJ, et al. (2000) Functional and genomic analyses reveal an essential

coordination between the unfolded protein response and ER-associated

degradation. *Cell* 101(3):249–258.

3. Hetz C (2012) The unfolded protein response: Controlling cell fate decisions

under ER stress and beyond. *Nat Rev Mol Cell Biol* 13(2):89–102.

4. Fonseca SG, Burcin M, Gromada J, Urano F (2009) Endoplasmic reticulum

stress in β-cells and development of diabetes. *Curr Opin Pharmacol* 9(6):763–

770.

5. Kaufman RJ (2002) Orchestrating the unfolded protein response in health and

disease. *J Clin Invest* 110(10):1389–1398.

6. Nikawa J, Yamashita S (1992) IRE1 encodes a putative protein kinase

containing a membrane-spanning domain and is required for inositol

phototrophy in Saccharomyces cerevisiae. *Mol Microbiol* 6(11):1441–1446.

7. Kimata Y, Kohno K (2011) Endoplasmic reticulum stress-sensing mechanisms

in yeast and mammalian cells. *Curr Opin Cell Biol* 23(2):135–142.

8. Cox JS, Shamu CE, Walter P (1993) Transcriptional induction of genes

encoding endoplasmic reticulum resident proteins requires a transmembrane

protein kinase. *Cell* 73(6):1197–1206.

9. Sidrauski C, Walter P (1997) The transmembrane kinase Ire1p is a site-specific

endonuclease that initiates mRNA splicing in the unfolded protein response.

Cell 90(6):1031–1039.

10. Mori K, Ma W, Gething MJ, Sambrook J (1993) A transmembrane protein with

a cdc2+/CDC28-related kinase activity is required for signaling from the ER to

the nucleus. *Cell* 74(4):743–756.

11. Karagöz GE, et al. (2017) An unfolded protein-induced conformational switch

activates mammalian IRE1. *Elife* 6:e30700.

12. Gardner BM, Walter P (2011) Unfolded proteins are Ire1-activating ligands that

directly induce the unfolded protein response. *Science* 333(6051):1891–1894.

13. Adams CJ, Kopp MC, Larburu N, Nowak PR, Ali MMU (2019) Structure and

Molecular Mechanism of ER Stress Signaling by the Unfolded Protein

Response Signal Activator IRE1. *Front Mol Biosci* 6:11.

14. Amin-Wetzel N, et al. (2017) A J-Protein Co-chaperone Recruits BiP to
Monomerize IRE1 and Repress the Unfolded Protein Response. *Cell*
171(7):1625-1637.e13.
15. Bertolotti A, Zhang Y, Hendershot LM, Harding HP, Ron D (2000) Dynamic
interaction of BiP and ER stress transducers in the unfolded-protein response.
Nat Cell Biol 2(6):326–332.
16. Kimata Y, et al. (2007) Two regulatory steps of ER-stress sensor Ire1 involving
its cluster formation and interaction with unfolded proteins. *J Cell Biol*
179(1):75–86.
17. Korennykh A V, et al. (2009) The unfolded protein response signals through
high-order assembly of Ire1. *Nature* 457(7230):687–93.
18. Mori K, et al. (1992) A 22 bp cis-acting element is necessary and sufficient for
the induction of the yeast KAR2 (BiP) gene by unfolded proteins. *EMBO J*
11(7):2583–2593.
19. Promlek T, et al. (2011) Membrane aberrancy and unfolded proteins activate
the endoplasmic reticulum stress sensor Ire1 in different ways. *Mol Biol Cell*
22(18):3520–3532.
20. Volmer R, van der Ploeg K, Ron D (2013) Membrane lipid saturation activates
endoplasmic reticulum unfolded protein response transducers through their
transmembrane domains. *Proc Natl Acad Sci U S A* 110(12):4628–4633.
21. Surma MA, et al. (2013) A lipid E-MAP identifies Ubx2 as a critical regulator of
lipid saturation and lipid bilayer stress. *Mol Cell* 51(4):519–530.
22. Ho N, Xu C, Thibault G (2018) From the unfolded protein response to
metabolic diseases – lipids under the spotlight. *J Cell Sci* 131(3):jcs199307.
23. Hou NS, et al. (2014) Activation of the endoplasmic reticulum unfolded protein
response by lipid disequilibrium without disturbed proteostasis in vivo. *Proc*
Natl Acad Sci U S A 111(22):E2271-2280.
24. Pineau L, Ferreira T (2010) Lipid-induced ER stress in yeast and β cells:
Parallel trails to a common fate. *FEMS Yeast Res* 10(8):1035–1045.
25. Halbleib K, et al. (2017) Activation of the Unfolded Protein Response by Lipid
Bilayer Stress. *Mol Cell* 67(4):673-684.e8.
26. Ernst R, Ballweg S, Levental I (2018) Cellular mechanisms of physicochemical
membrane homeostasis. *Curr Opin Cell Biol* 53:44–51.
27. Li H, Korennykh A V, Behrman SL, Walter P (2010) Mammalian endoplasmic

reticulum stress sensor IRE1 signals by dynamic clustering. *Proc Natl Acad Sci*
U S A 107(37):16113–16118.

28. van Anken E, et al. (2014) Specificity in endoplasmic reticulum-stress signaling
in yeast entails a step-wise engagement of HAC1 mRNA to clusters of the
stress sensor Ire1. *Elife* 3:e05031.

29. Costantini LM, et al. (2015) A palette of fluorescent proteins optimized for
diverse cellular environments. *Nat Commun* 6(1):7670.

30. Orm M, et al. (1996) Crystal Structure of the Aequorea victoria Green
Fluorescent Protein. *Science (80-)* 273(5280):1392–1395.

31. Cohen N, et al. (2017) Iron affects Ire1 clustering propensity and the amplitude
of endoplasmic reticulum stress signaling. *J Cell Sci* 130(19):3222–3233.

32. Kobashi K (1968) Catalytic oxidation of sulfhydryl groups by o-phenanthroline
copper complex. *BBA - Gen Subj* 158(2):239–245.

33. Ghaemmaghami S, et al. (2003) Global analysis of protein expression in yeast.
Nature 425(6959):737–741.

34. Lajoie P, Moir RD, Willis IM, Snapp EL (2012) Kar2p availability defines distinct
forms of endoplasmic reticulum stress in living cells. *Mol Biol Cell* 23(5):955–
64.

35. Brooks AJ, et al. (2014) Mechanism of Activation of Protein Kinase JAK2 by
the Growth Hormone Receptor. *Science (80-)* 344(6185):1249783–1249783.

36. Matthews EE, et al. (2011) Thrombopoietin receptor activation: transmembrane
helix dimerization, rotation, and allosteric modulation. *FASEB J* 25(7):2234–
2244.

37. Cho H, et al. (2019) Intrinsic Structural Features of the Human IRE1α
Transmembrane Domain Sense Membrane Lipid Saturation. *Cell Rep*
27(1):307-320.e5.

38. Covino R, Hummer G, Ernst R (2018) Integrated Functions of Membrane
Property Sensors and a Hidden Side of the Unfolded Protein Response. *Mol*
Cell 71(3):458–467.

39. Radanović T, Reinhard J, Ballweg S, Pesek K, Ernst R (2018) An emerging
group of membrane property sensors controls the physical state of organellar
membranes to maintain their identity. *BioEssays* 40(5):e1700250.

40. Koh JH, Wang L, Beaudoin-Chabot C, Thibault G (2018) Lipid bilayer stress-
activated IRE-1 modulates autophagy during endoplasmic reticulum stress. *J*

Cell Sci 131(22):jcs217992.

41. Thakur PC, Davison JM, Stuckenholz C, Lu L, Bahary N (2014) Dysregulated
phosphatidylinositol signaling promotes endoplasmic-reticulum-stress-
mediated intestinal mucosal injury and inflammation in zebrafish. *Dis Model*
Mech 7(1):93–106.

42. Kitai Y, et al. (2013) Membrane lipid saturation activates IRE1α without
inducing clustering. *Genes to Cells* 18(9):798–809.

43. Tam AB, et al. (2018) The UPR Activator ATF6 Responds to Proteotoxic and
Lipotoxic Stress by Distinct Mechanisms. *Dev Cell* 46(3):327-343.e7.

44. Kono N, Amin-Wetzel N, Ron D (2017) Generic membrane-spanning features
endow IRE1α with responsiveness to membrane aberrancy. *Mol Biol Cell*
28(17):2318–2332.

45. Covino R, et al. (2016) A eukaryotic sensor for membrane lipid saturation. *Mol*
Cell 63(1):49–59.

46. Ballweg S, Sezgin E, Wunnicke D, Haenelt I, Ernst R (2019) Regulation of lipid
saturation without sensing membrane fluidity.
doi:https://doi.org/10.1101/706556.

47. Korennykh A, Walter P (2012) Structural Basis of the Unfolded Protein
Response. *Annu Rev Cell Dev Biol* 28(1):251–77.

48. Sepulveda D, et al. (2018) Interactome Screening Identifies the ER Luminal
Chaperone Hsp47 as a Regulator of the Unfolded Protein Response
Transducer IRE1α. *Mol Cell* 69(2):238-251.e8.

49. Liu CY, Schroder M, Kaufman RJ (2000) Ligand-independent dimerization
activates the stress response kinases IRE1 and PERK in the lumen of the
endoplasmic reticulum. *J Biol Chem* 275(32):24881–24885.

50. Zhang K, Kaufman RJ (2008) From endoplasmic-reticulum stress to the
inflammatory response. *Nature* 454(7203):455–462.

51. Hu CCA, Dougan SK, McGehee AM, Love JC, Ploegh HL (2009) XBP-1
regulates signal transduction, transcription factors and bone marrow
colonization in B cells. *EMBO J* 28(11):1624–1636.

52. Rutkowski DT, Hegde RS (2010) Regulation of basal cellular physiology by the
homeostatic unfolded protein response. *J Cell Biol* 189(5):783–794.

53. Michell RH (2018) Do inositol supplements enhance phosphatidylinositol
supply and thus support endoplasmic reticulum function? *Br J Nutr* 120(3):301–

316.

54.   McQuin C, et al. (2018) CellProfiler 3.0: Next-generation image processing for
842   biology. *PLoS Biol* 16(7):e2005970.

843   55.   Abraham MJ, et al. (2015) Gromacs: High performance molecular simulations
844   through multi-level parallelism from laptops to supercomputers. *SoftwareX* 1–
845   2:19–25.

846   56.   Humphrey W, Dalke A, Schulten K (1996) VMD: Visual molecular dynamics. *J*
847   *Mol Graph* 14(1):33–38.

848   57.   Wu EL, et al. (2014) CHARMM-GUI membrane builder toward realistic
849   biological membrane simulations. *J Comput Chem* 35(27):1997–2004.

850   58.   Lee J, et al. (2016) CHARMM-GUI Input Generator for NAMD, GROMACS,
851   AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the
852   CHARMM36 Additive Force Field. *J Chem Theory Comput* 12(1):405–413.

853   59.   Klauda JB, et al. (2010) Update of the CHARMM All-Atom Additive Force Field
854   for Lipids: Validation on Six Lipid Types. *J Phys Chem B* 114(23):7830–7843.

855   60.   Best RB, et al. (2012) Optimization of the additive CHARMM all-atom protein
856   force field targeting improved sampling of the backbone φ, ψ and side-chain
857   χ(1) and χ(2) dihedral angles. *J Chem Theory Comput* 8(9):3257–3273.

858   61.   Huang J, et al. (2016) CHARMM36m: An improved force field for folded and
859   intrinsically disordered proteins. *Nat Methods* 14(1):71–73.

860   62.   PLUMED consortium (2019) Promoting transparency and reproducibility in
861   enhanced molecular simulations. *Nat Methods* 16(8):670–673.

862   63.   Tribello GA, Bonomi M, Branduardi D, Camilloni C, Bussi G (2014) PLUMED 2:
863   New feathers for an old bird. *Comput Phys Commun* 185(2):604–613.

864   64.   Essmann U, et al. (1995) A smooth particle mesh Ewald method. *J Chem Phys*
865   103(19):8577–8593.

866   65.   Bussi G, Donadio D, Parrinello M (2007) Canonical sampling through velocity
867   rescaling. *J Chem Phys* 126(1):014101.

868   66.   Berendsen HJC, Postma JPM, Van Gunsteren WF, Dinola A, Haak JR (1984)
869   Molecular dynamics with coupling to an external bath. *J Chem Phys*
870   81(8):3684–3690.

871   67.   Parrinello M, Rahman A (1981) Polymorphic transitions in single crystals: A
872   new molecular dynamics method. *J Appl Phys* 52(12):7182–7190.

873   68.   Hess B, Bekker H, Berendsen HJC, Fraaije JGEM (1998) LINCS: A Linear

874   Constraint Solver for molecular simulations. *J Comput Chem* 18(12):1463–
875   1472.
876

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (SFB807 'Transport and

Communication across Biological Membranes' to R.E. and G.H.; SFB894 'Ca^{2+}-Signals:

Molecular Mechanisms and Integrative Functions' to R.E). R.C. and G.H. were supported by

the Max Planck Society. We thank Carsten Mattes for excellent technical support and

especially Kristina Halbleib for her important contributions during the early phase of the project.

883
884

Fig. 1: Cysteine-less Ire1 expressed from its endogenous locus is functional. (*A*)
Schematic representations of the knock-in construct *IRE1*$_{3xHA-GFP}$ indicating the
position of cysteine residues and topology. All twelve cysteines of Ire1 and C48 ^{yeGFP}
of yeGFP were substituted to serine to generate a cysteine-less variant. Cysteine 70
of yeGFP (C70 ^{yeGFP}) is functionally relevant (29) and remained in the final construct.
The two cysteines in the signal sequence of Ire1 have not been replaced, as the signal
sequence is removed upon ER-translocation. (*B*) Resistance of the indicated yeast
strains to chronic ER-stress. Stationary overnight cultures of the indicated yeast strains
were used to inoculate a fresh culture in full or minimal media to an OD$_{600}$ of 0.2. After
cultivation for 5 to 7 h at 30°C the cells were diluted with pre-warmed full or minimal
media to an OD$_{600}$ of 0.1. Cells were cultivated for 18 h at 30°C in the indicated media
and stressed with DTT. The density of the resulting culture was determined using the
OD$_{620}$ or OD$_{600}$. (*C*) Relative levels of spliced *HAC1* mRNA in acutely stressed cells
normalized to the degree of *HAC1* slicing in stressed cells expressing *IRE1*$_{3xHA-GFP}$
wildtype. Exponentially growing cells of the indicated strains were used to inoculated
fresh YPD media to an OD$_{600}$ of 0.2, cultivated in YPD and acutely stressed with either
4 mM DTT (left panel) or 1.0 µg/ml Tunicamycin (right panel) for 1 h. The relative level
of spliced *HAC1* in these cells was analyzed by RT-qPCR and quantitated using the
comparative $\Delta\Delta$CT method using normalization to *ACT1* levels. The data were
normalized to the splicing of *HAC1* in stressed cells carrying the *IRE1*$_{3xHA-GFP}$ wildtype
construct. All error bars in this figure represent the mean $\pm$ SEM of at least three
independent experiments. Significance was tested by an unpaired, two-tailed
Student's t test. ***p<0.001, **p<0.01, *p<0.05.

A Overview of experimental setup

SCD complete

OD 0.2 → OD 0.7 → 1h + 2 mM DTT / + 1.5 µg/ml TM → ≈ OD 1 → microsome preparation

B Overview of prepared microsomes used for crosslinking

microsomes unstressed cells

microsomes stressed cells

C Cysteine crosslinking in microsomes with CuSO$_4$

F544C   cys-less

Fig. 2: The crosslinking of Ire1 via single cysteines in microsomes requires
CuSO$_4$ and pre-formed clusters. (*A*) Overview of the cultivation of yeast cells for
cysteine crosslinking. Stationary cells were used to inoculate a fresh culture in SCD
complete media to an OD$_{600}$ of 0.2. After cultivation at 30°C to an OD$_{600}$ of 0.7, Ire1-
clustering was induced either by DTT (1h, 2 mM, SCD) or TM (1h, 1.5 µg/ml, SCD).
After harvesting the unstressed and stressed cells, cells were lysed and used for
microsomal membrane preparation by differential centrifugation. (*B*) Overview of
cysteine crosslinking with CuSO$_4$. Microsomes from unstressed cells and microsomes
from stressed cells containing Ire1 clusters were incubated with CuSO$_4$ to induce
cysteine crosslinking of Ire1. (*C*) Crosslinking of single cysteine variants of Ire1 in
microsomes. The indicated strains were cultivated in the presence and absence of ER-
stressors. Cells were cultivated and treated as described in (*A*). 80 OD equivalents of
cells were harvested and disrupted prior to the preparation of crude microsomes by
differential centrifugation. The crosslinking of juxta-posed cysteines was induced by 10
mM CuSO$_4$ on ice and an incubation for 5 min. 8 µl microsomes with a typical protein
concentration of 1 mg/ml were incubated with 2 µl 50 mM CuSO$_4$. The reaction was
stopped by the addition of 2 µl 1 M NEM, 2 µl 0.5 M EDTA and 4 µl membrane sample
buffer. The resulting samples were subjected to SDS-PAGE corresponding to 0.25 OD
equivalents and analyzed by immunoblotting using a monoclonal anti-HA antibody.

930
Figure 3: Systematic crosslinking of cysteines in the TMH region of Ire1 reveals
a specific configuration during ER stress. (*A*) Primary structure of TMH region of
Ire1 including the adjacent ER-luminal amphipathic helix (25). Almost all residues of
the short TMH (shown in bold) were systematically mutated to cysteines for $CuSO_4$
crosslinking and are illustrated as helical wheel representation in (*B*). (*B*) Helical wheel
representation of the TMH residues (Ire1$^{540-552}$) subjected to the cysteine scanning
approach. (*C*) Crosslinking of single cysteine variants of Ire1 in microsomes. Stationary
cells were used to inoculate fresh SCD complete media to an OD_{600} of 0.2. After
cultivation to an OD_{600} of 0.7 at 30°C, cells were treated either with 2 mM DTT or 1.5
µg/ml TM in SCD medium for 1 h. Microsomes from unstressed cells (no) and stressed
cells were prepared by differential centrifugation and subjected to a $CuSO_4$-mediated
crosslinking procedure. 8 µl microsomes were incubated with 2 µl 50 mM $CuSO_4$ (final
concentration 10 mM) for 5 min on ice. After the reaction was stopped with NEM, EDTA
and membrane sample buffer, the samples (0.25 OD equivalents) were subjected to
SDS-PAGE followed by immunoblotting using anti-HA antibodies. (*D*) Quantification of
tested UPR stress conditions. Cells were cultivated and treated as described in (*C*).
For inositol depletion, exponentially growing cells were inoculated to an OD_{600} of 0.5
and cultivated for 3h in inositol depleted media. Cells were harvested and treated as
described in (*C*). Immunoblots of crosslinked microsomes from cells depleted from
inositol are shown in Fig. S3. The percentage of crosslinked dimer from at least three
independent experiments was determined using the densiometric signals of the bands
illustrated in (*C*) (ER stressors: DTT and TM) and Fig S3. (ER stressors: inositol
depletion) corresponding to the monomeric and covalently dimeric protein, which were
determined using ImageJ (n > 4, mean ± SEM).

956
Figure 4: Structural model of the TMH region of Ire1.
Configuration of a model TMH dimer obtained from atomistic molecular dynamics
simulations. Protomers are shown as an orange ribbon, with the residues F544
highlighted in red. POPC lipids and their phosphate moiety are shown in gray and
purple, respectively. Cholesterol molecules and their hydroxyl groups are shown in
light green and red, respectively. Water is shown in a transparent surface
representation.
964

965
Figure 5: The impact of mutations in the TMH and the AH of Ire1 on its
functionality and crosslinking propensity. (*A*) The ER-stress resistance of cells
expressing the F544A variant of Ire1$_{3xHA-GFP}$ containing the native cysteine 552 was
scored using a yeast growth assay. Stationary overnight cultures of the indicated yeast
strains were used to inoculate a fresh culture to an OD$_{600}$ of 0.2. After cultivation for 5
to 7 h at 30°C, the cells were diluted with fresh media to an OD$_{600}$ of 0.1. Cells were
cultivated for 18 h at 30°C in the presence of the indicated concentrations of DTT. The
density of the resulting culture was determined using the OD$_{600}$. (*B*) The impact of the
F544A mutation in the TMH of Ire1 on the degree of crosslinking via cysteine 552 was
determined using the microsome-based crosslinking assay. Stationary cultures were
used to inoculate fresh SCD complete media to an OD$_{600}$ of 0.2 and cultivated to an
OD$_{600}$ of 0.7. Cells were stressed by the addition of 2 mM DTT or 1.5 µg/ml TM for 1
h. Microsomes of 80 OD units of the indicated yeast strains were prepared by
differential centrifugation and used for cysteine crosslinks. 8 µl microsomal
membranes were incubated with 2 µl CuSO$_4$ (final concentration 10 mM) and
incubated for 5 min on ice. The reaction was stopped by adding 8 µl stopping buffer.
Samples corresponding to 0.25 OD equivalents were subjected to SDS PAGE and a
subsequent immunoblotting with anti-HA antibodies. (*C*) The ER-stress resistance of
cells expressing the AH-disrupting F531R variant of Ire1$_{3xHA-GFP}$ containing the native
cysteine 552 was scored using a yeast growth assay. The indicated cells were
cultivated and treated as in (*A*). (*D*) The impact of the AH-disrupting F531R mutation
of Ire1 on the degree of crosslinking via cysteine 552 was determined using the
microsome-based crosslinking assay. Cells were cultivated and further treated as
described in (*B*). The data are represented as the mean ± SEM and are derived from
at least three independent experiments. Significance was tested by an unpaired, two-
tailed Student's t test. **p<0.01.
992

993
Fig. S1: Protein levels of cysteine-less Ire1 and characterization of its membrane
association. (*A*) Protein levels of cells expressing either *IRE1*$_{3xHA-GFP}$ WT or the
cysteine-less (cysteine-less) variant. The isogenic wildtype strain BY4741 that does
not express a HA-tagged variant of IRE1 was used as a specificity control (NC).
Stationary overnight cultures were used to inoculate a fresh culture in SCD complete
to an OD$_{600}$ of 0.2 and cultivated until an OD$_{600}$ of 1 was reached. 0.1 OD equivalents
of cell lysates were immunoblotted using anti-HA and anti-Pgk1 antibodies. (*B*)
Subcellular fractionation of exponentially growing cells expressing cysteine-less
IRE1$_{3xHA-GFP}$ by differential centrifugation at 5,000 x g and 100,000 x g. Stationary
overnight cultures were used to inoculate a fresh culture in SCD complete to an OD$_{600}$
of 0.2 and cultivated until an OD$_{600}$ of 1 was reached. 80 OD$_{600}$ equivalents were
harvested and used for microsomal membrane preparation. The individual supernatant
and pellet fractions were analyzed by immunoblotting using anti-HA, anti-Pgk1 and
anti-Dpm1 antibodies by loading 0.4 OD equivalents. (*C*) Extraction assay of
microsomes. Carbonate and urea extraction validate proper membrane integration of
cysteine-less *IRE1*$_{3xHA-GFP}$ (cysteine-less). Samples of each step corresponding to 0.2
OD equivalents were analyzed by immunoblotting using anti-HA, anti-Pgk1 and anti-
Dpm1 antibodies. (*D*) PDI levels in acutely stressed cells normalized to the fold change
of unstressed cells expressing *IRE1*$_{3xHA-GFP}$ wildtype. Exponentially growing cells of the
indicated strains were used to inoculated fresh YPD media to an OD$_{600}$ of 0.2,
cultivated in YPD and acutely stressed with either 4 mM DTT (left panel) or 1.0 μg/ml
Tunicamycin (right panel) for 1 h. The relative level of *PDI* in these cells was analyzed
by RT-qPCR and quantitated using the comparative ΔΔCT method using normalization
to *ACT1* levels. The data were normalized to the *PDI* level in unstressed cells carrying
the *IRE1*$_{3xHA-GFP}$ wildtype construct. All error bars in this figure represent the mean ±
SEM of at least three independent experiments. Significance was tested by an
unpaired, two-tailed Student's t test. **p<0.01, *p<0.05. (E) Analysis of fluorescent
clusters by microscopy in the indicated cells expressing either *IRE1*$_{3xHA-GFP}$ WT or the
cysteine-less construct. The number of clusters per cell was determined for the
indicated conditions and plotted in the pie chart. The total number of analyzed cells is
given for each condition.

1025

1026

1027
Fig. S2. Co-immunoprecipitation validates the formation of homo-dimers of Ire1
in crosslinking experiments. A crosslinking experiment using $CuSO_4$ was performed
with microsomes prepared from cells expressing a HA-tagged variant of Ire1 from
endogenous locus (*IRE1*3xHA-GFP) and a Flag-tagged variant (*IRE1*3xFlag-GFP) from a
CEN-based plasmid. A yeast culture in selective SCD-LEU was inoculated to an OD_{600}
of 0.2 from a stationary overnight culture and cultivated at 30°C until an OD_{600} of 0.7
was reached. The cells were either stressed with 2 mM DTT or left untreated and were
further cultivated for 1 h. 80 OD_{600} equivalents from these cultures were harvested by
centrifugation. Microsomal membranes were isolated by differential centrifugation.
Microsomes prepared from cells expressing only one of the two tagged variants of Ire1
served as controls. Both constructs contained a single cysteine in the TMH region at
the position 552 (C552). After incubation of the microsomes with 10 mM $CuSO_4$ on ice
for 5 min, the crosslinking reaction was stopped by the addition of NEM in a final
concentration of 111 mM and EDTA in a final concentration of 50 mM. The microsomes
were then solubilized using 2% Triton X-100 and subjected to an IP using anti-Flag
beads. Both the input and IP samples were analyzed by immunoblotting using anti-
Flag and anti-HA antibodies.

1045
Figure S3: Functionality of cysteine mutants and their crosslinking potential in
lipid bilayer stress conditions. (*A*) The resistance to ER-stress was investigated for
the indicated yeast strains. Stationary overnight cultures of the indicated yeast strains
were used to inoculate a fresh culture in full or minimal media to an OD_{600} of 0.2. After
cultivation for 5 to 7 h at 30°C the cells were diluted with fresh minimal media to an
OD_{600} of 0.1. Cells were cultivated for 18 h at 30°C and stressed with DTT. The density
of the resulting culture was determined using the OD_{620} or OD_{600}. The error bars
represent the mean ± SEM of at least two independent clones. (*B*) Protein levels of
cells expressing different *IRE1*$_{3xHA-GFP}$ variants. The lysates of exponentially growing
cells were immunoblotted using anti-HA and anti-Pgk1 antibodies. (*C*) Crosslinking of
single cysteine variants of Ire1 in microsomes derived from cells grown in lipid bilayer
stress conditions. Exponentially growing cells in SCD complete media were washed
and used to inoculate a fresh media w/o inositol to an OD_{600} of 0.5. To induce lipid
bilayer stress, the cells were grown in inositol depleted SCD complete media for 3 h.
80 OD equivalents were harvested and used for microsomal membrane preparation.

CuSO$_4$ induced crosslink was performed by incubating 8 µl of microsomes with 2 µl of
50 mM CuSO$_4$ for 5 min on ice. After stopping the reaction with NEM and EDTA,
samples were subjected to SDS-PAGE with a subsequent immunoblotting with anti-
HA antibody.
1065

1066
1067
Figure S4: The dimeric TMH region of Ire1 deforms the membrane.
Configuration of a model TMH dimer obtained from atomistic molecular dynamics
simulations. Protomers are shown as an orange ribbon, with the residues F544
highlighted in red. The phosphate moieties of POPC are shown as purple beads. Rhe
hydroxyl groups of cholesterol molecules are shown as red beads. Water is shown in
a transparent surface representation.

1074

1075
Fig. S5: The impact of the AH mutation of Ire1 on its functionality and
crosslinking propensity. (*A*) The ER-stress resistance of cells expressing the F544A
variant of Ire1$_{3xHA-GFP}$ containing the native cysteine 552 was scored using the yeast
growth assay as described in Figure 5. (*B*) The impact of the F544A mutation in the
TMH of Ire1 on the degree of crosslinking via cysteine 552 was determined using the
microsome-based crosslinking assay. The data are represented as the mean ± SEM
and are derived from at least three independent experiments. Significance was tested
by an unpaired, two-tailed Student's t test. *p<0.05.
1084

1085   **Supplementary Movie: A structural model of the TMH region of Ire1 highlights**
1086   **membrane thinning and water penetration into the bilayer.**

 Table 1: Yeast strains of used in this study

Strain No,	Description	Genotype	Source
RE001	BY4741	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0	Euroscarf
RE046	*ΔIRE1*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; ura3Δ0; *ire1*Δ::kanMX4	Euroscarf
RE127	*ΔIRE1ΔIRE1promotor*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA pUG72	(25)
RE425	*IRE1*-3xHA-yeGFP	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE451	(25)
RE343	*IRE1*-3xHA-yeGFP *cysteine-less*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE375	This paper
RE342	*IRE1*-3xHA-yeGFP C552	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE374	This paper
RE725	*IRE1*-3xHA-yeGFP cysteine-less + CEN IRE1-3xFLAG-yeGFP *cysteine-less*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE375 *IRE1*-3xFLAG-yeGFP::LEU pRE699	This paper
RE726	*IRE1*-3xHA-yeGFP C552 + *CEN IRE1*-3xFLAG-yeGFP C552	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE374 *IRE1*-3xFLAG-yeGFP::LEU pRE700	This paper
RE530	*IRE1*-3xHA-yeGFP E540C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE575	This paper
RE531	*IRE1*-3xHA-yeGFP T541C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE576	This paper
RE532	*IRE1*-3xHA-yeGFP G542C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE577	This paper
RE533	*IRE1*-3xHA-yeGFP V543C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE578	This paper
RE534	*IRE1*-3xHA-yeGFP F544C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE579	This paper
RE522	*IRE1*-3xHA-yeGFP L545C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE570	This paper
RE535	*IRE1*-3xHA-yeGFP L546C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE581	This paper
RE717	*IRE1*-3xHA-yeGFP L547C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE691	This paper
RE718	*IRE1*-3xHA-yeGFP F548C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE692	This paper
RE719	*IRE1*-3xHA-yeGFP L549C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE693	This paper
RE720	*IRE1*-3xHA-yeGFP I550C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE694	This paper
RE721	*IRE1*-3xHA-yeGFP F551C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE695	This paper
RE722	*IRE1*-3xHA-yeGFP F544A C552 *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE696	This paper
RE723	*IRE1*-3xHA-yeGFP F531R C552 *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE698	This paper
RE724	*IRE1*-3xHA-yeGFP F531R F544C *single cysteine*	BY4741 MATa; *his3*Δ1; *leu2*Δ0; *met15*Δ0; *ura3*Δ0; *ire1*Δ::URA *IRE1*-3xHA-yeGFP::HIS pRE697	This paper

1088

1089

1090 **Table 2: Plasmids used in this study**

Number	Description	Recombinant DNA	Source
pRE451	*IRE1*-3xHA-yeGFP	pcDNA3.1 *IRE1*-3xHA-yeGFP WT	(25)
pEv200	pRS315 *IRE1*-yeGFP-HA	pRS315 *IRE1*-yeGFP-HA	(28)
pRE375	*IRE1*-3xHA-yeGFP *cysteine-less*	pcDNA3.1 *IRE1*-3xHA-yeGFP *cysteine-less*	This paper
pRE374	*IRE1*-3xHA-yeGFP C552 *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP C552 *single cysteine*	This paper
pRE699	*CEN IRE1*-3xFLAG-yeGFP *cysteine-less*	pRS315 *IRE1*-3xFLAG-yeGFP *cysteine-less*	This paper
pRE700	*CEN IRE1*-3xFLAG-yeGFP C552 *single cysteine*	pRS315 *IRE1*-3xFLAG-yeGFP C552 *single cysteine*	This paper
pRE575	*IRE1*-3xHA-yeGFP E540C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP E540C *single cysteine*	This paper
pRE576	*IRE1*-3xHA-yeGFP T541C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP T541C *single cysteine*	This paper
pRE577	*IRE1*-3xHA-yeGFP G542C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP G542C *single cysteine*	This paper
pRE578	*IRE1*-3xHA-yeGFP V543C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP V543C *single cysteine*	This paper
pRE579	*IRE1*-3xHA-yeGFP F544C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP F544C *single cysteine*	This paper
pRE570	*IRE1*-3xHA-yeGFP L545C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP L545C *single cysteine*	This paper
pRE581	*IRE1*-3xHA-yeGFP L546C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP L546C *single cysteine*	This paper
pRE691	*IRE1*-3xHA-yeGFP L547C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP L547C *single cysteine*	This paper
pRE692	*IRE1*-3xHA-yeGFP F548C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP F548C *single cysteine*	This paper
pRE693	*IRE1*-3xHA-yeGFP L549C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP L549C *single cysteine*	This paper
pRE694	*IRE1*-3xHA-yeGFP I550C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP I550C *single cysteine*	This paper
pRE695	*IRE1*-3xHA-yeGFP F551C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP F551C *single cysteine*	This paper
pRE696	*IRE1*-3xHA-yeGFP F544A C552 *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP F544A C552 *single cysteine*	This paper
pRE697	*IRE1*-3xHA-yeGFP F531R F544C *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP F531R F544C *single cysteine*	This paper
pRE698	*IRE1*-3x-HA-yeGFP F531R C552 *single cysteine*	pcDNA3.1 *IRE1*-3xHA-yeGFP F531R C552 *single cysteine*	This paper

1091