TITLE: An African origin for *Mycobacterium bovis*

Chloé Loiseau[1,2*], Fabrizio Menardo[1,2*], Abraham Aseffa[3], Elena Hailu[3], Balako Gumi[4],

Gobena Ameni[5], Stefan Berg[6], Leen Rigouts[7,8,9], Suelee Robbe-Austerman[10], Jakob

Zinsstag[1,2], Sebastien Gagneux[1,2*] and Daniela Brites[1,2*]

[1]Swiss Tropical and Public Health Institute, Basel, Switzerland

[2]University of Basel, Basel, Switzerland

[3]Armauer Hansen Research Centre, Addis Ababa, Ethiopia

[4]Bule Hora University, Department of Animal Science and Range Management, Bule Hora

Town, Ethiopia

[5]Addis Ababa University, Aklilu Lemma Institute of Pathobiology, Addis Ababa, Ethiopia

[6]Animal & Plant Health Agency (APHA), Bacteriology Department, Weybridge, Surrey,

United Kingdom

[7]Mycobacteriology Unit, Department of Biomedical Sciences, Institute of Tropical Medicine,

Antwerp, Belgium

[8]Collection of Mycobacterial Cultures (BCCM/ITM), Institute of Tropical Medicine,

Antwerp, Belgium

[9]Department of Biomedical Sciences, Antwerp University, Antwerp, Belgium

[10]National Veterinary Services Laboratories, United States Department of Agriculture, Ames,

Iowa, USA

Corresponding authors:d.brites@swisstph.ch; sebastien.gagneux@swisstph.ch

*equal contribution

ABSTRACT

Background and objectives

Mycobacterium bovis and *Mycobacterium caprae* are two of the most important agents of

tuberculosis (TB) in livestock and the most important causes of zoonotic TB in humans.

However, little is known about the global population structure, phylogeography and

evolutionary history of these pathogens.

Methodology

We compiled a global collection of 3364 whole-genome sequences from *M. bovis* and *M.*

caprae originating from 35 countries and inferred their phylogenetic relationships, geographic

origins and age.

Results

Our results resolved the phylogenetic relationship among the four previously defined clonal

complexes of *M. bovis*, and another eight newly described here. Our phylogeographic analysis

showed that *M. bovis* likely originated in East Africa. While some groups remained restricted

to East- and West Africa, others have subsequently dispersed to different parts of the world.

Conclusions and implications

Our results allow a better understanding of the global population structure of *M. bovis* and its

evolutionary history. This knowledge can be used to define better molecular markers for

epidemiological investigations of *M. bovis* in settings where whole genome sequencing

cannot easily be implemented.

BACKGROUND AND OBJECTIVES

Tuberculosis (TB) remains an important burden for global health and the economy [1]. TB is the number one cause of human death due to infection globally, with an estimated 10.0 million new cases and 1.5 million deaths occurring every year [1]. TB is caused by members of the *Mycobacterium tuberculosis* complex (MTBC), which includes seven human-adapted lineages, and several animal-adapted ecotypes including *M. bovis* and *M. caprae.* Animal TB complicates the control of human TB due to the zoonotic transfer of TB bacilli from infected animals to exposed human populations e.g. through the consumption of unpasteurized milk or handling of contaminated meat [2]. *M. bovis* and *M. caprae* are the most important agents of TB in livestock and the most important agents of zoonotic TB in humans, causing an estimated 147 000 new human cases and 12 500 human deaths yearly [1, 3]. Zoonotic TB caused by *M. bovis* also poses a challenge for patient treatment, due to its natural resistance to pyrazinamide (PZA), one of the four first-line drugs used in the treatment of TB. In addition, TB in livestock accounts for an estimated loss of three billion US dollars per year [4]. In Africa, the prevalence of *M. bovis* is highest in peri-urban dairy belts of larger cities and remains at low levels in rural areas [5], often also threatening wildlife populations [6]. During the last few years, analyses of large globally representative collections of whole genome sequences (WGS) from the human-adapted MTBC lineages have enhanced our understanding of the global population structure, phylogeography and evolutionary history of these pathogens [7]. By contrast, little corresponding data exist for the various animal-adapted ecotypes of the MTBC such as *M. bovis*.

Current knowledge about global *M. bovis* populations stems mostly from spoligotyping [8, 9]. This method has been highly valuable for showing that *M. bovis* populations vary by geography, and defining strain families based on the presence or absence of spacers in the Direct Repeat region of the MTBC genome [8]. However, the discriminatory capacity of spoligotyping is limited since diversity is measured at a single locus prone to convergent evolution and phylogenetic distances cannot be reliably inferred [10].

In addition to spoligotyping, other genomic markers such as deletions [11-14] and single nucleotide polymorphisms (SNPs) [14], have given insights into the biogeography of *M. bovis*. These markers have been used to define four major groups of genotypes within *M. bovis*, known as clonal complexes European 1 and 2 (Eu1, Eu2) and African 1 and 2 (Af1 and Af2) [11-14]. Bovine TB in West Africa and East Africa is mainly caused by the clonal complexes Af1 and Af2, respectively [11, 12]. Bovine TB in Europe and in the Americas is caused by clonal complex Eu1, which affects mostly the British Islands and former trading

countries of the UK [13], while Eu2 is prevalent mostly in the Iberian Peninsula and Brazil

[14].

More recently, studies based on WGS have brought deeper insights into the population

dynamics of *M. bovis* and showed that unlike *M. tuberculosis*, wild animals can act as *M.*

bovis reservoirs in different regions of the world [15-18]. However, most studies using WGS

have aimed at investigating local epidemics, and little is known about the global population

structure and evolutionary history of *M. bovis*. Recently, we suggested a scenario for the

evolution of the animal-adapted MTBC, in which we propose that *M. caprae* and *M. bovis*

might have originally come out of Africa [19]. Here we gathered 3356 *M. bovis* and *M.*

caprae WGS from the public domain, to which we added eight new *M. bovis* sequences from

strains isolated in East Africa. Our results provide a phylogenetic basis to better understand

the global population structure of *M. bovis*. Moreover, they point to East Africa as the most

likely origin of contemporary *M. bovis*.

METHODS

Data collection

A total of 3929 *M. bovis* genomes were retrieved from EBI: 3834 BioSamples were registered

on EBI with the taxon id 1775 (corresponding to "*Mycobacterium tuberculosis* variant bovis")

and downloaded on the 11[th] of March 2019 and 95 *M. bovis* genomes were registered under

taxon id 1765 (corresponding to "*Mycobacterium tuberculosis*").

Of these, 457 were excluded because they were part of pre-publications releases from the

Wellcome Trust Sanger Institute, 130 were excluded because they were registered as BCG –

Bacille Calmette Guérin, the vaccine strain derived from *M. bovis,* one genome was excluded

because it was wrongly classified as *M. bovis*, and three samples were excluded because they

corresponded to RNA-seq libraries.

In addition, we added 81 publically available *M. caprae* genomes and eight previously

unpublished sequences from *M. bovis* isolated in Ethiopia (n=7) and Burundi (n=1). The

sequencing data has been deposited in the European Nucleotide Archive (EMBL-EBI) under

the study ID PRJEB33773.

From this total of 3427 genomes, 63 sequences were excluded because they did not meet our

criteria for downstream analyses (average whole-genome coverage below 7, ratio of

heterogeneous SNPs to fixed SNPs above 1), yielding a final dataset of 3364 genomes (Fig.

S1, Table S1). Geographical origin of the isolates, date of isolation and host metadata were

recovered from EBI (Table S1).

Whole genome sequence analysis

All samples were subject to the same whole-genome sequencing analysis pipeline, as

described in [20]. In brief, reads were trimmed with Trimmomatic v0.33 [21]. Only reads

larger than 20 bp were kept for the downstream analysis. The software SeqPrep

(https://github.com/jstjohn/SeqPrep) was used to identify and merge any overlapping paired-

end reads. The resulting reads were aligned to the reconstructed ancestral sequence of the

MTBC [22] using the MEM algorithm of BWA v0.7.13 [23] with default parameters.

Duplicated reads were marked using the MarkDuplicates module of Picard v2.9.1

(https://github.com/broadinstitute/picard). The RealignerTargetCreator and IndelRealigner

modules of GATK v 3.4.0 were used to perform local realignment of reads around InDels

[24]. Finally, SNPs were called with Samtools v1.2 mpileup [25] and VarScan v2.4.1 [26]

using the following thresholds: minimum mapping quality of 20, minimum base quality at a

position of 20, minimum read depth at a position of 7X, maximum strand bias for a position

90%. Only SNPs considered to have reached fixation within an isolate (frequency within-

isolate ≥90%) were considered. For SNPs with ≤10% frequency, the ancestor state was called.

SNPs were annotated using snpEff v4.1144 [27], using the *M. tuberculosis* H37Rv reference

annotation (NC_000962.3) as the genome of *M. bovis* (AF2122/97) has no genes absent from

H37Rv except for TbD1 (contains *mmpS6* and the 5′ region of *mmpL6*) [28].

***In silico* spoligotyping, genomic deletions and previously defined clonal complexes**

The spoligotype pattern of the 3364 genomes was determined *in silico* using KvarQ [29]. The

results were submitted to the *Mycobacterium bovis* spoligotype database

https://www.mbovis.org/ [30] and SB numbers obtained.

All 3364 genomes were screened *in silico* for the presence of molecular markers defining the

previously described *M. bovis* clonal complexes; i.e. for the presence or absence of the

genomic deletions RDAf1, RDAf2, RDEu1 (also known as RD17) [11-13], and in the case of

Eu2 [14], for the presence of SNP 3813236 G to A with respect to the H37Rv

(NC_000962.3). Other deletions, RD4, RDpan and N-RD17, previously used to genotype *M.*

bovis lineages were also screened for [31-33]. The genomic coordinates in H37Rv

(NC_000962.3) used to determine each deletion were the following; RDAf1 (664254-

669601); RDAf2 (680337-694429); RDEu1 (1768074-1768878); RD4 (1696017-1708748);

RDpan (4371020-4373425); N-RD17 (3897069-3897783). A genomic region was considered

deleted if the average coverage over the region was below two.

Phylogenetic analyses

All phylogenetic trees were inferred with RAxML (v.8.2.12) using alignments containing

only polymorphic sites. A position was considered polymorphic if at least one genome had a

SNP at that position with a minimum percentage of reads supporting the call of 90%.

Deletions and positions not called according to the minimum threshold of 7, were encoded as

gaps. We excluded positions with more than 10% missing data, positions falling in PE/PPE

genes, phages, insertion sequences and in regions with at least 50 bp identity to other regions

in the genome [34]. Positions falling in drug resistance-related genes were also excluded. The

alignment used to produce Figure 2 comprised 22 492 variable positions and the alignment

used to produce Figure S2 comprised 45 981 variable positions.

Maximum likelihood phylogenies were computed using the general time-reversible model of sequence evolution (-m GTRCAT -V options), 1,000 rapid bootstrap inferences, followed by a thorough maximum-likelihood search performed through CIPRES [35]. All phylogenies were rooted using a *M. africanum* Lineage (L) 6 genome from Ghana (SAMEA3359865).

Obtaining a representative dataset of *M. bovis* genomes - Subsampling 1

Our phylogenetic reconstruction indicated that sequences belonging to clonal complex Eu1 and Eu2 were over-represented in the initial 3364 genome dataset, particularly from the USA, Mexico, New Zealand and the UK. To obtain a smaller dataset with a more even representation of the different phylogenetic groups, we pruned the 3364 genomes using the following criteria: 1) we removed all genomes with non-available country metadata (n=739), which resulted in 2625 genomes; 2) we used Treemmer v0.2 [20] with the option *-RTL 99* to keep 99% of the original tree length and the option *–lm* to include a list of taxa to protect from pruning. This list included all genomes belonging to clonal complexes Af1 and Af2, as well as any genome belonging to any unclassified clade; 3) we visually identified monophyletic clades with all taxa from the same country and used Treemmer v0.2 [20], using options *-lmc* and *–lm*, to only keep a few representatives of each of these clades. To have representatives of the BCG clade, we kept 11 BCG genomes from [36]. This selection process rendered a dataset of 476 genomes.

Ancestral reconstruction of geographic ranges - Subsampling 2

To infer the geographic origin of the ancestors of the main groups of *M. bovis* and *M. caprae*, we used the 476 genomes dataset (see subsampling 1) and excluded all BCG genomes and all *M. bovis* from human TB cases or from unknown hosts, if the strains were isolated in a low incidence TB country (Europe, North America, Oceania). This is justified by the fact that the majority of such cases correspond to immigrants from high incidence countries that were infected in their country of origin, i.e. country of isolation does not correspond to the native geographic range of the strain and is thus not informative for the geographic reconstruction. *M. bovis* from patients in high incidence countries were kept (Table S1). The resulting dataset was composed of 392 genomes.

For the ancestral reconstruction of geographic ranges, we used the geographic origin of the strains and the phylogenetic relationships of the 392 genomes. Geographic origin was treated

as a discrete character to which 13 states, corresponding to UN-defined regions, were
assigned. To select the best model of character evolution, the function fitMk from the package
phytools 0.6.60 in R 3.5.0 [37] was used to obtain the likelihoods of the models ER (equal-
rates), SYM (symmetrical) and ARD (all rates different) [38]. A Likelihood Ratio Test (LRT)
was used to compare the different log-Likelihoods obtained. According to the former, the best
fitting model was SYM, a model that allows states to transition at different rates in a
reversible way, i.e. reverse and forward transitions share the same parameters (Table S2). The
function *make.simmap* in phytools package 0.6.60 in R 3.5.0 [37, 39] was used to apply
stochastic character mapping as implemented in SIMMAP [40] on the 392 genomes
phylogeny inferred from the best-scoring ML tree rooted on L6, using the SYM model with
100 replicates. We summarized the results of the 100 replicates using the function *summary* in
phytools package 0.6.60 in R [37].

Molecular Dating of *M. bovis* and *M. caprae* – Subsampling 3

For the molecular clock analyses, we considered only genomes for which the date of isolation
was known (n=2058). For the eight genomes sequenced in this study, the date of isolation was
retrieved at a later point, and these strains were not included in the dating analysis (Table S1).
We used a pipeline similar to that reported in [20]. We built SNP alignments including
variable positions with less than 10% of missing data (alignment length; 24 828 variable
positions). We added an L6 strain as outgroup (SAMEA3359865) and inferred the Maximum
Likelihood tree as described above. Since the alignment contained only variable positions, we
rescaled the branch lengths of the trees: rescaled_branch_length = ((branch_length *
alignment_length) / (alignment_length + invariant_sites)). To evaluate the strength of the
temporal signal, we performed root-to-tip regression using the R package ape [41].
Additionally, we used the least square method implemented in LSD v0.3-beta [42] to estimate
the molecular clock rate in the observed data and performed a date randomization test with
100 randomized datasets. To do this, we used the quadratic programming dating (QPD)
algorithm and calculated the confidence interval (options -f 100 and -s).
We also estimated the molecular clock rates using a Bayesian analysis. For this, we reduced
the dataset to 300 strains with Treemmer v0.2 in the following way: we randomly subsampled
strains, maintaining the outgroup and at least one representative of four small clades of the
tree that would have disappeared with simple random subsampling strategy (Af2 clonal

complex: G42133; Af1 clonal complex: G02538; *PZA_ sus_unknown1*: G04143, G04145, G04147; *M. caprae*: G42152, G42153, G37371, G37372, G41838; Table S1). The resulting alignment included 13 012 variable sites (subset1).

We used jModelTest 2.1.10 v20160303 [43] to identify the best fitting nucleotide substitution model among 11 possible schemes, including unequal nucleotide frequencies (total models = 22, options -s 11 and -f). We performed Bayesian inference with BEAST2 [44]. We corrected the xml file to specify the number of invariant sites as indicated here: https://groups.google.com/forum/#!topic/beast-users/QfBHMOqImFE, and used the tip sampling years to calibrate the molecular clock.

We used the uncorrelated lognormal relaxed clock model [45], the best fitting nucleotide substitution model according to the results of jModelTest (all criteria selected the transversional model (TVM) as the best model), and three different coalescent priors: constant population size, exponential population growth and the Bayesian Skyline [46]. We chose a 1/x prior for the population size [0- 10^9], a 1/x prior for the mean of the lognormal distribution of the clock rate [10^{-10} – 10^{-5}], and the standard Gamma distribution as prior for the standard deviation of the lognormal distribution of the clock rate [0 – infinity]. For the exponential growth rate prior, we used the standard Laplace distribution [-infinity – infinity]. For all analyses, we ran two runs and used Tracer 1.7.1 [47] to evaluate convergence among runs and to calculate the estimated effective sample size (ESS). We stopped the runs when they reached convergence, and the ESS of the posterior and of all parameters were larger than 200. The number of generations ranged from 150 to 300 million depending on the run. We used Tracer [48] to identify and exclude the burn-in, which ranged from 4 to 30 million generations, depending on the run.

Since the BEAST analysis was based on a sub-sample of the data, we tested the robustness of the sub-sampling by repeating twice the sub-sampling with Treemmer [20]. This resulted in two alignments of 13 272 and 12 820 SNPs, respectively (subset2 and subset3). We then repeated all the BEAST analyses described above on these two additional datasets. For the BEAST analyses, all trees were summarized in a maximum clade credibility tree with the software Treeannotator (part of the BEAST package), after removing the burn-in and sub-sampling one tree every 10 000 generations.

RESULTS AND DISCUSSION

Phylogenetic inference of *M. bovis* and *M. caprae* populations

The phylogenetic reconstruction of all *M. bovis* and *M. caprae* sequences obtained (n=3364) confirmed that these two ecotypes correspond to two monophyletic groups, despite infecting similar hosts [49, 50] (Fig. S3). The range of host species from which *M. bovis* was isolated is broad, confirming that *M. bovis* can cause infection in many different mammalian species (Table S1). Our collection of *M. caprae* included genomes from Japan (isolated in elephants from Borneo) [51], China (isolated in primates, Table S1) and Peru (host information unavailable, but possibly human [52]), suggesting that the host and geographic distribution of this ecotype ranges well beyond Southern- and Central Europe [53, 54]. One group of *M. caprae* genomes with origin in Germany contained a deletion of 36-38 kb, which encompasses the region of difference RD4 (Table S1). This is in agreement with a previous study reporting Alpine *M. caprae* isolates as RD4 deleted, using conventional RD typing [55]. For all *M. bovis* genomes, we determined *in silico* clonal complexes Eu1, Eu2, Af1 and Af2 and spoligotypes, and mapped them on the phylogenetic tree and onto a world map (Fig. 1, Fig. S2-S3, Table S1). All previously described clonal complexes corresponded to monophyletic groups in our genome-based phylogeny (Fig. S3). The phylogenetic tree also revealed *M. bovis* representatives that did not fall into any of the previously described clonal complexes (n=175, 5.3%, Fig. 1, Fig. S2-S3, Table S1). These belonged to eight monophyletic clades with unknown classification and to a few singleton branches (Fig. S3). The tree topology showed a strong bias towards closely related strains, in particular among Eu1, which reflects the different sampling and WGS efforts in the different geographic regions (Fig. 1, Fig. S3). Closely related genomes inform the local epidemiology but not the middle/long term evolutionary history of the strains and were thus excluded from further analysis (Subsampling 1, see Methods).

Two deep divergence events in *M. bovis* populations were notorious: one giving rise to an unclassified lineage we named *M. bovis PZA_sus_unknown1* (RD4 deleted as other *M. bovis*), which included five samples from Uganda (isolated from *Bos taurus* cattle, Table S1), three from Malawi (isolated from humans, Table S1) and one isolated from an antelope in Germany (Table S1). These *M. bovis* isolates lacked the *PncA* H57D mutation that is responsible for the intrinsic pyrazinamide resistance of canonical *M. bovis* as reported previously [56] (Fig. 2). This clade, retained the region of difference N-RD17, unlike all remaining *M. bovis*, and is

probably related to a group of strains previously isolated from cattle in Tanzania and reported

as "ancestral" by [31] (Fig. 2). In agreement with that, our *PZA_sus_unknown1* clade had

other deletions reported as specific to these Tanzanian isolates; a larger deletion

encompassing RDpan (RDbovis(a)_Δpan) and RDbovis(a)_kdp [31]. The second deep

branching lineage included all other *M. bovis* strains descendent from an ancestor that

acquired the *PncA* H57D mutation and therefore encompasses all previously described clonal

complexes [8, 14], as well as the other previously unclassified clades we describe here.

From the *M. bovis* PZA resistant ancestor strains, two main splits occurred; one split led to the

ancestor of Af2 and its previously unclassified sister clade which we called *unknown2* and

which contains the BCG vaccine strains (Fig. 2). *M. bovis* strains with spoligotyping patterns

similar to BCG have previously been referred to as "BCG-like". However, our genome-based

phylogeny shows that BCG-like spoligotyping patterns are present in several clades and have

thus little discriminatory power [10] (Fig. 2, Table S4). In Af2 and *unknown2* the region of

difference RDpan is present [31] (Fig. 2). Otherwise, RDpan is deleted in all other *M. bovis*

except for the *unknown3* clade, in which it is polymorphic (Fig. 2). The other split led to the

ancestor, from which all remaining *M. bovis* strains evolved, i.e. Af1, Eu2 and Eu1 as well as

other groups (Fig. 2). Interestingly, Af1 does not share a MRCA with Af2 but with Eu1 and

Eu2 as well as with another unclassified group, which we called *unknown3* (Fig. 2). Clonal

complexes Eu1 and Eu2 share a MRCA together with five other *unknown* clades (*unknown 4*

– *unknown 8*). Eu2 is more closely related to clades *unknown4* and *5,* than to Eu1 (Fig. 2).

Eu1 in turn shares a common ancestor with three other clades *unknown6*, *7* and *8* (Fig. 2).

The temporal and geographic origin of *M. bovis*

Our reconstruction of ancestral geographical ranges points to East Africa as the most likely

origin for the ancestor of all *M. bovis* (Fig. 2, Fig. S4). This is supported by the fact that the

basal clade *M. bovis - PZA_sus_unknown1* has an exclusively East African distribution and is

pyrazinamide susceptible. Pyrazinamide susceptibility in *M. bovis* is probably an ancestral

character given that all other lineages of the MTBC are pyrazinamide susceptible.

Alternatively, the ancestral *M. bovis* pyrazinamide susceptible populations could have had a

much broader geographic distribution, which later became restricted to East Africa. For *M.*

caprae, the sampling was too small and biased (Table S1) and no conclusions can be

confidently drawn. We performed tip-dating calibration using the isolation dates of the strains

with both Bayesian methods and LSD (see methods). Both the tip-to-root regression and the

randomization tests performed indicated a temporal signal in the data (Fig. S5). We estimated

a clock rate of between 6.66×10^{-8} and 1.26×10^{-7} for the BEAST analyses (Table S3), and

between 6.10×10^{-8} and 8.29×10^{-8} for the LSD analysis. These results are in line with the

results of previous studies [15, 57]. The common ancestor of *M. bovis* was estimated to have

evolved between the years 256 and 1125 AD by the Bayesian analysis (cumulative range of

the 95% Highest Posterior Density (HPD) of all nine BEAST analyses) and in the year 388

AD by LSD (Fig. 3, TreeS1-S10). Together, these estimates suggest that *M. bovis* has

emerged in East Africa sometime during the period spanning the 3rd to the 12th century AD

(Fig. 3). However, the credibility intervals of the different analysis spanned several centuries.

This was due to the intrinsic uncertainty of dating ancestral nodes when the clock calibration

is based on the sampling time of recently sampled tips (all strains considered in this study

have been sampled in the last 40 years). Moreover, by relying exclusively on recent

calibration points to date older nodes, we ignored the issue of the time-dependency of the

estimated clock rates [58]. According to the time-dependency hypothesis, evolutionary rate

estimates depend on the age of the calibration points, with older calibration points resulting in

lower rates. In MTBC, this topic was discussed at length in other publications, and the

available data to date does not allow to confidently accept or reject that hypothesis [57, 59-

62]. Our analysis assumes that rates of evolution do not depend on the age of the calibration

points. Therefore, we potentially underestimate the age of the older nodes of the tree.

Molecular archaeological evidence suggests indeed that our dating analyses possibly

underestimate the age of the MRCA of *M. bovis*. In particular, 2000 years old *M. bovis* DNA

reported as having the RD4 and RD17 deletions, was found in human remains in Siberia [63].

The region of difference RD17 corresponds to RDEu1 [13], and defines the clonal complex

Eu1 of *M. bovis*, which we estimate has evolved between the years 1236 and 1603 AD (Fig.

3).

As discussed elsewhere [57], both tip-dating and the analysis of ancient DNA have potential

pitfalls, and these discrepancies cannot be reconciled without additional data. Nevertheless,

the tip-dating calibration provided accurate results for the emergency of BCG strains and for

the introduction of *M. bovis* to New Zealand [64, 65](TreeS1-S10), indicating that the method

can reliably infer divergence times at least for events occurred in the last 200 years.

Insights into the detailed population structure of *M. bovis* around the world

Understanding the evolutionary history of the *M. bovis* populations requires understanding

their geographic distribution at a continental scale. Our WGS data set has limited

geographical resolution due to the biased sampling of certain regions of the world, and to the

partial unavailability of associated metadata such as the origin of foreign-born TB patients

from Western countries. To get more insights into the geographical ranges of the different *M.*

bovis clades, we used the spoligotype patterns inferred from the WGS data and searched for

references describing the prevalence of those in different regions of the world (Table S4).

Patterns SB0120 and SB0134, known as "BCG-like" and reported to be relatively prevalent

[9], as well as SB0944, are phylogenetically uninformative; SB0120 is present in several

clades, and SB0134 and SB0944 have evolved independently in two different *M. bovis*

populations (Fig. 2, Table S4).

Our results suggest that the sister clade of all contemporary pyrazinamide resistant *M. bovis*,

PZA_sus_unknown1, is restricted to East Africa. The same holds true for Af2, which is in

accordance with previous reports [8, 12]. Our findings further suggest that the geographical

distribution of the Af2 sister clade *unknown2* includes East Africa (Eritrea, Ethiopia), but also

Southern Europe (Spain and France). Informative spoligotypes of the *unknown2* clade show

that it also circulates in North Africa (Fig. 2, Table S4). Of note, the original strain, from

which all BCG vaccine strains were derived, was isolated in France [66]. Our inferences

suggest that a common ancestor of Af2 and *unknown2* evolved in East Africa, and while Af2

remained geographically restricted, its sister clade *unknown2* has subsequently dispersed (Fig.

2).

All remaining *M. bovis* descended from a common ancestor, for which the geographical

origin was impossible to infer reliably with our data. However, the tree topology showed that

from this ancestor several clades have evolved which are important causes of bovine TB

today in different regions of the world (i.e., the clonal complexes Eu1, Eu2 and Af1; Fig.2).

The most basal clade within this group is *unknown3,* which contained 25 genomes mostly

isolated from humans (Table S1). The *in silico* derived spoligotypes suggest that the

geographical spread of *unknown3* ranges from Western Asia to Eastern Europe, but also

includes East Africa (Fig. 2, Table S4). The next split in our phylogeny corresponds to Af1,

which has been characterized extensively using the deletion RDAf1 and spoligotyping, and

shown to be most prevalent in countries from West- and Central Africa [11]. Here, we could

only compile nine Af1 genomes, of which five originated in Ghana [67], and the remaining

had either a European or an unknown origin. The small diversity of Af1 spoligotypes found in

our WGS dataset [11] indicates strong undersampling (Fig. 2, Table S4). Nevertheless, it was

possible to estimate the divergence of the Af1 clade from the remaining *M. bovis* to a period

ranging from the year 921 to 1449 AD (Fig. 3), making it unlikely that Af1 was originally

brought to West Africa by Europeans [68].

The next split comprises clades *unknown4*, *unknown5* and Eu2. Clade *unknown4* was

composed of 33 genomes with little geographic information and for which the most common

spoligotyping pattern was the uninformative SB0120 (n=19). Additional *unknown4*

spoligotypes indicate that strains belonging to this clade circulate in Southern Europe,

Northern and Eastern Africa (Fig. 2, Table S4), supporting dispersion events between Africa

and Southern Europe. Clade *unknown5* comprised only nine genomes isolated mostly from

Zambian cattle. Its corresponding spoligotype is also SB0120, limiting further geographical

inferences.

In contrast to the strains from clades *unknown4* and *unknown5*, among the 323 Eu2 genomes,

no genomes of East African origin were found, and Africa was only represented by nine

South African genomes [69]. By far, most Eu2 were isolated in the Americas. Previous

studies have shown that Eu2 dominates in Southern Europe, particularly in the Iberian

Peninsula [14], thus possibly the source of Eu2 in the Americas. There were no

representatives of Eu2 from the Iberian Peninsula in our dataset. However, our molecular

dating analysis revealed that the common ancestor of Eu2 evolved during the period 1416 to

1705 AD (Fig. 3), which would be compatible with an introduction from Europe into the

Americas.

Clonal complex Eu1, *unknown6*, *unknow7* and *unknown8*, form a sister group to the

previously described. Eu1 has previously been characterized based on the RDEu1 deletion

and spoligotyping, showing that it is highly prevalent in regions of the world that were former

trading partners of the UK [8, 13]. That geographic range is well represented in our dataset,

including many genomes from the UK (n=215) and Ireland (n=45) (Table S1). The latter were

very closely related, suggesting that there was probably fixation of just a few genotypes in

this region as previously proposed [8]. In contrast, most branching events within Eu1

correspond to *M. bovis* isolated in North- and Central America as well as New Zealand,

resulting from the expansion of clonal families not seen in the British Islands. Consequently,

most of the genetic diversity of Eu1 exists outside of its putative region of origin. Our

molecular dating is compatible with this view, indicating that the ancestor of Eu1 is likely to

have emerged between the years 1236 to 1603 AD (Fig. 3), with several Eu1 sub-clades

evolving in the last 200-300 years (TreeS1-S10).

The closest relative to Eu1 is a genome from Ethiopia (*unknown8*) with the spoligotyping

pattern SB1476, commonly found in Ethiopia [12]. *Unknown6* comprised seven genomes

from North America (Fig. 2, Table S1, Table S4), whereas *unknown7* included eight genomes,

four of which were isolated in Western Europe and another four without country of origin

available. Spoligotyping patterns indicate that identical strains are common in Southern

Europe, Northern and Eastern Africa expanding the geographic range of *unknown7*.

CONCLUSIONS AND IMPLICATIONS

We screened the public repositories and compiled 3364 genome sequences of *M. bovis* and *M.*

caprae from 35 countries. Despite the biased geographic distribution of our samples, our

results provide novel insights into the phylogeography of *M. bovis* and *M. caprae.* Our whole-

genome based phylogeny showed that although certain spoligotypes are associated with

specific monophyletic groups, prevalent patterns such as the so-called "BCG-like" should not

be used to infer phylogenetic relatedness. Moreover, our data extend the previously known

phylogenetic diversity of *M. bovis* by eight previously uncharacterized clades in addition to

the four clonal complexes described previously. Among those, *Pza_sus_unknown 1* shares a

common ancestor with the rest of *M. bovis,* has an exclusively East African distribution and

does not share the PncA mutation H57D, conferring intrinsic resistance to PZA.

Our further inferences suggest that *M. bovis* evolved in East Africa. The evolutionary success

of *M. bovis* is linked to the fact that it can infect and transmit very efficiently in cattle. Cattle

have been domesticated twice independently; once in the Near East (*Bos taurus*) and once in

the Indus Valley (*Bos indicus*) approximately 10 000 year ago, and both were introduced to

Africa at different time points and various locations, subsequently interbreeding with local

wild species [70]. Whereas *B. taurus* was introduced probably during the 6 millennium BC

possibly through Egypt, *B. indicus* was most likely introduced twice, first during the second

millennium BC and later during the Islamic conquests [71]. *M. bovis* could have emerged

after the introduction of cattle, benefiting from the development of African pastoralism and

expanding within the continent. The timing of these events is difficult to estimate; the initial

introductions of cattle predate by several thousands of years our inferred temporal origin of

M. bovis. But as discussed, our estimates are possibly affected by the current uncertainty in dating deeper evolutionary events within the MTBC. Alternatively, *M. bovis* could have emerged in the Near East and been introduced to Africa together with cattle. We cannot test this hypothesis formally, as the Near East is poorly represented in our dataset. However, this scenario is difficult to reconcile with the restricted East African distribution of the *Pza_sus_unknown_1* clade, as the Near East was also the origin of taurine cattle both in Europe and Asia, where no clade retaining the ancestral characteristic of pyrazinamide susceptibility was found.

While some *M. bovis* groups remained restricted to East Africa, others have dispersed to different parts of the world. The contemporary geographic distribution of *M. bovis* clades suggest that East- and North Africa, Southern Europe and Western Asia have played an important role in shaping the population structure of these pathogens. However, these regions were not well represented in our dataset. Thus, more *M. bovis* genomes from these regions are necessary to generate better insights, particularly given the central role of these regions in the history of cattle domestication [72]. From a more applied perspective, our work provides a global phylogenetic framework that can be further exploited to find better molecular markers for studying *M. bovis* in settings where genome sequencing cannot be easily implemented.

ACKNOWLEDGEMENTS

Calculations were performed at sciCORE (http://scicore.unibas.ch/) scientific computing core facility at University of Basel. This work was supported by the Swiss National Science Foundation (grants 310030_166687, 310030_188888, IZRJZ3_164171, IZLSZ3_170834 and CRSII5_177163), the European Research Council (309540-EVODRTB) and SystemsX.ch.

REFERENCES

1.   WHO; Global tuberculosis report 2018. Geneva: World Health Organization, 2018.

2.   Olea-Popelka F, Muwonge A, Perera A, et al.; Zoonotic tuberculosis in human beings caused by Mycobacterium bovis-a call for action. *Lancet Infect Dis* 2017;**17**(1):e21-e25. doi: 10.1016/S1473-3099(16)30139-6.

3.   Muller B, Durr S, Alonso S, et al.; Zoonotic Mycobacterium bovis-induced tuberculosis in humans. *Emerg Infect Dis* 2013;**19**(6):899-908. doi: 10.3201/eid1906.120543.

4.   Waters WR, Palmer MV, Buddle BM, et al.; Bovine tuberculosis vaccine research:
historical perspectives and recent advances. *Vaccine* 2012;**30**(16):2611-22.

5.   Tschopp R, Hattendorf J, Roth F, et al.; Cost estimate of bovine tuberculosis to
Ethiopia. *Current topics in microbiology and immunology* 2013;**365**:249-68.

6.   Michel AL, Muller B, van Helden PD; Mycobacterium bovis at the animal-human
interface: a problem, or not? *Veterinary microbiology* 2010;**140**(3-4):371-81.

7.   Gagneux S; Ecology and evolution of Mycobacterium tuberculosis. *Nat Rev Microbiol*
2018;**16**(4):202-213. doi: 10.1038/nrmicro.2018.8.

8.   Smith NH; The global distribution and phylogeography of Mycobacterium bovis
clonal complexes. *Infect Genet Evol* 2012;**12**(4):857-65. doi: 10.1016/j.meegid.2011.09.007.

9.   Ghavidel M, Mansury D, Nourian K, et al.; The most common spoligotype of
Mycobacterium bovis isolated in the world and the recommended loci for VNTR typing; A
systematic review. *Microbial Pathogenesis* 2018;**118**:310-315.

10.   Comas I, Homolka S, Niemann S, et al.; Genotyping of genetically monomorphic
bacteria: DNA sequencing in mycobacterium tuberculosis highlights the limitations of current
methodologies. *PLoS ONE* 2009;**4**(11):e7815. doi: 10.1371/journal.pone.0007815.

11.   Muller B, Hilty M, Berg S, et al.; African 1; An Epidemiologically Important Clonal
Complex of Mycobacterium bovis Dominant in Mali, Nigeria, Cameroon and Chad. *J*
Bacteriol 2009;**191**(6): 1951-1960. doi: 10.1128/JB.01590-08.

12.   Berg S, Garcia-Pelayo MC, Muller B, et al.; African 2, a clonal complex of
Mycobacterium bovis epidemiologically important in East Africa. *J Bacteriol*
2011;**193**(3):670-8. doi: 10.1128/JB.00750-10.

13.   Smith NH, Berg S, Dale J, et al.; European 1: a globally important clonal complex of
Mycobacterium bovis. *Infect Genet Evol* 2011;**11**(6):1340-51. doi:
10.1016/j.meegid.2011.04.027.

14.   Rodriguez-Campos S, Schurch AC, Dale J, et al.; European 2--a clonal complex of
Mycobacterium bovis dominant in the Iberian Peninsula. *Infect Genet Evol* 2012;**12**(4):866-
72. doi: 10.1016/j.meegid.2011.09.004.

15.   Crispell J, Zadoks RN, Harris SR, et al.; Using whole genome sequencing to
investigate transmission in a multi-host system: bovine tuberculosis in New Zealand. *BMC*
genomics 2017;**18**(1):180.

16.   Orloski K, Robbe-Austerman S, Stuber T, et al.; Whole Genome Sequencing of
Mycobacterium bovis Isolated From Livestock in the United States, 1989-2018. *Front Vet Sci*
2018;**5**:253. doi: 10.3389/fvets.2018.00253.

17.   Salvador LCM, O'Brien DJ, Cosgrove MK, et al.; Disease management at the wildlife-
livestock interface: Using whole-genome sequencing to study the role of elk in
Mycobacterium bovis transmission in Michigan, USA. *Mol Ecol* 2019;**28**(9):2192-2205. doi:
10.1111/mec.15061.

18.   Price-Carter M, Brauning R, de Lisle GW, et al.; Whole Genome Sequencing for
Determining the Source of Mycobacterium bovis Infections in Livestock Herds and Wildlife
in New Zealand. *Front Vet Sci* 2018;**5**:272. doi: 10.3389/fvets.2018.00272.

19.   Brites D, Loiseau C, Menardo F, et al.; A New Phylogenetic Framework for the
Animal-Adapted Mycobacterium tuberculosis Complex. *Frontiers in microbiology*
2018;**9**:2820.

20.   Menardo F, Loiseau C, Brites D, et al.; Treemmer: a tool to reduce large phylogenetic
datasets with minimal loss of diversity. *BMC Bioinformatics* 2018;**19**(1):164. doi:
10.1186/s12859-018-2164-8.

21.   Bolger AM, Lohse M, Usadel B; Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 2014;**30**(15):2114-20. doi: 10.1093/bioinformatics/btu170.

22.   Comas I, Chakravartti J, Small PM, et al.; Human T cell epitopes of Mycobacterium
tuberculosis are evolutionarily hyperconserved. *Nat Genet* 2010;**42**(6):498-503. doi:
10.1038/ng.590.

23.   Li H, Handsaker B, Wysoker A, et al.; The Sequence Alignment/Map format and
SAMtools. *Bioinformatics* 2009;**25**(16):2078-9. doi: btp352 [pii]
10.1093/bioinformatics/btp352.

24.   McKenna A, Hanna M, Banks E, et al.; The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Res*
2010;**20**(9):1297-303. doi: 10.1101/gr.107524.110.

25.   Li H; A statistical framework for SNP calling, mutation discovery, association
mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*
2011;**27**(21):2987-93. doi: 10.1093/bioinformatics/btr509.

26.   Koboldt DC, Zhang Q, Larson DE, et al.; VarScan 2: somatic mutation and copy
number alteration discovery in cancer by exome sequencing. *Genome Res* 2012;**22**(3):568-76.
doi: 10.1101/gr.129684.111.

27.   Cingolani P, Platts A, Wang le L, et al.; A program for annotating and predicting the
effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)* 2012;**6**(2):80-92. doi:
10.4161/fly.19695.

28.   Garnier T, Eiglmeier K, Camus JC, et al.; The complete genome sequence of
Mycobacterium bovis. *Proc Natl Acad Sci U S A* 2003;**100**(13):7877-82.

29.   Steiner A, Stucki D, Coscolla M, et al.; KvarQ: targeted and direct variant calling from
fastq reads of bacterial genomes. *BMC Genomics* 2014;**15**:881. doi: 10.1186/1471-2164-15-
881.

30.   Smith NH, Upton P; Naming spoligotype patterns for the RD9-deleted lineage of the
Mycobacterium tuberculosis complex; www.Mbovis.org. *Infection, genetics and evolution :*
journal of molecular epidemiology and evolutionary genetics in infectious diseases
2012;**12**(4):873-6.

31.   Mostowy S, Inwald J, Gordon S, et al.; Revisiting the evolution of Mycobacterium
bovis. *J Bacteriol* 2005;**187**(18):6386-95.

32.   Brosch R, Gordon SV, Garnier T, et al.; Genome plasticity of BCG and impact on
vaccine efficacy. *Proc Natl Acad Sci U S A* 2007.

33.   Salamon H, Kato-Maeda M, Small PM, et al.; Detection of deleted genomic DNA
using a semiautomated computational analysis of GeneChip data. *Genome Res*
2000;**10**(12):2044-54.

34.   Stucki D, Brites D, Jeljeli L, et al.; Mycobacterium tuberculosis lineage 4 comprises
globally distributed and geographically restricted sublineages. *Nat Genet* 2016;**48**(12):1535-
1543. doi: 10.1038/ng.3704.

35.   Miller MA, Pfeiffer W, Schwartz T; Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. *Proceedings of the Gateway Computing Environments*
Workshop (GCE). New Orleans, LA, 2010, 1-8.

36.   Copin R, Coscolla M, Efstathiadis E, et al.; Impact of in vitro evolution on antigenic
diversity of Mycobacterium bovis bacillus Calmette-Guerin (BCG). *Vaccine*
2014;**32**(45):5998-6004. doi: 10.1016/j.vaccine.2014.07.113.

37.   Revell LJ; phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* 2011;**3**(2):217-223.

38.   Lewis PO; A likelihood approach to estimating phylogeny from discrete
morphological character data. *Syst Biol* 2001;**50**(6):913-25. doi:
10.1080/106351501753462876.

39.   Team RC; R: A language and environment for statistical computing. R Foundation for
Statistical Computing. Vienna, Austria, 2019.

40.   Bollback JP; SIMMAP: stochastic character mapping of discrete traits on phylogenies.
BMC Bioinformatics 2006;**7**:88. doi: 10.1186/1471-2105-7-88.

41.   Paradis E, Schliep K; ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* 2019;**35**(3):526-528. doi:
10.1093/bioinformatics/bty633.

42.   To TH, Jung M, Lycett S, et al.; Fast Dating Using Least-Squares Criteria and
Algorithms. *Systematic Biology* 2016;**65**(1):82-97. doi: 10.1093/sysbio/syv068.

43.   Darriba D, Taboada GL, Doallo R, et al.; jModelTest 2: more models, new heuristics
and parallel computing. *Nature Methods* 2012;**9**(8):772-772. doi: DOI 10.1038/nmeth.2109.

44.   Bouckaert R, Heled J, Kuhnert D, et al.; BEAST 2: a software platform for Bayesian
evolutionary analysis. *PLoS Comput Biol* 2014;**10**(4):e1003537. doi:
10.1371/journal.pcbi.1003537.

45.   Drummond AJ, Ho SYW, Phillips MJ, et al.; Relaxed phylogenetics and dating with
confidence. *Plos Biology* 2006;**4**(5):699-710. doi: ARTN e88
10.1371/journal.pbio.0040088.

46.   Drummond AJ, Rambaut A, Shapiro B, et al.; Bayesian coalescent inference of past
population dynamics from molecular sequences. *Mol Biol Evol* 2005;**22**(5):1185-92. doi:
msi103 [pii]
10.1093/molbev/msi103.

47.   Rambaut A, Drummond AJ, Xie D, et al.; Posterior Summarization in Bayesian
Phylogenetics Using Tracer 1.7. *Systematic Biology* 2018;**67**(5):901-904. doi:
10.1093/sysbio/syy032.

48.   Rambaut A, Drummond AJ, Xie D, et al.; Posterior Summarization in Bayesian
Phylogenetics Using Tracer 1.7. *Syst Biol* 2018;**67**(5):901-904. doi: 10.1093/sysbio/syy032.

49.   Rodriguez S, Bezos J, Romero B, et al.; Mycobacterium caprae Infection in Livestock
and Wildlife, Spain. *Emerging Infectious Diseases* 2011;**17**(3):532-535. doi:
10.3201/eid1703.100618.

50.   Prodinger WM, Brandstatter A, Naumann L, et al.; Characterization of
Mycobacterium caprae isolates from Europe by mycobacterial interspersed repetitive unit
genotyping. *J Clin Microbiol* 2005;**43**(10):4984-92.

51.   Yoshida S, Suga S, Ishikawa S, et al.; Mycobacterium caprae Infection in Captive
Borneo Elephant, Japan. *Emerg Infect Dis* 2018;**24**(10):1937-1940. doi:
10.3201/eid2410.180018.

52.   Consortium C, the GP, Allix-Beguec C, et al.; Prediction of Susceptibility to First-
Line Tuberculosis Drugs by DNA Sequencing. *N Engl J Med* 2018;**379**(15):1403-1415. doi:
10.1056/NEJMoa1800474.

53.   Aranaz A, Cousins D, Mateos A, et al.; Elevation of Mycobacterium tuberculosis
subsp. caprae Aranaz et al. 1999 to species rank as Mycobacterium caprae comb. nov., sp.
nov. *Int J Syst Evol Microbiol* 2003;**53**(Pt 6):1785-9.

54.   Broeckl S, Krebs S, Varadharajan A, et al.; Investigation of intra-herd spread of
Mycobacterium caprae in cattle by generation and use of a whole-genome sequence. *Vet Res*
Commun 2017;**41**(2):113-128. doi: 10.1007/s11259-017-9679-8.

55.   Domogalla J, Prodinger WM, Blum H, et al.; Region of difference 4 in alpine
Mycobacterium caprae isolates indicates three variants. *J Clin Microbiol* 2013;**51**(5):1381-8.
doi: 10.1128/JCM.02966-12.

56.   Loiseau C, Brites D, Moser I, et al.; Revised Interpretation of the Hain Lifescience
GenoType MTBC To Differentiate Mycobacterium canettii and Members of the
Mycobacterium tuberculosis Complex. *Antimicrobial Agents and Chemotherapy* 2019;**63**(6).
doi: ARTN e00159-19
10.1128/AAC.00159-19.

57.   Menardo F, Duchene S, Brites D, et al.; The molecular clock of Mycobacterium
tuberculosis. *PLoS Pathog* 2019;**15**(9):e1008067. doi: 10.1371/journal.ppat.1008067.

58.   Ho SY, Phillips MJ, Cooper A, et al.; Time dependency of molecular rate estimates
and systematic overestimation of recent divergence times. *Mol Biol Evol* 2005;**22**(7):1561-8.
doi: 10.1093/molbev/msi145.

59.   Comas I, Coscolla M, Luo T, et al.; Out-of-Africa migration and Neolithic
coexpansion of Mycobacterium tuberculosis with modern humans. *Nat Genet*
2013;**45**(10):1176-82. doi: 10.1038/ng.2744.

60.   Pepperell CS, Casto AM, Kitchen A, et al.; The role of selection in shaping diversity
of natural M. tuberculosis populations. *PLoS Pathog* 2013;**9**(8):e1003543. doi:
10.1371/journal.ppat.1003543.

61.   Bos KI, Harkins KM, Herbig A, et al.; Pre-Columbian mycobacterial genomes reveal
seals as a source of New World human tuberculosis. *Nature* 2014;**514**(7523):494-7. doi:
10.1038/nature13591.

62.   Eldholm V, Pettersson JH, Brynildsrud OB, et al.; Armed conflict and population
displacement as drivers of the evolution and dispersal of Mycobacterium tuberculosis. *Proc*
Natl Acad Sci U S A 2016;**113**(48):13881-13886. doi: 10.1073/pnas.1611283113.

63.   Taylor GM, Murphy E, Hopkins R, et al.; First report of Mycobacterium bovis DNA
in human remains from the Iron Age. *Microbiology* 2007;**153**(Pt 4):1243-9.

64.   Behr MA, Small PM; A historical and molecular phylogeny of BCG strains. *Vaccine*
1999;**17**(7-8):915-922.

65.   Binney BM, Biggs PJ, Carter PE, et al.; Quantification of historical livestock
importation into New Zealand 1860-1979. *N Z Vet J* 2014;**62**(6):309-14. doi:
10.1080/00480169.2014.914861.

66.   Oettinger T, Jorgensen M, Ladefoged A, et al.; Development of the Mycobacterium
bovis BCG vaccine: review of the historical and biochemical evidence for a genealogical tree.
Tuber Lung Dis 1999;**79**(4):243-50. doi: 10.1054/tuld.1999.0206.

67.   Otchere ID, van Tonder AJ, Asante-Poku A, et al.; Molecular epidemiology and whole
genome sequencing analysis of clinical Mycobacterium bovis from Ghana. *PLoS One*
2019;**14**(3):e0209395. doi: 10.1371/journal.pone.0209395.

68.   Muwonge A, Franklyn E, Mark B, et al.; Molecular Epidemiology of Mycobacterium
bovis in Africa. In: B. DA, J. KNP, O. TCs (eds). *Tuberculosis in Animals: An African*
Perspective. Switzerland: Springer, 2019, 127-170.

69.   Dippenaar A, Parsons SDC, Miller MA, et al.; Progenitor strain introduction of
Mycobacterium bovis at the wildlife-livestock interface can lead to clonal expansion of the
disease in a single ecosystem. *Infect Genet Evol* 2017;**51**:235-238. doi:
10.1016/j.meegid.2017.04.012.

70.   Loftus RT, MacHugh DE, Bradley DG, et al.; Evidence for two independent
domestications of cattle. *Proc Natl Acad Sci U S A* 1994;**91**(7):2757-61.

71.   Verdugo MP, Mullin VE, Scheu A, et al.; Ancient cattle genomics, origins, and rapid
turnover in the Fertile Crescent. *Science* 2019;**365**(6449):173-176. doi:
10.1126/science.aav1002.

72.   Decker JE, McKay SD, Rolf MM, et al.; Worldwide Patterns of Ancestry, Divergence,
and Admixture in Domesticated Cattle. *Plos Genetics* 2014;**10**(3). doi: ARTN e1004254
10.1371/journal.pgen.1004254.

73.   Rambaut A; FigTree. Edinburgh: Institute of Evolutionary Biology, University of
Edinburgh, 2010.
701
702

FIGURE LEGENDS

Figure 1 – Geographic distribution of the *M. bovis* samples used in this study according to

isolation country. The circles correspond to pie charts and are coloured according to clonal

complexes.

Figure 2 – Maximum likelihood phylogeny of 476 of the 3364 genomes included in this

study (redundant genomes were removed), and inferred from 22 492 variable positions. The

scale bar indicates the number of substitutions per polymorphic site. The phylogeny is rooted

on a *M. tuberculosis* Lineage 6 genome from Ghana (not shown) and bootstrap values are

shown for the most important splits. The coloured bars on the side of the phylogeny show the

different clonal complexes. Other "unknown" monophyletic clades are coloured in black and

additionally the branches of the eight clades are coloured to show their phylogenetic position

more precisely. The pie charts mapped on the tree represent the summary posterior

probabilities (from 100 runs) of the reconstructed ancestral geographic states and are coloured

according to geographical UN region. Inferred spoligotype patterns from WGS described in

M. bovis spoligotype database [30] are indicated for the unknown clades. The red circles at

the tips correspond to the eight newly sequenced genomes. Regions of difference (RD) as in

[31] are indicated; superscript + and - refers to presence of the region or its deletion,

respectively.

Figure 3 – The inferred age of main monophyletic clades according to LSD and BEAST

dating analyses. For BEAST we report the results of the two analyses that resulted in the

lowest clock rate (subset1 Bayesian Skyline) and in the highest clock rate (subset3

exponential population growth). The confidence intervals reported correspond to the merged

HPD interval of the two BEAST analyses mentioned above. The BEAST analysis was based

on 300 genomes and the LSD analysis was based on 2058 genomes (see methods section for

subsampling strategy). Only one genome from the Af1 clonal complex was included in the

dating analyses and therefore the dates reported correspond to the node where Af1 diverged.

Supplementary Figures:

Figure S1 – Flow chart showing the selection of genomes.

Figure S2 - Geographic distribution of the *M. bovis* samples with unknown classification

used in this study according to isolation country.

Figure S3 – Maximum likelihood phylogeny of all 3364 genomes, based on 45 981 variable

positions. The scale bar indicates the number of substitutions per polymorphic site. The

phylogeny is rooted on a *M. tuberculosis* Lineage 6 genome from Ghana. The outer ring

indicates the geographical region from which the strains were isolated. The four clonal

complexes are highlighted on the tree. Branches corresponding to BCG genomes are coloured

in grey and the *PncA* mutation H57D is indicated by a yellow star.

Figure S4 – Phylogeographic reconstruction of *M. bovis* and *M. caprae*, inferred from 392 genomes. Thirteen UN-defined geographic regions were assigned to the discrete character geographic origin, and mapped onto the phylogeny. Pie charts at internal nodes represent the summary posterior probabilities (from 100 runs) of the reconstructed ancestral geographic states and are coloured according to geographical UN region**.**

Figure S5 – A) Tip-to-root regression and B) Date randomization tests (DTR). The confidence interval of the clock rate estimate for the observed data does not overlap with the confidence intervals of the clock rate estimates obtained from the randomized sets.

Supplemental Tables:

Table S1 - List of genomes included in this study along with metadata used for the analyses.

Table S2 - Comparison of models for discrete character evolution using likelihood ratio tests.

Table S3 – Results of all BEAST analyses.

Table S4 - Spoligotype patterns determined *in silico* for different clonal complex groups with reference to other studies.

Supplementary files:
TreeS1-S10 Ten time-calibrated trees resulted from the molecular clock analyses. The file names indicate the software used (LSD or BEAST), the subsample, and the coalescent population prior (BSP: Bayesian Skyline; exponential: exponential population growth; constant: constant population size). Tip labels are present in Table S1. Ages in years before present can be visualized as well as the 95% High Posterior Density (HPD) (BEAST trees) using FigTree [73].