Vinculin targeting by *Shigella* IpaA promotes stable cell adhesion independent of

mechanotransduction

Cesar Valencia-Gallardo[1-4]*, Daniel-Isui Aguilar-Salvador[1-4]*, Hamed Khakzad[5, 7]§, Charles Bou-Nader[8, 9]§, Christian Malosse[10], Diogo Borges Lima[10], Chakir Bello[1-4], Benjamin Cocom-Chan[1-4], Nicole Quenech'Du[1-4], Bilal Mazhar[1-4], Delphine Javelaud[11-13], Jacques Fattaccioli[14, 15], Alain Mauviel[11-13], Marc Fontecave[8, 9], Atef Asnacios[16], Julia Chamot-Rooke[10], Lars Malmström[5, 7, 17], Guy Tran Van Nhieu[1-4]¶*

[1]Equipe Communication Intercellulaire et Infections Microbiennes, Centre de Recherche Interdisciplinaire en Biologie (CIRB), Collège de France, 75005 Paris, France

[2]Institut National de la Santé et de la Recherche Médicale U1050, 75005 Paris, France

[3]Centre National de la Recherche Scientifique UMR7241, 75005 Paris, France

[4]MEMOLIFE Laboratory of excellence and Paris Science Lettre

[5]Institute for Computational Science, University of Zürich, Zürich, Switzerland.

[6]S3IT, University of Zürich, Zürich, Switzerland.

[7]Swiss Institute of Bioinformatics, Lausanne, Switzerland.

[8]Laboratoire de Chimie des Processus Biologiques, Collège De France, 75005 Paris, France.

[9]Centre National de la Recherche Scientifique UMR8229, 75005 Paris, France

[10] Mass Spectrometry for Biology Utechs, Institut Pasteur, USR 2000, CNRS, 75015 Paris, France

[11]Institut Curie, PSL Research University, INSERM U1021, CNRS UMR3347, Team "TGF-ß and Oncogenesis", Equipe Labellisée LIGUE 2016, F-91400, Orsay, France.

[12]Université Paris-Sud, F-91400, Orsay, France.

[13]Centre National de la Recherche Scientifique UMR 3347, 91400 Orsay, France

[14]PASTEUR, Département de Chimie, École Normale Supérieure, PSL University, Sorbonne Université,

CNRS, 75005 Paris, France

[15]Institut Pierre-Gilles de Gennes pour la Microfluidique, 75005 Paris, France

[16]Laboratoire Matière et Systèmes Complexes, UMR 7057 CNRS & Université Paris Diderot, Sorbonne

Paris Cité, Paris, France

[17]Division of Infection Medicine, Department of Clinical Sciences, Lund University, Lund, Sweden.

*, §equal contribution

¶For correspondence:

E-mail: guy.tran-van-nhieu@college-de-france.fr. Tel: 33-1-44-27-14-89. FAX: 33-1-44-27-14-19

Shigella, the causative agent of bacillary dysentery, invades epithelial cells by injecting type

III effectors that locally reorganize the actin cytoskeleton[1]. The type III effector IpaA targets the

focal adhesion protein vinculin to induce bacterial adhesion associated with the recruitment of

mature focal adhesion markers, despite the inability of bacteria to exert significant counter-forces[2,

3]. Here, we show that three vinculin-binding sites (VBSs) exposed at the IpaA C-terminus act in a

cooperative manner to trigger a yet unreported mode of vinculin activation through specific

interactions with sites in the vinculin head sub-domain D2. Structural modeling based on the mass

spectrometry identification of interacting residues by cross-linking indicated that upon IpaA

binding to vinculin D2, vinculin head sub-domains undergo major conformational changes leading

to higher order heterocomplexes and vinculin homo-trimers. IpaA-mediated vinculin activation

induces the formation of large and stable focal adhesions resisting the action of actin relaxing

drugs. This property enables IpaA to rapidly elicit strong cell adhesion to fibronectin-coated

substrates. While *Shigella* IpaA promotes strong adhesion in the absence of mechanotransduction,

its mode of vinculin activation likely reflects a key step during maturation of cell adhesions driven

by acto-myosin contractility.

Integrin-mediated cell adhesion critically depends on the cytoskeletal linkers talin and vinculin[4-6]. During mechanotransduction, talin acts as a mechanosensor by exposing its VBSs as a function of the stretching force generated by acto-myosin and dependent on substrate stiffness[7]. Talin exposed VBSs recruit and activate vinculin, reinforcing anchorage to the actin cytoskeleton in response to mechanical load[8, 9]. Vinculin contains three repetitions (D1-D3) of a conserved domain consisting of two bundles of four helices, and a fourth D4 domain containing only one helical bundle connected to a proline-rich unstructured region and the carboxyterminal F-actin binding domain[10]. Under its inactive folded form, intramolecular interactions between the vinculin head and tail domains prevent ligand binding.

The *Shigella* type III invasion effector IpaA contains 3 VBSs located in its carboxyterminal moiety[11-13]. IpaA VBS1, as for all VBSs described to date to promote vinculin activation, interacts with the first helical bundle of the D1 domain, promoting major conformational changes that disrupt the D1-tail intramolecular interactions and free the vinculin F-actin binding region[11] (Fig. 1a). IpaA VBS2, in contrast, interacts with the second helical bundle of D1[12] (Fig. 1a) and its association with IpaA VBS1 results in a very high affinity and stable IpaA VBS1-2:D1 complex ,with an estimated K_D in the femtoM range[12]. While functional evidence indicates that IpaA VBS3 cooperates with IpaA VBS1-2 to stimulate bacterial invasion, the isolated peptide acts as IpaA VBS1 in promoting vinculin activation through interaction with the vinculin D1 first helical bundle and also forms a folded bundle with the talin H1-H4 helices[13, 14].

Here, we studied the effects of the IpaA subdomains containing VBS1-2 (A524) or VBS1-3 (A483) on vinculin activation (Fig. 1b). We performed SEC-MALS (Size Exclusion Chromatography-Multi-Angle Light Scattering) to analyze complexes formed upon incubation of A483 with HV_{1-834} containing the D1-D4 domains, corresponding to full-length human vinculin (HV) devoid of the

carboxyterminal F-actin binding domain (Fig. 1b). In addition to the 1:1 D1D4: A483 complex, larger

complexes were observed corresponding to a 2:1 heterocomplex and a 3:0 D1-D4 homo-trimer (Fig.

1c). Similar 2:1 and 3:0 complexes were observed when A483 was incubated with HV_{1-484} containing

only the vinculin D1 and D2 domains (D1D2) (Fig. 1d), indicating that vinculin homo-trimerization

occurs exclusively through these vinculin head sub-domains. By contrast, when A524 was incubated

with D1D2, 1:1 and 2:1 D1D2:A524 complexes were detected, but not the D1D2 homo-trimer

suggesting a role for IpaA VBS3 in vinculin trimerization (Fig. 1d). Consistently, higher order D1D2

homo-complexes devoid of A483 and D1D2:A524 hetero-complexes were visualized by native PAGE

(Fig. 1e and Suppl. Figs. 3c, d). These results suggest that binding of IpaA VBSs to vinculin triggers

conformational changes in vinculin leading to the formation of an IpaA VBS3-dependent vinculin

homo-trimer.

To further investigate initial interactions responsible for vinculin trimer formation, w e performed

binding assays with HV derivatives immobilized onto a solid phase to restrict conformational changes.

These assays indicated that A483 and A524 bound to HV with a similar affinity, as estimated by their

EC50 (95% confidence interval) of 6.1 (4.2-9.0) and 3.7 (1.7-8.1) nM, respectively (Suppl. Fig. 1a).

Strikingly, a large difference was observed in the binding plateau, indicating that HV presented more

binding sites for A483 than for A524 (Suppl. Fig. 1a). Also, D1D2 presented more binding sites for

HV_{1-258} containing the D1 domain only, suggesting the presence of additional sites on the D2 domain

(Suppl. Fig. 1b). Consistently, BN-PAGE showed the formation of 1:1, as well as a 1:2 D1D2:A483

complex, observed with increasing A483 molar ratio (Suppl. Fig. 1c). In contrast, single 1:1 complexes

were observed for D1:A483, D1:A524 or D1D2-A524 (Suppl. Figs. 1c-h), indicating that IpaA VBS3 was

required to reveal additional sites on the D2 domain. Of note, D1D2 homo-trimers observed in the

SEC-MALS and native gel analysis (Figs. 1d, e and Ext Data Figs. 3c, d) were not detected in BN-PAGE,

suggesting that Coomassie brilliant blue interfered with the formation of higher order D1D2

complexes. Together, these results suggested that the formation of vinculin trimers triggered by

A483 required the IpaA VBS3 dependent exposure of binding sites on D2. These findings were

| 96 | unexpected, since vinculin activating ligands have been described to bind to a single site on the D1 |
| 97 | domain of vinculin. |

To map interactions of A524 and A483 with D1D2, complexes were cross-linked, subjected to

proteolysis and analyzed using Liquid Chromatography coupled to Mass Spectrometry (LC-MS)

(Methods). Intermolecular links were identified from the characterization of cross-linked peptides

(Suppl. Tables 1-3; Methods) and along with identified intramolecular links, used to produce

structural models (Methods). The A524:D1 complex showed links consistent with a "canonical"

conformer expected from established structures[11, 12, 15] (Suppl. Fig. 2). Similar links were identified for

the A524:D1D2 complex, with a majority of links observed with the D1 domain (Fig. 2a). For both

complexes, the structure shows interactions between IpaA VBS1 and VBS2 with the D1 first and

second bundles, respectively, leading to helical bundle reorganization of D1 associated with vinculin

activation (Figs. 2c, d; Suppl. Fig. 2 and Suppl. Tables 1, 2). For the A483:D1D2 complex, structural

modeling shows 2 major conformers accounting for the majority of links. In a first "closed"

conformer, IpaA VBS1 and VBS2 interact with the D1 bundles in a similar manner as for A524, where

the relative positioning of D1 and D2 is globally conserved compared to apo D1D2 in the A524-D1D2

complex (compare Figs. 2d and 2e). In this "closed" conformer, IpaA VBS3 interacts with an interface

formed by the H5 (residues 128-149) and H8 (residues 222-250) helices in the second bundle of D1

and the H13 (residues 373-397) helix in the second bundle of the D2 domain (Figs. 2e and f). The

second "open" conformer, however, shows a major re-orientation of D1 and D2 subdomains with

their major axis forming an angle value of ca 82° compared to the 25° observed in the native vinculin

structure or the first conformer (Fig. 2g), with IpaA VBS3 docking sidewise through extensive

interaction with the H5 (residues 128-149) and H8 (residues 222-250) helices of the vinculin D1

domain. Because this latter conformer leads to major changes in bundle exposure in D1 and D2, we

posit that it is involved in the formation of higher order D1D2 complexes and trimer induced by A483.

To test this, we engineered a structural clamp by substituting residue Q68 in the first D1 bundle and

A396 in the second D2 bundle for cysteine residues, expected to prevent the formation of the open

conformer upon disulfide bridge formation (Suppl. Figs. 3a, b). Consistently, cysteine clamped D1D2-Q68C A396C did not prevent the exposure of additional sites on D2 or 1:1 complex formation induced by A524 or A483, but prevented the formation of higher order complexes (Fig. 1 e and Suppl. Figs. 3c-e). We coined "supra-activation" the mode of vinculin activation induced by A483 involving major conformational changes in the vinculin head, to distinguish it from the canonical activation associated with the dissociation of vinculin head-tail domains.

We then characterized the effects of A524 and A483 expression by performing immunofluorescence staining of vinculin-containing adhesion structures. As shown in Figs 3a-c, C2.7 cells transfected with GFP-A524 formed more numerous and larger peripheral adhesions as well as actin-rich ruffles compared to control cells (Figs. 3a-d). Strikingly, GFP-A483 transfected cells formed even larger and more numerous adhesions than GFP-A524 transfected cells, but significantly less actin ruffles than GFP-A524 transfectants (Figs. 3a-d). These observations were confirmed by live TIRF microscopy showing the extreme stability of adhesions in GFP-A483 transfectants, with a median duration of at least 84 min, while GFP-A524 and control cells showed adhesions with a comparable median duration of less than 25 min (Fig. 3e; Suppl. movie 1). GFP-A524 and GFP-A483 transfectants showed slightly slower median instant rates of adhesion assembly than control cells (Suppl. Fig. 4; Suppl. movie 1) but significant 1.6-fold and 2-fold decrease in median instant rates of disassembly relative to control cells, respectively (Suppl. Fig. 4). The stability of GFP-A483-induced FAs was independent of acto-myosin contraction. Indeed, GFP-A483-induced FAs resisted the action of the Rho-kinase inhibitor Y27632 relaxing actin-myosin, with a five- and four-times slower median rate of disassembly of adhesions relative to control cells and GFP-A524 transfectants, respectively (Figs. 3f-i; Suppl. movie 2). Furthermore, large adhesions formed in GFP-A483 transfectants following addition of the inhibitor (Figs. 3f, h), a process that was not observed for the other samples, including cells transfected with GFP-A524 or GFP fused to the vinculin D1 domain (vD1) reported to delay talin refolding following stretching [16-18] (Figs. 3g-i; Suppl. movie 2). In line with the stabilization of mature FAs through vinculin supra-activation, GFP-A483 but not GFP-vD1 also delayed the Y27632-induced

removal of the late adhesion marker VASP (Suppl. Fig. 5; Suppl. movie 3). Consistent with A483 bypassing mechanotransduction, GFP-A483 expression induced 5- and 1.6-fold higher yield of short term (≤ 10 min) cell adhesion to fibronectin substrates compared to control GFP and GFP-A524 transfectants, respectively (Suppl. Fig. 6a). By contrast, little difference in adhesion yield was detected at 15 min suggesting that canonical activation of vinculin also resulted in its supra-activation during mechanotranduction (Suppl. Fig. 6a). In line with this, MEF vinculin-null cells transfected with the clamped vinculin version formed fewer and smaller FAs than cells transfected with wild-type vinculin but more than mock-transfected cells (Suppl. Figs. 6b-d).

To confirm and extend these findings, cell adhesion strength was assessed measuring their detachment upon application of a controlled shear stress in a microfluidic chamber. When 1205Lu melanoma cells were allowed to adhere to fibronectin-coated surfaces for more than 25 min, little difference in resistance to shear stress could be detected among samples (Suppl. Fig. 7a). In contrast, when shear stress was applied less than 20 min following cell incubation, GFP-A483-transfected cells showed significantly higher resistance to shear stress up to 22.2 dynes.cm^{-2} than GFP-A524- or GFP-transfected cells, with 1.7 ± 0.2 -and 0.9 ± 0.14-fold enrichment ± SD of adherent cells for GFP-A483 and GFP-A524-transfected cells versus control GFP-transfected cells, respectively (Fig. 4b; Suppl. movie 4). In addition, similar to melanoma cells depleted for by siRNA treatment, cells transfected with the clamped vinculin version showed a decreased ability to adhere in comparison to WT vinculin-transfected cells (Suppl. Figs. 7a, b). These results are in full agreement with effects observed on adhesion structures and suggest that A483 interaction with vinculin leads to the bypass of mechanotransducing steps to promote strong adhesion.

Vinculin is paradoxically described as a prognostic marker favoring the migration of cancer cells or as a tumor suppressor stimulating cell anchorage [19-21]. These contradictory findings reflect its complex and poorly understood regulation, as well as different roles in 2D or 3D systems [22, 23]. Also, an increase of the total pool of vinculin may not correlate with increased vinculin activation. We took advantage of the unique property of A483 to study the effects of vinculin supra-activation on the

motility and invasion of melanoma cells. In time-lapse microscopy experiments in 2D-chambers, GFP-

A524 inhibited melanocyte motility compared to control cells, with a rate of Root Median Square

Displacement (rMSD) of 3.16 and 15.6 µm.min, respectively (Figs. 4c, d). An even stronger inhibition

was observed for GFP-A483-transfected cells (rMSD = 2.3 µm.min) (Figs. 4c, d). Transmigration of

melanocytes in 3D-matrigels was similarly inhibited by A524 and A483 (Fig. 4e).

Bacteria invading through a triggering mode rely on a discrete number of T3SS-dependent

contacts for which cytoskeletal tethering is likely critical for invasion[2]. As opposed to physiological

substrates, bacteria cannot sustain the range of counter-forces associated with integrin-mediated

adhesion to the substrate. The *Shigella* type III effector IpaA provides an elegant solution to this

problem by promoting strong adhesion without requirement for mechanotransduction. Through the

joint action of its VBSs, IpaA induces major conformational changes of the vinculin head sub-

domains. Understanding how these major vinculin conformational changes regulate the composition

and properties of cell adhesions will bring important insights into cell adhesion processes and will be

the focus of future investigations.

REFERENCES

1   Ogawa, M., Handa, Y., Ashida, H., Suzuki, M. & Sasakawa, C. The versatility of *Shigella* effectors.

Nat Rev Microbiol **6**, 11-16, doi:10.1038/nrmicro1814 (2008).

2   Valencia-Gallardo, C. M., Carayol, N. & Tran Van Nhieu, G. Cytoskeletal mechanics during *Shigella*

invasion and dissemination in epithelial cells. *Cellular microbiology* **17**, 174-182,

doi:10.1111/cmi.12400 (2015).

3   Tran Van Nhieu, G., Ben-Ze'ev, A. & Sansonetti, P. J. Modulation of bacterial entry into epithelial

cells by association between vinculin and the *Shigella* IpaA invasin. *The EMBO journal* **16**, 2717-

2729, doi:10.1093/emboj/16.10.2717 (1997).

4   Atherton, P., Stutchbury, B., Jethwa, D. & Ballestrem, C. Mechanosensitive components of

integrin adhesions: Role of vinculin. *Exp Cell Res*, doi:10.1016/j.yexcr.2015.11.017 (2015).

5   Yan, J., Yao, M., Goult, B. T. & Sheetz, M. P. Talin Dependent Mechanosensitivity of Cell Focal

Adhesions. *Cellular and molecular bioengineering* **8**, 151-159, doi:10.1007/s12195-014-0364-5

(2015).

6   Burridge, K. & Guilluy, C. Focal adhesions, stress fibers and mechanical tension. *Exp Cell Res*,

doi:10.1016/j.yexcr.2015.10.029 (2015).

7   Parsons, J. T., Horwitz, A. R. & Schwartz, M. A. Cell adhesion: integrating cytoskeletal dynamics

and cellular tension. *Nat Rev Mol Cell Biol* **11**, 633-643, doi:10.1038/nrm2957 (2010).

8   Humphries, J. D. *et al.* Vinculin controls focal adhesion formation by direct interactions with talin

and actin. *J Cell Biol* **179**, 1043-1057, doi:10.1083/jcb.200703036 (2007).

9   Lavelin, I., Wolfenson, H., Patla, I., Henis, Y. I., Medalia, O., Volberg, T., Livne, A., Kam, Z., Geiger,

B. Differential effect of actomyosin relaxation on the dynamic properties of focal adhesion

proteins. PLoS One. 8:e73549 (2013).

10   Bakolitsa, C. *et al.* Structural basis for vinculin activation at sites of cell adhesion. *Nature* **430**,

583-586, doi:10.1038/nature02610 (2004).

11   Izard, T., Tran Van Nhieu, G. & Bois, P. R. Shigella applies molecular mimicry to subvert vinculin

and invade host cells. *J Cell Biol* **175**, 465-475, doi:10.1083/jcb.200605091 (2006).

12   Tran Van Nhieu, G. & Izard, T. Vinculin binding in its closed conformation by a helix addition

mechanism. *The EMBO journal* **26**, 4588-4596, doi:10.1038/sj.emboj.7601863 (2007).

13   Park, H., Valencia-Gallardo, C., Sharff, A., Tran Van Nhieu, G. & Izard, T. Novel vinculin binding

site of the IpaA invasin of *Shigella*. *J Biol Chem* **286**, 23214-23221, doi:10.1074/jbc.M110.184283

(2011).

14   Valencia-Gallardo C, Bou-Nader C, Aguilar D, Carayol N, Quenech'Du N, Pecqueur L, Park HJ,

Fontecave M, Izard T, Tran Van Nhieu G. *Shigella* IpaA binding to talin stimulates filopodial

capture and cell adhesion. *Cell Reports* **26**, 921-932. doi: 10.1016/j.celrep.2018.12.091 (2019).

15   Izard, T. *et al.* Vinculin activation by talin through helical bundle conversion. *Nature* **427**, 171-

175, doi:10.1038/nature02281 (2004).

16   del Rio, A. *et al.* Stretching single talin rod molecules activates vinculin binding. *Science* **323**, 638-

641, doi:10.1126/science.1162912 (2009).

17   Margadant, F. *et al.* Mechanotransduction in vivo by repeated talin stretch-relaxation events

depends upon vinculin. *PLoS Biol* **9**, e1001223, doi:10.1371/journal.pbio.1001223 (2011).

18   Carisey, A. *et al.* Vinculin regulates the recruitment and release of core focal adhesion proteins in

a force-dependent manner. *Curr Biol* **23**, 271-281, doi:10.1016/j.cub.2013.01.009 (2013).

19   Goldmann, W. H. Role of vinculin in cellular mechanotransduction. *Cell Biol Int* **40**, 241-256,

doi:10.1002/cbin.10563 (2016).

20   Labernadie, A. *et al.* A mechanically active heterotypic E-cadherin/N-cadherin adhesion enables

fibroblasts to drive cancer cell invasion. *Nat Cell Biol* **19**, 224-237, doi:10.1038/ncb3478 (2017).

21   Hamidi, H. & Ivaska, J. Every step of the way: integrins in cancer progression and metastasis. *Nat*

Rev Cancer, doi:10.1038/s41568-018-0038-z (2018).

22   Mierke, C. T. *et al.* Vinculin facilitates cell invasion into three-dimensional collagen matrices. *J*

Biol Chem **285**, 13121-13130, doi:10.1074/jbc.M109.087171 (2010).

23   Gulvady, A. C., Dubois, F., Deakin, N. O., Goreczny, G. J. & Turner, C. E. Hic-5 expression is a

major indicator of cancer cell morphology, migration, and plasticity in three-dimensional

matrices. *Mol Biol Cell* **29**, 1704-1717, doi:10.1091/mbc.E18-02-0092 (2018).

ACKNOWLEDGEMENTS

The authors thank Gauthier Mercante for technical help, Philippe Mailly for help with image analysis and René-Marc Mège for insightful discussions and reading of the manuscript. This work was supported by grants from Inserm, CNRS and Collège de France to the CIRB, as well as grant from the PSL Idex project "Shigaforce". DIAS and BCC are recipients of a PhD fellowship from a CONACYT scholarship. CV-G and DIAS also received support from the Memolife Labex. HK and LM were supported by Swiss National Science Foundation (grant no. SNF 200021 160188) and LM by the Knut And Alice Wallenberg Foundation (grant no. KAW 2016.0023).

AUTHOR CONTRIBUTIONS

CV-G and DIAS conceived and performed most of the experimental works, data analysis and

wrote the manuscript. BCC and BM analyzed TIRF experiments. CB-N performed the SEC-MALS

analysis. CB and NQD performed and analyzed experiments with melanocytes with the help of DJ and

AM. BCC, AA and JF provided technical help for cell adhesion and microfluidics experiments. CM and

JCR designed and performed the LC-MS analysis. DBL analyzed the cross-linked mass spectrometry

data. HK and LM generated structural models. GTVN designed the project and wrote the manuscript.

261   **Figure 1 – IpaA reveals binding sites in vinculin head subdomains**

262   **a,** Scheme of folded vinculin (HV). The binding sites and corresponding ligands are indicated. **b,**

263   Scheme of HV and IpaA constructs. HV domains and IpaA VBSs are depicted as boxes. The numbers

264   indicate the start residue of each domain. **c, d,** SEC elution profiles of complexes formed between A483

265   (green) or A524 (purple) and vinculin D1D4 (**c**) or D1D2 (**d**). The indicated complex stoichiometry was

266   inferred from the molecular weight estimated by MALS. **e**, native gel analysis of vinculin D1D2 and IpaA

267   derivatives. D1D2 or cysteine-clamped D1D2 (S-S) were incubated with the indicated IpaA derivatives

268   and analyzed by native PAGE followed by Coomassie staining. Arrows: 1:1 complexes. *: higher order

269   complexes. Note the absence of higher order complexes for cysteine-clamped D1D2.

Figure 2 – Characterization of IpaA contact sites on vinculin.

a, **b**, EDC cross-link map from mass spectrometry analysis (LC-MS/MS) of vinculin D1D2-A524 (**a**) and

D1D2-A483 (**b**) following extraction of 1:1 complexes from BN-PAGE. Blue lines: inter-molecular links.

Red lines: intra-molecular links. Note the links between IpaA VBS3 and the D2 second bundle. Cross-

linked residues are detailed in Suppl. Table 1. **c-e**) Structural models of D1D2 (**c**), D1D2-A524 (**d**), D1D2-

A483 "closed" conformer (**e**), D1D2-A483 "open" conformer (**g**). **f**, **h**, higher magnification of the IpaA

VBS3-D1D2 interaction in (**e**) and (**g**) showing the distance between residues in Å. IpaA VBS1-3 were

docked on the surface of Vinculin D1D2 using MS cross-link constraints. TX-MS protocol in combination

with MS constraints was used to unify and adjust the final model, which justifies over 100 cross-links.

Figure 3 – A483 stabilizes vinculin in cell adhesions in the absence of mechanotransduction

a-c, Immunofluorescence analysis of vinculin adhesions. CTRL: C2.7 cells. A524: GFP-A524

transfectants. A483: GFP-A483 transfectants. **a,** representative fluorescence micrographs. Arrows:

adhesions; arrowheads: ruffles. Green: GFP; red: vinculin; cyan: actin. **b**, percent of cells with ruffles ±

SEM. Cells with: no ruffles (empty bars); small ruffles (light grey bars); large ruffles (dark grey bars). *:

Pearson's Chi-squared test (N=3, n > 30, p = 0.036). **c, d**, vinculin containing FAs were detected using a

semi-automated program. **c**, median number of FAs per cell; **d**, FA size. (N=3, n > 30). Dunn's multiple

comparisons test. *: $p < 0.05$;***: $p < 0.005$. **e**, TIRF microscopy of C2.7 cells transfected with mCherry-

vinculin alone (CTRL) or co-transfected with GFP-A524 (A524), GFP-vD1 (vD1), or GFP-A483 (A483). The

duration of vinculin-containing adhesions was determined from time-lapse acquisitions. **f-i**, TIRF

microscopy analysis of cells treated with 100 μM Y-27632. **f**, Representative time series acquisitions.

Numbers indicate the elapsed time in seconds, with the inhibitor added at t = 0. Scale bar = 5 μm. **g, h**,

F/F_0: normalized average fluorescence intensity of adhesions as a function of time. Representative traces

corresponding to single adhesions for: **h**, control cells: blue, A524: red, vD1: green; **g**, A483: purple and

orange. **i,** initial rates of adhesion disassembly inferred from linear fits. N = 5. Number of adhesions

analyzed: CTRL: 84; vD1: 75; A524: 140; A483: 97. Dunn's multiple comparisons test. *: $p < 0.05$; **: $p <$

0.01;***: $p < 0.005$; ****: $p < 0.001$.

Figure 4. A483-mediated vinculin supra-activation stimulates strong cell adhesion and inhibits tumor cell invasion.

a, b, 1205Lu melanoma cells were transfected with the indicated constructs, labeled with calcein (methods) and mixed with the same ratio of control cells. Cells were perfused in a microfluidic chamber and allowed to adhere for 20 min prior to shear stress application. **a,** representative fields. The number indicates the elapsed time after shear stress application. **b,** Scatterplot of fold enrichment of: A483 (N = 4, n = 610) or A524 (N = 4, n = 433) transfected cells vs control cells (1594 cells, N = 4). Unpaired t test. *: p = 0.0229. **c, d,** cells were transfected with the indicated constructs, and analyzed by time-lapse videomicroscopy. **c,** representative single cell migration 20-hour tracks for indicated samples. **d,** Root of Median Square of displacement over time for control- (61 cells, N = 3), GFP-A524- (61 cells, N = 3) and GFP-A483 transfectants (64 cells, N = 3). ***p = 0.0007. The slopes were analyzed using a covariance test and found to be statistically different (ANCOVA, $p < 2 \times 10^{-16}$). **e,** 5×10^{4} cells were seeded in matrigel

chambers. The percent of transmigrated cells is indicated. (N = 3). Kruskal-Wallis test with Dunn's

multiple comparisons test. *: $p < 0.05$.

METHODS

Generation of constructs

Human vinculin constructs were generated by polymerase chain reaction using the forward primer

5' GCGCATATGCCAGTGTTTCATACG-3' and reverse primer 5'-CGTCGACTCACCAGGCATCTTCATCGGC-3' for

D1 (residues 1-258) or 5'-CGTCGACTCAGTGTACAGCTGCTTTG-3' for D2 (residues 1-492) using a plasmid

containing full-length octahistidine-tagged human vinculin (residues 1–1,066), as template[10], and cloned

into the NdeI-SalI sites of pet15b (Novagen). The IpaA constructs GFP-A524 and GFP-A483 were

generated by polymerase chain reaction (PCR) and cloning into pcDNA3.1 NT-GFP Topo TA (Invitrogen)

using the 5'-TCAAAGGACATTACAAAATCC-3' and 5'-GCGATATCATGGCCAGCAAAGG-3' forward primers,

respectively, and the 5'-GCGCGGCCGCTTAATCCTTATTGATATTC-3' reverse primer. The GST-A483A

construct was generated by PCR using 5'-GGCGAATTCCCGGAGACACATATTTAACACG-3' forward and 5'-

GCCGTCGACTTAATCCTTATTGATATTCT-3' reverse primers and cloning into the *EcoRI-SalI* ofpGEX-4T-2

(GE Lifesciences). pGST-A524 was previously described[24]. The pGFP-vD1 plasmid was generated by

polymerase chain reaction using the forward primer 5'-ACCCGGGATCCCGCC-3' and reverse primer 5'-

ACCCGGGACCAGGCA-3', and cloned into peGFP. The pmCherry-human vinculin (HV) and pmCherry-VASP

plasmids were from Addgene. Stealth siRNA anti-human vinculin was from Invitrogen (reference number

1299001).

Protein purification

BL21 (DE3) chemically competent *E. coli* (Life Technologies) was transformed with the expression

constructs. D1 and D1D2 were purified essentially as described[11, 25]. For the IpaA derivatives, bacteria

grown until OD_{600nm} = 1.0 were induced with 0.5 mM IPTG and incubated for another 3 hrs. Bacteria

were pelleted and washed in binding buffer 25 mM Tris PH 7.4, 100 mM NaCl and 1 mM beta-

mercaptoethanol, containing Complete[TM] protease inhibitor. Bacterial pellets were resuspended in

1/50th of the original culture volume and lyzed using a cell disruptor (One shot model, Constant System

Inc.). Proteins were purified by affinity chromatography using a GSTrap HP affinity column (GE

Healthcare) and size exclusion chromatography (HiLoad S200, Ge Healthcare). Samples were stored

aliquoted at -80°C at concentrations ranging from 1 to 10 mg/ml.

Protein complex formation analysis

Proteins were incubated at a concentration of 30 µM in binding buffer for 60 min at 4°C. Samples

were analyzed by SEC-MALS (Wyatt Technology Europe) using a 24 ml Superdex 200 Increase 10/300 GL

filtration column and a MiniDAWN TREOS equipped with a quasi-elastic light scattering module and a

refractometer Optilab T-rEX (Wyatt Technology). Data were analyzed using the ASTRA 6.1.7.17 software

(Wyatt Technology Europe). Protein complex formation was visualized by PAGE under non-denaturing

conditions using à 7.5% polycrylamide gel, followed by Coomassie blue staining.

Solid-phase binding assay

96–well Maxisorp (Nunc) ELISA plates were coated with 30 nM of full-length vinculin, vinculin

constructs or IpaA proteins at the indicated concentrations in binding buffer (25 mM Tris PH 7.4, 100

mM NaCl and 1 mM β-mercaptoethanol). Samples were blocked with PBS-BSA 2%, washed and

incubated with IpaA or vinculin proteins in binding buffer containing 0.2% BSA at room temperature for

one hour. After incubation, the plates were washed and incubated with an anti-IpaA (dilution 1/2000[e])

polyclonal primary antibody[3] or anti-vinculin (dilution 1/2000[e]) Vin11Vin.5 monoclonal antibody (Sigma-

Aldrich) in binding buffer containing 0.2% BSA for one hour at room temperature. Plates were washed

and incubated with an HRP-coupled secondary anti-rabbit or anti-mouse IgG antibody (1/32000[e])

(Jackson ImmunoResearch) for one hour. The reaction was revealed by adding 100 µl of

tetramethylbenzidine (Sigma-Aldrich) for 15 min, stopped by adding 50 µl of 0.66N H_2SO_4 and the

absorbance was read at 450 nm (Dynatech MR400).

BN-PAGE (Blue Native – Polyacrylamide Gel Electrophoresis) protein native gel analysis and

complex cross-linking

25 μM of vinculin constructs were incubated with different molar ratios of IpaA proteins in a 1X BN-PAGE buffer (250 mM ε-aminicaprionic acid and 25mM Bis-Tris PH 7,0) at 4°C for one hour. The protein mixtures were separated in a one-dimension native BN-PAGE electrophoresis as described[26]. For vinculin-IpaA protein ratio assay, vinculin-IpaA bands containing the complexes separated by BN-PAGE were cut, sliced and boiled in 2 x Laemmli SDS buffer followed by SDS-PAGE. The second dimension SDS-PAGE gels were stained (colloidal Coomassie staining) and the density of the bands was determined using Image J. The normalized vinculin:IpaA ratio of the complexes was compared using a non-parametric Kruskal-Wallis rank sum test (R statistical software).

For crosslinking vinculin-IpaA complex, bands containing the complexes were cut, sliced and electroeluted in native conditions (15 mM Bis-Tris pH 7.0 and 50 mM Tricine) inside a closed dialysis membrane (SpectraPor). The soluble complexes were recovered and their buffer exchanged twice into an amine-free cross-link buffer in 25 mM HEPES pH 7.0 containing 100 mM NaCl using 10MWCO ZEBA desalting columns (Thermo Scientific). The fractions containing the complexes were incubated for 1 hr at 4°C with 10 mM N-hydroxysulfosuccinimide and 5 mM EDC (Sigma-Aldrich) following the manufacturer's recommendations. The cross-linking reaction was stopped by adding 50 mM Tris pH 7.4 and incubating for 20 minutes. Samples were denaturated in 2x SDS Laemmli buffer for 5 min at 95°C and complexes were eluted from gel slices following SDS-PAGE.

Liquid Chromatography Mass spectrometry (LC-MS)

Complexes obtained after the cross-linking step were loaded onto a 4-20% polyacrylamide gradient gels and Coomassie stained. The bands containing the complexes were cut and submitted to tryptic digestion[27]. The experiments were performed in duplicates for the 3 complexes D1:A524, D1D2:A524 and D1D2:A483. Peptides obtained after tryptic digestion were analyzed on a Q Exactive Plus instrument (Thermo Fisher Scientific, Bremen) coupled with an EASY nLC 1 000 chromatography system (Thermo

Fisher Scientific, Bremen). Sample was loaded on an in-house packed 50 cm nano-HPLC column (75 µm

inner diameter) with C18 resin (1.9 µm particles, 100 Å pore size, Reprosil-Pur Basic C18-HD resin, Dr.

Maisch GmbH, Ammerbuch-Entringen, Germany) and equilibrated in 98 % solvent A (H2O, 0.1 % FA) and

2 % solvent B (ACN, 0.1 % FA). A 120 minute-gradient of solvent B at 250 nL.min-1 flow rate was applied

to separate peptides. The instrument method for the Q Exactive Plus was set up in DDA mode (Data

Dependent Acquisition). After a survey scan in the Orbitrap (resolution 70 000), the 10 most intense

precursor ions were selected for HCD fragmentation with a normalized collision energy set up to 28.

Charge state screening was enabled, and precursors with unknown charge state or a charge state of 1

and >7 were excluded. Dynamic exclusion was enabled for 35 or 45 seconds respectively.

Data analysis

The identification of cross-linked peptides from LC-MS data was performed using SIM-XL v. 1.3[28],

with the following search parameters: EDC as cross-linker, a tolerance of 20 ppm for precursor and

fragment ions, trypsin fully specific digestion with up to three missed cleavages. Carbamidomethylation

of cysteines was considered as a fixed modification. All initial identification of cross-linked peptides

required a primary score of SIM-XL greater than 2.5 for inter-links and 2.0 for intra-links or loop-links. As

single incorrect cross-link identification might lead to a different model, a manual post-validation of the

search engine results at the MS2 level was thus performed. A 2D-map showing the protein-protein

interaction was generated as an output (Figs. 2a,b). Only peptides present in the 2 replicates are

gathered in Supplementary Tables 1-3 and were used for the modeling.

Modeling

We used the constraints obtained from the cross-linking MS data (Suppl. Tables 1-3) to guide the

protein structure modeling using the TX-MS protocol as described by Hauri, Khakzad et al.[29]. In short,

TX-MS uses the Rosetta comparative modeling protocol (RosettaCM)[30], and the flexible backbone

docking protocol (RosettaDock)[31] to generate models and evaluate how well each model explains the MS

constraints using a novel scoring function. Here, a total of 100,000 models was generated, of which the

highest-scoring model is displayed in (Fig2e), supported by a total of 100 inter and intra-molecular cross-

links.

Cell lines

C2.7 myoblasts[32] and MEF vinculin null cells[8] were routinely grown in DMEM 1 g / L glucose

containing 10 % FCS in a 37°C incubator containing 10 % CO_2. 1205Lu melanoma cells[33] were grown in

RPMI + Glutamax medium (RPMI1640) supplemented with 10 % fetal calf serum (FCS) and non-essential

aminoacids in a 37°C incubator with 5 % CO_2.

Immunofluorescence staining

C2.7 mice myoblasts cells were seeded at 2.5×10^4 cells in 25 mm-diameter coverslips. Cells were

transfected with 3 μg of pGFP-A524 or pGFP-A483 plasmids with 6 μl JetPEI transfection reagent

(Polyplus) for 16 hours following the manufacturer's recommendations. Cells were fixed in PBS

containing 3.7% paraformaldehyde for 20 min at 21°C and permeabilized with 0.1% Triton X-100 for 4

min at 21°C. Cells were processed for immunofluorescence staining using the Vin11.5 anti-vinculin

monoclonal antibody (ref. V4505, Sigma-Aldrich) and anti-mouse IgG antibody coupled to Alexa 546

(Jackson Research) and Phalloidin-Alexa 633 (Invitrogen), as described previously[3]. Samples were

analyzed using an Eclipse Ti inverted microscope (Nikon) equipped with a 60 x objective, a CSU-X1

spinning disk confocal head (Yokogawa), and a Coolsnap HQ2 camera (Roper Scientific Instruments),

controlled by the Metamorph 7.7 software.

TIRF (Total Internal Reflection Microscopy) analysis

C2.7 cells were transfected with pmCherry-HV or pmCherry-VASP and the indicated plasmids as

described above. Samples were mounted onto a TIRF microscopy chamber on a stage of an Eclipse Ti

inverted microscope (Nikon) equipped with an Apo TIRF 100 x N.A. 1.49 oil objective heated at 37°C. TIRF

analysis was performed using the Roper ILAS module and an Evolve EM-CCD camera (Roper Scientific

Instruments). When mentioned, Y-27632 was used at 100 µM. Image acquisition was performed every

12.5 seconds for 30 to 90 minutes.

Live cell tracking

1205Lu melanoma cells were transfected with IpaA constructs or GFP alone (control) and

transferred in microscopy chamber on a 37°C 5%-CO_2 stage in RPMI1640 medium containing 25 mM

HEPES. For cell tracking, samples were analyzed using and inverted Leica DRMIBe microscope and a 20 X

phase contrast objective. Image acquisitions were performed every 3 min for 200 hrs. The mean velocity

of migration was measured for all tracks followed for at least 5 hours. The root square of MdSD over

time was plotted over time and fitted by linear regression. The slopes of the linear fit were compared

using an ANCOVA test (linear model). The median cell surface was quantified as the mean of the surface

for three time points (25%, 50% and 75%) of the whole cell track and dispersion measured by the

Median absolute dispersion (MAD).

Invasion assays

Tissue culture Transwell inserts (8 µm pore size; Falcon, Franklin Lakes, NJ) were coated for

3 hours with 10 µg of Matrigel following the manufacturer's instructions (Biocoat, BD Biosciences, San

Jose, CA). Inserts were placed into 24-well dishes containing 500 µl of RPMI medium supplemented

with 1% fetal calf serum. 5×10^4 melanoma cells were added to the upper chamber in 250 µls of serum-

free RPMI medium. After 24 hours, transmigrated cells were scored by bright field microscopy.

Experiments were performed at least three times, each with duplicate samples.

Image processing and statistical analysis

For the quantification of the number of adhesion structures in C2.7 cells, a semi-automated

protocol was developed using Icy software[34]. Spinning-disk fluorescent microscopy planes were used to

detect vinculin structures using HK means thresholding and overlaid binary masks obtained from the

threshold projections of F-actin labeled images (Max-entropy method). FAs were detected as spots

positive for both vinculin mCherry and actin structures using Wavelet Spot Detector. The number of

adhesions was analyzed using Dunn's multiple comparisons test. The statistical analysis of cell motility

was performed in the R software. Medians were compared using a Wilcoxon rank sum test and

dispersion by Median absolute dispersion (MAD) parameter.

Microfluidics cell adhesion assay

Analysis of cell detachment under shear stress was based on previous works[35]. 1205Lu

melanocytes were transfected with the indicated constructs, then labeled with 2 μ⃞ calcein-AM (Life

Technologies) in serum-free DMEM for 20 minutes. Cells were detached by incubation with 2

μ⃞ Cytochalasin D (Sigma-Aldrich) for 40 minutes to disassemble FAs, followed by incubation in PBS

containing 10 mM EDTA for 20 minutes. Cells were washed in EM buffer (120 mM NaCl, 7 mM KCl, 1.8

mM $CaCl_2$, 0.8 mM $MgCl_2$, 5 mM glucose and 25 mM HEPES at pH 7.3) by centrifugation and resuspended

in the same buffer at a density of 1.5 x 10^6 cells/ml. Calcein-labeled transfected cells and control

unlabeled cells were mixed at a 1:1 ratio and perfused onto a 25 mm-diameter glass coverslips

(Marienfeld) previously coated with 20 μg/ml fibronectin and blocked with PBS containing 2% BSA

(Sigma-Aldrich) in a microfluidic chamber on a microscope stage at 37°C. We used a commercial

microfluidic setup (Flow chamber system 1C, Provitro) and a Miniplus3 peristaltic pump (Gilson) to

adjust the flow rate in the chamber. Microscopy analysis was performed using a LEICA DMRIBe inverted

microscope equipped with a Cascade 512B camera and LED source lights (Roper Instruments), driven by

the Metamorph 7.7 software (Universal imaging). Cells were allowed to settle for the indicated time

prior to application of a 4 ml/min, flow corresponding to a wall shear stress of 22.2 dyn/cm^2 (2.22 Pa).

Acquisition was performed using a 20 X objective using phase contrast and fluorescence illumination

(excitation 480 ± 20 nm, emission 527 ± 30 nm). Fluorescent images were acquired before and after

flushing to differentiate between target and control cells. Phase contrast images were acquired every

200 ms. Fold enrichment was defined as the ratio between of attached labeled and unlabeled cells.

REFERENCES TO METHODS

24   Ramarao, N. *et al.* Capping of actin filaments by vinculin activated by the Shigella IpaA carboxyl-

terminal domain. *FEBS letters* **581**, 853-857, doi:10.1016/j.febslet.2007.01.057 (2007).

25   Papagrigoriou, E. *et al.* Activation of a vinculin-binding site in the talin rod involves

rearrangement of a five-helix bundle. *The EMBO journal* **23**, 2942-2951,

doi:10.1038/sj.emboj.7600285 (2004).

26   Eubel, H. & Millar, A. H. Systematic monitoring of protein complex composition and abundance

by blue-native PAGE. *Cold Spring Harb Protoc*, pdb prot5221, doi:10.1101/pdb.prot5221 (2009).

27   Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M. Nat Protoc. In-gel digestion for mass

spectrometric characterization of proteins and proteomes. **1**(6):2856-60 (2006).

28   Lima, D. B. et al. SIM-XL: A powerful and user-friendly tool for peptide cross-linking analysis. J

Proteomics **129**, 51-55, doi:10.1016/j.jprot.2015.01.013 (2015).

29   Hauri, S., Khakzad, H., Happonen, L., Teleman, J., Malmström, J., & Malmström, L. Rapid

determination of quaternary protein structures in complex biological samples. Nature

communications, **10**(1), 192. doi:10.1038/s41467-018-07986-1 (2019).

30   Song, Y. *et al.* High-resolution comparative modeling with RosettaCM. *Structure* **21**, 1735-1742,

doi:10.1016/j.str.2013.08.005 (2013).

31   Gray, J. J. High-resolution protein-protein docking. *Current opinion in structural biology* **16**, 183-

193, doi:10.1016/j.sbi.2006.03.003 (2006).

32   Mitrossilis, D. et al. Single-cell response to stiffness exhibits muscle-like behavior. *Proc. Nat.*

Acad. Sci USA. **106**(43), 18243-18248; https://doi.org/10.1073/pnas.0903994106 (2009).

33   Smalley, K. S., et al. Increased cyclin D1 expression can mediate BRAF inhibitor resistance in

BRAF V600E-mutated melanomas. *Mol Cancer Ther.,* **7**(9):2876-83. doi: 10.1158/1535-

7163.MCT-08-0431 (2008).

34   de Chaumont, F. *et al.* Icy: an open bioimage informatics platform for extended reproducible

research. *Nat Methods* **9**, 690-696, doi:10.1038/nmeth.2075 (2012).

35   Gutierrez, E. *et al.* Microfluidic devices for studies of shear-dependent platelet adhesion. *Lab*

512   *Chip* **8**, 1486-1495, doi:10.1039/b804795b (2008).

Supplementary Fig. 1. IpaA VBS3 reveals multiple binding sites on vinculin.

a, b, Solid phase binding assays. **a,** coating: HV; ligands: A483 (solid circles); A524 (solid squares).

b, coating: D1 (solid circles) or HVD1D2 (solid squares); ligand: A483. **c, f-h,** BN-PAGE in 6-18%

polyacrylamide gradient gels and Coomassie staining analysis of D1D2:A483 (**c**), D1D2:A524 (**f**),

D1:A524 (**g**) or D1:A483 (**h**) complexes. The molar ratio is indicated above each lane.

Arrowheads indicate protein alone, or complex migration at the indicated molar ratio. **d,** bands

were recovered from BN-PAGE and analyzed in a second dimension SDS-PAGE in a 15% poly-

acrylamide gel and Coomassie staining. Bands were analyzed by densitometry. **e,** ratio of

density values for the A524-D1D2 complex (empty bar) and A483-D1D2 complexes

corresponding to the upper (light grey bar) or lower (dark grey bar) shifts.

Supplementary Figure 2. Structural models of vD1:A524. a, Structure predicted from the resolved vD1:

IpaA VBS1: and vD1:IpaA VBS2 crystal structures[11, 12]. **b,** Structural model of vD1:A524. The model was

established using RosettaCM protocol and accounts for 19 inter and intra-molecular cross-links out of 24

identified (Suppl. Table 1).

Supplementary Figure 3. Cysteine-clamped vinculin D1D2 does not form high order complexes. **a,** Structural model of cystein-clamped vinculin. Blue: D1 domain. Green: D2 domain in the closed conformer. Grey: D2 domain in the open conformer. Black: C68-C396 cystein clamp preventing the switch from closed to open conformers. **b-e,** Coomassie blue staining. **b,** Disulfide bridge formation in D1D2. D1D2: wild-type sequence. SS: D1D2 Q68C A396C. SDS-PAGE analysis using a 10 % polyacrylamide gel. + β-metOH: samples were boiled in Laemmli sample loading buffer containing 5 mM beta-mercaptoethanol prior to SDS-PAGE. The molecular weight markers in kDa are indicated. The black and red bars point the respective migration of unreduced and reduced D1D2 Q68C A396C, respectively. **c, d,** gel strips of the native-PAGE corresponding to the A524 + D1D2 (**c**) or A483 + D1D2 (**d**) as shown in **Fig. 1e** were analyzed in a second dimension by SDS-PAGE using a 10 % polyacrylamide gel as depicted. D1D2, SS, A483 and A524 are indicated. **e,** BN-PAGE analysis in a 6-18% polyacrylamide gradient gel of

D1D2:A483 (D1D2) and SS:A483 (SS) complexes. The molar ratio is indicated above each lane.

Arrowheads indicate protein alone, or complex migration

Supplementary Figure 4. TIRF analysis of FA dynamics. C2. 7 cells were transfected with HV-mCherry

(HV), HV-mCherry and GFP-A524 (A524) or HV and GFP-A483 (A483). The dynamics of HV-mCherry-

labeled FAs were analyzed by TIRF microscopy. **a**, traces correspond to the variations of average

fluorescence intensity of a representative single FA (F) normalized to its maximal average fluorescence

intensity over the analyzed period in seconds (F_{max}). Blue: HV; green: HV+A524; red: HV+A483. **b**, **c**,

instant assembly (**b**) or disassembly (**c**) rates were inferred from the slopes of linear fits as depicted in a),

with a Pearson correlation value > 0.85. HV: n = 41, N = 3; HV+A524: n = 31, N = 2; HV+A483: n = 55, N =

3. Mann-Whitney U test. *: $p < 0.05$

Supplementary Figure 5. A483 stabilizes VASP-containing cell adhesions in the absence of mechanotransduction

TIRF microscopy of C2.7 cells transfected with mCherry-VASP alone (CTRL) or co-transfected with GFP-A524 (A524), GFP-vD1 (vD1), or GFP-A483 (A483). Adhesion kinetic parameters were determined from time-lapse acquisitions of cells co-transfected with mCherry-VASP and the indicated construct following treatment with 100 µM Y-27632. **a**) Representative time series acquisitions. Numbers indicated the elapsed time in seconds, with the inhibitor added at t = 0. Scale bar = 5 µm. **b**, $\Delta F/F_0$: normalized average fluorescence intensity of adhesions as a function of time. Representative traces corresponding to single adhesions for the indicate samples. **c**, initial rates of adhesion disassembly inferred from linear

Supplementary Figure 6. Effects of A483 and vinculin supra-activation on cell adhesion kinetics

a, 1205Lu melanoma cells were transfected with GFP alone (blue), GFP-A524 (red) or GFP-A483 (green), lifted

up by trypsinization and plated for the indicated time on Fn-coated coverslips. Samples were washed, fixed and

adherent cells were scored microscopically. The total number of adherent cells scored is indicated. GFP: 3223

cells, N = 4; GFP-A524: n=7418, N = 4; GFP-A483: n = 5668, N = 4. Chi square corrected with Bonferroni mutliple

comparison correction. ****: p < 0.001. **b-d**, MEF vinculin null cells were co-transfected with GFP-talin (CTRL)

and full length vinculin-mCherry (HV) or HV Q68C A396C-mCherry (SS). Samples were fixed and processed for

fluorescence staining of F-actin. **b,** Representative fluorescence micrographs. Red: Vinculin-mCherry; green:

GFP-talin; cyan: F-actin. Scale bar = 1 µm. **c**, **d**, FAs were scored using a semi-automatic detection program

(Methods) and their number per cell (left) and size (right) are shown. Bar: median size. FAs labeled with: **c**,

GFP-talin; **d**, HV or SS. CTRL: n=28, N=3; HV: n = 25 , N = 3; SS: n = 25, N = 3. Mann-Whitney test with

Bonferroni multiple comparison correction. * p<0.05; ** p<0.01; *** p<0.005; **** p<0.001.

Supplementary Figure 7. Role of vinculin supra-activation on cell adhesion.

a, 1205Lu melanoma cells were transfected with the indicated constructs, labeled with calcein (methods) and

mixed with the same ratio of control cells. Cells were perfused in a microfluidic chamber and allowed to

adhere for 30 min (large solid circles) or 15 min (small solid circles) prior to shear stress application. Scatterplot

of the ratio of adherent cells: A483 (N = 3, n = 557); GFP (N = 3, n = 490); HV: vinculin mCherry (N = 3, n = 481);

SS: vinculin Q68C A396C-mCherry (N = 3, n = 259); siRNA: cells treated with anti-vinculin siRNA (N = 3, n = 395).

Unpaired t test. ***: p < 0.005. **b**, 1205Lu melanoma cells were mock-transfected (CTRL) or treated with anti-

vinculin siRNA (siRNA, Methods). Right: anti-vinculin Western blot analysis. Left: average HV band intensity

normalized to that of control cells. Unpaired t test. ***: p = 0.005.

#	Exp. MH⁺	Primary Score	Peptide Sequence	A524	Hv D1
32670	4457.1	3.28	IDDTSAELLTDDISDLKNNN**D**ITAENNNIYK - MA**K**MIDER	D604	K173
32798	1924.16	2.61	INN**K**LK - **E**LLPVLISAMK	K540	E200
33042	4613.23	3.70	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - NLGPGMT**K**MAK	D594	K170
33042	4613.23	3.70	IDDTSAELLTD**D**ISDLKNNNDITAENNNIYK - NLGPGMT**K**MAK	D595	K170
33042	4613.23	4.11	IDDTSAELLTDDISDLKNNN**D**ITAENNNIYK - NLGPGMT**K**MAK	D604	K170
33322	5310.64	4.48	IDDTSAELLTDDISDL**K**NNNDITAENNNIYK - VGKETVQTTE**D**QILKR	K600	D67
33334	2347.28	2.51	DVTTSLS**K**VLK - MSA**E**INEIIR	K625	E240
33432	2138.26	2.88	VL**K**NINKD - **E**LLPVLISAMK	K628	E200
33496	2315.32	3.09	AA**K**DVTTSLSK - **E**LLPVLISAMK	K617	E200
34822	2642.55	2.68	AKEVSSALS**K**VLSK - **E**LLPVLISAMK	K579	E200
38218	3685.86	3.87	NIN**K**D - TIESILEPVAQQISHLVIMHEEG**E**VDGK	K632	E31
38218	3685.86	3.66	NIN**K**D - TIESILEPVAQQISHLVIMH**E**EGEVDGK	K632	E28
#	Exp. MH⁺	Primary Score	Peptide Sequence	Hv D1	
34075	2342.29	3	**E**LLPVLISAMK - NLGPGMT**K**MAK	E200	K170

36135	3088.67	2.62	NFTVE**K**MSAEINEIIR - **E**LLPVLISAMK	K236	E200

#	Exp. MH$^+$	Primary Score	Peptide Sequence	A524	
14281	3051.56	3.52	NYVT**E**TNADTIDKNHAIYEK - INN**K**LK	E554	K540
15477	3342.67	2.74	NYVT**E**TNADTIDKNHAIYEK - A**K**EVSSALSK	E554	K571
17603	3141.54	2.71	NYVTETNADTID**K**NHAIYEK - **E**VSSALSK	K562	K572
31359	4483.2	2.64	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - A**K**EVSSALSK	E590	K571
31996	2965.5	2.94	IDDTSAELLTDDIS**D**LK - AA**K**DVTTSLSK	D598	K617
31996	2965.5	2.51	IDDTSAELLT**D**DISDLK - AA**K**DVTTSLSK	D594	K617
32705	4586.24	4.1	IDDTSAELLTDDISDLKNNNDITA**E**NNNIYK - AA**K**DVTTSLSK	E608	K617
33253	3291.74	3.81	IDDTSAELLTD**D**ISDLK - AKEVSSALS**K**VLSK	D595	K579
33997	5328.54	4.76	IDDTSAELLTDDISDL**K**NNNDITAENNNIYK - ID**D**TSAELLTDDISDLK	K600	D586
34007	4655.33	4.78	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - DVTTSLS**K**VLK	D594	K625

Suppl. Table 1. Cross-linked residues characterized in the D1:A524 complex (XL-amino acids

of each protein are bolded in the sequences).

#	Exp. MH$^+$	Primary	Peptide Sequence	A524	Hv D1D2

		Score			
20480	3470.68	3.86	NYVTE TNADTIDKNHAIYEK - NLGPGMT K MAK	E554	K170
23021	3883.92	3.71	NYVTETNADTID K NHAIYEK - E TVQTTEDQILKR	K562	E60
24707	3117.73	5.00	AKEVSSALSKVLSK - VGKETVQTT E DQILK	K579	E66
36724	4612.22	4.82	IDDTSAELLTDDISDLKNNN D ITAENNNIYK - NLGPGMT K MAK	D604	K170
39712	5255.49	3.79	IDDTSAELLTDDIS D LKNNNDITAENNNIYK - NLGPGMTKMA K MIDER	D598	K173
39893	5162.6	4.51	IDDTSA E LLTDDISDLKNNNDITAENNNIYK - ALA K QVATALQNLQTK	E590	K464
40011	2641.56	4.45	AKEVSSALS K VLSK - E LLPVLISAMK	K579	E200
41002	3047.64	3.99	L K VTDANIR - GILSGTSDLLLTFD E AEVR	K542	E128
41440	3944.01	3.86	IDDTSAELLTDDIS D LK - AQQVSQGLDVLTA K VENAAR	D598	K366
42699	3685.87	4.03	NIN K D -TIESILEPVAQQISHLVIMHEEG E VDGK	K632	E31

#	Exp. MH+	Primary Score	Peptide Sequence	Hv D1D2	
25162	3323.71	4.54	LNQA K GWLRDPSASPGDAGEQAIR - ALASI D SK	K281	D274
30025	3169.64	4.03	GQGSSPVAMQ K AQQVSQGLDVLTAK - V E NAAR	K352	E368
30045	3278.66	3.69	VLQLTSWDE D AWASK - K LEAMTNSKQSIAK	D275	K381
30201	3695.87	3.82	LNQA K GWLRDPSASPGDAGEQAIR - MSA E INEIIR	K281	E240
31264	3251.77	5.03	ALA K QVATALQNLQTKTNR - RQGKGDSP E AR	K464	E458
31264	3251.77	4.50	ALA K QVATALQNLQTKTNR - RQGKG D SPEAR	K464	D455

#	Exp. MH⁺	Primary Score	Peptide Sequence		
31620	2516.36	3.44	ALASIDS**K**LNQAK - MSA**E**INEIIR	K295	E240
35256	3787.88	4.12	GILSGTSDLLLTFD**E**AEVR - GSSHHHHHHSS**G**LVPR	E128	G12
38115	3310.75	4.29	GILSGTSDLLLTFD**E**AEVR - AVANSRPA**K**AAVH	E147	K507
39508	3685.90	3.56	GQGSSPVAMQ**K**AQQVSQGLDVLTAK - MSA**E**INEIIR	K352	E240
40134	3129.61	2.19	NQGIEEALKNRNFTVEKMSAEINEIIR	E224	K236
40285	3184.71	4.81	AQQVSQGLDVLTA**K**VENAAR - SLG**E**ISALTSK	K366	E437
40296	2280.32	3.69	**E**LLPVLISAMK - LNQA**K**GWLR	E200	K281
40351	3255.72	4.00	AQQVSQGLDVLTA**K**VENAAR - MSA**E**INEIIR	K366	E240
40571	4536.51	4.44	G**D**SPEARALAKQVATALQNLQTK - SLGEISALTS**K**LADLRRQGK	D455	K444
40881	3704.92	4.32	**K**IDAAQNWLADPNGGPEGEEQIR - **E**LLPVLISAMK	K387	E200
40923	3100.63	3.94	AQQVSQGLDVLTAKV**E**NAAR - **K**LEAMTNSK	E368	K373
41473	3723.01	4.34	GQGSSPVAMQ**K**AQQVSQGLDVLTAK - **E**LLPVLISAMK	K352	E200
41482	4633.41	3.54	TI**E**SILEPVAQQISHLVIMHEEGEVDGK - **K**LEAMTNSKQSIAK	E28	K381
42201	3294.83	3.66	AQQVSQGLDVLTA**K**VENAAR - **E**LLPVLISAMK	K366	E200
#	Exp. MH⁺	Primary Score	Peptide Sequence	A524	
15424	3050.56	3.57	NYVT**E**TNADTIDKNHAIYEK - INN**K**LK	E554	K540
16531	3340.67	3.55	NYVTETNA**D**TIDKNHAIYEK - A**K**EVSSALSK	E558	K571

16865	3341.67	4.11	NYVT**E**TNADTIDKNHAIYEK - A**K**EVSSALSK	E554	K571
34602	4584.25	4.05	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - AA**K**DVTTSLSK	E590	K617
35202	4586.25	5.2	IDDTSAELLTDDISDLKNNN**D**ITAENNNIYK - AA**K**DVTTSLSK	D594	K617
35824	4583.26	3.43	IDDTSAELLTDDISDLKNNNDITA**EN**NNIYK - AA**K**DVTTSLSK	E608	K617
39340	4911.48	4.37	IDDTSAELLTD**D**ISDLKNNNDITAENNNIYK - A**K**EVSSALS**K**VLSK	D595	K579
39340	4911.48	4.06	IDDTSAELLTDDIS**D**LKNNNDITAENNNIYK - A**K**EVSSALS**K**VLSK	D598	K579
39340	4911.48	3.83	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - A**K**EVSSALS**K**VLSK	E590	K579
39730	3291.74	3.64	IDDTSAELLTD**D**ISDLK - A**K**EVSSALSKVLSK	D595	K571
40244	4655.32	4.35	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - DVTTSLS**K**VLK	D594	K625
40244	4655.32	4.35	IDDTSAELLTD**D**ISDLKNNNDITAENNNIYK - DVTTSLS**K**VLK	D595	K625

625   **Suppl. Table 2**. Cross-linked residues characterized in the D1D2:A524 complex (XL-amino

626   acids of each protein are bolded in the sequences).

#	Exp. MH$^+$	Primary Score	Peptide Sequence	A483	Hv D1D2
13237	3014.59	3.29	DIT**K**STTEHR - VGK**E**TVQTTEDQILKR	K530	E66
14671	2518.38	2.49	SKDIT**K** - VGK**E**TVQTTEDQILKR	K526	E66
14791	2556.44	2.35	INN**K**LK - VGK**E**TVQTTEDQILKR	K540	E66
17806	2621.29	3.10	**G**SPGIPGDTYLTR - QQ**E**LTHQEHR	G517	E181
20039	2571.40	3.08	L**K**VTDANIR **- E**TVQTTEDQILKR	K542	E66
20526	3651.79	3.37	NYVT**E**TNADTIDKNHAIYEK - NS**K**NQGIEEALK	E554	K219
20985	3469.67	3.01	NYVTETNA**D**TIDKNHAIYEK - NLGPGMT**K**MAK	D558	K170
22936	3468.66	2.62	NYVT**E**TNADTIDKNHAIYEK - NLGPGMT**K**MAK	E554	K170
23137	4011	3.29	NYVTETNADTI**D**KNHAIYEK - VG**K**ETVQTTEDQILK	D561	K59
25418	3725.82	3.27	NYVTETNADTID**K**NHAIYEK - ETVQTT**E**DQILK	K562	E66
25418	3725.82	3.72	NYVTETNADTID**K**NHAIYEK - **E**TVQTTEDQILK	K562	E60
27618	3161.66	3.21	**G**SPGIPGDTYLTR - VGKETVQTT**E**DQILKR	G517	E60
27618	3161.66	3.61	**G**SPGIPGDTYLTR - VGKETVQTTE**D**QILKR	G517	D67
28366	2918.61	3.18	EVSSALS**K**VLSK - VGK**E**TVQTTEDQILK -	K579	E66
29996	1977.96	3.06	**G**SPGIPGDTYLTR - MI**D**ER	G517	D176
36497	2772.45	3.78	**G**SPGIPGDTYLTR - AQQVSQGL**D**VLTAK	G517	D361
36615	4613.23	3.68	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - NLGPGMT**K**MAK	D594	K170
36632	2214.27	3.22	A**K**EVSSALSK **- E**LLPVLISAMK -	K571	E200
37536	3824.86	2.64	**G**SPGIPGDTYLTR - KI**D**AAQNWLADPNGGPEGEEQIR	G517	D389

#	Exp. MH⁺	Primary Score	Peptide Sequence		
40263	3413.78	3.79	GSPGIPG**D**TYLTR - AQQVSQGLDVLTA**K**VENAAR	D484	K366
40432	5992.94	5.1	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - GQGSSPVAMQKAQQVSQGLDVLTA**K**	E590	K362
40679	2708.42	3.25	S**K**DITK - GILSGTSDLLLTFD**E**AEVR	K526	E128
40994	3074.69	3.12	**K**VTNSLSNLISLIGTK - ETVQTT**E**DQILK	K498	E66
41187	4612.22	2.86	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - NLGPGMT**K**MAK	E590	K170
41816	2387.41	3.65	DVTTSLS**K**VLK - **E**LLPVLISAMK	K625	E200
43326	3137.65	3.41	AA**K**DVTTSLS**K** - GILSGTSDLLLTFD**E**AEVR	K625	E128
43537	3542.88	3.63	IDDTSAELLT**D**DISDLK - ALA**K**QVATALQNLQTK	D594	K464
43612	3604.91	3.34	VTNSLSNLISLIGT**K**SGTQER - ETVQTT**E**DQILK	K513	E66
45081	3350.71	3.48	**G**SPGIPGDTYLTR - GILSGTS**D**LLLTFDEAEVR	G517	D121
53474	5745.03	4.09	NIN**K**D - TIESILEPVAQQISHLVIMHEEG**E**VDGKAIPDLTAPV AAVQAAVSNLVR	K632	E31

#	Exp. MH⁺	Primary Score	Peptide Sequence	Hv D1D2	
17582	1670.89	3.51	VG**K**ETVQTT**E**DQILK	K59	E66
17582	1670.89	3.38	VG**K**ETVQTTE**D**QILK	K59	D67
26000	3322.71	4.03	LNQA**K**GWLRDPSASPGDAGEQAIR - ALASI**D**SK	K281	D374
30901	3169.64	3.79	GQGSSPVAMQ**K**AQQVSQGLDVLTAK - V**E**NAAR	K352	E368
31031	3695.87	3.81	LNQA**K**GWLRDPSASPGDAGEQAIR - MSA**E**INEIIR	K281	E240
35827	3665.8	3.75	**K**IDAAQNWLADPNGGPEGEEQIR - MSA**E**INEIIR	K387	E240

#	Exp. MH+	Primary Score	Peptide Sequence		
38215	3040.71	3.43	VGKETVQTTEDQILKR - ELLPVLISAMK	K59	K200
39734	3614.9	4.15	GQGSSPVAMQKAQQVSQGLDVLTAK - SLGEISALTSK	K352	E437
40218	3580.72	3.88	VLQLTSWDEDAWASKDTEAMK - MSAEINEIIR	K261	E240
40418	2881.63	3.77	QVATALQNLQTKTNR - ELLPVLISAMK	K476	E200
40559	3184.71	4.71	AQQVSQGLDVLTAKVENAAR - SLGEISALTSK	K366	E437
40560	2280.32	3.35	ELLPVLISAMK - LNQAKGWLR	E200	K281
40604	2553.46	3.78	ALASIDSKLNQAK - ELLPVLISAMK	K276	E200
41140	2892.69	3.97	ALAKQVATALQNLQTK - ELLPVLISAMK	K464	E200
41598	4173.16	5.49	GQGSSPVAMQKAQQVSQGLDVLTAKVENAAR - KLEAMTNSK	K366	E375
41598	4173.16	5.8	GQGSSPVAMQKAQQVSQGLDVLTAKVENAAR - KLEAMTNSK	E368	K373
41598	4173.16	4.98	GQGSSPVAMQKAQQVSQGLDVLTAK - VENAARKLEAMTNSK	D361	K373
41682	3726.02	3.87	GQGSSPVAMQKAQQVSQGLDVLTAK - ELLPVLISAMK	K352	E200
42387	3293.83	4.63	AQQVSQGLDVLTAKVENAAR - ELLPVLISAMK	K366	E200
42387	3293.83	3.3	AQQVSQGLDVLTAKVENAARK - LNQAKGWLR	E368	K281

#	Exp. MH+	Primary Score	Peptide Sequence	A483	
17398	3339.66	3.31	NYVTETNADTIDKNHAIYEK - AKEVSSALSK	D561	K571
17630	3341.68	3.44	NYVTETNADTIDKNHAIYEK - AKEVSSALSK	D558	K571
17630	3341.68	4.19	NYVTETNADTIDKNHAIYEK - AKEVSSALSK	E554	K571

20071	3141.53	3.53	NYVTETNADTID**K**NHAIYEK - AK**E**VSSALSK	K562	E572
20150	3349.69	4.44	NYVTETNA**D**TIDKNHAIYEK - L**K**VTDANIR	D558	K542
20150	3349.69	4.03	NYVTETNADTI**D**KNHAIYEK - L**K**VTDANIR	D561	K542
35163	4584.24	3.59	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - AA**K**DVTTSLSK	D594	K617
35163	4584.24	3.49	IDDTSAELLTD**D**ISDLKNNNDITAENNNIYK - AA**K**DVTTSLSK	D595	K617
35382	4586.24	3.35	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - AA**K**DVTTSLSK	E590	K617
35967	4584.24	4.42	IDDTSAELLTDDIS**D**LKNNNDITAENNNIYK - AA**K**DVTTSLSK	D598	K617
35967	4584.24	4.95	IDDTSAELLTDDISDLKNNNDITA**E**NNNIYK - AA**K**DVTTSLSK	E608	K617
35967	4584.24	4.52	IDDTSAELLTDDISDL**K**NNNDITAENNNIYK - AA**K**DVTTSLSK	K600	D618
35967	4584.24	4.22	IDDTSAELLTDDISDLKNNNDITAENNNIY**K** - AA**K**DVTTSLSK	K614	D618
35967	4584.24	4.94	IDDTSAELLTDDISDLKNNN**D**ITAENNNIYK - AA**K**DVTTSLSK	D604	K617
39881	2845.52	4.73	VTNSLSNLISLIGT**K**SGTQER - **E**LQEK	K513	E520
39881	2845.52	4.62	VTNSLSNLISLIGT**K**SGTQER - ELQ**E**K	K513	E523
40044	4655.33	5.07	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - DVTTSLSKVL**K**	D594	K625
40044	4655.33	4.86	IDDTSAELLTD**D**ISDLKNNNDITAENNNIYK - DVTTSLSKVL**K**	D595	K625

40044	4655.33	4.45	IDDTSAELLTDDIS**D**LKNNNDITAENNNIYK - DVTTSLSKVL**K**	D598	K625
40044	4655.33	4.44	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - DVTTSLSKVL**K**	E590	K625
40065	3092.6	3.51	IDDTSAELLT**D**DISDLK - EVSSALSKVLS**K**	D594	K579
40065	3092.6	3.38	IDDTSAELLTD**D**ISDLK - EVSSALSKVLS**K**	D594	K579
40069	3093.61	3.36	IDDTSAELLTDDIS**D**LK - EVSSALSKVLS**K**	D598	K579
40095	2988.61	4.48	VTNSLSNLISLIGT**K**SGTQER - VT**D**ANIR	K513	D545
40448	3252.74	3.81	VTNSLSNLISLIGT**K**SGTQER - ETIF**E**ASKK	K513	E494
40603	5153.62	4.36	IDDTSAELLT**D**DISDLKNNNDITAENNNIYK - **K**VTNSLSNLISLIGTK	D594	K498
40603	5153.62	4.46	IDDTSAELLTD**D**ISDLKNNNDITAENNNIYK - **K**VTNSLSNLISLIGTK	D595	K498
40603	5153.62	4.56	IDDTSAELLTDDIS**D**LKNNNDITAENNNIYK - **K**VTNSLSNLISLIGTK	D598	K498
40603	5153.62	3.99	IDDTSAELLTDDISDLKNNN**D**ITAENNNIYK - **K**VTNSLSNLISLIGTK	D604	K498
40606	5153.62	3.53	IDDTSA**E**LLTDDISDLKNNNDITAENNNIYK - **K**VTNSLSNLISLIGTK	E590	K498
41116	2593.44	3.85	**K**VTNSLSNLISLIGTK - **E**TIFEASK	K498	E490
41142	3909.09	4	**E**TIFEASKKVTNSLSNLISLIGTK - **G**SPGIPGDTYLTR	E490	G477

638   **Suppl. Table 3**. Cross-linked residues characterized in the D1D2 -A483 1:1 complex (XL-amino

639   acids of each protein are bolded in the sequences)

Suppl. movie 1. TIRF analysis of HV-mCherry expressing C2.7 cells co-transfected with a GFP fusion to

the indicated construct. The time is indicated in seconds.

Suppl. movie 2. TIRF analysis of HV-mCherry expressing C2.7 cells co-transfected with a GFP fusion to

the indicated construct. The time is indicated in seconds. At time "0", addition of the Rho kinase inhibitor

Y-27632 was added at 100 μM final concentration.

Suppl. movie3. TIRF analysis of VASP-mCherry expressing C2.7 cells co-transfected with a GFP fusion to

the indicated construct. The time is indicated in seconds. At time "0", addition of the Rho kinase inhibitor

Y-27632 was added at 100 μM final concentration.

Suppl. Movie 4. 1205Lu melanoma cells 1205Lu melanoma cells were transfected with the indicated

constructs. Cells were perfused in a microfluidic chamber and allowed to adhere for 20 min prior to

application of shear stress reaching 22.2 dynes.cm^{-2}. The elapsed time is indicated in seconds.