1

To be submitted to: *Scientific Reports*

Running title: Plastid phylogenomics of the orchid family

Plastid phylogenomics resolves ambiguous relationships within the orchid

family and provides a solid timeframe for biogeography and macroevolution

Maria Alejandra Serna-Sánchez[1,2+], Oscar A. Pérez-Escobar[3*+], Diego Bogarín[4,5+], María Fernanda

Torres[6], Astrid Catalina Alvarez-Yela[7], Juliana E. Arcila[1], Climbie F. Hall[8], Fábio de Barros[8],

Fábio Pinheiro[9], Steven Dodsworth[10], Mark W. Chase[3], Alexandre Antonelli[3,6,11], Tatiana

Arias[1,7,12*]

[1] Laboratorio de Biología Comparativa. Corporación para Investigaciones Biológicas (CIB), Cra. 72 A No.

78 B 141, Medellín, Colombia.

[2] Biodiversity, Evolution and Conservation. EAFIT University, Cra. 49, No. 7 sur 50, Medellín,

Colombia.

[3] Royal Botanic Gardens, Kew, TW9 3AE, London, UK.

[4] Jardín Botánico Lankester, Universidad de Costa Rica, P. O. Box 302-7050, Cartago, Costa Rica.

[5] Naturalis Biodiversity Center, Endless Forms group, P.O. Box 9517, 2300 RA Leiden, The

Netherlands; Herbario UCH, Universidad Autónoma de Chiriquí, David, Panamá.

[6] Gothenburg Global Biodiversity Centre, Department of Biological and Environmental Sciences,

University of Gothenburg, 405 30 Gothenburg, Sweden.

[7] Centro de Bioinformática y Biología Computacional (BIOS). Ecoparque Los Yarumos Edificio BIOS,

Manizales, Colombia.

[8] Instituto de Botânica, Núcleo de Pesquisa Orquídario do Estado, 68041, São Paulo.

[9] Universidade Estadual de Campinas, Instituto de Biologia, Departamento de Biologia Vegetal, 13083-862,

Campinas-SP, Brazil

[10] School of Life Sciences, University of Bedfordshire, University Square, Luton, LU1 3JU, UK.

[11] Department of Plant Sciences, University of Oxford, South Parks Road, OX1 3RB Oxford, United

Kingdom

[12] Tecnológico de Antioquia, Calle 78B NO. 72A - 220 Medellín- Colombia

[*] Corresponding Authors: T.A. (tatiana.arias48@tdea.edu.co) & O.A.P.E. (o.perez-escobar@kew.org)

[+] These authors contributed equally to the study

All data have been deposited in Bioproject (XXXXXXX) and SRA (XXXXXXX).

ABSTRACT [291 words]

Recent phylogenomic analyses based on the maternally inherited plastid organelle have enlightened evolutionary relationships between the subfamilies of Orchidaceae and most of the tribes. However, uncertainty remains within several subtribes and genera for which phylogenetic relationships have not ever been tested in a phylogenomic context. To address these knowledge-gaps, we here provide the most extensively sampled analysis of the orchid family to date, based on 78 plastid coding genes representing 264 species, 117 genera, 18 tribes and 28 subtribes. Divergence times are also provided as inferred from strict and relaxed molecular clocks and birth-death tree models. Our taxon sampling includes 51 newly sequenced plastid genomes produced by a genome skimming approach. We focus our sampling efforts on previously unplaced clades within tribes Cymbidieae and Epidendreae. Our results confirmed phylogenetic relationships in Orchidaceae as recovered in previous studies, most of which were recovered with maximum support (209 of the 262 tree nodes). We provide for the first time a clear phylogenetic placement for Codonorchideae within subfamily Orchidoideae, and Podochilieae and Collabieae within subfamily Epidendroideae. We also identify relationships that have been persistently problematic across multiple studies, regardless of the different details of sampling and genomic datasets used for phylogenetic reconstructions. Our study provides an expanded, robust temporal phylogenomic framework of the Orchidaceae that paves the way for biogeographical and macroevolutionary studies.

Key words: Cymbidieae, High-throughput sequencing, Orchidaceae, Orchidoideae, Phylogenomics, Whole Plastid Genome

1. Introduction

Orchidaceae, with *ca*. 25,000 species and ~800 genera[1,2] are one of two of the most diverse and widely distributed flowering plant families on Earth and have captivated the attention of scientists for centuries[3]. The family has a striking morphological and ecological diversity and

evolved complicated interactions with fungi, animal and other plants[4,5] and a diverse array of sexual systems[6–8]. Numerous efforts have been made to understand the natural history, evolution and phylogenetic relationships within the family[2,7,9–13]. To date, there are seven nuclear genome sequences available, i.e., *Apostasia shenzhenica*[14], *Dendrobium catenatum*[15], *D. officinale*[16], *Gastrodia elata*[17], *Phalaenopsis equestris*[18], a *Phalaenopsis* hybrid cultivar[19], *P. aphrodite*[20], *Vanilla planifolia*[21], 221 complete plastid genomes and 2,678 sequence read archives for Orchidaceae in NCBI (accessed 22 August 2020) .

Phylogenomic approaches have been implemented to infer relationships between major orchids clades in deep and recent time[2,10,12,13,22,23], but extensive uncertainties remain regarding the phylogenetic placement of several subtribes. This knowledge-gap stems from the large gaps in both taxonomic and genomic sampling efforts that would be required to comprehensively cover all major orchid clades (subtribes/groups of genera). Givnish et al.[2] published the first well-supported analysis of Orchidaceae based on plastid phylogenomics. They performed a maximum likelihood (ML) analysis of 75 genes from the plastid genome of 39 orchid species, covering 22 subtribes, 18 tribes and five subfamilies. This robust but taxonomically under-sampled study agreed corroborated relationships of the subfamilies and tribes, observed in previous studies[10–13].

Previous orchid studies have failed to resolve relationships in rapidly diversifying clades[24–26] because of reduced taxon and data sampling[27]. This is particularly true for Cymbidieae and Pleurothallidinae, the two most species-rich groups in which generic relationships are largely the product of rapid diversification[28] that is difficult to resolve using only a few loci[25,29]. Cymbidieae comprise 10 subtribes, ~145 genera and nearly 3,800 species[1], 90% of which occur in the Neotropics [28]. Four of these subtribes are among the most species-rich in the Andean and Chocoan region (Maxillariinae, Oncidiinae, Stanhopeinae and Zygopetaliinae[30,31]). Pleurothallidinae include ~ 5,500 exclusively Neotropical species in 47 genera. Pleurothallid orchids are one of the most prominent components of the cloud forest flora in the northern and central Andes and Central America[32].

Another group in which phylogenetic relationships are unresolved is Orchidoideae [1,33]. This group comprises four mostly terrestrial tribes, 25 subtribes and over 3,600 species. The subfamily occurs on all continents except the Antarctic. Previous efforts to disentangle the phylogenetic relationships in the subfamily have mostly relied on a small set of nuclear and plastid markers[34], and more recently on extensive plastid coding sequence data[2].

The wide geographical range of these groups in the tropics and temperate regions and their

striking vegetative and reproductive morphological variability make them ideal model clades for

disentangling the contribution of abiotic and biotic drivers of orchid diversification across biomes.

Occurring from alpine ecosystems to grasslands, they have conquered virtually all ecosystems

available in any elevational gradient[35–37], showing independent transitions to terrestrial, rupicolous

and epiphytic habit. Moreover, they have evolved a diverse array of pollination systems[38–40],

rewarding species offering scent, oil and nectar, and even food- and sexual deceptive species[41,42].

However, the absence of a solid phylogenetic framework has precluded the study of how such

systems evolved and the diversification dynamics of Cymbidieae, Pleurothallidinae and

Orchidoideae more broadly.

Phylogenetic analyses are crucial to understanding the drivers of diversification in orchids,

including the mode and tempo of morphological evolution[30,43]. High-throughput sequencing and

modern comparative methods have enabled the production of massive molecular datasets to

reconstruct evolutionary histories and thus provide unrivalled knowledge on plant phylogenetics[44].

Here, we present the most densely sampled plastid analysis of Orchidaceae, including data from 51

newly sequenced plastid genomes,. We apply two general approaches: a) maximum likelihood

phylogenetic analysis conducted on 78 plastid coding regions to inform relationships; b) Bayesian

inference in combination with strict and relaxed molecular clocks and a birth-death model applied

to a subset of the plastid coding regions to produce a temporal framework of the orchid family. Our

study expands the current generic representation for the Orchidaceae and clarifies previously

unresolved phylogenetic relations within the Cymbidieae, Pleurothallidinae and Orchidoideae. The

results reported here provide a robust framework for the orchid family and new insights into

relationships at both deep and shallow phylogenetic levels.

2. Results

2.1 Phylogenetic relationships and divergence times in the orchid family

The ML tree derived from the 78 plastid genes is provided in Fig. 1. Two hundred-and-thirty-one nodes were recovered as strongly supported (i.e. likelihood bootstrap percentage [LBP] = 85-100), of which 209 attained maximum support. Only 26 nodes recovered LBPs between 25 and 84 (Fig. 1, inset). Unsupported relationships were restricted to Epidendroideae and Orchidoideae but were more frequent in Epidendroideae and often linked to low levels of sequence variation. Here, poorly supported relationships occurred mostly towards the backbone of the tribes Arethuseae, Cymbidieae, Epidendreae and Neottieae and Tropidieae + Nervilieae and the most recent common ancestor (MRCA) of Arethuseae, Malaxideae, Podochilieae, Collabieae, Epidendreae, Vandeae and Cymbidieae. Intrageneric relationships were robustly supported, with only two instances for which few nodes were recovered as poorly supported (*Dendrobium*: 3; *Cymbidium:* 1; Fig. S1).

Absolute times of divergence under strict and relaxed clocks for Orchidaceae, subfamilies and most tribes are provided in Table 1 (phylogenetic trees with mean ages and intervals of confidence produced under both clock models are provided on Figs S2, S3). Strict and relaxed molecular clocks revealed similar ages of divergence for the majority of the MRCAs of main orchid clades, although we found stark differences in the length of the 95% highest posterior density intervals (HPD) derived from both models are obvious, with the relaxed clock producing larger HPDs (Tab. 1; Fig. S2, S3; Tab. S1, S2). Under the strict and relaxed clocks, Orchidaceae diversified first during the late Cretaceous (88.1 my ± 3; 89.1 my ±9, respectively). The largest differences on the MRCA ages occurred in Epidendroideae (44 my ± 2 vs 60 my ± 10 under a strict and relaxed clock models, respectively) and Vanilloideae (80 my ± 4 vs 67 my ± 9). A complete account of mean and median ages, HPDs, branch lengths and rate values estimated for all nodes of chronograms estimated strict and relaxed molecular clock models are provided on Tab. S1, S2.

2.2. Phylogenetic informativeness of plastid genes

Phylogenetic informativeness plots are provided on Fig. S4 (see Tab. S3,S4 for a detailed account of PI per-site and net values for each assessed locus). Per-site and net phylogenetic informativeness (PI) analyses recovered both *ycf*1 as the most informative locus, which attained the highest values at a reference time (phylogenetic depth) of 0.51. On average, plastid loci attained their highest PI value at a reference time of 0.85 (SD=0.16). In contrast, the highest PI values of

the 10 most informative loci occurred at an average reference time of 0.63 (SD=0.11) and 0.80

(SD=0.17) for per-site and net PI calculations.

3. Discussion

3.1 A robust temporal phylogenomic framework for the orchid family

Previous phylogenomic studies of the orchid family included up to 74 species representing

18 tribes, 19 subtribes and 66 genera[27]. Our study sampled 264 species from all subfamilies,

representing 18 tribes (out of 22), 28 subtribes (out of 46) and 74 genera (~10% of the currently

recognised genera; Fig. 2). In general, our phylogenomic frameworks are in agreement with

previously published family-wide orchid analyses either inferred from dozens of markers[2,13] or

from a handful of loci[29]. Here, representativeness of Cymbidieae and Epidendreae, two of the most

prominent tropical Epidendroideae[45]clades, have increased from eight to 32 genera and six to 30,

respectively[2,27]. In particular, relationships inferred from extensive plastid data within

Zygopetaliinae (Cymbidieae) and Pleurothallidinae (Epidendreae) are presented for the first time.

Our 78-coding sequence plastid ML analysis led to similar results as reported by Givnish et al.[2],

Niu et al.[13] and Li et al.[27] but with an overall clear increase in support: 22% of nodes with LBS <

85 in Givnish et al.[2] and 21% in Li et al.[27] *vs* 11.5% in this study. This is particularly evident in

relationships inferred within Orchidoideae (see section 3.5 of *Discussion*) and Cymbidieae,

Epidendreae (see sections 3.3 and 3.4 of *Discussion,* respectively) and Collabieae. For the last, for

the first time we provide high support for the previously unresolved relationship of

Podochilieae+Collabieae[2,27].

The absolute age estimates derived from our strict and relaxed molecular clocks and five of

the most informative plastid loci are in line with previous nuclear-plastid multi-locus and

phylogenomic plastid-only chronograms[2,46,47]. Nonetheless, our ML tree also identifies intricate

relationships that have been consistently recovered as unsupported in several studies. These include

unsupported basal nodes in Epidendroideae representing Sobralieae, Nervilieae and Tiphoreae[27,48],

Arundina+remainder of Arethuseae[27], and the position of Eulophiinae in the Cymbidieae[25,28,49]

(Fig. 2). Uncertainty around the phylogenetic position of these clades might be due to limited taxon

sampling in this and previous studies. Alternatively, intragenomic conflict[50–52] and lack of

phylogenetic informativeness required to sort out relationships derived from rapid

diversifications[22,53,54] in plastid DNA sequences (regardless of whether whole plastid genome

datasets are employed[55]) might hamper the phylogenetic placement of clades with robust support.

3.3 Improved support of phylogenetic relationships within Cymbidieae

Multiple studies have inferred evolutionary relationships in Cymbidieae from morphological and molecular characters[28,29]. Relationships among subtribes have recently been estimated using the plastid genes *psaB, rbcL, matK* and *ycf1* combined with the low-copy nuclear gene *Xdh*[25]. Here, Cymbidiinae was sister to the remainder of Cymbidieae. Poorly supported and incongruent relationships were found among Catasetinae, Eulophiinae and Eriopsidinae, however, when compared with the topologies obtained by Whitten et al.[29], Freudenstein & Chase[48] and Pérez-Escobar et al.[7]

The most complete taxonomic sampling conducted to date under a plastid phylogenomic framework[2] included 8 of 11 subtribes of Cymbidieae, but some inter-subtribal relationships were unresolved: Stanhopeinae (20 genera), Maxillariinae (12 genera), Zygopetalinae (36 genera), Oncidiinae (65 genera) and Eulophiinae (13 genera). A clade formed by Stanhopeinae and Maxillariinae had poor support (LBP=62) and their relationship to Zygopetaliinae also had low support (LBP=72). The relationship between Eulophiinae and a clade of Stanhopeinae, Maxillariinae, Zygopetalinae and Oncidiinae also had poor support (LBP=42). One of the outcomes of our expanded sampling (nine subtribes) is the improvement of support in Cymbidieae, more specifically for nodes of some groups involved in rapid diversifications that historically have been problematic to resolve[2,29]. In particular, Maxillariinae+Stanhopeinae and Catasetinae+Cyrtopodiinae are now both strongly supported (LBP=100). In addition, our results also support the placement of *Dipodium* (Dipodiinae) is supported as sister to the rest of Cymbidieae, a relationship which was previously recovered from a few loci[25]. However, our plastid phylogenomic framework is still incomplete due to absence of representatives of Eriopsidiinae and Coleopsidinae.

One other novelty of our study is the inference of relationships in Zygopetalinae, a subtribe in which relationships have previously been poorly understood[56]. The most extensively sampled analysis of Zygopetalinae inferred from plastid markers (*mat*K-*ycf1*)[29] included 60 species and 27 genera, but relationships between most genera attained only low support. Our expanded molecular, but taxonomically reduced, matrix (i.e. 20 genera and 21 species) produced greater support for the backbone relationships in the subtribe, including the radiation of the *Huntleya* clade (*Dichaea, Huntleya, Chaubardia* and the *Chondrorhyncha* complex[56,57]). Nonetheless, relationships between the *Huntleya* grade (i.e. *Huntleya* clade + *Cryptarrhena*) and the remainder of Zygopetalinae still

remains unresolved.

Our phylogenetic analyses further place for the first time in the orchid tree of life

Cheiradenia and *Hoehneella* with moderate to strong support (Fig. 1, S1). *Cheiradenia* is a

monospecific genus restricted to the lowland wet forests of Venezuela and Guyana whereas

Hoehneella includes two species exclusively distributed in the Brazilian evergreen wet forests of

the Brazilian states of Espirito Santo and São Paulo[58]. Referring to the similarity of both vegetative

and floral reproductive characters, Pupulin [58] hypothesised that *Cheiradenia* should be closely

related to members of the *Zygopetalum* clade (e.g. *Koellensteina*, *Paradisanthus*), with *Hoehneella*

being related to the *Huntleya* clade (i.e. *Huntleya* and *Chaubardia*). Our ML tree supports both

assumptions, placing *Cheiradenia* as sister to *Paradisanthus* with maximum support and

Hoehneella as sister to *Chaubardia* in a moderately supported clade (83 LBP: Fig. 1, S1).

Koellensteina kellneriana (the taxonomic type of the genus) clustered with *Acacallis* and not with

Otostylis and *Paradisanthus*, and therefore we confirm that *Koellensteina* in the strict sense is

related to *Acacallis*. In addition, *Otostylis* is recovered as sister to *Warrea* and not to

Paradisanthus as previously suggested by Williams et al.[56] based on a weakly supported

placement. Our results also highlight the extensive and independent terrestrial and epiphytic habit

transitions occurring in this clade, as most sister genera shows different habit types.

3.5. Novel and robust relationships in the most rapidly diversifying subtribe Pleurothallidinae

One of the most spectacular Neotropical plant diversifications is perhaps that of the

Pleurothallidinae, for it involves the evolution of ~5,000 species that have conquered virtually all

biogeographical regions in the American tropics[32,45]. The rapid radiation of Pleurothallidinae

occurring in the last ~20 Myrs[28] is associated with the evolution of a diverse suite of pollination

systems ranging from food deception[59] to pseudocopulation[60] linked to dipterans[61,62] and a complex

array of reproductive and vegetative morphologies[22,32]. Understanding of relationships in the

subtribe has relied mostly on relatively small number of markers[63–65], which have informed with

some confidence the phylogenetic placement and monophyly of genera in Pleurothallidinae, yet

basal nodes in these trees have often lacked good support.

Several attempts have been conducted to estimate generic relationships in the subtribe,

most of which have relied on nuclear rITS and plastid *matK* markers[66]. A synthesis of the

phylogenetic relationships in the subtribe based on such studies was conducted by Karremans[67].

Here, a cladogram depicting the commonest topologies of relationships between genera was

provided and nine clades were defined (termed "affinities" by the author) but without considering

the magnitude of the support for these (see Figure 2 in Karremans[67]). Our plastid phylogenomic

analysis recovered well-supported relationships in Pleurothallidinae that are mostly in line with

previously published studies [28,63,68]. However, these previous trees based on a handful of DNA

nuclear and plastid markers yielded poor resolution and low support for backbone nodes as well as

infrageneric relationships. In contrast, our plastid phylogenomic inferences recovered high support

along the backbone, thus recovering novel placements. Some of these noteworthy well-supported

relationships are the position of *Acianthera* as sister to *Myoxanthus* and *Dresslerella* as sister to

Barbosella+Restrepia (Fig. 1, S1).

Acianthera includes over 300 species distributed throughout the American tropics and

subtropics[64,69,70], is often retrieved as sister to the remainder of Pleurothallidinae with moderate

support[68]. Karremans[67] used a series of "affinities" to describe to groups of genera affiliated with a

core genus of these group and thus described the "*Acianthera* affinity" as the frequent clustering of

several Central American genera with *Acianthera*[64]. Our study contradicts Karreman's[67] concept of

the *Acianthera* affinity by placing with high support *Acianthera* in the *Restrepia* affinity as sister to

Myoxanthus. Dresslerella was previously recovered with low support as sister to the remaining

genera in the *Restrepia* affinity (*Barbosella*, *Echinosepala*, *Myoxanthus*, *Restrepia*, *Restrepiella*

and *Restrepiopsis*). In contrast, our analysis robustly places *Dreslerella* as sister to *Restrepia* and

Barbosella, a result that does not support the monophyly of the *Restrepia* affinity.

Although estimates of the ancestral distribution of the Pleurothallidinae are still uncertain,

most of the early divergent Pleurothallidinae and their sister groups are found in the Antilles or

Brazil[28]. The remarkable relationship recovered here for *Acianthera+Myoxanthus* could yield more

clues about the biogeographic history and evolution of the subtribe because Brazil harbours a high

species diversity of *Acianthera* and some of the early divergent clades in *Myoxanthus* (particularly

the species close to *M. lonchophyllus*), whereas *Myoxanthus* is notably absent in the Antilles. In

addition, other early divergent clades such as *Octomeria* and *Barbosella* are more diverse in Brazil.

These early diverging clades share the lack of stem annulus as a morphological symplesiomorphy,

a character that later appears in more diverse clades such as *Masdevallia+Dracula*, *Lepanthes*, and

Pleurothallis+Stelis[71]. Members of these clades probably diversified after a migration to the

mountainous areas of the northern Andes ca 16 ± 5 Ma and together account for almost 80% of the

species in the subtribe[28]. The modern range extends mostly along the Andean and Central

American mountain ranges. Here, another noteworthy relationship is that the less diverse

Specklinia clade (*Scaphosepalum+Platystele*) was recovered as sister to the most species-rich

clades of the subtribe (*Masdevallia*, *Lepanthes*, and *Pleurothallis*). In previous phylogenetic

analyses *Specklinia* clade was recovered as sister to just *Pleurothallis*[28].

Likewise, relationships between early divergent members in the *Lepanthes* affinity

(*Anathallis*, *Draconanthes*, *Epibator*, *Lepanthes*, *Opilionanthe*, *Trichosalpinx* and *Tubella*) were

largely weakly supported, demonstrating the need for increased taxon sampling, principally in

Lepanthopsis and *Tubella*[32]. In particular, the early diversification of the *Lepanthes* affinity (>1500

spp.), inferred to have occurred around 8 Ma, has been linked to colonisation of newly formed

environments in the Andean Cordillera, a product of accelerated mountain uplift and specific

pollination systems (pseudocopulation and food mimicry[60]).

Another novel placement concerns *Teagueia* (diverse in Colombia, Ecuador and Peru[72–74]),

which resembles *Platystele*[75]. Karremans[76] had suggested a close relationship between *Teagueia*

and *Scaphosepalum*, but our results place *Teagueia* as sister to *Platystele* with high support, thus

corroborating the long-standing hypotheses of their sister relationship based on the of their

reproductive structures[74,75].

3.5. Evolutionary relationships in Orchidoideae

Our study provides a well-supported tree for Orchidoideae. Our ML inference supports the

findings of Pridgeon et al.[35] in which Diurideae is sister to Cranichideae and Codonorchideae to

Orchideae. Our findings differ from Givnish et al.[2] and Salazar et al.[34], in which

Diurideae/Cranichideae are sister to Codonorchideae, with Orchideae sister to all these (Fig. 2).

Givnish et al.[2] included all four tribes but only six of 21 subtribes of Orchidoideae, and the

relationship of Diurideae to Cranichideae was poorly supported.

Conclusions

This study presents a well-resolved, more densely sampled and strongly supported analysis of

Orchidaceae and their absolute times of divergence than all previous such studies. For deep

branches and recent diversifications in Cymbidieae and Epidendreae, support is improved, yet

several recalcitrant nodes that historically have been challenging to resolve were also found (e.g.

early divergent taxa in the Epidendroideae, initial radiation of the *Lepanthes* affinity in

Pleurothallidinae). Similarly, our analyses provide the a well-supported result for Orchidoideae.

Although taxon sampling was sufficient to resolve the relationships between the major clades in

the family, sampling of unrepresented genera and representatives of Eriopsidiinae, Goodyerinae,

and Coleopsidinae would further enhance our understanding of phylogenetic relationships.

Material and methods

2.1 Sampling, DNA extraction and sequencing

Two-hundred and sixty-four species representing 117 genera, 28 subtribes and 18 tribes were sampled in this study. For 51 species plastid genomes were sequenced. Table S5 provides accession numbers of plastid genomes sourced from NCBI and GenBank numbers of those newly generated. Fresh leaves were stored in silica gel for subsequent DNA extraction using a CTAB method[77]. Total DNA was purified with silica columns and then eluted in Tris-EDTA[78]. DNA samples were adjusted to 50 ng/uL and sheared to fragments of approximately 500 bp.

High-throughput sequencing

The library preparation, barcoding and sequencing (Illumina HiSeqX) were conducted at Rapid Genomics LLC (Gainesville, FL, USA) and Genewiz GmbH (Leipzig, Germany). Pair-end reads of 150 bp were obtained for fragments with insert size of 300-600 bp. Overhangs were blunt ended using T4 DNA polymerase, Klenow fragment and T4 polynucleotide kinase. Subsequently, a base 'A' was added to the 3' end of the phosphorylated blunt DNA fragments. DNA fragments were ligated to adapters, which have a thymine (T) overhang. Ligation products were gel-purified by electrophoresis to remove all unbound adapters or split adapters that were ligated together. Ligation products were then selectively enriched and amplified by PCR. For each sample, between one and 10 million paired-end reads were generated.

Plastid genome assembly and annotation

Raw sequences were quality filtered using Trimmomatic[79] in order to eliminate sequencing artefacts, improve uniformity in the read length (>40 bp) and ensure quality (>20) for further analysis. Filtered sequences were processed with BBNorm[80] to normalize coverage by down-sampling reads over high-depth areas of the genomes (maximum depth coverage 900x and minimum depth 6x). This step creates a flat coverage distribution in order to improve read assembly. Subsequently, overlapping reads were merged into single reads using BBmerge[81] in order to accelerate the assembly process. Overlapping of paired reads was evaluated with Flash[82] to reduce redundancy. Merged reads were used to carry out the whole genome de novo assembly with SPAdes (hash length 33,55,77)[83].

To produce contiguous, linear plastid genome sequences we relied on a refence-based and

de-novo approaches. The reference based approach was conducted on MIRA v. 4[84], a software that

maps read data against a consensus sequence of a reference assembly (simple mapping). MIRA has

been useful for assembling complicated genomes with many repetitive sequences[85–87]. MIRA

produces BAM files as output, which were subsequently used to generate consensus sequences in

SAMTOOLS[88]. We sourced 11 reference plastomes from the NCBI repository that represent

related species, namely: *Cattleya crispata, Goodyera fumata, Masdevallia picturata*, *M. coccinea*,

Oncidium sphacelatum and *Sobralia callosa*. The *de-novo* assembly approach relied on

GetOrganelle[89], using the recommended default settings for assemblies of green-plant plastid

genomes.

Newly sequenced and datamined plastid genomes were annotated through the Chlorobox

portal of the Max Planck Institute[90]. Sequences were uploaded as fasta files, and running

parameters were established as follow: BLAST protein search identity=65%, BLAST rRNA,

tRNA, DNA search identity=85%, genetic code = bacterial/plant plastid, max intron length=3,000,

options= allow overlaps. *Apostasia wallichii, Masdevallia picturata, Oncidium sphacelatum,*

Sobralia callosa and *Goodyera fumata* were set as the 'Server Reference' and *Cattleya liliputana*

was set as the 'Custom Reference' for CDS and tRNA, rRNA, primer, other DNA or RNA

specifications.

Phylogenetic analysis

A set of 78 plastid genes was used to reconstruct phylogenetic relationships in Orchidaceae.

These were aligned[91] using MAFFT 7[92] and subsequently concatenated (proportions of missing

data per species is provided on Tab. S5). This step was performed at the supercomputing centre

APOLO, EAFIT University, Medellín, Colombia. Phylogenetic reconstruction based on maximum

likelihood (ML) was implemented in RAxML v. 8.0[93], using 1,000 bootstrap replicates and the

GTR+GAMMA model. Absolute age estimation analyses relied on fossil and secondary calibration

points, strict and molecular clocks and a birth/death model implemented in BEAST v. 1.8[94]. The

fossil constraint was added to the MRCA of *Dendrobium* following Xiang et al.[95] using a normal

distribution with mean value of 21.07 and a standard deviation (SD) of 3.0. Following Givnish et

al.[2], the two secondary calibration points were added to the root of the tree and MRCA of the

Orchidaceae, using a normal distribution and mean values of 123.48 (SD=2.0) and 90 (SD=2.0).

Because dating analyses conducted on dozens of gene alignments and hundreds of terminals are

extremely computationally greedy, we estimated absolute ages on the five most phylogenetically

informative genes (see below) and by constraining the tree topology to the ML tree derived from

RAxML. For each clock model, we conducted two MCMC analyses with 250 million generations

each with a sampling frequency of 10000 generations. The convergence of the strict and relaxed

molecular clocks parameters was confirmed on the software TRACER v1.6.

(http://tree.bio.ed.ac.uk/software/tracer/). Maximum clade credibility trees were summarised from

the MCMC trees in the program TreeAnnotator v.1.8. of the software BEAST.

Phylogenetic informativeness profiles

To estimate the phylogenetic informativeness (PI) of plastid genes we calculated the per-

site and net values for each assessed locus with the HyPhy substitution rates algorithm for DNA

sequences[96] using in the web application PhyDesign http://phydesign. townsend.yale.edu/). The

input files were the consensus ML ultrametric tree converted with the function *chronos* of the R-

package APE (http://ape-package.ird.fr/) using an smoothing rate of 1 and a relaxed clock model,

and the partitioned concatenated gene alignments.

Acknowledgments

We would like to thank Esteban Urrea for helping with the bioinformatics pipelines. We thank

Norris Williams and the late Mark Whitten (University of Florida) for collecting and preparing the

specimens. Kurt Neubig from Southern Illinois University provided the sequences of 11 new

samples. We also thank Janice Valencia for critical feedback on the paper, Juan David Pineda

Cardenas for advising about computational resources used through EAFIT and Juan Carlos Correa

for computational advices at BIOS. The University of Costa Rica provided access to the genetic

material for the projects B8257 and B6140. Finally, we would like to thank IDEA WILD for

supporting with photographic equipment and Sociedad Colombiana de Orquideología for

supporting M. A. Serna-Sánchez with a grant to conduct her undergraduate studies.

O.A.P.E. is supported by the Swiss Orchid Foundation and the Lady Sainsbury Orchid Fellowship.

A.A. acknowledges financial support from the Swedish Research Council (2019-05191), the

Swedish Foundation for Strategic Research (FFL15-0196) and the Royal Botanic Gardens, Kew.

References

1. Chase, M. W. *et al.* An updated classification of Orchidaceae: Updated Classification of Orchidaceae. *Bot. J. Linn. Soc.* **177,** 151–174 (2015).

2. Givnish, T. J. *et al.* Orchid phylogenomics and multiple drivers of their extraordinary diversification. *Proc. R. Soc. B Biol. Sci.* **282,** 1553 (2015).

3. Darwin, Ch., *On the various contrivances by which british and foreign orchids are fertilised by insects, and on the good effects of intercrossing. London: John Murray. 1st ed., 1st issue.* (1862).

4. Fay, M. F. & Chase, M. W. Orchid biology: from Linnaeus via Darwin to the 21st century. *Ann. Bot.* **104,** 359–364 (2009).

5. Ramírez, S. R. *et al.* Asynchronous diversification in a specialized plant-pollinator mutualism. *Science* **333,** 1742–1746 (2011).

6. Borba, E. L., Barbosa, A. R., Melo, M. C. de, Gontijo, S. L. & Oliveira, H. O. de. Mating systems in the Pleurothallidinae (Orchidaceae): evolutionary and systematic implications. *Lankesteriana* (2011). doi:10.15517/lank.v11i3.18275

7. Pérez-Escobar, O. A. *et al.* Multiple geographical origins of environmental sex determination enhanced the diversification of Darwin's favourite orchids. *Sci. Rep.* **7,** 12878 (2017).

8. Pérez-Escobar, O. A., Gottschling, M., Whitten, W. M., Salazar, G. & Gerlach, G. Sex and the Catasetinae (Darwin's favourite orchids). *Mol. Phylogenet. Evol.* **97,** 1–10 (2016).

9. Bateman, R. & Rudall, P. Evolutionary and Morphometric Implications of Morphological Variation Among Flowers Within an Inflorescence: A Case-Study Using European Orchids. *Ann. Bot.* **98,** 975–93 (2006).

10. Dong, W.-L. *et al.* Molecular Evolution of Chloroplast Genomes of Orchid Species: Insights into Phylogenetic Relationship and Adaptive Evolution. *Int. J. Mol. Sci.* **19,** 716 (2018).

11. Freudenstein, J. V. & Chase, M. W. Phylogenetic relationships in Epidendroideae (Orchidaceae), one of the great flowering plant radiations: progressive specialization and diversification. *Ann. Bot.* **115,** 665–681 (2015).

12. Luo, J. *et al.* Comparative Chloroplast Genomes of Photosynthetic Orchids: Insights into Evolution of the Orchidaceae and Development of Molecular Markers for Phylogenetic Applications. *PLoS One* **9,** e99016 (2014).

13. Niu, Z. *et al.* The Complete Plastome Sequences of Four Orchid Species: Insights into the

440   Evolution of the Orchidaceae and the Utility of Plastomic Mutational Hotspots. *Front. Plant*

441   *Sci.* **8,** (2017).

442 14. Zhang, G.-Q. *et al.* The *Apostasia* genome and the evolution of orchids. *Nature* **549,** 379–

443   383 (2017).

444 15. Zhang, G.-Q. *et al.* The *Dendrobium catenatum* Lindl. genome sequence provides insights

445   into polysaccharide synthase, floral development and adaptive evolution. *Sci. Rep.* **6,** 19029

446   (2016).

447 16. Yan, L. *et al.* The Genome of *Dendrobium officinale* Illuminates the Biology of the

448   Important Traditional Chinese Orchid Herb. *Mol. Plant* **8,** 922–934 (2015).

449 17. Yuan, Y. *et al.* The *Gastrodia elata* genome provides insights into plant adaptation to

450   heterotrophy. *Nat. Commun.* **9,** (2018).

451 18. Cai, J. *et al.* The genome sequence of the orchid *Phalaenopsis equestris*. *Nat. Genet.* **47,** 65–

452   72 (2014).

453 19. Huang, J.-Z. *et al.* The genome and transcriptome of *Phalaenopsis* yield insights into floral

454   organ development and flowering regulation. *PeerJ* **4,** e2017 (2016).

455 20. Chao, Y.-T. *et al.* Chromosome-level assembly, genetic and physical mapping of

456   *Phalaenopsis aphrodite* genome provides new insights into species adaptation and resources

457   for orchid breeding. *Plant Biotechnol. J.* **16,** 2027–2041 (2018).

458 21. Hu, Y. *et al.* Genomics-based diversity analysis of Vanilla species using a *Vanilla planifolia*

459   draft genome and Genotyping-By-Sequencing. *Sci. Rep.* **9,** 3416 (2019).

460 22. Bogarín, D. *et al.* Anchored Hybrid Enrichment generated nuclear, plastid and mitochondrial

461   markers resolve the *Lepanthes horrida* (Orchidaceae: Pleurothallidinae) species complex.

462   *Mol. Phylogenet. Evol.* (2018). doi:10.1016/j.ympev.2018.07.014

463 23. Pérez-Escobar, O. A. *et al.* Resolving relationships in an exceedingly young Neotropical

464   orchid lineage using Genotyping-by-sequencing data. *Mol. Phylogenet. Evol.* **144,** 106672

465   (2020).

466 24. Jin, W.-T. *et al.* Phylogenetics of subtribe Orchidinae s.l. (Orchidaceae; Orchidoideae) based

467   on seven markers (plastid matK, psaB, rbcL, trnL-F, trnH-psba, and nuclear nrITS, Xdh):

468   implications for generic delimitation. *BMC Plant Biol.* **17,** (2017).

469 25. Li, M.-H., Zhang, G.-Q., Liu, Z.-J. & Lan, S.-R. Subtribal relationships in Cymbidieae

470   (Epidendroideae, Orchidaceae) reveal a new subtribe, Dipodiinae, based on plastid and

471   nuclear coding DNA. *Phytotaxa* **246,** 37 (2016).

26.   Nauheimer, L., Schley, R. J., Clements, M. A., Micheneau, C. & Nargar, K. Australasian orchid biogeography at continental scale: Molecular phylogenetic insights from the Sun Orchids (*Thelymitra*, Orchidaceae). *Mol. Phylogenet. Evol.* **127,** 304–319 (2018).

27.   Li, Y.-X. *et al.* Phylogenomics of Orchidaceae based on plastid and mitochondrial genomes. *Mol. Phylogenet. Evol.* **139,** 106540 (2019).

28.   Pérez-Escobar, O. A. *et al.* Recent origin and rapid speciation of Neotropical orchids in the world's richest plant biodiversity hotspot. *New Phytol.* **215,** 891–905 (2017).

29.   Whitten, W. M., Neubig, K. M. & Williams, N. H. Generic and subtribal relationships in neotropical Cymbidieae (Orchidaceae) based on matK/ycf1 plastid data. *Lankesteriana* **13,** 375–392 (2014).

30.   Pridgeon, A. *Genera Orchidacearum Vol. 5, Vol. 5,.* (Oxford University Press, 2009).

31.   Pérez-Escobar, O. A. *et al.* The origin and diversification of the hyperdiverse flora in the Chocó biogeographic region. *Front. Plant Sci.* **10,** 1328 (2019).

32.   Bogarín, D. *et al.* Phylogenetic comparative methods improve the selection of characters for generic delimitations in a hyperdiverse Neotropical orchid clade. *Sci. Rep.* **9,** 1–17 (2019).

33.   Górniak, M., Paun, O. & Chase, M. W. Phylogenetic relationships within Orchidaceae based on a low-copy nuclear coding gene, Xdh: Congruence with organellar and nuclear ribosomal DNA results. *Mol. Phylogenet. Evol.* **56,** 784–795 (2010).

34.   Salazar, G. A., Chase, M. W., Soto Arenas, M. A. & Ingrouille, M. Phylogenetics of Cranichideae with emphasis on Spiranthinae (Orchidaceae, Orchidoideae): evidence from plastid and nuclear DNA sequences. *Am. J. Bot.* **90,** 777–795 (2003).

35.   Bone, R. E., Cribb, P. J. & Buerki, S. Phylogenetics of Eulophiinae (Orchidaceae: Epidendroideae): evolutionary patterns and implications for generic delimitation: Evolutionary patterns in Eulophiinae. *Bot. J. Linn. Soc.* **179,** 43–56 (2015).

36.   Pérez-Escobar, O. A. *et al.* Andean mountain building did not preclude dispersal of lowland epiphytic orchids in the Neotropics. *Sci. Rep.* **7,** 4919 (2017).

37.   Salazar, G. A. *et al.* Phylogenetic systematics of subtribe spiranthinae (Orchidaceae: Orchidoideae: Cranichideae) based on nuclear and plastid DNA sequences of a nearly complete generic sample. *Bot. J. Linn. Soc.* **186,** 273–303 (2018).

38.   Martins, A. *et al.* From tree tops to the ground: Reversals to terrestrial habit in *Galeandra* orchids (Epidendroideae: Catasetinae). *Mol. Phylogenet. Evol.* **127,** (2018).

39.   Nunes, C. *et al.* More than euglossines: the diverse pollinators and floral scents of

504    Zygopetalinae orchids. *Sci. Nat.* **104,** (2017).

505 40. Pansarin, L., Pansarin, E., Gerlach, G. & Sazima, M. The Natural History of *Cirrhaea* and

506    the Pollination System of Stanhopeinae (Orchidaceae). *Int. J. Plant Sci.* 000–000 (2018).

507    doi:10.1086/697997

508 41. Cisternas, M. A. *et al.* Phylogenetic analysis of Chloraeinae (Orchidaceae) based on plastid

509    and nuclear DNA sequences. *Botanical Journal of the Linnean Society* (2012).

510 42. Ramirez, S. R., Roubik, D. W., Skov, C. & Pierce, N. E. Phylogeny, diversification patterns

511    and historical biogeography of euglossine orchid bees (Hymenoptera: Apidae). *Biol. J. Linn.*

512    *Soc.* **100,** 552–572 (2010).

513 43. Cingel, N. A. Van Der. *An Atlas of Orchid Pollination: European Orchids*. (CRC Press,

514    2001).

515 44. Weitemier, K. *et al.* Hyb-Seq: Combining target enrichment and genome skimming for plant

516    phylogenomics. *Appl. Plant Sci.* **2,** 1400042 (2014).

517 45. Pridgeon, A. M., Cribb, P. J., Chase, M. W. & Rasmussen, F. N. *Genera Orchidacearum:*

518    *Vol. 5. Epidendroideae (part two)*. (Oxford University Press, 2009).

519 46. Chomicki, G. *et al.* The velamen protects photosynthetic orchid roots against UV-B damage,

520    and a large dated phylogeny implies multiple gains and losses of this function during the

521    Cenozoic. *New Phytol.* **205,** 1330–1341 (2015).

522 47. Ramírez, S. R., Gravendeel, B., Singer, R. B., Marshall, C. R. & Pierce, N. E. Dating the

523    origin of the Orchidaceae from a fossil orchid with its pollinator. *Nature* **448,** 1042–1045

524    (2007).

525 48. Freudenstein, J. V. & Chase, M. W. Phylogenetic relationships in Epidendroideae

526    (Orchidaceae), one of the great flowering plant radiations: progressive specialization and

527    diversification. *Ann. Bot.* (2015). doi:10.1093/aob/mcu253

528 49. Pérez-Escobar, O. A. Molecular phylogenetics, evolution of sexual systems and historical

529    biogeography of Darwin's favorite orchids (Catasetinae) and Swan orchids (Cycnoches

530    Lindl.). (Ludwig-Maximilians Universität, 2016).

531 50. Soltis, D. E. & Kuzoff, R. K. Discordance between Nuclear and Chloroplast Phylogenies in

532    the Heuchera Group Author (Saxifragaceae). *Evolution (N. Y).* **49,** 727–742 (1995).

533 51. van der Niet, T. & Peter Linder, H. Dealing with incongruence in the quest for the species

534    tree: a case study from the orchid genus *Satyrium*. *Mol. Phylogenet. Evol.* **47,** 154–74

535    (2008).

52.   Pérez-Escobar, O. A., Balbuena, J. A. & Gottschling, M. Rumbling Orchids: How to Assess

Divergent Evolution between Chloroplast Endosymbionts and the Nuclear Host. *Syst. Biol.*

65, (2016).

53.   Fragoso-Martínez, I. *et al.* A pilot study applying the plant Anchored Hybrid Enrichment

method to New World sages (Salvia subgenus Calosphace; Lamiaceae). *Mol. Phylogenet.*

Evol. **117,** 124–134 (2017).

54.   Léveillé-Bourret, É., Starr, J. R., Ford, B. A., Moriarty Lemmon, E. & Lemmon, A. R.

Resolving Rapid Radiations within Angiosperm Families Using Anchored Phylogenomics.

Syst. Biol. **67,** 94–112 (2018).

55.   Zhang, R. *et al.* Exploration of Plastid Phylogenomic Conflict Yields New Insights into the

Deep Relationships of Leguminosae. *Syst. Biol.* **69,** 613–622 (2020).

56.   Mark Whitten, W., Williams, N. H., Dressler, R. L., Gerlach, G. & Pupulin, F. Generic

relationships of Zygopetalinae (Orchidaceae: Cymbidieae): Combined molecular evidence.

Lankesteriana **5,** 87–107 (2005).

57.   Neubig, K. M., Williams, N. H., Whitten, W. M. & Pupulin, F. Molecular phylogenetics and

the evolution of fruit and leaf morphology of *Dichaea* (Orchidaceae: Zygopetalinae). *Ann.*

Bot. **104,** 457–467 (2009).

58.   Pupulin, F. in *Genera Orchidacearum. Vol. 5. Epidendroideae (Part Two)* (eds. Pridgeon,

A. M., Cribb, P. J., Chase, M. W. & Rasmussen, F. N.) **5,** 456–546 (Oxford University

Press, Oxford., 2009).

59.   Bogarín, D. *et al.* Pollination of *Trichosalpinx* (Orchidaceae: Pleurothallidinae) by biting

midges (Diptera: Ceratopogonidae). *Bot. J. Linn. Soc.* (2018).

doi:10.1093/botlinnean/box087

60.   Blanco, M. A. & Barboza, G. Pseudocopulatory pollination in *Lepanthes* (Orchidaceae:

Pleurothallidinae) by fungus gnats. *Ann. Bot.* **95,** 763–772 (2005).

61.   Damon, A. *et al.* A survey of pollination in remnant orchid populations in Soconusco ,

Chiapas , Mexico. **48,** 1–14 (2007).

62.   Policha, T. *et al.* Disentangling visual and olfactory signals in mushroom-mimicking

Dracula orchids using realistic three-dimensional printed flowers. *New Phytol.* (2016).

doi:10.1111/nph.13855

63.   Pridgeon, A. M., Solano, R. & Chase, M. W. Phylogenetic relationships in Pleurothallidinae

(Orchidaceae): combined evidence from nuclear and plastid DNA sequences. *Am. J. Bot.* **88,**

2286–2308 (2001).

64. Karremans, A. P. *et al.* Phylogenetic reassessment of *Acianthera* (Orchidaceae:

Pleurothallidinae). *Harvard Pap. Bot.* **21,** 171–187 (2016).

65. Bogarín, D., Karremans, A. P. & Fernández, M. Genus-level taxonomical changes in the

Lepanthes affinity (Orchidaceae, pleurothallidinae). *Phytotaxa* **340,** 128–136 (2018).

67. Karremans, A. P. Genera Pleurothallidinarum: an updated phylogenetic overview of

Pleurothallidinae. *Lankesteriana* **16,** 219–241 (2016).

68. Pridgeon, A. in *Genera Orchidacearum Volume 4: Epidendroideae (Part 1)* 319–422

(2005).

69. Damián, A., Mitidieri, N. & Chiron, G. A taxonomic synopsis of *Acianthera* (Orchidaceae:

Pleurothallidinae) in Peru, including two new species. *An. del Jard. Bot. Madrid* **75,** 1–21

(2018).

70. Luer, C. A. A Systematic Method of Classification of the Pleurothallidinae Versus a Strictly

Phylogenetic Method. *Selbyana* **23,** 57–110 (2002).

71. Stern, W. W. L., Pridgeon, A. & Luer, C. Stem structure and its bearing on the systematics

of Pleurothallidinae (Orchidaceae). *Bot. J. Linn. Soc.* **91,** 457–471 (1985).

72. Jost, L. & Shepard, A. Two new species of Teagueia (Orchidaceae: Pleurothallidinae) from

east-central Ecuador. *Lankesteriana* **11,** 9–14 (2011).

73. Luer, C. A. Icones Pleurothallidinarum VIII. Systematics of *Lepanthopsis*, *Octomeria*

subgenus Pleurothallopsis, *Restrepiella*, *Restrepiopsis*, *Salpistele*, and *Teagueia*. Addenda to

Platystele, *Porroglossum*, and *Scaphosepalum*. *Monogr. Syst. Bot. from Missouri Bot. Gard.*

39, 1–161 (1991).

74. Luer, C. A. Icones Pleurothallidinarum XXIV. A First Century Of New Species of *Stelis* of

Ecuador part one. Addenda to *Lepanthes* of Ecuador. Addenda to *Barbosella*, *Dracula*,

Dresslerella, *Lepanthopsis*, *Platystele*, *Pleurothallis*, *Restrepia*, *Scaphosepalum*, *Teagueia*

and . *Monogr. Syst. Bot. from Missouri Bot. Gard.* **88,** 1–122 (2002).

75. Luer, C. A. Icones Pleurothallidinarum VII. Systematics of *Platystele* (Orchidaceae).

Monogr. Syst. Bot. from Missouri Bot. Gard. **38,** 1–115 (1990).

76. Karremans, A. P. *et al.* Phylogenetic reassessment of *Specklinia* and its allied genera in the

pleurothallidinae (Orchidaceae). *Phytotaxa* **272,** 1–36 (2016).

77. Doyle, J. & Doyle, J. Genomic plant DNA preparation from fresh tissue-CTAB method.

Phytochem Bull **19,** 11–15 (1987).

78.   Neubig, K. M. *et al.* in *DNA Banking for the 21st Century: Proceedings of the US Workshop*
on DNA Banking (eds. Applequist, W. & Campbell, L.) 81–112 (William L. Brown Center,
Missouri Botanical Garden, 2014).

79.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30,** 2114–2120 (2014).

80.   Bushnell. BBMap/BBTools. (2017).

81.   Bushnell, B., Rood, J. & Singer, E. BBMerge – Accurate paired shotgun read merging via
overlap. *PLoS One* **12,** e0185056 (2017).

82.   Magoc, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics* **27,** 2957–2963 (2011).

83.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to
Single-Cell Sequencing. *J. Comput. Biol.* **19,** 455–477 (2012).

84.   Chevreux, B., Wetter, T. & Suhai, S. Genome sequence assembly using trace signals and
additional sequence information. in *German conference on bioinformatics* **99,** 45–56
(Citeseer, 1999).

85.   Cock, P. J. A., Grüning, B. A., Paszkiewicz, K. & Pritchard, L. Galaxy tools and workflows
for sequence analysis with applications in molecular plant pathology. *PeerJ* **1,** e167 (2013).

86.   Parakhia, M. V. *et al.* Draft Genome Sequence of the Endophytic Bacterium *Enterobacter*
spp. MR1, Isolated from Drought Tolerant Plant (Butea monosperma). *Indian J. Microbiol.*
54, 118–119 (2014).

87.   Ward, J. A., Ponnala, L. & Weber, C. A. Strategies for transcriptome analysis in nonmodel
plants. *Am. J. Bot.* **99,** 267–276 (2012).

88.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–
2079 (2009).

89.   Jin, J. J. *et al.* GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of
organelle genomes. *Genome Biol.* (2020). doi:10.1186/s13059-020-02154-5

90.   Tillich, M. *et al.* GeSeq – versatile and accurate annotation of organelle genomes. *Nucleic*
Acids Res. **45,** W6–W11 (2017).

91.   Chan, C. X. & Ragan, M. A. Next-generation phylogenomics. *Biol. Direct* **8,** (2013).

92.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.* **30,** 772–780 (2013).

93.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30,** 1312–1313 (2014).

94.   Drummond, A. J. & Rambaut, A. BEAST: Bayesian evolutionary analysis by sampling

trees. *BMC Evol. Biol.* **7,** 214 (2007).

95.   Xiang, X. G. *et al.* Biogeographical diversification of mainland Asian *Dendrobium*

(Orchidaceae) and its implications for the historical dynamics of evergreen broad-leaved

forests. *J. Biogeogr.* **43,** 1310–1323 (2016).

96.   Kosakovsky Pond, S. L., Frost, S. D. W. & Muse, S. V. HyPhy: Hypothesis testing using

phylogenies. *Bioinformatics* (2005). doi:10.1093/bioinformatics/bti079

Author Contributions Statement

M.A.S.S., O.A.P.E., and T.A. designed research; O.A.P.E., M.A.S.S., T.A., C.H. and A.A.

generated new data; M.A.S.S., O.A.P.E., M.F.T., A.C.A.Y., J.A. and S.D. performed all analyses;

O.A.P.E., D.B., M.W.C., M.A.S.S., S.D. and T.A. wrote the manuscript, with contributions from

all authors.

Legends

Table 1. Absolute ages and confidence intervals of main orchid lineages as inferred under a strict

and relaxed molecular clocks and a Birth-Death model.

Figure 1. Maximum Likelihood phylogeny of the orchid family inferred from 78 coding plastid

genes. Likelihood bootstrap support values (LBS) < 85% at nodes are highlighted in red together

with their corresponding subtending branches. Orchid genera, tribes and subfamilies are indicated

in the phylogeny together with photographs of selected representative species per subfamily.

(Inset): Bar plot showing the frequency of LBS values at nodes as computed by bin intervals of 5

units.

Figure 2. A comparison of the main plastid topologies of the orchid family published to date. A)

Givnish et al.[2'] inference based on 75 plastid genes and 39 orchid species; B) Li et al.[27'] inference

based on 76 plastid genes and 76 orchid species; C) This study: 78 plastid and 264 orchid species.

LBP at nodes are highlighted in red together with their corresponding subtending branches. (Inset):

trees witih branch lengths proportional to substitutions/site. Photos: O. Pérez-Escobar.

Supplementary materials

Table S1. Detailed absolute ages, confidence intervals and rates for the orchid family as inferred

under a strict molecular clock and a birth-death model. The table contains 528 rows and 16

columns and is available at https://doi.org/10.6084/m9.figshare.13185008.v1.

Table S2. Detailed absolute ages, confidence intervals and rates for the orchid family as inferred

under a relaxed molecular clock and a birth-death model. The table contains 528 rows and 16

columns and is available at https://doi.org/10.6084/m9.figshare.13185008.v1.

Table S3. Phylogenetic informativeness per-site.

Table S4. Phylogenetic net informativeness.

Table S5. Voucher information and proportion of missing data in gene alignments.

Figure S1. Detailed maximum likelihood tree of the orchid family inferred from 78 plastid genes.

LBP <100 are shown at nodes, with LBP <85 highlighted in red together with their corresponding

subtending branches.

Figure S2. Chronogram of the orchid family as inferred from a strict molecular clock and a birth-

death model. LBP at nodes <85 are highlighted in red together with their corresponding subtending

branches. Blue bars at nodes denote 95% high density probability (HDP) absolute age intervals.

Figure S3. Chronogram of the orchid family as inferred from a relaxed molecular clock and a

birth-death model. LBP at nodes <85 are highlighted in red together with their corresponding

subtending branches. Blue bars at nodes denote 95% high density probability (HDP) absolute age

intervals.

Figure S4. Phylogenetic informativeness (PI) of 78 plastid gene alignments used in this study to

infer orchid relationships. A) Chronogram of Orchidaceae as inferred by PATH8 from the ML tree

derived from RAxML; B) Per-site PI; C) Net PI.

Fig. 1. ML tree inferred from 78g and 264 terminals