A Meta-Analysis of Hippocampal Transcriptional Profiling Studies in a Selectively-Bred Rat Model Provides Converging Evidence with Genetic Sequencing to Implicate Specific Candidate Genes and Pathways in the Liability for Internalizing and Externalizing Psychiatric Disorders

Isabelle Birt[1*], Megan Hagenauer, Ph.D.[1*], Sarah Clinton, Ph.D.[2], Cigdem Aydin, Ph.D.[1], Peter Blandino, Ph.D.[1], John Stead, Ph.D.[3], Kathryn Hilde, Ph.D. [1], Fan Meng, Ph.D.[1], Robert Thompson, Ph.D.[1], Huzefa Khalil, Ph.D. [1], Alex Stefanov [1], Pamela Maras, Ph.D. [1], Zhifeng Zhou, Ph.D.[4], David Goldman[4], Stanley Watson, Jr., Ph.D.[1] and Huda Akil, Ph.D.[1]

[1]University of Michigan, Ann Arbor, MI; [2]Virginia Tech, Blacksburg, VA; [3]Carleton University, Ottawa, Ontario, CA; [4] National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda, MD

*Shared first authorship

Correspondence:
Megan Hastings Hagenauer, Ph.D.
e-mail: hagenaue@umich.edu
Molecular Behavioral Neuroscience Institute (MBNI)
205 Zina Pitcher Pl.
Ann Arbor, MI 48109

Abstract

For over 16 years, we have selectively-bred rats to react differently to a novel, anxiety-inducing environment, exhibiting either high or low exploratory activity. These "bred High Responder" (bHR) and "bred Low Responder" (bLR) rats serve as a general model for extreme manifestations of behavioral inhibition and temperament, showing large differences in a variety of internalizing and externalizing behaviors relevant to both mood and substance abuse disorders. The current study elucidated persistent differences in gene expression related to bHR/bLR phenotype across development (P7, P14, P21) and adulthood within the hippocampus, a brain structure critical for emotional regulation. To do this, we meta-analyzed eight transcriptional profiling datasets (microarray and RNA-Seq) spanning 43 generations of selective breeding (n=2-6 rats per group per dataset; total n per meta-analysis: adult: n=46, P7: n=22, P14: n=49, P21: n=21). By cross-referencing these results with a concurrent exome sequencing study performed on our colony, we pinpointed several genes that are strongly implicated in bHR/bLR behavioral phenotype, including two genes previously associated with energy metabolism and mood: Thyrotropin releasing hormone receptor (Trhr) and the mitochondrial protein Uncoupling protein 2 (Ucp2). Our meta-analysis also highlighted robust bHR/bLR functional differences in the hippocampus, including a network essential for neurodevelopmental programming, cell proliferation, and differentiation, which centered on the hub genes Bone morphogenetic protein 4 (Bmp4) and the canonical Marker of proliferation (Mki67). Another functional theme was microglial activation and phagocytosis, including differential expression of pro-inflammatory Complement C1q A chain (C1qa) and the anti-inflammatory Milk fat globule-EGF factor 8 (Mfge8), situated within a chromosomal loci implicated by our concurrent genetic study. Given the newly-discovered role of microglia in synaptic pruning in relationship to neuronal activity during both development and adulthood, we propose that these functional pathways have the capability to not only direct bHR and bLR rats along a different developmental trajectory, but to set the stage for a widely-different reactivity to the environment.

Keywords: Anxiety, Depression, Hyperactivity, Impulsivity, Limbic, Neonatal, Postnatal, Trhr, Ucp2, Ncan, Bmp4, Mki67, C1qa, Mfge8, Etv4

Introduction

Both mood disorders and substance abuse disorders affect approximately 8-10% of adults in the United States each year (1, 2). Due to high comorbidity, these disorders are often classified as either internalizing and externalizing. Internalizing disorders are associated with neuroticism, anxiety, and depression, whereas externalizing disorders are associated with greater risk-taking and novelty-seeking, as seen in mania, substance abuse, and impulse-control disorders (3). This pattern of comorbidity is thought to represent a spectrum of latent liability, which arises from a complex interplay of genetic risk and environmental factors, such as stress and childhood adversity (3–5).

Within our laboratory, we model the genetic contributions underlying both extremes of this spectrum by selectively breeding rats that react differently when introduced to a novel, anxiety-inducing environment. Our "bred high responder" (bHR) rats are highly exploratory, and exhibit multiple behavioral manifestations of a disinhibited, novelty-seeking temperament, including hyperactivity, aggression, and drug-seeking. The "bred low responder" (bLR) rats are highly-inhibited, and show both reduced locomotor activity and general neuroticism, including anxiety and depressive-like behaviors (6–15). Notably, these behavioral propensities are robust and stable, beginning early in development (16, 17) similar to temperament in humans (18).

This highly-differentiated phenotype makes our selectively-bred bHR and bLR rats ideal for observing the developmental programming and adult manifestation of the neurochemistry and morphology underlying internalizing and externalizing tendencies (10, 13, 19). For our current study, we focused on the hippocampus, a region deep in the temporal lobe important for emotion and behavioral inhibition (20–22). The hippocampus regulates reactivity to the environment, including contextual conditioning (20) and the release of glucocorticoids following a stressful event (23, 24). Its volumetric size decreases in response to anxiety and depression due to reductions in cell growth (25). Within the bHR/bLR model, we have previously observed multiple molecular and morphological differences within the hippocampus, including glucocorticoid receptor and growth factor expression, histone methylation, cell proliferation, survival, and overall volume (7, 11, 16, 26, 27).

In our current study, we characterized both the developmental trajectory and stable global differences in gene expression within the hippocampus of bHR and bLR rats by performing a meta-analysis of eight transcriptional profiling datasets (microarray and RNA-Seq) collected across four developmental stages (P7, P14, P21, adulthood) and spanning 43 generations of selective breeding. The top results identified by this meta-analysis included genes with known connections to affective behavior and highlighted functional pathways broadly related to mood and neurodevelopment. Concurrently, we performed a genetic study on our colony that discovered a minimum of seven chromosomal regions containing bHR/bLR segregating variants that are likely to contribute to exploratory locomotor phenotype (28). By comparing across these two studies, we were able to identify differentially-expressed genes situated within implicated chromosomal regions, pinpointing promising candidates that could mediate the influence of selective breeding on exploratory locomotion. Further, we determined which of those candidates were potentially key-players in the hippocampal functional pathways implicated in bHR/bLR phenotype by our transcriptional profiling meta-analysis. We highlight these candidates in our results and discussion below.

Methods

The bHR/bLR Rat Colony

Selective Breeding: We began selectively-breeding bHR and bLR rats in the Molecular Behavioral Neuroscience Institute (MBNI) at the University of Michigan in 2003 (protocol: (13)).

For the first generation, we chose 120 Sprague–Dawley rats (Charles River, Inc) that exhibited
extreme locomotion scores within a novel environment (the top and bottom 20%, respectively).
Since then, we have maintained 12 breeding families in each line. For generations F1-F41, we
amplified the differences between the two lines while minimizing inbreeding by mating the males
and females with the most extreme locomotor scores from each family. At generation F42, the
breeding strategy at MBNI changed to target a phenotypic locomotor score (~1800 beam breaks
for bHR and ~150 beam breaks for bLR) to prevent the lines from becoming more extreme (**Fig
1**). During generation F30, a second colony was begun at the University of Alabama-
Birmingham using bHR/bLR rats from the MBNI colony.
95
96   [FIGURE 1]
97
98   The transcriptomic datasets used in this meta-analysis were derived from male bHR/bLR
99   rats that spanned a wide range of generations (F4 to F43). Using breeders from generation F37,
100 we also produced an bHRxbLR cross ("Intermediate Responder" (bIR) rats). These bIRs were
101 included in the transcriptional profiling and behavioral analysis associated with the
102 *MBNI_RNASeq_F37* dataset.
103
Animal Care: On postnatal day 1 (PND1), litters were reduced to 12 pups (6 males, 6
females) and raised by their mothers. Following weaning, the rats were housed 2-3 per cage in-
house and maintained on a 12:12 light:dark schedule, with access to food and water *ad libitum*.
All experiments were approved by the University Committee on the Use and Care of Animals
(UCUCA) at the University of Michigan or at the University of Alabama at Birmingham in
accordance with the National Institute of Health (NIH) Guide for the Care and Use of Laboratory
Animals, dictated by the National Research Council in 1996.
111
Behavioral Analysis: In each generation, locomotor response to a novel environment
was assessed between ages P50–75 (protocol: (13)). In association with the
MBNI_RNASeq_F37 dataset, we also measured anxiety-like behavior in adulthood (ages P160-
P167 for bHR and bLR rats, P65-75 for bIR rats) as indexed using the percent time spent in the
open arms of an Elevated Plus Maze (EPM) during a 5 min test (protocol: (29)). The number of
fecal boluses excreted during the EPM test served as a measure of anxiety independent from
spontaneous exploration (30). In association with the *MBNI_RNASeq_F43* dataset, we
measured social interaction in adulthood (P92, protocol: (6)) after 15 minutes of exposure to the
anxiogenic EPM (protocol: (31)).
121
122 **Hippocampal Gene Expression Analyses**
123
124 ***Broad Overview of the Datasets:*** Within our meta-analysis of hippocampal
125 transcriptional profiling studies, we used eight datasets collected from bHR/bLR rats from four
126 age groups: P7, P14, P21, and adult. We refer to these datasets according to their respective
127 institution, transcriptomic profiling platform, and generation (**Table 1**).
128
129 [TABLE 1]
130
When considering the results, please note that the transcriptional profiling platforms
changed with time: earlier generations (F4, F6, F15) were profiled using microarray, whereas
later generations were primarily profiled using RNA-Seq (F29, F37, F43). In general, the tissue
was derived from baseline animals although adults from two studies were exposed to behavioral
testing or repeated saline injections. In all studies, the rats were sacrificed by rapid decapitation
without anesthesia. Within seven datasets, the whole hippocampus was dissected immediately

after brain extraction, whereas one dataset performed tissue punches on sliced frozen tissue
from the dorsal hippocampus alone. Details for each dataset can be found in **Table 1** and the
Suppl. Methods. Analysis code is available at *https://github.com/isabellie4/PhenotypeProject*
and *https://github.com/hagenaue/bHRbLR_MetaAnalysisProject*.
141
Broad Overview of the Data Preprocessing: We analyzed the datasets individually
and then performed a collective meta-analysis (statistical software: R Studio v.3.3.2, R v.3.2.2).
The pre-processing steps for each study varied according to the platform and data format
(**Suppl. Methods**), but included many common steps, including re-annotation, normalization to
reduce the influence of technical variation, and quality control. Microarray data were typically
summarized into expression sets using the Robust Multi-Array Average method (RMA: (32)),
which adjusted for background signal, corrected technical variation using quantile normalization,
and log(2)-transformed signal values. Similarly, gene-level RNA-Seq read count summaries
were converted to log(2) fragments per million (FPM). Quality control for both data types
included an examination of the overall data distribution for all genes and the pattern of sample-
sample correlations (*e.g.,* principal components analysis). When applicable, transcript data was
averaged by gene symbol to get a single expression value per sample per gene.
154
Cell Type Data Deconvolution: After preprocessing, the data underwent a cell type-
based matrix deconvolution procedure using previously-documented cell type-specific gene
expression (R package *BrainInABlender*; (33)). For this analysis, we excluded the small
MBNI_RNASeq_F29 dataset (n=2 per group), because the raw data was inaccessible at the
time of analysis. To estimate relative cell type balance, z-score normalized data for genes
previously-identified as having cell type specific expression were averaged for each cell type for
each subject (33). The cell type categories included astrocyte, endothelial, microglia, mural,
oligodendrocyte, immature oligodendrocyte (progenitor), and red blood cell. We excluded
cortically-derived neuronal categories, due to their questionable applicability to hippocampal
tissue.
165
Calculating Effect Size: Within each individual dataset, we calculated the effect size
(Cohen's d and variance of d) for the effect of bHR/bLR phenotype on both log(2) gene
expression (either hybridization signal or FPM) and on the estimates of relative cell type
expression ("cell type indices") within each age group (R package *compute.es* (34)). For the
small MBNI_RNASeq_F29 dataset, the raw data were not accessible at the time of analysis, but
we were able to re-derive estimated effect sizes from the t-statistic output from the original
CuffDiff analysis (35), averaging the small subset of results representing duplicated gene
symbols.
174
Meta-Analysis: The Cohen's d and variance of d for each dataset were grouped
according to age (P7, P14, P21, and adult), and aligned using official gene symbol. To be
included in the meta-analysis, a gene was required to be present in at least two datasets. As a
result, the P14 and Adult meta-analyses contained a much larger number of genes because
they contained five datasets, whereas the P7 and P21 meta-analyses each contained two
datasets (**Figure 2**).
181
182   [FIGURE 2]
183
The meta-analysis was performed on the collective Cohen's d and variance of d for each
age group for each gene using the *rma.mv()* function and visualized using forest plots created
using the *forest.rma()* function (R Package *metafor* (36)). False discovery rate (FDR) was
calculated via the *multtest* package in R (37) using the Benjamini-Hochberg method. Additional

exploratory analyses revealed that including generation as a co-variate in the meta-analysis
provided little additional insight (generation: all genes FDR>0.3), most likely due to lack of
power. The cell type meta-analyses followed similar methods. However, the resulting *p*-values
did not undergo a multiple comparison correction because there were a small number of cell
types present within each analysis.
193
Gene Set Enrichment Analysis: We ran Gene Set Enrichment Analysis (GSEA) (38,
39) to elucidate overall functional trends within the two largest sets of results: P14 and Adult.
The gene lists were ranked according to the beta coefficient values (β) representing the
magnitude and direction of effect estimated by the meta-analysis. We performed GSEA using
the *fgsea()* function (R package *fgsea* (40); settings: 100,000 permutations, maxSize=1000).
We ran GSEA using gene set matrix files (.gmt) containing standard gene ontology
biological processes gene sets for rats (go2msig.org; (41)), and then created a custom .gmt file
(**Suppl. Table 1**) focused on hippocampal-specific gene sets that we curated from three
sources: *1)* human hippocampal co-expression modules derived from a large sample of post-
mortem and freshly-resected tissue (42); *2)* mouse hippocampal co-expression modules
identified using a large sample (n=100) of behaviorally-profiled animals from the hybrid mouse
diversity panel (43); *3)* sets of genes with expression specific to hippocampal neuronal subtypes
or regions (*Hipposeq:* (44)).
207
Protein-protein Interaction (PPI) Networks: We further explored the hippocampal-
specific gene sets that were enriched for bHR/bLR differential expression by constructing
predicted PPI networks using string-db.org (45) (confidence setting=0.15 due to pre-existing
evidence suggesting interaction). We also constructed a PPI network using all differentially
expressed genes from the adult meta-analysis (FDR<0.10) and a higher confidence setting
(0.40).
214
Positional Gene Enrichment Analysis: We evaluated whether the top bHR/bLR
differentially expressed genes clustered around particular chromosomal locations to explore
which genes may be either co-regulated or in linkage disequilibrium with a causal genetic
variant. To perform this analysis, we extracted top results from the P14 and adult meta-analyses
(nominal p<0.01), and removed duplicated EntrezIDs. Enrichment at particular chromosomal
loci were then evaluated using Positional Gene Enrichment analysis (PGE,
http://silico.biotoul.fr/pge/; (46)). Top clusters (p<0.001, FDR<0.01) were cross-referenced with
a newer assembly (Rnor_6.0, https://www.ncbi.nlm.nih.gov/genome/gdv/?org=rattus-
norvegicus).
224
Overlap with Previously-Identified Genetic QTLs: Our concurrent genetic study
performed exome sequencing to identify bHR/bLR segregating variants, and then used a
sampling of those segregating variants to pinpoint QTLs for exploratory locomotion using an
bHRxbLR F2 intercross (28). For our current study, we identified all genes within +/-1MB of
those QTL peaks (LOD>3) using Rnor_6 annotation from the R package *org.Rn.eg.db (47)* and
the NCBI Genome Remapping Service (https://www.ncbi.nlm.nih.gov/genome/tools/remap,
accessed 8/8/2019 using the first 100 bp of each gene and default parameters). We then
evaluated whether our top adult meta-analysis results (FDR<0.05) where enriched for these
QTL genes using Fisher's Exact Test. We also evaluated overlap with additional QTLs relevant
to non-locomotor aspects of the bHR/bLR behavioral phenotype (from the Rat Genome
Database (48), accessed 08/08/2019 using the key words "Anxiety", "Stress", and "Despair",
and Rnor6 chromosomal coordinates).

Examining the Relationship Between Gene Expression and Behavior: For the two
adult datasets that contained associated behavioral data (*discussed above*), we used simple
bivariate linear models to explore the relationship between gene expression and behavior: total
locomotor score, percent time in the open arm of the EPM, fecal boli produced during the EPM
test, and percentage of time spent interacting socially following exposure to a single mild
stressor. A general effect of phenotype on these variables was evaluated using ANOVA (Type
3).

qPCR Validation: Using pups from later generations (18 litters: generations F44, F51,
F54, F55, and F57), hippocampal tissue from 6-10 bHR and bLR males was collected at each of
the following time points: P2, P7, P14, P21, P60, P90 (**Suppl. Fig 1**) using methods similar to
those previously described. Following cDNA synthesis using RT-PCR, Bmp4 transcript was
quantified using qPCR and custom-designed primers (ACC# NM_012827.2; forward primer: 5'-
CCCTGGTCAACTCCGTTAAT-3', start = 1214; reverse primer: 5'-
AACACCACCTTGTCGTACTC-3', start = 1319) using the reference gene glyceraldehyde-3-
phosphate dehydrogenase (Gapdh) (ACC# NM_017008.4; forward primer: 5'-
GTTTGTGATGGGTGTGAACC-3', start = 459; reverse primer: 5'-
TCTTCTGAGTGGCAGTGATG-3', start = 628). The calibration curves revealed efficiencies
close to 1 for each probe (Bmp4: R^2=0.98, GAPDH: R^2=0.99, **Suppl. Fig 1**), therefore the qPCR
data for each developmental time point was analyzed using the traditional Livak method (49),
with group differences in ΔC_q assessed using Welch's two sample t-test ((50); further details:
Suppl. Methods).

259   **Results**
260
Selective Breeding Amplifies the Propensity for Internalizing vs. Externalizing Behavior
Over sixteen years of selective breeding, bHR/bLR differences in exploratory locomotor
activity grew increasingly prominent (**Fig 1A**). This divergence happened rapidly in a manner
implying oligogenic inheritance (28), a conclusion supported by our recent genetic study (28).
As shown previously (13, 16, 51, 52), these differences in exploratory locomotion were
accompanied by differences in affective behavior. For example, in the behavioral data
accompanying the *MBNI_RNASeq_F37* dataset, we observed not only a large effect of
phenotype on total locomotor score ($F_{(2,15)}$=444.47, *p*=4.46E-14), but also an effect on anxiety
as measured by the percent time spent in the open arms ($F_{(2,15)}$=6.72, *p*=8.25E-03) or the
number of fecal boli produced in the EPM ($F_{(2,15)}$=6.40, *p*=9.79E-03, *data not shown*), with
bHRs showing less anxiety than bLRs, and bIRs showing an intermediate behavioral phenotype
(**Fig 1B**). In the behavioral data accompanying the *MBNI_RNASeq_F43* dataset, bLRs spent
less time interacting socially following exposure to a stressor than bHRs ($F_{(1,8)}$=5.86,
p=0.0418). Therefore, selective breeding for exploratory locomotion amplified traits that are
broadly associated with internalizing and externalizing tendencies. For this reason, we expected
that an examination of gene expression across bHR/bLR generations would reveal a
convergence of effects within pathways essential to affective behavior and general reactivity to
the environment.
279
A Meta-analysis of the Effects of bHR/bLR Phenotype on Hippocampal Gene Expression
Reveals Promising Candidates: Overview
During the many years that we have been selectively breeding bHR/bLR rats, we
conducted eight transcriptomic studies profiling the hippocampus of bHR/bLR rats at four ages:

P7, P14, P21, and adult (**Table 1).** The results from these small studies were often noisy, and
produced few reliable results when analyzed individually. For example, the pairwise correlations
between the effect sizes for the individual studies were weak (ranging between R=-0.22 to
R=0.17), exhibiting almost no overlap amongst the top findings until the very latest generations
(F37 vs. F43: R=0.40, **Suppl. Figs 2-3**). Nevertheless, a formal meta-analysis of these datasets
revealed multiple robust and interesting candidate genes with consistent differential expression
across multiple generations both in adulthood and development (**Fig 2, Table 2, Suppl. Table
2**). These results can be explored interactively at
https://y.mbni.org/bHRbLR_HippocampalMetaResults/.
293
294   [TABLE 2]
295
Adult: The adult meta-analysis included expression data from a total of 16,269 unique
gene symbols, 4,381 of which were represented in all five datasets. The effect of bHR/bLR
phenotype on gene expression was significant for 74 genes (FDR<0.05, **Table 2**), 32 of which
were found in at least four datasets. In general, we observed equal distributions of positive and
negative estimated effect sizes (β) in the meta-analyses from all age groups (**Figure 2**). We
found that the estimated effect sizes (β's) tended to be more extreme for genes with expression
data in fewer datasets, most likely because these genes were only present in the RNA-Seq
datasets from later generations. For that reason, as a point of comparison we ran a meta-
analysis using just the adult RNA-Seq data from the two most recent generations (F37 and
F43), and confirmed that similar differentially-expressed genes and pathways were identified
(**Suppl. Fig 4**). Therefore, the differentially-expressed genes that were identified within the full
meta-analysis using data from a greater number of datasets are of particular interest because
we have evidence that their expression began diverging during the earliest generations in
addition to clear evidence that they are differentially expressed in recent generations.
310
Development: The effects of bHR/bLR phenotype on gene expression in the
developmental meta-analyses were less robust, most likely due to a greater dependency on
data from earlier generations. The P14 meta-analysis included expression data from a total of
15,682 unique gene symbols, 2,353 of which were represented in all five datasets. None of
these genes showed an effect of bHR/bLR phenotype that survived false detection correction
(FDR<0.05), but the top gene, Bone morphogenetic protein 4 (Bmp4), was consistently
expressed at lower levels in bHRs than bLRs within both the P14 (β=-1.49, *p*=9.40E-06,
FDR=1.47E-01) and adult datasets since the F4 generation (β=-1.04, *p*=1.01E-03, FDR=9.38E-
02, **Fig 3**) and was highlighted in many of our functional analyses (*discussed below*). We
confirmed this differential expression across development (P2, P7, P14, P21, P60, P90) using
qPCR and two separate cohorts of rats (**Fig 4**, **Suppl. Fig 1**; Welch's t-test: *P2*: Log(2)FC=-
1.11, T(17.0)=-2.52, p=0.0219; *P7*: Log(2)FC=-1.68, T(14.2)=-3.34, p=0.00483; *P14*:
Log(2)FC=-3.74, T(5.60)=-6.10, p=0.00115; *P21*: Log(2)FC=-1.59, T(7.72)=-2.76, p=0.0257;
P60: Log(2)FC=-2.04, T(5.99)=-3.22, p=0.0182; *P90*: T(8.74)=-6.87, p=8.44E-05).
325
326   [FIGURE 3, FIGURE 4]
327
The P7 and P21 meta-analyses included a much smaller sample size (only two datasets
in each) and expression data from a much smaller number of unique gene symbols (3,527).
Despite these constraints, within the P7 meta-analysis one gene, Neurocan (Ncan), survived
FDR correction. Ncan was more highly expressed in bLRs since as early as the F6 generation,
showing an unusually strong effect size (β= -4.16, *p*=1.38E-07, FDR= 4.86E-04; **Fig 5**). Ncan
also showed nominally higher expression in bLRs within the P14 meta-analysis (β= -0.89,

p=0.010, FDR=0.72), and the trend persisted in the adult meta-analysis (*β*= -0.57, *p*=0.067,
FDR=0.47).
336   [FIGURE 5]
337
Many of the Top Differentially-Expressed Genes Are Located Within Quantitative Trait
Loci for Exploratory Locomotion Identified by Our Concurrent Genetic Study
A concurrent genetic study performed on our colony used exome sequencing to identify
genetic variants segregating selectively-bred bHR and bLR rats, and then used a sampling of
those variants to perform a whole-genome analysis identifying QTLs for exploratory locomotor
activity in an bHRxbLR F2 intercross (28). They identified seven genome-wide significant QTL
peaks (LOD>4), and six additional promising candidates (LOD>3) that are likely to contribute to
bHR/bLR differences in exploratory locomotion. These 13 candidate loci also overlapped
extensively with other previously-identified QTLs present in the Rat Genome Database (48)
relevant to non-locomotor aspects of the bHR/bLR behavioral phenotype, including QTLs for
anxiety (53–58), stress response (59–61), and behavioral despair (62). We found that eight of
the top differentially-expressed genes within our adult meta-analysis (FDR<0.05) were located
within these loci (**Figure 6A**), including two genes previously associated with internalizing and
externalizing behaviors (Trhr, Ucp2; **Fig 7&8**, (63–68)), and two genes highlighted by our
functional analyses (Trhr, Mfge8, **Fig 7&9,** *discussed below*). In general, the top differentially
expressed genes from our adult meta-analysis (FDR<0.05) were 4.5x more likely to be located
within a QTL for exploratory locomotion (LOD>4) than the rest of the genes included in our
analysis (**Figure 6B,** Fisher's exact test: *p*=0.00353). This pattern of results fits our expectation
that the influence of bHR/bLR segregating genetic variants on exploratory locomotion is at least
partially mediated by effects on gene expression within the hippocampus.
358
359   [FIGURE 6]
360
Positional Gene Enrichment Analysis Specifies Narrower Chromosomal Regions
Contributing to bHR/bLR Phenotype
Using our transcriptomic data, Positional Gene Enrichment (PGE) identified 132
chromosomal regions that showed a significant enrichment (FDR<0.05) of differentially
expressed genes within the P14 and adult meta-analyses. We focused on the top 13 regions,
which showed FDR<0.001 (**Figure 6C**). Ten of these top loci included genes with differential
expression that survived FDR<0.10 in the adult meta-analysis, and most loci (10/13) were also
confirmed using a PGE analysis of adult meta-analysis results using only RNA-Seq data from
the two most recent generations (F37 & F43), ruling out any bias towards regions
overrepresented on older microarray platforms. Notably, three of the top loci were narrow
regions within previously-identified QTLs for exploratory locomotor activity in a bred bHRxbLR
F2 intercross (28) (LOD>4; *discussed above*), encompassing two differentially-expressed genes
previously associated with internalizing and externalizing behavior (Ucp2, Trhr; **Fig 7 & 8**, (63–
68)). Several of the loci also contained differentially expressed genes highlighted by our
functional analyses below (Trhr, Etv4, C1qa, **Fig 7 & 9, Suppl. Fig 7**). Provocatively, even
though the top loci identified by PGE were much narrower than the regions identified as QTLs
for exploratory locomotion in our concurrent genetic study (measured in KB instead of MB), they
overlapped with a strikingly high percentage of the QTLs relevant to externalizing and
internalizing behaviors identified within the Rat Genome Database (RGD: (48)), including 13%
of the QTLs for anxiety (6/45, (53–56)), 21% of the QTLs for stress-related responses (8/38,
(59–61, 69–71)), and 23% of the QTLs for behavioral despair (3/13, (62)), suggesting that these
loci could represent variants contributing to internalizing/externalizing aspects of the bHR/bLR
behavioral phenotype beyond exploratory locomotion.
384

385   [FIGURE 7, FIGURE 8, FIGURE 9]
386
387   **bHR/bLR Differential Expression is Enriched within Particular Functional Pathways**
388
bHR/bLR Phenotype is Associated with Proliferation and Differentiation: When
running GSEA using traditional functional ontology gene sets, eight gene sets (out of 2,761)
showed an enrichment of differential expression within the P14 meta-analysis (FDR<0.05), all of
which were upregulated in bLRs (**Fig 3, Suppl. Table 3, Suppl. Fig 5**) and predominantly
related to neurogenesis, neuron differentiation, and nervous system development. Within the
adult meta-analysis, 2 of the 4 top gene sets showing an enrichment of differential expression
(FDR<0.1, out of 2,761 total) were similarly related to proliferation and development, but
upregulated in bHRs. This pattern continued to be present within the adult meta-analysis results
when only considering RNA-Seq data from the latest generations (F37 and F43, **Suppl. Table**
3).
In general, the top genes in these pathways in development and adulthood included
Sox9, Bmp4, Farp1, Bex1, Cd24, Mfge8, Apln, Cav1, Hes5, Htra1, and Glul. Indeed, a PPI
network constructed using the top genes from the adult meta-analysis (192 genes with
FDR<0.10) included a dominant subnetwork that highlighted many of these same genes (**Fig 3**),
and centered on hub genes including the top result from the P14 meta-analysis, Bmp4, and the
canonical marker of proliferation Mki67. This network also included Sox9, which is located within
a chromosomal loci that is highly enriched for differential expression overlapping QTLs related
to anxiety and stress-response ((56, 70, 71), **Figure 6C**). A literature search confirmed the PPI
interactions within this subnetwork and highlighted their role in proliferation and differentiation
within the brain, as well as their re-activation in response to injury (72–84).
409
bHR/bLR Phenotype is Associated with the Dentate Gyrus: The functional ontology
results suggested that hippocampal development and renewal might progress differently in bHR
and bLR rats. When performing GSEA using a set of 69 gene sets custom-designed to reflect
hippocampal-specific cell types and networks **(Suppl. Table 1**), we similarly observed an
enrichment of differential expression within two gene sets related to the dentate gyrus (**Fig 7**,
Suppl. Fig 6, Suppl. Table 3), the location of neural proliferation within the hippocampus.
Within the P14 meta-analysis, bLRs showed an upregulation of genes with enriched expression
in the dentate gyrus as compared to the Cornu Ammonis (CA) regions (44) (NES<0;
FDR<0.05). Within the adult meta-analysis, bHRs showed an upregulation of genes with
enriched expression in the ventral compared to the dorsal dentate gyrus (44) (NES<0;
FDR<0.05), including one of the top meta-analysis genes, Trhr (FDR<0.05, **Fig 6**) which is
located within a chromosomal loci that was highly enriched for differential expression and
overlaps a QTL for exploratory locomotion (*discussed above*). We confirmed that Trhr is a
strong marker for the ventral dentate gyrus using mouse *in situ* hybridization images publically-
available in the Allen Brain Atlas (85). Interestingly, the differentially-expressed genes
associated with the dentate gyrus-related results were largely independent from those driving
the functional associations with proliferation and cell differentiation.
427
bHR/bLR Phenotype is Associated with Hippocampal Co-expression Networks
Related to Synaptic Signaling: Co-expression modules can capture regionally-important cell
types and functions that remain undocumented in traditional ontology databases (86). We
observed an enrichment of bHR/bLR effects within six hippocampal co-expression modules
within the P14 meta-analysis (FDR<0.05) and within five co-expression modules within the adult
meta-analysis (FDR<0.05, **Fig 7, Suppl. Fig 6, Suppl. Table 3**). For the sake of conciseness,
we have highlighted the results from two of these modules which include genes that were

associated with chromosomal loci enriched for differential expression. A more detailed
description can be found in the **Suppl. Results.**
The first co-expression module of interest was *lightcyan*, a large module (695 genes)
previously-identified in the mouse hippocampus (43). This module showed elevated expression
in bLRs relative to bHRs (NES<0) in both development (FDR<0.05) and adulthood (FDR<0.10)
and contained three genes with robust bHR/bLR effects in adulthood (FDR<0.05, Etv4, Rltpr,
Slc27a1). One of these genes, Etv4, was also found within a chromosomal loci highly enriched
for differential expression (**Suppl. Fig 7**) overlapping QTLs related to anxiety and stress-
response (56, 70, 71). Etv4 is a transcription factor required for proper hippocampal dendrite
development (87) and, unlike the rest of the co-expression module, was more highly expressed
in bHRs (β=2.02, p=3.30E-05, FDR=1.92E-02). When we constructed a PPI network (medium
confidence=0.40) using the genes from this co-expression module that showed bHR/bLR
differential expression in adulthood (n=74, p<0.05), we found an enrichment of genes related to
cell projections, neurons, synapses, and cation binding (FDR<0.05).
The second module of interest was *sienna3*, which showed elevated expression in bHRs
(NES>0; FDR<0.05) in the adult meta-analysis. The top gene in this module was Trhr, which, as
previously discussed, has elevated expression in the ventral dentate gyrus, and was located
within a chromosomal loci overlapping QTLs for exploratory locomotor activity (28), anxiety and
stress-response (55, 59). A PPI network constructed using all 39 genes in this small co-
expression module centered on its ligand, thyrotropin releasing hormone (Trh; **Fig 7**), and
included a wide variety of signaling molecules with well-known connections to reward behavior,
including Cartpt (88), Oxt (89) and Drd1a (90, 91).
457
bHR/bLR Phenotype is Associated with Microglia and Endothelial-Specific Gene
Expression: The functional ontology and co-expression gene set enrichment results suggested
that there might be overall differences in cell type balance in the hippocampus. Therefore, we
applied a more traditional cell type deconvolution method to assess expression specific to a
variety of non-neuronal cell types. At P14 and adulthood, we found that bLRs showed relatively
higher microglial-specific expression compared to bHRs (Adult: β=-0.65, *p*=4.47E-02, P14: β=-
0.69, *p*=2.90E-02; **Fig 9A&C**). At P14, bLRs also showed greater endothelial-specific
expression (β=-0.89, *p*=5.75E-03 (**Fig 9B**). These effects could reflect overall differences in cell
type balance or activation state. Notably, two of the top differentially-expressed genes in the
adult meta-analysis are well-known regulators of microglial state: Milk fat globule-EGF factor 8
(Mfge8: β=1.50, p=2.70E-05, FDR=1.73E-02, **Fig 9 D&E**), which promotes alternative (M2)
activation, and Complement component C1q A Chain (C1qa: β=-1.40, p=6.57E-05, FDR=2.61E-
02, **Fig 9 F&G**), which promotes classical activation (92). The differential expression for both of
these genes was also plausibly driven by genetic variants segregating bHR and bLR rats in our
colony: C1q was located in a region of chromosome 5 that we found was enriched with
differentially expressed genes and overlapped a previously-identified QTL for anxiety-related
behavior ((55), *discussed above*), and Mfge8 was located within a QTL on chromosome 1 for
exploratory locomotor activity within our concurrent bHR/bLR genetic study that just barely
missed the threshold for genome-wide significance (28) and overlapped previously-identified
QTLs for anxiety-related behavior (53, 54) and stress response ((60), *discussed above*).
Beyond these findings, a comparison of our meta-analysis results (FDR<0.10) with the
new mousebrain.org database (93) indicated that our top bHR/bLR differentially expressed
genes are highly expressed in a variety of hippocampal cell types (**Suppl. Fig 8**). Therefore,
bHR/bLR phenotype correlates with either the activation or balance of cell types within the
hippocampus, as well as gene expression within many kinds of cells, including pathways related
to cell proliferation, differentiation, regional functionality, and synaptic signaling.
484
485

Discussion

By selectively-breeding rats for over 16 years, we have produced a robust, genetic model of the co-occurrence of common internalizing and externalizing behaviors, including extremes in exploratory activity, anxiety, stress-susceptibility, social interaction, risk-taking, reward-seeking, and drug self-administration (6–17). Such large differences in behavior would be expected to be accompanied by similarly strong differences in gene expression in areas of the brain responsible for affective behavior. However, prior to our meta-analysis, we found that many of the effects of bHR/bLR phenotype on hippocampal gene expression could not be reliably identified within noisy, small individual transcriptional profiling datasets. By performing a formal meta-analysis across multiple datasets, we managed to greatly increase the effective sample size in our study and consequently provided more substantial insight into phenotypic differences in hippocampal gene expression across development and adulthood. Further, by cross-referencing these results with a concurrent exome sequencing study performed within our colony, we were able to hone in on several strong candidates for mediating the influence of selective breeding on behavior. Through exploration of these candidate genes and their interactions we hope to provide initial insight into the gene expression underlying susceptibility to internalizing and externalizing disorders.

Cross-referencing our meta-analysis results with a concurrent genetic study performed on our colony identifies two strong candidates for contributing to bHR/bLR behavioral phenotype: Trhr and Ucp2

A concurrent exome sequencing study performed on our colony identified genetic variants segregating bred bHR and bLR rats, and then used a sampling of those variants to identify seven genome-wide significant QTL peaks for exploratory locomotor activity in an bHRxbLR F2 intercross (28). These findings offer a first glimpse at the genetic factors contributing to our selectively-bred model. However, the QTLs are broad, encompassing several hundred genes, and it is difficult to predict how the variants contained within these regions influence gene expression and behavior. By cross-referencing the genes encompassed by these QTLs with our differential expression findings within a brain region important for behavioral inhibition and emotional reactivity we identified two particularly strong candidate genes. These candidates have been previously-associated with mood and energetics, and thus seemed capable of mediating the influence of selective breeding on exploratory locomotion as well as other internalizing vs. externalizing behaviors.

The first gene, Thyrotropin Releasing Hormone Receptor (Trhr, or Trhr1), exhibited consistently higher expression in bHRs since at least the F4 generation in the adult meta-analysis. Within our concurrent genetic study (28) we found that Trhr was located on chromosome 7 adjacent to a genetic variant that fully segregated bHR/bLR rats in our colony and within a QTL for exploratory locomotor activity in the bHRxbLR F2 intercross. This region of chromosome 7 overlaps with previously-identified QTLs for anxiety (55) and stress response (59). Within our transcriptomic meta-analysis, we found that a narrow part (1.4 MB) of this locus was enriched for bHR/bLR differentially expressed genes, including Trhr.

Trhr has been studied for years due to its involvement in the hypothalamic-pituitary-thyroid (HPT) axis, which is important for regulating both energetics and mood (94, 95). In previous studies, Trhr1-KO mice displayed increased anxiety and depressive-like behavior (68) similar to our bLRs. Likewise, the Trh analog, Glu-Trh, decreased anxiety in rats (65). We found that Trhr was the top gene within a bHR-upregulated gene set associated with the ventral dentate gyrus (44), a region of the hippocampus that may be particularly important for emotional regulation and stress (20). Its ligand, Trh, was also a hub gene within an bHR-upregulated hippocampal-specific co-expression and protein-protein interaction (PPI) network that included many other genes with well-known connections to reward behavior, including Cartpt (88), Oxt

(89) and Drd1a (90, 91). Together, this evidence strongly suggests that genetic variation within
the QTL on chromosome 7 is driving differential expression of Trhr in the hippocampus in a
manner that may directly contribute to bHR/bLR differences in exploratory locomotion as well as
their propensity for other internalizing vs. externalizing behaviors.
The second gene, mitochondrial protein Uncoupling Protein 2 (Ucp2), also showed
consistently higher expression in bHRs in the adult meta-analysis across many generations of
selective breeding. Ucp2 is located on chromosome 1 adjacent to a genetic variant that fully
segregated bHR/bLR rats in our colony and within a QTL for exploratory locomotor activity in the
bHRxbLR F2 intercross (28). This region of chromosome 1 overlaps with previously-identified
QTLs for anxiety (53, 54), and stress response (60). Within our transcriptomic meta-analysis, we
found that a narrow part (1.4 MB) of this region was enriched for bHR/bLR differentially-
expressed genes, including Ucp2. In the literature, similar to our bLR rats, Ucp2-KO mice
showed higher anxiety-like behavior and lower locomotor activity at baseline, and lowered
resilience to stress (63, 64, 66, 67). Therefore, it seems likely that genetic variation on
chromosome 1 is driving differential expression in Ucp2 in a manner that may directly contribute
to bHR/bLR differences in exploratory locomotion and the propensity for internalizing behaviors.
As a mitochondrial protein Ucp2 plays a clear role in energetics and metabolism (96)
similar to Trhr. Therefore, both candidates imply that it might be useful to perform a general
thyroid and metabolic panel on our bHR/bLR rats, as well as local measurements of energy
metabolism within the hippocampus. There is also evidence that Trhr and Ucp2 could interact
(96). However, our concurrent genetic study found little evidence of epistatic interaction
between the QTLs housing Trhr and Ucp2 on chromosomes 1 and 7 (28). This indicates that
their relevance to bHR/bLR behavioral phenotype may be largely independent, explaining a
small but meaningful portion of the difference in exploratory locomotion between the phenotypes
(<10%, approximately 200 locomotor counts, (28)) – a magnitude akin to the bHR/bLR
difference in locomotor score present in the F1 generation.
563
The top genes identified in the developmental meta-analyses suggest that bHR/bLR
differences in hippocampal structure arise early in development alongside differences in
behavior: Ncan and Bmp4
The different propensity of bHR/bLR rats towards externalizing or internalizing behavior
is evident at a young age (16, 17), and likely arises from alterations in hippocampal
development (16). Our meta-analyses encompassed three postnatal ages (P7, P14, and P21) to
provide insight into this neurodevelopmental trajectory. However, these developmental meta-
analyses depended on data derived from earlier generations and older transcriptomic platforms
and thus produced fewer significant results than the adult meta-analysis. Despite this limitation,
the top candidates from both the P7 and P14 meta-analyses are compelling, showing previous
associations with a variety of internalizing and externalizing behaviors as well as clear functional
roles in hippocampal development.
The top result from the earliest postnatal age (P7) was Neurocan (Ncan). It was the only
gene that had an effect of phenotype that survived multiple comparisons correction (FDR<0.05),
exhibiting a strikingly large effect size (bHR<bLR, ß=4.16) as early as generation F6. Similar to
previous studies, we found that Ncan was also generally upregulated during early brain
development (97, 98). As part of the extracellular matrix, Ncan modulates cell adhesion and
migration and is capable of binding growth factors (99). Ncan has been linked to bipolar disorder
(BD) in humans (97), and, similar to our bHR rats, greater externalizing behavior is observed in
Ncan knock-out mice, which display higher locomotor activity, less anxiety-like behavior,
increased risk-taking and hedonia, and amphetamine hypersensitivity (98). Within our
concurrent genetic study, Ncan was located on chromosome 16 adjacent to a variant that fully
segregated bHR/bLR rats in our colony, but this loci was not found to be a QTL for exploratory
locomotion in the bHRxbLR F2 intercross (28). It did overlap previously-identified QTLs for

despair-related behavior (62). Therefore, Ncan may contribute to aspects of the selectively-bred
bHR/bLR behavioral phenotype that are not directly captured by exploratory locomotor score as
well as to long-term structural differences within the hippocampus.
The top result within our P14 meta-analysis, Bone Morphogenetic Protein 4 (Bmp4), had
elevated expression in bLRs that did not survive multiple comparisons correction (FDR<0.15),
but this differential expression appeared to continue into adulthood (FDR<0.10), and was
evident as early as the F4 generation, and therefore seemed worthy of additional investigation.
Using additional tissue collected from two separate cohorts and six developmental time points,
we confirmed with qPCR that Bmp4 had consistently higher expression in bLRs across
postnatal development and adulthood (P2-P90), with the largest differences observed at P14.
These results are consistent with the association between Bmp4 and anxiety and depressive-
like behaviors in the literature. Blocking Bmp4 expression in mice reduces anxiety and
depressive-like behavior and increases the effectiveness of certain anti-depressant treatments
(100). Similarly, transgenic mice that are deficient for Bmp receptors have extremely low levels
of anxiety and diminished fear conditioning (101). Therefore, Bmp4 could contribute to the
propensity of bHR/bLR rats for internalizing vs. externalizing behavior.
Bmp4 is also a strong candidate for driving long-term structural change in the
hippocampus in a manner that could produce stable differences in temperament. Within our
meta-analysis results, Bmp4 was one of the leading genes driving an enrichment of bHR/bLR
effects within gene sets related to proliferation and development, and was also a central hub in
a related protein-protein interaction (PPI) network constructed using the top genes from the
adult meta-analysis (*discussed below*). Notably, the direction of effect in these gene sets
changed between P14 and adulthood – at P14, gene sets related to proliferation and
differentiation were enriched with elevated expression in bLRs, whereas by adulthood elevated
expression was seen in bHRs. This pattern fits the changing role of Bmp4 across development:
Bmp4 is important for neural induction during early development (102, 103), but later
suppresses neurogenesis (100, 104–106) and promotes other cell fates (102, 107). For
example, Bmp4 can a promote endothelial proliferation (84), which matches the elevated
endothelial-specific gene expression that we observed in the bLR hippocampus at P14. Most
importantly, Bmp signaling promotes dorsal cell type identity in the developing telencephalon
and is essential for the proper development of the dentate gyrus (101), fostering the production
of granule cell neurons during the peak period of neurogenesis. Animals lacking Bmp signaling
have a smaller dentate gyrus in adulthood and reduced fear and anxiety (101). These findings
match our results indicating an enrichment of gene expression related to the dentate gyrus in
bLR rats at P14, followed by an enrichment of gene expression normally found in the dorsal (vs.
ventral) dentate gyrus in bLRs in adulthood.
Therefore, altogether, elevated expression of Bmp4 in bLR rats during development and
adulthood has the potential to explain a surprising number of our other findings. However, it is
not clear what is driving the differential expression of Bmp4 in our bHR/bLR rats. In our
concurrent genetic study Bmp4 was not located near any of the variants segregating bHR and
bLR rats in our colony nor a QTL for exploratory locomotion (28), although it did overlap with a
previously-identified QTL for despair-related behavior (62). It is possible that a relevant variant
located close to Bmp4 was missed due to the focus on exomes in our original genetic
sequencing study or the differential expression of Bmp4 may be driven by variation within a
gene upstream in its signaling pathway.
633
Functional analyses implicate hippocampal proliferation and differentiation in bHR/bLR
phenotype
As discussed above, one of the most prominent themes amongst the top results in both
our developmental and adult meta-analyses were functions related to cell proliferation and
neuron differentiation, as revealed using both gene set enrichment analyses and co-

expression/PPI networks. Indeed, among our top meta-analysis results (FDR<0.05), we found
that the canonical Marker of Proliferation Ki-67 (Mki67) was more highly expressed in bLRs in
adulthood, similar to the upregulation observed histologically in development in earlier
generations (16) and potentially in adulthood (11, 108). bLRs also exhibited greater expression
of ubiquitin-like PHD ring finger-1 (Uhrf1), a regulator of methylation and promoter of cell
survival that is highly expressed in neural stem cells and rapidly down-regulated following
differentiation (109). These findings confirm that, at least within our model, the relationship
between internalizing behavior and cell proliferation is unlikely to be as simple as a general
stunting of growth-related processes, as might be hypothesized based on the neurotrophic
model of stress-related mood disorders (110).
Many of our top differentially-expressed genes were also important regulators of cell
fate. In addition to Bmp4, Sox9, Sox2, Hes Family BHLH Transcription Factor 5 (Hes5), Cd24,
and Tek regulate functions such as the developmental progression of neural differentiation,
gliogenesis, and endothelial proliferation (72, 80, 81, 83, 111–113). Differential expression of
genes within these pathways is clearly relevant to neurodevelopmental programming (*discussed
above*), but many of these genes were differentially expressed in adulthood – indeed, some of
the effects appeared more robust. The role of these genes in adulthood is less well-studied, but
several have been found to regulate growth and plasticity in response to neural activity and
injury (75, 114–116). Therefore, differential expression within these pathways could explain
observations from our previous morphological study, in which we found that cell differentiation
progressed differently in the adult hippocampus in bHR and bLR rats under conditions of mild
stress. Under these conditions, a higher percentage of newly-born cells in adult bLRs did not
differentiate into either neurons or astrocytes, whereas adult bHRs showed an increased rate of
differentiation into astrocytes (11). Together, these findings raise the interesting possibility that
differential expression within neurodevelopmental programming pathways could provide a
general mechanism by which environmental stimuli, such as stress or drugs, produces divergent
changes in the structure of the hippocampus in bHR and bLR animals.
666
Functional analyses implicate microglial activation and phagocytosis in bHR/bLR
phenotype
The second theme within our results was microglial function, namely phagocytosis. We
found that microglial-specific gene expression was upregulated at both P14 and adulthood in
bLRs and several top candidate genes were important regulators of microglial function. For
example, within our adult meta-analysis, bLRs exhibited greater expression of C1qa since at
least generation F4, and this expression was associated with decreased exploratory locomotor
activity. The C1q genes promote classical activation and increase microglial phagocytosis (92)
and have been implicated in phagocytosis-driven synaptic pruning (92, 117, 118).
Two of the other top candidate genes involved in microglial function were Ucp2 and Milk
fat globule-EGF factor 8 (Mfge8). Unlike the C1q family, these genes showed greater
expression in bHRs. Ucp2, discussed previously for its role in affective behavior, also has an
anti-inflammatory function through its involvement in the production of reactive oxygen species
(ROS) (63, 64, 66, 119, 120). Mfge8 is similarly associated with reduced pro-inflammatory
factors (121) as well as alternative (M2) activation of microglia (122), playing an important role
in microglial phagocytosis (122–124). Notably, within our concurrent genetic study, both Ucp2
and Mfge8 were near genetic variants that fully segregated bHR/bLR rats in our colony and
located within probable QTLs for exploratory locomotor activity (28), suggesting that genetic
variation could contribute to their differential expression in the hippocampus.
Together, this set of differentially-expressed genes would seem to suggest that bLR rats
exhibit not only basally elevated microglial-related gene expression but perhaps also an inability
to turn off acute immune response during injury or stress, which would fit pro-inflammatory
theories of anxiety and depression (125). However, within our results we found little evidence of

bHR/bLR differences in the expression of traditional inflammatory markers (e.g., Tnf, Il6, Il1b,
Il1a, Ifng) or inflammatory gene sets under basal conditions. Therefore, it seems more likely that
the large bHR/bLR differences in microglial phagocytosis genes are tied to less traditional, non-
immune roles for microglia within the brain. Given the enrichment of bHR/bLR differentially-
expressed genes within pathways related to proliferation, neural differentiation, and
development, the role of microglia and complement activation in both neurogenesis and cell
survival (126) and synaptic pruning (118, 127) seems compelling. Strikingly, these signaling
pathways are dependent on neuronal activity, with microglia preferentially phagocytosing less
active synapses (118). Therefore, in many ways microglial phagocytosis could be considered a
multi-faceted tool to tailor plasticity either during development, or in response to environmental
stimuli like stress or drugs of abuse.
701
Conclusion and Future Directions
In conclusion, by comparing across exome sequencing findings and hippocampal gene
expression studies conducted in development and adulthood spanning 43 generations of
selective breeding in our bHR/bLR colony, we implicate a diverse and compelling array of genes
whose effects may converge to promote internalizing and externalizing behavior. Moreover,
these candidate genes were enriched within two functional pathways that have the capability to
not only guide bHR and bLR rats along a different developmental trajectory, but to set the stage
for a widely-different reactivity to the environment.
Due to dependence on older platforms and exclusively male rats, we cannot claim to
have identified all relevant candidates. However, many of our findings have already inspired
promising new avenues of research in our model (128–130). First, the differentially-expressed
genes involved in proliferation, cell differentiation, and growth include many genes that play
different roles across development in association with a variety of cell types. Their full story may
only be fully understood by conducting a more detailed, cell-type specific analysis of the
developmental trajectory of the bHR/bLR hippocampus, along with direct manipulation to assess
relevance to behavioral phenotype. Second, the evidence supporting increased microglial
phagocytosis begs for experiments of a more morphological and dynamic nature, tracking
microglial activation and synaptic pruning under conditions of normal development and
adulthood and under stressful challenge. Finally, our top candidates, Trhr and Ucp2, suggest
that our model could provide insight into long-standing questions regarding the relationship
between the propensity for internalizing and externalizing behaviors and energy metabolism.
723

724 **Acknowledgements**
725
The work that was performed at the Molecular Behavioral Neuroscience Institute was
supported by the following grants: Hope for Depression Research Foundation (HDRF; RGA#
DTF Phase II (D) (McEwen- PI)), Pritzker Neuropsychiatric Disorders Research Consortium,
National Institute on Drug Abuse Grants 5P01 DA021633-02, NIDA (5RO1 DA013386), Office of
Naval Research Grant N00014-12-1-0366, Office of Naval Research (ONR) N00014-09-1-0598,
Office of Naval Research (ONR) N00014-02-1-0879, NIMH Conte Center Grant #L99MH60398,
and NIMH Program Project Grant #P01MH42251-01. The University of Alabama study was
funded by NIMH 4R00MH085859-02 (S.M.C.). Research training for I.B. was supported by the
Undergraduate Research Opportunity Program at the University of Michigan. We would also
like to acknowledge everyone involved in the studies included in the meta-analysis: James
Stewart, Angela Koelsch, Sharon Burke, Mary Hoverstein, Jennifer Fitzpatrick, Hui Li, FeiLi, and
Amy Tang, Elaine Hebda-Bauer, the University of Michigan Advanced Genomics Core, and
HudsonAlpha Institute for Biotechnology Genomics Services Lab. Finally, we thank everyone
who has provided feedback on the manuscript: Dr. Shelly
Flagel, Alek Pankonin, Pouya Mandi Gholami, Dr. Aram Parsegian, and Leah Johnson.
741

Works Cited

1. Substance Abuse and Mental Health Services Administration (2017): Mental and Substance Use Disorders. Text. Retrieved June 1, 2018, from https://www.samhsa.gov/disorders.

2. The National Institute of Mental Health Information Resource Center (2017): NIMH: Health Information > Statistics > Any Mood Disorder. . Retrieved June 1, 2018, from https://www.nimh.nih.gov/health/statistics/any-mood-disorder.shtml.

3. Krueger RF, Markon KE (2006): Reinterpreting comorbidity: a model-based approach to understanding and classifying psychopathology. *Annu Rev Clin Psychol*. 2: 111–133.

4. Smoller JW (2016): The Genetics of Stress-Related Disorders: PTSD, Depression, and Anxiety Disorders. *Neuropsychopharmacology*. 41: 297–319.

5. Wray NR, Ripke S, Mattheisen M, Trzaskowski M, Byrne EM, Abdellaoui A, *et al.* (2018): Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet*. 50: 668–681.

6. Aydin C, Frohmader K, Akil H (2015): Revealing a latent variable: individual differences in affective response to repeated injections. *Behav Neurosci*. 129: 679–682.

7. Chaudhury S, Aurbach EL, Sharma V, Blandino P, Turner CA, Watson SJ, Akil H (2014): FGF2 is a target and a trigger of epigenetic mechanisms associated with differences in emotionality: Partnership with H3K9me3. *Proc Natl Acad Sci U S A*. 111: 11834–11839.

8. Clinton SM, Vazquez DM, Kabbaj M, Kabbaj M-H, Watson SJ, Akil H (2007): Individual differences in novelty-seeking and emotional reactivity correlate with variation in maternal behavior. *Horm Behav*. 51: 655–664.

9. Flagel SB, Robinson TE, Clark JJ, Clinton SM, Watson SJ, Seeman P, *et al.* (2010): An animal model of genetic vulnerability to behavioral disinhibition and responsiveness to reward-related cues: implications for addiction. *Neuropsychopharmacology*. 35: 388–400.

10. Flagel SB, Waselus M, Clinton SM, Watson SJ, Akil H (2014): Antecedents and consequences of drug abuse in rats selectively bred for high and low response to novelty. *Neuropharmacology*. 76 Pt B: 425–436.

11. Perez JA, Clinton SM, Turner CA, Watson SJ, Akil H (2009): A New Role for FGF2 as an Endogenous Inhibitor of Anxiety. *J Neurosci*. 29: 6379–6387.

12. Prater KE, Aurbach EL, Larcinese HK, Smith TN, Turner CA, Blandino P, *et al.* (2017): Selectively Bred Rats Provide a Unique Model of Vulnerability to PTSD-Like Behavior and Respond Differentially to FGF2 Augmentation Early in Life. *Neuropsychopharmacology*. 42: 1706–1714.

13. Stead JDH, Clinton S, Neal C, Schneider J, Jama A, Miller S, *et al.* (2006): Selective breeding for divergence in novelty-seeking traits: Heritability and enrichment in spontaneous anxiety-related behaviors. *Behav Genet*. 36: 697–712.

14. Stedenfeld KA, Clinton SM, Kerman IA, Akil H, Watson SJ, Sved AF (2011): Novelty-seeking behavior predicts vulnerability in a rodent model of depression. *Physiol Behav*. 103: 210–216.

15. Turner CA, Clinton SM, Thompson RC, Watson SJ, Akil H (2011): Fibroblast growth factor-2 (FGF2) augmentation early in life alters hippocampal development and rescues the anxiety phenotype in vulnerable animals. *Proc Natl Acad Sci U S A*. 108: 8021–8025.

16. Clinton SM, Stead JDH, Miller S, Watson SJ, Akil H (2011): Developmental underpinnings of differences in rodent novelty-seeking and emotional reactivity. *Eur J Neurosci*. 34: 994–1005.

17. Turner CA, Hagenauer MH, Aurbach EL, Maras PM, Fournier CL, Blandino P, *et al.* (2019): Effects of early-life FGF2 on ultrasonic vocalizations (USVs) and the mu-opioid receptor in male Sprague-Dawley rats selectively-bred for differences in their response to novelty. *Brain Res*. 1715: 106–114.

793   18. Robinson JL, Kagan J, Reznick JS, Corley R (1992): The heritability of inhibited and
794   uninhibited behavior: A twin study. *Developmental Psychology*. 28: 1030–1037.
795   19. Turner CA, Flagel SB, Blandino P, Watson SJ, Akil H (2017): Utilizing a unique animal
796   model to better understand human temperament. *Curr Opin Behav Sci*. 14: 108–114.
797   20. Fanselow MS, Dong H-W (2010): Are the dorsal and ventral hippocampus functionally
798   distinct structures? *Neuron*. 65: 7–19.
799   21. Gray JA (1982): *The neuropsychology of anxiety: An enquiry into the functions of the septo-*
800   *hippocampal system*. The neuropsychology of anxiety: An enquiry into the functions of
801   the septo-hippocampal system. New York, NY, US: Clarendon Press/Oxford University
802   Press.
803   22. Papez JW (1937): A proposed mechanism of emotion. *Archives of Neurology & Psychiatry*.
804   38: 725–743.
805   23. Campbell S, Macqueen G (2004): The role of the hippocampus in the pathophysiology of
806   major depression. *J Psychiatry Neurosci*. 29: 417–426.
807   24. Gerlach JL, McEwen BS (1972): Rat brain binds adrenal steroid hormone: radioautography
808   of hippocampus with corticosterone. *Science*. 175: 1133–1136.
809   25. Dranovsky A, Hen R (2006): Hippocampal neurogenesis: regulation by stress and
810   antidepressants. *Biol Psychiatry*. 59: 1136–1143.
811   26. Clinton S, Miller S, Watson SJ, Akil H (2008): Prenatal stress does not alter innate novelty-
812   seeking behavioral traits, but differentially affects individual differences in
813   neuroendocrine stress responsivity. *Psychoneuroendocrinology*. 33: 162–177.
814   27. Kabbaj M, Devine DP, Savage VR, Akil H (2000): Neurobiological correlates of individual
815   differences in novelty-seeking behavior in the rat: differential expression of stress-related
816   molecules. *J Neurosci*. 20: 6983–6988.
817   28. Zhou Z, Blandino P, Yuan Q, Shen P-H, Hodgkinson CA, Virkkunen M, *et al.* (2019):
818   Exploratory locomotion, a predictor of addiction vulnerability, is oligogenic in rats
819   selected for this phenotype. *Proc Natl Acad Sci USA*. 116: 13107–13115.
820   29. Aurbach EL, Inui EG, Turner CA, Hagenauer MH, Prater KE, Li JZ, *et al.* (2015): Fibroblast
821   growth factor 9 is a novel modulator of negative affect. *Proc Natl Acad Sci USA*. 112:
822   11953–11958.
823   30. Clinton SM, Watson SJ, Akil H (2014): High novelty-seeking rats are resilient to negative
824   physiological effects of the early life stress. *Stress*. 17: 97–107.
825   31. Isgor C, Kabbaj M, Akil H, Watson SJ (2004): Delayed effects of chronic variable stress
826   during peripubertal-juvenile period on hippocampal morphology and on cognitive and
827   stress axis functions in rats. *Hippocampus*. 14: 636–648.
828   32. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP
829   (2003): Exploration, normalization, and summaries of high density oligonucleotide array
830   probe level data. *Biostatistics*. 4: 249–264.
831   33. Hagenauer MH, Schulmann A, Li JZ, Vawter MP, Walsh DM, Thompson RC, *et al.* (2018):
832   Inference of cell type content from human brain transcriptomic datasets illuminates the
833   effects of age, manner of death, dissection, and psychiatric diagnosis. *PLoS ONE*. 13:
834   e0200003.
835   34. Re AD (2014): *compute.es: Compute Effect Sizes*. . Retrieved May 23, 2018, from
836   https://CRAN.R-project.org/package=compute.es.
837   35. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, *et al.* (2012): Differential gene
838   and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks.
839   *Nat Protoc*. 7: 562–578.
840   36. Viechtbauer W (2010): Conducting Meta-Analyses in R with The metafor Package. *Journal*
841   *of Statistical Software*. 36. doi: 10.18637/jss.v036.i03.
842   37. Pollard KS, Dudoit S, Laan MJ van der (2005): Multiple Testing Procedures: the multtest
843   Package and Applications to Genomics. *Bioinformatics and Computational Biology*

Solutions Using R and Bioconductor, Statistics for Biology and Health. Springer, New
York, NY, pp 249–271.
38. Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, *et al.* (2003):
PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. *Nat Genet*. 34: 267–273.
39. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, *et al.* (2005):
Gene set enrichment analysis: a knowledge-based approach for interpreting genome-
wide expression profiles. *Proc Natl Acad Sci USA*. 102: 15545–15550.
40. Sergushichev A (2016): An algorithm for fast preranked gene set enrichment analysis using
cumulative statistic calculation. *bioRxiv*. . doi: 10.1101/060012.
41. Powell JAC (2014): GO2MSIG, an automated GO based multi-species gene set generator
for gene set enrichment analysis. *BMC Bioinformatics*. 15: 146.
42. Johnson MR, Shkura K, Langley SR, Delahaye-Duriez A, Srivastava P, Hill WD, *et al.*
(2016): Systems genetics identifies a convergent gene network for cognition and
neurodevelopmental disease. *Nat Neurosci*. 19: 223–232.
43. Park CC, Gale GD, de Jong S, Ghazalpour A, Bennett BJ, Farber CR, *et al.* (2011): Gene
networks associated with conditional fear in mice identified using a systems genetics
approach. *BMC Syst Biol*. 5: 43.
44. Cembrowski MS, Wang L, Sugino K, Shields BC, Spruston N (2016): Hipposeq: a
comprehensive RNA-seq database of gene expression in hippocampal principal
neurons. *Elife*. 5: e14997.
45. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, *et al.* (2015):
STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic
Acids Res*. 43: D447-452.
46. De Preter K, Barriot R, Speleman F, Vandesompele J, Moreau Y (2008): Positional gene
enrichment analysis of gene sets for high-resolution identification of overrepresented
chromosomal regions. *Nucleic Acids Res*. 36: e43.
47. Carlson M (2017): *org.Rn.eg.db: Genome wide annotation for Rat*. R package. Retrieved
May 23, 2018, from http://bioconductor.org/packages/org.Rn.eg.db/.
48. Shimoyama M, De Pons J, Hayman GT, Laulederkind SJF, Liu W, Nigam R, *et al.* (2015):
The Rat Genome Database 2015: genomic, phenotypic and environmental variations
and disease. *Nucleic Acids Res*. 43: D743-750.
49. Livak KJ, Schmittgen TD (2001): Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 25: 402–408.
50. Yuan JS, Reed A, Chen F, Stewart CN (2006): Statistical analysis of real-time PCR data.
BMC Bioinformatics. 7: 85.
51. Calvo N, Cecchi M, Kabbaj M, Watson SJ, Akil H (2011): Differential effects of social defeat
in rats with high and low locomotor response to novelty. *Neuroscience*. 183: 81–89.
52. Kabbaj M, Evans S, Watson SJ, Akil H (2004): The search for the neurobiological basis of
vulnerability to drug abuse: using microarrays to investigate the role of stress and
individual differences. *Neuropharmacology*. 47 Suppl 1: 111–122.
53. Albert FW, Carlborg O, Plyusnina I, Besnier F, Hedwig D, Lautenschläger S, *et al.* (2009):
Genetic architecture of tameness in a rat model of animal domestication. *Genetics*. 182:
541–554.
54. Terenina-Rigaldie E, Moisan M-P, Colas A, Beaugé F, Shah KV, Jones BC, Mormède P
(2003): Genetics of behaviour: phenotypic and molecular study of rats derived from high-
and low-alcohol consuming lines. *Pharmacogenetics*. 13: 543–554.
55. Conti LH, Jirout M, Breen L, Vanella JJ, Schork NJ, Printz MP (2004): Identification of
quantitative trait Loci for anxiety and locomotion phenotypes in rat recombinant inbred
strains. *Behav Genet*. 34: 93–103.

56. Baum AE, Solberg LC, Churchill GA, Ahmadiyeh N, Takahashi JS, Redei EE (2006): Test-
and behavior-specific genetic factors affect WKY hypoactivity in tests of emotionality.
Behav Brain Res. 169: 220–230.
57. Johannesson M, Lopez-Aumatell R, Stridh P, Diez M, Tuncel J, Blázquez G, *et al.* (2009): A
resource for the simultaneous high-resolution mapping of multiple quantitative trait loci in
rats: the NIH heterogeneous stock. *Genome Res*. 19: 150–158.
58. Fernández-Teruel A, Escorihuela RM, Gray JA, Aguilar R, Gil L, Giménez-Llort L, *et al.*
(2002): A quantitative trait locus influencing anxiety in the laboratory rat. *Genome Res*.
12: 618–626.
59. Solberg LC, Baum AE, Ahmadiyeh N, Shimomura K, Li R, Turek FW, *et al.* (2006): Genetic
analysis of the stress-responsive adrenocortical axis. *Physiol Genomics*. 27: 362–369.
60. Yamazato M, Ohya Y, Nakamoto M, Sakima A, Tagawa T, Harada Y, *et al.* (2006):
Sympathetic hyperreactivity to air-jet stress in the chromosome 1 blood pressure
quantitative trait locus congenic rats. *Am J Physiol Regul Integr Comp Physiol*. 290:
R709-714.
61. Llamas B, Contesse V, Guyonnet-Duperat V, Vaudry H, Mormède P, Moisan M-P (2005):
QTL mapping for traits associated with stress neuroendocrine reactivity in rats. *Mamm*
Genome. 16: 505–515.
62. Solberg LC, Baum AE, Ahmadiyeh N, Shimomura K, Li R, Turek FW, *et al.* (2004): Sex- and
lineage-specific inheritance of depression-like behavior in the rat. *Mamm Genome*. 15:
648–662.
63. Du R-H, Wu F-F, Lu M, Shu X-D, Ding J-H, Wu G, Hu G (2016): Uncoupling protein 2
modulation of the NLRP3 inflammasome in astrocytes and its implications in depression.
Redox Biol. 9: 178–187.
64. Gimsa U, Kanitz E, Otten W, Aheng C, Tuchscherer M, Ricquier D, *et al.* (2011): Alterations
in anxiety-like behavior following knockout of the uncoupling protein 2 (ucp2) gene in
mice. *Life Sci*. 89: 677–684.
65. Pekary AE, Sattin A, Lloyd RL (2015): Ketamine modulates TRH and TRH-like peptide
turnover in brain and peripheral tissues of male rats. *Peptides*. 69: 66–76.
66. Sun X-L, Liu Y, Dai T, Ding J-H, Hu G (2011): Uncoupling protein 2 knockout exacerbates
depression-like behaviors in mice via enhancing inflammatory response. *Neuroscience*.
192: 507–514.
67. Wang D, Zhai X, Chen P, Yang M, Zhao J, Dong J, Liu H (2014): Hippocampal UCP2 is
essential for cognition and resistance to anxiety but not required for the benefits of
exercise. *Neuroscience*. 277: 36–44.
68. Zeng H, Schimpf BA, Rohde AD, Pavlova MN, Gragerov A, Bergmann JE (2007):
Thyrotropin-releasing hormone receptor 1-deficient mice display increased depression
and anxiety-like behavior. *Mol Endocrinol*. 21: 2795–2804.
69. Xiao B, Harada Y, Kawakami K, Nabika T (2011): A 1.8-Mbp fragment on chromosome 1
affects sympathetic response to stress: evaluation in reciprocal congenic strains
between stroke-prone spontaneously hypertensive rat and Wistar-Kyoto rat. *J*
Hypertens. 29: 257–265.
70. Klimes I, Weston K, Gasperíková D, Kovács P, Kvetnanský R, Jezová D, *et al.* (2005):
Mapping of genetic determinants of the sympathoneural response to stress. *Physiol*
Genomics. 20: 183–187.
71. Potenza MN, Brodkin ES, Joe B, Luo X, Remmers EF, Wilder RL, *et al.* (2004): Genomic
regions controlling corticosterone levels in rats. *Biol Psychiatry*. 55: 634–641.
72. Belvindrah R, Rougon G, Chazal G (2002): Increased neurogenesis in adult mCD24-
deficient mice. *J Neurosci*. 22: 3594–3607.
73. Cerezo A, Guadamillas MC, Goetz JG, Sánchez-Perales S, Klein E, Assoian RK, del Pozo
MA (2009): The absence of caveolin-1 increases proliferation and anchorage-

independent growth by a Rac-dependent, Erk-independent mechanism. *Mol Cell Biol*. 29: 5046–5059.

74. Galach M (2014): Molecular mechanisms involved in epithelial differentiation of human induced pluripotent stem cells. Dissertation. doi: DOI:10.11588/heidok.00017870.

75. Head BP, Peart JN, Panneerselvam M, Yokoyama T, Pearn ML, Niesman IR, *et al.* (2010): Loss of caveolin-1 accelerates neurodegeneration and aging. *PLoS ONE*. 5: e15697.

76. Hossain MB, Shifat R, Li J, Luo X, Hess KR, Rivera-Molina Y, *et al.* (2017): TIE2 Associates with Caveolae and Regulates Caveolin-1 To Promote Their Nuclear Translocation. *Mol Cell Biol*. 37. doi: 10.1128/MCB.00142-17.

77. Jo A, Denduluri S, Zhang B, Wang Z, Yin L, Yan Z, *et al.* (2014): The versatile functions of Sox9 in development, stem cells, and human diseases. *Genes Dis*. 1: 149–161.

78. Li X-L, Liu L, Li D-D, He Y-P, Guo L-H, Sun L-P, *et al.* (2017): Integrin β4 promotes cell invasion and epithelial-mesenchymal transition through the modulation of Slug expression in hepatocellular carcinoma. *Sci Rep*. 7: 40464.

79. Liu C, Goswami M, Talley J, Chesser-Martinez PL, Lou C-H, Sater AK (2012): TAK1 promotes BMP4/Smad1 signaling via inhibition of erk MAPK: a new link in the FGF/BMP regulatory network. *Differentiation*. 83: 210–219.

80. Panaliappan TK, Wittmann W, Jidigam VK, Mercurio S, Bertolini JA, Sghari S, *et al.* (2018): Sox2 is required for olfactory pit formation and olfactory neurogenesis through BMP restriction and Hes5 upregulation. *Development*. 145. doi: 10.1242/dev.153791.

81. Poché RA, Furuta Y, Chaboissier M-C, Schedl A, Behringer RR (2008): Sox9 is expressed in mouse multipotent retinal progenitor cells and functions in Müller glial cell development. *J Comp Neurol*. 510: 237–250.

82. Shakiba N, White CA, Lipsitz YY, Yachie-Kinoshita A, Tonge PD, Hussein SMI, *et al.* (2015): CD24 tracks divergent pluripotent states in mouse and human cells. *Nat Commun*. 6: 7329.

83. Stolt CC, Lommes P, Sock E, Chaboissier M-C, Schedl A, Wegner M (2003): The Sox9 transcription factor determines glial fate choice in the developing spinal cord. *Genes Dev*. 17: 1677–1689.

84. Suzuki Y, Montagne K, Nishihara A, Watabe T, Miyazono K (2008): BMPs promote proliferation and migration of endothelial cells via stimulation of VEGF-A/VEGFR2 and angiopoietin-1/Tie2 signalling. *J Biochem*. 143: 199–206.

85. Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, *et al.* (2007): Genome-wide atlas of gene expression in the adult mouse brain. *Nature*. 445: 168–176.

86. Oldham MC, Konopka G, Iwamoto K, Langfelder P, Kato T, Horvath S, Geschwind DH (2008): Functional organization of the transcriptome in human brain. *Nat Neurosci*. 11: 1271–1282.

87. Fontanet PA, Ríos AS, Alsina FC, Paratcha G, Ledda F (2018): Pea3 Transcription Factors, Etv4 and Etv5, Are Required for Proper Hippocampal Dendrite Development and Plasticity. *Cereb Cortex*. 28: 236–249.

88. Jaworski JN, Jones DC (2006): The role of CART in the reward/reinforcing properties of psychostimulants. *Peptides*. 27: 1993–2004.

89. Young LJ, Wang Z (2004): The neurobiology of pair bonding. *Nat Neurosci*. 7: 1048–1054.

90. Higa KK, Young JW, Ji B, Nichols DE, Geyer MA, Zhou X (2017): Striatal dopamine D1 receptor suppression impairs reward-associative learning. *Behav Brain Res*. 323: 100–110.

91. Milienne-Petiot M, Groenink L, Minassian A, Young JW (2017): Blockade of dopamine D1-family receptors attenuates the mania-like hyperactive, risk-preferring, and high motivation behavioral profile of mice with low dopamine transporter levels. *J Psychopharmacol (Oxford)*. 31: 1334–1346.

92. Kettenmann H, Kirchhoff F, Verkhratsky A (2013): Microglia: new roles for the synaptic
stripper. *Neuron*. 77: 10–18.
93. Zeisel A, Hochgerner H, Lonnerberg P, Johnsson A, Memic F, Zwan J van der, *et al.* (2018):
Molecular architecture of the mouse nervous system. *bioRxiv*. 294918.
94. Fischer S, Ehlert U (2018): Hypothalamic-pituitary-thyroid (HPT) axis functioning in anxiety
disorders. A systematic review. *Depress Anxiety*. 35: 98–110.
95. Maes M, Vandewoude M, Maes L, Schotte C, Cosyns P (1989): A revised interpretation of
the TRH test results in female depressed patients. Part I: TSH responses. Effects of
severity of illness, thyroid hormones, monoamines, age, sex hormonal, corticosteroid
and nutritional state. *J Affect Disord*. 16: 203–213.
96. Lanni A, Moreno M, Lombardi A, Goglia F (2003): Thyroid hormone and uncoupling
proteins. *FEBS Lett*. 543: 5–10.
97. Cichon S, Mühleisen TW, Degenhardt FA, Mattheisen M, Miró X, Strohmaier J, *et al.* (2011):
Genome-wide association study identifies genetic variation in neurocan as a
susceptibility factor for bipolar disorder. *Am J Hum Genet*. 88: 372–381.
98. Miró X, Meier S, Dreisow ML, Frank J, Strohmaier J, Breuer R, *et al.* (2012): Studies in
humans and mice implicate neurocan in the etiology of mania. *Am J Psychiatry*. 169:
982–990.
99. Rauch U, Feng K, Zhou XH (2001): Neurocan: a brain chondroitin sulfate proteoglycan. *Cell*
Mol Life Sci. 58: 1842–1856.
100. Brooker SM, Gobeske KT, Chen J, Peng C-Y, Kessler JA (2017): Hippocampal bone
morphogenetic protein signaling mediates behavioral effects of antidepressant
treatment. *Mol Psychiatry*. 22: 910–919.
101. Caronia G, Wilcoxon J, Feldman P, Grove EA (2010): Bone morphogenetic protein
signaling in the developing telencephalon controls formation of the hippocampal dentate
gyrus and modifies fear-related behavior. *J Neurosci*. 30: 6291–6301.
102. Cole AE, Murray SS, Xiao J (2016): Bone Morphogenetic Protein 4 Signalling in Neural
Stem and Progenitor Cells during Development and after Injury. *Stem Cells Int*. 2016:
9260592.
103. Harland R (2000): Neural induction. *Curr Opin Genet Dev*. 10: 357–362.
104. Bond AM, Bhalala OG, Kessler JA (2012): The dynamic role of bone morphogenetic
proteins in neural stem cell fate and maturation. *Dev Neurobiol*. 72: 1068–1084.
105. Gobeske KT, Das S, Bonaguidi MA, Weiss C, Radulovic J, Disterhoft JF, Kessler JA
(2009): BMP signaling mediates effects of exercise on hippocampal neurogenesis and
cognition in mice. *PLoS ONE*. 4: e7506.
106. Meyers EA, Gobeske KT, Bond AM, Jarrett JC, Peng C-Y, Kessler JA (2016): Increased
bone morphogenetic protein signaling contributes to age-related declines in
neurogenesis and cognition. *Neurobiol Aging*. 38: 164–175.
107. Gomes WA, Mehler MF, Kessler JA (2003): Transgenic overexpression of BMP4 increases
astroglial and decreases oligodendroglial lineage commitment. *Dev Biol*. 255: 164–177.
108. García-Fuster MJ, Perez JA, Clinton SM, Watson SJ, Akil H (2010): Impact of cocaine on
adult hippocampal neurogenesis in an animal model of differential propensity to drug
abuse. *Eur J Neurosci*. 31: 79–89.
109. Ramesh V, Bayam E, Cernilogar FM, Bonapace IM, Schulze M, Riemenschneider MJ, *et*
al. (2016): Loss of Uhrf1 in neural stem cells leads to activation of retroviral elements
and delayed neurodegeneration. *Genes Dev*. 30: 2199–2212.
110. Duman RS, Monteggia LM (2006): A neurotrophic model for stress-related mood disorders.
Biol Psychiatry. 59: 1116–1127.
111. Bansod S, Kageyama R, Ohtsuka T (2017): Hes5 regulates the transition timing of
neurogenesis and gliogenesis in mammalian neocortical development. *Development*.
144: 3156–3167.

112. Sakai D, Suzuki T, Osumi N, Wakamatsu Y (2006): Cooperative action of Sox9, Snail2 and PKA signaling in early neural crest development. *Development*. 133: 1323–1333.

113. Sun W, Cornwell A, Li J, Peng S, Osorio MJ, Aalling N, *et al.* (2017): SOX9 Is an Astrocyte-Specific Nuclear Marker in the Adult Brain Outside the Neurogenic Regions. *J Neurosci*. 37: 4493–4507.

114. Ming G-L, Song H (2011): Adult neurogenesis in the mammalian brain: significant answers and significant questions. *Neuron*. 70: 687–702.

115. Fuller ML, DeChant AK, Rothstein B, Caprariello A, Wang R, Hall AK, Miller RH (2007): Bone morphogenetic proteins promote gliosis in demyelinating spinal cord lesions. *Ann Neurol*. 62: 288–300.

116. Yan H, Zhu X, Xie J, Zhao Y, Liu X (2016): β-amyloid increases neurocan expression through regulating Sox9 in astrocytes: A potential relationship between Sox9 and chondroitin sulfate proteoglycans in Alzheimer's disease. *Brain Res*. 1646: 377–383.

117. Schafer DP, Stevens B (2015): Microglia Function in Central Nervous System Development and Plasticity. *Cold Spring Harb Perspect Biol*. 7: a020545.

118. Wu Y, Dissing-Olesen L, MacVicar BA, Stevens B (2015): Microglia: Dynamic Mediators of Synapse Development and Plasticity. *Trends Immunol*. 36: 605–613.

119. Vogler S, Pahnke J, Rousset S, Ricquier D, Moch H, Miroux B, Ibrahim SM (2006): Uncoupling protein 2 has protective function during experimental autoimmune encephalomyelitis. *Am J Pathol*. 168: 1570–1575.

120. Arsenijevic D, Onuma H, Pecqueur C, Raimbault S, Manning BS, Miroux B, *et al.* (2000): Disruption of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species production. *Nat Genet*. 26: 435–439.

121. Albus E, Sinningen K, Winzer M, Thiele S, Baschant U, Hannemann A, *et al.* (2016): Milk Fat Globule-Epidermal Growth Factor 8 (MFG-E8) Is a Novel Anti-inflammatory Factor in Rheumatoid Arthritis in Mice and Humans. *J Bone Miner Res*. 31: 596–605.

122. Spittau B, Rilka J, Steinfath E, Zöller T, Krieglstein K (2015): TGFβ1 increases microglia-mediated engulfment of apoptotic cells via upregulation of the milk fat globule-EGF factor 8. *Glia*. 63: 142–153.

123. Fricker M, Neher JJ, Zhao J-W, Théry C, Tolkovsky AM, Brown GC (2012): MFG-E8 mediates primary phagocytosis of viable neurons during neuroinflammation. *J Neurosci*. 32: 2657–2666.

124. Fuller AD, Van Eldik LJ (2008): MFG-E8 regulates microglial phagocytosis of apoptotic neurons. *J Neuroimmune Pharmacol*. 3: 246–256.

125. Jo WK, Zhang Y, Emrich HM, Dietrich DE (2015): Glia in the cytokine-mediated onset of depression: fine tuning the immune response. *Front Cell Neurosci*. 9: 268.

126. Sato K (2015): Effects of Microglia on Neurogenesis. *Glia*. 63: 1394–1405.

127. Mastellos DC (2014): Complement emerges as a masterful regulator of CNS homeostasis, neural synaptic plasticity and cognitive function. *Exp Neurol*. 261: 469–474.

128. Maras PM, Daugherty J, Hebda-Bauer E, Watson SJ, Akil H (2018): Minocycline treatment reduces anxiety- and depressive-like behaviors in a genetic model of internalizing mood disorders. *Neuroscience Meeting Planner*. Presented at the Society for Neuroscience, San Diego, CA, p Program No. 234.03.

129. Hilde KL, Hagenauer MH, Stefanov AV, Birt I, Hebda-Bauer E, Watson SJ, Akil H (2018): Mechanisms of hippocampal development: Molecular and cellular phenotypes in the selectively bred high-responder/low-responder model of affective disorders. *Neuroscience Meeting Planner*. Presented at the Society for Neuroscience, San Diego, CA, p Program No. 233.04.

130. O'Connor AM, Pardo T, Birt I, Hagenauer MH, Maras PM, Prater KE, *et al.* (2018): The hippocampal glycome contributes to behavioural phenotype in a novel rodent model of

1096   mood disorders. *Neuroscience Meeting Planner*. Presented at the Society for
1097   Neuroscience, San Diego, CA, p Program No. 233.01.
1098   131. Clinton SM, Stead JDH, Miller S, Watson SJ, Akil H (2011): Developmental underpinnings
1099   of differences in rodent novelty-seeking and emotional reactivity. *Eur J Neurosci*. 34:
1100   994–1005.
1101   132. Cohen JL, Glover ME, Pugh PC, Fant AD, Simmons RK, Akil H, *et al.* (2015): Maternal
1102   Style Selectively Shapes Amygdalar Development and Social Behavior in Rats
1103   Genetically Prone to High Anxiety. *Dev Neurosci*. 37: 203–214.
1104

Tables and Table Legends

	MBNI_AffymetrixRae230_F4	MBNI_AffymetrixRgU34A_F6	MBNI_AffymetrixRae230_F15	MBNI_IlluminaRatRef12v1_F15	MBNI_RNASeq_F29	Alabama_Nimblegen_F34	MBNI_RNASeq_F37	MBNI_RNASeq_F43
Generation	4	6	15	15	29	34	37	43
Laboratory	MBNI	MBNI	MBNI	MBNI	MBNI	Alabama	MBNI	MBNI
Lead Scientist	Dr. John Stead	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Peter Blandino	Dr. Cigdem Aydin
Age	Adult	P7, P14, P21	P14	P14	P14, Adult	P7, P14, P21, Adult	Adult	Adult
n per group	6	6	6	6	2	5	6	5
Platform	Affymetrix Rat Expression Set 230 A	Affymetrix Rat Genome (RG) U34A GeneChips	Affymetrix Rat Expression Set 230 A	Illumina RatRef- 12v1 Beadchip	RNA-Seq	NimbleGen Rat Gene Expression 12x135	RNA-Seq	RNA-Seq
Tissue Extraction	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Dorsal Hippocampus Tissue Punch	Whole Hippocampus	Whole Hippocampus
Exposure	Basal	Basal	Basal	Basal	Basal	Basal	Basal	Vehicle Injections
Behavioral Testing	Basal	Basal	Basal	Basal	Basal	Basal	Locomotor, Anxiety	Social Interaction After Mild Stress
Data Release	TBA	GSE29552	TBA	TBA	TBA	GSE88874	TBA	TBA
Citation		Clinton *et al.* 2011				Cohen *et al.* 2015		

Table 1. An overview of the eight transcriptional profiling studies included in our current meta-analyses of differential gene expression in the bHR and bLR hippocampus at four developmental time points: P7, P14, P21, and adulthood. Citations: (131, 132)

Adult Meta-Analysis: Top Results (FDR<0.05)										
Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets	Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets
1	Tmem144	3.04E-08	4.95E-04	-3.57	3	38	Trappc6a	6.16E-05	2.61E-02	-2.06	3
2	Asb15	4.05E-07	2.32E-03	2.82	3	39	Cav1	6.47E-05	2.61E-02	1.37	5
3	Kif15	4.27E-07	2.32E-03	-2.21	4	40	Robo3	6.54E-05	2.61E-02	-2.00	3
4	Pkhd1l1	6.13E-07	2.49E-03	-3.17	3	41	C1qa	6.57E-05	2.61E-02	-1.40	5
5	Car9	7.87E-07	2.56E-03	2.78	3	42	Tubg1	6.88E-05	2.65E-02	-1.40	5
6	Klhl34	1.04E-06	2.81E-03	-2.39	4	43	Zfp952	7.17E-05	2.65E-02	2.12	2
7	Mvb12b	2.03E-06	4.32E-03	-2.95	2	44	Slc9a3r1	7.31E-05	2.65E-02	1.32	5
8	Amer2	2.12E-06	4.32E-03	2.10	3	45	Egfem1	7.45E-05	2.65E-02	-2.11	2
9	Rpl17	2.86E-06	5.17E-03	-2.16	4	46	Epb41l4a	7.49E-05	2.65E-02	-2.10	2
10	Nudt4	4.27E-06	6.51E-03	-1.58	5	47	Trmt10a	8.18E-05	2.83E-02	-1.75	3
11	Pkib	4.40E-06	6.51E-03	1.70	5	48	Oard1	8.81E-05	2.98E-02	1.65	3
12	Fxyd7	5.22E-06	6.62E-03	-1.74	4	49	Selenop	1.00E-04	3.33E-02	2.21	2
13	Scube2	5.29E-06	6.62E-03	-2.68	2	50	Ezr	1.17E-04	3.72E-02	1.27	5
14	Ddx20	5.77E-06	6.62E-03	-1.93	4	51	Acox3	1.17E-04	3.72E-02	1.26	5
15	R3hdm4	6.10E-06	6.62E-03	-2.03	3	52	Ints8	1.21E-04	3.74E-02	2.11	2
16	Hmgn5b	7.33E-06	6.94E-03	2.60	2	53	Pld4	1.22E-04	3.74E-02	-1.45	4
17	Slc39a12	7.35E-06	6.94E-03	2.62	3	54	Ucp2	1.27E-04	3.83E-02	1.29	5
18	Acss2	7.68E-06	6.94E-03	-1.87	4	55	Rbm3	1.29E-04	3.83E-02	-1.52	4
19	Rhpn2	8.16E-06	6.98E-03	2.33	3	56	Ankdd1b	1.37E-04	3.93E-02	-2.00	2
20	Exosc7	9.65E-06	7.85E-03	1.80	4	57	Aar2	1.38E-04	3.93E-02	2.02	2
21	Rnls	1.04E-05	8.06E-03	1.92	3	58	Fn3k	1.42E-04	3.97E-02	-1.47	4
22	Slc19a3	1.50E-05	1.11E-02	-2.36	3	59	Slc27a1	1.45E-04	4.01E-02	1.24	5
23	Prss55	1.91E-05	1.35E-02	-2.43	2	60	Afmid	1.52E-04	4.11E-02	1.98	2
24	Apln	2.07E-05	1.41E-02	1.48	5	61	Slc4a11	1.54E-04	4.11E-02	-1.78	3
25	Mfge8	2.70E-05	1.73E-02	1.50	5	62	Zfp110	1.64E-04	4.31E-02	1.24	5
26	C2cd3	2.84E-05	1.73E-02	2.06	3	63	Tek	1.70E-04	4.33E-02	-1.79	3
27	Dnaaf3	2.87E-05	1.73E-02	-2.51	2	64	Fmo5	1.70E-04	4.33E-02	1.42	4
28	Etv4	3.30E-05	1.92E-02	2.02	3	65	Blvra	1.74E-04	4.36E-02	1.52	4
29	Tmem2	3.72E-05	1.99E-02	-2.29	2	66	Prdm5	2.03E-04	4.94E-02	-1.76	3
30	Tdg	3.84E-05	1.99E-02	-1.38	5	67	Samd5	2.09E-04	4.94E-02	1.95	3
31	Mki67	4.03E-05	1.99E-02	-2.01	3	68	Ntn4	2.10E-04	4.94E-02	1.94	2
32	Tnnt1	4.06E-05	1.99E-02	-1.51	5	69	Chd1l	2.11E-04	4.94E-02	1.45	4
33	Zfp90	4.13E-05	1.99E-02	-1.59	4	70	Ccdc137	2.13E-04	4.94E-02	1.97	2
34	LOC363337	4.16E-05	1.99E-02	-2.82	2	71	Trhr	2.16E-04	4.94E-02	1.19	5
35	Uhrf1	4.95E-05	2.29E-02	-1.67	3	72	Zfp821	2.22E-04	4.98E-02	-1.48	3
36	Tuba8	5.06E-05	2.29E-02	-1.44	4	73	Myh6	2.23E-04	4.98E-02	-1.25	5
37	Rltpr	5.58E-05	2.45E-02	-2.28	2	74	Sp3	2.26E-04	4.98E-02	1.93	3

P7: Top Result (FDR<0.05)						P14: Top Result (FDR<0.15)				
Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets	Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets
1	Ncan	1.38E-07	4.86E-04	-4.16	2	1	Bmp4	9.40E-06	1.47E-01	-1.49	5

Table 2. The top differentially expressed genes within the bHR/bLR hippocampal gene expression meta-analyses. P-value=nominal p-value, FDR=false detection rate, Estimate=estimated effect size (i.e., the difference in expression between bHR and bLR rats in units of standard deviation, green/positive=higher expression in bHRs, red/negative= higher expression in bLRs), # of datasets=number of datasets included in the meta-analysis for that gene.

Figures and Figure Legends

Figure 1. Selectively-bred high responder (bHR) and low responder (bLR) rats model an extreme propensity for internalizing vs. externalizing behavior. A) Over the course of 56 generations of selective breeding (F1-F56), the bHR rats (green) have developed increasingly elevated exploratory activity in a novel field (*y-axis:* average total locomotor score), whereas the bLR rats (red) have become less exploratory. These trends plateaued after F42, when our breeding strategy changed to deaccelerate divergence. Arrows indicate the generations during which hippocampal transcriptomic profiling datasets were collected, along with a name indicating the respective laboratory, platform, and generation for each dataset. **B)** bLR rats have been highly anxious since the initiation of our breeding colony. The example above is from the behavioral data accompanying the MBNI_RNASeq_F37 transcriptomic dataset showing bLRs spending a smaller percentage of time in the anxiogenic open arms of the elevated plus maze than bHR rats (effect of phenotype: *p*<0.01, boxes=first quartile, median, and third quartile,

whiskers = range). **C)** bLR rats are more reactive to stressors. This example is from the behavioral data accompanying the MBNI_RNASeq_F43 transcriptomic dataset showing bLR rats spending a smaller percentage of time interacting socially following exposure to a single mild stressor ($p<0.05$).

A.

Age	Number of Genes in Meta-Analysis	Number of Datasets in Meta-Analysis	Number of Significant Genes (p-value < 0.05)	Number of Significant Genes after FDR (<0.05)
P7	3527	2	392	1
P14	15682	5	921	0
P21	3527	2	263	0
Adult	16269	5	1881	74

Figure 2. Meta-analysis uncovers hippocampal gene expression related to bHR/bLR phenotype: An overview. *A)* A table overviewing the results of the meta-analyses conducted for each of the four age groups. Genes were only included in the meta-analysis if they were represented in at least two datasets. The most differentially-expressed genes were detected within the adult meta-analysis, which predominantly included data from later generations (F29-F43). **B-G.** Volcano plots illustrating the distribution of the results for all genes included in the meta-analysis for each age group. The x-axis is the estimated effect size for each gene (i.e., the difference in expression between bHR and bLR rats in units of standard deviation, positive=higher expression in bHRs), the y-axis is the -log(10) nominal p-value from the meta-analysis (larger=more significant). These plots are colored to either illustrate: **B-C & E-F)** the number of genes in each meta-analysis with an estimated effect size greater than one (red) or with p-values surpassing false detection correction (FDR<0.05, blue), **D&G)** The number of datasets included in the meta-analysis for each gene (for the P14 and adult meta-analyses). We found that the estimated effect sizes tended to be more extreme for genes that were present in fewer datasets, and less extreme for genes present in all five datasets, most likely because many genes were only represented in the RNA-Seq datasets from later generations.

A. Top Results Are Enriched with Genes Related to Cell Proliferation, Differentiation, and Development

Top Pathways in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
protein_folding(6)	2.32	1.14	**FKBP4**,HSPA8,**CCT4**,CHORDC1,PDIA6,HSP90AA1,CLU,DNAJC10,TCP1
positive_regulation_of_cell_proliferation(4)	1.59	-1.30	SOX9,**MFGE8**,**APLN**,GPR183,**CAV1**,**CD24**,**HES5**,**HTRA1**,**GLUL**,LEP
nephron_development(4)	2.08	-0.94	**SOX9**,**CD24**,FOXD1,PTPRO,ANGPT2,SERPINB7,WNK4,SOX8,LHX1,TACSTD2
chaperone_mediated_protein_folding(7)	2.17	1.32	**FKBP4**,HSPA8,**CCT4**,CHORDC1,CLU,PDIA4,CANX,PPID,ST13,HSPD1

Top Pathways at P14 (FDR<0.05):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
generation_of_neurons(7)	0.94	-1.56	FARP1,**BMP4**,BEX1,**SOX9**,ANKRD27,SHC1,SNX3,TNC,FBXO31,TCF12
negative_regulation_of_multicellular_organismal_process(3)	1.11	-1.62	**BMP4**,BEX1,**SOX9**,LRP1,XYLT1,SRGAP2,SERPINF1,NPFF,GIMAP5,HDAC7
neurogenesis(6)	0.98	-1.51	FARP1,**BMP4**,BEX1,**SOX9**,ANKRD27,SHC1,SNX3,TNC,FBXO31,TCF12
regulation_of_neurogenesis(6)	0.78	-1.60	**BMP4**,BEX1,**SOX9**,SNX3,FBXO31,TCF12,LRP1,XYLT1,SRGAP2,SERPINF1
neuron_differentiation(6)	0.86	-1.56	FARP1,**BMP4**,BEX1,**SOX9**,ANKRD27,SHC1,SNX3,TNC,FBXO31,TCF12
negative_regulation_of_cellular_macromolecule_biosynthetic_process(6)	0.92	-1.58	**BMP4**,**SOX9**,UXT,CHD8,YBX1,GFPT1,RARA,ZEB1,HDAC7,KHDRBS1
regulation_of_neuron_differentiation(7)	-0.70	-1.59	**BMP4**,BEX1,**SOX9**,SNX3,FBXO31,TCF12,LRP1,XYLT1,SERPINF1,RARA
regulation_of_nervous_system_development(5)	0.83	-1.57	**BMP4**,BEX1,**SOX9**,SNX3,FBXO31,TCF12,LRP1,XYLT1,SRGAP2,SERPINF1

B. An Unbiased PPI-Network Constructed from the Top Results Highlights Genes Involved in Cell Proliferation and Differentiation

Up in bHR

Up in bLR

Network nodes: Cd24, Hes5, Itgb4, Cav1, Sp3, Mki67, Tek, Sox9, Bmp4, Map3k7, Sox2, Hipk4

C. Marker of Proliferation Ki-67 is More Highly Expressed in bLRs in Adulthood

bLR — bHR

Investigator & Study	Cohen's D [95% CI]
MBNI_RNASeq_F29	-0.81 [-2.85, 1.23]
MBNI_RNASeq_F37	-2.14 [-3.55, -0.73]
MBNI_RNASeq_F43	-2.67 [-4.38, -0.96]
RE Model	-2.01 [-2.97, -1.05]

Observed Outcome

D. Marker of Proliferation Ki-67 Correlates with Exploratory Activity

R-squared= 0.41 , P-value= 0.0042

Mki67: Log(2) FPM vs Locomotor Score

Legend: bHR, bIR, bLR

E. The Top Result at P14 is a Master Regulator of Development, Bmp4

bLR — bHR

Investigator & Study	Cohen's D [95% CI]
MBNI_AffymetrixRgU34A_F6	-1.12 [-2.34, 0.10]
MBNI_AffymetrixRae230_F15	-0.62 [-1.83, 0.59]
MBNI_IlluminaRatRef12v1_F15	-2.38 [-3.86, -0.90]
MBNI_RNASeq_F29	-5.28 [-9.43, -1.13]
Alabama_NimbleGen_F34	-1.97 [-3.48, -0.46]
RE Model	-1.49 [-2.15, -0.83]

Observed Outcome

F. Bmp4 Remains More Highly Expressed in bLRs in Adulthood

bLR — bHR

Investigator & Study	Cohen's D [95% CI]
MBNI_AffymetrixRae230_F4	-1.28 [-2.64, 0.08]
MBNI_RNASeq_F29	-2.47 [-5.07, 0.13]
Alabama_NimbleGen_F34	-0.82 [-2.11, 0.47]
MBNI_RNASeq_F37	-0.81 [-1.99, 0.37]
MBNI_RNASeq_F43	-0.97 [-2.28, 0.34]
RE Model	-1.04 [-1.66, -0.42]

Observed Outcome

G. Bmp4 Correlates with Anxiety

R-squared= 0.28 , P-value= 0.024

Bmp4: Log(2) FPM vs % Time in the Open Arm

Legend: bHR, bIR, bLR

Figure 3. The top bHR vs. bLR differential expression results are enriched with genes related to cell proliferation, differentiation, and development, including Bone morphogenetic protein 4 (Bmp4) and the canonical Marker of Proliferation (Mki67). A) A table of the top functional ontology gene sets identified as enriched for bHR/bLR differentially expressed genes by GSEA (FDR: False Detection Rate; NES: Normalized Enrichment Score, with positive scores (green) indicated higher expression in bHRs and negative scores (red) indicating higher expression in bLRs, bold: $p<0.05$ in GSEA results, bold+italics: FDR<0.05 in GSEA results). The top 10 "Leading Edge" genes for each gene set are shown (bold+italics: FDR<0.05 in meta-analysis). These genes have large estimated effect sizes within the meta-analysis and help drive the enrichment of effects within these gene sets. **B)** A PPI network constructed using the top genes from the adult meta-analysis (192 genes with FDR<0.10) included a dominant subnetwork that had Bmp4 and Mki67 as hub genes. Many of these genes are related to cell proliferation and differentiation within the brain. **C)** A forest plot showing that Mki67 was consistently elevated in bLR rats in adulthood (boxes=Cohen's D from each study +/- 95% confidence interval, adult: $\beta=-2.01$, $p=4.03E-05$, FDR=1.99E-02). **D)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, we found that Mki67 (units=log(2) fragments per million (FPM)) showed a negative relationship with locomotor score ($\beta=-0.000249$, $R^2= 0.41$, $p=0.0042$). **E-F)** Two forest plots showing that Bmp4 was consistently elevated in bLR rats in both development (P14) and adulthood (P14: $\beta=-1.49$, $p=9.40E-06$, FDR=1.47E-01; adult: $\beta=-1.04$, $p=1.01E-03$, FDR=9.38E-02). This direction of effect mirrors findings in the literature that show that blocking the expression of Bmp4 in mice reduces anxiety and depressive-like behavior (100). **G)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, we found that Bmp4 showed a negative relationship with percent time in the open arms ($\beta=-0.034$, $R^2= 0.28$, $p=0.024$) and a positive relationship with the number of fecal boli produced on the EPM ($\beta=0.32$, $R^2= 0.29$, $p=0.020$, *data not shown*).

Figure 4. Validation using qPCR in additional cohorts: Bmp4 is more highly expressed in bLRs across development and adulthood. We confirmed that bLRs showed greater *Bmp4* expression than bHRs at P14 and adulthood (P60, P90) as well as at other developmental time points (P2, P7, P21) using hippocampal tissue from later generations (P2=F44/F57, P7-P90= F54-F55). Log(2) fold change in *Bmp4* expression was calculated using the Livak method (-ΔΔCq, (49)), using *Gapdh* as the reference housekeeping gene and bLRs as the reference group (therefore the bLR mean is set to 0 in all panels). *p<0.05, **p<0.005 (Welch's t-test: *P2*: Log(2)FC=-1.11, T(17.0)=-2.52, p=0.0219; *P7*: Log(2)FC=-1.68, T(14.2)=-3.34, p=0.00483; *P14*: Log(2)FC=-3.74, T(5.60)=-6.10, p=0.00115; *P21*: Log(2)FC=-1.59, T(7.72)=-2.76, p=0.0257; *P60*: Log(2)FC=-2.04, T(5.99)=-3.22, p=0.0182; *P90*: T(8.74)=-6.87, p=8.44E-05). Please note that the group difference at P21 may be underestimated due to differential expression of the reference housekeeping gene, which showed higher raw expression (lower Cq) in bLRs (T(9.26)=-3.01, p=0.0143).

Figure 5. Neurocan (Ncan) has elevated expression in bLR rats at age P7. A-B) Boxplots illustrating the effect of age and bHR/bLR phenotype on Ncan expression (log(2) signal) in two microarray studies (boxes=median and interquartile range, whiskers=range, red=bLR, green=bHR). The effect of phenotype is most obvious at an age when Ncan is elevated in development (P7). **C-D)** Forest plots showing that Ncan was more expressed in bLRs (boxes=Cohen's D from each study +/-95% confidence interval) **C)** in the P7 meta-analysis (FDR<0.05), **D)** and nominally in the P14 meta-analysis (p<0.05).

A. Overlap with Our Previous Genetic Findings:

	QTLs from bHR/bLR F2 Intercross: Exploratory Locomotion		Overlapping QTLs from Rat Genome Database			
Chr	QTL range in MB* (LOD>3, +/-1MB)	LOD	Genes w/ FDR<0.1 in Meta-Analysis	Anxiety	Stress Response	Behavioral Despair
1	28-31	3.13			[59]	[62]
1	39-65	**4.75**	*Ezr*	[53]	[59]	
1	98-104	3.42	Dbp	[53],[54]	[59]	
1	115-132	**7.76**	*Hmgn5b*	[53],[54]	[59],[60]	
1	139-142	3.14	*Mfge8*	[53],[54]	[60]	
1	156-171	**4.32**	Olr35, Spcs2, *C2cd3, Ucp2*	[53],[54]	[60]	
2	43-70	**4.26**	Pdzd2, *Selenop*	[55]	[61]	
2	92-95	3.02			[61], [61]	
3	43-65	**4.71**	Ttc30a1	[55]	[59]	[62]
5	98-102	3.34		[57],[58]		
7	67-85	**5.14**	*Trhr, Pkhd1l1*	[55]	[59]	
17	157-177	3.82	Chrm3, Akr1cl			
18	71-81	**4.99**		[56]		

B. The Top Meta-Analysis Genes Often Overlap QTLs for Exploratory Locomotion

C. Top Genomic Regions Enriched with Differentially Expressed Genes:

	Genetic Loci Enriched with bHR/bLR Differentially Expressed Genes (PGE)						Overlapping QTLs:			
Chr	Location (Rnor_6.0)	P-value	FDR	Size of Region (kb)	Enrichment Score	Genes w/ FDR<0.1 in Meta-Analysis	Exploratory Locomotion	Anxiety	Stress Response	Behavioral Despair
1	2782900-4653300	7.76E-04	1.02E-02	1,870	0.6	*Samd5*				[62]
1	164225800-165684000	4.97E-05	4.68E-03	1,458	0.2	Olr35, Spcs2, *C2cd3, Ucp2*	[28]	[53], [54]	[60]	
1	174385400-174620200	1.77E-04	5.69E-03	235	1	*Scube2*		[54]	[69]	
3	62800900-63455200	7.76E-04	1.02E-02	654	0.6	Ttc30a1	[28]	[55]	[59]	[62]
3	119361600-119677400	1.22E-04	5.69E-03	316	0.6	*Blvra*		[59],[61]	[62]	
3	151032900-151150600	1.77E-04	5.69E-03	118	1	*Acss2,* Myh7b			[61]	
3	176280000-176526000	4.08E-04	7.97E-03	246	0.8					
5	155246400-155264100	1.77E-04	5.69E-03	18	1	*C1qa,* C1qc		[55]		
7	81919900-83358500	2.02E-04	6.05E-03	1,439	0.5	*Trhr*	[28]	[55]	[59]	
10	88392200-103730100	6.80E-05	4.68E-03	15,338	0.1	*Slc9a3r1* , Sox9, *Tubg1* ,Ghdc, Tmem101, Acbd4, Ddx42, *Etv4*		[56]	[70], [71]	
12	37444100-39302600	4.30E-06	2.96E-03	1,859	0.3	Eif2b1				
16	81035000-81209500	3.35E-05	4.68E-03	175	0.8				[61]	
19	32182900-42110300	2.85E-05	4.68E-03	9,927	0.1	Lcat, Ist1, Zfp821, Zfp612, *Zfp90*				

Figure 6. Many of the top genes identified by our bHR/bLR meta-analyses are located within quantitative trait loci (QTLs) for exploratory locomotion identified by our concurrent genetic study. A. Several of the top genes identified by our meta-analysis overlap quantitative trait loci (QTL) identified by our concurrent genetic study (28)**.** Our concurrent genetic study used exome sequencing to identify variants that segregated bHR/bLR rats in our colony, and then used a sampling of those segregating variants to identify QTLs for exploratory locomotor activity in a novel field using an bHR/bLR F2 intercross (28). Shown are the top genes from our meta-analyses (FDR<0.1, bold+italic=FDR<0.05) that overlap (+/-1 MB) significant (LOD>4) and putative (LOD>3) QTLs identified by this study. Also depicted is overlap with QTLs identified in the Rat Genome Database (48) for the following behaviors relevant to the bHR/bLR phenotype: anxiety (53–58), stress-related responses (59–61), and behavioral despair (62). **B.** The top meta-analysis genes (FDR<0.05) were 4.5x more likely to overlap (+/-1MB) QTLs for exploratory locomotion than other genes included in our meta-analysis (Fisher's exact test: *p*=0.0035). **C.** The top chromosomal loci enriched for bHR/bLR differentially expressed genes overlap with previously-identified QTLs relevant to externalizing and internalizing behaviors. The top chromosomal loci enriched for bHR/bLR differentially expressed genes were identified using positional gene enrichment analysis (PGE): location (full coordinates), p-value=nominal p-value, FDR=false detection rate, enrichment score= ratio of genes with p<0.01 out of all genes in the region. Also depicted is the overlap (+/-1 MB) of these enriched chromosomal loci with QTLs for exploratory locomotor activity (28), as well as with QTLs identified in the Rat Genome Database (48) for the following behaviors relevant to the

bHR/bLR phenotype: anxiety (53–56), stress-related responses (59–61, 69–71), and behavioral despair (62).

A. Hippocampal-Specific Gene Sets Enriched with bHR/bLR Differential Expression

Top Co-expression Networks in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
From mouse: "paleturquoise"	*1.94*	-1.08	**ADAMTS2**,FOXC1,FBLN1,MRC1,OMD,ANGPT2,EMP3,BGN,CD248,HSPB1
From human: "M11"	*-1.72*	*-1.84*	**FBXO31**,NPHP4,PPME1,ARHGAP39,FAM131B,PLXNA1,NCOR2,PPFIA4,DMTN,STRN4
From mouse: "darkred"	*-1.68*	0.98	MTF2,DUSP11,SNHG11,DDX26B,LRRC29,RYR3,NFKBIZ,EML5,TCF25,NRXN3
From human: "M24"	*1.54*	-0.97	**EZR**,ITSN1,IDNK,PPAP2B,FAM167A,GFPT2,**ITGB4**,PSD2,PITPNC1,PHGDH
From mouse: "sienna3"	*1.91*	-0.64	OTOF,**TRHR**,SMOC1,ZBTB7C,GLRA2,GRB10,MEIS1,GPR83,ARHGAP6
From mouse: "lightcyan"	*-1.27*	*-1.47*	**RLTPR**,ATMIN,KAT2A,JPH3,ACAD8,REV1,USE1,B9D2,RAB3D,PWWP2B

Top Co-expression Networks at P14 (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
From mouse: "lightyellow"	-1.13	*-1.56*	ZFP277,ACTA2,DMWD,SEMA7A,UQCRC1,CNNM3,ATM,NPAS2,NCAN,BRD4
From human: "M11"	*-1.72*	*-1.84*	DTX3,FBXO31,GFOD1,HYOU1,ADCY9,LRFN4,DYRK1B,PTMS,UBA1,WBSCR17
From mouse: "lightcyan"	*-1.27*	*-1.47*	H1FX,ELFN2,ZCRB1,APBA2,SLC2A13,POLK,SNUPN,ATG5,HYOU1,SNAPIN
From human: "M1"	NA	*-1.36*	EID2,CHST15,RIMBP2,EMX1,NT5C3B,ELFN2,FAM65A,SRSF7,ZDHHC22,APBA2
From mouse: "skyblue"	0.86	*-1.84*	TSC22D4,MOBP,ADSSL1,SLC35B2,OPALIN,MAG,EFHD1,GJC2,ITGB4
From human: "M7"	-1.45	*-1.70*	ARHGEF4,ST6GALNAC6,DHRS11,AGAP3,B3GAT1,ARHGEF25,LRP3,PLSCR3,LMO4,NCS1

Top Regional Markers in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
"dg_v__VS__dg_d"	*2.05*	0.88	**TRHR**,TOX3,PDZRN3,FBLN1,TSHZ1,STK32B,ZFP423,NCAPG,LCORL

Top Regional Markers at P14 (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
"dg_d-dg_v__VS__ca4-ca3_d-ca3_v-ca2-ca1_d-ca1_v"	-0.88	*-1.76*	SCN3A,ARHGAP20,MARCKS,SV2C,PDE7B,TRPC6,SLC4A4,BTG2,IGFBP5,RFX3

B. Trhr is the Top Result in a Section of Chr 7 Enriched for Differentially-Expressed Genes

Chr7: 81,919,000-83,359,000

Gene Symbol	Adult_estimate	P14_estimate
Rspo2	-1.35	NA
Emc2	1.04	NA
Trhr	*1.19*	0.35
Eny2	-1.01	0.14

C. Trhr is the Top Result Within a Hippocampal Co-expression Network Enriched for Differentially-Expressed Genes

D. Trhr is More Highly Expressed in bHR Rats in Adulthood

bLR		bHR
Investigator & Study		Cohen's D [95% CI]
MBNI_AffymetrixRae230_F4		1.15 [-0.19, 2.49]
MBNI_RNASeq_F29		1.03 [-1.05, 3.11]
Alabama_NimbleGen_F34		0.96 [-0.35, 2.27]
MBNI_RNASeq_F37		1.08 [-0.13, 2.29]
MBNI_RNASeq_F43		1.77 [0.30, 3.24]
RE Model		1.19 [0.56, 1.82]

Observed Outcome

P-value: 2.2 E-04   *FDR*: 4.9E-02

D. Trhr Correlates with Exploratory Activity

R-squared= 0.22 , P-value= 0.051

Figure 7. Thyrotropin releasing hormone receptor (Trhr) was the top gene within two hippocampal specific gene sets and within a region of Chromosome 7 enriched for bHR/bLR differential expression. A) The top hippocampal-specific gene sets identified as enriched for bHR/bLR differentially expressed genes by GSEA (FDR: False Detection Rate; NES: Normalized Enrichment Score, with positive scores (green) indicating greater expression in bHRs and negative scores (red) indicating greater expression in bLRs, bold: p<0.05 in GSEA results, bold+italics: FDR<0.05 in GSEA results). The top 10 "Leading Edge" genes for each gene set are shown (bold+italics: FDR<0.05 in meta-analysis). These genes have large estimated effect sizes within the meta-analysis and help drive the enrichment of effects within these gene sets. Note that Trhr is the strongest leading gene in two of the gene sets. **B)** Trhr was the strongest result within a segment of Chromosome 7 enriched for differentially expressed genes and containing a QTL for exploratory locomotor activity (28). The table illustrates the differentially expressed genes within this region: estimate=estimated effect size (green/positive=greater expression in bHRs), bold=p<0.05, bold+italic=FDR<0.05. **C)** The ligand for Trhr, Trh, is a hub gene within a hippocampal specific co-expression network that includes Trhr and is enriched for bHR-upregulated genes. Genes within this network with known protein-protein interactions are illustrated above. Many of these genes have documented associations with reward behavior. **D)** A forest plot showing that Trhr expression was consistently elevated in bHR rats since generation F4 (5 datasets, boxes=Cohen's D from each study +/-95% confidence interval, β=1.19, p=2.16E-04, FDR= 4.94E-02). This direction of effect mirrors findings in the literature that show Trhr-KO mice exhibit greater anxiety and depressive-like behavior (68). **E)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Trhr (units: log(2) fragments per million (FPM)) showed a trend towards a positive relationship with exploratory locomotor activity (β=4.48E-04, R^2= 0.22, p=5.08E-02).

A. C2cd3 and Ucp2 are the Top Results in a Region of Chr 1 Enriched for Differentially-Expressed Genes

B. C2cd3 is More Highly Expressed in bHRs in Adulthood

C. C2cd3 Correlates with Exploratory Activity

D. Ucp2 is More Highly Expressed in bHR Rats in Adulthood

E. Ucp2 Correlates with Anxiety

Figure 8. A region on chromosome 1 that contains a QTL for exploratory locomotor activity is enriched for bHR/bLR differential expression and contains two genes important for brain function and development, C2 Calcium Dependent Domain Containing 3 (C2cd3) and Uncoupling Protein 2 (Ucp2). A) C2cd3 and Ucp2 were the top results within a segment of Chromosome 1 enriched for differentially expressed genes and containing a QTL for exploratory locomotor activity (28). The table illustrates the differentially expressed genes within this region: estimate=estimated effect size (green/positive=higher expression in bHRs), bold=p<0.05, bold+italic=FDR<0.05. **B)** A forest plot showing that C2cd3 had higher expression in bHRs in three adult datasets included in the adult meta-analysis (boxes=Cohen's D from each study +/-95% confidence interval, effect of phenotype: β=2.06, p=2.84E-05, FDR=1.73E-02). **C)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, C2cd3 (units: log(2) fragments per million (FPM)) showed a positive relationship with exploratory locomotor activity (β= 0.000109, R^2=0.26, p=3.20E-02). **D)** A forest plot showing that Ucp2 had higher expression in bHRs in four out of the five adult datasets included in the adult meta-analysis (effect of phenotype: β=1.29, p=1.27E-04, FDR=3.83E-02). This direction of effect mirrors findings in the literature showing that Ucp2 knockout mice have higher anxiety-like behavior and lower locomotor activity, as well as greater sensitivity to stress (63, 64, 66, 67), much like our bLR rats. **E)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Ucp2 showed a trend towards a positive relationship with percent of time spent in the anxiogenic open arms of the EPM (β=0.03, R^2=0.22, p=5.16E-02).

bLRs Exhibit Overall Elevated Endothelial and Microglia Specific Gene Expression:

Microglia Activation State May Differ In bHRs and bLRs:

Figure 9. There is an overall upregulation of endothelial- and microglial-specific gene expression in bLR rats. A) A heatmap illustrating the effect of bHR/bLR phenotype on cell type specific gene expression, which often reflects overall cell type balance (33): green=upregulated in bHRs, red=upregulated in bLRs, *=p<0.05. **B-C)** Forest plots showing an upregulation of endothelial-specific (**B**) and microglial-specific (**C**) gene expression in bLRs in development (P14) and adulthood (p<0.05, boxes=Cohen's D from each study +/-95% confidence interval). **D, F)** Forest plots illustrating that two regulators of microglial activation state are differentially expressed in bHRs and bLRs (boxes=Cohen's D from each study +/-95% confidence interval). **D)** In all five adult datasets, Milk fat globule-EGF factor 8 (Mfge8), a gene that promotes alternative (M2) activation of microglia, has elevated expression in bHRs in adulthood (β=1.50, p=2.70E-05, FDR=1.73E-02). **E)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Mfge8 (units=log(2) fragments per million (FPM)) showed a positive relationship with total locomotor score (β=0.000146, R^2=0.62, p=1.10E-04) as well as the percent time in the open arms of the EPM (β=0.00603, R^2=0.28, p=2.30E-02). **F)** Complement component C1q A Chain (C1qa), a gene that promotes classical activation of microglia, had elevated expression in bLRs in five adult transcriptomic studies (C1qa: β=-1.40, p=6.57E-05, FDR=2.61E-02). The other two complement component C1q genes showed a similar trend (C1qb: β =-1.07, p=3.02E-03, FDR=1.50E-01, C1qc: β=-1.20, p=3.57E-04, FDR=5.43E-02, *data not shown*). **G)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, C1qa showed a negative relationship with total locomotor score (C1qa: β=-5.17E-04, R^2=0.29, p=2.09E-02).