Genetic liability for internalizing versus externalizing behavior manifests in the developing and adult hippocampus: Insight from a meta-analysis of transcriptional profiling studies in a selectively-bred rat model

Isabelle A. Birt[1*], Megan H. Hagenauer, Ph.D.[1*], Sarah M. Clinton, Ph.D.[2], Cigdem Aydin, Ph.D.[1], Peter Blandino, Jr., Ph.D.[1], John D. H. Stead, Ph.D.[3], Kathryn L. Hilde, Ph.D.[1], Fan Meng, Ph.D.[1], Robert C. Thompson, Ph.D.[1], Huzefa Khalil, Ph.D.[1], Alex Stefanov[1], Pamela Maras, Ph.D.[1], Zhifeng Zhou, Ph.D.[4], Elaine K. Hebda-Bauer, Ph.D.[1], David Goldman, Ph.D.[4], Stanley J. Watson, Jr., Ph.D.[1] and Huda Akil, Ph.D.[1]

[1]University of Michigan, Ann Arbor, MI, USA; [2]Virginia Tech University, Blacksburg, VA, USA; [3]Carleton University, Ottawa, Ontario, CA; [4] National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda, MD, USA

*Shared first authorship

Correspondence:

Megan Hastings Hagenauer, Ph.D.

e-mail: hagenaue@umich.edu

Molecular Behavioral Neuroscience Institute (MBNI)

205 Zina Pitcher Pl.

Ann Arbor, MI 48109

Abstract

Background: For over 16 years, we have selectively bred rats to show either high or

low levels of exploratory activity within a novel environment. These "bred High Responder"

(bHR) and "bred Low Responder" (bLR) rats serve as a model for temperamental extremes,

exhibiting large differences in many internalizing and externalizing behaviors relevant to mood

and substance abuse disorders.

Methods: Our study elucidated persistent differences in gene expression related to

bHR/bLR phenotype across development and adulthood in the hippocampus, a region critical

for emotional regulation. We meta-analyzed eight transcriptional profiling datasets (microarray,

RNA-Seq) spanning 43 generations of selective breeding (adult: *n*=46, P7: *n*=22, P14: *n*=49,

P21: *n*=21; all male). We cross-referenced these results with exome sequencing performed on

our colony to pinpoint candidates likely to mediate the effect of selective breeding on behavioral

phenotype.

Results*:* Genetic and transcriptional profiling results converged to implicate two genes

with previous associations with metabolism and mood: Thyrotropin releasing hormone receptor

and Uncoupling protein 2. Our results also highlighted bHR/bLR functional differences in the

hippocampus, including a network essential for neurodevelopmental programming, proliferation,

and differentiation, containing hub genes Bone morphogenetic protein 4 and Marker of

proliferation ki-67. Finally, we observed differential expression related to microglial activation,

which is important for synaptic pruning, including two genes within implicated chromosomal

regions: Complement C1q A chain and Milk fat globule-EGF factor 8.

Conclusion*:* These candidate genes and functional pathways have the capability to

direct bHR/bLR rats along divergent developmental trajectories and promote a widely different

reactivity to the environment.

Keywords: Anxiety, Depression, Hyperactivity, Impulsivity, Limbic, Neonatal, Postnatal, Trhr,

Ucp2, Ncan, Bmp4, Mki67, C1qa, Mfge8, Etv4

Introduction

The strong pattern of comorbidity amongst psychiatric disorders is believed to be

generated by a spectrum of latent liability (1), arising from a complex interplay of genetic risk

and environmental factors, such as stress and childhood adversity (1–3). At one end of this

spectrum are internalizing disorders, which are associated with neuroticism, anxiety, and

depression. At the other end of the spectrum are externalizing disorders, which are associated

with risk-taking and novelty-seeking, as seen in mania, substance abuse, and impulse-control

disorders (1).

We model the genetic contributions underlying both extremes of this spectrum by

selectively breeding rats that react differently to a novel environment. "Bred High Responder"

(bHR) rats are highly exploratory with a disinhibited, novelty-seeking temperament, including

hyperactivity, aggression, and drug-seeking. "Bred Low Responder" (bLR) rats are highly-

inhibited, exhibiting reduced locomotor activity and anxious and depressive-like behavior (4–13).

These behavioral propensities are robust and stable, beginning early in development (14,15)

similar to temperament in humans (16).

This highly-differentiated phenotype makes bHR/bLR rats ideal for observing the

developmental programming and adult manifestation of neurological factors underlying

internalizing and externalizing tendencies (8,11,17). This study focused on the hippocampus, a

region important for emotional regulation, behavioral inhibition (18–20), and reactivity to the

environment (18), including stress-related glucocorticoid release (21,22). In the bHR/bLR model,

we previously observed differences in hippocampal glucocorticoid receptor and growth factor

expression, histone methylation, cell proliferation, survival, and overall volume (5,9,14,23,24).

Our current study characterized hippocampal gene expression in bHR/bLR rats across

development and adulthood using a meta-analysis of eight transcriptional profiling datasets

spanning 43 generations of selective breeding. Concurrently, we discovered chromosomal

regions containing bHR/bLR segregating variants that are likely to contribute to exploratory

locomotor phenotype (25). By comparing across these studies, we identified differentially-

expressed (DE) genes situated within implicated chromosomal regions and hippocampal

functional pathways essential for mood and development (**Fig 1**). These genes are promising

candidates for mediating the influence of selective breeding on behavioral phenotype.

64   [Figure 1]

67   **Methods**

69   Full methods for the individual experiments and analyses are provided in the

70   supplement. The associated datasets have been released on GEO (**Table 1**) or FigShare

71   (https://doi.org/10.1101/774034). Analysis code (R Studio v.1.0.153, R v.3.2.2) is available at

72   *https://github.com/isabellie4/PhenotypeProject* and

73   *https://github.com/hagenaue/bHRbLR_MetaAnalysisProject*.

75   [Table 1]

The bHR/bLR Rat Colony

All experiments were approved by the local University Committee on the Use and Care

of Animals, in accordance with the National Institutes of Health Guide for the Care and Use of

Laboratory Animals.

Selective Breeding: We began selectively-breeding bHR/bLR rats in the Molecular

Behavioral Neuroscience Institute (MBNI) at the University of Michigan in 2003 (protocol: (11)).

Later, a second colony was begun at the University of Alabama-Birmingham using generation

F30 bHR/bLR rats from MBNI. Our meta-analyses use datasets derived from male bHR/bLR

rats spanning generations F4-F43. We refer to these datasets according to their respective

institution, transcriptional profiling platform, and generation (**Table 1**). *MBNI_RNASeq_F37* also

included a bHRxbLR cross ("Intermediate Responder" (bIR) rats)).

Behavioral Testing: For each generation, locomotor response to a novel environment

was assessed between P50–75 (11). For *MBNI_RNASeq_F37*, we measured anxiety-like

behavior in adulthood (bHR/bLR: P160-P167; bIR: P65-75) using the percent time spent in the

open arms of an Elevated Plus Maze (EPM; 5 min test, protocol: (26)). For

MBNI_RNASeq_F43, we measured social interaction in adulthood (P92, protocol: (4)) after 15

min exposure to the anxiogenic open arms of the EPM (protocol: (27)).

Hippocampal Gene Expression Analyses

Broad Overview of the Datasets: Our meta-analyses included eight datasets from

bHR/bLR rats aged P7, P14, P21, and adult (**Table 1**). The rats were housed in standard

conditions with minimal intervention besides behavioral testing or saline injections, and

sacrificed by rapid decapitation without anesthesia. The whole hippocampus was dissected,

except in *Alabama_Nimblegen_F34*, where dorsal hippocampal tissue punches were performed

on sliced frozen tissue (28). The extracted RNA was profiled using microarray (earlier

generations: F4, F6, F15, F34) or RNA-Seq (later generations: F29, F37, F43).

Broad Overview of the Data Preprocessing: The pre-processing steps for each study

varied according to platform, but included common steps, including re-annotation, normalization

to reduce technical variation, and quality control. Microarray data were typically summarized into

log2-transformed expression sets using Robust Multi-Array Average (RMA: (29)). Gene-level

RNA-Seq read count summaries were converted to log(2) fragments per million (FPM). When

applicable, transcript data were averaged by gene symbol to obtain a single expression value

per sample per gene.

Meta-Analyses: Within each dataset, we calculated the effect size (Cohen's d and

variance of d) of bHR/bLR phenotype on the expression data for each gene within each age

group. This output was aligned across datasets using official gene symbol. The meta-analysis

for each age group was performed using *rma.mv()* (package *metafor* (30)) and corrected for

false discovery rate (FDR; Benjamini-Hochberg method in *multtest*: (31)). Including generation

as a co-variate provided little additional insight (*generation*: all genes FDR>0.3).

Overlap with Previously-Identified Genetic Variants: Our exome sequencing study

identified bHR/bLR segregating variants (single-nucleotide polymorphisms, or SNPs), and used

a sampling of those variants to pinpoint quantitative trait loci (QTLs) for exploratory locomotion

using an bHRxbLR F2 intercross (25). In our current study, we identified all DE genes nearby

(+/- 1 MB) the segregating variants and QTL peaks (LOD>3) and determined overlap with

additional QTLs relevant to bHR/bLR behavioral phenotype from Rat Genome Database ((32),

accessed 08/08/2019, keywords: "Anxiety", "Stress", and "Despair").

Positional Gene Enrichment Analysis: To explore which genes might be either co-

regulated or in linkage disequilibrium with a causal genetic variant, we evaluated the clustering

of top bHR/bLR DE genes within chromosomal regions using Positional Gene Enrichment

analysis (*PGE*, http://silico.biotoul.fr/pge/, (33)) and the top results from the P14 and adult meta-

analyses (nominal p<0.01, removing duplicated EntrezIDs).

Gene Set Enrichment Analysis (GSEA) & Protein-Protein Interaction (PPI)

Networks*: To elucidate overall functional trends within the largest sets of results (P14 and

Adult) we used GSEA ((34,35); package *fgsea* (36)) and gene set matrix files (.gmt) containing

standard gene ontology for rats (go2msig.org; (37)), or customized hippocampal-specific gene

sets **(Table S1)** including previously-identified co-expression modules (38,39) and expression

specific to hippocampal neuronal subtypes or subregions (*Hipposeq:* (40)). We further explored

top DE genes (adult meta-analysis FDR<0.10) and implicated hippocampal gene sets using

predicted PPI networks (string-db.org (41)).

Cell Type Data Deconvolution: To interrogate the relative cell type composition of our

samples, we used the *BrainInABlender* method (42). Data for genes previously identified as

having cell type specific expression was extracted, normalized, and averaged to produce a "cell

type index". For this analysis, we excluded the small MBNI_RNASeq_F29 dataset (n=2/group).

We then performed a meta-analysis of the effects of bHR/bLR phenotype on these cell type

indices using aforementioned methods.

qPCR Validation: Hippocampal tissue from 6 bHR and 6 bLR males was collected at

ages P14 (generation F55) and P90 (generation F51; **Fig S1**). Following cDNA synthesis, Bone

morphogenetic protein 4 (Bmp4) was quantified using qPCR and custom-designed primers,

using the Livak method (43) and glyceraldehyde-3-phosphate dehydrogenase (Gapdh) as

reference. Group differences in -ΔC_q at each age were assessed using Welch's two-sample t-

test (44).

158   **Results**

Selective Breeding Amplifies the Propensity for Internalizing vs. Externalizing Behavior

The divergence of bHR/bLR exploratory activity in response to selective breeding

happened rapidly (**Fig 2A)**, implying oligogenic inheritance (25). This divergence was

accompanied by a general amplification of internalizing and externalizing tendencies

(11,14,45,46). For example, in the behavioral data accompanying our transcriptional profiling

datasets, bLRs showed more anxiety-like behavior than bHRs (**Fig 2B),** and spent less time

interacting socially following a stressful challenge (**Fig 2C**). Therefore, we expected that

examining gene expression across bHR/bLR generations would reveal a convergence of effects

within implicated chromosomal regions and pathways essential to affective behavior and

reactivity to the environment.

[FIGURE 2]

Selective Breeding for Exploratory Locomotion Alters Hippocampal Gene Expression

Between generations F4-43, we conducted eight exploratory studies transcriptionally

profiling the hippocampus of bHR/bLR rats at four ages (P7, P14, P21, and adult)**.** These small

studies individually produced few reliable results (**Figs S2-S3**). Nevertheless, a formal meta-

analysis revealed multiple genes with consistent DE across generations (**Fig 3, Table S2**).

These results can be explored interactively at

https://mbni.org/dashboard/huzefak/hrlrMetaAnalysis/.

[FIGURE 3]

Adulthood: The effect of bHR/bLR phenotype on gene expression was significant for 74

genes (FDR<0.05, out of 16,269; **Fig 3**). In general, the estimated effect sizes (β's) tended to be

more extreme for genes exclusively represented in RNA-Seq data from later generations (**Fig

S4), but due to smaller sample size, these effects were not overrepresented in the top results.

To rule out bias due to platform, we ran a meta-analysis using only recent RNA-Seq data

(F37/F43) and confirmed that similar DE genes and pathways were identified (**Fig S5**). In the

following discussion we emphasize candidate genes for which there is evidence that expression

diverged during the earliest generations.

Development: The developmental meta-analyses were more dependent on data from

earlier generations and produced less robust results**.** However, the top genes had consistent

effects across age groups. Within the P7 meta-analysis, only one gene (out of 3,257), Neurocan

(Ncan), was differentially expressed, with higher expression in bLRs than bHRs since an early

generation (**Fig 4B-C**), in a manner that nominally persisted at P14 (**Fig 4D)**. Within the P14

and P21 meta-analysis, none of the effects survived multiple comparison correction (for 15,682

and 3,257 genes, respectively). However, the top gene at P14, Bmp4, was consistently

expressed at higher levels in bLRs than bHRs within both the P14 (**Fig 5A**) and adult datasets

since the F4 generation (**Fig 5B**). We confirmed DE at these ages using qPCR (**Fig 5C**, **Fig**

S1).

203   [FIGURE 4, FIGURE 5]

Many bHR/bLR Differentially Expressed Genes Are Located Within Implicated

Chromosomal Regions

Our exome sequencing study identified bHR/bLR segregating genetic variants, and then

used a sampling of those variants to identify chromosomal regions that are likely to contribute to

exploratory locomotor phenotype (QTL peaks). These implicated chromosomal regions

overlapped extensively with QTLs relevant to internalizing/externalizing behavior (32), including

anxiety (47–52), stress response (53–55), and behavioral despair (56). In our current study,

68% of the genes DE in adulthood (FDR<0.05) were within +/-1 MB of a bHR/bLR segregating

variant, and 21% were within +/-100 kB of a highly segregating variant (Bonferonni-corrected

α=5.00E-05), a 4.7x enrichment compared to non-DE genes (**Fig 6A,** Fischer's exact test:

p=3.50E-06). DE genes were also 4.5x more likely to be located within a QTL for exploratory

locomotion (LOD>4; **Fig 6B**). This overlap included two genes previously associated with

internalizing and externalizing behaviors (Thyrotropin Releasing Hormone Receptor (Trhr),

Uncoupling Protein 2 (Ucp2), **Fig 6C,** (57–62)). These results fit our expectation that the

influence of bHR/bLR segregating genetic variants on exploratory locomotion is at least partially

mediated by effects on gene expression within the hippocampus.

223   [FIGURE 6]

Positional Gene Enrichment Analysis Specifies Narrower Chromosomal Regions

Contributing to bHR/bLR Phenotype

Positional Gene Enrichment (PGE) identified 132 chromosomal regions with a significant

enrichment (FDR<0.05) of DE genes within the P14 and adult meta-analyses. We focused on

the top regions (FDR<0.001; **Figure 6D**). We confirmed that most of these top loci (10/13) could

be identified using recent RNA-Seq data (F37/F43), ruling out bias towards regions

overrepresented on older microarray platforms. These 13 top loci were narrow chromosomal

regions (measured in KB), but overlapped a strikingly high percentage of QTLs relevant to

bHR/bLR phenotype. This overlap included three QTLs for exploratory activity (25),

encompassing two DE genes associated with internalizing and externalizing behavior (Ucp2,

Trhr; **Fig 7&8**), as well as 13% of the QTLs for anxiety (6/45, (47–50)), 21% of the QTLs for

stress-related responses (8/38, (53–55,63–65)), and 23% of the QTLs for behavioral despair

(3/13, (56)) in the Rat Genome Database (32). Therefore, these enriched loci could contain

genetic variants contributing to internalizing/externalizing aspects of the bHR/bLR behavioral

phenotype beyond exploratory locomotion.

241   [FIGURE 7, FIGURE 8]

243   **bHR/bLR Differential Expression is Enriched within Hippocampal Functional Pathways**

245   ***bHR/bLR Phenotype is Associated with Proliferation and Differentiation:*** Eight

functional ontology gene sets (out of 2,761) showed an enrichment of DE within the P14 meta-

analysis (FDR<0.05), all of which were upregulated in bLRs (**Fig 9A, Table S3, Fig S6**) and

predominantly related to neurogenesis, differentiation, and brain development. Within the adult

meta-analysis, 2 of the 4 top gene sets enriched with DE (FDR<0.1, out of 2,761 total) were

similarly related to proliferation and development, but upregulated in bHRs. This pattern was

confirmed using recent RNA-Seq data (F37/F43, **Table S3**), ruling out bias towards gene

families overrepresented on microarray platforms.

253   A PPI network constructed using the top genes from the adult meta-analysis (FDR<0.10:

254   192 genes) had a dominant subnetwork highlighting many of these same genes (**Fig 9B**),

255   including hubs Bmp4 (*discussed above*) and the canonical marker of proliferation Mki67 (**Fig**

256   **9C-E**). Literature review confirmed the PPI interactions within this subnetwork and their role in

257   proliferation and differentiation in the brain (66–78).

259   [FIGURE 9]

261   ***bHR/bLR Phenotype is Associated with the Dentate Gyrus (DG):*** Similarly, when

performing GSEA using 69 gene sets custom-designed to reflect hippocampal-specific cell

types and networks **(Table S1)**, we observed an enrichment of DE related to the DG (**Fig 8E**,

Fig S7, Table S3), the location of neural proliferation within the hippocampus. At P14, bLRs

showed an upregulation of genes with enriched expression in the DG as compared to the Cornu

Ammonis (CA) regions (40) (FDR<0.05). In adulthood, bHRs showed an upregulation of genes

with enriched expression in the ventral compared to the dorsal DG (40) (FDR<0.05), including

Trhr (FDR<0.05, *discussed above*).

bHR/bLR Phenotype is Associated with Hippocampal Co-expression Networks

Related to Synaptic Signaling: Co-expression modules can capture regionally-important cell

types and functions that remain undocumented in traditional ontology databases (79). We

observed an enrichment of bHR/bLR effects within six previously identified hippocampal co-

expression modules within the P14 meta-analysis (FDR<0.05) and five within the adult meta-

analysis (FDR<0.05, **Fig 8E**, **Fig S7, Table S3**). Two of these modules included genes within

implicated chromosomal regions.

The first was a large co-expression module (695 genes) previously identified in the

mouse hippocampus ((39), "*lightcyan*") which showed elevated expression in bLRs relative to

bHRs at P14 (FDR<0.05) and adulthood (FDR<0.10). One of the top DE genes in this module,

ETS Variant 4 (Etv4, FDR<0.05, bHR>bLR), is a transcription factor required for proper

hippocampal dendrite development (80) located within an implicated chromosomal region (**Fig**

S8). A PPI network constructed using the DE genes in this module (n=74, adult p<0.05), was

enriched with genes related to cell projections, neurons, synapses, and cation binding

(FDR<0.05).

The second was a small co-expression module (39 genes) previously identified in the

mouse hippocampus ((39), *sienna3*), which showed elevated expression in bHRs in adulthood

(FDR<0.05). The top gene in this module was Trhr (*discussed above*). A PPI network

constructed using all 39 genes in this module centered on Trhr and its ligand, thyrotropin

releasing hormone (Trh; **Fig 8G**), and included many reward-related signaling molecules,

including CART prepropeptide (Cartpt (81)), Oxytocin/Neurophysin I Prepropeptide (Oxt (82)),

and Dopamine Receptor D1a (Drd1a (83,84)).

bHR/bLR Phenotype is Associated with Microglial and Endothelial-Specific Gene

Expression: Our results suggested that bHR/bLR rats might have differences in hippocampal

cell type composition. A cell type deconvolution analysis focused on well-characterized non-

neuronal cell type categories revealed that bLRs had greater microglial-specific expression than

bHRs at P14 and adulthood (**Fig 10A,C-D**). At P14, bLRs also showed greater endothelial-

specific expression (**Fig 10B**). These effects could reflect differences in cell type composition or

activation state. Notably, two of the top DE genes that were located within implicated

chromosomal regions are regulators of microglial state: Milk fat globule-EGF factor 8 (Mfge8,

Fig 10E-H) promotes alternative (M2) activation (85), and Complement component C1q A Chain

(C1qa, **Fig 10I-L**) promotes classical activation (86). To interrogate less well-characterized

hippocampal cell types, we compared our meta-analysis results to the new mousebrain.org

database (87), and found that the top DE genes (FDR<0.10) were highly expressed in a variety

of cell types, including neuronal subcategories (**Fig S9**), mirroring the diversity of hippocampal

functions implicated in bHR/bLR phenotype.

308   [FIGURE 10]

310   **Discussion**

By selectively-breeding rats for over 16 years, we have produced a robust, genetic

model of the co-occurrence of common internalizing and externalizing behaviors. Such large

differences in behavior would be expected to be accompanied by similarly strong differences in

gene expression in affective circuitry. By performing a formal meta-analysis across small,

exploratory datasets, we provide insight into bHR/bLR differences in hippocampal gene

expression across development and adulthood. Further, by cross-referencing these results with

our concurrent genetic study (25), we pinpoint strong candidates for mediating the influence of

selective breeding on hippocampal function and internalizing/externalizing behavior.

Transcriptional profiling converges with genetic results to identify two strong candidates

for contributing to bHR/bLR behavioral phenotype: Trhr and Ucp2

Our exome sequencing study offered a first glimpse of the genetic factors contributing to

our selectively-bred phenotypes (25). However, the implicated chromosomal regions

encompass several hundred genes, making their specific effects on gene expression and

function difficult to predict. By cross-referencing these findings with our DE results, we

discovered two strong candidates: Trhr and Ucp2. These candidates are located near bHR/bLR

fully-segregating genetic variants (25) within narrow chromosomal regions highly enriched for

DE genes, and overlap QTLs for exploratory activity (25), anxiety (47–49), and stress response

(53,54). Trhr was also the top gene within a bHR-upregulated gene set associated with the

ventral DG (40), a region important for proliferation and emotional regulation (18), and was a

hub in a bHR-upregulated hippocampal network containing reward-related signaling molecules.

Trhr and Ucp2 are both important for energy metabolism and extensively linked to

internalizing and externalizing behavior (88–90). Knocking out Ucp2 produces a bLR-like

phenotype: higher anxiety-like behavior, lower locomotor activity, and reduced stress resilience

(57,58,60,61). Trhr is an important component of the hypothalamic-pituitary-thyroid (HPT) axis

and regulates anxiety and depressive-like behavior (59,62,91). In our exome-sequencing study,

the variants associated with Trhr and Ucp2 explained a moderate portion of exploratory

locomotor behavior (<10%, approximately 200 locomotor counts, (25)) – a magnitude akin to the

bHR/bLR difference in locomotor score present in the F1 generation. Altogether, this evidence

suggests that bHR/bLR segregating genetic variants are driving DE of Ucp2 and Trhr in a

manner that could meaningfully contribute to bHR/bLR differences in hippocampal function and

internalizing/externalizing behavior.

The top genes identified in the developmental meta-analyses suggest that bHR/bLR

differences in hippocampal structure arise early in development: Ncan and Bmp4

The different propensity of bHR/bLR rats towards externalizing or internalizing behavior

is evident at a young age (14,15), paralleling hippocampal development (14). Our meta-

analyses encompassed three postnatal ages (P7, P14, and P21) to provide insight into this

neurodevelopmental trajectory. These meta-analyses depended on data from earlier

generations and older transcriptional profiling platforms yet identified two strong candidates with

clear associations with internalizing/externalizing behavior and hippocampal development.

The top P7 result was Ncan, exhibiting a strikingly large effect size (bHR<bLR) as early

as generation F6. Ncan is located adjacent to a bHR/bLR segregating variant (25), overlapping

a QTL for despair-related behavior (56). As part of the extracellular matrix, Ncan is upregulated

during early brain development (92,93) and modulates cell adhesion, migration, and growth

factor binding (94). Ncan has been linked to Bipolar Disorder (92), and knocking-out Ncan

enhances locomotor activity, risk-taking, hedonia, and amphetamine hypersensitivity (93).

Therefore, lower levels of Ncan in early development in bHRs could promote externalizing

behavior as well as divergent hippocampal development.

The top P14 result was Bmp4, which was elevated in bLRs since the earliest

generations in a manner that appeared to persist into adulthood. As a regulator of development,

Bmp4 is initially important for neural induction (95,96), but later suppresses neurogenesis (97–

101) and promotes other cell fates (78,95,102). Bmp4 was an important driver in bHR/bLR-

enriched gene sets related to proliferation and differentiation at P14 and a hub in a related PPI

network constructed from top adult DE genes. In the hippocampus, Bmp signaling promotes

dorsal cell type identity and is essential for DG formation (103), matching our results indicating

bLR-enrichment of gene expression related to the DG at P14 and adulthood.

Moreover, blocking Bmp signaling can produce a bHR-like phenotype, reducing anxiety,

fear conditioning, and depressive-like behavior (98,103). Therefore, Bmp4 is a strong candidate

for driving long-term structural differences in the hippocampus capable of producing stable

differences in temperament. However, Bmp4 was not located near a bHR/bLR segregating

variant in our exome sequencing study (25), implying that impactful variation may be located in

a nearby non-coding region or within a gene upstream in Bmp4's signaling pathway.

Functional analyses implicate hippocampal proliferation and differentiation in bHR/bLR

phenotype

One of the most prominent themes in our results were functions related to cell

proliferation and differentiation. Indeed, we found that Mki67 itself contained two bHR/bLR

segregating genetic variants, and was more highly expressed in bLRs in adulthood (FDR<0.05),

matching the upregulation in bLRs observed histologically in development (14) and maybe

adulthood (9,104). These findings confirm that the relationship between internalizing behavior

and cell proliferation in our model is unlikely to be as simple as a general stunting of growth-

related processes, as suggested by the neurotrophic model of stress-related mood disorders

(105).

Many of our top DE genes were also important regulators of cell fate. Bmp4, SRY-box 9

(Sox9), SRY-box 2 (Sox2), Hes Family BHLH Transcription Factor 5 (Hes5), CD24 Molecule

(Cd24), and TEK Receptor Tyrosine Kinase (Tek) regulate functions such as the developmental

progression of neural differentiation, gliogenesis, and endothelial proliferation (66,74,75,77,106–

108). Their role in adulthood includes growth and plasticity in response to neural activity and

injury (69,109–111). Therefore, these results could explain previous morphological findings

indicating that cell differentiation progressed differently in the adult hippocampus in bHRs and

bLRs under conditions of mild stress (9). Together, these findings raise the interesting possibility

that DE within neurodevelopmental programming pathways could provide a general mechanism

by which environmental stimuli, such as stress or drugs, produces divergent changes in

hippocampal structure in bHR/bLR animals.

Functional analyses implicate microglial activation in bHR/bLR phenotype

Microglial-specific gene expression was upregulated at both P14 and adulthood in bLRs,

suggesting either an increased proportion of microglia cells or microglial activation. Several top

candidate genes were important regulators of microglial function. bLRs had greater expression

of C1qa than bHRs since the earliest generations. The C1q genes promote classical microglial

activation (86) and are implicated in phagocytosis-driven synaptic pruning (86,112,113). In

contrast, Mfge8 was more expressed in bHRs. Mfge8 is associated with reduced pro-

inflammatory factors (114) as well as alternative (M2) microglial activation (85), playing an

important role in phagocytosis (85,115,116). Ucp2, discussed previously, has anti-inflammatory

function (57,58,60,117,118) and was more highly expressed in bHRs than bLRs. Both Mfge8

and Ucp2 contain bHR/bLR segregating genetic variants within probable QTLs for exploratory

activity (25), suggesting that genetic variation could contribute to their DE in the hippocampus.

Together, these results seem to fit pro-inflammatory theories of internalizing disorders

(119). However, we found little evidence of bHR/bLR differences in the expression of traditional

inflammatory markers. Therefore, it seems more likely that bHR/bLR differences in microglial

activation genes are tied to non-immune roles for microglia within the brain, including the

regulation of neurogenesis, cell survival (120), and synaptic pruning (113,121) in response to

neuronal activity (113). Therefore, microglial phagocytosis could be serving as a multi-faceted

tool to tailor plasticity either during development, or in response to environmental stimuli like

stress or drugs of abuse.

Conclusion and Future Directions

By comparing exome sequencing findings to hippocampal differential expression

patterns during development and adulthood across many generations of selective breeding in

our bHR/bLR colony, we implicate a diverse and compelling array of genes whose effects may

converge to promote internalizing and externalizing behavior. Due to a dependence of our

results on older platforms and exclusively male rats, we cannot claim to have identified all

relevant candidates, nor have we highlighted all promising results in our text. However, we

implicate two functional pathways with the capability to guide bHRs and bLRs along a divergent

developmental trajectory and set the stage for a widely different reactivity to the environment.

These findings will inspire new avenues of research (122–124), including cell type specific

morphological analyses and the manipulation of candidate pathways to assess relevance to

behavioral and neurological phenotype in our model.

434   **Acknowledgements**

The work that was performed at the Molecular Behavioral Neuroscience Institute was

supported by the following grants: Hope for Depression Research Foundation (HDRF; RGA#

DTF Phase II (D) (McEwen- PI)), Pritzker Neuropsychiatric Disorders Research Consortium,

National Institute on Drug Abuse Grants 5P01 DA021633-02, NIDA (5RO1 DA013386), Office of

Naval Research Grant N00014-12-1-0366, Office of Naval Research (ONR) N00014-09-1-0598,

Office of Naval Research (ONR) N00014-02-1-0879, NIMH Conte Center Grant #L99MH60398,

and NIMH Program Project Grant #P01MH42251-01. The University of Alabama study was

funded by NIMH 4R00MH085859-02 (S.M.C.). Research training for I.B. was supported by the

Undergraduate Research Opportunity Program at the University of Michigan. We would also

like to acknowledge everyone involved in the studies included in the meta-analysis: James

Stewart, Angela Koelsch, Sharon Burke, Mary Hoverstein, Jennifer Fitzpatrick, Hui Li, Fei Li,

and Amy Tang, the University of Michigan Advanced Genomics Core, and HudsonAlpha

Institute for Biotechnology Genomics Services Lab. Finally, we thank everyone who has

provided feedback on the manuscript: Dr. Shelly Flagel, Alek Pankonin, Pouya Mandi Gholami,

Dr. Aram Parsegian, and Leah Johnson.

Disclosures

These results were previously published in pre-print format (*bioRxiv* 774034; doi:

https://doi.org/10.1101/774034). Several of the authors are members of the Pritzker

Neuropsychiatric Disorders Research Consortium (MHH, RCT, FM, HA, SJW), which is

supported by Pritzker Neuropsychiatric Disorders Research Fund, LLC. A shared intellectual

property agreement exists between the academic and philanthropic entities of the consortium.

The authors have declared that no competing interests exist.

Works Cited

1. Krueger RF, Markon KE (2006): Reinterpreting comorbidity: a model-based approach to understanding and classifying psychopathology. *Annu Rev Clin Psychol* 2: 111–133.

2. Smoller JW (2016): The Genetics of Stress-Related Disorders: PTSD, Depression, and Anxiety Disorders. *Neuropsychopharmacology* 41: 297–319.

3. Wray NR, Ripke S, Mattheisen M, Trzaskowski M, Byrne EM, Abdellaoui A, *et al.* (2018): Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet* 50: 668–681.

4. Aydin C, Frohmader K, Akil H (2015): Revealing a latent variable: individual differences in affective response to repeated injections. *Behav Neurosci* 129: 679–682.

5. Chaudhury S, Aurbach EL, Sharma V, Blandino P, Turner CA, Watson SJ, Akil H (2014): FGF2 is a target and a trigger of epigenetic mechanisms associated with differences in emotionality: Partnership with H3K9me3. *Proc Natl Acad Sci U S A* 111: 11834–11839.

6. Clinton SM, Vazquez DM, Kabbaj M, Kabbaj M-H, Watson SJ, Akil H (2007): Individual differences in novelty-seeking and emotional reactivity correlate with variation in maternal behavior. *Horm Behav* 51: 655–664.

7. Flagel SB, Robinson TE, Clark JJ, Clinton SM, Watson SJ, Seeman P, *et al.* (2010): An animal model of genetic vulnerability to behavioral disinhibition and responsiveness to reward-related cues: implications for addiction. *Neuropsychopharmacology* 35: 388–400.

8. Flagel SB, Waselus M, Clinton SM, Watson SJ, Akil H (2014): Antecedents and consequences of drug abuse in rats selectively bred for high and low response to novelty. *Neuropharmacology* 76 Pt B: 425–436.

9. Perez JA, Clinton SM, Turner CA, Watson SJ, Akil H (2009): A New Role for FGF2 as an Endogenous

Inhibitor of Anxiety. *J Neurosci* 29: 6379–6387.

10. Prater KE, Aurbach EL, Larcinese HK, Smith TN, Turner CA, Blandino P, *et al.* (2017): Selectively Bred

Rats Provide a Unique Model of Vulnerability to PTSD-Like Behavior and Respond Differentially

to FGF2 Augmentation Early in Life. *Neuropsychopharmacology* 42: 1706–1714.

11. Stead JDH, Clinton S, Neal C, Schneider J, Jama A, Miller S, *et al.* (2006): Selective breeding for

divergence in novelty-seeking traits: Heritability and enrichment in spontaneous anxiety-related

behaviors. *Behav Genet* 36: 697–712.

12. Stedenfeld KA, Clinton SM, Kerman IA, Akil H, Watson SJ, Sved AF (2011): Novelty-seeking behavior

predicts vulnerability in a rodent model of depression. *Physiol Behav* 103: 210–216.

13. Turner CA, Clinton SM, Thompson RC, Watson SJ, Akil H (2011): Fibroblast growth factor-2 (FGF2)

augmentation early in life alters hippocampal development and rescues the anxiety phenotype

in vulnerable animals. *Proc Natl Acad Sci U S A* 108: 8021–8025.

14. Clinton SM, Stead JDH, Miller S, Watson SJ, Akil H (2011): Developmental underpinnings of

differences in rodent novelty-seeking and emotional reactivity. *Eur J Neurosci* 34: 994–1005.

15. Turner CA, Hagenauer MH, Aurbach EL, Maras PM, Fournier CL, Blandino P, *et al.* (2019): Effects of

early-life FGF2 on ultrasonic vocalizations (USVs) and the mu-opioid receptor in male Sprague-

Dawley rats selectively-bred for differences in their response to novelty. *Brain Res* 1715: 106–

114.

16. Robinson JL, Kagan J, Reznick JS, Corley R (1992): The heritability of inhibited and uninhibited

behavior: A twin study. *Developmental Psychology* 28: 1030–1037.

17. Turner CA, Flagel SB, Blandino P, Watson SJ, Akil H (2017): Utilizing a unique animal model to better

understand human temperament. *Curr Opin Behav Sci* 14: 108–114.

18. Fanselow MS, Dong H-W (2010): Are the dorsal and ventral hippocampus functionally distinct

structures? *Neuron* 65: 7–19.

19. Gray JA (1982): *The Neuropsychology of Anxiety: An Enquiry into the Functions of the Septo-*

Hippocampal System. New York, NY, US: Clarendon Press/Oxford University Press.

20. Papez JW (1937): A proposed mechanism of emotion. *Archives of Neurology & Psychiatry* 38: 725–

743.

21. Campbell S, Macqueen G (2004): The role of the hippocampus in the pathophysiology of major

depression. *J Psychiatry Neurosci* 29: 417–426.

22. Gerlach JL, McEwen BS (1972): Rat brain binds adrenal steroid hormone: radioautography of

hippocampus with corticosterone. *Science* 175: 1133–1136.

23. Clinton S, Miller S, Watson SJ, Akil H (2008): Prenatal stress does not alter innate novelty-seeking

behavioral traits, but differentially affects individual differences in neuroendocrine stress

responsivity. *Psychoneuroendocrinology* 33: 162–177.

24. Kabbaj M, Devine DP, Savage VR, Akil H (2000): Neurobiological correlates of individual differences in

novelty-seeking behavior in the rat: differential expression of stress-related molecules. *J*

Neurosci 20: 6983–6988.

25. Zhou Z, Blandino P, Yuan Q, Shen P-H, Hodgkinson CA, Virkkunen M, *et al.* (2019): Exploratory

locomotion, a predictor of addiction vulnerability, is oligogenic in rats selected for this

phenotype. *Proc Natl Acad Sci USA* 116: 13107–13115.

26. Aurbach EL, Inui EG, Turner CA, Hagenauer MH, Prater KE, Li JZ, *et al.* (2015): Fibroblast growth

factor 9 is a novel modulator of negative affect. *Proc Natl Acad Sci USA* 112: 11953–11958.

27. Isgor C, Kabbaj M, Akil H, Watson SJ (2004): Delayed effects of chronic variable stress during

peripubertal-juvenile period on hippocampal morphology and on cognitive and stress axis

functions in rats. *Hippocampus* 14: 636–648.

28. Cohen JL, Glover ME, Pugh PC, Fant AD, Simmons RK, Akil H, *et al.* (2015): Maternal Style Selectively

533   Shapes Amygdalar Development and Social Behavior in Rats Genetically Prone to High Anxiety.

534   *Dev Neurosci* 37: 203–214.

535   29. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP (2003):

536   Exploration, normalization, and summaries of high density oligonucleotide array probe level

537   data. *Biostatistics* 4: 249–264.

538   30. Viechtbauer W (2010): Conducting Meta-Analyses in R with The metafor Package. *Journal of*

539   *Statistical Software* 36. https://doi.org/10.18637/jss.v036.i03

540   31. Pollard KS, Dudoit S, Laan MJ van der (2005): Multiple Testing Procedures: the multtest Package and

541   Applications to Genomics. *Bioinformatics and Computational Biology Solutions Using R and*

542   *Bioconductor*. Springer, New York, NY, pp 249–271.

543   32. Shimoyama M, De Pons J, Hayman GT, Laulederkind SJF, Liu W, Nigam R, *et al.* (2015): The Rat

544   Genome Database 2015: genomic, phenotypic and environmental variations and disease.

545   *Nucleic Acids Res* 43: D743-750.

546   33. De Preter K, Barriot R, Speleman F, Vandesompele J, Moreau Y (2008): Positional gene enrichment

547   analysis of gene sets for high-resolution identification of overrepresented chromosomal regions.

548   *Nucleic Acids Res* 36: e43.

549   34. Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, *et al.* (2003): PGC-1alpha-

550   responsive genes involved in oxidative phosphorylation are coordinately downregulated in

551   human diabetes. *Nat Genet* 34: 267–273.

552   35. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, *et al.* (2005): Gene set

553   enrichment analysis: a knowledge-based approach for interpreting genome-wide expression

554   profiles. *Proc Natl Acad Sci USA* 102: 15545–15550.

36. Sergushichev A (2016): An algorithm for fast preranked gene set enrichment analysis using

cumulative statistic calculation. *bioRxiv*. https://doi.org/10.1101/060012

37. Powell JAC (2014): GO2MSIG, an automated GO based multi-species gene set generator for gene set

enrichment analysis. *BMC Bioinformatics* 15: 146.

38. Johnson MR, Shkura K, Langley SR, Delahaye-Duriez A, Srivastava P, Hill WD, *et al.* (2016): Systems

genetics identifies a convergent gene network for cognition and neurodevelopmental disease.

Nat Neurosci 19: 223–232.

39. Park CC, Gale GD, de Jong S, Ghazalpour A, Bennett BJ, Farber CR, *et al.* (2011): Gene networks

associated with conditional fear in mice identified using a systems genetics approach. *BMC Syst*

Biol 5: 43.

40. Cembrowski MS, Wang L, Sugino K, Shields BC, Spruston N (2016): Hipposeq: a comprehensive RNA-

seq database of gene expression in hippocampal principal neurons. *Elife* 5: e14997.

41. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, *et al.* (2015): STRING v10:

protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 43:

D447-452.

42. Hagenauer MH, Schulmann A, Li JZ, Vawter MP, Walsh DM, Thompson RC, *et al.* (2018): Inference of

cell type content from human brain transcriptomic datasets illuminates the effects of age,

manner of death, dissection, and psychiatric diagnosis. *PLoS ONE* 13: e0200003.

43. Livak KJ, Schmittgen TD (2001): Analysis of relative gene expression data using real-time quantitative

PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402–408.

44. Yuan JS, Reed A, Chen F, Stewart CN (2006): Statistical analysis of real-time PCR data. *BMC*

Bioinformatics 7: 85.

45. Calvo N, Cecchi M, Kabbaj M, Watson SJ, Akil H (2011): Differential effects of social defeat in rats

with high and low locomotor response to novelty. *Neuroscience* 183: 81–89.

46. Kabbaj M, Evans S, Watson SJ, Akil H (2004): The search for the neurobiological basis of vulnerability

to drug abuse: using microarrays to investigate the role of stress and individual differences.

Neuropharmacology 47 Suppl 1: 111–122.

47. Albert FW, Carlborg O, Plyusnina I, Besnier F, Hedwig D, Lautenschläger S, *et al.* (2009): Genetic

architecture of tameness in a rat model of animal domestication. *Genetics* 182: 541–554.

48. Terenina-Rigaldie E, Moisan M-P, Colas A, Beaugé F, Shah KV, Jones BC, Mormède P (2003): Genetics

of behaviour: phenotypic and molecular study of rats derived from high- and low-alcohol

consuming lines. *Pharmacogenetics* 13: 543–554.

49. Conti LH, Jirout M, Breen L, Vanella JJ, Schork NJ, Printz MP (2004): Identification of quantitative trait

Loci for anxiety and locomotion phenotypes in rat recombinant inbred strains. *Behav Genet* 34:

93–103.

50. Baum AE, Solberg LC, Churchill GA, Ahmadiyeh N, Takahashi JS, Redei EE (2006): Test- and behavior-

specific genetic factors affect WKY hypoactivity in tests of emotionality. *Behav Brain Res* 169:

220–230.

51. Johannesson M, Lopez-Aumatell R, Stridh P, Diez M, Tuncel J, Blázquez G, *et al.* (2009): A resource

for the simultaneous high-resolution mapping of multiple quantitative trait loci in rats: the NIH

heterogeneous stock. *Genome Res* 19: 150–158.

52. Fernández-Teruel A, Escorihuela RM, Gray JA, Aguilar R, Gil L, Giménez-Llort L, *et al.* (2002): A

quantitative trait locus influencing anxiety in the laboratory rat. *Genome Res* 12: 618–626.

53. Solberg LC, Baum AE, Ahmadiyeh N, Shimomura K, Li R, Turek FW, *et al.* (2006): Genetic analysis of

the stress-responsive adrenocortical axis. *Physiol Genomics* 27: 362–369.

54. Yamazato M, Ohya Y, Nakamoto M, Sakima A, Tagawa T, Harada Y, *et al.* (2006): Sympathetic

hyperreactivity to air-jet stress in the chromosome 1 blood pressure quantitative trait locus

congenic rats. *Am J Physiol Regul Integr Comp Physiol* 290: R709-714.

55. Llamas B, Contesse V, Guyonnet-Duperat V, Vaudry H, Mormède P, Moisan M-P (2005): QTL mapping

604   for traits associated with stress neuroendocrine reactivity in rats. *Mamm Genome* 16: 505–515.

605   56. Solberg LC, Baum AE, Ahmadiyeh N, Shimomura K, Li R, Turek FW, *et al.* (2004): Sex- and lineage-

606   specific inheritance of depression-like behavior in the rat. *Mamm Genome* 15: 648–662.

607   57. Du R-H, Wu F-F, Lu M, Shu X-D, Ding J-H, Wu G, Hu G (2016): Uncoupling protein 2 modulation of the

608   NLRP3 inflammasome in astrocytes and its implications in depression. *Redox Biol* 9: 178–187.

609   58. Gimsa U, Kanitz E, Otten W, Aheng C, Tuchscherer M, Ricquier D, *et al.* (2011): Alterations in anxiety-

610   like behavior following knockout of the uncoupling protein 2 (ucp2) gene in mice. *Life Sci* 89:

611   677–684.

612   59. Pekary AE, Sattin A, Lloyd RL (2015): Ketamine modulates TRH and TRH-like peptide turnover in brain

613   and peripheral tissues of male rats. *Peptides* 69: 66–76.

614   60. Sun X-L, Liu Y, Dai T, Ding J-H, Hu G (2011): Uncoupling protein 2 knockout exacerbates depression-

615   like behaviors in mice via enhancing inflammatory response. *Neuroscience* 192: 507–514.

616   61. Wang D, Zhai X, Chen P, Yang M, Zhao J, Dong J, Liu H (2014): Hippocampal UCP2 is essential for

617   cognition and resistance to anxiety but not required for the benefits of exercise. *Neuroscience*

618   277: 36–44.

619   62. Zeng H, Schimpf BA, Rohde AD, Pavlova MN, Gragerov A, Bergmann JE (2007): Thyrotropin-releasing

620   hormone receptor 1-deficient mice display increased depression and anxiety-like behavior. *Mol*

621   *Endocrinol* 21: 2795–2804.

622   63. Xiao B, Harada Y, Kawakami K, Nabika T (2011): A 1.8-Mbp fragment on chromosome 1 affects

623   sympathetic response to stress: evaluation in reciprocal congenic strains between stroke-prone

624   spontaneously hypertensive rat and Wistar-Kyoto rat. *J Hypertens* 29: 257–265.

625   64. Klimes I, Weston K, Gasperíková D, Kovács P, Kvetnanský R, Jezová D, *et al.* (2005): Mapping of

626   genetic determinants of the sympathoneural response to stress. *Physiol Genomics* 20: 183–187.

65. Potenza MN, Brodkin ES, Joe B, Luo X, Remmers EF, Wilder RL, *et al.* (2004): Genomic regions

controlling corticosterone levels in rats. *Biol Psychiatry* 55: 634–641.

66. Belvindrah R, Rougon G, Chazal G (2002): Increased neurogenesis in adult mCD24-deficient mice. *J

Neurosci* 22: 3594–3607.

67. Cerezo A, Guadamillas MC, Goetz JG, Sánchez-Perales S, Klein E, Assoian RK, del Pozo MA (2009): The

absence of caveolin-1 increases proliferation and anchorage- independent growth by a Rac-

dependent, Erk-independent mechanism. *Mol Cell Biol* 29: 5046–5059.

68. Galach M (2014): Molecular mechanisms involved in epithelial differentiation of human induced

pluripotent stem cells [Dissertation]. https://doi.org/DOI:10.11588/heidok.00017870

69. Head BP, Peart JN, Panneerselvam M, Yokoyama T, Pearn ML, Niesman IR, *et al.* (2010): Loss of

caveolin-1 accelerates neurodegeneration and aging. *PLoS ONE* 5: e15697.

70. Hossain MB, Shifat R, Li J, Luo X, Hess KR, Rivera-Molina Y, *et al.* (2017): TIE2 Associates with

Caveolae and Regulates Caveolin-1 To Promote Their Nuclear Translocation. *Mol Cell Biol* 37.

https://doi.org/10.1128/MCB.00142-17

71. Jo A, Denduluri S, Zhang B, Wang Z, Yin L, Yan Z, *et al.* (2014): The versatile functions of Sox9 in

development, stem cells, and human diseases. *Genes Dis* 1: 149–161.

72. Li X-L, Liu L, Li D-D, He Y-P, Guo L-H, Sun L-P, *et al.* (2017): Integrin β4 promotes cell invasion and

epithelial-mesenchymal transition through the modulation of Slug expression in hepatocellular

carcinoma. *Sci Rep* 7: 40464.

73. Liu C, Goswami M, Talley J, Chesser-Martinez PL, Lou C-H, Sater AK (2012): TAK1 promotes

BMP4/Smad1 signaling via inhibition of erk MAPK: a new link in the FGF/BMP regulatory

network. *Differentiation* 83: 210–219.

74. Panaliappan TK, Wittmann W, Jidigam VK, Mercurio S, Bertolini JA, Sghari S, *et al.* (2018): Sox2 is

required for olfactory pit formation and olfactory neurogenesis through BMP restriction and

Hes5 upregulation. *Development* 145. https://doi.org/10.1242/dev.153791

75. Poché RA, Furuta Y, Chaboissier M-C, Schedl A, Behringer RR (2008): Sox9 is expressed in mouse

multipotent retinal progenitor cells and functions in Müller glial cell development. *J Comp*

Neurol 510: 237–250.

76. Shakiba N, White CA, Lipsitz YY, Yachie-Kinoshita A, Tonge PD, Hussein SMI, *et al.* (2015): CD24 tracks

divergent pluripotent states in mouse and human cells. *Nat Commun* 6: 7329.

77. Stolt CC, Lommes P, Sock E, Chaboissier M-C, Schedl A, Wegner M (2003): The Sox9 transcription

factor determines glial fate choice in the developing spinal cord. *Genes Dev* 17: 1677–1689.

78. Suzuki Y, Montagne K, Nishihara A, Watabe T, Miyazono K (2008): BMPs promote proliferation and

migration of endothelial cells via stimulation of VEGF-A/VEGFR2 and angiopoietin-1/Tie2

signalling. *J Biochem* 143: 199–206.

79. Oldham MC, Konopka G, Iwamoto K, Langfelder P, Kato T, Horvath S, Geschwind DH (2008):

Functional organization of the transcriptome in human brain. *Nat Neurosci* 11: 1271–1282.

80. Fontanet PA, Ríos AS, Alsina FC, Paratcha G, Ledda F (2018): Pea3 Transcription Factors, Etv4 and

Etv5, Are Required for Proper Hippocampal Dendrite Development and Plasticity. *Cereb Cortex*

28: 236–249.

81. Jaworski JN, Jones DC (2006): The role of CART in the reward/reinforcing properties of

psychostimulants. *Peptides* 27: 1993–2004.

82. Young LJ, Wang Z (2004): The neurobiology of pair bonding. *Nat Neurosci* 7: 1048–1054.

83. Higa KK, Young JW, Ji B, Nichols DE, Geyer MA, Zhou X (2017): Striatal dopamine D1 receptor

suppression impairs reward-associative learning. *Behav Brain Res* 323: 100–110.

84. Milienne-Petiot M, Groenink L, Minassian A, Young JW (2017): Blockade of dopamine D1-family

receptors attenuates the mania-like hyperactive, risk-preferring, and high motivation behavioral

profile of mice with low dopamine transporter levels. *J Psychopharmacol (Oxford)* 31: 1334–

1346.

85. Spittau B, Rilka J, Steinfath E, Zöller T, Krieglstein K (2015): TGFβ1 increases microglia-mediated

engulfment of apoptotic cells via upregulation of the milk fat globule-EGF factor 8. *Glia* 63: 142–

153.

86. Kettenmann H, Kirchhoff F, Verkhratsky A (2013): Microglia: new roles for the synaptic stripper.

Neuron 77: 10–18.

87. Zeisel A, Hochgerner H, Lonnerberg P, Johnsson A, Memic F, Zwan J van der, *et al.* (2018): Molecular

architecture of the mouse nervous system. *bioRxiv* 294918.

88. Fischer S, Ehlert U (2018): Hypothalamic-pituitary-thyroid (HPT) axis functioning in anxiety disorders.

A systematic review. *Depress Anxiety* 35: 98–110.

89. Maes M, Vandewoude M, Maes L, Schotte C, Cosyns P (1989): A revised interpretation of the TRH

test results in female depressed patients. Part I: TSH responses. Effects of severity of illness,

thyroid hormones, monoamines, age, sex hormonal, corticosteroid and nutritional state. *J Affect*

Disord 16: 203–213.

90. Lanni A, Moreno M, Lombardi A, Goglia F (2003): Thyroid hormone and uncoupling proteins. *FEBS*

Lett 543: 5–10.

91. Choi J, Kim J, Kim T-K, Park J-Y, Lee J-E, Kim H, *et al.* (2015): TRH and TRH receptor system in the

basolateral amygdala mediate stress-induced depression-like behaviors. *Neuropharmacology*

97: 346–356.

92. Cichon S, Mühleisen TW, Degenhardt FA, Mattheisen M, Miró X, Strohmaier J, *et al.* (2011): Genome-

wide association study identifies genetic variation in neurocan as a susceptibility factor for

bipolar disorder. *Am J Hum Genet* 88: 372–381.

93. Miró X, Meier S, Dreisow ML, Frank J, Strohmaier J, Breuer R, *et al.* (2012): Studies in humans and

mice implicate neurocan in the etiology of mania. *Am J Psychiatry* 169: 982–990.

94. Rauch U, Feng K, Zhou XH (2001): Neurocan: a brain chondroitin sulfate proteoglycan. *Cell Mol Life*

Sci 58: 1842–1856.

95. Cole AE, Murray SS, Xiao J (2016): Bone Morphogenetic Protein 4 Signalling in Neural Stem and

Progenitor Cells during Development and after Injury. *Stem Cells Int* 2016: 9260592.

96. Harland R (2000): Neural induction. *Curr Opin Genet Dev* 10: 357–362.

97. Bond AM, Bhalala OG, Kessler JA (2012): The dynamic role of bone morphogenetic proteins in neural

stem cell fate and maturation. *Dev Neurobiol* 72: 1068–1084.

98. Brooker SM, Gobeske KT, Chen J, Peng C-Y, Kessler JA (2017): Hippocampal bone morphogenetic

protein signaling mediates behavioral effects of antidepressant treatment. *Mol Psychiatry* 22:

910–919.

99. Gobeske KT, Das S, Bonaguidi MA, Weiss C, Radulovic J, Disterhoft JF, Kessler JA (2009): BMP

signaling mediates effects of exercise on hippocampal neurogenesis and cognition in mice. *PLoS*

ONE 4: e7506.

100. Meyers EA, Gobeske KT, Bond AM, Jarrett JC, Peng C-Y, Kessler JA (2016): Increased bone

morphogenetic protein signaling contributes to age-related declines in neurogenesis and

cognition. *Neurobiol Aging* 38: 164–175.

101. Mira H, Andreu Z, Suh H, Lie DC, Jessberger S, Consiglio A, *et al.* (2010): Signaling through BMPR-IA

regulates quiescence and long-term activity of neural stem cells in the adult hippocampus. *Cell*

Stem Cell 7: 78–89.

102. Gomes WA, Mehler MF, Kessler JA (2003): Transgenic overexpression of BMP4 increases astroglial

and decreases oligodendroglial lineage commitment. *Dev Biol* 255: 164–177.

103. Caronia G, Wilcoxon J, Feldman P, Grove EA (2010): Bone morphogenetic protein signaling in the

developing telencephalon controls formation of the hippocampal dentate gyrus and modifies

fear-related behavior. *J Neurosci* 30: 6291–6301.

104. García-Fuster MJ, Perez JA, Clinton SM, Watson SJ, Akil H (2010): Impact of cocaine on adult

hippocampal neurogenesis in an animal model of differential propensity to drug abuse. *Eur J*

Neurosci 31: 79–89.

105. Duman RS, Monteggia LM (2006): A neurotrophic model for stress-related mood disorders. *Biol*

Psychiatry 59: 1116–1127.

106. Bansod S, Kageyama R, Ohtsuka T (2017): Hes5 regulates the transition timing of neurogenesis and

gliogenesis in mammalian neocortical development. *Development* 144: 3156–3167.

107. Sakai D, Suzuki T, Osumi N, Wakamatsu Y (2006): Cooperative action of Sox9, Snail2 and PKA

signaling in early neural crest development. *Development* 133: 1323–1333.

108. Sun W, Cornwell A, Li J, Peng S, Osorio MJ, Aalling N, *et al.* (2017): SOX9 Is an Astrocyte-Specific

Nuclear Marker in the Adult Brain Outside the Neurogenic Regions. *J Neurosci* 37: 4493–4507.

109. Ming G-L, Song H (2011): Adult neurogenesis in the mammalian brain: significant answers and

significant questions. *Neuron* 70: 687–702.

110. Fuller ML, DeChant AK, Rothstein B, Caprariello A, Wang R, Hall AK, Miller RH (2007): Bone

morphogenetic proteins promote gliosis in demyelinating spinal cord lesions. *Ann Neurol* 62:

288–300.

111. Yan H, Zhu X, Xie J, Zhao Y, Liu X (2016): β-amyloid increases neurocan expression through

regulating Sox9 in astrocytes: A potential relationship between Sox9 and chondroitin sulfate

proteoglycans in Alzheimer's disease. *Brain Res* 1646: 377–383.

112. Schafer DP, Stevens B (2015): Microglia Function in Central Nervous System Development and

Plasticity. *Cold Spring Harb Perspect Biol* 7: a020545.

113. Wu Y, Dissing-Olesen L, MacVicar BA, Stevens B (2015): Microglia: Dynamic Mediators of Synapse

Development and Plasticity. *Trends Immunol* 36: 605–613.

114. Albus E, Sinningen K, Winzer M, Thiele S, Baschant U, Hannemann A, *et al.* (2016): Milk Fat Globule-

Epidermal Growth Factor 8 (MFG-E8) Is a Novel Anti-inflammatory Factor in Rheumatoid

Arthritis in Mice and Humans. *J Bone Miner Res* 31: 596–605.

115. Fricker M, Neher JJ, Zhao J-W, Théry C, Tolkovsky AM, Brown GC (2012): MFG-E8 mediates primary

phagocytosis of viable neurons during neuroinflammation. *J Neurosci* 32: 2657–2666.

116. Fuller AD, Van Eldik LJ (2008): MFG-E8 regulates microglial phagocytosis of apoptotic neurons. *J

Neuroimmune Pharmacol* 3: 246–256.

117. Vogler S, Pahnke J, Rousset S, Ricquier D, Moch H, Miroux B, Ibrahim SM (2006): Uncoupling protein

2 has protective function during experimental autoimmune encephalomyelitis. *Am J Pathol* 168:

1570–1575.

118. Arsenijevic D, Onuma H, Pecqueur C, Raimbault S, Manning BS, Miroux B, *et al.* (2000): Disruption

of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species

production. *Nat Genet* 26: 435–439.

119. Jo WK, Zhang Y, Emrich HM, Dietrich DE (2015): Glia in the cytokine-mediated onset of depression:

fine tuning the immune response. *Front Cell Neurosci* 9: 268.

120. Sato K (2015): Effects of Microglia on Neurogenesis. *Glia* 63: 1394–1405.

121. Mastellos DC (2014): Complement emerges as a masterful regulator of CNS homeostasis, neural

synaptic plasticity and cognitive function. *Exp Neurol* 261: 469–474.

122. Maras PM, Daugherty J, Hebda-Bauer E, Watson SJ, Akil H (2018): Minocycline treatment reduces

anxiety- and depressive-like behaviors in a genetic model of internalizing mood disorders.

Neuroscience Meeting Planner Program No. 234.03.

123. Hilde KL, Hagenauer MH, Stefanov AV, Birt I, Hebda-Bauer E, Watson SJ, Akil H (2018): Mechanisms

of hippocampal development: Molecular and cellular phenotypes in the selectively bred high-

responder/low-responder model of affective disorders. *Neuroscience Meeting Planner* Program

No. 233.04.

124. O'Connor AM, Pardo T, Birt I, Hagenauer MH, Maras PM, Prater KE, *et al.* (2018): The hippocampal

glycome contributes to behavioural phenotype in a novel rodent model of mood disorders.

Neuroscience Meeting Planner Program No. 233.01.

125. Clinton SM, Stead JDH, Miller S, Watson SJ, Akil H (2011): Developmental underpinnings of

differences in rodent novelty-seeking and emotional reactivity. *Eur J Neurosci* 34: 994–1005.

Tables and Table Legends

	MBNI_AffymetrixRae230_F4	MBNI_AffymetrixRgU34A_F6	MBNI_AffymetrixRae230_F15	MBNI_IlluminaRatRef12v1_F15	MBNI_RNASeq_F29	Alabama_Nimblegen_F34	MBNI_RNASeq_F37	MBNI_RNASeq_F43
Generation	4	6	15	15	29	34	37	43
Laboratory	MBNI	MBNI	MBNI	MBNI	MBNI	Alabama	MBNI	MBNI
Lead Scientist	Dr. John Stead	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Sarah Clinton	Dr. Peter Blandino	Dr. Cigdem Aydin
Age	Adult	P7, P14, P21	P14	P14	P14, Adult	P7, P14, P21, Adult	Adult	Adult
n per group	6	6	6	6	2	5	6	5
Platform	Affymetrix Rat Expression Set 230 A	Affymetrix Rat Genome (RG) U34A GeneChips	Affymetrix Rat Expression Set 230 A	Illumina RatRef- 12v1 Beadchip	RNA-Seq	NimbleGen Rat Gene Expression 12x135	RNA-Seq	RNA-Seq
Tissue Extraction	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Whole Hippocampus	Dorsal Hippocampus Tissue Punch	Whole Hippocampus	Whole Hippocampus
Exposure	Basal	Basal	Basal	Basal	Basal	Basal	Basal	Vehicle Injections
Behavioral Testing	Basal	Basal	Basal	Basal	Basal	Basal	Locomotor, Anxiety	Social Interaction After Mild Stress
GEO Data Release	GSE140596	GSE29552	GSE140595	GSE140594	GSE140597	GSE88874	GSE140598	GSE140287
Citation		Clinton *et al.* 2011				Cohen *et al.* 2015		

Table 1. An overview of the eight transcriptional profiling studies included in our current meta-analyses of differential gene expression in the bHR and bLR hippocampus at four developmental time points: P7, P14, P21, and adulthood. Citations: (28,125)

Figures and Figure Legends

Figure 1. An overview of the experimental and analytical workflow used to identify top candidate genes for mediating the effects of selective breeding on bHR/bLR phenotype.

Left: Many generations of selective breeding based on exploratory locomotor behavior drove segregation of genetic variants that contribute to internalizing and externalizing behavior within our bHR and bLR rats. The effect of these variants on behavior is mediated by alterations in gene expression and cellular function, which produce local changes in cell type balance and structure within brain regions responsible for affective behavior, such as the hippocampus.

Right: Our concurrent genetic study used exome sequencing to identify genetic variants that segregated bHR/bLR rats in our colony, and then used a sampling of those variants to locate regions of the genome (quantitative trait loci – QTLs) that might contribute to exploratory locomotor activity (25). Our current study used meta-analyses of hippocampal transcriptional profiling studies to identify bHR/bLR differentially-expressed (DE) genes, pathways, cell types, and networks in development and adulthood. In our results, we emphasize DE genes that were *1)* consistently DE across multiple developmental stages, *2)* central to DE pathways, cell types, and networks, *3)* located near genetic variants that segregated bHR/bLR rats in our colony and/or within QTLs for exploratory locomotion. These genes are the top candidates for mediating the effects of selective breeding on bHR/bLR phenotype.

Figure 2. Selectively-bred high responder (bHR) and low responder (bLR) rats model an extreme propensity for internalizing vs. externalizing behavior. A) Over the course of 56 generations of selective breeding (F1-F56), the bHR rats (green) have developed increasingly elevated exploratory activity in a novel field (*y-axis:* average total locomotor score), whereas the bLR rats (red) have become less exploratory. These trends plateaued after F42, when our breeding strategy changed to deaccelerate divergence. Arrows indicate the generations during which hippocampal transcriptomic profiling datasets were collected, along with a name indicating the respective laboratory, platform, and generation for each dataset. **B)** bLR rats have

been highly anxious since the initiation of our breeding colony. The example above is from the

behavioral data accompanying the MBNI_RNASeq_F37 transcriptomic dataset showing bLRs

spending a smaller percentage of time in the anxiogenic open arms of the elevated plus maze

than bHR rats (effect of phenotype: $F(2,15)=6.72$, $p=8.25E-03$, boxes=first quartile, median, and

third quartile, whiskers = range). **C)** bLR rats are more reactive to stressors. This example is

from the behavioral data accompanying the MBNI_RNASeq_F43 transcriptomic dataset

showing bLR rats spending a smaller percentage of time interacting socially following exposure

to a single mild stressor ($F(1,8)=5.86$, $p=0.0418$).

Adult Meta-Analysis: Top Results (FDR<0.05)										
Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets	Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets
1	Tmem144	3.04E-08	4.95E-04	-3.57	3	38	Trappc6a	6.16E-05	2.61E-02	-2.06	3
2	Asb15	4.05E-07	2.32E-03	2.82	3	39	Cav1	6.47E-05	2.61E-02	1.37	5
3	Kif15	4.27E-07	2.32E-03	-2.21	4	40	Robo3	6.54E-05	2.61E-02	-2.00	3
4	Pkhd1l1	6.13E-07	2.49E-03	-3.17	3	41	C1qa	6.57E-05	2.61E-02	-1.40	5
5	Car9	7.87E-07	2.56E-03	2.78	3	42	Tubg1	6.88E-05	2.65E-02	-1.40	5
6	Klhl34	1.04E-06	2.81E-03	-2.39	4	43	Zfp952	7.17E-05	2.65E-02	2.12	2
7	Mvb12b	2.03E-06	4.32E-03	-2.95	2	44	Slc9a3r1	7.31E-05	2.65E-02	1.32	5
8	Amer2	2.12E-06	4.32E-03	2.10	3	45	Egfem1	7.45E-05	2.65E-02	-2.11	2
9	Rpl17	2.86E-06	5.17E-03	-2.16	4	46	Epb41l4a	7.49E-05	2.65E-02	-2.10	2
10	Nudt4	4.27E-06	6.51E-03	-1.58	5	47	Trmt10a	8.18E-05	2.83E-02	-1.75	3
11	Pkib	4.40E-06	6.51E-03	1.70	5	48	Oard1	8.81E-05	2.98E-02	1.65	3
12	Fxyd7	5.22E-06	6.62E-03	-1.74	4	49	Selenop	1.00E-04	3.33E-02	2.21	2
13	Scube2	5.29E-06	6.62E-03	-2.68	2	50	Ezr	1.17E-04	3.72E-02	1.27	5
14	Ddx20	5.77E-06	6.62E-03	-1.93	4	51	Acox3	1.17E-04	3.72E-02	1.26	5
15	R3hdm4	6.10E-06	6.62E-03	-2.03	3	52	Ints8	1.21E-04	3.74E-02	2.11	2
16	Hmgn5b	7.33E-06	6.94E-03	2.60	2	53	Pld4	1.22E-04	3.74E-02	-1.45	4
17	Slc39a12	7.35E-06	6.94E-03	2.62	3	54	Ucp2	1.27E-04	3.83E-02	1.29	5
18	Acss2	7.68E-06	6.94E-03	-1.87	4	55	Rbm3	1.29E-04	3.83E-02	-1.52	4
19	Rhpn2	8.16E-06	6.98E-03	2.33	3	56	Ankdd1b	1.37E-04	3.93E-02	-2.00	2
20	Exosc7	9.65E-06	7.85E-03	1.80	4	57	Aar2	1.38E-04	3.93E-02	2.02	2
21	Rnls	1.04E-05	8.06E-03	1.92	3	58	Fn3k	1.42E-04	3.97E-02	-1.47	4
22	Slc19a3	1.50E-05	1.11E-02	-2.36	3	59	Slc27a1	1.45E-04	4.01E-02	1.24	5
23	Prss55	1.91E-05	1.35E-02	-2.43	2	60	Afmid	1.52E-04	4.11E-02	1.98	2
24	Apln	2.07E-05	1.41E-02	1.48	5	61	Slc4a11	1.54E-04	4.11E-02	-1.78	3
25	Mfge8	2.70E-05	1.73E-02	1.50	5	62	Zfp110	1.64E-04	4.31E-02	1.24	5
26	C2cd3	2.84E-05	1.73E-02	2.06	3	63	Tek	1.70E-04	4.33E-02	-1.79	3
27	Dnaaf3	2.87E-05	1.73E-02	-2.51	2	64	Fmo5	1.70E-04	4.33E-02	1.42	4
28	Etv4	3.30E-05	1.92E-02	2.02	3	65	Blvra	1.74E-04	4.36E-02	1.52	4
29	Tmem2	3.72E-05	1.99E-02	-2.29	2	66	Prdm5	2.03E-04	4.94E-02	-1.76	3
30	Tdg	3.84E-05	1.99E-02	-1.38	5	67	Samd5	2.09E-04	4.94E-02	1.95	3
31	Mki67	4.03E-05	1.99E-02	-2.01	3	68	Ntn4	2.10E-04	4.94E-02	1.94	2
32	Tnnt1	4.06E-05	1.99E-02	-1.51	5	69	Chd1l	2.11E-04	4.94E-02	1.45	4
33	Zfp90	4.13E-05	1.99E-02	-1.59	4	70	Ccdc137	2.13E-04	4.94E-02	1.97	2
34	LOC363337	4.16E-05	1.99E-02	-2.82	2	71	Trhr	2.16E-04	4.94E-02	1.19	5
35	Uhrf1	4.95E-05	2.29E-02	-1.67	3	72	Zfp821	2.22E-04	4.98E-02	-1.48	3
36	Tuba8	5.06E-05	2.29E-02	-1.44	4	73	Myh6	2.23E-04	4.98E-02	-1.25	5
37	Rltpr	5.58E-05	2.45E-02	-2.28	2	74	Sp3	2.26E-04	4.98E-02	1.93	3

P7: Top Result (FDR<0.05)						P14: Top Result (FDR<0.15)				
Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets	Rank	Gene Symbol	P-value	FDR	Estimate	# of Datasets
1	Ncan	1.38E-07	4.86E-04	-4.16	2	1	Bmp4	9.40E-06	1.47E-01	-1.49	5

Figure 3. The top DE genes within the bHR/bLR hippocampal gene expression meta-analyses. P-value=nominal p-value, FDR=false detection rate, Estimate=estimated effect size (i.e., the difference in expression between bHR and bLR rats in units of standard deviation,

green/positive=higher expression in bHRs, red/negative= higher expression in bLRs), # of

datasets=number of datasets included in the meta-analysis for that gene.

Figure 4. The extracellular matrix constituent Neurocan (Ncan) has elevated expression in bLR rats at age P7. A) A genetic variant on Chr 16 nearby Ncan (<100 kb) segregated bHR and bLR rats (Fisher's exact test: *p*=1.63E-07). **B)** Boxplots illustrating the effect of age and bHR/bLR phenotype on Ncan expression (log(2) signal) in two microarray studies (boxes=median and interquartile range, whiskers=range, red=bLR, green=bHR). The effect of phenotype is most obvious at an age when Ncan is elevated in development (P7). **C-D)** Forest

plots showing that Ncan was more expressed in bLRs than bHRs (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis) **C)** in the P7 meta-analysis (β=-4.16, p=1.38E-07, *FDR*=4.86E-04), **D)** and nominally in the P14 meta-analysis (β=-0.90, p=1.01E-02, *FDR*=7.24E-01).

Figure 5. A regulator of proliferation and differentiation, Bone morphogenetic protein 4 (Bmp4), is more highly expressed in bLR rats than bHR rats at P14 and adulthood. A-B) Two forest plots showing that Bmp4 was consistently elevated in bLR rats (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis) at **A)** P14 (β=-1.49, p=9.40E-06, *FDR*=1.47E-01) and **B)** adulthood (adult: β=-1.04, p=1.01E-03, *FDR*=9.38E-02). This direction of effect mirrors findings in the literature that show that blocking the expression of Bmp4 in mice reduces anxiety

and depressive-like behavior (98). **C)** Using qPCR, we confirmed that bLRs showed greater

Bmp4 expression than bHRs at P14 and adulthood (P90) using hippocampal tissue from later

generations (F51, F55). Log(2) fold change in *Bmp4* expression was calculated using the Livak

method ($-\Delta\Delta$Cq, (43)), using *Gapdh* as the reference housekeeping gene and bLRs as the

reference group (therefore the bLR mean is set to 0 in all panels). **$p<0.005$ (Data release: doi:

10.6084/m9.figshare.10321658; *P14*: *Log(2)FC*=-3.74, $T(5.60)$=-6.10, *p*=0.00115; *P90*:

$T(8.74)$=-6.87, *p*=8.44E-05). **D)** Within the behavioral data accompanying the

MBNI_RNASeq_F37 dataset, we found that Bmp4 showed a negative relationship with percent

time in the open arms (β=-0.034, R^2= 0.28, *p*=0.024) and a positive relationship with the number

of fecal boli produced on the EPM (β=0.32, R^2= 0.29, *p*=0.020, *data not shown*).

A. bHR/bLR Differentially-Expressed Genes Are Often Located Close to Genetic Variants (SNPs) that Segregate bHR and bLR Rats

B. bHR/bLR Differentially-Expressed Genes Often Overlap QTLs for Exploratory Locomotion

p=0.0035

Adult Meta-Analysis

C. Differentially-Expressed Genes within QTLs for Exploratory Locomotion:

	QTLs from bHR/bLR F2 Intercross: Exploratory Locomotion			Overlapping QTLs from Rat Genome Database		
Chr	QTL range in MB* (LOD>3, +/-1MB)	LOD	Genes w/ FDR<0.1 in Meta-Analysis	Anxiety	Stress Response	Behavioral Despair
1	28-31	3.13			[53]	[56]
1	39-65	4.75	*Ezr*	[47]	[53]	
1	98-104	3.42	Dbp	[47],[48]	[53]	
1	115-132	7.76	*Hmgn5b*	[47],[48]	[53],[54]	
1	139-142	3.14	*Mfge8*	[47],[48]	[54]	
1	156-171	4.32	Olr35, Spcs2, *C2cd3*, *Ucp2*	[47],[48]	[54]	
2	43-70	4.26	Pdzd2, *Selenop*	[49]	[55]	
2	92-95	3.02			[55], [55]	
3	43-65	4.71	Ttc30a1	[49]	[53]	[56]
5	98-102	3.34		[51],[52]		
7	67-85	5.14	*Trhr*, *Pkhd1l1*	[49]	[53]	
17	157-177	3.82	Chrm3, Akr1cl			
18	71-81	4.99		[50]		

D. Top Genomic Regions Enriched with Differentially Expressed Genes:

	Genetic Loci Enriched with bHR/bLR Differentially Expressed Genes (PGE)					Overlapping QTLs:			
Chr	Location (Rnor_6.0)	P-value	FDR	Size of Region (kb)	Enrichment Score	Genes w/ FDR<0.1 in Meta-Analysis	Exploratory Locomotion	Anxiety	Stress Response	Behavioral Despair
1	2782900-4653300	7.76E-04	1.02E-02	1,870	0.6	*Samd5*				[56]
1	164225800-165684000	4.97E-05	4.68E-03	1,458	0.2	Olr35, Spcs2, *C2cd3*, *Ucp2*	[25]	[47], [48]	[54]	
1	174385400-174620200	1.77E-04	5.69E-03	235	1	*Scube2*		[48]	[63]	
3	62800900-63455200	7.76E-04	1.02E-02	654	0.6	Ttc30a1	[25]	[49]	[53]	[56]
3	119361600-119677400	1.22E-04	5.69E-03	316	0.6	*Blvra*			[53],[55]	[56]
3	151032900-151150600	1.77E-04	5.69E-03	118	1	*Acss2,* Myh7b			[55]	
3	176280000-176526000	4.08E-04	7.97E-03	246	0.8					
5	155246400-155264100	1.77E-04	5.69E-03	18	1	*C1qa*, C1qc		[49]		
7	81919900-83358500	2.02E-04	6.05E-03	1,439	0.5	*Trhr*	[25]	[49]	[53]	
10	88392200-103730100	6.80E-05	4.68E-03	15,338	0.1	*Slc9a3r1*, Sox9, *Tubg1*,Ghdc, Tmem101, Acbd4, Ddx42, *Etv4*		[50]	[64], [65]	
12	37444100-39302600	4.30E-06	2.96E-03	1,859	0.3	Eif2b1				
16	81035000-81209500	3.35E-05	4.68E-03	175	0.8				[55]	
19	32182900-42110300	2.85E-05	4.68E-03	9,927	0.1	*Zfp90*				

Figure 6. Many of the top differentially expressed genes are located near genetic variants that segregate bHR/bLR rats within quantitative trait loci (QTLs) for exploratory locomotor activity. A. Our concurrent genetic study used exome sequencing to identify variants (SNPs) that segregated bHR/bLR rats in our colony (25). A high percentage of bHR/bLR differentially-expressed genes (adult meta-analysis: *FDR*<0.05) were found within +/- 100 kb, 250 kb, 500 kb, or 1 MB of these segregating variants, using either traditional (Bonferonni-corrected *p*<0.05) or more stringent (Bonferonni-corrected *p*<5.00E-5) criteria to define segregation. Asterisks (*) designate enrichment (Fisher's exact test *p*<0.0001) in comparison to the non-differentially expressed genes in our meta-analysis (*FDR*>0.05). **B.** Our concurrent genetic study used a sampling of bHR/bLR segregating variants to identify QTLs for exploratory locomotor activity in a novel field using an bHR/bLR F2 intercross (25). bHR/bLR differentially-expressed genes (adult meta-analysis: *FDR*<0.05) were 4.5x more likely to overlap (+/- 1MB) QTLs for exploratory locomotion than other genes included in our meta-analysis (Fisher's exact test: *p*=0.0035). **C.** A table illustrating the top genes from our meta-analyses (*FDR*<0.1, bold+italic=*FDR*<0.05) that overlap (+/-1 MB) significant (LOD>4) and putative (LOD>3) QTLs for exploratory locomotion identified by our concurrent genetic study (25). Also depicted is overlap with QTLs identified in the Rat Genome Database (32) for the following behaviors relevant to the bHR/bLR phenotype: anxiety (47–52), stress-related responses (53–55), and behavioral despair (56). **D.** The top chromosomal loci enriched for bHR/bLR DE genes overlap previously-identified QTLs relevant to externalizing and internalizing behaviors. The top chromosomal loci enriched for bHR/bLR DE genes were identified using Positional Gene Enrichment analysis (PGE): location (full coordinates), p-value=nominal p-value, *FDR*=false detection rate, enrichment score= ratio of genes with *p*<0.01 out of all genes in the region. Also depicted is the overlap (+/-1 MB) of these enriched chromosomal loci with QTLs for exploratory locomotor activity (25), as well as with QTLs identified in the Rat Genome Database (32) for anxiety (47–50), stress-related responses (53–55,63–65), and behavioral despair (56).

Figure 7. A region on chromosome 1 implicated in bHR/bLR phenotype contains two genes important for brain function and development, Uncoupling Protein 2 (Ucp2) and C2 Calcium Dependent Domain Containing 3 (C2cd3). A) Genetic variants on Chr 1 within Ucp2 and C2cd3 segregate bHR and bLR rats in our colony (Rnor5 coordinates, Fisher's exact test: SNP 1_171712284: *p*=1.66E-09; SNP 1_171672484: *p*=1.27E-13). **B)** Ucp2 and C2cd3 are located on Chr 1 within a QTL for exploratory locomotor activity. An example of the correlation between genetic variation in this region and behavior is illustrated using the sequencing results

from a nearby SNP and exploratory locomotor activity measured in a bHRxbLR F2 intercross (*n*=310; *adj.R^2*=0.049, *p*=2.00E-04, *FDR*=0.0048). **C)** C2cd3 and Ucp2 were the top DE genes (*FDR*<0.05) within a segment of Chr 1 enriched for DE genes and containing a QTL for exploratory locomotor activity (25). The table illustrates the DE genes within this region: estimate=estimated effect size (green/positive=higher expression in bHRs), bold=*p*<0.05, bold+italic=*FDR*<0.05. **D)** A forest plot showing that Ucp2 had higher expression in bHRs in four out of the five adult datasets included in the adult meta-analysis (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis, effect of phenotype: *β*=1.29, *p*=1.27E-04, *FDR*=3.83E-02). This direction of effect mirrors findings in the literature showing that Ucp2 knockout mice have higher anxiety-like behavior and lower locomotor activity, as well as greater sensitivity to stress (57,58,60,61), much like our bLR rats. **E)** A forest plot showing that C2cd3 had higher expression in bHRs in three adult datasets included in the adult meta-analysis (effect of phenotype: *β*=2.06, *p*=2.84E-05, *FDR*=1.73E-02). **F)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, C2cd3 (units: log(2) fragments per million (FPM)) showed a positive relationship with exploratory locomotor activity (*β*= 0.000109, *R^2*=0.26, *p*=3.20E-02). **G)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Ucp2 showed a trend towards a positive relationship with percent of time spent in the anxiogenic open arms of the EPM (*β*=0.03, *R^2*=0.22, *p*=5.16E-02).

A. Genetic Variant Near Trhr (<200 kb distant) Segregates bHR and bLR Rats

Nearby SNP (in Nudcd1): 7_83331058 (Rnor5)

	CC	CT	TT
bHR	0	0	12
bLR	12	0	0

B. Trhr is Located within a QTL on Chr 7 for Exploratory Locomotor Activity

Nearby SNP in QTL: 7_83331058 (Rnor5)

C. Trhr is More Highly Expressed in bHR Rats in Adulthood

Investigator & Study	Cohen's D [95% CI]
MBNI_AffymetrixRae230_F4	1.15 [-0.19, 2.49]
MBNI_RNASeq_F29	1.03 [-1.05, 3.11]
Alabama_NimbleGen_F34	0.96 [-0.35, 2.27]
MBNI_RNASeq_F37	1.08 [-0.13, 2.29]
MBNI_RNASeq_F43	1.77 [0.30, 3.24]
Model	1.19 [0.56, 1.82]

D. Trhr is the Top Result in a Section of Chr 7 Enriched for Differentially-Expressed Genes

Chr7: 81,919,000-83,359,000

Gene Symbol	Adult_estimate	P14_estimate
Rspo2	-1.35	NA
Emc2	1.04	NA
Trhr	1.19	0.35
Eny2	-1.01	0.14

F. Trhr Correlates with Exploratory Activity

R-squared= 0.22 , P-value= 0.051

G. Trhr is the Top Result Within a Hippocampal Co-expression Network Enriched for Differentially-Expressed Genes

E. Trhr is a Top Result in Two Hippocampal-Specific Gene Sets Enriched with bHR/bLR Differential Expression

Top Co-expression Networks in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
From mouse: "paleturquoise"	1.94	-1.08	**ADAMTS2**,FOXC1,FBLN1,MRC1,OMD,ANGPT2,EMP3,BGN,CD248,HSPB1
From human: "M11"	-1.72	-1.84	**FBXO31**,NPHP4,PPME1,ARHGAP39,FAM131B,PLXNA1,NCOR2,PPFIA4,DMTN,STRN4
From mouse: "darkred"	-1.68	0.98	MTF2,DUSP11,SNHG11,DDX26B,LRRC29,RYR3,NFKBIZ,EML5,TCF25,NRXN3
From human: "M24"	1.54	-0.97	**EZR**,ITSN1,IDNK,PPAP2B,FAM167A,GFPT2,**ITGB4**,PSD2,PITPNC1,PHGDH
From mouse: "sienna3"	1.91	-0.64	OTOF,**TRHR**,SMOC1,ZBTB7C,GLRA2,GRB10,MEIS1,GPR83,ARHGAP6
From mouse: "lightcyan"	-1.27	-1.47	**RLTPR**,ATMIN,KAT2A,JPH3,ACAD8,REV1,USE1,B9D2,RAB3D,PWWP2B

Top Co-expression Networks at P14 (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
From mouse: "lightyellow"	-1.13	-1.56	ZFP277,ACTA2,DMWD,SEMA7A,UQCRC1,CNNM3,ATM,NPAS2,NCAN,BRD4
From human: "M11"	-1.72	-1.84	DTX3,FBXO31,GFOD1,HYOU1,ADCY9,LRFN4,DYRK1B,PTMS,UBA1,WBSCR17
From mouse: "lightcyan"	-1.27	-1.47	H1FX,ELFN2,ZCRB1,APBA2,SLC2A13,POLK,SNUPN,ATG5,HYOU1,SNAPIN
From human: "M1"	NA	-1.36	EID2,CHST15,RIMBP2,EMX1,NT5C3B,ELFN2,FAM65A,SRSF7,ZDHHC22,APBA2
From mouse: "skyblue"	0.86	-1.84	TSC22D4,MOBP,ADSSL1,SLC35B2,OPALIN,MAG,EFHD1,GJC2,ITGB4
From human: "M7"	-1.45	-1.70	ARHGEF4,ST6GALNAC6,DHRS11,AGAP3,B3GAT1,ARHGEF25,LRP3,PLSCR3,LMO4,NCS1

Top Regional Markers in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
"dg_v__VS__dg_d"	2.05	0.88	**TRHR**,TOX3,PDZRN3,FBLN1,TSHZ1,STK32B,ZFP423,NCAPG,LCORL

Top Regional Markers at P14 (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
"dg_d-dg_v__VS__ca4-ca3_d-ca3_v-ca2-ca1_d-ca1_v"	-0.88	-1.76	SCN3A,ARHGAP20,MARCKS,SV2C,PDE7B,TRPC6,SLC4A4,BTG2,IGFBP5,RFX3

Figure 8. Thyrotropin releasing hormone receptor (Trhr) was the top gene within two hippocampal specific gene sets and within a region of Chromosome 7 implicated in bHR/bLR phenotype. A) A genetic variant on Chr 7 near Trhr (<200 kb distant) segregates bHR and bLR rats in our colony (Fisher's exact test: p=6.20E-14). **B)** Trhr is located on Chr 7 within a QTL for exploratory locomotor activity. An example of the correlation between genetic variation in this region and behavior is illustrated using the sequencing results from a SNP nearby Trhr (discussed above) and exploratory locomotor activity measured in a bHRxbLR F2 intercross (n=315, $adj.R^2$=0.061, p=5.79E-05, FDR=0.00178). **C)** A forest plot showing that Trhr expression was consistently elevated in bHR rats since generation F4 (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis, β=1.19, p=2.16E-04, FDR=4.94E-02). This direction of effect mirrors findings in the literature that show Trhr-KO mice exhibit greater anxiety and depressive-like behavior (62). **D)** Trhr was the strongest result within a segment of Chromosome 7 enriched for DE genes and overlapping a QTL for exploratory locomotor activity. The table illustrates the DE genes within this region: estimate=estimated effect size (green/positive=greater expression in bHRs), bold=p<0.05, bold+italic=FDR<0.05. **E)** Trhr was a leading gene in two of the top hippocampal-specific gene sets identified as enriched for bHR/bLR DE genes by GSEA (FDR: False Detection Rate; NES: Normalized Enrichment Score, with positive scores (green) indicating greater expression in bHRs and negative scores (red) indicating greater expression in bLRs, bold: p<0.05 in GSEA results, bold+italics: FDR<0.05 in GSEA results). The top 10 "Leading Edge" genes for each gene set are shown (bold+italics: FDR<0.05 in meta-analysis). These genes have large estimated effect sizes within the meta-analysis and help drive the enrichment of effects within these gene sets. Regional marker gene sets use the following abbreviations: "dg"=dentate gyrus, "v"=ventral, "d"=dorsal, "ca"=Cornu Ammonis subregion. **F)** In the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Trhr (units: log(2) fragments per million (FPM)) showed a trend towards a positive relationship

with exploratory locomotor activity (β=4.48E-04, R^2= 0.22, p=5.08E-02). **G)** Trhr and its ligand,

Thyrotropin releasing hormone (Trh), are hub genes within a hippocampal specific co-

expression network that is enriched for bHR-upregulated genes. Genes within this network with

known protein-protein interactions are illustrated above (STRINGdb: confidence setting=0.15

due to hippocampal co-expression already suggesting potential interaction). Many of these

genes have documented associations with reward behavior.

A. Top Results Are Enriched with Genes Related to Cell Proliferation, Differentiation, and Development

Top Pathways in Adults (FDR<0.10):	Adult NES	P14 NES	Leading Edge Genes in Adults (Bold = FDR<0.10 in meta-analysis):
protein_folding(6)	2.32	1.14	**FKBP4**,HSPA8,**CCT4**,CHORDC1,PDIA6,HSP90AA1,CLU,DNAJC10,TCP1
positive_regulation_of_cell_proliferation(4)	1.59	-1.30	**SOX9,MFGE8,APLN**,GPR183,**CAV1,CD24,HES5,HTRA1,GLUL**,LEP
nephron_development(4)	2.08	-0.94	**SOX9,CD24**,FOXD1,PTPRO,ANGPT2,SERPINB7,WNK4,SOX8,LHX1,TACSTD2
chaperone_mediated_protein_folding(7)	2.17	1.32	**FKBP4**,HSPA8,**CCT4**,CHORDC1,CLU,PDIA4,CANX,PPID,ST13,HSPD1

Top Pathways at P14 (FDR<0.05):	Adult NES	P14 NES	Leading Edge Genes at P14 (Bold = FDR<0.10 in meta-analysis):
generation_of_neurons(7)	0.94	-1.56	FARP1,BMP4,BEX1,SOX9,ANKRD27,SHC1,SNX3,TNC,FBXO31,TCF12
negative_regulation_of_multicellular_organismal_process(3)	1.11	-1.62	BMP4,BEX1,SOX9,LRP1,XYLT1,SRGAP2,SERPINF1,NPFF,GIMAP5,HDAC7
neurogenesis(6)	0.98	-1.51	FARP1,BMP4,BEX1,SOX9,ANKRD27,SHC1,SNX3,TNC,FBXO31,TCF12
regulation_of_neurogenesis(6)	0.78	-1.60	BMP4,BEX1,SOX9,SNX3,FBXO31,TCF12,LRP1,XYLT1,SRGAP2,SERPINF1
neuron_differentiation(6)	0.86	-1.56	FARP1,BMP4,BEX1,SOX9,ANKRD27,SHC1,SNX3,TNC,FBXO31,TCF12
negative_regulation_of_cellular_macromolecule_biosynthetic_process(6)	0.92	-1.58	BMP4,SOX9,UXT,CHD8,YBX1,GFPT1,RARA,ZEB1,HDAC7,KHDRBS1
regulation_of_neuron_differentiation(7)	-0.70	-1.59	BMP4,BEX1,SOX9,SNX3,FBXO31,TCF12,LRP1,XYLT1,SERPINF1,RARA
regulation_of_nervous_system_development(5)	0.83	-1.57	BMP4,BEX1,SOX9,SNX3,FBXO31,TCF12,LRP1,XYLT1,SRGAP2,SERPINF1

B. An Unbiased PPI-Network Constructed from the Top Results Highlights Genes Involved in Cell Proliferation and Differentiation

Up in bHR (green arrow)   Up in bLR (red arrow)

Network nodes: Cd24, Hes5, Mki67, Itgb4, Cav1, Sp3, Sox9, Tek, Bmp4, Map3k7, Sox2, Hipk4

C. Genetic Variants in Marker of Proliferation Ki-67 Segregate bHRs & bLRs

SNP within Mki67:
1_214939984 (Rnor5)

	AA	AG	GG
bHR	12	0	0
bLR	0	2	10

SNP within Mki67:
1_214940284 (Rnor5)

	CC	CT	TT
bHR	12	0	0
bLR	0	2	10

D. Mki-67 is More Highly Expressed in bLRs in Adulthood

bLR   bHR

Investigator & Study		Cohen's D [95% CI]
MBNI_RNASeq_F29		-0.81 [-2.85, 1.23]
MBNI_RNASeq_F37		-2.14 [-3.55, -0.73]
MBNI_RNASeq_F43		-2.67 [-4.38, -0.96]
Model		-2.01 [-2.97, -1.05]

Observed Outcome: -6 -4 -2 0 2

E. Mki-67 Correlates with Exploratory Activity

R-squared= 0.41 , P-value= 0.0042

Mki67: Log(2) FPM vs Locomotor Score (0, 500, 1000, 1500, 2000, 2500)

Legend: bHR, bLR, bLR

Figure 9. The top bHR vs. bLR DE results are enriched with genes related to cell proliferation, differentiation, and development, including the canonical Marker of Proliferation (Mki67). A) A table of the top functional ontology gene sets identified as enriched for bHR/bLR DE genes by GSEA (FDR: False Detection Rate; NES: Normalized Enrichment Score, with positive scores (green) indicating higher expression in bHRs and negative scores (red) indicating higher expression in bLRs, bold: *p*<0.05 in GSEA results, bold+italics: *FDR*<0.05 in GSEA results). The top 10 "Leading Edge" genes for each gene set are shown (bold+italics: *FDR*<0.05 in meta-analysis). These genes have large estimated effect sizes within the meta-analysis and help drive the enrichment of effects within these gene sets. **B)** A PPI network constructed using the top genes from the adult meta-analysis (192 genes with *FDR*<0.10, STRINGdb: confidence setting=0.40) had a dominant subnetwork that included Bmp4 and Mki67 as hub genes. Many of these genes are related to cell proliferation and differentiation within the brain. **C)** Two genetic variants on Chr 1 within Mki-67 fully segregated bHR and bLR rats in our colony (Rnor5 coordinates, Fisher's exact test: SNP 1_214939984: *p*=2.02E-11, SNP 1_214940284*: p*=2.02E-11). **D)** A forest plot showing that Mki67 was consistently elevated in bLR rats in adulthood (boxes=Cohen's D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95% confidence intervals provided by the meta-analysis, adult: *β*=-2.01, *p*=4.03E-05, *FDR*=1.99E-02). **E)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, Mki67 (units=log(2) fragments per million (FPM)) showed a negative relationship with locomotor score (*β*=-0.000249, R^2=0.41, *p*=0.0042).

Figure 10. Microglial-related gene expression differentiates bHR and bLR rats. A) A

heatmap illustrating the effect of bHR/bLR phenotype on cell type specific gene expression,

which can reflect overall cell type balance (42) or activation: green=upregulated in bHRs,

red=upregulated in bLRs, asterisks(*) indicate *p*<0.05. Note that only well-characterized non-

neuronal cell type categories were included in this analysis. **B-D)** Forest plots (boxes=Cohen's

D from each study +/-95% confidence intervals, "Model"=estimated effect size +/-95%

confidence intervals provided by the meta-analysis) showing an upregulation in bLRs of (**B**)

endothelial-specific gene expression at P14 (*β*=-0.89, *p*=5.75E-03) and (**C-D**) microglial-specific

gene expression **C)** at P14 (*β*=-0.69, *p*=2.90E-02) and **D)** adulthood (*β*=-0.65, *p*=4.47E-02). **E)**

A genetic variant on Chr 1 in Milk fat globule-EGF factor 8 (Mfge8) segregates bHR and bLR

rats in our colony (Fisher's exact test: *p*=7.53E-08). Mfge8 promotes alternative (M2) activation

of microglia. **F)** A forest plot illustrating that Mfge8 is more highly expressed in bHRs than bLRs

in the adult meta-analysis in all five adult datasets (*β*=1.50, *p*=2.70E-05, *FDR*=1.73E-02). **G)**

Mfge8 is located on Chr 1 within a QTL for exploratory locomotor activity. An example of the

correlation between genetic variation in this region and behavior is illustrated using the

sequencing results from a nearby SNP (Rnor5 coordinates 1_ 141117448) and exploratory

locomotor activity measured in a bHRxbLR F2 intercross (*n*=317, *adj.R²*=0.061, *p*=1.81E-05,

FDR=0.002). **H)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset,

Mfge8 (units=log(2) fragments per million (FPM)) showed a positive relationship with total

locomotor score (*β*=0.000146, *R²*=0.62, *p*=1.10E-04) as well as the percent time in the open

arms of the EPM (*β*=0.00603, *R²*=0.28, *p*=2.30E-02). **I)** A genetic variant on Chr 5 near

Complement component C1q A Chain (500 kb distant) segregates bHR and bLR rats in our

colony (Rnor5 coordinates 5_159525078, Fisher's exact test: *p*=1.09E-07). C1qa promotes

classical activation of microglia via the complement cascade. **J)** A forest plot illustrating that

C1qa is more highly expressed in bLRs than bHRs in all five datasets in the adult meta-analysis

(*β*=-1.40, *p*=6.57E-05, *FDR*=2.61E-02). **K)** C1qa is the top DE gene (FDR<0.05) within a

segment of Chr 5 enriched for DE genes. The table illustrates the DE genes within this region: estimate=estimated effect size (red/negative=higher expression in bLRs), bold=p<0.05, bold+italic=FDR<0.05. **L)** Within the behavioral data accompanying the MBNI_RNASeq_F37 dataset, C1qa showed a negative relationship with total locomotor score (C1qa: β=-5.17E-04, R^2=0.29, p=2.09E-02).