Astrovirus in Reunion Free-tailed Bat (*Mormopterus francoismoutoui*)

Léa Joffrin, Axel O. G. Hoarau, Erwan Lagadec, Patrick Mavingui, Camille Lebarbenchon*

Université de La Réunion, UMR Processus Infectieux en Milieu Insulaire Tropical (PIMIT),
INSERM 1187, CNRS 9192, IRD 249, GIP CYROI, 2 rue Maxime Rivière, Saint Denis, La
Réunion, France.

* camille.lebarbenchon@univ-reunion.fr

Abstract

Astroviruses (AstVs) are RNA viruses responsible for infection of a large diversity of avian and

mammalian species, including bats, livestock, and humans. We investigated AstV infection in a

free-tailed bat species, *Mormopterus francoismoutoui*, endemic to Reunion Island. A total of 190

guano samples were collected in a maternity colony during 19 different sampling sessions, between

June 2016 and June 2017. Biological material was tested for the presence of the AstV RNA-

dependent RNA-polymerase (RdRp) gene with a pan-AstV semi-nested polymerase chain reaction

assay. In total, 15 guano samples (7.9%) tested positive, with high genetic diversity of the partial

RdRp gene sequences among positive samples. A phylogenetic analysis further revealed that the

detected viruses were genetically related to AstVs reported in reptiles, dogs, and pigs, but did not

cluster with AstVs commonly found in bats. Although more investigation need to be conducted to

assess the level of infected bats in the studied population, our findings suggest that Reunion free-

tailed bats are exposed to AstV, and that cross-species transmission may occur with other hosts

sharing the same habitat.

Introduction

Astroviruses (AstVs) are positive-sense single-stranded RNA viruses and have been detected in a large diversity of mammalian and avian species (De Benedictis, Schultz-Cherry, Burnham, & Cattoli, 2011). In humans, eight serotypes have been described worldwide, accounting for 2%–9% of all acute non-bacterial gastro-enteritis cases in children (Bosch, Pinto, & Guix, 2014). AstVs are also responsible for diseases in livestock, poultry and domestic pets (De Benedictis et al., 2011). In wild animals, they have been mostly described in bats (Fischer, dos Reis, & Balkema-Buschmann, 2017) and in aquatic birds (Chu et al., 2012), although detection in other hosts has been reported, such as in marine mammals (Rivera, Nollens, Venn-Watson, Gulland, & Wellehan, 2010) and non-human primates (Karlsson et al., 2015). In addition to this large host diversity, the evolutionary history of AstVs is characterized by frequent cross-species transmission events (Mendenhall, Smith, & Dhanasekaran, 2015), supporting limited host specificity and a risk of spillover and emergence in humans (Bosch et al., 2014).

AstVs have been reported in African insectivorous bats, in particular in species of the Miniopteridae and Rhinonycteridae families (Hoarau et al., 2018; Rougeron et al., 2016; Waruhiu et al., 2017; Yinda et al., 2018). Detection rates varies greatly depending on the study design and the tested species (Fischer et al., 2017), and the prevalence of individual shedding AstVs within bat populations is also highly variable. For instance, we estimated prevalence of shedding bats ranging from 20% to 81% in *Triaenops afer* in Mozambique, depending on the time of sample collection (Hoarau et al., 2018). Such a strong seasonality has been reported in both tropical and temperate regions and could be associated to bat population dynamics, in particular in maternity colonies (*e.g.* increase in the population size and density during the formation of the colony (Drexler et al., 2011). Body condition and co-infection with other viruses have also been identified as risk factors for AstV infection in bats (Mendenhall et al., 2017; Seltmann et al., 2017).

The islands of the Western Indian Ocean host eleven of the eighteen bat families described worldwide, with a highly contrasted diversity pattern between islands. Madagascar shelters over 46 bat species, of which nearly 80% are endemic. In contrast, the small oceanic islands surrounding Madagascar are inhabited by a limited number of bat species. For instance, only three species are present on Reunion Island and only one of them, the Reunion free-tailed bat (RFTB; *Mormopterus francoismoutoui*; Molossidae), is endemic to the island.

RFTB is also the most abundant bat species on Reunion Island. This small insectivorous species roost in large colonies in natural caves and synanthropic habitats such as bridges and houses (Goodman et al., 2008). Based on guano sampling in a RFTB maternity colony, we investigated the temporal variation in AstV shedding during 13 consecutive months, with a particular emphasis on the parturition period. Partial sequencing of the AstV RNA-dependent RNA-polymerase (RdRp) gene and phylogenetic analyses were performed to assess the diversity and the origin of the detected viruses.

Material and methods

The study was conducted in the main RFTB colony known to date on Reunion island, a natural cave (30 m^3) in the West coast of the island (21,111°S, 55,259°E). The colony is monospecific and mostly composed of adult females during the early stages of the breeding season (October to December), and of newborns and juveniles, after the parturition period, from mid-December to mid-January onward (Dietrich et al., 2015). Between 40 000 and 50 000 flying adult bats have been estimated before parturition; however, after the breeding season, the cave remains totally empty from bats (between May and September) (Dietrich et al., 2015).

A non-invasive sampling strategy was setup to avoid direct manipulation of bats and to limit colony disturbance. During each sampling session, we collected guano samples in the narrowest part of the cave entrance, in ten different points separated by 50 cm along a 4.5 m transect (Supplementary Figures 1 and 2). Sterile open-ended 2 mL syringes were used to collect a core sample, in order to obtain *c.* 130 mg of guano. Each guano sample was immediately mixed in 1.5 mL of Virus Transport Media (VTM; *e.g.* Lebarbenchon et al., 2017). Samples were maintained in a cooler with ice packs in the field and stored in a -80°C freezer within two hours.

In total, 190 samples were collected during 19 different sampling sessions between June 21st 2016, and June 29th 2017. During each sampling session, the colony was visually monitored in order to estimate the relative population size (*i.e.* no bat, less than half of the cave covered with bats, more than half of the cave covered), and to assess the population age structure and timing of parturition based on the presence of adults, newborns, and juveniles.

Samples were mixed and centrifuged at 1 500 g for 15 min. RNA extraction was performed on 140 µL of supernatant, with the QIAamp Viral RNA Mini Kit (QIAGEN, Valencia, CA, USA). Reverse-

transcription was done on 10 µL of RNA using the ProtoScript II Reverse Transcriptase and Random Primer 6 (New England BioLabs, Ipswich, MA, USA) following a previously published protocol (Lebarbenchon et al., 2017). cDNAs were tested for the presence of the AstV RdRp gene using a pan-AstV semi-nested polymerase chain reaction (PCR) assay (Chu, Poon, Guan, & Peiris, 2008). PCR products of the expected size were submitted for direct Sanger sequencing (Genoscreen, Lille, France). Chi-square tests were conducted to investigate the effects of the sampling date and of the sampling location in the transect, on the probability of successful detection of AstV RdRp gene. Statistical analyses were conducted with R 3.4.4 (R Core Team, 2018).

A phylogenetic analysis was performed on the RdRp gene sequences obtained in this study (GenBank accession numbers MK966904 to MK966918) and 79 reference RdRp sequences of AstVs detected in a large diversity of bat species, in particular from Africa (Madagascar, Mozambique, Gabon), as well as in other type of hosts (*e.g.* avian, human, livestock). Sequences were aligned with CLC Sequence Viewer 8.0 (CLC Bio, Aarhus, Denmark). A maximum-likelihood analysis was done with the software PhyML 3.1 (Guindon et al., 2010), with an evolutionary model selected by Model Generator 0.85 (GTR + I + Γ; I = 0.14, α = 0.83; (Keane, Creevey, Pentony, Naughton, & Mclnerney, 2006), and 1 000 bootstraps.

Results and discussion

In total, 15 guano samples (7.9 %) tested positive for the presence of AstVs, without difference between sampling location along the transect (χ^2 = 10.5; P = 0.31). Significant differences in the number of positive samples between sampling dates were detected (χ^2 = 46.7; P < 0.001), with AstVs mostly detected after the parturition period (late December), in February and March, when the cave was occupied by juveniles (Figure 1). This pattern suggests a potential effect of the population age structure (Mendenhall et al., 2017) rather than changes in size and density associated to the formation of the colony (Drexler et al., 2011). A previous investigation on paramyxovirus and *Leptospira* infection dynamics in the same bat colony revealed two peaks of infection: one during the colony formation and the other two months after the birth pulse (Dietrich et al., 2015). The second peak may coincide with the detection of most AstVs in February and March, and the first one with the three positive samples detected during our two sampling session in December. Although these results provide support for AstV RNA in the RFTB environment, our sampling design precludes further conclusion on the prevalence of shedding bats in the population. Bat

individual sampling is needed to further support AstV infection in RFTB and identify population-related factors involved in the temporal and spatial transmission dynamics of AstVs.

Figure 1. Number of Astrovirus positive guano samples as per sampling session (brown marks). Green shading indicates the relative bat population size: light green: no bat inside the cave, mid-green: less than half of the cave covered with bats, dark green: more than half of the cave covered with bats. Presence of adults, newborns, and juveniles bats is represented by gray bars.

Two samples collected before the breeding season (in July and September), when the cave was free of bats, also tested positive for AstV, and suggest a possible environmental persistence of AstV RNA in bat guano. AstV persistence in the environment could potentially favor bat infection during cave colonization at the beginning of the breeding season, but also create opportunity for virus spillover to other hosts. Rats (*Rattus rattus*) have been sighted in and around the cave during the study period, and previous studies have shown that paramyxoviruses may be transmitted between rats and RFTB in the studied colony (Dietrich et al., 2015). Bat guano is also used as a fertilizer for agricultural crops, worldwide, and is regularly harvested on a small scale by local farmer for personal use (Mildenstein & de Jong, 2011). At our study site, bat guano was indeed harvested in the cave entrance throughout the course of the breeding season. The detection of viral RNA do not provide evidence of the maintenance of infectious particles in bat guano and virus isolation would be required to experimentally assess the infectious ability of bat AstVs outside their hosts. However, transmission routes of bat viruses to humans often remain unknown (Joffrin, Dietrich, Mavingui, & Lebarbenchon, 2018) and given the high propensity of AstVs for cross-species transmission

(Mendenhall et al., 2015) our finding underlines that the risk of AstV spillover from bats to human should be fully investigated.

Analysis of the 15 RdRp partial sequences revealed a high genetic diversity (up to 49% of partial RdRp sequence differences). This finding is consistent with previous studies performed in Madagascar (Lebarbenchon et al., 2017), continental Africa (Hoarau et al., 2018; Rougeron et al., 2016; Waruhiu et al., 2017; Yinda et al., 2018), and other bat populations, worldwide. Such a high level of sequence diversity in a single bat species is nevertheless surprising and higher than sequence diversity measured for paramyxovirus and *Leptospira* in the same colony (Dietrich et al., 2015). Current classification of AstV based on host species origin has been largely challenged for the past years (Donato & Vijaykrishna, 2017). First, because AstVs isolated from different animal species can be genetically similar (Hoarau et al., 2018; Lebarbenchon et al., 2017; Waruhiu et al., 2017), but also because genetically different viruses can be isolated from a single species (Waruhiu et al., 2017; Yi et al., 2018). To improve classification of AstV, the International Committee on Taxonomy of Virus now recommends to compare AstV RdRp gene to capsid gene sequences (Bosch et al., 2011).

The phylogenetic analysis further revealed that the detected viruses were genetically related to AstVs found in reptiles, dogs, and pigs, but did not cluster in a single bat AstVs clade (Figure 2). This result could reflect the limited number of AstV nucleotide sequences available from Molossid bats in public database and, more broadly, from bats in tropical areas. Because our study was based on environmental sampling, however, we cannot exclude that the detected viruses came from other hosts than RFTB. As previously mentioned, rats, but also feral dogs (*Canis lupus familiaris*), and zebu (*Bos taurus indicus*), have been regularly sighted close to the cave entrance. Avian species such as house sparrow (*Passer domesticus*), Madagascar turtle-dove (*Nesoenas picturatus*), and common myna (*Acridotheres tristis*) have also been observed near to the cave, and can be in direct contact with bat guano. Further investigations involving the sampling of individual bats and other host species in the cave surroundings, as well as sequencing of the AstV capsid gene, could provide information on the level of potential cross-species transmission in this ecosystem.

Figure 2. Maximum likelihood consensus tree derived from 94 Astrovirus (AstV) RNA-dependent RNA-polymerase partial nucleotide sequences (393 bp). Black dots indicate nodes with bootstrap values higher than 50. Red: sequences generated in this study. Scale bar: mean number of nucleotide substitutions per site.

Conflicts of interests

The authors have no conflict of interests.

Acknowledgments

We thank Muriel Dietrich and Nicolas Diotel for their assistance in the field. Léa Joffrin is supported by a "Région Réunion, European Regional Development Funds (FEDER 2014-2020)" PhD fellowship, Axel Hoarau by a "Ministère de l'Enseignement supérieur, de la Recherche et de l'Innovation" PhD fellowship, and Camille Lebarbenchon by a "Chaire Mixte Institut National de la Santé et de la Recherche Médicale (INSERM) – Université de La Réunion".

References

Bosch, A., Guix, S., Krishna, N. K., Méndez, E., Monroe, S. S., Pantin-Jackwood, M., & Schultz-Cherry, S. (2011). Virus Taxonomy: Ninth Report ofthe International Committee on Taxonomy ofViruses. In A. M. Q. King, E. Lefkowitz, M. J. Adams, & E. Carstens (Eds.), *Virus Taxonomy: Ninth Report ofthe International Committee on Taxonomy of Viruses*. San Diego: Elsevier.

Bosch, A., Pinto, R. M., & Guix, S. (2014). Human Astroviruses. *Clinical Microbiology Reviews*, 27(4), 1048–1074.

Chu, D. K. W., Leung, C. Y. H., Perera, H. K. K., Ng, E. M., Gilbert, M., Joyner, P. H., … Poon, L. L. M. (2012). A novel group of avian astroviruses in wild aquatic birds. *Journal of Virology*, 86(24), 13772–13778.

Chu, D. K. W., Poon, L. L. M., Guan, Y., & Peiris, J. S. M. (2008). Novel astroviruses in insectivorous bats. *Journal of Virology*, 82(18), 9107–9114.

De Benedictis, P., Schultz-Cherry, S., Burnham, A., & Cattoli, G. (2011). Astrovirus infections in humans and animals - molecular biology, genetic diversity, and interspecies transmissions. *Infection, Genetics and Evolution*, 11(7), 1529–1544.

Dietrich, M., Wilkinson, D. A., Benlali, A., Lagadec, E., Ramasindrazana, B., Dellagi, K., & Tortosa, P. (2015). Leptospira and paramyxovirus infection dynamics in a bat maternity enlightens pathogen maintenance in wildlife. *Environmental Microbiology*, 17(11), 4280–4289.

Donato, C., & Vijaykrishna, D. (2017). The broad host range and genetic diversity of mammalian and avian astroviruses. *Viruses*, 9(5), 1–18.

Drexler, J. F., Corman, V. M., Wegner, T., Tateno, A. F., Zerbinati, R. M., Gloza-Rausch, F., … Drosten, C. (2011). Amplification of emerging viruses in a bat colony. *Emerging Infectious Diseases*, 17(3), 449–456.

Fischer, K., dos Reis, V. P., & Balkema-buschmann, A. (2017). Bat Astroviruses: towards understanding the transmission dynamics of a neglected virus family. *Viruses*, 9, 34.

Goodman, S. M., Jansen van Vuuren, B., Ratrimomanarivo, F., Probst, J., Bowie, R., Group, E. G., & Re, L. (2008). Specific status of populations in the mascarene islands referred to Mormopterus acetabulosus (chiroptera: molossidae), with description of a new species. *Journal of Mammalogy*, 89(5), 1316–1327.

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., … Gascuel, O.
(2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing
the performance of PhyML 3.0. *Systematic Biology*, 59(3), 307–21.

Hoarau, F., Le Minter, G., Joffrin, L., Schoeman, M. C., Lagadec, E., Ramasindrazana, B., …
Lebarbenchon, C. (2018). Bat astrovirus in Mozambique. *Virology Journal*, 15(1), 1–5.

Joffrin, L., Dietrich, M., Mavingui, P., & Lebarbenchon, C. (2018). Bat pathogens hit the road: But
which one? *PLOS Pathogens*, 14(8), e1007134.

Karlsson, E. A., Small, C. T., Freiden, P., Feeroz, M. M., Matsen, F. A., San, S., … Schultz-Cherry,
S. (2015). Non-human primates harbor diverse mammalian and avian astroviruses including
those associated with human infections. *PLoS Pathogens*, 11(11), e1005225.

Keane, T. M., Creevey, C. J., Pentony, M. M., Naughton, T. J., & Mclnerney, J. O. (2006).
Assessment of methods for amino acid matrix selection and their use on empirical data shows
that ad hoc assumptions for choice of matrix are not justified. *BMC Evolutionary Biology*, 6,
29.

Lebarbenchon, C., Ramasindrazana, B., Joffrin, L., Bos, S., Lagadec, E., Le Minter, G., …
Mavingui, P. (2017). Astroviruses in bats, Madagascar. *Emerging Microbes & Infections*, 6,
e58.

Mendenhall, I. H., Skiles, M. M., Neves, E. S., Borthwick, S. A., Low, D. H. W., Liang, B., …
Smith, G. J. D. (2017). Influence of age and body condition on astrovirus infection of bats in
Singapore: An evolutionary and epidemiological analysis. *One Health*, 4, 27–33.

Mendenhall, I. H., Smith, G. J. D., & Dhanasekaran, V. (2015). Ecological drivers of virus
evolution: Astrovirus as a case study. *Journal of Virology*, 89(14), 6978–6981.

Mildenstein, T., & de Jong, C. (2011). Natural history, ecological and socio-economic value of bats.
In S. H. Newman, H. Field, J. Epstein, & C. de Jong (Eds.), *Investigating the role of bats in*
emerging zoonoses (FAO Animal, pp. 15–28). Rome: Food and Agriculture Organisation of the
United Nations.

R Core Team. (2018). R: A language and environment for statistical computing. *R Foundation for*
Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Rivera, R., Nollens, H. H., Venn-Watson, S., Gulland, F. M. D., & Wellehan, J. F. X. (2010). Characterization of phylogenetically diverse astroviruses of marine mammals. *Journal of General Virology*, 91(1), 166–173.

Rougeron, V., Suquet, E., Maganda, G. D., Jiolle, D., Mombo, I. M., Bourgarel, M., … Leroy, E. M. (2016). Characterization and phylogenetic analysis of new bat astroviruses detected in Gabon, Central Africa. *Acta Virologica*, 60, 386–392.

Seltmann, A., Corman, V., Rasche, A., Drosten, C., Czirjak, G., Bernard, H., … Voigt, C. (2017). Seasonal fluctuations of astrovirus, but not coronavirus shedding in bats inhabiting human-modified tropical forests. *Ecohealth*, 14, 272–284.

Waruhiu, C., Ommeh, S., Obanda, V., Agwanda, B., Gakuya, F., Ge, X. Y., … Shi, Z. L. (2017). Molecular detection of viruses in Kenyan bats and discovery of novel astroviruses, caliciviruses and rotaviruses. *Virologica Sinica*, 32(2), 101–114.

Yi, S., Niu, J., Wang, H., Dong, G., Guo, Y., Dong, H., … Hu, G. (2018). Molecular characterization of feline astrovirus in domestic cats from Northeast China. *PLOS ONE*, 13(10), e0205441.

Yinda, C. K., Ghogomu, S. M., Conceição-Neto, N., Beller, L., Deboutte, W., Vanhulle, E., … Matthijnssens, J. (2018). Cameroonian fruit bats harbor divergent viruses, including rotavirus H, bastroviruses, and picobirnaviruses using an alternative genetic code. *Virus Evolution*, 4(1), 1–15.