1   **DIVERSITY IN CRISPR-BASED IMMUNITY PROTECTS SUSCEPTIBLE**

2   **GENOTYPES BY RESTRICTING PHAGE SPREAD AND EVOLUTION**

3

4   **Abstract**

5

6   Diversity in host resistance often associates with reduced pathogen spread. This

7   may result from ecological and evolutionary processes, likely with feedback between

8   them. Theory and experiments on bacteria-phage interactions have shown that

9   genetic diversity of the bacterial adaptive immune system can limit phage evolution

10   to overcome resistance. Using the CRISPR-Cas bacterial immune system and lytic

11   phage, we engineered a host-pathogen system where each bacterial host genotype

12   could be infected by only one phage genotype. With this model system, we explored

13   how CRISPR diversity impacts the spread of phage when they can overcome a

14   resistance allele, how immune diversity affects the evolution of the phage to increase

15   its host range, and if there was feedback between these processes. We show that

16   increasing CRISPR diversity benefits susceptible bacteria via a dilution effect, which

17   limits the spread of the phage. We suggest that this ecological effect impacts the

18   evolution of novel phage genotypes, which then feeds back into phage population

19   dynamics.

Introduction

Introduction

Genetic diversity is a key determinant of the ecology and evolution of host-pathogen

systems. Various studies of wild organisms have shown that genetic diversity within

a host species often affects pathogen prevalence. The importance of diversity for

disease prevalence in wild populations has been observed in numerous species, for

example: cheetahs (O'Brien *et al.*, 1985); Italian agile frogs (Pearman & Garner,

2005); crimson rosella parrots (Eastwood *et al.*, 2017); inbred black-footed ferret

populations (Thorne & Williams, 1988); inbred California sea lions (Acevedo-

Whitehouse *et al.*, 2003); young island populations of deer mice (Meagher, 1999);

and Galapagos hawks (Whiteman *et al.*, 2007). The importance of diversity for

limiting disease in agricultural contexts has long been recognised (Elton, 1958), for

example in rice (Zhu *et al.*, 2000) and hybridising populations of honeybees (López-

Uribe *et al.*, 2017). In laboratory environments, more genetically diverse populations

of *Daphnia magna* are more resistant to parasites (Altermatt & Ebert, 2008), an

effect that depends on the genetic architecture of resistance (Luijckx *et al.*, 2013). In

microbial systems, *Pseudomonas aeruginosa* PA14 and *Streptococcus thermophilus*

with diverse immunity alleles were shown to be more resistant against lytic

bacteriophage (van Houte *et al.*, 2016b; Morley *et al.*, 2017). Two recent meta-

analyses have shown that host diversity is a robust defence against pathogens both

in agricultural (Reiss & Drinkwater, 2018) and natural populations (Ekroth *et al.*,

2019).

The suggested reasons for the effect of host diversity on pathogen spread can be

broadly divided into ecological and evolutionary effects. The ecological effect of

diversity may manifest through a dilution effect (Ostfeld & Keesing, 2012; Civitello *et*

al., 2015). The dilution effect, in terms of within-species immune diversity, is

suggested to arise because increasing the number of resistant or low-quality

individuals in a host population decreases the proportion of susceptible individuals

in that population. This can reduce contact rates between free-living pathogens and

susceptible hosts, which in turn limits the basic reproduction number of the pathogen

(R_0) (Dobson, 2004; Gandon, 2004; Lively, 2010). There is much observational

support for the role of a dilution effect in multi-species host-pathogen systems

(reviewed in (Civitello *et al.*, 2015)), and some experimental work has suggested that

dilution of susceptible host genotypes can limit pathogen spread in single host

species systems (Dennehy *et al.*, 2007; Common & Westra, 2019). Evolutionary

effects of diversity may manifest through limitation of the evolutionary emergence of

novel pathogen genotypes (Sasaki, 2000; Ohtsuki & Sasaki, 2006). Importantly

though, the evolutionary and ecological effects of host diversity are interdependent

as the basic reproductive value of a pathogen will influence its ability to evolve to

overcome host resistance (Antia *et al.*, 2003). Because increasing host diversity

dilutes susceptible hosts, pathogens may reach smaller population sizes and

generate fewer novel variants on which selection can act (Antia *et al.*, 2003;

Dennehy *et al.*, 2006; Yates *et al.*, 2006). Simultaneously, increasing host diversity

can increase selection for novel variants. These two effects can maximise

evolutionary emergence at intermediate host diversity (Benmayor *et al.*, 2009;

Chabas *et al.*, 2018b).

The interaction between lytic bacteriophage (phage) and the bacterial CRISPR-Cas

(Clustered Regularly Interspaced Short Palindromic Repeats; CRISPR-associated)

immune system is a tractable model system to study the evolutionary ecology of

infectious diseases, including the role of host diversity (van Houte *et al.*, 2016b;

Westra *et al.*, 2017; Chabas *et al.*, 2018b). CRISPR-Cas immune systems can

incorporate short DNA fragments (spacers) of about 30 base pairs derived from the

phage genome into CRISPR loci on the host genome (Horvath *et al.*, 2008).

Processed CRISPR transcripts guide Cas immune complexes to identify and cleave

the invading phage genome, preventing successful re-infections (Brouns *et al.*, 2008;

Marraffini & Sontheimer, 2008; Garneau *et al.*, 2010; Datsenko *et al.*, 2012; Westra

et al., 2012). In turn, phage can evolve to overcome CRISPR immunity by point

mutation in the sequence targeted by the spacer (protospacer) or in the protospacer-

adjacent motif (PAM), which flanks the protospacer and functions in self/non-self

discrimination (Deveau *et al.*, 2008; Mojica *et al.*, 2009; Semenova *et al.*, 2011;

Westra *et al.*, 2013). Phage evolution to overcome CRISPR immunity can lead to

CRISPR-phage coevolution (Paez-Espino *et al.*, 2013; Paez-Espino *et al.*, 2015; Sun

et al., 2016; Common *et al.*, 2019). CRISPR loci in both natural and experimental

populations can be highly diverse (Andersson & Banfield, 2008; Paez-Espino *et al.*,

2013; Westra *et al.*, 2015; Common *et al.*, 2019), due to different bacteria in the

population acquiring different spacers (Westra *et al.*, 2017). Diversity has important

implications for the coevolutionary interaction, as CRISPR diversity can provide

increased resistance by limiting the ability of phage to acquire the mutations needed

to overcome CRISPR immunity (van Houte *et al.*, 2016b; Morley *et al.*, 2017;

Common *et al.*, 2019). Phage mutation is limited by mutation supply (Lenski & Levin,

1985; Levin *et al.*, 2013; Chabas *et al.*, 2018a), and full phage infectivity requires

mutations in all the protospacers targeted by the host CRISPR array. This becomes

more difficult as individual-and population-level CRISPR diversity increases (Levin *et*

al., 2013; van Houte *et al.*, 2016a), and can drive rapid phage extinction (van Houte

et al., 2016b; Morley *et al.*, 2017; Chabas *et al.*, 2018b).

Apart from this evolutionary effect, theory predicts that even if a phage mutant

evolved that can overcome one CRISPR resistance allele in the population, its ability

to amplify will be reduced in a more diverse host population (Lively, 2010). The

resulting smaller phage population sizes are in turn predicted to reduce the ability of

the phage to evolve to overcome other CRISPR resistance alleles in the population

(Antia *et al.*, 2003; Chabas *et al.*, 2018b). However, these predictions remain

untested. We therefore set out to explicitly test how host diversity could limit

pathogen population size through a dilution effect, if and how this affects phage

evolution, and the degree of interdependence between these processes. Using the

bacteria *Pseudomonas aeruginosa* and its lytic phage DMS3vir, we performed an

experiment where we manipulated the degree of CRISPR diversity in the host

population by mixing bacterial genotypes that each carried a different CRISPR

spacer. We then infected these host populations with a phage that was pre-evolved

to infect only one host genotype. We then tracked changes in host fitness, and

phage population dynamics and evolution, over 3 days.

Results

To explore how population-level immune diversity would influence the population

dynamics and evolution of an infective phage and its susceptible host genotype, we

first developed a library of 24 *P. aeruginosa* PA14 bacteriophage-insensitive mutants

(BIMs), and a corresponding library of 24 DMS3vir mutants, each of which was pre-

evolved to infect only one of the 24 host BIMs (escape phage; **Fig. 1A**). We then set

up an experiment where we mixed 1, 3, 6, 12 or 24 BIMs, and inoculated each

mixture with ~10^6 of a single escape phage (**Fig. 1B**). Because this experimental

design increases the proportion of resistant bacterial hosts while decreasing the

proportion of the susceptible host (i.e. the one that could be infected by the escape

phage) (**Figure S1**), it enables us to explicitly test how this dilution effect is related to

the benefits of host CRISPR allele diversity. The susceptible clone always carried a

lacZ reporter gene, so we could follow its population dynamics and competitive

performance during the experiment (**Fig. 1B**). *P. aeruginosa* Δ*pilA*, an isogenic

mutant which does not have CRISPR immunity and resists phage infection via

surface receptor modification, was added to each host mixture (**Fig. 1B**). *P.*

aeruginosa Δ*pilA* was resistant to all phage, and allowed us to compare the fitness of

the CRISPR immune populations across the CRISPR allele diversity levels. We also

included 1- and 24-clone treatments inoculated with ancestral phage to which the

whole bacterial population was resistant. These treatments allowed us to examine

bacteria and phage dynamics at the two extremes of host diversity we tested in host

populations with no pre-existing susceptibility. We then compared those dynamics

with host mixtures that included a susceptible clone (van Houte *et al.*, 2016b)(**Fig.**

1B). We then monitored population dynamics and evolution of the phage, of the

clones with CRISPR immunity, and the susceptible fraction of the CRISPR

population over 3 days.

We hypothesised that if CRISPR diversity limits phage epidemics, then phage

density would be negatively correlated with CRISPR diversity. Phage densities

decreased as CRISPR diversity increased ($F_{6,342}$ = 48.30, p < 2.2 x 10^{-16}; **Fig. 2**),

phage population sizes decreased over the course of the experiment (regression of

ln plaque-forming units [pfu] ml^{-1} over days post-infection [dpi]: β [95% CI] = -1.32 [-

1.74, -0.94], $t(2)_{344}$ = -6.40, p = 5.06 x 10^{-10}), and phage declined faster with

increasing CRISPR diversity (interaction between CRISPR diversity and days post-

infection: $F_{6,314}$ = 24.10, p < 2.2 x 10^{-16}; **Fig. 2A & B**). This is consistent with an effect

of CRISPR diversity that protects the bacterial population from phage. Further,

phage titre in the 1-clone treatment infected with ancestral phage ["1-clone

(ancestral)" in **Fig. 2B**] is statistically similar to that observed in the 1-clone treatment

infected with an escape phage ["Intercept" in **Fig. 2B**], which is in line with previous

data showing that monocultures composed of hosts with CRISPR immunity allow

phage persistence due to rapid evolution of escape phage that overcome the

CRISPR resistance allele (van Houte *et al.*, 2016b). The evolution of escape phage

in our monoculture treatments likely caused the observed fluctuations in the density

of hosts with CRISPR immunity (**Figure S1**). When we explored the model further,

we found that phage titres in the 24-clone (ancestral phage) treatment were lower

than the 24-clone treatment infected with escape phage (difference in ln pfu ml^{-1}: β

[95% CI] = -2.40 [-4.67, -0.33], $t(2)_{342}$ = -2.25, p = 0.03; **Fig. 2B**). This is consistent

with a modest epidemic being able to establish by replicating on the susceptible

fraction of the host population in the 24-clone treatment infected with escape phage.

Next, we hypothesised that if phage population size across our CRISPR diversity

treatments was related to phage evolution, the proportion of phage that evolved an

expanded infectivity range would correlate with CRISPR diversity. Given that our

experiment was designed such that the proportion of the host population resistant to

the escape phage increased with CRISPR diversity, there would likely have been

strong selection to acquire mutations in other protospacers and PAMs to infect other

hosts in the population. The evolution of range expansion did depend on diversity

(Likelihood Ratio = 6.60, p = 0.01), and was most likely in the 6-clone treatment,

particularly at 3 days post-infection (dpi) (**Fig. 3**). This result suggests that

intermediate host diversity in the 6-clone treatment increased the likelihood that

novel escape phage with an expanded host range would evolve (Benmayor *et al.*,

2009; Chabas *et al.*, 2018b). The fact that these novel phage could infect more hosts

likely explains why phage population size was more consistent in the 6-clone

compared to the 3-clone treatment (**Fig. 2A**), and there was no significant difference

in phage population size between these treatments (**Fig. 2B**). Together, these data

suggest that the increased likelihood of evolutionary emergence at intermediate host

diversity balances against the potential limitation of reproduction caused by the

dilution effect. These novel phage variants could then maintain phage population

size.

Additionally, the presence of phage in some treatments was promoted by evolution

of a host shift, where infectivity on the original host is lost but phage evolved to infect

a new host. This is a less likely event as it involves two mutations: back-mutation to

the ancestral state at the original protospacer followed by mutation at the new

protospacer, all while incurring the cost of loss of infectivity on the original host. We

observed phage in two out of eight replicates of the 24-clone treatment had evolved

a host shift. Sequence data confirmed that single nucleotide polymorphisms (SNPs)

in the PAM that conferred infectivity to the original host had been lost, and novel

SNPs or deletions had emerged in protospacers that conferred infectivity to a new

host present in the population (**Table 4 & 5 in S1 Text**). Recall that phage titres in

the 24-clone (ancestral phage) treatment were lower than the 24-clone treatment

infected with escape phage (**Fig. 2B**). Together with the presence of phage that

evolved by host shift, these data suggest that the establishment of a modest phage

epidemic in this treatment may have facilitated the evolutionary emergence of novel

phage.

Given that CRISPR diversity negatively affected phage population size, we expected

that this would lead to enhanced fitness of all bacterial hosts with CRISPR immunity

relative to hosts without CRISPR immunity (van Houte *et al.*, 2016b). We therefore

hypothesised that the selection rate (Lenski, 1991; Travisano & Lenski, 1996) of all

CRISPR clones (both resistant and susceptible) compared to the Δ*pilA* strain (which

does not use CRISPR immunity) would be positively related to CRISPR diversity.

Despite some variation among replicates, we did find that CRISPR clones in all

polyclonal host populations had higher selection rates compared to clonal

populations during infection with escape phage (**Figs. 4A & B**). In 3, 3, and 4

replicates in the 12-, 24-, and 24-clone (ancestral phage) treatments, respectively,

the density of the Δ*pilA* strain dropped to zero (**Figure S1)**, suggesting that hosts

with CRISPR immunity could occasionally outcompete those without. However, the

selection rate of hosts with CRISPR immunity did not differ among the polyclonal

treatments ($F_{4,113}$ = 1.73, p = 0.15; **Fig. 4C**). Selection rate also did not change

notably over time (regression of selection rate over dpi: β [95% CI] = 0.03 [-0.04,

0.11], $t(2)_{118}$ = 0.94, p = 0.35) even though phage densities decreased with time.

The absence of a detectable relationship between the level of CRISPR diversity and

the selection rate hosts with CRISPR immunity relative to those without might be

because escape phage dynamics were dependent on proportion of sensitive bacteria

in the host population. We hypothesised that the selection rate of susceptible

CRISPR clones relative to resistant CRISPR clones would be positively related to

the level of CRISPR diversity, because the relative frequency of sensitive hosts

declines as CRISPR diversity increases. Indeed, in some replicates in the 3- and 6-

clone treatments the selection rate of the susceptible CRISPR clone was <-6,

indicating that they were driven extinct by phage (**Fig. 5A**). Further, the selection rate

of susceptible CRISPR clones in these treatments was on average less than zero

(mean selection rate [95% CI]; 3-clone: -1.53 [-2.61, -0.45]; 6-clone: -0.99 [-2.07, -

0.09]), indicating they tended to be outcompeted by resistant CRISPR clones.

Turning to the 12- and 24-clone treatments, the selection rate of susceptible clones

was statistically similar to zero (12-clone: 0.12 [-0.96, 1.21]; 24-clone: 0.15 [-0.93,

1.24]; **Fig. 5B & C**), and higher than the 3- and 6-clone treatments (**Fig. 5B & C**).

Together, these data show that the selection rate of susceptible CRISPR clones

increased with CRISPR diversity. There was also an increase in selection rate of

susceptible CRISPR clones over time among polyclonal treatments (regression of

selection rate over dpi: β [95% CI] = 0.41 [0.03, 0.80], $t(2)_{118}$ = 2.17, p = 0.03; **Fig.**

5A & B), which parallels the decline in phage population sizes over time in the same

treatments (regression of ln pfu ml^{-1} over dpi: β [95% CI] = -2.58 [-3.23, -1.84],

$t(2)_{114}$ = -7.64, p = 7.27 x 10^{-12}). These data suggest that susceptible hosts are

protected by population-level CRISPR diversity, and that this protection is mediated

by a reduced ability of phage to replicate on increasingly dilute susceptible hosts

(Payne *et al.*, 2018)(**Figure S1**).

Discussion

Previous studies have shown that CRISPR diversity can limit the evolution of phage

to overcome host resistance (van Houte *et al.*, 2016b; Chabas *et al.*, 2018b). In

those studies, bacterial populations were infected with ancestral phage that had not

been previously exposed to resistant hosts. Here, we examined the consequences of

CRISPR diversity once a phage has already evolved to overcome one of the

CRISPR resistance alleles in the population. This tractable system enabled us to

closely monitor how the level of CRISPR diversity influenced the population and

evolutionary dynamics of the phage, as well as the evolutionary dynamics of the

host.

These analyses show that phage population sizes were larger and persisted longer

in populations with lower CRISPR diversity, which is consistent with previous studies

(van Houte *et al.*, 2016b). Our results indicate that phage population size is

increasingly limited by host diversity, because increasing host diversity reduces the

proportion of susceptible individuals in the host population . This dilution effect of

host resistance diversity is an important factor in explaining why genetically diverse

host populations often have reduced pathogen loads (Keesing *et al.*, 2006; Ostfeld &

Keesing, 2012; Civitello *et al.*, 2015). The finding that resistant hosts can protect the

susceptible fraction of the population is also consistent with the epidemiological

concept of "herd immunity" (Fine *et al.*, 2011). While herd immunity has been

extensively studied in eukaryotes, recent work has suggested it may be important in

understanding bacteria-phage dynamics (Payne *et al.*, 2018), a suggestion that for

which our results provide further experimental support. This result also qualitatively

matches theoretical predictions of how host diversity should affect pathogen spread

in a system with matching allele genetic architecture (Lively, 2010).

In theory, this kind of ecological effect can in turn shape the evolutionary dynamics of

the bacteria-phage interaction, since smaller phage population sizes will decrease

the mutation supply and hence the evolutionary potential of the phage (Antia *et al.*,

2003; Dennehy *et al.*, 2006; Yates *et al.*, 2006). We did observe that the evolution of

phage host range expansion peaked at intermediate levels of CRISPR diversity. This

is likely because increasing CRISPR diversity dilutes susceptible hosts, which results

in smaller phage population size and hence less genetic variation on which selection

can act. Simultaneously, increasing host diversity can increase selection for escape

mutations. These two effects in dynamic tension can maximise evolutionary

emergence at intermediate host diversity (Chabas *et al.*, 2018b). The kind of

interdependence between ecological and evolutionary dynamics of phage suggested

by this study provides important context for other factors that can affect – and are

affected by - evolutionary emergence. For example, the ability of a phage to escape

a particular CRISPR spacer can influence host population structure by determining

colony heterogeneity (Pyenson & Marraffini, 2020). Another example is when

CRISPR immunity is incomplete, because phage with anti-CRISPR (*Acr*) genes can

vary in the degree to which they inhibit the CRISPR immune response (Landsberger

et al., 2018). Differences in the degree of CRISPR immunity because of *Acr* diversity

have been shown to affect the evolutionary emergence of escape phage

(Chevallereau *et al.*, 2020). Understanding of how these factors interact with

CRISPR allele diversity to affect the emergence and spread of escape phage will be

important in future studies of CRISPR ecology and evolution.

In this study, we focussed on host populations where different CRISPR resistance

genotypes were at equal starting frequencies, but natural communities are often

composed of a few very common and many rare variants (Pachepsky *et al.*, 2001;

McGill *et al.*, 2007). This likely matters for the observed dynamics, since the

proportion of susceptible hosts has a large impact on the probability of evolutionary

emergence of pathogens (Chabas *et al.*, 2018b). Also, we focussed our analysis on

the simple case where a diverse host population is infected by a clonal pathogen

population. In nature, pathogen populations will frequently be genetically diverse as

well (Hudson *et al.*, 2006; Telfer *et al.*, 2010), and increased levels of pathogen

diversity may affect the benefits of host diversity (Ganz & Ebert, 2010). Indeed,

previous studies of CRISPR-phage interactions suggest that infection by two

different phages can increase bacteria-phage coexistence compared to infections

with a single phage (Paez-Espino *et al.*, 2013; Paez-Espino *et al.*, 2015). Further,

infection by diverse phage can select against CRISPR immunity in favour of surface-

based resistance at the level of the host population, but also select for individual

CRISPR clones with a more diverse array of spacers (Broniewski *et al.*, 2020). The

empirical system used in this study offers both a unique ability to link genotypes and

phenotypes, tight experimental control over the infectivity matrix of the host-phage

interaction, and a clear link with theoretical predictions. These features will make it

an ideal system for more detailed studies to understand how the composition of the

host population and the relative diversity levels of the phage and host shape

coevolutionary interactions.

Materials & Methods

Bacterial strains and phage

Evolution experiments were carried out using *Pseudomonas aeruginosa* UCBPP-PA14 (which has two CRISPR loci, CRISPR1 and CRISPR2), UCBPP-PA14 Δ*pilA* (this strain lacks the pilus, which is the phage DMS3 receptor, and therefore displays surface-based instead of CRISPR-based resistance) and phage DMS3vir (Zegans *et al.*, 2009). We used *P. aeruginosa* UCBPP-PA14 *csy3::lacZ* (Cady *et al.*, 2012), which carries an inactive CRISPR-Cas system, for phage amplification, and for top lawns in phage spot and plaque assays. *P. aeruginosa* PA14 *csy3::lacZ, Escherichia coli* DH5α (NEB), *E. coli* CC118 λpir (NEB), and *E. coli* MFD*pir* (Ferrieres *et al.*, 2010) were used for molecular cloning.

Co-culture experiment

To control the levels of CRISPR diversity in our evolution experiments, we established a library of 24 *P. aeruginosa* PA14 clones each carrying a single spacer in CRISPR2 (bacteriophage-insensitive mutants; BIMs). We also independently evolved 24 phage mutants that could infect each one BIM (escape phage).

We designed 5 treatments in which we manipulated the level of CRISPR spacer diversity, based on the BIM library: monocultures (1-clone), or polycultures consisting of 3, 6, 12 and 24 clones. In order to monitor the population dynamics and relative fitness of individual bacterial clones within the polyculture treatments over

the course of the experiment, we transformed 8 BIMs to carry a *lacZ* reporter gene.

The *LacZ* gene encodes the β-galactosidase enzyme that hydrolyses X-gal, resulting

in the production of a blue pigment. For each of the polyclonal treatments, a single

BIM carrying the *lacZ* reporter gene was included. 8 BIMs were chosen for

transformation so that a single clone could be monitored in each of the 3-clone

mixtures (BIMs 1, 4, 7, 10, 13, 16, 19, and 22; see **Table S1**)

From fresh overnight cultures of each BIM, we made mixes of equal proportion of

each clone corresponding to the diversity treatments. To monitor the population

dynamics and competitive performance of all bacterial clones with an active CRISPR

system, we also added PA14 Δ*pilA* (which is fully resistant to phage DMS3vir via

surface-based resistance and has a distinct "smooth" colony morphology) to each

mix in equal proportion to the CRISPR-carrying fraction of the population. We then

inoculated 6ml of M9 minimal media (supplemented with 0.2% glucose; M9m) with

bacteria from the mixed overnight cultures at a dilution of 1:100. Approximately

1×10^6 pfu ml^{-1} of the escape phage targeting the labelled BIM were then added to

each vial. We also established 1- and 24-clone treatments with ancestral phage as

controls. Polyclonal treatments consisted of 8 biological replicates (*N*=8) to ensure

that both BIMs and phage were equally represented across treatments, while the 1-

clone treatments consisted of 24 biological replicates (*N*=24). Glass vials were

incubated at 37˚C while shaking at 180rpm. At 1, 2, and 3 days-post infection (dpi),

the sampling of the phage and bacterial culture was repeated as described below.

Cultures were transferred 1:100 to fresh media after sampling had been carried out.

The experiment was terminated at 3 dpi.

Sampling proceeded as follows. Each day 180µl of culture was taken from each vial

and phage was extracted using chloroform. Phage titres were determined by serially

diluting extracted phage in 1x M9 salts, and then spotting 5µl of each dilution on a

top lawn of *P. aeruginosa* PA14 *csy3::lacZ*, which was then incubated at 37°C for

24hrs. Phage titres were calculated from this assay. The detection limit of phage

spot assays is 10^2 pfu ml^{-1}. To monitor bacterial densities, culture samples were

serially diluted in 1x M9 salts, and then plated on LB agar + 40µg/ml X-gal + 0.1mM

IPTG, and incubated for 48hrs at 37°C. The density of SM, CRISPR and the labelled

BIM was then calculated. SM were differentiated from CRISPR clones by their

"smooth" colony morphology, and the labelled BIM was identified by the blue:white

screen.

We assessed the competitive performance of the all BIMs relative to the Δ*pilA* strain,

and the labelled BIM relative to non-labelled BIMs by calculating selection rate (*r*)

each day (*t*) from 1-3 (*n*) dpi (r_n = (ln [density strain A at t_n/density strain A at t_{n-1}] –

ln[density strain B at t_n/density strain B at t_{n-1}])/day) (Lenski, 1991; Travisano &

Lenski, 1996), which expresses competitive performance as the natural log of the

relative change in density of one competitor against another. We decided to

calculate selection rate instead of relative fitness (*W*) as used in Westra *et al.* (2015),

because *W* is very sensitive to starting densities and hence becomes increasingly

skewed at later timepoints. Applied to our data, it consequently produced impossibly

high values.

Phage evolution

We examined phage evolution during the experiment by sampling 12 individual

plaques from each replicate that had detectable levels of phage from 1 to 3 dpi,

which were amplified on PA14 *csy3::lacZ* overnight in LB, at 37˚C and shaking at

180rpm. Phage were extracted using chloroform, and then diluted 1000-fold.

Samples of each phage were then applied on lawns of each of the 24 BIMs and

PA14 *csy3::lacZ*. A successful infection was indicated by a clear lysis zone on the

top lawn. Phage were classified according to whether they had expanded their

infectivity range (could infect the original susceptible clone and a new clone in the

BIM library). Of the phages that had undergone a host shift (lost infectivity to the

original clone and could only infect a new clone), we confirmed their expanded

infectivity range by sequencing the old and new protospacers on the evolved phage

genome (SourceBioscience, UK). We also sequenced the relevant protospacers of

the pre-evolved escape phage from the BIM-phage library and ancestral DMS3vir.

Primers are given in Table S3.

Statistical analyses

All statistical analyses were carried out in R v3.5.3 (R Core Team, 2018).

Generalised linear mixed models (GLMMs) were used throughout, and replicate was

treated as a random effect in all models. Days post-infection (dpi) was treated as a

continuous variable and CRISPR allele diversity as a categorical variable. Model

selection followed a nested approach, where full versus reduced models were

compared using information criteria (Burnham & Anderson, 2003, 2004), as well as

focussed comparisons of observed and predicted values. The overall statistical

significance of the single and interaction effects of fixed effects (that is, treatment

and dpi) was assed using ANOVA, which gives an F- and associated p-value. Simple

Bayesian GLMMs from the MCMCglmm (Hadfield, 2010) package were used to

analyse phage evolution to overcome model conversion issues encountered with this

dataset. These models used a probit transformation (the inverse standard normal

distribution of the probability) and a flat prior. The overall statistical significance of

the single and additive effects of fixed effects in these models was assessed using

Likelihood Ratio (LR) tests and their associated p-values using the VCVglmm

package (Brown, 2019). Where possible, exact p-values are given, but R is unable to

give exact values when $p < 2.2 \times 10^{-16}$. When phage titre was considered as the

response variable, data was log-transformed to improve model fit. Credibility

intervals around model coefficients and predictions were calculated to the 95%, 89%

and 67% level to give the reader a clearer indication of effect size. All R code used

for analyses and figures is available at this manuscript's GitHub page.

Acknowledgements

We thank Meaghan Castledine for help constructing the pBAM1(Gm)_lacZ plasmid.

 Figure captions

Fig 1. Experimental design

(A) Infectivity matrix of the library of bacteriophage-insensitive mutants (BIMs) and

escape phages. The identity of BIMs and escape phage (1-24) are shown in the first

row and column, respectively. Black squares represent infectivity, white indicates no

infectivity. Infectivity of the wild-type DMS3vir is shown, as well as the infectivity of

each escape phage on wild-type *P. aeruginosa* PA14. (B) Design of the co-culture

experiment. The library of BIMs and escape phage is shown at the top. For each

CRISPR allele diversity treatment, the relevant numbers of BIMs and escape phage

were taken from the library and used to start the experiment, as shown by the

arrows. The number of BIMs is shown next to the arrows. Blue text indicates that the

BIM was engineered to carry a *lacZ* reporter gene, and that it was susceptible to the

escape phage. Black text indicates that the BIMs were resistant to the phage. The

number of replicates (*N*) is given underneath each treatment. SM represents the

surface mutant strain PA14 $\Delta pilA$.

Fig 2. Increasing CRISPR diversity limited phage population size

(A) Population dynamics of phage at different levels of CRISPR diversity in the host

population (panels). Black lines show the phage density expressed as plaque-

forming units (pfu) ml^{-1} in individual replicates at each day post-infection. The limit of

phage detection is 200 pfu ml^{-1}. (B) Coefficients from a GLMM of the natural log of

phage density (ln[pfu ml^{-1}]), with CRISPR allele diversity treatment and days post-

infection (dpi) as fixed effects. The intercept is the mean phage titre in the 1-clone

treatment at 0 dpi. The β differences from the intercept are shown for the remaining

levels of both fixed effects. The dotted line at zero indicates no difference from the

intercept. Means are shown as white points with 67, 89% and 95% credibility

intervals given in decreasing width.

Fig 3. Phage evolutionary emergence was maximised at intermediate CRISPR

diversity

Estimated proportion of phage that had evolved to infect a second CRISPR clone in

addition to the original CRISPR clone they were pre-evolved to infect at each day

post-infection in each polyclonal CRISPR diversity treatment. Because these

estimates are derived from a Bayesian GLMM, they are posterior modes - not means

– and are shown as white points. Similarly, 67, 89 and 95% credibility intervals are

highest probability density intervals, and are given in decreasing width around mode

estimates. Raw values from each replicate are shown as points.

Fig 4. Selection rate of bacterial hosts with CRISPR immunity did not change

with any level of CRISPR diversity

Selection rate of hosts with CRISPR immunity relative to a strain without CRISPR

immunity in (A) the 1-clone CRISPR diversity treatments with infective and ancestral

phage and (B) the remaining CRISPR diversity treatments. Days post-infection is

indicated by colour. Selection rate is the natural log of the relative change in density

of bacteria with CRISPR immunity (both susceptible and resistant to phage) against

a strain without. The dotted line at zero indicates no difference in density change i.e.

both are equally fit. Boxplots show the median, 25^{th} and 75^{th} percentile, and the

interquartile range. Raw values from each replicate are shown as points. (C)

Predicted mean selection rate is calculated from a GLMM that statistically controls

for the effect of time. Means are shown as white points with 67, 89 and 95%

credibility intervals given in decreasing width.

Fig 5. The selection rate of susceptible CRISPR clones increased with CRISPR

diversity and over time

Selection rate of the susceptible CRISPR clones relative to the resistant CRISPR

clones in (A) the 3- and 6-clone CRISPR diversity treatments and (B) the remaining

CRISPR diversity treatments. Days post-infection is indicated by colour. Selection

rate is the natural log of the relative change in density of CRISPR clones that were

susceptible to phage against CRISPR clones that were resistant to phage. The

dotted line at zero indicates no difference in density change i.e. both are equally fit.

Boxplots show the median, 25^{th} and 75^{th} percentile, and the interquartile range. Raw

values from each replicate are shown as points. The 1-clone treatments, both with

infective and ancestral phage, have been excluded because all bacterial clones with

CRISPR-based resistance in this treatment were susceptible, and the data is

therefore present in Fig. 4A. (C) Predicted mean selection rate is calculated from a

GLMM that statistically controls for the effect of time. Means are shown as white

points with 67, 89 and 95% credibility intervals given in decreasing width.

Fig. 1

 Fig. 2

A

Days post-infection

B

 Fig. 3

507   **Fig. 4**

 Fig. 5

References

Acevedo-Whitehouse, K., Gulland, F., Greig, D., & Amos, W. (2003). Inbreeding: disease susceptibility in California sea lions. *Nature, 422*(6927), 35.

Altermatt, F., & Ebert, D. (2008). Genetic diversity of Daphnia magna populations enhances resistance to parasites. *Ecol Lett, 11*(9), 918-928. doi:10.1111/j.1461-0248.2008.01203.x

Andersson, A. F., & Banfield, J. F. (2008). Virus population dynamics and acquired virus resistance in natural microbial communities. *Science, 320*(5879), 1047-1050. doi:10.1126/science.1157358

Antia, R., Regoes, R. R., Koella, J. C., & Bergstrom, C. T. (2003). The role of evolution in the emergence of infectious diseases. *Nature, 426*(6967), 658.

Benmayor, R., Hodgson, D. J., Perron, G. G., & Buckling, A. (2009). Host mixing and disease emergence. *Curr Biol, 19*(9), 764-767.

Broniewski, J. M., Meaden, S., Paterson, S., Buckling, A., & Westra, E. R. (2020). The effect of phage genetic diversity on bacterial resistance evolution. *The ISME Journal*, 1-9.

Brouns, S. J. J., Jore, M. M., Lundgren, M., Westra, E. R., Slijkhuis, R. J. H., Snijders, A. P. L., . . . van der Oost, J. (2008). Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes. *Science, 321*(5891), 960-964. doi:10.1126/science.1159689

Brown, M. (2019). VCVglmm: Manipulating model outputs from MCMCglmm and lme4 objects, usually involving variance-covariance matrices (Version 0.0.0.9). Retrieved from https://github.com/Euphrasiologist/VCVglmm

Burnham, K. P., & Anderson, D. R. (2003). *Model selection and multimodel inference: a practical information-theoretic approach*. New York: Springer.

Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: understanding AIC
and BIC in model selection. *Sociol Methods Res, 33*(2), 261-304.

Cady, K. C., Bondy-Denomy, J., Heussler, G. E., Davidson, A. R., & O'Toole, G. A.
(2012). The CRISPR/Cas Adaptive Immune System of *Pseudomonas*
aeruginosa Mediates Resistance to Naturally Occurring and Engineered
Phages. *J Bacteriol, 194*(21), 5728-5738. doi:10.1128/jb.01184-12

Chabas, H., Lion, S., Nicot, A., Meaden, S., van Houte, S., Moineau, S., . . . Gandon,
S. (2018a). Evolutionary emergence of infectious diseases in heterogeneous
host populations. *bioRxiv*, 317099.

Chabas, H., Lion, S., Nicot, A., Meaden, S., van Houte, S., Moineau, S., . . . Gandon,
S. (2018b). Evolutionary emergence of infectious diseases in heterogeneous
host populations. *PLoS Biol, 16*(9), e2006738.
doi:10.1371/journal.pbio.2006738

Chevallereau, A., Meaden, S., Fradet, O., Landsberger, M., Maestri, A., Biswas, A., .
. . Westra, E. R. (2020). Exploitation of the cooperative behaviors of anti-
CRISPR phages. *Cell host & microbe, 27*(2), 189-198. e186.

Civitello, D. J., Cohen, J., Fatima, H., Halstead, N. T., Liriano, J., McMahon, T. A., . .
. Young, S. (2015). Biodiversity inhibits parasites: broad evidence for the
dilution effect. *Proc Natl Acad Sci USA, 112*(28), 8667-8671.

Common, J., Morley, D., van Houte, S., & Westra, E. R. (2019). CRISPR-Cas
immunity leads to a coevolutionary arms race between *Streptococcus*
thermophilus and lytic phage. *Philos Trans Royal Soc B*.
doi:10.1098/rstb.2018.0098

Common, J., & Westra, E. R. (2019). CRISPR evolution and bacteriophage
persistence in the context of population bottlenecks. *RNA Biol*.

Datsenko, K. A., Pougach, K., Tikhonov, A., Wanner, B. L., Severinov, K., &
Semenova, E. (2012). Molecular memory of prior infections activates the
CRISPR/Cas adaptive bacterial immunity system. *Nat Commun, 3*, 945.

doi:http://www.nature.com/ncomms/journal/v3/n7/suppinfo/ncomms1937_S1.h

tml

Dennehy, J. J., Friedenberg, N. A., Holt, R. D., & Turner, P. E. (2006). Viral ecology

and the maintenance of novel host use. *Am Nat, 167*(3), 429-439.

Dennehy, J. J., Friedenberg, N. A., Yang, Y. W., & Turner, P. E. (2007). Virus

population extinction via ecological traps. *Ecol Lett, 10*(3), 230-240.

Deveau, H., Barrangou, R., Garneau, J. E., Labonté, J., Fremaux, C., Boyaval, P., . .

. Moineau, S. (2008). Phage Response to CRISPR-Encoded Resistance in

Streptococcus thermophilus. J Bacteriol, 190(4), 1390-1400.

doi:10.1128/jb.01412-07

Dobson, A. (2004). Population dynamics of pathogens with multiple host species.

Am Nat, 164(S5), S64-S78.

Eastwood, J. R., Ribot, R. F., Rollins, L. A., Buchanan, K. L., Walder, K., Bennett, A.

T., & Berg, M. L. (2017). Host heterozygosity and genotype rarity affect viral

dynamics in an avian subspecies complex. *Sci Rep, 7*(1), 13310.

Ekroth, A. K. E., Rafaluk-Mohr, C., & King, K. C. (2019). Host genetic diversity limits

parasite success beyond agricultural systems: a meta-analysis. *Proceedings*

of the Royal Society B: Biological Sciences, 286(1911), 20191811.

doi:doi:10.1098/rspb.2019.1811

Elton, C. S. (1958). *The ecology of invasions by animals and plants*. London:

Methuen & Co.

Ferrieres, L., Hémery, G., Nham, T., Guérout, A.-M., Mazel, D., Beloin, C., & Ghigo,

J.-M. (2010). Silent mischief: bacteriophage Mu insertions contaminate

products of *Escherichia coli* random mutagenesis performed using suicidal

transposon delivery plasmids mobilized by broad-host-range RP4 conjugative

machinery. *J Bacteriol, 192*(24), 6418-6427.

Fine, P., Eames, K., & Heymann, D. L. (2011). "Herd immunity": a rough guide. *Clin*

Infect Dis, 52(7), 911-916.

Gandon, S. (2004). Evolution of multihost parasites. *Evolution, 58*(3), 455-469.

Ganz, H. H., & Ebert, D. (2010). Benefits of host genetic diversity for resistance to
infection depend on parasite diversity. *Ecology, 91*(5), 1263-1268.
doi:10.1890/09-1243.1

Garneau, J. E., Dupuis, M.-E., Villion, M., Romero, D. A., Barrangou, R., Boyaval, P.,
. . . Moineau, S. (2010). The CRISPR/Cas bacterial immune system cleaves
bacteriophage and plasmid DNA. *Nature, 468*(7320), 67-71.
doi:http://www.nature.com/nature/journal/v468/n7320/abs/nature09523.html -
supplementary-information

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed
Models: The MCMCglmm R Package. *Journal of Statistical Software, 33*(2), 1-
22.

Horvath, P., Romero, D. A., Coute-Monvoisin, A. C., Richards, M., Deveau, H.,
Moineau, S., . . . Barrangou, R. (2008). Diversity, activity, and evolution of
CRISPR loci in *Streptococcus thermophilus*. *J Bacteriol, 190*(4), 1401-1412.
doi:10.1128/JB.01415-07

Hudson, P. J., Dobson, A. P., & Lafferty, K. D. (2006). Is a healthy ecosystem one
that is rich in parasites? *Trends Ecol Evol, 21*(7), 381-385.

Keesing, F., Holt, R. D., & Ostfeld, R. S. (2006). Effects of species diversity on
disease risk. *Ecol Lett, 9*(4), 485-498.

Landsberger, M., Gandon, S., Meaden, S., Rollie, C., Chevallereau, A., Chabas, H., .
. . van Houte, S. (2018). Anti-CRISPR phages cooperate to overcome
CRISPR-Cas immunity. *Cell, 174*(4), 908-916. e912.

Lenski, R. E. (1991). Quantifying fitness and gene stability in microorganisms.
Biotechnology, 15, 173-192.

Lenski, R. E., & Levin, B. R. (1985). Constraints on the coevolution of bacteria and
virulent phage: a model, some experiments, and predictions for natural
communities. *Am Nat, 125*(4), 585-602.

Levin, B. R., Moineau, S., Bushman, M., & Barrangou, R. (2013). The population and
evolutionary dynamics of phage and bacteria with CRISPR–mediated
immunity. *PLoS genetics, 9*(3), e1003312.

Lively, C. M. (2010). The Effect of Host Genetic Diversity on Disease Spread. *Am
Nat, 175*(6), E149-E152. doi:10.1086/652430

López-Uribe, M. M., Appler, R. H., Youngsteadt, E., Dunn, R. R., Frank, S. D., &
Tarpy, D. R. (2017). Higher immunocompetence is associated with higher
genetic diversity in feral honey bee colonies (Apis mellifera). *Conserv Genet,
18*(3), 659-666.

Luijckx, P., Fienberg, H., Duneau, D., & Ebert, D. (2013). A Matching-Allele Model
Explains Host Resistance to Parasites. *Current Biology*.

Marraffini, L. A., & Sontheimer, E. J. (2008). CRISPR interference limits horizontal
gene transfer in staphylococci by targeting DNA. *Science, 322*(5909), 1843-
1845.

McGill, B. J., Etienne, R. S., Gray, J. S., Alonso, D., Anderson, M. J., Benecha, H.
K., . . . He, F. (2007). Species abundance distributions: moving beyond single
prediction theories to integration within an ecological framework. *Ecol Lett,
10*(10), 995-1015.

Meagher, S. (1999). Genetic diversity and Capillaria hepatica (Nematoda)
prevalence in Michigan deer mouse populations. *Evolution, 53*(4), 1318-1324.

Mojica, F. J., Diez-Villasenor, C., Garcia-Martinez, J., & Almendros, C. (2009). Short
motif sequences determine the targets of the prokaryotic CRISPR defence
system. *Microbiology, 155*(Pt 3), 733-740. doi:10.1099/mic.0.023960-0

Morley, D., Broniewski, J. M., Westra, E. R., Buckling, A., & van Houte, S. (2017).
Host diversity limits the evolution of parasite local adaptation. *Mol Ecol, 26*(7),
1756-1763.

O'Brien, S. J., Roelke, M. E., Marker, L., Newman, A., Winkler, C., Meltzer, D., . . . Wildt, D. E. (1985). Genetic basis for species vulnerability in the cheetah. *Science, 227*(4693), 1428-1434.

Ohtsuki, A., & Sasaki, A. (2006). Epidemiology and disease-control under gene-for-gene plant–pathogen interaction. *J Theor Biol, 238*(4), 780-794.

Ostfeld, R. S., & Keesing, F. (2012). Effects of host diversity on infectious disease. *Annu Rev Ecol Evol Syst, 43*.

Pachepsky, E., Crawford, J. W., Bown, J. L., & Squire, G. (2001). Towards a general theory of biodiversity. *Nature, 410*(6831), 923.

Paez-Espino, D., Morovic, W., Sun, C. L., Thomas, B. C., Ueda, K.-i., Stahl, B., . . . Banfield, J. F. (2013). Strong bias in the bacterial CRISPR elements that confer immunity to phage. *Nat Commun, 4*, 1430.

Paez-Espino, D., Sharon, I., Morovic, W., Stahl, B., Thomas, B. C., Barrangou, R., & Banfield, J. F. (2015). CRISPR immunity drives rapid phage genome evolution in Streptococcus thermophilus. *MBio, 6*(2). doi:10.1128/mBio.00262-15

Payne, P., Geyrhofer, L., Barton, N. H., & Bollback, J. P. (2018). CRISPR-based herd immunity can limit phage epidemics in bacterial populations. *eLife, 7*, e32035.

Pearman, P. B., & Garner, T. W. (2005). Susceptibility of Italian agile frog populations to an emerging strain of Ranavirus parallels population genetic diversity. *Ecol Lett, 8*(4), 401-408.

Pyenson, N. C., & Marraffini, L. A. (2020). Co-evolution within structured bacterial communities results in multiple expansion of CRISPR loci and enhanced immunity. *eLife, 9*.

R Core Team. (2018). R: A language and environment for statistical computing (Version 3.5.2 "Eggshell Igloo"). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from http://www.r-project.org/

Reiss, E. R., & Drinkwater, L. E. (2018). Cultivar mixtures: a meta-analysis of the
effect of intraspecific diversity on crop yield. *Ecol Appl, 28*(1), 62-77.

Sasaki, A. (2000). Host-parasite coevolution in a multilocus gene-for-gene system.
Proceedings of the Royal Society B: Biological Sciences, 267(1458), 2183-
2188.

Semenova, E., Jore, M. M., Datsenko, K. A., Semenova, A., Westra, E. R., Wanner,
B., . . . Severinov, K. (2011). Interference by clustered regularly interspaced
short palindromic repeat (CRISPR) RNA is governed by a seed sequence.
Proc Natl Acad Sci USA, 108(25), 10098-10103.
doi:10.1073/pnas.1104144108

Sun, C. L., Thomas, B. C., Barrangou, R., & Banfield, J. F. (2016). Metagenomic
reconstructions of bacterial CRISPR loci constrain population histories. *ISME*
J, 10(4), 858-870. doi:10.1038/ismej.2015.162

Telfer, S., Lambin, X., Birtles, R., Beldomenico, P., Burthe, S., Paterson, S., &
Begon, M. (2010). Species interactions in a parasite community drive infection
risk in a wildlife population. *Science, 330*(6001), 243-246.

Thorne, E. T., & Williams, E. S. (1988). Disease and endangered species: the
black-footed ferret as a recent example. *Conserv Biol, 2*(1), 66-74.

Travisano, M., & Lenski, R. E. (1996). Long-term experimental evolution in
Escherichia coli. IV. Targets of selection and the specificity of adaptation.
Genetics, 143(1), 15-26.

van Houte, S., Ekroth, A. K., Broniewski, J. M., Chabas, H., Ashby, B., Bondy-
Denomy, J., . . . Buckling, A. (2016a). The diversity-generating benefits of a
prokaryotic adaptive immune system. *Nature, 532*(7599), 385.

van Houte, S., Ekroth, A. K. E., Broniewski, J. M., Chabas, H., Ashby, B., Bondy-
Denomy, J., . . . Westra, E. R. (2016b). The diversity-generating benefits of a
prokaryotic adaptive immune system. *Nature, 532*(7599), 385-388.
doi:10.1038/nature17436

Westra, E. R., Semenova, E., Datsenko, K. A., Jackson, R. N., Wiedenheft, B.,
Severinov, K., & Brouns, S. J. (2013). Type IE CRISPR-cas systems
discriminate target from non-target DNA through base pairing-independent
PAM recognition. *PLoS Genet, 9*(9), e1003742.

Westra, E. R., Sünderhauf, D., Landsberger, M., & Buckling, A. (2017). Mechanisms
and consequences of diversity-generating immune strategies. *Nature Reviews
Immunology, 17*(11), 719.

Westra, E. R., van Erp, P. B., Künne, T., Wong, S. P., Staals, R. H., Seegers, C. L., .
. . Severinov, K. (2012). CRISPR immunity relies on the consecutive binding
and degradation of negatively supercoiled invader DNA by Cascade and
Cas3. *Mol Cell, 46*(5), 595-605.

Westra, E. R., van Houte, S., Oyesiku-Blakemore, S., Makin, B., Broniewski,
Jenny M., Best, A., . . . Buckling, A. (2015). Parasite Exposure Drives
Selective Evolution of Constitutive versus Inducible Defense. *Curr Biol, 25*(8),
1043-1049. doi:10.1016/j.cub.2015.01.065

Whiteman, N. K., Kimball, R. T., & Parker, P. G. (2007). Co-phylogeography and
comparative population genetics of the threatened Galápagos hawk and three
ectoparasite species: ecology shapes population histories within parasite
communities. *Mol Ecol, 16*(22), 4759-4773.

Yates, A., Antia, R., & Regoes, R. R. (2006). How do pathogen evolution and host
heterogeneity interact in disease emergence? *Proceedings of the Royal
Society B: Biological Sciences, 273*(1605), 3075-3083.

Zegans, M. E., Wagner, J. C., Cady, K. C., Murphy, D. M., Hammond, J. H., &
O'Toole, G. A. (2009). Interaction between bacteriophage DMS3 and host
CRISPR region inhibits group behaviors of *Pseudomonas aeruginosa. J
Bacteriol, 191*(1), 210-219.

Zhu, Y., Chen, H., Fan, J., & Wang, Y. (2000). Genetic diversity and disease control
in rice. *Nature, 406*(6797), 718.