**Differential effects of farming practice on cuckoo bumblebee communities in relation to their hosts**

Charlotte E. Howard[1], Alexander J. Austin[1] and James D. J. Gilbert[1*]

[1]Department of Biological and Marine Science, University of Hull, Cottingham Road, Hull HU6 7RX, UK

*To whom correspondence should be addressed: james.gilbert@hull.ac.uk

## ABSTRACT

1. Bees are important for vital pollination of wild and crop plants, but are in decline worldwide. Intensification of agriculture is a major driver of bee decline. Organic farming practices are designed to limit environmental impacts of agriculture and can increase bee abundance and species diversity. However, studies have been heavily focused towards some guilds of bees, overlooking others. This includes social brood parasites, cuckoo bumblebees, an understudied bee lineage. Little is known about bumblebee host and cuckoo population dynamics, and the effects of farming practice on cuckoo bumblebees have never previously been evaluated.

2. To compare the effects of farming practice (organic vs conventional) on the abundance, species diversity, and community dissimilarity of cuckoo bumblebees and their hosts, we compared host and cuckoo community metrics across ten matched pairs of organic and conventional farms in Yorkshire, UK.

3. As found by many previous studies, host bumblebees were more abundant on organic farms than on conventional farms. Despite this, cuckoo bumblebees were equally abundant on both farm types. Contrary to prediction, community dissimilarity and species diversity were unaffected by farm type for both host and cuckoo communities.

4. **Synthesis and applications:** Results suggest that cuckoo bumblebee community metrics are not solely driven by host community metrics, and that cuckoos may respond differently from their hosts to differences among farming practices. This could, in turn, indicate that a unified management practice is not sufficient to conserve all bumblebee species.

## KEYWORDS

Bee conservation, Cuckoo Bumblebee, Community Metrics, Conventional farming, Nutrition, Organic farming, Pollinator decline, Social Brood Parasite.

## INTRODUCTION

Human induced habitat degradation has led to worldwide biodiversity declines (Collen et al., 2009). For example, intensive farming practices have led to landscape simplification (Adhikari et al., 2019), increased pesticide applications (Whitehorn et al., 2012), and reductions in non-crop plant diversity and semi-natural habitat (Goulson et al., 2015). Sustainable farming systems, such as organic farming, are designed to reduce the environmental impacts of agriculture and aim to reduce biodiversity declines (Hole et al., 2005). Organic farming is less intensive (Kennedy et al., 2013; Hole et al., 2005; Power et al., 2012; Whitehorn et al., 2012), tending to use smaller sized crop fields (Kennedy et al., 2013; Hole et al., 2005), alternatives to agrochemicals, and sympathetic management of non-crop habitat, such as field margins (Hole et al., 2005; Adhikari et al., 2019). However, studies investigating the effects upon biodiversity of less intensive farming practices, such as organic farming, have often given mixed results which vary between taxa, habitat types, and landscapes (Aude et al., 2004; Bengtsson et al., 2005; Hole et al., 2005; Fuller et al., 2005; Holzschuh et al., 2008; Le Féon et al., 2010; Brittain et al., 2010). The reasons behind these differences are still poorly understood, with few studies investigating this in detail (e.g. Fuller et al., 2005; Tuck et al., 2014).

Pollinators are economically and ecologically important as they provide crucial pollination services to both wild and crop plants (Power and Stout, 2011; Scriven et al., 2015; Goulson et al., 2015; Adhikari et al., 2019), but, as with many other animal groups, are vulnerable to landscape change (Hole et al., 2005; Ceballos et al., 2015). Modern agricultural intensification plays a major role in the decline of bees, some of the most important pollinators (Holzschuh et al., 2007; Holzschuh et al., 2008; Le Féon et al., 2010; Power and Stout, 2011; Adhikari et al., 2019). This creates large expanses of resource-poor areas for bees (Carvell et al., 2006; Ollerton et al., 2014; Goulson et al., 2015; Vaudo et al., 2015; Tscharntke et al., 2012; Vasseur et al., 2013; Henckel et al., 2015), which are wholly dependent on plants for their nutritional, energetic, and developmental requirements (Goulson et al., 2011).

Organic farming is sometimes associated with more diverse, species rich, and abundant bee communities compared to conventional farming. This difference has been attributed to increased flower cover and availability of floral resources, such as non-crop plants (Aude et al., 2004; Holzschuh et al., 2007; Rundlöf et al., 2008; Le Féon et al., 2010; Power and Stout, 2011). These are essential for bee nutrition (Brodschneider and Crailsheim, 2010; Whitehorn et al., 2012; Di Pasquale et al., 2013; Adhikari et al., 2019). However, studies have often grouped all bee guilds together (Holzschuh et al., 2007; Holzschuh et al., 2010;

Power and Stout, 2011) or studied guilds in isolation (Wintermantel et al., 2019; Nooten and

Rehan, 2019). The extent to which organic farming practices can benefit bees varies

between guilds; for example among honeybees, social bumblebees, and solitary bees (Le

Féon et al., 2010; Holzschuh et al., 2008), and at species-level within the *Bombus* genus

(Rundlöf et al., 2008). It has been suggested that this is due to varied foraging distances

(Holzschuh et al., 2008; Rundlöf et al., 2008) and nesting resource requirements of different

bee guilds (Rundlöf et al., 2008). Given bees' highly varied ecology (Goulson, 2003; Goulson

and Darvill, 2004; Lhomme and Hines, 2018), the assumption that all bee guilds will be

affected in the same way by farming and management practices may therefore not be valid

(Rundlöf et al., 2008). To our knowledge there have been very few studies of the mechanism

underlying differential effects of farming practices upon different guilds (e.g. Kennedy et al.,

2013), in line with a paucity of such studies generally.

To date, studies investigating the effects of organic farming practices have been heavily

focused on specific bee groups, such as honeybees and social bumblebees (Serrano and

Guerra-Sanz, 2006; Velthuis and Van Doorn, 2006; Holzschuh et al., 2008; Le Féon et al.,

2010; De Luca et al., 2013). As a result, some bee guilds in particular are underrepresented

in the literature (Wood et al., 2016; Lhomme and Hines, 2018). Among these

underrepresented guilds, obligate social brood parasites, cuckoo bumblebees, are amongst

the least well studied (Power and Stout, 2011; Waters et al., 2011; Lhomme and Hines,

2018). Cuckoo bumblebees are a monophyletic subgenus of the *Bombus* genus, called

*Psithyrus*, yet despite being closely related, their ecology varies a great deal from that of

their host (Cameron et al., 2007; Lhomme and Hines, 2018). Unlike other bumblebees,

*Psithyrus* species have lost the ability to produce workers and collect pollen (Lhomme and

Hines, 2018). Emergence of cuckoo bumblebees from dormancy is delayed in order to allow

time for the host to establish a colony and produce a worker generation (Lhomme and Hines,

2018). A cuckoo queen sneaks into a host colony, matching her chemical signals to those of

the target nest to avoid detection by the host workers (Lhomme and Hines, 2018). Following

this, the cuckoo attacks the host queen, most often killing her, then ejects the host eggs from

the nest and enslaves the host workers to care for the parasitic brood (Lhomme and Hines,

2018). *Psithyrus* species are important pollinators, and evidence suggests that they are

more at risk of extinction than their hosts (Kosior et al., 2007; Suhonen et al., 2015; Lhomme

and Hines, 2018).

Host-parasite relationships in hymenopterans, such as bees, can be influenced by factors

such as nutrition (Goodell, 2003; Di Pasquale et al., 2013; Maure et al., 2016), pesticides

(Pettis et al., 2012), and host availability (Mahmoudi et al., 2010; Kaspi et al., 2011), which

can affect parasitism success rates, and host recovery (Goodell, 2003; Di Pasquale et al.,
2013; Maure et al., 2016) among other effects. Since all these factors can be affected by
farming practice, it is likely that farming practice can mediate the host-parasite relationship
between cuckoo and host bumblebees. Despite this, the effect of farming practice on cuckoo
bumblebee diversity and abundance has never previously been investigated.
139
In this study we compare the effects of farming practice, organic and conventional, on the
community metrics of cuckoo bumblebees and their hosts, using captured samples of
bumblebees from ten matched pairs of organic and conventional farms in Yorkshire. Since
farming practice can affect the abundance, diversity, and species richness of bumblebees
(Rundlöf et al., 2008; Adhikari et al., 2019), it can be expected that this could also cause
knock-on effects to the cuckoo bumblebee community (Arneberg et al., 1998; Steffan-
Dewenter and Schiele, 2008; Suhonen et al., 2015). Firstly, we predicted that species
richness, diversity, and abundance of hosts, and cuckoos, will be higher on organic farms
than on conventional farms. Obligate brood parasites such as cuckoo bumblebees, are
entirely dependent on their hosts for reproduction, thus the host can be considered a
resource (Steffan-Dewenter and Schiele, 2008; Suhonen et al., 2016). Therefore, elements
of their community structure are likely to be driven by those of their host (Murray et al., 2009;
Suhonen et al., 2015; Suhonen et al., 2016). Therefore, we also predicted that cuckoo
bumblebee community metrics would be driven by the community metrics of their hosts.
154
155
156 **METHODS**
157
**Study sites**
Ten organic farms were identified in the East Riding of Yorkshire and North Yorkshire, UK.
Sampling had a paired design to reduce confounding factors due to variation in the local
environment (Holzschuh et al., 2008; Holzschuh et al., 2010; Kennedy et al., 2013). Organic
and conventional farms were paired together based on crop type, soil type, and locality
(Power et al. 2012) (see Figure 1 for farm locations), since these factors can influence the
effects of farming practice (Rundlöf et al., 2008; Power and Stout, 2011; Tuck et al., 2014).
All paired sites were at least four kilometres apart in order to minimise the risk of any bees
foraging at both locations (Zurbuchen et al. 2010). Sample collection took place in the arable
portion of each farm, except on the two pairs of pastoral farms where sampling took place
within the pasture. A single field was chosen as the sampling site for each area. Crop or crop
type of the sampling area was matched for each pair.
170

171
172
173  **Figure 1:** Study sites for transect sweep-net sampling of bees in East Riding of Yorkshire
174  and North Yorkshire, UK. Each site location is marked with a circle. Eight of the ten farm
175  pairs were arable or mixed arable/pastoral farms, two pairs were dairy/beef farms. Cereals
176  were the primary crop at all farm sites. Farms were matched by locality and soil type and
177  were at least 4km apart to minimise risks of any individuals foraging at both sites.
178
179

**Sample collection**

Samples were collected from May to September, 2016, between 09:00 and 16:00, on dry,

sunny days with wind speeds up to moderate (Forup and Memmott, 2005; Power and Stout,

2011). There were single hour-long visits to each site per month. This spread the sample

collection, minimising the effects on the environment and bee communities. Each sampling

site was visited five times in total. The order in which each farm within a pair was visited was

alternated every sampling session to ensure that a) farms within pairs were sampled at the

same time of day, and b) each farm was sampled during both the morning and afternoon

(Power and Stout, 2011). These methods reduced the effects of confounds such as weather

and time of day. At each visit of every site, sampling was carried out using a sweep net

along a transect line, pausing the stopwatch for sample processing (ensuring a full hour of
sampling effort). The transect was 'E' shaped, with the spine (20x1m) running along the
hedgerow/margin and the three arms (20x1m) stretching into the crop. This was walked at a
constant pace. This shape was chosen in order to sample a representative amount of each
habitat type; hedgerow, margin and crop. For example, as the largest proportion of the fields
were the crop, the largest proportion of sampling time was spent here. The location of the
first transect at each site was decided with a random number generator, and subsequent
visits were done in a clockwise fashion, with each margin sampled at least once. All bees
were stored individually in a 50ml falcon tube in a cool bag with ice to reduce stress. Bees
were euthanised by placing into a freezer at -20°C, and stored at -80°C. Bees were thawed
in the fridge before manual identification to species level under a light microscope using Falk
(2015).
202
Manual identification was used since the sample size was too large to use molecular
techniques for the entire sample. However, cuckoo bumblebee species and some host
bumblebee species are difficult to distinguish morphologically (lucorum complex: Bossert et
al., 2016; Psithyrus spp.: Falk, 2015). Therefore, DNA barcoding (Supplementary Methods)
of a subset of samples was used to confirm the accuracy of the manual identification to
improve confidence in the accuracy of the data used for the community statistics.
209
**Community statistics**
Community statistics were carried out using R software (R Core Team, 2017) with the
following packages; iNEXT (Hsieh et al. 2019), lme4 (Bates et al. 2015). This included
calculation of species richness, abundance, and Simpson's diversity index. Simpson's
diversity index was chosen as it is a good general-purpose statistic (Magurran, 2003) and is
a more compatible index to use with the iNEXT package than the Shannon's index.
Sample-size integrations of rarefaction and extrapolation of data with 'iNEXT' (Appendix 2)
were used to standardise and compare species richness and diversity of hosts and cuckoos
across farm type (Hsieh et al., 2016). Generalised linear mixed models (GLMMs) were used
to assess the effects of farm type and month on cuckoo (n=89) and host (n=1,247)
community metrics. Separate models were fitted with 'abundance', 'species richness' and
'Simpson's diversity index' as response variables, respectively. Each model had 'farm type'
and 'visit month' as predictors, and 'farm pairs' as a random effect. A non-metric
multidimensional scaling ordination (NMDS) plot on Bray-Curtis distance metrics was used
to visualise community dissimilarity.
225
226

# RESULTS

Out of the sample tested, DNA barcoding confirmed accurate identification of all host individuals except for two (96%, n=50) and all cuckoo bumblebees but 5 (94%, n=87). One sample matched to a species of water mould, *Peronospora flava*, and was discarded due to contamination.

## Abundance

We found 8 species of host bumblebee (n=1,247), along with 7 species of cuckoo bumblebee (n=89). Abundance of hosts and cuckoos varied over the months in the season (Figure 2B). For example, on organic farms, abundance of hosts peaked in July, but on conventional farms, appeared to increase linearly from May to September (GLMM containing farm type and visit month as predictors with farm pairs as a random factor, dropping the farm:month interaction, $\chi^2_{4,16}$=10.83, p< 0.05; dropping bee type:month, $\chi^2_{4,16}$=35.12,p<0.001; Figure 2B). Total abundance of cuckoos did not vary across farm type. Abundance of hosts was higher on organic farms than conventional farms (Figure 2A). Cuckoo abundance was driven mainly by two months in particular, July and August. Abundance of cuckoos was higher on organic farms in July, and higher on conventional farms in August, despite host abundance being lower on conventional farms in August. Cuckoos were less abundant than hosts in all five months (Figure 2B).

## Species richness and Simpson's diversity index

Species richness and Simpson's diversity index broadly reflected similar patterns through the months of the season for hosts and cuckoos (Figure 3A and 3C). Total richness and diversity of hosts communities were similar on organic and conventional farms, and the same was true for cuckoo communities (Figure 3B and 3D). Species richness and Simpson's diversity index differed between hosts and cuckoos, and changed across the months (species richness: GLMM containing farm type and visit month as predictors with farm pairs as a random factor, dropping the main effect of visit month, $\chi^2_{4,8}$=25.115, P<0.001; and bee type, $\chi^2_{1,9}$=24.033, P<0.001) (Simpson's diversity index: GLMM containing farm type and visit month as predictors with farm pairs as a random factor, dropping the main effect of visit month, $\chi^2_{4,8}$=24.087, P<0.001; and bee type, $\chi^2_{1,8}$=10.335, P=0.001; Figure 3). Temporal patterns in richness and diversity did not differ between farm types. On conventional farms, cuckoo richness and diversity peaked in August (Figure 3A and C). In contrast, hosts were most species-rich in July, but were more diverse in August (Figure 3A and 3C). On organic farms, richness and diversity of both hosts and cuckoos peaked in July (Figure 3A and 3C). In total across the whole season, hosts were more diverse and species-rich than the cuckoos (Figure 3B and 3D).

**Figure 2:** Abundance of cuckoo and host bumblebees found on organic and conventional farms in total (A) and by month (B). For clarity, a larger-scaled section of cuckoo bumblebee abundance across the months is shown in the top left of graph B for their two key months of activity, July and August.

273
274
275 **Figure 3:** Extrapolated species richness of cuckoo and host bumblebees found on organic
276 and conventional farms by month (A), and in total (B). Extrapolated Simpson's diversity
277 index of cuckoo and host bumblebees found on organic and conventional farms by month
278 (C), and in total (D). Sample-size integrations of rarefaction and extrapolation of data with
279 'iNEXT' (see text) were used to standardise and compare species richness and diversity.
280
281

## **Community Dissimilarity**

The NMDS plot shows that the species composition was very similar on organic

farms and conventional farms for all bumblebees, both hosts and cuckoos (Figure 4). For

cuckoos, there was almost a complete overlap of the bumblebee communities at each farm

type, as with the host communities (Figure 4). Cuckoo communities on organic farms

showed a very small amount of variation compared to cuckoo communities on conventional

farms (Figure 4B). The species composition of cuckoos on conventional farms was a subset

of the species composition of cuckoos on organic farms (Figure 4B). Splitting the data by

month revealed no particular patterns (data not shown).

294
295    **Figure 4:** A non-metric multidimensional scaling ordination (NMDS) plot on Bray-Curtis
296    distance metrics to visualise community dissimilarity of A) cuckoo bumblebees, B) host
297    bumblebees, collected from organic and conventional farms. Dark blue area shows
298    communities on organic farms, and the pale blue area shows communities on conventional
299    farms.
300

## **DISCUSSION**

There have been few previous studies that investigate why organic farming should not be universally beneficial to farmland wildlife; and none investigating the effect of farm type on the community structure of cuckoo bumblebees (Power and Stout, 2011; Lichtenberg et al., 2017). Interestingly, and contrary to prediction, cuckoo abundance was *not* primarily driven by the abundance of their hosts. As predicted, hosts were more abundant on organic farms than on conventional farms,but despite this, cuckoos were equally abundant across both farming practices. In both hosts and cuckoos, both diversity and species richness were similar on organic and conventional farms; both metrics showed broadly similar patterns across the months within the season, and the two guilds showed very similar species composition across farm types.

Perhaps the reason why cuckoos were equally abundant on both organic and conventional farms, despite their hosts being more abundant on organic farms, was due to decreased health and nutrition of hosts on conventional farms, and hence increased susceptibility to cuckoo usurpation. For example, the increased use of pesticides on conventional farms could have negative effects on bee health; pesticides, for instance, can impair immune response (Pettis et al., 2012), colony growth (Rundlöf et al., 2015), queen mortality (Scholer and Krischik, 2014), susceptibility to parasites (Pettis et al., 2012), and foraging behaviour (Yang et al., 2008; Henry et al., 2012). Hosts emerge earlier from overwintering than cuckoos, hence are active for a longer period of time within the season, thus are more likely to be exposed to insecticides than cuckoos (Botías et al., 2015; Lhomme and Hines, 2018). Similarly, unlike cuckoos, the hosts will forage for pollen in addition to eating nectar (Lhomme and Hines, 2018).

Additionally, since organic farms generally offer increased diversity and abundance of floral resources (Rundlöf et al., 2008; Le Féon et al., 2010; Power and Stout, 2011), bees on organic farms can benefit from a more nutritionally diverse diet (Carvell et al., 2006; Ollerton et al., 2014; Goulson et al., 2015; Vaudo et al., 2015). Not all plants produce pollen of sufficient nutritional quality to meet the requirements of bees (Filipiak, 2017a,b). Since bee larvae require specific nutrients for growth and development (Austin & Gilbert 2018), large quantities of food can not compensate for poor quality (Filipiak, 2018). Bumblebees are wholly dependent on flowers for their nutritional, energetic, and developmental requirements (Goulson et al., 2011). In bees, nutrition affects body size, immune defence (Alaux et al., 2010; DeGrandi-Hoffman et al., 2010), fecundity, longevity, individual health, colony health, and colony productivity (Brodschneider and Crailsheim, 2010; Di Pasquale et al., 2013).

Therefore, perhaps nutrition and health are linked with a host queen's ability to defend

herself from attacks by the large, heavily armoured *Psithyrus* queen (Lhomme and Hines,

2018; Fisher and Sampson, 1992). One study has investigated this idea in cuckoo

bumblebees (Cartar and Dill, 1991), and found that food scarcity increased susceptibility of

bumblebee hosts to brood parasitism by *Psithyrus insularis* (Cartar and Dill, 1991). While

this idea in cuckoo bumblebees has received almost no attention in the literature since,

similar results have been found for other parasitic Hymenoptera (Goodell, 2003; Di Pasquale

et al., 2013; Maure et al., 2016). Future studies could further investigate the effects of health

and nutrition on invasion success of *Psithyrus* species, for example using controlled diets

under laboratory conditions (Goodell, 2003). Such research would be critical to

understanding the complex, differential ecology and communities of hosts and cuckoos.

Studies investigating the effects of organic or other less intensive farming practices on bee

richness and diversity have found mixed results (Bengtsson et al., 2005; Hole et al., 2005;

Le Féon et al., 2010; Aude et al., 2004; Brittain et al., 2010). The nesting strategies and

specialisation of guilds means that the availability of the correct quantity and quality of

resources, spatially and temporally, are the key determinants for which species a landscape

can support (Tscharntke et al., 2005). Many studies have found increased species richness,

and diversity of bees and bumblebees on organic farms, and as with abundance, have

mainly attributed this to increased floral resources (Holzschuh et al., 2007; Holzschuh et al.,

2008; Le Féon et al., 2010). However, the results of this study found no difference in species

richness, diversity, or community dissimilarity across farm type. An explanation for this

finding could lay in which specific host species are found in the area. The majority of  host

bumblebees found in the UK are generalist feeders (Falk, 2015). Hence, perhaps increased

floral richness would not attract additional host species to the area, and therefore would not

attract additional cuckoo species either. All of the host species found on the conventional

farms were also found on the organic farms. Additionally, landscape effects must be

considered (Holzschuh et al., 2008; Rundlöf et al., 2008; Weibull et al., 2000), for example,

organic farming may not always show the same benefits to pollinators, if isolated within a

mosaic of conventional or other farms (Brittain et al., 2010). Organic farms in Yorkshire are

few, and are thus relatively isolated. Hence, perhaps the effects of organic farming practices

are masked by the surrounding landscape. Future studies could include landscape effects in

the analyses, for example by separating farm pairs into groups depending on habitat type

(Power and Stout, 2011).

Host bumblebees were more abundant on organic farms than on conventional farms. This is

supported by a wide body of literature (Aude et al., 2004; Morandin and Winston, 2005;

Rundlöf et al., 2008; Le Féon et al., 2010; Power and Stout, 2011). This result is commonly
assigned to reduced abundance and diversity of floral resources on conventional farms
(Aude et al., 2004; Holzschuh et al., 2007; Rundlöf et al., 2008; Le Féon et al., 2010; Power
and Stout, 2011; Whitehorn et al., 2012; Adhikari et al., 2019), often resulting from less
intensive agricultural practices (Bengtsson et al. 2005, Hole et al. 2005, Roschewitz et al.
2005; Gabriel et al. 2006; Holzschuh et al. 2007; Holzschuh et al., 2008; Austin and Gilbert,
2019). Hosts were also more abundant than cuckoos in total across the season. Generally,
macro-parasites have a lower population size than their host (Murray et al., 2009) and, in the
case of cuckoo bumblebees, emerge later in the season once the hosts have already
established a colony and produced workers (Erler and Lattorff, 2010; Suhonen et al., 2015;
Lhomme and Hines, 2018). Similarly, cuckoo bumblebees do not produce workers, hence
are naturally rarer than their hosts (Lhomme and Hines, 2018).
388
Cuckoo abundance across the season did not follow the same patterns as host abundance.
Despite a lower abundance of hosts on conventional farms, cuckoo bumblebees were
equally as abundant on organic farms as they were on conventional farms. This was not
expected, since for obligate parasites such as cuckoo bumblebees, the host can be
considered a resource (Suhonen et al., 2016), hence elements of their community structure
are likely driven by those of their host (Murray et al., 2009; Suhonen et al., 2015; Suhonen et
al., 2016). Kleptoparasites benefit from the resources gathered and constructed by their host
(Murray et al., 2009), therefore the cuckoo presumably gains a fitness benefit from an
increase in the resources available to their hosts. There have been very few studies on
cuckoo bumblebees (Lhomme and Hines, 2018), hence future studies could shed more light
into the relationship between cuckoo and host communities.
400
Despite receiving little attention in the literature (Williams and Osborne, 2009; Lhomme and
Hines, 2018), cuckoo bumblebees are important pollinators (Kosior et al., 2007; Lecocq et
al., 2011). As such, further research is needed to fully understand their ecology (Lhomme
and Hines, 2018) and evidence suggests that they are even more at risk of extinction than
their hosts (Suhonen et al., 2015). Whilst the ecology and community metrics of hosts will
affect cuckoo community metrics due to their reproductive dependency on their host
(Suhonen et al., 2016), the results suggest that perhaps cuckoos can respond differently
from their hosts to environmental factors, such as farming practice. Differential effects of
farming practices have previously been found among groups within the bee superfamily, for
example among honeybees, bumblebees and solitary bees (Le Féon et al., 2010; Holzschuh
et al., 2008) and are even found within the *Bombus* genus (Rundlöf et al., 2008). The results
of this study support the idea that perhaps hosts and cuckoo bumblebees need to be

considered and targeted separately when designing conservation measures and
management practices. There are more than two hundred species of bumblebee (Williams et
al., 2008), with 25 resident in the UK, within which there is great variation in colony size
(Rundlöf et al., 2008), feeding ecology, habitat, and reproduction (Goulson, 2003; Goulson
and Darvill, 2004; Lhomme and Hines, 2018). As such, the assumption that all guilds of bees
will be affected in the same way by farming and management practices may not be valid,
and the assumption that a unified management practice is sufficient to conserve all
bumblebee species may be naïve (Rundlöf et al., 2008).
421

422 ## **ACKNOWLEDGEMENTS**

423 The authors thank L. Handley for useful discussions on the manuscript and G. Sellers for
424 help with lab work.
425

**Authors' contributions**

All authors initially conceived the idea for this work. JG and AA designed the overall
infrastructure underpinning the study, while CH and AA designed the specifics of this
investigation. CH performed all molecular work, conducted all data analysis and wrote the
manuscript. AA performed all fieldwork and collected all field samples. All authors edited,
contributed to and approved the final manuscript.
432

433 **Ethics Statement**

434 This study was ethically approved by the University of Hull Faculty of Science & Engineering
435 Ethics Committee under licence FEC_55_2017_U_FEC.

## REFERENCES

ADHIKARI, S., BURKLE, L. A., O'NEILL, K. M., WEAVER, D. K. & MENALLED, F. D. 2019. Dryland organic farming increases floral resources and bee colony success in highly simplified agricultural landscapes. *Agriculture, Ecosystems & Environment,* 270**,** 9-18.

ALAUX, C., DUCLOZ, F., CRAUSER, D. & LE CONTE, Y. 2010. Diet effects on honeybee immunocompetence. *Biology letters,* 6**,** 562-565.

ARNEBERG, P., SKORPING, A., GRENFELL, B. & READ, A. F. 1998. Host densities as determinants of abundance in parasite communities. *Proceedings of the Royal Society of London. Series B: Biological Sciences,* 265**,** 1283-1289.

AUDE, E., ET AL. 2004. Conservation value of the herbaceous vegetation in hedgerows–does organic farming make a difference? *Biological conservation,* 118**,** 467-478.

AUSTIN, A., GILBERT, J.D.J. 2018. The geometry of dependence: solitary bee larvae prioritize carbohydrate over protein in parentally provided pollen. BioRxiv, https://doi.org/10.1101/397802

BATES, D., ET AL. 2015. Package 'lme4'. *Convergence,* 12.

BENGTSSON, J., AHNSTRÖM, J. & WEIBULL, A. C. 2005. The effects of organic agriculture on biodiversity and abundance: a meta-analysis. *Journal of applied ecology,* 42**,** 261-269.

BOSSERT, S., GEREBEN-KRENN, B.-A., NEUMAYER, J., SCHNELLER, B. & KRENN, H. W. 2016. The cryptic Bombus lucorum complex (Hymenoptera: Apidae) in Austria: Phylogeny, distribution, habitat usage and a climatic characterization based on COI sequence data. *Zool Stud,* 55**,** 2016.55-13.

BOTÍAS, C., BASLEY, K., NICHOLLS, E. & GOULSON, D. 2019. Impact of pesticide use on the flora and fauna of field margins and hedgerows. *The Ecology of Hedgerows and Field Margins***,** 90.

BUTCHART, S.H., ET AL. 2010. Global biodiversity: indicators of recent declines. *Science*, 328(5982), pp.1164-1168.

BRITTAIN, C., BOMMARCO, R., VIGHI, M., SETTELE, J. & POTTS, S. G. 2010. Organic farming in isolated landscapes does not benefit flower-visiting insects and pollination. *Biological Conservation,* 143**,** 1860-1867.

BRODSCHNEIDER, R. & CRAILSHEIM, K. 2010. Nutrition and health in honey bees*. *Apidologie,* 41**,** 278-294.

486      CAMERON, S., HINES, H. & WILLIAMS, P. 2007. A comprehensive phylogeny of the
487           bumble bees (*Bombus*). *Biological Journal of the Linnean Society,* 91**,** 161-
488           188.
489      CARTAR, R. V. & DILL, L. M. 1991. Costs of energy shortfall for bumble bee
490           colonies: predation, social parasitism, and brood development. *The Canadian*
491           *Entomologist,* 123**,** 283-293.
492      CARVELL, C., ET AL. 2006. Declines in forage availability for bumblebees at a
493           national scale. *Biological conservation,* 132**,** 481-489.
494      CEBALLOS, G., ET AL. 2015. Accelerated modern human–induced species losses:
495           Entering the sixth mass extinction. *Science advances,* 1**,** e1400253.
496      DARVILL, B., ET AL. 2010. Cryptic differences in dispersal lead to differential
497           sensitivity to habitat fragmentation in two bumblebee species. *Molecular*
498           *Ecology,* 19**,** 53-63.
499      DE LUCA, P. A., ET AL. 2013. Variability in bumblebee pollination buzzes affects the
500           quantity of pollen released from flowers. *Oecologia,* 172**,** 805-816.
501      DEGRANDI-HOFFMAN, G., CHEN, Y., HUANG, E. & HUANG, M. H. 2010. The
502           effect of diet on protein concentration, hypopharyngeal gland development
503           and virus load in worker honey bees (Apis mellifera L.). *Journal of insect*
504           *physiology,* 56**,** 1184-1191.
505      DI PASQUALE, G., ET AL. 2013. Influence of pollen nutrition on honey bee health:
506           do pollen quality and diversity matter? *PloS one,* 8**,** e72016.
507      ELMQVIST, T., ET AL. 2003. Response diversity, ecosystem change, and resilience.
508           *Frontiers in Ecology and the Environment,* 1**,** 488-494.
509      ERLER, S. & LATTORFF, H. M. G. 2010. The degree of parasitism of the bumblebee
510           (*Bombus terrestris*) by cuckoo bumblebees (*Bombus (Psithyrus) vestalis*).
511           *Insectes sociaux,* 57**,** 371-377.
512      FALK, S. 2015. Field guide to the bees of Great Britain and Ireland, (ed.) Bloomsbury
513           Publishing Plc.
514      FILIPIAK, M. ET AL. 2017. Ecological stoichiometry of the honeybee: Pollen diversity
515           and adequate species composition are needed to mitigate limitations imposed
516           on the growth and development of bees by pollen quality. *PLoS ONE,* 12, 8,
517           e0183236.
518      FILIPIAK, M.; WEINER, J. 2017. Plant-insect interactions: The role of ecological
519           stoichiometry. *Acta Agrobot*. 70, 1.
520      FISHER, R. M. & SAMPSON, B. J. 1992. Morphological specializations of the
521           bumble bee social parasite *Psithyrus ashtoni* (Cresson)(Hymenoptera:
522           Apidae). *The Canadian Entomologist,* 124**,** 69-77.

FOLMER, O., BLACK, M., HOEH, W., LUTZ, R. & VRIJENHOEK, R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular marine biology and biotechnology,* 3**,** 294-299.

FORUP, M. L. & MEMMOTT, J. 2005. The restoration of plant–pollinator interactions in hay meadows. *Restoration Ecology,* 13**,** 265-274.

FULLER, R., ET AL. 2005. Benefits of organic farming to biodiversity vary among taxa. *Biology letters,* 1**,** 431-434.

GILL, R. J. & RAINE, N. E. 2014. Chronic impairment of bumblebee natural foraging behaviour induced by sublethal pesticide exposure. *Functional Ecology,* 28**,** 1459-1471.

GOODELL, K. 2003. Food availability affects *Osmia pumila* (Hymenoptera: Megachilidae) foraging, reproduction, and brood parasitism. *Oecologia,* 134**,** 518-527.

GOULSON, D. 2003. *Bumblebees: their behaviour and ecology*, Oxford University Press, USA.

GOULSON, D. & DARVILL, B. 2004. Niche overlap and diet breadth in bumblebees; are rare species more specialized in their choice of flowers? *Apidologie,* 35**,** 55-63.

GOULSON, D., NICHOLLS, E., BOTÍAS, C. & ROTHERAY, E. L. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science,* 347**,** 1255957.

GOULSON, D., RAYNER, P., DAWSON, B. & DARVILL, B. 2011. Translating research into action; bumblebee conservation as a case study. *Journal of Applied Ecology,* 48**,** 3-8.

HENCKEL, L., BÖRGER, L., MEISS, H., GABA, S. & BRETAGNOLLE, V. 2015. Organic fields sustain weed metacommunity dynamics in farmland landscapes. *Proceedings of the Royal Society B: Biological Sciences,* 282**,** 20150002.

HENRY, M., ET AL. 2012. A common pesticide decreases foraging success and survival in honey bees. *Science,* 336**,** 348-350.

HOLDAWAY, R. J., ET AL. 2017. Using DNA metabarcoding to assess New Zealand's terrestrial biodiversity. *New Zealand Journal of Ecology,* 41**,** 251-262.

HOLE, D., ET AL. 2005. Does organic farming benefit biodiversity? *Biological conservation,* 122**,** 113-130.

HOLZSCHUH, A., STEFFAN-DEWENTER, I., KLEIJN, D. & TSCHARNTKE, T. 2007.
Diversity of flower-visiting bees in cereal fields: effects of farming system,
landscape composition and regional context. *Journal of Applied Ecology,* 44**,**
41-49.

HOLZSCHUH, A., STEFFAN-DEWENTER, I. & TSCHARNTKE, T. 2008. Agricultural
landscapes with organic crops support higher pollinator diversity. *Oikos,* 117**,**
354-361.

HOLZSCHUH, A., STEFFAN-DEWENTER, I. & TSCHARNTKE, T. 2010. How do
landscape composition and configuration, organic farming and fallow strips
affect the diversity of bees, wasps and their parasitoids? *Journal of Animal*
*Ecology,* 79**,** 491-500.

HSIEH, T., MA, K., CHAO, A. & HSIEH, M. T. 2019. Package 'iNEXT'.

HSIEH, T. C., MA, K. H. & CHAO, A. 2016. iNEXT: an R package for rarefaction and
extrapolation of species diversity (Hill numbers). *Methods in Ecology and*
*Evolution,* 7**,** 1451-1456.

KASPI, R., YUVAL, B. & PARRELLA, M. P. 2011. Anticipated host availability
increases parasitoid host attack behaviour. *Animal behaviour,* 82**,** 1159-1165.

KENNEDY, C. M., ET AL. 2013. A global quantitative synthesis of local and
landscape effects on wild bee pollinators in agroecosystems. *Ecology letters,*
16**,** 584-599.

KLEIJN, D., ET AL. 2006. Mixed biodiversity benefits of agri-environment schemes in
five European countries. *Ecology letters,* 9**,** 243-254.

KOSIOR, A., ET AL. 2007. The decline of the bumble bees and cuckoo bees
(Hymenoptera: Apidae: *Bombini*) of Western and Central Europe. *Oryx,* 41**,**
79-88.

LE FÉON, V., ET AL. 2010. Intensification of agriculture, landscape composition and
wild bee communities: a large scale study in four European countries.
*Agriculture, Ecosystems & Environment,* 137**,** 143-150.

LHOMME, P. & HINES, H. M. 2018. Ecology and Evolution of Cuckoo Bumble Bees.
*Annals of the Entomological Society of America*.

LICHTENBERG, E. M., ET AL. 2017. A global synthesis of the effects of diversified
farming systems on arthropod diversity within fields and across agricultural
landscapes. *Global change biology,* 23**,** 4946-4957.

MA, X., CHEN, L., ZHU, Q., CHEN, Y. & LIU, Y.-G. 2015. Rapid decoding of
sequence-specific nuclease-induced heterozygous and biallelic mutations by
direct sequencing of PCR products. *Molecular plant,* 8**,** 1285-1287.

MAES, J., PARACCHINI, M., ZULIAN, G., DUNBAR, M. & ALKEMADE, R. 2012.
Synergies and trade-offs between ecosystem service supply, biodiversity, and
habitat conservation status in Europe. *Biological conservation,* 155**,** 1-12.
MAGURRAN, A. E. 2003. Measuring biological diversity. Oxford :: Blackwell.
MAHMOUDI, M., SAHRAGARD, A. & SENDI, J. J. 2010. Effects of age and host
availability on reproduction of *Trioxys angelicae Haliday* (Hymenoptera:
Braconidae) parasitizing *Aphis fabae Scopoli* (Hemiptera: Aphididae). *Journal*
*of pest science,* 83**,** 33-39.
MAURE, F., THOMAS, F., DOYON, J. & BRODEUR, J. 2016. Host nutritional status
mediates degree of parasitoid virulence. *Oikos,* 125**,** 1314-1323.
MOFFAT, C., ET AL. 2015. Chronic exposure to neonicotinoids increases neuronal
vulnerability to mitochondrial dysfunction in the bumblebee (*Bombus*
*terrestris*). *The FASEB Journal,* 29**,** 2112-2119.
MORANDIN, L. A. & WINSTON, M. L. 2005. Wild bee abundance and seed
production in conventional, organic, and genetically modified canola.
*Ecological applications,* 15**,** 871-881.
MURRAY, T. E., KUHLMANN, M. & POTTS, S. G. 2009. Conservation ecology of
bees: populations, species and communities. *Apidologie,* 40**,** 211-236.
NYSTRÖM, M. 2006. Redundancy and response diversity of functional groups:
implications for the resilience of coral reefs. *AMBIO: A Journal of the Human*
*Environment,* 35**,** 30-35.
OLLERTON, J., ERENLER, H., EDWARDS, M. & CROCKETT, R. 2014. Extinctions
of aculeate pollinators in Britain and the role of large-scale agricultural
changes. *Science,* 346**,** 1360-1362.
PETTIS, J. S., JOHNSON, J. & DIVELY, G. 2012. Pesticide exposure in honey bees
results in increased levels of the gut pathogen *Nosema*. *Naturwissenschaften,*
99**,** 153-158.
POWER, E. F., KELLY, D. L. & STOUT, J. C. 2012. Organic farming and landscape
structure: effects on insect-pollinated plant diversity in intensively managed
grasslands. *PLoS One,* 7**,** e38073.
POWER, E. F. & STOUT, J. C. 2011. Organic dairy farming: impacts on insect–
flower interaction networks and pollination. *Journal of Applied Ecology,* 48**,**
561-569.
R CORE TEAM. 2017. R: A language and environment for statistical computing. R
Foundation for    Statistical Computing, Vienna, Austria. URL https://www.R-
project.org/.

ROSCHEWITZ, I., GABRIEL, D., TSCHARNTKE, T. & THIES, C. 2005. The effects
of landscape complexity on arable weed species diversity in organic and
conventional farming. *Journal of Applied Ecology,* 42**,** 873-882.

Roulston, T. H., Cane, J. H., & Buchmann, S. L. 2000. What governs protein content
of pollen: Pollinator preferences, pollen-pistil interactions, or phylogeny?
*Ecological Monographs*, 70(4), 617–643.

RUNDLÖF, M., ET AL. 2015. Seed coating with a neonicotinoid insecticide
negatively affects wild bees. *Nature,* 521**,** 77.

RUNDLÖF, M., NILSSON, H. & SMITH, H. G. 2008. Interacting effects of farming
practice and landscape context on bumble bees. *Biological Conservation,*
141**,** 417-426.

SANGER, F., NICKLEN, S. & COULSON, A. R. 1977. DNA sequencing with chain-
terminating inhibitors. *Proceedings of the national academy of sciences,* 74**,**
5463-5467.

SCHOLER, J. & KRISCHIK, V. 2014. Chronic exposure of imidacloprid and
clothianidin reduce queen survival, foraging, and nectar storing in colonies of
Bombus impatiens. *PLoS One,* 9**,** e91573.

SCRIVEN, J. J., ET AL. 2015. Revealing the hidden niches of cryptic bumblebees in
Great Britain: implications for conservation. *Biological Conservation,* 182**,**
126-133.

SERRANO, A. R. & GUERRA-SANZ, J. M. 2006. Quality fruit improvement in sweet
pepper culture by bumblebee pollination. *Scientia Horticulturae,* 110**,** 160-
166.

SIMON-DELSO, N., ET AL. 2015. Systemic insecticides (neonicotinoids and fipronil):
trends, uses, mode of action and metabolites. *Environmental Science and*
*Pollution Research,* 22**,** 5-34.

STEFFAN-DEWENTER, I. & SCHIELE, S. 2008. Do resources or natural enemies
drive bee population dynamics in fragmented habitats. *Ecology,* 89**,** 1375-
1387.

STEWART, S. D., ET AL. 2014. Potential exposure of pollinators to neonicotinoid
insecticides from the use of insecticide seed treatments in the mid-southern
United States. *Environmental science & technology,* 48**,** 9762-9769.

SUHONEN, J., RANNIKKO, J. & SORVARI, J. The rarity of host species affects the
co-extinction risk in socially parasitic bumblebee Bombus (Psithyrus) species.
Annales Zoologici Fennici, 2015. BioOne, 236-242.

SUHONEN, J., RANNIKKO, J. & SORVARI, J. 2016. Species richness of cuckoo
bumblebees is determined by the geographical range area of the host
bumblebee. *Insect Conservation and Diversity,* 9**,** 529-535.
TRUETT, G., HEEGER, P., MYNATT, R., TRUETT, A., WALKER, J. & WARMAN, M.
2000. Preparation of PCR-quality mouse genomic DNA with hot sodium
hydroxide and tris (HotSHOT). *Biotechniques,* 29**,** 52-54.
TSCHARNTKE, T., KLEIN, A. M., KRUESS, A., STEFFAN-DEWENTER, I. & THIES,
C. 2005. Landscape perspectives on agricultural intensification and
biodiversity–ecosystem service management. *Ecology letters,* 8**,** 857-874.
TSCHARNTKE, T., ET AL. 2012. Landscape moderation of biodiversity patterns and
processes-eight hypotheses. *Biological reviews,* 87**,** 661-685.
TSVETKOV, N., ET AL. 2017. Chronic exposure to neonicotinoids reduces honey
bee health near corn crops. *Science,* 356**,** 1395-1397.
TUCK, S. L., ET AL. 2014. Land-use intensity and the effects of organic farming on
biodiversity: a hierarchical meta-analysis. *Journal of Applied Ecology,* 51**,**
746-755.
VASSEUR, C., ET AL. 2013. The cropping systems mosaic: how does the hidden
heterogeneity of agricultural landscapes drive arthropod populations?
*Agriculture, ecosystems & environment,* 166**,** 3-14.
VAUDO, A. D., TOOKER, J. F., GROZINGER, C. M. & PATCH, H. M. 2015. Bee
nutrition and floral resource restoration. *Current Opinion in Insect Science,*
10**,** 133-141.
VELTHUIS, H. H. & VAN DOORN, A. 2006. A century of advances in bumblebee
domestication and the economic and environmental aspects of its
commercialization for pollination. *Apidologie,* 37**,** 421-451.
WARNER, P. A., OPPEN, M. J. H. & WILLIS, B. L. 2015. Unexpected cryptic species
diversity in the widespread coral Seriatopora hystrix masks spatial-genetic
patterns of connectivity. *Molecular Ecology,* 24**,** 2993-3008.
WATERS, J., DARVILL, B., LYE, G. C. & GOULSON, D. 2011. Niche differentiation
of a cryptic bumblebee complex in the Western Isles of Scotland. *Insect*
*Conservation and Diversity,* 4**,** 46-52.
WEIBULL, A. C., BENGTSSON, J. & NOHLGREN, E. 2000. Diversity of butterflies in
the agricultural landscape: the role of farming system and landscape
heterogeneity. *Ecography,* 23**,** 743-750.
WHITEHORN, P. R., O'CONNOR, S., WACKERS, F. L. & GOULSON, D. 2012.
Neonicotinoid pesticide reduces bumble bee colony growth and queen
production. *Science***,** 1215025.

WICKHAM, H., CHANG, W. & WICKHAM, M. H. 2016. Package 'ggplot2'. *Create*
*Elegant Data Visualisations Using the Grammar of Graphics. Version,* 2**,** 1-
189.
WILLIAMS, P. H., AN, J. & HUANG, J. 2011. The bumblebees of the subgenus
Subterraneobombus: integrating evidence from morphology and DNA
barcodes (Hymenoptera, Apidae, Bombus). *Zoological Journal of the Linnean*
*Society,* 163**,** 813-862.
WILLIAMS, P. H., ET AL. 2012. Unveiling cryptic species of the bumblebee
subgenus Bombus s. str. worldwide with COI barcodes (Hymenoptera:
Apidae). *Systematics and Biodiversity,* 10**,** 21-56.
WILLIAMS, P. H., BYVALTSEV, A., SHEFFIELD, C. & RASMONT, P. 2013. *Bombus*
*cullumanus*—an extinct European bumblebee species? *Apidologie,* 44**,** 121-
132.
WILLIAMS, P. H., CAMERON, S. A., HINES, H. M., CEDERBERG, B. & RASMONT,
P. 2008. A simplified subgeneric classification of the bumblebees (genus
*Bombus*). *Apidologie,* 39**,** 46-74.
WILLIAMS, P. H. & OSBORNE, J. L. 2009. Bumblebee vulnerability and
conservation world-wide. *Apidologie,* 40**,** 367-387.
WOOD, T.J., HOLLAND, J.M., GOULSON, D., 2016. Providing foraging resources
for solitary bees on farmland: current schemes for pollinators benefit a limited
suite of species. *J. Appl. Ecol.,* 54: 323-333.
YANG, E., CHUANG, Y., CHEN, Y. & CHANG, L. 2008. Abnormal foraging behavior
induced by sublethal dosage of imidacloprid in the honey bee (Hymenoptera:
Apidae). *Journal of economic entomology,* 101**,** 1743-1748.
ZURBUCHEN, A., ET AL. 2010. Maximum foraging ranges in solitary bees: only few
individuals have the capability to cover long foraging distances. *Biological*
*Conservation,* 143**,** 669-676.