Title: Loss of Piccolo function in rats induces Pontocerebellar Hypoplasia type 3-like phenotypes

Abbreviated title: *Piccolo loss of function triggers PCH3 phenotypes*

Author names and affiliations:

Joanne Falck[1], Christine Bruns[1], Sheila Hoffmann[1], Isabelle Straub[2], Erik J. Plautz[3], Marta Orlando[4], Humaira Munawar[5], Marion Rivalan[4,5], York Winter[4,5], Zsuzsanna Izsvák[6], Dietmar Schmitz[4], F. Kent Hamra[7], Stefan Hallermann[2], Craig Garner[1*] and Frauke Ackermann[1*]

1) German Center for Neurodegenerative Diseases (DZNE), Charité Medical University, Charitéplatz 1, 10117 Berlin, Germany

2) Carl-Ludwig Institute for Physiology, 04103, Leipzig, Germany

3) Department of Neurology and Neurotherapeutics, University of Texas Southwestern, Dallas Texas 75390

4) Charité Medical University Berlin, corporate member of Free University Berlin and Humboldt University Berlin; Berlin Institute of Health, NeuroCure Cluster of Excellence, Charitéplatz 1, 10117 Berlin, Germany

5) Department of Biology, Humboldt University, Philippstr. 13, 10099 Berlin, Germany

6) Max Delbrück Center for Molecular Medicine in the Helmholtz Society, 13125, Berlin, Germany

7) Department of Obstetrics and Gynecology, Cecil H. & Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern, Dallas Texas 75390

*equal contribution

Corresponding author email address: Frauke Ackermann: frauke.ackermann@dzne.de and Craig Garner: craig.garner@dzne.de, German Center for Neurodegenerative Diseases (DZNE), Charité Medical University, Charitéplatz 1, 10117 Berlin, Germany.

Conflict of interest statement: The authors declare no competing financial interests.

Number of pages: 46

Number of figures: 9

Number of words: abstract 248, introduction 609, discussion 1499

Acknowledgments: We would like to thank Susanne Wegmann and Eckart Gundelfinger for discussion and valuable comments on the manuscript; Anny Kretschmer and Katja Czieselsky for technical assistance. The work was supported by German Center for Neurodegenerative Diseases (DZNE), the Federal Government of Germany (DFG) SFB958 to CCG and (DFG) EXC 257 for the Center of Excellence NeuroCure to YW. Work to generate $Pclo^{gt/gt}$ rats was supported by National institute of Health R24RR03232601 & R24OD011108 to FKH. Neurological analyses on Piccolo mutant rats were conducted by The Neuro-Models Facility (EJP, LBG) at UT Southwestern Medical Center, and supported by the Haggerty Center for Brain Injury and Repair.

Author contributions: J. Falck performed the majority of the experiments and analyzed data. J. Falck, F. Ackermann, F. K. Hamra, S. Hallermann and C. C. Garner designed experiments. M. Orlando performed electron microscopy studies, I. Straub performed electrophysiology experiments and C. Bruns performed Western Blot studies and a portion of the immunohistochemical stainings. Behavioral experiments were performed by E. J. Plautz; Behavioral experiments were performed by E. J. Plautz. and F. K. Hamra. Behavioral experiments and analysis in the OptiMan setup were performed by H. Munawar, M. Rivalan and

Y. Winter, Z. Izsvák generated *Pclo*^{gt/gt} rat. J. Falck, S. Hoffmann, C.C. Garner and F. Ackermann wrote the manuscript.

Abstract

Piccolo, a presynaptic active zone protein, is best known for its role in the regulated

assembly and function of vertebrate synapses. Genetic studies suggest a further link to several

psychiatric disorders as well as Pontocerebellar Hypoplasia type 3 (PCH3), although a causal

relationship is lacking. We have characterized recently generated knockout ($Pclo^{gt/gt}$) rats.

Analysis revealed a dramatic reduction in brain size compared to wildtype ($Pclo^{wt/wt}$) animals,

attributed to a decrease in the size of the cerebral cortical, cerebellar and pontine regions.

Analysis of the cerebellum and brainstem revealed a reduced granule cell (GC) layer and a

reduction in size of pontine nuclei. Moreover, the maturation of mossy fiber (MF) afferents from

pontine neurons and the expression of the α6 $GABA_A$ receptor subunit at the MF-GC synapse are

perturbed, as well as the innervation of Purkinje cells by cerebellar climbing fibers (CFs).

Ultrastructural and functional studies revealed a reduced size of MF boutons, with fewer synaptic

vesicles and altered synaptic transmission. These data imply that Piccolo is required for the

normal development, maturation and function of neuronal networks formed between the

brainstem and cerebellum. Consistently, behavioral studies demonstrated that adult $Pclo^{gt/gt}$ rats

display impaired motor coordination, despite adequate performance in tasks that reflect muscle

strength and locomotion. Together these data suggest that loss of Piccolo function in patients with

PCH3 could be causal for many of the observed anatomical and behavioral symptoms, and that

the further analysis of these animals could provide fundamental mechanistic insights into this

devastating disorder.

Significance Statement: Pontocerebellar Hypoplasia type 3 is a devastating developmental

disorder associated with severe developmental delay, progressive microcephaly with

brachycephaly, optic atrophy, seizures and hypertonia with hyperreflexia. Recent genetic studies

have identified non-sense mutations in the coding region of the Piccolo gene, suggesting a

functional link between this disorder and the presynaptic active zone. Our analysis of Piccolo

knockout rats supports this hypothesis, formally demonstrating that anatomical and behavioral

phenotypes seen in patients with PCH3 are also exhibited by these Piccolo deficient animals.

Introduction

Pontocerebellar Hypoplasia is a rare and highly heterogeneous group of neurological

disorders, often with a genetic origin, characterized by an abnormally small cerebellum and pons

(Rajab et al., 2003; Namavar et al., 2011). In type 3 Pontocerebellar Hypoplasia (PCH3) - also

known as Cerebellar Atrophy with Progressive Microcephaly (CLAM) - patients suffer from

severe developmental delay, progressive microcephaly with brachycephaly, seizures, hypertonia

with hyperreflexia and short stature (Rajab et al., 2003; Namavar et al., 2011). Additional features

include the presence of craniofacial dysmorphisms and optic atrophy (Durmaz et al., 2009;

Rudnik-Schoneborn et al., 2014).

Previous studies mapped the PCH phenotype to chromosome 7q11-21 (Rajab et al., 2003;

Durmaz et al., 2009). More recently, a single nucleotide polymorphism (SNP) in the human *Pclo*

gene - indeed, located on chromosome 7 at position 21.11 - has been found in patients with

PCH3. This non-sense mutation is predicted to eliminate the C-terminus of the longest Piccolo

isoforms including its PDZ and C2 domains (Ahmed et al., 2015) and perhaps destabilize the

protein, leading to the hypothesis that Piccolo loss of function is responsible for the phenotypes

seen in this neurodevelopmental/neurodegenerative disorder.

Piccolo is a very large (560kDa) multidomain presynaptic scaffold protein and core

component of the cytoskeletal matrix assembled at active zones (CAZ) (Cases-Langhoff et al.,

1996). A range of studies suggest that Piccolo uses its multidomain structure to scaffold not only

other CAZ proteins critical for the regulated release of neurotransmitters, but also proteins

involved in the dynamic assembly of F-actin, synaptic vesicle (SV) recycling and synapse

integrity (Gundelfinger et al., 2015; Ackermann et al., 2019). Intriguingly, Piccolo is present at nearly every synaptic subtype including glutamatergic, GABAergic, cholinergic and dopaminergic synapses within the central (CNS) and peripheral nervous system (PNS) (Cases-Langhoff et al., 1996; Fenster et al., 2000; Fenster and Garner, 2002) and is highly expressed in the cerebrum, hippocampus, cerebellum and olfactory bulb, among others (Cases-Langhoff et al., 1996; Human Protein Atlas, 2015) . It is one of the very first active zone (AZ) proteins recruited to nascent synapses *in vitro* as well as in the developing brain (Zhai et al., 2001). For example, Piccolo appears at emerging synapses formed between mossy fiber boutons and cerebellar granule cells as well as between parallel fiber boutons and Purkinje cell dendrites during the earliest stages of cerebellar development (Zhai et al., 2001). The large size of Piccolo and the complexity of the *Pclo* gene has thwarted most efforts to elucidate its function, though critical roles in retinal ribbon synapse formation and visual function (Regus-Leidig et al., 2014; Muller et al., 2019) as well as the integrity of hippocampal synapses has been identified (Waites et al., 2013). What remains unclear is how Piccolo contributes to cerebellar development and whether, as suggested by genetic studies, it has a primary role in the etiology of PCH3. The recent generation of a Piccolo knockout rat ($Pclo^{gt/gt}$) using transposon mutagenesis (Medrano et al., 2019) provides an opportunity to explore this potential relationship.

In the current study, we have assessed the contribution of Piccolo to cerebellar structure and function through the anatomical, functional and behavioral characterization of adult $Pclo^{gt/gt}$ rats. Our analysis reveals a striking number of similarities to patients with PCH3, including a smaller cerebral cortex, a reduced volume of the cerebellum and pons as well as impaired motor control and the presence of seizures. Our analysis has also uncovered changes in the anatomical and ultrastructural features of mossy fiber terminals and electrophysiological properties of these synapses. Together, these phenotypes are predicted to not only alter the functionality of the cerebellum but to contribute to motor and perhaps also behavioral dysfunctions seen in PCH3 affected individuals.

Results

Brain morphology is changed in Piccolo knockout brains.

The recent analysis of a pair of boys with PCH3 identified a non-sense mutation in the coding

region of the human *Pclo* gene (chr7:82579280 G>A), predicted to eliminate the C-terminal third

of the longest Piccolo isoforms (Ahmed et al., 2015) and likely its expression. These individuals

have profound cognitive and motor impairment as well as atrophy of the cerebrum, cerebellum

and pons (Ahmed et al., 2015). A fundamental question is whether Piccolo loss of function is

causal for this disorder. To explore this possibility, we have characterized a recently generated

line of rats ($Pclo^{gt/gt}$) wherein transposon mutagenesis was used to disrupt the Piccolo gene via an

insertion into exon 3 (Figure 1A). This insertion is predicted to cause a frame shift in the reading

frame and thus disrupt the expression of full-length Piccolo (560kDa) and most of its alternatively

spliced lower molecular weight isoforms (70-350kDa) . Schematic is adapted from Ackermann et

al. (Ackermann et al., 2019; Medrano et al., 2019). Just after birth (P0-P2) Piccolo pups were

found born in normal Mendelian numbers (Figure 1B). Western blot analysis of brain lysates

from postnatal day 0-2 (P0-P2) pups demonstrates the loss of nearly all isoforms in homozygous

knockout animals (Figure 1C). To assess whether Piccolo loss of function adversely affects brain

development, we performed an anatomical characterization of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ animals;

$Pclo^{gt/gt}$ pups were smaller and weighed significantly less than their $Pclo^{wt/wt}$ littermates (Figure

1D, E and F). However, brain weights were not significantly altered between $Pclo^{gt/gt}$ and $Pclo^{wt/wt}$

littermates at P0-2 and display similar brain morphology (Figure 1G, H and I), suggesting that

changes in brain size develop postnatally. However, the overall size and weight of $Pclo^{gt/gt}$ brains

was significantly reduced in 3 month-old adult rats compared to $Pclo^{wt/wt}$ brains (Figure 1J and

K). To assess whether this was associated with an overall loss in brain volume or due to

reductions in specific brain regions, serial sagittal and coronal sections from 3 month-old animals

were collected and stained with Nissl to visualize brain morphology (Figure 1L). Qualitative

analysis revealed that thalamic, cerebellar and brainstem regions are dramatically reduced in size

in 3 month-old $Pclo^{gt/gt}$ animals. Some thinning of the cerebral cortex can also be observed,

however, no obvious changes in the hippocampus are detectable. Furthermore, ventricles (V1,

V2, V3 and V4) in $Pclo^{gt/gt}$ brains were all notably larger than in $Pclo^{wt/wt}$ littermates (Figure 1L).

Intriguingly, the observed morphological changes in brains of 3 month-old $Pclo^{gt/gt}$

animals are remarkably similar to those reported for patients with PCH3, who exhibit

microcephaly, a reduced size of the cerebrum, cerebellum and pons as well as larger ventricles

(Maricich et al., 2011). As hypoplasia of the pons and cerebellum as well as a reduced thickness

of the cortex are the most dramatic features of PCH3 patients, we examined in more detail

changes occurring in these regions of $Pclo^{gt/gt}$ brains. Here, we found that the thickness of the

cortex is significantly reduced in brains lacking Piccolo (Figure 2A and D). The area of the pons

is dramatically reduced in $Pclo^{gt/gt}$ brain slices stained with Nissl (Figure 2B). As the density of

neurons was not changed (Figure 2B, zoom), these data indicate a loss of the total number of

neurons within the pons. This conclusion is supported by data from brainstem sections stained

with antibodies against the synaptic vesicle (SV) protein VGluT1, prominently expressed in pons

neurons, which reveals a smaller area occupied by these neurons (Figure 2C and E).

Analysis of sagittal sections through the cerebellum of adult rats reveals that, whilst the

cerebellum is smaller in $Pclo^{gt/gt}$ animals, the overall anatomy is not altered (Figure 3A). For

example, there are no remarkable defects in foliation of $Pclo^{gt/gt}$ animals, with all lobes present

and appearing to be formed normally at the vermis (Figure 3A and E). Our analysis of the granule

cell layer (GCL), using DAPI to stain granule cell (GC) nuclei, reveals that this layer is

significantly reduced in size in $Pclo^{gt/gt}$ compared to $Pclo^{wt/wt}$ controls (Figure 3A and C).

However, this decrease was not associated with a proportional increase in GC packing density, as

the number of GCs per GCL area is only very slightly increased (Figure 3B and D). These data

indicate an overall loss of GCs in $Pclo^{gt/gt}$ cerebella. Conceptually, this is predicted to reduce the

total number of GC parallel fibers innervating PC dendrites in the molecular layer (ML), a

situation that could lead to a thinner ML and perhaps an altered packing density of Purkinje cells (PCs).

Sections immunostained with antibodies against Calbindin, which specifically labels PCs, reveals that the organization of the ML and the PC layer (PCL) appear to be intact and that the dendritic arbors of PCs are correctly orientated (Figure 3E, F and I). However, quantifying the average area per unit length of the ML reveals a dramatic reduction in the size of this layer in $Pclo^{gt/gt}$ brains (Figure 3G). Furthermore, cells in the PCL appear overcrowded in $Pclo^{gt/gt}$ cerebella (Figure 3E, F and I), a conclusion supported by data showing an increased packing density of PCs in $Pclo^{gt/gt}$ brains (Figure 3H).

Taken together, these data indicate a reduced number of GCs and a higher packing density of PCs in $Pclo^{gt/gt}$ cerebella compared to $Pclo^{wt/wt}$ controls.

Loss of Piccolo alters climbing fiber and mossy fiber innervation in the cerebellum.

Anatomically, the cerebellum receives its major excitatory afferents in the form of climbing fibers (CFs) and mossy fibers (MFs) that project from the inferior olive or the brainstem/spinal cord, respectively (Leto et al., 2016). Both form glutamatergic synapses but terminate in different layers of the cerebellum (Apps and Garwicz, 2005). For example, CFs form excitatory synapses onto the proximal branches of PC dendritic arbors, modulating the dynamic firing properties of PCs and motor learning (Hashimoto and Kano, 1998). In contrast, MFs terminate on the dendrites of GCs, which then provide direct excitatory input to PCs via their parallel fiber (PF) axons. Both also extend collaterals to the deep cerebellar nuclei before projecting into the cerebellar cortex (Shinoda et al., 1992).

In previous studies, we observed that Piccolo was present in the boutons of each of these excitatory synapses (Cases-Langhoff et al., 1996). This was confirmed by immunostaining cerebellar sections of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ with antibodies against Piccolo and the SV protein VGlut1 (Fig. 4A and B). This revealed that Piccolo immuno-reactivity was indeed present at each

of these synaptic types and that this immuno-reactivity was lost in cerebella from $Pclo^{gt/gt}$

animals. To explore whether deficiencies in either could contribute to the anatomical and

functional changes in the cerebellum, we initially analyzed potential differences in the CF input

into the ML. Synaptic input from CFs onto PC dendrites, immuno-positive for Calbindin, was

visualized with antibodies against VGluT2 (Miyazaki et al., 2003). Here our analysis of sagittal

sections revealed the presence of a large number of VGluT2 positive puncta decorating Calbindin

positive dendrites in $Pclo^{gt/gt}$ and $Pclo^{wt/wt}$ sections (Figure 4C and D). These data indicate that the

loss of Piccolo does not affect the ability of CFs to project into the ML and form synapses with

PC soma and dendrites. Qualitatively, VGluT2 positive puncta in $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ sections

were of similar size and beautifully decorated both primary and tertiary PC dendrites, though the

total number of puncta appeared more numerous in $Pclo^{gt/gt}$ sections. Quantification of the total

area of VGluT2 per ML supports this impression (Figure 4D). Additionally, we compared the

distribution of the CFs to the synaptic inputs from GCs onto PCs, the parallel fiber (PF) axons

which project into the ML. The latter synapses were identified with antibodies against VGluT1

(Miyazaki et al., 2003). Although the ML is thinner in $Pclo^{gt/gt}$ cerebella (Figure 3A), we

observed no overt changes in the intensity or distribution of VGluT1 positive puncta throughout

the ML of $Pclo^{gt/gt}$ animals compared to $Pclo^{wt/wt}$ controls (Figure 4C). This implies that PF axons

project normally and form a robust number of synapses with PCs. Given that fewer GCs are

formed in $Pclo^{gt/gt}$ cerebella (Figure 3A and B), we postulate that the thinner ML is most likely

due to fewer PFs and less total synaptic input on PC dendrites.

GCs are known to receive their excitatory input from MFs arising from afferent axons

from a number of distinct nuclei in the brainstem including the pontine nuclei (Sillitoe, 2012).

These collaterals form large glomerular structures with multiple AZs, forming a rosette of

synapses with claws from dendrites of multiple GCs (Jakab and Hamori, 1988; Xu-Friedman and

Regehr, 2003). Given the smaller size of the pons, we thus explored whether the boutons from the

remaining cells properly reached the cerebellum and formed robust MF terminals. As afferent

fibers from the pons primarily innervate cerebellar lobes VI to IX, we examined sagittal sections

of these lobes stained with antibodies against the somatodendritic marker MAP2 and VGluT2 in

lobe VII. In $Pclo^{wt/wt}$ sections, multiple large VGluT2 positive puncta are seen packed tightly

together within a dense meshwork of MAP2 positive dendrites projecting from a ring of GCs.

These puncta represent subclusters of SVs with large 100-200 μm^3 terminals (Jakab and Hamori,

1988). In sections from $Pclo^{gt/gt}$ animals, each bouton appeared smaller in size and less organized,

though they still appear to contact GC dendrites (Figure 5A). This impression is further supported

by sections immunostained with antibodies against VGluT1 and VGluT2. In these images,

distinct VGluT1 and/or VGluT2 positive clusters are observed in both $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$

animals, consistent with individual boutons or glomeruli (Figure 5B). Here again, MF terminals

from $Pclo^{gt/gt}$ animals appeared to have much smaller VGluT1 or VGluT2 positive clusters

(Figure 5B, zoom). Quantifying the average size of the VGluT1 and VGluT2 positive clusters

revealed that clusters of $Pclo^{wt/wt}$ cerebella were more than double the size compared to the same

lobes in $Pclo^{gt/gt}$ animals (Figure 5B and C). However, this effect was not lobe-specific as it was

observed in all lobes and not just in those receiving pontocerebellar afferents (Figure 5B). These

findings suggest that defects in the formation of large robust MF glomeruli in $Pclo^{gt/gt}$ cerebella is

a common feature shared by MF afferents arising from the pons and other brainstem nuclei, and

may reflect aberrant signaling during development between GCs and these neurons.

Quantitatively, the spread of distribution of VGluT1 and VGluT2 cluster sizes is far more

shifted towards smaller cluster sizes in $Pclo^{gt/gt}$, with mutant cerebellar displaying approximately

45 % more small synaptic clusters of 5 μm^2 in size than $Pclo^{wt/wt}$ counterparts (Figure 5D). At the

larger end of the scale, cluster sizes over 50 μm^2 were much more frequent for $Pclo^{wt/wt}$ than

$Pclo^{gt/gt}$ cerebella (Figure 5E).

In addition to excitatory input, MF glomeruli are also modulated by GABAergic

inhibition via cerebellar Golgi cells, which offer regulatory feedback to the complex, as they

themselves are excited by GCs (Maex and De Schutter, 1998). In principle, smaller MFs

terminals seen in Piccolo KO animals could represent less excitatory input into GCs. This may

also reduce excitatory drive onto PCs via PFs from GCs, as well as perhaps inhibitory drive via

the Golgi cells. One of the dominant receptors mediating inhibitory input from Golgi cells to GCs

are the α6 subunit-expressing GABA$_A$ receptors, which are highly concentrated within MF

glomeruli (Nusser et al., 1996). Sagittal sections immunostained with antibodies against VGluT2

and α6 subunits revealed that the GABA$_A$ α6 receptor subunit is only weakly expressed in $Pclo^{gt/gt}$

MF rosettes, whereas in $Pclo^{wt/wt}$, it nicely localizes to the synaptic complex (Figure 6A).

Antibody staining of the granule cell layer demonstrates higher intensity of GABA$_A$ α6 antibody

staining in $Pclo^{wt/wt}$ compared to $Pclo^{gt/gt}$ (Figure 6B). However, knockout models of GABA$_A$α6

(Homanics et al., 1997) do not display PCH3 phenotypes or alterations in the anatomy of the

cerebellum. Therefore, GABA$_A$α6 downregulation can be attributed to Piccolo loss and not for

the PCH3 phenotypes.

These data suggest that the loss of Piccolo not only affects the size of MF inputs into the

cerebellum, but also gene expression and therefore inhibitory drive within each glomerulus, a

condition that could relate to the maturation of these structures and/or their functionality, a

situation that could adversely affect cerebellar function.

Ultrastructural analysis of mossy fiber glomeruli in $Pclo^{gt/gt}$ cerebellum.

The reduced area of VGluT1 and VGluT2 clusters within $Pclo^{gt/gt}$ MF terminals (Figure

6) could be due to a reduction in size of MF glomeruli themselves and/or in the number of

VGluT1/2 positive SVs per bouton. To explore these options, we investigated MF glomeruli in

$Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ cerebella using electron microscopy (EM). Analysis of ultrathin cerebellar

brain sections from 3 month-old rats revealed that the average size of $Pclo^{gt/gt}$ glomeruli was

significantly smaller than $Pclo^{wt/wt}$ glomeruli (Figure 7A and C). Furthermore, the complexity of

the glomeruli indicated by a P2A value, measuring the ratio of perimeter per area, was

significantly reduced (Figure 7E). The size of the presynaptic area was also reduced in $Pclo^{gt/gt}$

cerebellar sections (Figure 7B and D). However, the number of active zones (AZs) present at

each glomerulus was still proportional to their size, as $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ had a similar number

of AZs per glomerular area (Figure 7B and F). Given that $Pclo^{gt/gt}$ boutons are smaller, this

indicates that the overall output of the MF glomeruli could be reduced in $Pclo^{gt/gt}$ cerebellum.

Intriguingly, we also noticed an accumulation of clathrin-coated vesicles (CCVs) in the terminals

of $Pclo^{gt/gt}$ MFs (Figure 7G). This phenotype resembles recent findings from hippocampal

synapses from $Pclo^{gt/gt}$ animals, which revealed defects in the formation of endosomal membranes

and an overall reduction in SV number (Ackermann et al., 2019) and suggest possible changes in

the recycling of SVs within MF boutons post fusion with the plasma membrane.

Piccolo loss alters GC properties and mossy fiber to GC synaptic transmission

The anatomical and morphological changes observed in MF boutons lacking Piccolo are

predicted to not only represent altered afferent input into the cerebellum from the pons and other

brainstem nuclei, but also altered cerebellar function. As an initial test of this hypothesis, we

performed whole-cell current clamp recordings of cerebellar GCs from acute P90 rat cerebellar

slices. A two-photon image of a typical cerebellar GC from $Pclo^{gt/gt}$ filled with ATTO dye reveals

a normal radial arrangement of its dendrites as they project their claws into MF glomeruli (Figure

8A). An analysis of the intrinsic biophysical properties of these cells revealed that the GC

properties differed between $Pclo^{gt/gt}$ and $Pclo^{wt/wt}$ animals. Specifically, no changes were detected

in either the capacitance or the membrane potential of these cells, but the input resistance was

significantly increased (Figure 8B). Since these experiments were performed in the presence of

$GABA_A$ receptor blockers, the decreased shunting inhibition mediated by tonic activation of α6-

subunit-containing $GABA_A$ receptors (Brickley et al., 1996; Nusser et al., 1998) is expected to

further increase the difference in input resistance. Yet, there was also no change in the amplitude,

threshold of activation or duration of action potentials fired by these cells (Figure 8B), indicating

unaltered active membrane properties. Examining the frequency and amplitude of spontaneous

miniature excitatory postsynaptic currents (mEPSCs) of these GCs revealed a dramatic increase in the frequency of these events in $Pclo^{gt/gt}$ slices with no change in mEPSC amplitudes (Figure 8C). These data suggest that on average each excitatory synapse formed on to these GC dendrites has normal levels of postsynaptic AMPA-type glutamate receptors. The change in frequency could either be due to an increase in the number of MF boutons contacting GC dendrites and/or an increase in the release probability of MF boutons. Consistently, the average amplitude of the evoked excitatory postsynaptic currents (EPSCs) was increased in $Pclo^{gt/gt}$ animals, with no change in the weighted time constant (τ_w) (Figure 8D). Given the amplitudes of the mEPSCs are not changed, these results suggest that there is a higher number of synaptic connections between MF boutons and the dendrites of GCs, consistent with the larger number of smaller SV clusters/rosette in $Pclo^{gt/gt}$ animals (Figure 6). Taken together, these data indicate that, in addition to changes in cell number, the loss of Piccolo in the cerebellum and brainstem is associated with changes in the morphology and function of GCs and their mossy fiber input.

Behavioral and motor defects in Piccolo knockout rats.

The motor difficulties reported (Zelnik et al., 1996; Durmaz et al., 2009) in humans with PCH3 and the anatomical changes observed in the cerebellum and brainstem of rats lacking Piccolo predict altered motor function in these animals. To test this hypothesis, $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats were monitored for their motor abilities. In rotarod tasks, $Pclo^{gt/gt}$ rat performance was significantly reduced compared to $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ rats (Figure 9A). Specifically, while $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ rats exhibited increasing performance levels regarding the ability to stay on the rotarod over time, $Pclo^{gt/gt}$ rats showed no indication of being able to adapt to the task. $Pclo^{gt/gt}$ rats were less adept at staying on the task apparatus once rod rotation was initiated (Figure 9A). Intriguingly, no differences in forelimb grip strength were scored between Piccolo genotypes (Figure 9B), indicating that $Pclo^{gt/gt}$ rats showed lack of motivation and/or impaired coordination.

In addition to deficits in motor ability, $Pclo^{gt/gt}$ rats displayed an increased frequency in

front and rear foot stepping errors during ladder rung walking compared to $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$

rats (Figure 9C). In open field tests, no significant differences were recorded between $Pclo^{wt/wt}$,

$Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats for peripheral line crossing or self-grooming events (Figure 9D).

However, $Pclo^{gt/gt}$ rats displayed 2- to 3-fold decreases in peripheral rearing events compared to

$Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ animals (Figure 9D). Deficits in $Pclo^{gt/gt}$ rat performance during rotarod,

ladder rung and open field tests (Figure 9A-D) reflected recessive traits, consistent with

dysfunction in proprioceptive sensation and motor control (Curzon et al., 2009).

Alongside traditional tests for motor coordination, we tested the behavior of $Pclo^{wt/wt}$,

$Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats in a home cage setup, the OptiMan (Operator Independent Motor-

analysis) system, where animals can be monitored without interference from experimenters. Rats

were tagged with a radio frequency identification chip that allowed for tracking of locomotor

activity while they were also required to complete an isometric pull task that requires precise and

finely controlled movements. In the home cage, the $Pclo^{gt/gt}$ rats were more active and covered

about twice the distance than $Pclo^{wt/wt}$ rats (Figure 9E) during each measurement. In the isometric

pull task, performance of $Pclo^{gt/gt}$ rats was significantly lower than the performance of $Pclo^{wt/wt}$

rats, quantified using four different force thresholds (Figure 9F). Taken together, $Pclo^{gt/gt}$ rats

show clear motor deficits, very similar to symptoms seen in PCH3 patients (Rajab et al., 2003).

Discussion

Our study demonstrates that Piccolo LOF causes alterations in brain anatomy. In

particular, the cerebrum, pons, brainstem and cerebellum are severely reduced in size whereas the

ventricles are increased (Figure 1). These changes are associated with reductions in cerebellar and

pontine cell numbers and perturbations in cerebellar CF and MF afferents (Figure 4 and 5). These

changes are predicted to adversely affect cerebellar function - supported by changes in synaptic

transmission and motor control in $Pclo^{gt/gt}$ rats. Interestingly, the changes in brain morphology

resemble changes in children with PCH3, recently linked to a SNP in the PCLO gene (Ahmed et

al., 2015). In addition to brain atrophy, children with PCH3 have, amongst other symptoms,

cognitive and motor deficits as well as seizures. Given that seizures were also observed in $Pclo^{gt/gt}$

rats (Medrano et al., 2019), we postulate that $Pclo^{gt/gt}$ rats represent a model to study the

underlying mechanisms of this devastating disease.

Similar to patients with PCH3, $Pclo^{gt/gt}$ rats have smaller cerebella, brainstem and pontine

nuclei. In the cerebellum, a striking change was the thickness of the ML and CGL (Figure 3). The

latter was associated with fewer total GCs, which could result in fewer PFs and a thinner ML.

The orientation of PCs and the ramification of their dendritic arbors were largely unchanged. At

present, it is unclear why there are fewer GCs in $Pclo^{gt/gt}$ cerebella. During normal development,

several factors including Sonic hedgehog (Wallace, 1999; Miyashita et al., 2017) and Notch

(Solecki et al., 2001) are known to control the proliferation of GCs. How Piccolo loss influences

these and related signaling pathways is unclear; though Piccolo and Bassoon have been shown to

regulate the activity-dependent translocation of c-terminal binding protein 1 (CTPB1) to the

nucleus and thereby the expression of neuronal genes (Ivana et al., 2015). CTPB1 and CTPB2 are

highly expressed in the cerebellum and their function and/or localization could be affected by

Piccolo LOF (Hubler et al., 2012; Ivanova et al., 2015). There is a clear rationale for exploring

this phenotype further.

In the ML, CF morphology and arrangement appear to be normal, though a higher

innervation of PC dendrites by CFs can be observed (Figure 4). One possible explanation for this

hyper-innervation is that homeostatic changes to the network, such as heterosynaptic competition

with PFs for PC dendrite territory (Hashimoto et al., 2009; Ichikawa et al., 2016) could contribute

to these differences in CF distribution and therefore the functionality of the cerebellum. This

could affect normal pruning mechanisms of CF during development.

Our study also revealed that the pontine nuclei was dramatically decreased in size in

$Pclo^{gt/gt}$ animals. This is note-worthy as afferent MFs from the pons and other brainstem nuclei

are the primary excitatory input onto GCs, forming elaborate rosette synapses (Voogd and

Glickstein, 1998). The smaller size of the pons suggests a net reduction in MF input into the

cerebellum that appears to correspond to the reduced number of GCs (Figure 3). However, it is

unclear how the loss of Piccolo could influence the number of neurons in the pons.

Surprisingly, our histological studies revealed that MF boutons across all lobes of the

cerebellum were reduced in size (Figure 5). This finding was supported by our EM studies

revealing that MF glomeruli are severely reduced in size, potentially resulting in smaller SV

clusters (Figure 7). These observations suggest that the development/maturation of MFs from the

pons and other brainstem nuclei are muted in $Pclo^{gt/gt}$ animals. Additionally, electrophysiological

changes (Figure 8) indicate that the network has triggered compensatory changes to overcome a

smaller size or impaired strength of MFs reaching the GC layer, for instance by increasing both

input resistance and release probability (Turrigiano, 2012).

Our electron micrographs also showed what could be disturbed synaptic integrity in

$Pclo^{gt/gt}$ MF boutons with more CCVs being present (Figure 7), suggesting that the loss of Piccolo

may also affect SV recycling. These observations are in accordance with recent findings by

Ackermann et al. (2019) of hippocampal synapses from $Pclo^{gt/gt}$ animals. Here, it was reported

that the loss of Piccolo had a dramatic effect on the recycling of SV proteins through a functional

block in the formation of early endosomes from endocytic vesicles, due to defects in the

activation of Rab5 via a Piccolo-dependent loss of Pra1 from synapses. Although beyond the

current study, it is likely that this endocytic defect in the recycling of SV proteins could also

contribute to altered size function of MF synapses, especially given their high frequency

transmission and therefore need for high SV turnover (Byczkowicz et al., 2018).

GCs also receive inhibitory modulation from Golgi cells (Eccles et al., 1966) and it is

well appreciated that GABA$_A$ receptors at the MF synapse contain α6 subunits (Nusser et al.,

1996). Intriguingly, $Pclo^{gt/gt}$ rats display a reduced expression of the GABA$_A$α6 subunit at MF

rosettes (Figure 6). This is in line with findings from Medrano et al. (2019), who performed RNA

sequencing on the $Pclo^{gt/gt}$ rats and found a severe reduction in GABA$_A$α6 subunit gene

expression.

At present, it is unclear why levels of these subunits are reduced. One possibility is that it

reflects a homeostatic change within cerebellar circuitry to compensate for a reduced excitatory

input from MFs (Figure 5). This concept is supported by electrophysiological data showing that

the input resistance of GCs is higher as is mEPSC frequency and EPSC amplitudes (Figure 8).

This condition might arise to compensate for the smaller MF terminals reaching the GCL, which

could initially have weaker output properties. Alternatively, reduced GABA$_A$α6 expression could

be due to lower levels of BDNF, which is secreted from precerebellar neurons, at MF terminals

(Chen et al., 2016) and is necessary to promote the formation of GABAergic synapses onto GCs.

Though a role of Piccolo in the secretion of BDNF has not been investigated, the expression of

Bassoon, which shares significant functional redundancy with Piccolo, is linked to presynaptic

levels of BDNF (Heyden et al., 2011). In this regard, the remaining Bassoon protein could

suppress BDNF secretion in MF terminals, altering the maturation of the glomeruli and

GABA$_A$α6 expression.

An important question not addressed by our studies is why MF boutons are smaller. By

several measures mentioned above, it would appear that the MF glomeruli are less mature. Key

regulators of MF maturation are members of the Wnt family, a group of target-derived factors

which accelerate neuronal maturation or directly induce synapse formation (Scheiffele, 2003;

Waites et al., 2005). Interestingly, mouse cerebellar MFs lacking Wnt7a show a similar reduction

in MF size and complexity, as we observed in $Pclo^{gt/gt}$ (Hall et al., 2000). However, it is important

to note that in the case of Wnt7a loss, MF synapse size catches up to WT size by age P15, which

points to a lag in maturation and does not quite seem to be the case in our $Pclo^{gt/gt}$ rats. Knockout

of Disheveled1 (Dvl1), a downstream target of Wnt (Salinas and Zou, 2008) also shows a

reduction in MF cluster size, indicating that presynaptic Dvl1 is a necessary step in the Wnt

signaling cascade, underscoring the importance of these proteins for MF-GC synapse formation

(Ahmad-Annuar et al., 2006). Like Dvl1, Piccolo is located at the AZ of the presynaptic terminal

and regulates F-actin assembly and synaptic transmission through its interaction with Daam1 and

Profilin (Wagh et al., 2015). Daam1 is a formin and a known regulator/interaction partner of Dvl1

(Gao and Chen, 2010). It is thus possible that, in the absence of Piccolo, Dvl1 is not properly

localized to presynaptic sites, preventing proper Wnt signaling and consequently causing the

formation of smaller and less organized MF-GC synapses. Furthermore, analysis of $Pclo^{gt/gt}$

transcripts reveals that Wnt expression is reduced (Medrano et al., 2019). Thus, there appears to

be at least two possible mechanisms that could contribute to smaller MF boutons in Piccolo

knockout rats: defects in SV recycling and Wnt signaling. Clearly further studies are needed to

explore these options.

A fundamental question raised by the anatomical, morphological and functional changes

within the cerebellum in rats lacking Piccolo is whether these changes affect the functionality of

the cerebellum. In behavioral tests, we observed that $Pclo^{gt/gt}$ rats performed significantly worse

than both $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ littermates at motor function tasks, highlighting the recessive

nature of the behavioral impairments. Our colleagues Medrano et al. (2019) also observed

epileptic seizures and increased aggression in $Pclo^{gt/gt}$ rats, and a failure to reproduce due to

impaired brain-gonad signaling.

Taken together, these data support the concept that Piccolo loss of function in patients

with PCH3 could be causal for many of the observed phenotypes, including changes in the

volume of brain structures and behavioral abnormalities such as impaired motor control and

epileptic seizures (Rajab et al., 2003; Namavar et al., 2011). With regard to reduced cerebellar

function, our studies highlight a prominent role for MF boutons which are not only smaller in size

but with altered synaptic properties. Mechanistic studies, which probe how Piccolo loss

contributes to these changes, should provide insights into the etiology of this devastating disease.

Materials and Methods

<u>Generation of Piccolo KO rats ($Pclo^{gt/gt}$)</u>

Generation of mutant Piccolo rat strains: Mutant rat strains harbored Sleeping Beauty β-Geo trap

transposons (Ivics et al., 2009), originally transmitted from a donor, recombinant rat

spermatogonial stem cell library (Izsvak et al., 2010). Recipient males were bred with wildtype

females to produce a random panel of mutant rat strains enriched with gene traps in protein

coding genes (Izsvak et al., 2010). All experiments were approved by the Institutional Animal

Care and Use Committee (IACUC) at UT-Southwestern Medical Center in Dallas, as certified by

the Association for Assessment and Accreditation of Laboratory Animal Care International

(AALAC) NIH OLAW Assurance # D16-00296.

<u>Characterization of pups and genotyping</u>

All procedures for experiments involving animals, were approved by the animal welfare

committee of Charité Medical University and the Berlin state government. P0-P2 rats were

weighed using Kern 440-43N scales and measured for approximate length with a ruler.

Genotyping of pups' cortices later revealed their genetic identity.

P0-P2 pups were genotyped using a PCR based reaction. In brief, brain tissue was

digested in lysis buffer (100 mM Tris-HCl (pH 8.0) with 10 mg/ml proteinase K, 100 mM NaCl)

for 5 minutes at 55 °C, before inhibiting Proteinase K by incubation at 99 °C for 10 minutes.

Samples were then centrifuged at 14,800 rpm for 2 minutes and 1 µl of supernatant was used for

the PCR reaction as outlined below.

For determination of genotype for adult rats, earpieces were taken and digested overnight

at 55 °C in SNET-buffer (400 mM NaCl, 1 % SDS, 200 mM Tris (pH 8.0), 5 mM EDTA)

containing 10 mg/ml proteinase K. Proteinase K enzyme reaction was stopped incubating the

samples for 10 min at 99 °C. The mixture was centrifuged for 2 min at 14,800 rpm. Supernatant

was transferred into a fresh tube and DNA was precipitated by addition of 100 % isopropanol.

Following samples were centrifuged for 15 min at 4 °C, 13,000 rpm. Precipitated DNA was

washed once with 70 % ethanol and centrifuged again for 5 min at 13,000 rpm. Supernatant was

discarded and the DNA pellet was air dried and resuspended in H2O. A PCR reaction with a

specific primer combination was performed on isolated DNA. The following primers were used:

Pclo KO F2: 3` gcaggaacacaaaccaacaa 5`; Pclo KO R1: 3` tgacctttagccggaactgt 5`; SBF2: 3`

tcatcaaggaaaccctggac 5`. The PCR reaction protocol was the following: 2 min 94 °C; 3 x (30 sec

94 °C, 60 °C 30 sec, 72 °C 30 sec); 35 x (94 °C 30 sec, 55 °C 30 sec, 72 °C 30 sec); 72 °C 10

min. Samples were mixed with a loading dye (New England BioLabs, MA, USA) and run on 2 %

agarose gel (Serva, Heidelberg, Germany) at 110 V for 45 min. The gel was imaged using

BioDocAnalyze UV transilluminator and BioDocAnalyze2.2 software.

<u>Western blot analysis</u>

Brains from P0 – P2 pups were lysed in Lysis buffer (50 mM Tris-HCl, 150 mM NaCl, 5 mM

EDTA, 1 % Triton X-100, 0.5 % Deoxycholate, protease inhibitor pH 7.5) and incubated on ice

for 5 min. Samples were centrifuged at 13,000 rpm for 10 min at 4 °C. Afterwards the

supernatant was transferred into a fresh tube and the protein concentration was determined using

a BCA protein assay kit (Thermo Fisher scientific, Waltham, Massachusetts, USA). The same

protein amounts for $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $\underline{Pclo^{gt/gt}}$ samples were separated by SDS-PAGE and

transferred onto nitrocellulose membranes (running buffer: 25 mM Tris, 190 mM Glycine, 0.1 %

SDS, pH 8.3; transfer buffer: 25 mM Tris, 192 mM Glycine, 1 % SDS, 10 % Methanol for small

proteins, 7 % Methanol for larger proteins pH 8.3). After the transfer, nitrocellulose membranes

were blocked with 5 % milk in TBST (20 mM Tris pH 7.5, 150 mM NaCl, 0.1 % Tween 20) and

incubated with primary antibodies in 3 % milk in TBST overnight at 4 °C. The following

antibodies were used: Piccolo (1:1000; rabbit; Synaptic Systems, Göttingen, Germany; Cat#

142002), GABA$_A$α6 receptor (1:300; Merck Darmstadt, Germany; Cat# 06-868) overnight.

Nitrocellulose membranes were washed 3 times for 10 min with TBST and incubated with HRP-

conjugated secondary antibodies for 1 h at RT (1:1000; Sigma-Aldrich, St. Louis, USA; Cat#

NA9310 Cat# NA934). Membranes were washed 3 times for 10 min with TBST, afterwards

secondary antibody binding was detected with ECL Western Blotting Detection Reagents

(Thermo Fisher Scientific, Waltham, USA) and a Fusion FX7 image and analytics system (Vilber

Lourmat, Collégien, France).

<u>Immunohistochemistry</u>

Immunohistochemistry was performed on brain tissue from rats perfused with 4 %

paraformaldehyde (Roth, Karlsruhe, Germany) dissolved in PB (80 mM Na2HPO4 (Roth,

Karlsruhe, Germany), 20 mM NaH2PO4 (Bernd Kraft, Duisburg, Germany)) (PFA). Brains were

extracted and placed in 4 % PFA overnight, cryoprotected in 15 % and then 30 % sucrose

solution (Sigma-Aldrich, St. Louis, USA; dissolved in PB) for 24 hours each. Brains were snap-

frozen by submersion in 2-methylbutane (Roth, Karlsruhe, Germany) cooled to -60 °C and then

stored at -20 °C until use. Brains were cut para-sagittally or coronally with a Leica cryostat to

either 20 μm thick sections and mounted on superfrost slides (Thermo Fisher Scientific,

Waltham, USA), or 50 μm sections which were processed free-floating. Slides were left to dry for

a minimum of one hour before storage at -20 °C and free-float sections were stored in antifreeze

solution (30 % Ethylene glycol, 30 % Glycerol (Roth, Karlsruhe, Germany), 30 % ddH2O, 10 %

0.244M PO4 buffer (NaOH, NaH2PO4, Roth, Karlsruhe, Germany)).

Prior to staining at least 4 slides (each containing two sections) from each animal were

left to equilibrate at room temperature (RT) for one hour. Sections were selected to encompass

the range of the axis we were investigating. A hydrophobic barrier was created around sections

using a DAKO pen (DAKO, Glostrup, Denmark) and sections were washed and permeabilized

with TBS (20 mM Tris pH 7.5, 150 mM NaCl, Roth, Karlsruhe, Germany) with 0.025 % Triton

X-100 (Roth, Karlsruhe, Germany) (TBST) for 3 x 5 min, prior to blocking with 10 % Normal

Goat Serum (NGS, Sigma-Aldrich, St. Louis, USA) with 1 % Bovine Serum Albumin (BSA,

Sigma-Aldrich, St. Louis, USA) in TBS. The following primary antibodies were used: Calbindin

(1:750; rabbit; Synaptic Systems, Göttingen, Germany; Cat# 214002), GABA$_A$ receptor α subunit

(1:250; rabbit; Sigma-Aldrich, St. Louis, USA; Cat# G5544), MAP2 (1:1000; chicken;

MilliporeSigma, Burlington, USA, Cat# AB5543), Piccolo (1:200; guinea pig; Synaptic Systems,

Göttingen, Germany; Cat# 142 104), VGluT1 (1:1000; rabbit; Abcam, Cambridge, UK; Cat#

ab104898), VGluT2 (1:250; guinea pig; Synaptic Systems, Göttingen, Germany; Cat#

13540419). VGluT2 (1:300; mouse monoclonal (epitope AA 566 to 582); Synaptic Systems,

Göttingen, Germany; Cat# 135 421). Antibodies diluted in TBS with 1 % BSA were applied and

left overnight at 4 °C. After 3 x 5 min washing with TBST, differently labeled secondary

antibodies were used from Invitrogen (Thermo Fisher Scientific, Waltham, USA, dilution

1:1000), again diluted in 1 % BSA in TBS antibody solution and then applied for 1 hour at RT.

Sections were then washed with TBS 2 x 10 min or, if desired, incubated in TBS with DAPI

(Roth, Karlsruhe, Germany) for 30 min before washing again 2 x 10 min. Slides were

coverslipped (24x50 mm coverslips, Menzel Gläser, Braunschweig, Germany) with Immu-Mount

(Shandon-Thermo Scientific, Cheshire, UK) and sealed with clear nail polish once hardened.

<u>Nissl stain</u>

Sections were washed 3 times in PBS then mounted onto superfrost slides and allowed to dry for

1-2 days. Slides were inserted into slide racks and passed through the following solutions: 95 %

EtOH (Ethanol from Roth, Karlsruhe, Germany; diluted as appropriate with ddH2O) x 15 min, 70

% EtOH, 50 % EtOH, ddH2O, 10 min Blue counterstain (TACS from Trevigen, MD, USA),

ddH2O and then dehydrated through 50 %, 70 % acid EtOH (1 % glacial acetic acid (Th. Geyer,

Renningen, Germany) in 70 % EtOH), 95 % and 100 % EtOH before clearing in Roti Histol

(Roth, Karlsruhe, Germany), coverslipped using Entellan mounting medium (Entellan,

Darmstadt, Germany) and left to dry for 24 hours under a fume hood.

<u>Confocal z-stack image acquisition and processing</u>

Images were acquired on a spinning disc confocal microscope (Zeiss Axio Observer.Z1 with

Andor spinning disc and cobolt, omricron, i-beam laser) using a 40x (1.3 NA) and 100x (1.4 NA)

Plan-Apochromat oil objective and an iXon ultra (Andor, Belfast, UK) camera controlled by iQ

software (Andor, Belfast, UK). Sections for $GABA_A\alpha6$ analysis (Figure 7) were imaged with a

Nikon Spinning Disk Confocal CSU-X using a 100x (1.45 NA) Plan Apo oil objective and an

EMCCD camera with Andor Revolution SD System(Andor, Belfast, UK).

<u>Tile scan overview images</u>

An upright microscope (Olympus BX61) was used to image fluorescently stained cerebellar

sections. A CCD color camera was used with a 10x or 5x lens for overview pictures. CellSens

software (Olympus, Hamburg, Germany) stitched multiple images together to give an overview

of the whole cerebellar region.

<u>Image analysis</u>

For image processing and analysis ImageJ/FIJI software was used (Schindelin et al., 2012). For

analysis of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ tissue sections were selected from the vermis and approximately

every 10th slide laterally (each slide containing 2 sections) was analyzed. The best quality section

per slide was imaged. For layer thickness, all lobes were measured and for closer analysis of MF

size, CF coverage and GC density, images were taken from lobes I, III, V, VII and IX where

possible. For $GABA_a\alpha6$ subunit expression, 6 images were taken per slide, 4 slides per animal.

Average signal intensity was measured for the whole field of view in the $GABA_a\alpha6$ antibody

channel and then normalized to the MAP2 channel for each image.

542   Layer 'thickness' was calculated per lobe by dividing the area of the layer by the inner

length of the layer for both GCL and ML. For Purkinje cell density, the number of PCs per lobe

was counted and divided by the length of the PC layer for that lobe.   Data points represent

'thickness' values from individual lobes.

Granule cell density was calculated from 100x magnification single plane images stained

with DAPI. GCs were counted per image by an experimenter blinded to the genotype of each

image, and this number divided by the area of each image. Data points represent each image. .

Mossy fiber cluster size was measured using a script within ImageJ to analyze particles

stained with VGluT1 and/or VGluT2. The area of each puncta was measured and an average

taken per image. Data points represent average cluster size of each image.

Climbing fiber innervation was assessed using VGluT2 and Calbindin staining. The

molecular layer area was defined in the Calbindin channel, consisting of Calbindin-positive PC

dendrites, and the VGluT2 channel was selected and pasted into a new imageJ file. The same

script as for MF analysis was run to calculate the total area stained with VGluT2, divided by the

area of the ML and x100 to give percentage coverage. Data presented represent percentage per

image.

Electron microscopy

Pclo$^{wt/wt}$ and *Pclo*$^{gt/gt}$ rats were anesthetized deeply with 33 mg/ml Ketamine (Inresa Arzneimittel

GmbH, Freiburg, Germany) 830 μg/ml Xylavet (cp-pharma, Burgdorf, Germany) and perfused

transcardially with 37 °C physiological saline for 3 to 4 min followed by a 0.1 M phosphate

buffer containing 4 % paraformaldehyde and 0.05 % glutaraldehyde. Brains were stored in

fixative overnight and subsequently sliced sagittally (350 μm) on a vibratome. Regions of interest

were cut into small pieces and post-fixed in 1 % OsO4 and 0.1 M sodium cacodylate and en-bloc

stained in 1 % uranyl acetate aqueous solution. Finally, samples were dehydrated and embedded

in epoxy resin (Epon 812; EMS). Ultrathin sections were cut using an Ultracut ultramicrotome

(UCT, Leica, Wetzlar, Germany) equipped with a diamond knife (Ultra 45, DiATOME, Hatfield,

USA) and collected on formvar-coated 200-mesh copper grids (EMS). Sections were imaged in a

Zeiss EM-900 Transmission Electron Microscope (Zeiss) operated at 80 keV and equipped with a

2K x 2K CCD camera. Data were analyzed blindly using the ImageJ software. Data points

represent analysis from individual images obtained from 3 rats per genotype. For histological and

EM data, a normality test was performed (D'Agostino-Pearson omnibus normality test). If

successful then a student's t-test was used to compare $Pclo^{wt/wt}$ to $Pclo^{gt/gt}$ rats, or alternatively a

test for non-normal data (Mann-Whitney U test) was used.

Electrophysiology methods

Cerebellar slices were prepared from 3-to-4 month-old $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ rats of either sex.

Animals were treated in accordance with the German Protection of Animals Act and with the

guidelines for the welfare of experimental animals issued by the European Communities Council;

local authorities approved the experiments. Animals were anesthetized with isoflurane (Baxter,

Deerfield, IL) followed by a rapid decapitation with a custom-built guillotine. The cerebellar

vermis was quickly removed and mounted in a chamber filled with chilled extracellular solution.

Parasagittal 300 μm slices were cut using a Leica VT1200 microtome (Leica Microsystems,

Wetzlar, Germany), transferred to an incubation chamber at ~35 °C for 30 min and subsequently

stored at RT. Artificial cerebrospinal fluid (ACSF) was used for slice cutting, storage, and

experiments. ACSF contained: 125 mM NaCl, 25 mM NaHCO3, 2.5 mM KCl, 1.25 mM

NaH2PO4, 1.1 mM CaCl2, 1 mM MgCl2, 3 mM Glucose, 17 mM Sucrose (~310 mOsm, pH 7.3

when bubbled with Carbogen (5% O2/95% CO2)). Patch pipettes were pulled from borosilicate

glass (Science Products, Hofheim, Germany) using a DMZ Puller (Zeitz-Instruments,

Martinsried, Germany). Patch pipettes had open-tip resistances of 6–9 MΩ. The intracellular

solution contained: 150 mM K-gluconate, 10 mM NaCl, 10 mM K-HEPES, 3 mM Mg-ATP, 0.3

mM Na-GTP (300–305 mOsm, pH adjusted to 7.3 with KOH). In some of the experiments, the

intracellular solution contained 10 μM of the fluorescence dye Atto594. Experiments were

performed at 35–37 °C and slices were continuously superfused with ACSF containing 20 μM

SR95531 and 40 μM D-(2R)-amino-5-phosphonovaleric acid (D-APV) to block Golgi-cell

inhibition and NMDA-receptors, respectively. Atto594 was obtained from Atto-Tec (Atto-Tec,

Siegen, Germany); all other chemicals were purchased from Sigma-Aldrich (St. Louis, MO).

<u>Current clamp recordings</u>.

To determine the input resistance, subthreshold current pulses were applied from -20 to +20 pA

in 2pA steps. Action potentials were evoked in current-clamp mode by current pulses (amplitude

20-400 pA, duration 300 ms). The resistance of the solution-filled patch-pipettes was 24.9 ± 1

MΩ and 23 ± 1 MΩ for $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ rats respectively. Patch-clamp recordings were made

using a HEKA EPC10/2 USB amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany). Data

were sampled at 200 kHz. Measurements were corrected for a liquid junction potential of +16

mV.

Excitatory postsynaptic currents (EPSCs)

To measure evoked EPSCs, GCs were held at a holding potential of – 80 mV and Mossy fiber

axons were stimulated at 1 Hz. The stimulation voltage ranged between 16 to 40 V for control

and $Pclo^{gt/gt}$ animals. For spontaneous EPSCs GCs were held at -80 mV for around 3 minutes.

Single events were detected using the Igor Pro extension NeuroMatic (Rothman and Silver,

2018b) tool for Event detection.

The current clamp data were analyzed using custom-made procedures in Igor Pro

software (WaveMetrics, Oregon, USA) as described previously (PMID:31379501). In short,

properties of action potentials of GCs were determined from the injected currents, that elicited the

largest number of action potentials (APs). The action potential threshold was defined as the

membrane voltage at which the first derivative exceeded 100 V s^{-1}, the minimal AP peak was set

as -20 mV and the minimal amplitude to 20 mV. All APs with a half-width smaller than 50 μs

and higher than 500 μs were excluded. AP frequency and AP half-width were calculated from the

first three APs. Membrane capacitance, resting membrane potential and series resistance were

read from the amplifier software (HEKA) after achieving the whole-cell configuration. Input

resistance (R_{in}) was analyzed by plotting the steady-state voltage elicited by the subthreshold

current injections against the injected current and a spline interpolation was performed to obtain

the slope at the holding membrane potential (0 pA current injection).

Excitatory postsynaptic currents

To measure evoked EPSCs, GCs were held at a holding potential of – 80 mV and mossy fiber

axons were stimulated extracellular with a second patch pipette at 1 Hz. For spontaneous EPSCs

GCs, were held at -80 mV for around 3 minutes. Single events were detected using the template

detection algorithm of the Igor Pro extension NeuroMatic (Rothman and Silver, 2018a).

EPSCs were analyzed with the Igor Pro software. The amplitude and the kinetics were

determined from the average of 25 individual single EPSCs. To obtain the decay kinetics, single

EPSCs were fitted with one or two exponentials. The weighted time constant was calculated as:

$$\tau w = \frac{Aslow\ \tau slow\ +\ Afast\ \tau fast}{Aslow\ +\ Afast}$$

Behavioral assessment

Rotarod task

The apparatus (IITC Life Science, Woodland Hills, CA) consisted of 5 semi-enclosed lanes and

an elevated metal rod (9.525 cm diameter, 29.21 cm elevation) with a fine textured finish to

enhance grip. For each trial, all rats were placed on the unmoving rod, allowed to stabilize their

posture, and then rod rotation was initiated. Test parameters were: rotation direction, toward

investigator to encourage rats to face away while walking; start speed, 4 rpm; top speed, 44 rpm;

acceleration rate, 0.2 rpm/s (200 sec from start to top speed); max test duration, 300 s. Each rat's

trial ended when it fell from the rod, triggering the fall-detection sensor. Data was automatically

recorded to a computer. Rats underwent 4 trials/day, with an inter-trial interval of at least 10 min

for 4 consecutive days.

<u>Ladder rung task</u>

Ladder rung tests were performed on cohorts of $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats by methods

previously described (Metz and Whishaw, 2009). Rats were trained to cross a plexiglass tunnel

about 1 m long with metal rods provided at regular intervals as steps. Each step that the rat took

was scored on the basis of paw placement on a 7 category scale with 0 being paw totally missing

the rung and 6 being correct paw placement. The average score per pair of limbs was used to

quantify step score. During the trial, the error per step was also quantified by number of low

scoring steps (0-2) divided by the total number of steps that the rat took during the test with each

pair of limbs (student's t-test). Data were derived from 3 trials with 3 cm rung separation

conducted on the same test day with at least a 10 min inter-trial interval, and group means were

compared using one-way ANOVA and Bonferroni post-hoc tests.

<u>Grip test</u>

To assess grip strength, rats were allowed to cling on to a support by forelimbs or hindlimbs and

pulled (Curzon et al., 2009). The support is attached to a transducer that can measure the pull

force being applied on the rat by the tester. During each trial, force that was necessary to be

applied by the tester to release the grip of the rat was recorded. Three trials per limb pair were

done and the means compared by student's t-test. To assess grip strength, the rat's torso was

supported ventrally while both forelimbs were allowed to grasp a metal support bar, then the rat

was pulled in a horizontal plane until the bar was released. Peak force was measured by a

transducer attached to the support bar (San Diego Instruments, San Diego, USA). Nine trials per

rat were conducted over two days (inter-trial interval at least 5 min) and group means were

compared using one-way ANOVA and Bonferroni post-hoc tests.

<u>Open field test</u>

The open field test (Curzon et al., 2009) was performed for each rat using a rectangular arena

(91.44 cm x 60.96 cm) divided into 9 equal sectors (1 center, 8 perimeter). At the start of the

experiment, the rat was placed in the center sector and allowed to explore for 10 min. The test

was recorded by digital video for later analysis. One test was conducted per rat. The behaviors

counted were crossing, rearing and grooming. Each crossing event was counted when all four

limbs of the test subject crossed the boundary of one sector into another. Rearing was defined as

the number of times the subject stood up on its hind limbs. Grooming was defined as the number

of times a rat licked or scratched itself while remaining stationary. Events that occurred in the

center versus perimeter sectors were tallied separately, and group means were analyzed for each

region by one-way ANOVA and Bonferroni post-hoc tests.

<u>OptiMan system</u>

Experiments were approved by the animal welfare committee of Charité Medical University and

the Berlin state government. Four groups (22 male & 4 female) of $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$

rats were used in this study. Rats carried subcutaneous radio-frequency identification (RFID) tags

in ventral location. Animals were group housed with 6 - 8 rats per group. Initially, all rats were

habituated to the OptiMan multi-cage environment with open sorter gates for voluntary

exploration. Then, automated sorting was activated so that only one rat could enter the operant

chamber at a time. During pull-task training, force thresholds and handle positions were adapted

every day for each session to the current skill level reached by a rat. Each session had 20 trials,

and two to three individual sessions were given per day with an intersession interval of 30 to 60

min. Within a session, the maximum time interval a rat was allowed to remain inactive between

trials was 6 - 8 min. A session ended when such inactivity occurred. The OptiMan (Operator

Independent Motor-Analysis, PhenoSys) system consists of two interconnected group home cages

(EU Type IV cage and 2000P) resting on an RFID sensor array plate that automatically tracks

RFID tag movement within the cage, and thus the locomotor activity of individual rats, day and

night. One of the home cages was connected via an electronic guillotine gate to a second cage

resting on a balance that automatically determined the body mass of a rat when inside this cage.

Individual animals voluntarily and self-motivated decided to visit the balance cage. Subsequent to

the balance cage, and again separated by an electronic guillotine gate, was a cage compartment

containing a horizontal ladder (1 m) with electronically monitored dual rungs on the left and right

side. From this ladder compartment, an animal entered a cage containing the isometric pull-task.

From there back to the home cage a rat passed over a force grid sensor array that sensed ground

forces exerted by the paws.

<u>Isometric pull task</u>

Rats were trained to pull a handle attached to a stationary force transducer with a predetermined

force threshold upon which they received a sugar pellet reward. Upon a rat's entry into the

isometric pull task cage a sugar pellet was delivered into the reward tray. A session started when

the rat retrieved this first reward, which led to the automatic closing of the entry gate, and to the

motorized slide-in appearance of the force handle to its predefined position. During each session,

5 different handle positions were presented to a rat. These positions varied from 11 mm inside the

cage wall (easiest position), to 7 mm outside of the cage wall (most difficult position). Handle

positions changed automatically during a session through a motorized slide. The difficulty level

within each handle position increased stepwise by increasing the force threshold for pellet release.

This started at 30 g pull-force and was increased to 40 g, 50 g and finally 60 g. A trial started

when a pull-force of 5 g was sensed. From then on, the pull-force was sampled for a duration of 4

sec. If a rat reached the force threshold within a 2 sec time interval then a trial was successful and

a reward delivered. The schedule then advanced to the next level of difficulty by either increasing

force threshold or moving the handle one position further towards the outside. Thus, a rat needed

a minimum of 20 trials to complete a session with 5 different handle positions and 4 different

force thresholds at each position. If the threshold was not reached then a trial was considered

unsuccessful and the rat had to continue with its next trial with unchanged conditions.

Experiments lasted for 15 experimental days with 2 - 3 daily sessions per individual.

Animal welfare

All animals were treated and cared for in accordance with national and institutional guidelines:

Generation of mutant Piccolo rat strains and behavioral experiments Figure 10 a-e: Rat

protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at UT-

Southwestern Medical Center in Dallas, as certified by the Association for Assessment and

Accreditation of Laboratory Animal Care International (AALAC), permit number: NIH OLAW

Assurance # D16-00296.

Western blotting, immunohistochemistry, electron microscopy and behavioral

experiments with Optiman setup (Figure 10 d and e): Animals were treated in accordance with

the German Protection of Animals Act (TierSchG §4 Abs. 3); all procedures for experiments

involving animals were approved by the animal welfare committee of Charité Medical University

and the Berlin state government, permit number: T 0036/14.

Electrophysiology: Animals were treated in accordance with the German Protection of Animals

Act (TierSchG §4 Abs. 3) and with the guidelines for the welfare of experimental animals issued

by the European Communities Council Directive of 24. November 1986 (86/609/EEC). The local

authorities approved the experiments (Landesdirektion Leipzig), permit number: T24/18

Experimental design and statistical analysis

GraphPad Prism (RRID:SCR_002798) was used to analyze and represent data. Statistical design,

sample sizes and tests for each experiment can be found in the figure legends.

References

Ackermann F, Schink KO, Bruns C, Izsvak Z, Hamra FK, Rosenmund C, Garner CC (2019)
753   Critical role for Piccolo in synaptic vesicle retrieval. Elife 8.
754   Ahmad-Annuar A, Ciani L, Simeonidis I, Herreros J, Fredj NB, Rosso SB, Hall A, Brickley S,
755   Salinas PC (2006) Signaling across the synapse: a role for Wnt and Dishevelled in
756   presynaptic assembly and neurotransmitter release. J Cell Biol 174:127-139.
757   Ahmed MY, Chioza BA, Rajab A, Schmitz-Abe K, Al-Khayat A, Al-Turki S, Baple EL, Patton
758   MA, Al-Memar AY, Hurles ME, Partlow JN, Hill RS, Evrony GD, Servattalab S,
759   Markianos K, Walsh CA, Crosby AH, Mochida GH (2015) Loss of PCLO function
760   underlies pontocerebellar hypoplasia type III. Neurology 84:1745-1750.
761   Apps R, Garwicz M (2005) Anatomical and physiological foundations of cerebellar information
762   processing. Nat Rev Neurosci 6:297-311.
763   Brickley SG, Cull-Candy SG, Farrant M (1996) Development of a tonic form of synaptic
764   inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA
765   receptors. J Physiol 497 (Pt 3):753-759.
766   Byczkowicz N, Ritzau-Jost A, Delvendahl I, Hallermann S (2018) How to maintain active zone
767   integrity during high-frequency transmission. Neurosci Res 127:61-69.
768   Cases-Langhoff C, Voss B, Garner AM, Appeltauer U, Takei K, Kindler S, Veh RW, De Camilli
769   P, Gundelfinger ED, Garner CC (1996) Piccolo, a novel 420 kDa protein associated with
770   the presynaptic cytomatrix. Eur J Cell Biol 69:214-223.
771   Chen AI, Zang K, Masliah E, Reichardt LF (2016) Glutamatergic axon-derived BDNF controls
772   GABAergic synaptic differentiation in the cerebellum. Sci Rep 6:20201.
773   Curzon P, Zhang M, Radek RJ, Fox GB (2009) The Behavioral Assessment of Sensorimotor
774   Processes in the Mouse: Acoustic Startle, Sensory Gating, Locomotor Activity, Rotarod,
775   and Beam Walking. In: Methods of Behavior Analysis in Neuroscience (nd, Buccafusco
776   JJ, eds). Boca Raton (FL).
777   Durmaz B, Wollnik B, Cogulu O, Li Y, Tekgul H, Hazan F, Ozkinay F (2009) Pontocerebellar
778   hypoplasia type III (CLAM): Extended phenotype and novel molecular findings. J Neurol
779   256:416-419.
780   Eccles JC, Llinás R, Sasaki K (1966) The mossy fibre-granule cell relay of the cerebellum and its
781   inhibitory control by Golgi cells. Experimental Brain Research 1:82-101.
782   Fenster SD, Garner CC (2002) Gene structure and genetic localization of the PCLO gene
783   encoding the presynaptic active zone protein Piccolo. Int J Dev Neurosci 20:161-171.
784   Fenster SD, Chung WJ, Zhai R, Cases-Langhoff C, Voss B, Garner AM, Kaempf U, Kindler S,
785   Gundelfinger ED, Garner CC (2000) Piccolo, a presynaptic zinc finger protein
786   structurally related to bassoon. Neuron 25:203-214.
787   Gao C, Chen YG (2010) Dishevelled: The hub of Wnt signaling. Cell Signal 22:717-727.
788   Gundelfinger ED, Reissner C, Garner CC (2015) Role of Bassoon and Piccolo in Assembly and
789   Molecular Organization of the Active Zone. Front Synaptic Neurosci 7:19.
790   Hall AC, Lucas FR, Salinas PC (2000) Axonal remodeling and synaptic differentiation in the
791   cerebellum is regulated by WNT-7a signaling. Cell 100:525-535.
792   Hashimoto K, Kano M (1998) Presynaptic origin of paired-pulse depression at climbing fibre-
793   Purkinje cell synapses in the rat cerebellum. J Physiol 506 (Pt 2):391-405.
794   Hashimoto K, Ichikawa R, Kitamura K, Watanabe M, Kano M (2009) Translocation of a
795   "winner" climbing fiber to the Purkinje cell dendrite and subsequent elimination of
796   "losers" from the soma in developing cerebellum. Neuron 63:106-118.
797   Heyden A, Ionescu MCS, Romorini S, Kracht B, Ghiglieri V, Calabresi P, Seidenbecher C,
798   Angenstein F, Gundelfinger ED (2011) Hippocampal enlargement in Bassoon-mutant
799   mice is associated with enhanced neurogenesis, reduced apoptosis, and abnormal BDNF
800   levels. Cell Tissue Res 346:11-26.

801 Homanics GE, Ferguson C, Quinlan JJ, Daggett J, Snyder K, Lagenaur C, Mi Z-P, Wang X-H,
802  Grayson DR, Firestone LL (1997) Gene Knockout of the α6 Subunit of the γ-
803  Aminobutyric Acid Type A Receptor: Lack of Effect on Responses to Ethanol,
804  Pentobarbital, and General Anesthetics. Molecular Pharmacology 51:588-596.
805 Hubler D, Rankovic M, Richter K, Lazarevic V, Altrock WD, Fischer KD, Gundelfinger ED,
806  Fejtova A (2012) Differential spatial expression and subcellular localization of CtBP
807  family members in rodent brain. PLoS One 7:e39710.
808 Human Protein Atlas (2015) PCLO. In.
809 Ichikawa R, Hashimoto K, Miyazaki T, Uchigashima M, Yamasaki M, Aiba A, Kano M,
810  Watanabe M (2016) Territories of heterologous inputs onto Purkinje cell dendrites are
811  segregated by mGluR1-dependent parallel fiber synapse elimination. Proc Natl Acad Sci
812  U S A 113:2282-2287.
813 Ivanova D, Dirks A, Montenegro-Venegas C, Schone C, Altrock WD, Marini C, Frischknecht R,
814  Schanze D, Zenker M, Gundelfinger ED, Fejtova A (2015) Synaptic activity controls
815  localization and function of CtBP1 via binding to Bassoon and Piccolo. EMBO J
816  34:1056-1077.
817 Ivics Z, Li MA, Mates L, Boeke JD, Nagy A, Bradley A, Izsvak Z (2009) Transposon-mediated
818  genome manipulation in vertebrates (vol 6, pg 415, 2009). Nat Methods 6:546-546.
819 Izsvak Z, Frohlich J, Grabundzija I, Shirley JR, Powell HM, Chapman KM, Ivics Z, Hamra FK
820  (2010) Generating knockout rats by transposon mutagenesis in spermatogonial stem cells.
821  Nat Methods 7:443-445.
822 Jakab RL, Hamori J (1988) Quantitative Morphology and Synaptology of Cerebellar Glomeruli in
823  the Rat. Anat Embryol 179:81-88.
824 Leto K et al. (2016) Consensus Paper: Cerebellar Development. Cerebellum 15:789-828.
825 Maex R, De Schutter E (1998) Synchronization of golgi and granule cell firing in a detailed
826  network model of the cerebellar granule cell layer. J Neurophysiol 80:2521-2537.
827 Maricich SM, Aqeeb KA, Moayedi Y, Mathes EL, Patel MS, Chitayat D, Lyon G, Leroy JG,
828  Zoghbi HY (2011) Pontocerebellar Hypoplasia: Review of Classification and Genetics,
829  and Exclusion of Several Genes Known to Be Important for Cerebellar Development.
830  Journal of Child Neurology 26:288-294.
831 Medrano GA, Singh M, Plautz EJ, Good LB, Chapman KM, Chaudhary J, Jaichander P, Powell
832  HM, Pudasaini A, Shelton JM, Richardson JA, Xie X-J, Ivics Z, Braun C, Ackermann F,
833  Garner CC, Izsvák Z, Hamra FK (2019) Mutant screen reveals depression-associated
834  *Piccolo's* control over brain-gonad cross talk and reproductive behavior.
835  bioRxiv:405985.
836 Metz GA, Whishaw IQ (2009) The ladder rung walking task: a scoring system and its practical
837  application. J Vis Exp.
838 Miyashita S, Adachi T, Yamashita M, Sota T, Hoshino M (2017) Dynamics of the cell division
839  orientation of granule cell precursors during cerebellar development. Mech Dev 147:1-7.
840 Miyazaki T, Fukaya M, Shimizu H, Watanabe M (2003) Subtype switching of vesicular
841  glutamate transporters at parallel fibre-Purkinje cell synapses in developing mouse
842  cerebellum. Eur J Neurosci 17:2563-2572.
843 Muller TM, Gierke K, Joachimsthaler A, Sticht H, Izsvak Z, Hamra FK, Fejtova A, Ackermann
844  F, Garner CC, Kremers J, Brandstatter JH, Regus-Leidig H (2019) A Multiple Piccolino-
845  RIBEYE Interaction Supports Plate-Shaped Synaptic Ribbons in Retinal Neurons. J
846  Neurosci 39:2606-2619.
847 Namavar Y et al. (2011) Clinical, neuroradiological and genetic findings in pontocerebellar
848  hypoplasia. Brain 134:143-156.
849 Nusser Z, Sieghart W, Somogyi P (1998) Segregation of different GABAA receptors to synaptic
850  and extrasynaptic membranes of cerebellar granule cells. J Neurosci 18:1693-1703.

Nusser Z, Sieghart W, Stephenson F, Somogyi P (1996) The alpha 6 subunit of the GABAA
receptor is concentrated in both inhibitory and excitatory synapses on cerebellar granule
cells. The Journal of Neuroscience 16:103-114.
Rajab A, Mochida GH, Hill A, Ganesh V, Bodell A, Riaz A, Grant PE, Shugart YY, Walsh CA
(2003) A novel form of pontocerebellar hypoplasia maps to chromosome 7q11-21.
Neurology 60:1664-1667.
Regus-Leidig H, Fuchs M, Lohner M, Leist SR, Leal-Ortiz S, Chiodo VA, Hauswirth WW,
Garner CC, Brandstatter JH (2014) In vivo knockdown of Piccolino disrupts presynaptic
ribbon morphology in mouse photoreceptor synapses. Front Cell Neurosci 8:259.
Rothman JS, Silver RA (2018a) NeuroMatic: An Integrated Open-Source Software Toolkit for
Acquisition, Analysis and Simulation of Electrophysiological Data. Front Neuroinform
12:14.
Rothman JS, Silver RA (2018b) NeuroMatic: An Integrated Open-Source Software Toolkit for
Acquisition, Analysis and Simulation of Electrophysiological Data. Front Neuroinform
12.
Rudnik-Schoneborn S, Barth PG, Zerres K (2014) Pontocerebellar hypoplasia. Am J Med Genet
C Semin Med Genet 166C:173-183.
Salinas PC, Zou YM (2008) Wnt signaling in neural circuit assembly. Annual Review of
Neuroscience 31:339-358.
Scheiffele P (2003) Cell-cell signaling during synapse formation in the CNS. Annual Review of
Neuroscience 26:485-508.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P,
Cardona A (2012) Fiji: an open-source platform for biological-image analysis. Nat
Methods 9:676-682.
Shinoda Y, Sugiuchi Y, Futami T, Izawa R (1992) Axon Collaterals of Mossy Fibers from the
Pontine Nucleus in the Cerebellar Dentate Nucleus. Journal of Neurophysiology 67:547-
560.
Sillitoe R, Fu, Y., Watson, C. (2012) Chapter 11 - Cerebellum: Academic Press.
Solecki DJ, Liu XL, Tomoda T, Fang Y, Hatten ME (2001) Activated Notch2 signaling inhibits
differentiation of cerebellar granule neuron precursors by maintaining proliferation.
Neuron 31:557-568.
Turrigiano G (2012) Homeostatic synaptic plasticity: local and global mechanisms for stabilizing
neuronal function. Cold Spring Harb Perspect Biol 4:a005736.
Voogd J, Glickstein M (1998) The anatomy of the cerebellum. Trends Neurosci 21:370-375.
Wagh D, Terry-Lorenzo R, Waites CL, Leal-Ortiz SA, Maas C, Reimer RJ, Garner CC (2015)
Piccolo Directs Activity Dependent F-Actin Assembly from Presynaptic Active Zones
via Daam1. PLoS One 10:e0120093.
Waites CL, Craig AM, Garner CC (2005) Mechanisms of vertebrate synaptogenesis. Annual
Review of Neuroscience 28:251-274.
Waites CL, Leal-Ortiz SA, Okerlund N, Dalke H, Fejtova A, Altrock WD, Gundelfinger ED,
Garner CC (2013) Bassoon and Piccolo maintain synapse integrity by regulating protein
ubiquitination and degradation. EMBO J 32:954-969.
Wallace VA (1999) Purkinje-cell-derived Sonic hedgehog regulates granule neuron precursor cell
proliferation in the developing mouse cerebellum. Curr Biol 9:445-448.
Xu-Friedman MA, Regehr WG (2003) Ultrastructural contributions to desensitization at
cerebellar mossy fiber to granule cell synapses. Journal of Neuroscience 23:2182-2192.
Zelnik N, Dobyns WB, Forem SL, Kolodny EH (1996) Congenital pontocerebellar atrophy in
three patients: clinical, radiologic and etiologic considerations. Neuroradiology 38:684-
687.

Zhai RG, Vardinon-Friedman H, Cases-Langhoff C, Becker B, Gundelfinger ED, Ziv NE, Garner
CC (2001) Assembling the presynaptic active zone: a characterization of an active one
precursor vesicle. Neuron 29:131-143.
904

 Figure Legends

Figure 1. Generation of $Pclo^{gt/gt}$ mutant animals.

A) Sleeping beauty transposon mutagenesis was used to generate gene trap (gt) piccolo knockout

rats. The transposon element inserted into exon 3 of the piccolo genomic sequence and caused a

stop in the reading frame. Adapted from Ackermann et al. (Ackermann et al., 2019).

B) Pairs of heterozygous ($Pclo^{wt/gt}$) males and females produced litters with Mendelian

distribution. Pie chart demonstrates the birth rates of homozygous wildtype ($Pclo^{wt/wt}$),

homozygous gene trap mutation ($Pclo^{gt/gt}$) and heterozygous ($Pclo^{wt/gt}$) pups.

C) Western blot analysis of brain lysates prepared from postnatal day 2 (P2) animals to confirm

the loss of full-length Piccolo protein from the brain. A band of the Piccolo-corresponding size of

560 kDa is detectable in lysates prepared from $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ animals, but is absent in

$Pclo^{gt/gt}$ brain lysates (data are representative of 3 independent experiments). However, smaller

bands between 100 and 70 kDA are still present in $Pclo^{gt/gt}$ brain lysates.

D-F) Image of postnatal day 1 (P1) littermates (D), E) Quantification of the body length of P0-P2

$Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ pups ($Pclo^{wt/wt}$ = 5.5 cm ± 0.077, n = 23; $Pclo^{gt/gt}$ = 5.15 cm ± 0.070, n = 27; 6

independent litters; unpaired t-test, p^{**} = 0.0014). F) Quantification of the body weight of P0-P2

$Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ pups ($Pclo^{wt/wt}$ = 8.09 g ± 0.203, n = 39; $Pclo^{gt/gt}$ = 7.31 g ± 0.166, n = 35; 12

independent litters; unpaired t-test, p^{**} = 0.0044).

G-H) Image of brains dissected from P1 $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ pups (G). H) Quantification of the

brain weight of P0-P2 $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ pups ($Pclo^{wt/wt}$ = 0.293 ± 0.00533, n = 7;

$Pclo^{gt/gt}$ = 0.289 g ± 0.00758 n = 3; 3 independent litters; unpaired t-test, p = 0.683)

I) Nissl stained sagittal sections from P2 rat brains show no overt differences between $Pclo^{gt/gt}$

and $Pclo^{wt/wt}$ pups. Scale bar = 0.5 cm.

J-L) Image of 4% PFA-perfused brains from $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ animals at 3 months of age (I),

K) Quantification of the brain weight showing that $Pclo^{gt/gt}$ brains are significantly lighter than

Pclo[wt/wt] brains (*Pclo*[wt/wt] = 2.098 g ± 0.074, *Pclo*[gt/gt] = 1.435 g ± 0.021; n = 6, Mann-Whitney U

test, $p^{**} = 0.0022$).

L) Nissl stained sagittal sections from 3 month-old rat brains reveal microcephaly in *Pclo*[gt/gt]

compared to *Pclo*[wt/wt]. Note, ventricles are larger and cerebellum, pons, cerebrum and subcortical

regions are smaller.

Scale bars: 1 cm (G and K); 0.5 cm (I). Error bars represent SEM.

Figure 2. *Pclo*[gt/gt] rats show cortical thinning and a smaller pons area compared to *Pclo*[wt/wt]

littermates.

A-B) Nissl stained somatosensory cortex (indicated by dashed lines) (A) and the brainstem

including pontine area (B) of *Pclo*[wt/wt] and *Pclo*[gt/gt] brains at 3 month of age. B) Zoom

demonstrates that pontine neurons are of similar density in *Pclo*[wt/wt] and *Pclo*[gt/gt].

C) Pontine area visualized by staining with antibodies against VGluT1 and subsequent DAB

conversion of *Pclo*[wt/wt] and *Pclo*[gt/gt] brains at 3 month of age.

D) Quantification of the thickness of the somatosensory cortex (*Pclo*[wt/wt] = 2.18 mm ± 0.045, n =

25 brain sections; *Pclo*[gt/gt] = 1.98 mm ± 0.041, n = 21 brain sections; n = 3 independent

experiments; unpaired t-test, $p^{**} = 0.0018$).

E) Quantification of the area of the pons (*Pclo*[wt/wt] = 12.24 mm^2 ± 0.620, n = 26 brain sections;

Pclo[gt/gt] = 5.58 mm^2 ± 0.333, n = 17 brain sections; n = 3 independent experiments; unpaired t-

test, $p^{****} < 0.0001$).

Scale bars: 200 μm (A and C), 500 μm (B and C) and 100 μm (B, zoom). Error bars represent

SEM.

Figure 3. Morphometric comparison of *Pclo*[gt/gt] and *Pclo*[wt/wt] cerebella.

A-D) Images of sagittal sections of *Pclo*[wt/wt] and *Pclo*[gt/gt] cerebella at 3 month of age. The densely

packed granule cell layer (GCL) is visualized by DAPI staining. B) Higher magnification images

of (A) demonstrating GC density in the GCL. C) Quantification of the GCL thickness ($Pclo^{wt/wt}$ =

200.8 μm ± 2.932, n = 160 lobes; $Pclo^{gt/gt}$ = 134.8 μm ± 2.859, n = 148 lobes; n = 4 independent

experiments; Mann-Whitney U test, $p^{****} < 0.0001$). D) Quantification of the number of GCs per

100 μm² ($Pclo^{wt/wt}$ = 1.702 ± 0.020, n = 111 images; $Pclo^{gt/gt}$ = 1.156 ± 0.022, n = 114 images; n =

3 independent experiments; unpaired t-test, p^{**} = 0.0042).

E-H) Images of sagittal sections of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ cerebella at 3 months of age. Purkinje

cells (PCs) stained with antibodies against Calbindin determine the molecular layer (ML) (lobes

I-III shown). F) Higher magnification images of (E). Note the closer packing of PCs in $Pclo^{gt/gt}$

compared to $Pclo^{wt/wt}$. G) Quantification of the ML thickness ($Pclo^{wt/wt}$ = 187.2 μm ± 2.719, n =

148 lobes; $Pclo^{gt/gt}$ = 127.2 μm ± 2.378, n = 125 lobes; n = 4 independent experiments; Mann-

Whitney U test, $p^{****} < 0.0001$). H) Quantification of the number of PCs per 100 μm length of

PC layer ($Pclo^{wt/wt}$ = 1.797 ± 0.036; n = 89 lobes; $Pclo^{gt/gt}$ = 2.554 ± 0.058; n = 65 lobes; n = 3

independent experiments; unpaired t-test, $p^{****} < 0.0001$).

I) Images of sagittal sections stained with antibodies against Calbindin showing that PCs migrate

correctly to their position in the ML and are correctly orientated.

Scale bars: 1cm (A), 20 μm (B and J), 50 μm (I) and 200 μm (F). Error bars represent SEM. Data

points represent images taken from lobes I, III, V, VII and IX; 4 sections per animal (B and D).

Figure 4. Abberant climbing fiber innervation of Purkinje cells in $Pclo^{gt/gt}$ rats compared to

$Pclo^{wt/wt}$ littermates.

A and B) Images of sagittal sections of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ cerebella at 3 months of age stained

with antibodies against VGluT1 and Piccolo demonstrate the loss of Piccolo in the ML (A) and

the GCL (B) in $Pclo^{gt/gt}$ rats.

C) Images of sagittal sections of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ cerebella at 3 months of age stained with

antibodies against Calbindin and VGluT2. Note that the climbing fiber synapses, immuno-

positive for VGluT2, are increased in the ML of $Pclo^{gt/gt}$ cerebella compared to $Pclo^{wt/wt}$ controls

(C and D). When Piccolo is absent, parallel fiber synapses in the ML - immuno-positive for

VGluT1 – do not appear overtly different between $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ (A). However, MF and CF

synapses are altered (4B-D, Figure 5).

D) Quantification of the percentage of the ML (indicated by dashed lines) immuno-positive for

VGluT2 from (C) ($Pclo^{wt/wt}$ = 3.521 ± 0.160, n = 128 images; $Pclo^{gt/gt}$ = 4.377 ± 0.241, n = 112

images; n = 3 independent experiments; Mann-Whitney U test, $p*$ = 0.0278).

Scale bars: 20 μm. Error bars represent SEM. Data points represent images taken from lobes I,

III, V, VII and IX; 4 sections per animal.

Figure 5. Cerebella from $Pclo^{gt/gt}$ animals have smaller mossy fiber rosettes.

A and B) Images of sagittal sections of $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ cerebella at 3 month of age stained

with antibodies against VGluT2, which is highly expressed at mossy fiber (MF) boutons, and the

somatodendritic marker MAP2 (A) or VGluT2 and VGluT1 (B). A) GCs extend their dendrites

into MF boutons in the $Pclo^{wt/wt}$ condition. However, in $Pclo^{gt/gt}$, whilst GC dendrites are still

juxtaposed to VGluT2-positive boutons, the boutons are smaller and therefore the arrangement is

less organized. B) Presynaptic MF glomeruli from lobes I (upper panel) and VII (lower panel) are

visualized by VGluT1 and VGluT2. Note that the reduction in MF size is consistent regardless of

the lobe. Note also that rosettes are generally labeled with either VGluT1 or VGluT2 and

occasionally with both markers consistent with them being innervated by a single synaptic input

from different neuronal cell types.

C) Quantification of the size of VGluT1/VGluT2 clusters ($Pclo^{wt/wt}$ = 22.73 μm^2 ± 0.896, n = 4

animals; $Pclo^{gt/gt}$ = 9.46 μm^2 ± 0.2899, n = 4 animals; unpaired t-test, $p****$ < 0.0001).

D-E) Histograms to show the distribution of puncta sized 5-50 μm^2 (D) and 50-260 μm^2 (E). The

shift of the data indicates $Pclo^{gt/gt}$ MFs have more smaller puncta (5 μm^2), whereas $Pclo^{wt/wt}$ MFs

have more larger puncta (up to 260 μm^2).

Scale bars: 50 μm (A and B) and 20 μm (B, zoom). Error bars represent SEM. Data points

represent average puncta size per animal from images taken from lobes I, III, V, VII and IX, 4

sections per animal (C, D and E).

1012

Figure 6. GABA$_A$α6 subunit expression is lower in *Pclo*$^{gt/gt}$ than in *Pclo*$^{wt/wt}$.

A) Representative images of sagittal sections of *Pclo*$^{wt/wt}$ and *Pclo*$^{gt/gt}$ cerebella at 3 months of age

stained with antibodies against the GABA$_A$ subunit α6,VGluT2 and MAP2. Note the decreased

levels of GABA$_A$α6 in the GC layer of *Pclo*$^{gt/gt}$ animals compared to *Pclo*$^{wt/wt}$ controls, quantified

in B).

B) Quantification of GABA$_A$ subunit α6, measured by the average intensity of antibody staining

in images taken from the GC of the cerebellum, normalized to MAP2 antibody intensities for the

same image (*Pclo*$^{wt/wt}$ = 0.661 ± 0.0479 arbitrary units (a.u.), n = 45 images from 3 individual

animals; *Pclo*$^{gt/gt}$ = 0.443 ± 0.26 a.u., n = 44 images from 3 individual animals; Mann-Whitney U

test, p*** = 0.0009).

Scale bars: 20 μm. Error bars represent SEM. Data points represent images taken from 4 sections

per animal.

1025

Figure 7. Glomerular rosettes are smaller and less complex in *Pclo*$^{gt/gt}$.

A, B and G) Electron microscopy images of the GCL of *Pclo*$^{wt/wt}$ and *Pclo*$^{gt/gt}$ cerebella at 3

months of age. Granule cells are indicated by 'gc' (A and B), cerebellar glomeruli are highlighted

in blue (*Pclo*$^{wt/wt}$) and green (*Pclo*$^{gt/gt}$) (A) and the presynaptic terminals of cerebellar MF are

highlighted in blue (*Pclo*$^{wt/wt}$) and green (*Pclo*$^{gt/gt}$) (B). Asterisks mark active zones (AZs) (B) and

the zoom in (G) emphasizes the presence of more clathrin-coated vesicles (CCVs) in *Pclo*$^{gt/gt}$ MF

boutons compared to than *Pclo*$^{wt/wt}$.

(C-F) Quantification of the size of glomeruli (C), the size of the MF presynapse (D), the

complexity (squared perimeter/presynaptic area) of the MF bouton (E) and the density of AZs

(F). Note the strong decrease in glomerulus and MF bouton size (C and D) (C) ($Pclo^{wt/wt}$ = 34.59

μm^2 ± 1.287, n = 130 images; $Pclo^{gt/gt}$ = 18.82 μm^2 ± 0.853, n = 103 images; n = 3 independent

experiments; Mann-Whitney U test, p**** < 0.0001. (D) ($Pclo^{wt/wt}$ = 17.73 μm^2 ± 0.603, n = 141

images; $Pclo^{gt/gt}$ = 11.69 μm^2 ± 0.497, n = 95 images; n = 3 independent experiments; Mann-

Whitney U test, p**** < 0.0001). (E) ($Pclo^{wt/wt}$ = 90.5 ± 4.089, n = 141 images; $Pclo^{gt/gt}$ = 58.1 ±

3.818, n = 84 images; n = 3 independent experiments; Mann-Whitney U test, p**** < 0.0001).

(F) ($Pclo^{wt/wt}$ = 0.625 ± 0.021, n = 121 images; $Pclo^{gt/gt}$ = 0.652 ± 0.032, n = 84 images; n = 3

independent experiments; Mann-Whitney U test, p = 0.7390).

Scale bars: 2.5 μm (A), 1μm (B), 500 nm (G) and 250 nm (G, zoom). Error bars represent SEM.

1044

Figure 8. Physiological assessment of mossy fiber boutons.

A) Example Two-photon image of a cerebellar granule cell (GC) from a $Pclo^{gt/gt}$ rat filled with

ATTO dye.

B) Average data of biophysical properties of GCs for $Pclo^{wt/wt}$ and $Pclo^{gt/gt}$ rats. The input

resistance of GCs was higher in $Pclo^{gt/gt}$ compared to $Pclo^{wt/wt}$ (Pclowt/wt = 722.6 MΩ ± 76.24, n

= 22 cells; $Pclo^{gt/gt}$ = 1160 MΩ ± 154.9, n = 23 cells; n = 3 rats per genotype; Mann-Whitney U

test, p* = 0.0462). Whereas no differences were found in capacitance ($Pclo^{wt/wt}$ = 3.717 pF ±

0.269, n = 21 cells; $Pclo^{gt/gt}$ = 3.643 pF ± 0.231, n = 23 cells; n = 3 rats per genotype; Mann-

Whitney U test, p = 0.912), resting membrane potential (Vm) ($Pclo^{wt/wt}$ = -99.96 mV ± 1.261, n =

18 cells, $Pclo^{gt/gt}$ = -100.8 mV ± 0.786, n = 23 cells; n = 3 rats per genotype; Mann-Whitney U

test, p = 0.612), the half-duration of the action potential ($Pclo^{wt/wt}$ = 179.7 μs ± 11.84, n = 18 cells;

$Pclo^{gt/gt}$ = 172.2 μs ± 7.366, n = 23 cells; n = 3 rats per genotype; Mann-Whitney U test, p =

0.866), the amplitude of the action potential ($Pclo^{wt/wt}$ = 67.85 mV ± 3.016, n = 18 cells; $Pclo^{gt/gt}$ =

63.92 mV ± 2.761, n = 23 cells; n = 3 rats per genotype; Mann-Whitney U test, p = 0.291) and the

voltage threshold to elicit an action potential ($Pclo^{wt/wt}$ = -51.27 mV ± 1.748, n = 18 cells; $Pclo^{gt/gt}$

= -48.6 mV ± 1.659, n = 23 cells; n = 3 rats per genotype; Mann-Whitney U test, p = 0.162).

C) Miniature excitatory postsynaptic currents from $Pclo^{gt/gt}$ GCs were not different in their

amplitude ($Pclo^{wt/wt}$ = -19.62 pA ± 1.682, n = 15 cells; $Pclo^{gt/gt}$ = -22.44 pA ± 1.765, n = 23 cells;

n = 3 rats per genotype; Mann-Whitney U test, p = 0.286) but in their frequency ($Pclo^{wt/wt}$ = 0.102

Hz ± 0.0167, n = 15 cells; $Pclo^{gt/gt}$ = 0.257 Hz ± 0.0481, n = 22 cells; n = 3 rats per genotype;

Mann-Whitney U test, $p*$ = 0.0392).

D) Excitatory postsynaptic currents from GCs measured after stimulation of single mossy fibers

were increased in $Pclo^{gt/gt}$ compared to $Pclo^{wt/wt}$ ($Pclo^{wt/wt}$ = 47.58 pA ± 12.12, n = 13 cells;

$Pclo^{gt/gt}$ = 67.62 pA ± 9.64, n = 15 cells; n = 3 rats per genotype; Mann-Whitney U test, $p*$ =

0.0356), whereas the decay of the EPSC was not alerted ($Pclo^{wt/wt}$ = 1.79 ms ± 0.258, n = 12 cells;

$Pclo^{gt/gt}$ = 1.404 ms ± 0.141, n = 14 cells; n = 3 rats per genotype; Mann-Whitney test U, $p*$ =

0.231). Right hand panel: example traces of evoked EPSCs, as quantified in D), in response 1 Hz

stimulation in the presence of 20 μM SR95531 and 40 μM D-(2R)-amino-5-phosphonovaleric

acid (D-APV).

Scale bar = 20 μm (A). Error bars represent SEM. Data points represent individual cells from 3

rats per genotype.

1076

Figure 10. Behavioral outcome of Piccolo loss of function resembles PCH3 symptoms.

A) Rotarod performance for $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats for 16 trials over 4 days. $Pclo^{gt/gt}$

rats fell significantly faster than $Pclo^{wt/wt}$ on trials 6 and 8 and faster than both $Pclo^{wt/wt}$ and

$Pclo^{wt/gt}$ on trials 9-16 (trial 6: $Pclo^{wt/wt}$ = 56.25 s ± 5.023, $Pclo^{gt/gt}$ = 3.5 ± 0.5, $p*$ = 0.0293; trial 8:

$Pclo^{wt/wt}$ = 65.25 s ± 18.688, $Pclo^{gt/gt}$ = 4.5 ± 0.957, $p**$ = 0.0042; trial 9: $Pclo^{wt/wt}$ = 63.25 s ±

16.163, $Pclo^{wt/gt}$ = 61.5 s ± 18.319, $Pclo^{gt/gt}$ = 3.25 ± 0.75, $p**$ = 0.0051, $p^{\#\#}$ = 0.0079 (where * =

p value between $Pclo^{wt/w}$ and $Pclo^{gt/gt}$ and # = p value between $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$); trial 10:

$Pclo^{wt/wt}$ = 84.5 s ± 6.225, $Pclo^{wt/gt}$ = 60.25 s ± 18.396, $Pclo^{gt/gt}$ = 2.5 ± 0.5, $p****$ = <0.0001, $p^{\#\#}$ =

0.0089; trial 11: $Pclo^{wt/wt}$ = 72 s ± 5.339, $Pclo^{wt/gt}$ = 76.75 s ± 12.419, $Pclo^{gt/gt}$ = 3 ± 0.408, $p***$ =

0.0005, $p^{\#\#\#}$ =0.001; trial 12: $Pclo^{wt/wt}$ = 71 s ± 6.178, $Pclo^{wt/gt}$ = 68.75 s ± 9.681, $Pclo^{gt/gt}$ = 4 ±

0.707, $p^{***} = 0.0008$, $p^{\#\#} = 0.0015$; trial 13: $Pclo^{wt/wt} = 59$ s $\pm$ 8.297, $Pclo^{wt/gt} = 57.75$ s $\pm$ 12.99,

$Pclo^{gt/gt} = 2.5 \pm 0.5$, $p^{*} = 0.0121$, $p^{\#} = 0.0164$; trial 14: $Pclo^{wt/wt} = 69.25$ s $\pm$ 12.479, $Pclo^{wt/gt} = 61$ s

$\pm$ 13.638, $Pclo^{gt/gt} = 2.5 \pm 0.645$, $p^{***} = 0.0009$, $p^{\#\#} = 0.0074$; trial 15: $Pclo^{wt/wt} = 60.5$ s $\pm$ 16.297,

$Pclo^{wt/gt} = 65$ s $\pm$ 18.757, $Pclo^{gt/gt} = 1.75 \pm 0.479$, $p^{**} = 0.0070$, $p^{\#\#} = 0.0022$; trial 16: $Pclo^{wt/wt} =$

56.5 s $\pm$ 16.775, $Pclo^{wt/gt} = 68.75$ s $\pm$ 17.109, $Pclo^{gt/gt} = 2.25 \pm 0.25$, $p^{*} = 0.0207$, $p^{\#\#\#\#} = 0.009$; n =

4 animals per genotype; two-way ANOVA with Bonferroni correction).

B) Grip strength task for $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats for 9 trials over 2 days. No differences

were found for forelimb grip strength between the groups. ($Pclo^{wt/wt} = 737.2 \pm 109.9$; $Pclo^{wt/gt} =$

621.2 ± 93.46, $Pclo^{gt/gt} = 757.5 \pm 66.74$; n = 4 rats per genotype, one-way ANOVA, $p = 0.549$).

C) Ladder walk task for $Pclo^{wt/wt}$, 859 $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats for 3 trials over 1 day. $Pclo^{gt/gt}$ rats

had a higher rate of stepping errors (ladder rung foot slips/misses) than $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ rats

(forelimb error/step: $Pclo^{wt/wt} = 0.09 \pm 0.00925$, $Pclo^{wt/gt} = 0.148 \pm 0.0263$, $Pclo^{gt/gt} = 0.297 \pm$

0.0145, n = 4, one-way ANOVA, $p^{**} = 0.002$, $p^{\#} = 0.0157$ (where $* = p$ value between $Pclo^{wt/w}$

and $Pclo^{gt/gt}$ and $^{\#} = p$ value between $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$); hindlimb error/step: $Pclo^{wt/wt} = 0.0563$

$\pm$ 0.0221, $Pclo^{wt/gt} = 0.0663 \pm 0.00877$, $Pclo^{gt/gt} = 0.238 \pm 0.0543$, n = 4; one-way ANOVA with

Bonferroni correction, $p^{*} = 0.0135$, $p^{\#} = 0.0186$).

D) Open field task for $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats for 1 trial each over 1 day. $Pclo^{gt/gt}$ rats

performed fewer rearing behaviors than $Pclo^{wt/wt}$ and significantly less than $Pclo^{wt/gt}$ rats in the

perimeter sectors of the arena ($Pclo^{wt/wt} = 64.25 \pm 18.31$ events; $Pclo^{wt/gt} = 76.25 \pm 11.3$ events;

$Pclo^{gt/gt} = 22 \pm 4.203$ events; n = 4 rats per genotype, one-way ANOVA with Bonferroni

correction, $p^{\#} = 0.0427$ (where $^{\#} = p$ value between $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$). Other behaviors such as

crossing the open field and grooming were not different between the three groups (crossing:

$Pclo^{wt/wt} = 178.3 \pm 29.68$ events; $Pclo^{wt/gt} = 164.5 \pm 45.51$ events; $Pclo^{gt/gt} = 235.5 \pm 27.11$ events,

n = 4 rats per genotype, one-way ANOVA with Bonferroni correction, $p = 0.828$; grooming:

$Pclo^{wt/wt} = 2.75 \pm 0.75$ events; $Pclo^{wt/gt} = 2 \pm 1.08$ events; $Pclo^{gt/gt} = 5 \pm 1.155$ events, n = 4 rats

per genotype, one-way ANOVA with Bonferroni correction, $p = 0.488$)

E). Locomotor activity of $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats during the 12h dark phase. $Pclo^{gt/gt}$

rats covered a more than 50% longer distance than $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$ rats ($Pclo^{wt/wt}$ = 135.7 ±

11.25, n = 10; $Pclo^{wt/gt}$ = 129.3 ± 10.44, n = 4; $Pclo^{gt/gt}$ = 217.8 ± 14.33, n = 12; one-way

ANOVA, p^{***} = 0.0002). Data points are individual means over 15 nights.

F) Performance of $Pclo^{wt/wt}$, $Pclo^{wt/gt}$ and $Pclo^{gt/gt}$ rats during the isometric pull-task (handle

position 11 mm inside the cage). Only 4 out of 11 $Pclo^{gt/gt}$ rats succeeded at the 60 g force

threshold and $Pclo^{gt/gt}$ rats pulled with a significantly lower success rate at all force thresholds as

compared to $Pclo^{wt/wt}$ and $Pclo^{wt/gt}$. (30g: $Pclo^{wt/wt}$ = 86.722 ± 2.3, n = 10; $Pclo^{wt/gt}$ = 84.374 ±

6.324, n=4; $Pclo^{gt/gt}$ = 61.044 ± 5.928, n = 11; 2 way ANOVA with Bonferroni correction, p^{**} =

0.0018; 40g: $Pclo^{wt/wt}$ = 83.603 ± 2.86, n = 10; $Pclo^{wt/gt}$ = 85.136 ± 7.373, n=4; $Pclo^{gt/gt}$ = 47.804 ±

6.897, n = 10; two-way ANOVA with Bonferroni correction, p^{****} < 0.0001, $p^{###}$ = 0.0007; 50g:

$Pclo^{wt/wt}$ = 80.49 ± 3.442, n = 10; $Pclo^{wt/gt}$ = 71.930 ± 8.695, n=4; $Pclo^{gt/gt}$ = 33.647 ± 6.802, n =

10; two-way ANOVA with Bonferroni correction, p^{****} < 0.0001, $p^{###}$ = 0.0005 60g: $Pclo^{wt/wt}$ =

69.186 ± 4.99, n = 10; $Pclo^{wt/gt}$ = 70.409 ± 10.181, n=3; $Pclo^{gt/gt}$ = 30.003 ± 11.359, n = 4; 2 way

ANOVA with Bonferroni correction, p^{***} = 0.0004, $p^{##}$ = 0.0056).

Error bars represent SEM. Data points represent individual rats.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a

| | GABA_Aα6 | VGLUT2 | MAP2 | VGLUT2/GABA_Aα6 |

Pclo^{wt/wt}

Pclo^{gt/gt}

b

Figure 7

Figure 9