1 Size matters, so does condition: the use of a body condition index reveals

2 the costs and benefits of structural body size in an insect.

3 Zanchi[1,2], Caroline*; Moret[2], Yannick; Gillingham[2,3], Mark A. F.*

4
5 [1] *Institute of Biology, Free University of Berlin, Königin-Luise-Straße 1-3, DE-14195 Berlin,*
6 *Germany.*
7
8 [2] *UMR CNRS 6282 BioGéoSciences, Équipe Écologie Évolutive, Université Bourgogne-Franche*
9 *Comté, Dijon, France*
10
11 [3] *Biodiversity Research Institute (CSIC, Oviedo University, Principality of Asturias), Campus of*
12 *Mieres, University of Oviedo, 33600 Mieres, Spain*
13

14 *Corresponding Authors:

15 Dr. Mark Gillingham
16 mark.alan.gillingham@gmail.com
17
18 Dr. Caroline Zanchi
19 caroline.zanchi@fu-berlin.de
20

22 **Keywords:** Body condition, body condition indices, Absolute Energy Demand Hypothesis,

23 Relative Efficiency Hypothesis, insect ecology, energy reserves, dietary restriction, *Tenebrio*

24 *molitor*.

Abstract

1. Insects are core actors for the balance of many earth ecosystems, as well as an alternative source of food and feed with a low ecological footprint. A comprehensive understanding of their life history requires reliable tools. Body condition constitutes the amount of energy reserves available to a fitness trait after maintenance costs have been accounted for. Body condition is standardly estimated using Body Condition Indexes (BCIs) in vertebrates. In insects the relevance of BCIs is frequently questioned on the basis that they might not accurately reflect neither energy reserves nor fitness.. However, to date no study has tested whether the very concept of body condition is relevant in insects, i.e. whether BCIs accurately reflect the relative energy reserves allocated to fitness traits.

2. We propose that the relevance of using BCIs in insects depends on whether their structural size has a fitness cost. If on the contrary insects only benefit from a larger body size, a simple measurement of body size or mass will predict fitness, but not a BCI. We experimentally manipulated food availability at the larval and adult stage and used total fecundity of females as a fitness proxy of *Tenebrio molitor*, an important model in physiology, ecology and evolution, and one of the first insects to be considered as a source of food and feed.

3. Our results support three key assumptions of the relevance of BCIs in insects: (i) a valid BCI correlated with energy reserves corrected for a given size (i.e. relative energy reserves) and not with absolute measures of energy reserves; (ii) both structural size and body condition positively predict different components of fitness; and, (iii) the effect of body condition was dependent on resource availability, whereby its effect was only apparent and large when food was unrestricted at the larval stage and restricted at the adult stage.

4. Overall we demonstrate the relevance of using BCIs in insects. Their use should be generalized to improve fitness readouts in evolution, ecological and physiological studies, as well as improve their husbandry for commercial purposes.

Introduction

Insects make up for the majority of species on the planet and might represent the highest biomass of terrestrial animals (Bar-On et al., 2018). The ease of their maintenance and manipulation coupled with their diverse life history has resulted in several insect species becoming prominent models in physiology (Scully & Bidochka, 2006; Schneider & Chambers, 2008; Keil & Steinbrecht, 2010; Adamski et al., 2019) and in ecology and evolution (Funk et al., 2002; Boggs, 2009; Bilton et al., 2019). Moreover, there is growing interest of cultivating insects as a source of food and feed (Kim et al., 2019; van Huis, 2021). These fields often require the measurement, sometimes repeated, of fitness traits during an insect's lifespan, which is likely to be dependent on the amount of energy reserves present in individuals. For instance, the tracking of temporal variations in the nutritional state of experimental individuals could also allow the optimization of mass-rearing conditions in an industrial context (Gligorescu et al., 2020; Adámková et al., 2020). The amount of reserves present in an animal can be estimated by the direct -and invasive- quantification of several body components from the organism, such as fat, glycogen and protein content, which prevents the assessment of their predictive power on fitness and to follow their evolution. To circumvent this, **Body Condition Indexes (BCIs)** can be used.

BCIs are expected to reflect **body condition**, which is classically defined as the amount of reserves present in the body of an animal after maintenance costs have been accounted for (Rowe & Houle, 1996; Barnett et al., 2015). It is also sometimes referred to as the "general health state", "physiological state", or "quality" of an animal (Jakob et al., 1996; Wilson & Nussey, 2010). The relationship between the amount of reserves and the "quality" of an animal is likely to be more complicated than stated in most studies (Raubenheimer & Simpson, 2012; Barnett et al., 2015), which is why Hill (2011) defined body condition as being "the relative capacity to maintain optimal functionality of essential cellular processes"

in an animal. In numerous cases, this capacity relies strongly on the amount of energy

reserves an individual has at its disposal (Schulte-Hostedde et al. 2005; Nandy et al., 2012),

which has been found to predict several fitness traits in animals such as dispersal

(Bargielowski et al., 2012; Evenden et al., 2014), resistance to diseases or chemical

contamination (Townsend et al., 2010; Reid & Purcell, 2011), overwintering success

(Sinclair, 2015) and fecundity (Arrese & Soulages, 2010).

Maintenance costs are assumed to stem from the cost of maintaining **structural size**,

which is to say the amount of mass which cannot be used as a source of energy. Indeed, the

Absolute Energy Demand Hypothesis (Calder, 1984) proposes that large animals require a

larger absolute amount of energy reserves to maintain their larger size, which is diverted from

fitness traits. This hypothesis therefore assumes that maintaining a larger body size is more

costly than maintaining a smaller one. This is the reason why BCIs are constructed from

several morphological measurements: typically, the measurement of a phenotypically plastic

morphological trait which varies according to the intake of resources from the environment

(e.g. mass), corrected for a measurement of the fixed structural body size (e.g. body length).

The use of BCIs is widely accepted in ecological studies involving vertebrates, where

ethical concerns impair the ability to sacrifice animals to invasively assess the state of the

resources present in their bodies (Stevenson et al., 2006; Warner et al., 2016). Their use in

invertebrates, and especially insects, is far less widespread, where authors often rely on a

simple measurement of size or mass (sometimes used interchangeably) to predict the amount

of energy reserves experimental insects contain (Rolff & Joop, 2002; Amarillo-Suárez et al.,

2011; Yoder et al., 2010; Sturm, 2016). The main assumption behind the use of body size

instead of body condition is that the larger the insect, the higher the nutrient content (Berger

et al., 2012). This assumption would be in line with the **Relative Efficiency Hypothesis**,

which states that animals with a large body size have a more efficient energy use because

mass specific metabolic rate decreases with body size (Kleiber, 1932; Thommen et al., 2019).

Consequently, a large body size is advantageous compared to a smaller one (Tammaru et al.,

2002).

A major limitation of this assumption is that insects undergo a complex life cycle in

which the next developmental stage is achieved by shedding a rigid exoskeleton, the cuticle,

therefore allowing further growth (Anderson, 1974; Reim, 2009; Knapp & Knappova, 2013,

Moya-Laraño et al., 2008). In holometabolous insects, which undergo a complete

metamorphosis between larval and adult stages, adult size is determined by the size reached

by the larva before entering the pupal stage (Chafino et al., 2019). Therefore, the cuticle of

the newly emerged insect is produced using resources of the previous developmental stage,

which means that a simple measurement of structural size will represent past developmental

conditions instead of current status (Anderson, 1974; Reim, 2009; Knapp & Knappova, 2013;

reviewed in Knapp & Uhnavá, 2014), whereas a simple measurement of mass will include the

mass of the exoskeleton, which can in some species be relatively heavy and costly to

maintain (Locke, 1991). On the one hand, adult structural size reflects the amount of

resources accumulated by the larvae and the absolute amount of resources available to the

adult. On the other hand, the quantity of resources accumulated during adult life can vary and

the relative amount of resources available for fitness traits may be constrained by structural

size.

Insect body size has been shown to be positively correlated with fitness traits on

numerous occasions (see Whitman, 2008, for review). Bigger females of the cricket

Teleogryllus commodus lay more eggs (Sturm, 2016), as well as females of the moth

Streblote panda (Calvo & Molina, 2005). Larger males sire a higher proportion of the

offspring of females of the field cricket *Gryllus firmus* (Saleh et al., 2013). Body size is even

assumed to be the main constraint on female fecundity (Honěk et al., 1993; Tammaru et al.,

2002). On the other hand, studies which found a cost to a large structural body size in insects

(reviewed in Blanckenhorn, 2000) identified mainly an increased predation risk but no direct

energetic cost. Teuschl et al. (2007) observed a developmental cost of the larvae associated with the selection for a larger structural body size in the fly *Scathophaga stercoraria*, and pointed out that while there is strong empirical evidence of selection for large body size in insects, there is little evidence of the contrary.

Another important aspect to consider is that the use of BCIs in insects has been criticized because of their lack of correlation with the absolute amount of reserves present in an animal's body and its fitness (see Wilder et al., 2016 for a review). Paradoxically, proponents of the use of BCIs argue that it is erroneous to invalidate BCIs on the basis of low correlation with absolute energy reserves since they are precisely designed to correlate with energy reserves corrected for a given size (i.e. relative energy reserves). Thus, their correlation with the absolute quantity of energetic reserves present in individuals is a sign that they do not properly control for structural size (DeBano, 2008; Peig and Green 2009, 2010).

We propose that the relevance of using BCIs in insects will depend on whether there is a significant cost to structural size. This cost would ultimately indicate the relative support for the Absolute Energy Demand or the Relative Efficiency Hypotheses in an organism in a given context. In this study we aimed to provide a thorough understanding of the biological consequences of interchangeably using body mass, structural size and body condition as a single proxy of energy reserves and predictor of fitness. We used an experimental approach using the mealworm beetle *Tenebrio molitor* (Coleoptera: Tenebrionidae), an insect massively reared as a source of food and for which breeder try to maximize body mass (Morales-Ramos et al., 2019; Costa et al., 2020). Fitness costs have been shown to increase in stressful environments (Schmid-Hempel and Schmid-Hempel 1998). We therefore assessed the cost of structural size on control and populations under food restriction at the larval and/or adult stage. First, we aimed to validate the use of structural size as a proxy of absolute energy reserves and BCIs as a proxy of relative energy reserves. This was done by sacrificing individuals, quantifying the lipids and sugars present in their bodies and correlating their

156 absolute and relative values with body mass, structural size and BCIs. Second, we

157 investigated the effect of body mass, structural size and body condition on the fitness

158 parameter fecundity across feeding treatments.

Materials and methods

Body mass and elytron size (as a measure of structural size) of adult female beetles

were taken. From these measurements, we calculated 2 body condition indexes chosen from

the literature, and explored their adequacy in controlling for the interdependence between a

phenotypically plastic morphometric measurement and structural size. We did so on groups

of female beetles having experienced food shortage or not at the larval and/or the adult stage

(4 experimental treatments). From these females, a subset was killed after measuring in order

to assess the quantity of energy reserves contained in their abdomens invasively, whereas a

subset was allowed to reproduce in order to assess their fecundity. This way, we could

evaluate the predictive power of body condition indexes on both the actual amount of

reserves and female fecundity.

Insect rearing and maintenance

Young larvae (~ 1cm) were collected from a mass rearing outbred stock kept in the

laboratory at the University of Burgundy in wheat bran (as a substrate, is also used as a

source of food) with *ad libitum* access to water, and regularly supplemented with piglet flour

(as a source of proteins) and a piece of apple (for carbohydrates and vitamins). Our insects

are kept at 25°C and in the dark. After collection, we reared *T. molitor* larvae either in a rich

or a poor larval feeding treatment, and subsequently maintained adults in a matching or

mismatching environment, i.e. similar or dissimilar to the larval feeding treatment

respectively.

Generation of larval feeding treatments

After retrieving the larvae from the stock, we maintained them at a density of 250

larvae in a plastic box (L xH x l = 30 x 22 x 25 cm) in 5 litres of wheat bran with *ad libitum*

access to water. We kept them in 2 separate containers according to the larval feeding

treatment we subjected them to while controlling for density:

- In "**rich larval feeding**" conditions, larvae were kept similarly to our stock, that is to

say in wheat bran + piglet flour + apple.

- In "**poor larval feeding**" conditions, larvae were simply kept in wheat bran with

unlimited access to water and no further addition of piglet flour or apple. Importantly,

larvae in this treatment took approximately 2 months longer to reach the pupal stage

(data not shown).

All experimental individuals originate from these 2 boxes, which we checked daily for the

presence of pupae.

The dynamics of pupation in our rearing conditions is such that a few larvae pupated

early, preceding a "pupation peak" over 3 days, after which some larvae kept pupating late.

Moreover, the growth of *T. molitor* is such that while getting closer to pupation, the food

uptake of larvae stops as the insects are undergoing the physiological changes preparing

metamorphosis (Connat et al., 1991). We retrieved pupae as soon as they were spotted, but

we used in this experiment only the pupae formed during this "pupation peak". This way, we

assumed that we minimized as much as possible the effect of the removal of conspecifics on

the accumulation of energy reserves by each individual larva. The resulting pupae were kept

separately in a plastic box and checked daily for the presence of emerged adults in the teneral

stage.

After emergence, adults were kept individually in Petri dishes supplied with wheat

bran and a piece of apple for 5 days, time during which they reach sexual maturity. Five days

after emergence the beetles were sclerotized enough to allow the determination of their sex

by observation of their genitalia. As our study focused on variation in females body condition

only, "poor larval" males were discarded, while "rich larval" males were transferred from

their Petri dish into new plastic boxes where they were kept together according to their

emergence date in wheat bran + *ad libitum* water + piglet flour + apple, waiting to be used in

the later copulation experiments.

Generation of adult female treatments

In this part of the experiment, wheat bran was replaced by thin flour in order to facilitate the

recovery of the eggs by sieving (600 µm). Five days after emergence, females were

maintained in Petri dishes according to the adult feeding treatment we subjected them to:

- "**poor adult feeding**" conditions, where females were kept in whole grain thin wheat

flour, with *ad libitum* access to water.

- "**rich adult feeding**" conditions, where females were kept similarly but with addition

of piglet flour and apple every second day.

The final sample sizes of the females of the four treatment combinations, aged 10

days old after emergence by the beginning of the experiment, were :

- 74 females in the "poor larval – poor adult" feeding treatment, 17 killed for

assessment of their reserves, 56 allowed to reproduce

- 71 females in the "poor larval – rich adult" feeding treatment, 16 killed for

assessment of their reserves, 55 allowed to reproduce

- 85 females in the "rich larval – poor adult" feeding treatment, 19 killed for

assessment of their reserves, 54 allowed to reproduce

- 97 females in the "rich larval – rich adult" feeding treatment, 37 killed for

assessment of their reserves, 56 allowed to reproduce

Measures of body condition

Measure of morphological parameters

The mass of each female was measured on a Sartorius balance to a precision of 10^{-5} g before the first reproduction. Elytron length, a proxy of abdomen size (place of storage of reserves), of the *T. molitor* females was taken with a digital calliper (Mitutoyo, Absolute) to the nearest 0.01 mm. Repeatability of elytron length measurements was assessed by taking them twice on the same individual in a non-consecutive way. The females used for assessment of the energy reserves were weighed and measured before being frozen.

Body condition indexes calculations

For the calculation of BCIs, the correction by structural body size can be performed in different ways, resulting in several body condition indexes to chose from. The most frequently used in the literature for vertebrates (Blem 1990; Brown 1996; Schulte-Hostedde et al. 2005) and invertebrates (Jakob et al., 1996; DeBano, 2008; Kasumovic et al., 2009; Dubois et al., 2010) is the residuals from an Ordinary Least Squares regression of log transformed mass in relation to log transformed body size (referred to as OLS_{resid} hereafter; Jakob et al., 1996; Hayes and Shonkwiler, 2001). Its use has been criticized because despite controlling for the lack of independence between mass and structural size, it fails to control for the lack of independence between structural size and mass (see Green 2001; Peig and Green 2009, 2010; and supplementary material). As a consequence the use of the Standardized Mass Index (SMI) has been suggested as a good alternative (Peig and Green 2009; 2010; see supplementary materials, Figure S1). Peig and Green (2009) used the Thope-Lleonar model of scaling to estimate body condition, **using the slope from an standardized major axis regression of mass and structural size as the scaling exponent**:

$$\bar{M}_i = M_i[T_0/T_i]^{bSMA}$$

where M_i and T_i are the mass and structural size (elytron length) of an individual respectively. T_0 is the population mean of the structural size, and bSMA is the slope of the

regression between the log of M_i and the log of T_i.

The benefit of Peig and Green's (2009) formula is that it retains the original mass

units. This results in a correction of our non-fixed morphological measure (i.e. mass) for

structural size (i.e. elytron length). We refer to size-corrected morphological measures as

"**scaled**", i.e. **Scaled Mass Index (SMI)**. Since standardized major axis regressions

standardise data, SMI can deal with Y and X variables measured on different scales. Its true

error variance assumptions are also more realistic than ordinary least squares regressions in

most cases (Warton et al., 2006; Green, 2001; see supplementary materials, Figure S1). We

compare first the validity of these two BCIs based on their ability to predict both the relative

amount of reserves present in our insect model system.

Quantification of the energy reserves

We used a dosage by colorimetry to assess the lipid content as well as the glycogen

and the circulating carbohydrates present in the abdomen of the beetles, by using a protocol

detailed in Rivero and Ferguson (2003). In brief, frozen beetles were dissected to remove

their prothorax, head, elytra and legs. The remaining abdomens were homogenized in

microcentrifuge tubes containing a 2% sodium sulfate solution and incubated in a 2:1 mix of

chloroform:methanol solution for 24 hours. They were then centrifuged, to separate the pellet

(containing glycogen) from the supernatant (containing fat and circulating sugars). The

supernatant was divided into 2 equal parts. One part we used to quantify lipids with vanillin-

phosphoric acid reagent. After a brief incubation, the optical density of the resulting solution

was read in a spectrophotometer (SpectraMax®, Molecular Devices) at 525 nm and

compared to a standard curve made of a serial dilution of sunflower seed oil. The other part

was used to quantify mono and disaccharides from the supernatant and the pellet (after

breaking down of the glycogen of the pellet), by incubating each of these fractions with

anthrone reagent. The optical density of the resulting solutions was read at 625 nm and

compared to a standard curve made of a serial dilution of glucose. As there was a strong correlation between circulating sugars and glycogen in the beetles (Pearson's correlation of log-transformed sugars and glycogen [95% CI], r = 0.811 [0.725, 0.872], Figure S2), we considered both compounds together in the results.

Mono- and disaccharides from the supernatant represent circulating sugars mostly in the form of trehalose, as it is the main sugar present in the hemolymph of insects and serves as a storage for the glucose used in locomotory activity (Candy, 1989; Thompson, 2003), whereas glycogen stored in the fat body helps replenishing consumed trehalose (Candy, 1989). Diacylglycerol is another fuel used in locomotory activity, which is stored in the fat body (Canavoso, 2001; Williams & Robertson, 2008).

Similarly to the procedure applied to the morphological measures, we corrected the absolute quantities of lipids and sugars for structural size (i.e. elytron length) using Peig and Green's (2009) formula. We refer to size-corrected body components as **scaled lipids** and **scaled sugars**.

A prerequisite to pooling data from different treatments when scaling metrics is that the relationship between log-transformed mass and log-transformed size does not change significantly according to our larval and adult feeding treatments (Peig & Green, 2009). This is the case in our dataset (standardized major axis regressions: larval feeding treatment: LRT = 2.558, *p-value* = 0.110; adult feeding treatment: LRT = 2.080, *p-value* = 0.149). Furthermore the relationship between all log-transformed energy reserves (lipids and sugars) and structural size also did not significantly differ between feeding treatments (standardized major axis regressions, all p-values > 0.05). Therefore we pooled all data together when calculating body condition indexes, scaled lipids and scaled sugars enabling comparisons between treatment groups.

Copulations and fecundity assessment

Dnervich et al. (2001) showed that the mating of *T. molitor* females every second day

prevented their sperm depletion. Our preliminary observations showed that females could lay eggs for a week following a single copulation event, and that when mated every second day they barely laid any eggs past 2 weeks after the first copulation. We thus designed our experiment in order to prevent any sperm depletion and capture most of the fitness of the females.

Ten days after emergence females were exposed to one male for 4 hours. During copulation, a male from the "rich larval" feeding treatment and a focal female were placed in an empty Petri dish containing only a piece of filter paper. We observed the pairs until we could record at least one copulation for each one of them, and we noticed that several events of copulation can take place in 4 hours. At the end of these 4 hours, the males were removed, put back together in a box and fed *ad libitum*, waiting to be reused 2 days later. Note that as males were not kept individually, they were randomly picked from their stock every second day, which reduces the chance that fecundity of the female would reflect the fitness of one male she would be paired with. Females on the other hand were individually placed in a new Petri dish supplemented with food according to their adult feeding treatment, and allowed to oviposit for 2 days until the next copulation event. The eggs of the previous Petri dish resulting from the previous reproduction event were sieved off of the flour and counted as a proxy for female fitness. There were in total 8 copulation sessions per female. Males that reached 17 days post emergence were removed from the experiment.

Statistical analyses

All analyses were carried out in the statistical software R (R core team, 2021). Unless stated otherwise, we used the information-theoretic approach to achieve model selection (Burnham and Anderson, 2002). The relative strength of support of all candidate models was assessed via Akaike's Information Criterion adjusted for small sample sizes (AICc) and AICc weights (ω), the adjusted R^2 as defined by the MuMIn R package (Bartoń, 2020). We also report the effect size Cohen's D for categorical variables and the partial r effect size for

continuous variables (Nakagawa and Cuthill, 2007) and estimate the 95% confidence intervals by bootstrap ($n = 10{,}000$) using the "boot" R package (Canty and Ripley, 2021).

We first set out to validate whether morphometric measures and BCIs were reliable indicators of energy reserves. To that effect, we first tested the repeatability of the morphometric measure elytron length using the R package "rptR" and the strength of its correlation with mass (Stoffel and Nakagawa, 2017), both important preconditions of valid BCIs using these metrics (Peig and Green 2009; 2010). Second, we investigated Pearson correlations between measures of energy reserves and morphometric measures as well as with BCIs. Prior to correlations all morphometric measures and energetic reserves were log transformed.

Second, we aimed to test if absolute measures of energy reserves (using elytron length and body mass as proxies), relative energy reserves (using SMI as a proxy) or the combination of both were the best predictors of a fitness trait (i.e fecundity) in our insert model. With this aim in mind, we built five general linear models (GLM) with a negative binomial distribution, a log link function and with fecundity (i.e. number of eggs) as a response variable. Elytron length is fixed after emergence, it reflects body size independently of body condition, whilst SMI reflects body condition independently of structural size. We were able to include both elytron length and SMI in the same model because SMI and elytron were only weakly negatively correlated (Pearson's correlation [95% confidence intervals]; $r = -0.315$ [-0.429; -0.191]) and the variance inflation factor was below 2, suggesting low collinearity when including both terms in a single model (Zuur et al. 2009). This allowed us to disentangle the relative support for the Relative Efficiency Hypothesis and the Absolute Energy Demand Hypothesis. In contrast, because of collinearity between elytron length and mass both parameters were always assessed in separate models. Thus, the five different models assessed were: fecundity according to body mass, fecundity according to structural size (elytron length), fecundity according to body condition (SMI), fecundity according to both structural size and body condition and the null model.

Third, we predicted that food limitation at the larval and/or the adult stage will accentuate a potential cost to structural size and thus will change the relationship between fitness and body condition. We investigated if food restriction resulted in a different relationship between fecundity and elytron length/body mass/SMI using a GLM with a negative binomial distribution, a log link function and with fecundity (i.e. number of eggs) as a response variable. We build sixteen models with the explanatory variables elytron length, body mass, SMI and interactions between feeding treatment and elytron length/body mass/SMI. Once again, the effect of mass was assessed in separate models from elytron length and SMI due to collinearity.

Since selection and/or environmental plasticity at the larval stage can result in differences in morphology of adults at pupal emergence (Teuschl, Reim & Blanckenhorn, 2006), we tested whether larval feeding treatment affected morphology and condition at emergence from pupation. We therefore assessed differences in mean elytron size, mass and SMI between larval treatments, after checking for homogeneity of variance using the Fligner-Killeen test. We then assessed mean differences between larval treatment groups using linear models for elytron length and mass and a general least square (GLS) model for SMI to control for significant heterogeneity (see results).

Results

1. Validation of structural size and body condition indexes as measure of absolute and relative energy reserves respectively

Our measures of elytron length were significantly repeatable ($p < 0.05$) with an R [95%CI] of 0.921 [0.902, 0.936] and were strongly correlated with mass (Pearson's correlation [95% confidence intervals]; $r = 0.776$ [0.729; 0.816]). Our results confirm that elytron length is an ideal indicator of structural size in our insect given its high repeatability

in measurement and its strong correlation with mass. Moreover, its size is fixed after hatching

from the pupal stage, since the cuticle gets tanned and sclerotized after emergence (i.e. it does

not vary during the life of the insect)

Both elytron length and mass were highly correlated with absolute values of lipids

and sugars (**Table 1**). In contrast, mass correlated poorly with the amounts of lipids and

sugars that were scaled for structural size (**Table 1**). OLS_{resid} and SMI on the other hand

significantly correlated with quantities of lipids and sugars which were scaled for body size

(**Table 1**). However, importantly, in contrast with SMI, OLS_{resid} additionally correlated

significantly with absolute values of lipids and sugars (**Table 1**), consistent with the fact that

OLS_{resid} is correlated with structural body size (see supplementary material, Figure S1, for

more details). Since BCIs have been introduced within the context of the Absolute Energy

Demand Hypothesis, they should not be expected to correlate with the absolute amount of

reserves, but with the amount of reserves relative to structural size instead (DeBano, 2008;

Peig & Green 2009; 2010). Thus our results confirm that OLS_{resid} is a biased index of body

condition since it does not control adequately for the codependence between structural size

and mass, whereas SMI does. We therefore excluded OLS_{resid} from all subsequent analyses,

and focus on elytron length, mass and SMI.

Table 1: Pearson's correlation between log-transformed components of body condition and log-transformed body measures or body condition indices.

	log(Lipids)	log(Glucose+glycogen)	log(Scaled lipids)	log(Scaled glucose+glycogen)
a. Body measures				
log(Elytron length)	*r = 0.612* (0.462; 0.727)	*r = 0.703* (0.580; 0.795)	*N/A*	*N/A*
log(Mass)	*r = 0.685* (0.556; 0.782)	*r = 0.740* (0.628; 0.821)	*r = -0.144* (-0.342; 0.066)	*r = -0.093* (-0.295; 0.118)
b. Body condition indices				
OLS residuals (OLS$_{resid}$)	*r = 0.374* (0.180; 0.541)	*r = 0.350* (0.153; 0.520)	*r = 0.326* (0.126; 0.500)	*r = 0.341* (0.143; 0.513)
Scaled mass index (SMI)	*r = 0.169* (-0.041; 0.364)	*r = 0.097* (-0.113; 0.299)	*r = 0.452* (0.269; 0.603)	*r = 0.424* (0.236; 0.581)

2. The relevance of structural size and body condition as predictors of fitness in an insect model

In this section, we assessed five different models of female fecundity: fecundity according to body mass, fecundity according to structural size (elytron length), fecundity according to body condition (SMI), fecundity according to both structural size and body condition and the null model. The latter was the model which was the best supported (**Table 1**; Δ**AICc** = 2.066; ω = 0.622; R^2 = 0.055), with both predictors positively associated with fecundity (**Figure 1**; elytron length: partial-*r* [95%CI] = 0.156 [0.020; 0.277]; SMI: partial-*r* [95%CI] = 0.223 [0.109; 0.326]). Among the other models, when predictors were assessed individually, models with body mass or SMI had stronger support than the null model (**Table 2**; **Figure 1**; body mass: R^2 = 0.037; partial-*r* [95%CI] = 0.197 [0.075; 0.315] and SMI: R^2 = 0.033; partial-*r* [95%CI] = 0.183 [0.071; 0.291]) but interestingly this was not the case for elytron length (**Table 2**; **Figure 2**; R^2 = 0.007; partial-*r* [95%CI] = 0.086 [-0.044; 0.212]). Therefore the effect of body size on fecundity was only apparent and statistically supported when accounting for body condition (**Figure 1**), and the effect of body condition was stronger and better supported when accounting for body size.

Table 2: Five candidate models of the effect of elytron length, body mass and SMI on fecundity according to a general linear model (GLM) with negative binomial distribution. Model rank, the model structure, model degrees of freedom (df), model log-likelihood (LogLik), model Aikaike information criterion for small sizes (AICc), AICc weight (ω) and model R^2 are shown.

Model Rank	Model	df	LogLik	AICc	ΔAICc	ω	R^2
1	*Fecundity ~ Elytron length + SMI*	4	-1133.662	2275.509	0.000	0.622	0.055
2	*Fecundity ~ body mass*	3	-1135.732	2277.575	2.066	0.221	0.037
3	*Fecundity ~ SMI*	3	-1136.192	2278.494	2.985	0.140	0.033
4	*Null model*	2	-1139.856	2283.767	8.258	0.010	0.000
5	*Fecundity ~ Elytron length*	3	-1139.106	2284.322	8.813	0.008	0.007

Figure 1: Effect size of the effect of mass (model 2; Table 2), elytron length (model 5; Table 2), elytron length when controlling for SMI (model 1; Table 2), SMI (model 3; Table 2) and SMI controlling for elytron length (model 1; Table 2) on total fecundity.

3. Is the cost of size on fitness dependent on resource availability during the larval and adult stages?

In this next section, we investigated whether the effect of food limitation at the larval and/or adult stage changes the relationship between fecundity and elytron length/mass/body condition. We expected that food limitation at the larval and/or the adult stage would accentuate a potential cost to structural size and we therefore expected statistical support for the effect of an interaction between body condition and feeding treatment on fecundity. Feeding treatment was an important predictor of fecundity (ΔAICc = 23.588; Table 3; Figure 2) and, as predicted, the effect of an interaction between food treatment and SMI was supported by model selection (ΔAICc = 3.107; Table 3; Figure 3). None of the other interactions were supported by model selection (ΔAICc < 2; Table 3). Therefore, the effect of SMI on female fecundity was dependent of the feeding treatment, but not mass or elytron length.

Contrary to our expectations, the restriction of food at the larval stage (in "poor larval-poor adult" and "poor larval-rich adult") did not increase the effect of SMI on fecundity. Instead, we found that the positive effect of SMI on fecundity was large and at its strongest within the rich larval and poor adult feeding groups ("rich larval-poor adult": partial *r* [95%CI] = 0.476 [0.268; 0.632]), whilst the effect was small and not statistically supported in the remaining feeding treatments (in decreasing order: "poor larval-rich adult": partial *r* [95%CI] = 0.206 [-0.114; 0.490]; "rich larval-rich adult": partial *r* [95%CI] = 0.148 [-0.125; 0.367]; "poor larval-poor adult": partial *r* [95%CI] = 0.126 [-0.104; 0.366]). This result can best be explained by investigating the variation in elytron length, mass and SMI within larval feeding treatment groups. Indeed, individuals originating from the poor larval feeding treatment emerged as adults with larger structural size (Cohen's d = 0.354 [0.125-0.583]; **Figure 3a.**) and heavier mass (Cohen's d = 0.287 [0.061-0.514]; **Figure 3b.**) on average than individuals originating from the rich larval feeding treatment. In contrast, adults originating

from both treatments have a similar body condition (SMI: Cohen's d = 0.100 [-0.018; 0.416]; **Figure 3c.**). However, the variability in SMI differed between the larval feeding treatments, adults originating from the poor larval feeding treatment showing less dispersion in the SMI scores than adults originating from the rich larval feeding treatment (Fligner-Killeen test: χ^2 = 9.712; *p-value* = 0.002; **Figure 3c.**). We do not observe this heterogeneity between larval feeding treatments on either structural size (elytron length) or mass (**Figure 3a. and b.**). It therefore appears that individuals form the poor larval food treatment emerged larger and more individuals were in peak body condition (lower variation in SMI around the mean) compared to the rich larval treatment. This reduced variation in SMI explains the weak effect of SMI on fecundity within adults originating from the poor larval feeding treatment (**Figure 3c. and e.**).

The fact that the effect of SMI was stronger in rich larval-poor adult feeding treatment compared to rich larval-rich adult feeding treatment is in line with our prediction that food restriction would accentuate the effect of body condition on fecundity. Further evidence of the stressful effect of food restriction on our animals comes from the fact that we also observed a higher mortality in females in the rich larval and poor adult feeding groups, whilst there was no mortality in the poor larval and rich adult feeding group (ΔAICc = 14.308; mortality: Rich larval/Poor adult = 18%; Rich larval/Rich adult = 7%; Poor larval/Poor adult = 2%; Poor larval/Rich adult = 0%). We found no support of effects of SMI, mass, elytron length or interactions on mortality (ΔAICc < 2; Table S1; Figure S3).

Table 3: Sixteen candidate models of the effect of elytron length, body mass, SMI and interactions between feeding treatment and elytron length/body mass/SMI on fecundity according to a general linear model (GLM) with negative binomial distribution. The effect of mass was assessed separately from elytron length and SMI due to collinearity. Models with interaction terms included the corresponding additive terms, but for simplicity the shortcut formula is presented (i.e. a model with y ~ x + z + x:z is presented as y ~ x * z). Model rank, the model structure, model degrees of freedom (df), model log-likelihood (LogLik), model aikaike information criterion for small sizes (AICc), AICc weight (ω) and model coefficient of determination (R^2) are shown.

Model Rank	Model	df	LogLik	AICc	ΔAICc	ω	R^2
1	Fecundity ~ feeding treatment * SMI + elytron length	10	-1115.436	2251.920	0.000	0.615	0.198
2	Fecundity ~ feeding treatment * elytron length + feeding treatment * SMI	13	-1113.635	2255.028	3.107	0.130	0.211
3	Fecundity ~ feeding treatment + elytron length + SMI	7	-1120.437	2255.401	3.480	0.108	0.161
4	Fecundity ~ feeding treatment * elytron length + SMI	10	-1117.702	2256.453	4.532	0.064	0.182
5	Fecundity ~ feeding treatment * SMI	9	-1119.304	2257.460	5.540	0.039	0.170
6	Fecundity ~ feeding treatment + mass	6	-1122.749	2257.892	5.971	0.031	0.143
7	Fecundity ~ feeding treatment + SMI	6	-1124.001	2260.395	8.474	0.009	0.134
8	Fecundity ~feeding treatment * mass	9	-1121.511	2261.876	9.955	0.004	0.153
9	Fecundity ~ feeding treatment + elytron length	6	-1126.790	2265.973	14.053	0.001	0.112
10	Fecundity ~ feeding treatment	5	-1128.006	2266.290	14.370	<0.001	0.102
11	Fecundity ~ feeding treatment * elytron length	9	-1124.490	2267.833	15.912	<0.001	0.130
12	Fecundity ~ elytron length + SMI	4	-1133.662	2275.509	23.588	<0.001	0.055
13	Fecundity ~ mass	3	-1135.732	2277.575	25.655	<0.001	0.037
14	Fecundity ~ SMI	3	-1136.192	2278.494	26.574	<0.001	0.033
15	Null model	2	-1139.856	2283.767	31.846	<0.001	0.000
16	Fecundity ~ elytron length	3	-1139.106	2284.322	32.402	<0.001	0.007

Figure 2: Boxplot of total fecundity according to larval and adult feeding treatments. Each

data represents a value from a female. Also shown are p-values from a Tukey's honest

significance test comparing the effect of different levels feeding treatments on fecundity

(controlling for elytron length and SMI) according to model 3 in Table 3 (the most

parsimonious model without interaction terms). Feeding restriction had a significant effect on

fecundity at the adult stage but not at the larval stage.

Figure 3: Density curves elytron length (a.), mass (b.) and SMI (c.) according to the larval

feeding treatment. The mean and 95%CI intervals are shown as a dashed line along and

shaded ribbons. Also shown are the output from the Fligner-Killeen Test of Homogeneity of

Variances to test differences in variation between larval feeding treatment groups. Finally

differences in mean between larval treatment groups were tested using linear models (LM)

for mass (a.) and elytron length (b.) and whilst general lest squares (GLS) were used for SMI

to control for heterogeneity between larval feeding groups. The effect of mass (d.), elytron

length (e.), and SMI (f.) on fecundity according to feeding treatment are plotted with fitted

lines and 95% confidence intervals.

Discussion

Many invertebrate studies use mass, size and body condition interchangeably as a single proxy of energy reserves and predictor of fitness (Amarillo-Suárez et al., 2011; Yoder et al., 2010; Sturm, 2016). Moreover, in both vertebrate and invertebrate studies, body condition indexes are frequently "invalidated" on the basis that they correlate poorly with absolute measures of energy reserves (Schulte-Hostedde et al. 2001; Labocha, Schutz & Hayes 2014; Wilder et al., 2016; McGuire et al. 2018). With this study, we aimed at shedding light on whether the concept of body condition, and thus the use of BCIs, is relevant in insects, or if on the contrary, absolute measures of energy reserves are more relevant. Indeed, in insects there is extensive evidence of a benefit to a larger structural size, but no evidence of a direct cost. These costs or benefits would ultimately indicate the relative support for the Absolute Energy Demand or the Relative Efficiency Hypotheses in an organism in a given context.

From a methodological point of view, our results confirm that simple measurements of size and mass are a good measure of the absolute amount of lipids and sugars present in *T. molitor* females. Regarding BCIs on the other hand, the correlation between SMI and the absolute amount of reserves was weak and not statistically supported. The fact that a BCI does not correlate or only weakly to an absolute amount of reserves has been criticized by several authors (Labocha, Schutz & Hayes 2014; Wilder et al., 2016). It is important to note however that since BCIs have been introduced within the context of the Absolute Energy Demand Hypothesis, they are supposed to account for energy expenditure relative to body size. Thus, they are not expected to correlate with the absolute amount of reserves, but with the amount of reserves relative to structural size instead (DeBano, 2008; Peig & Green 2009; 2010). In this case, it is precisely the strong correlation between SMI and relative energy reserves and the lack of correlation with absolute energy reserves that validates BCI as a good indicator of body condition.

In vertebrates, prior to Peig and Green's (2019) study, OLS$_{resid}$ was the most

prevalent BCI in use but has since largely been displaced by SMI as a consequence of

OLS$_{resid}$ biases in estimating body condition (Peig & Green 2009; 2010). Our study expands

the findings of the biases of OLS$_{resid}$ to an insect model, where OLS$_{resid}$ do not appropriately

control for the dependence of body mass on body structural size, and consequently correlates

with both absolute and relative body reserves. Furthermore, SMI are likely to be a superior to

OLS$_{resid}$ in most systems since the dependence of mass on structural size will violate OLS

error assumptions in most scenarios (DeBano, 2008; Peig & Green 2009, 2010; and, see

supplementary material, Figure S1). Therefore, we converge with several other authors who

advise on BCIs which are based on standardised major axis regressions rather than ordinary

least squares in any ecological study (Kotiaho, 1999, Green, 2001; Peig & Green, 2009,

2010).

From an ecological point of view, models with both elytron length and body condition

were better supported than models with a simple measurement of mass in *T. molitor* females.

Furthermore, the effect of body size on fitness was only discernable when controlling for the

effects of body condition. Thus, our results show that after accounting for the cost of body

size maintenance (consistent with the Absolute Energy Demand Hypothesis), larger

individuals have a higher fitness (consistent with the Relative Efficiency Hypothesis),

contradicting previous observations in several insect species that structural size does not

represent a significant cost (Tammaru et al., 2002; Calvo & Molina, 2005; Honěk et al., 1993;

Wilder et al., 2016).

As predicted, the observed effect of the interaction between SMI and feeding

treatment indicates that the relative weight of the Relative Efficiency Hypothesis and

Absolute Energy Demand Hypothesis is dependent on environmental resource availability.

An examination of the effects of larval feeding treatments on the morphological parameters

reveals that adult females originating from the poor larval feeding treatment show a similar

average but a reduced variance in their body condition compared to females originating from the rich larval feeding treatment. This reduced variance is likely to be responsible for the weak relationship between SMI and fecundity in females which went through food restriction at the larval stage. In contrast, the higher variation in body condition in females that had unrestricted access to food at the larval stage resulted in a strong and positive relationship between body condition and fitness but only when environmental food access was limited at the adult stage.

Unlike body condition, structural size and mass show a divergence in their mean values between larval feeding treatments but a similar dispersion: females originating from the poor larval feeding treatment reach a larger structural size than their well-fed counterparts. Our experimental design does not allow us to conclude whether the reduced variance in body condition and increased body size in the poor larval treatments is the result of developmental plasticity or differential mortality. On the one hand, although we did not find a higher larval mortality between our feeding treatments, we did not monitor mortality during metamorphosis. As a very costly life-history event which lives insects vulnerable to predation, or, in our setting, cannibalism, we could expect it to be an important culling point in *T. molitor*, as well as numerous insect species (Ichikawa & Kurauchi, 2009; Lowe et al., 2021). On the other hand, larvae reared in the poor larval feeding treatment took approximately two months extra to reach the pupation stage compared to larvae reared in the rich larval feeding treatment, which indicates a certain degree of developmental plasticity. *T. molitor* is indeed known to be very plastic in its larval developmental time depending on access to food among other parameters (Cotton & St. George, 1929).

It has been shown in several insect species that restricted access to food at the larval stage can sometimes lead to a delayed maturation at a smaller adult body size (Day & Rowe, 2002, Dmitriew et al., 2009). We observed the contrary, that adult females emerging from the poor larval feeding treatment were slightly bigger (2 % difference in the elytron size) than

females emerging from the rich larval feeding treatment. It is therefore likely that this

reduced variance in body condition is the result of both developmental plasticity and

differential mortality prior to adult emergence. Moreover, we found higher adult mortality for

insects well fed at the larval stage but food restricted at the adult stage, independently of body

condition. Thus, selection and/or developmental plasticity which resulted from the restriction

of food at the larval stage resulted in insects which were better able to cope with food

restriction at the adult stage, regardless of their starting body condition. In contrast,

emergence at suboptimal body condition for females with unrestricted access to food at the

larval stage resulted in reduced fitness when food was restricted at the adult stage. Thus, the

potential adaptive benefit of emerging earlier in favourable environmental conditions comes

at a fitness cost for females with low body condition if the environment changes

unfavourably at the adult stage.

The reduced variance in SMI but divergence in mean structural size and body mass

indicates that the amount of resources relative to structural size, instead of the absolute

amount of reserves, is optimized prior to adult emergence in *T. molitor* females. This means

that there is a cost to structural size in our insect model, consistent with the Absolute Energy

Demand Hypothesis. At the same time, the larger structural body size in females originating

from the poor larval feeding treatment could be the result of a maximisation of the mass-

specific metabolic rate during the larval development under food restriction, which is

consistent with the Relative Efficiency Hypothesis. Considering that our insect model is a

flightless burrowing insect, we can expect the cost of structural size due to locomotion is

reduced compared to other insect species in which flight is an additional source of costs

(Dixon & Kindlmann, 1999; Williams & Robertson, 2008).

Therefore, we urge authors to take into account body condition in the form of BCIs in

insect ecological studies in order to highlight fitness effects which might have been hidden by

the use of structural size only or a measure which is not independent from it. From a more

applied perspective, our study might help insect breeders maximize nutrient content of the

insects, by maximizing their body condition, over the production of components of structural

size (such as the cuticle) which have a poor availability to the vertebrate digestive system. In

contrast, the production of insect by-products such as chitin, the main component of the

insect cuticle, and thus, of insect structural size, might be improved by maximizing structural

size over body condition.

To conclude, our study demonstrates that the concept of body condition, and thus

correcting the amount of energy reserves for structural body size in invertebrates, is more

relevant than using a simple measurement of mass as a proxy for the amount of reserves

available for the allocation to fitness traits (Knapp & Knappova, 2013; Kasumovic &

Andrade, 2006; Kasumovic et al., 2009; Moya Laraño et al., 2008). Our analyses show that

this does not stem from stronger support of the Absolute Energy Demand Hypothesis over the

Relative Efficiency Hypothesis in our insect model. Instead, the use of an appropriate BCI

allowed us to detect both a cost (since higher body condition predicts fecundity) and a benefit

to structural body size (since larger females have a higher fitness at a given body condition)

in our insect model. This confirms, as stated by (Reim et al. 2006), that both the Absolute

Energy Demand Hypothesis and the Relative Efficiency Hypothesis are not mutually

exclusive, and instead that the fitness of an individual will be the result of their combined

action. Applying a similar approach in more studies on various animal models might reveal

the relative importance of both these processes in different ecological contexts.

Acknowledgements: We thank Maria Texeira Brandao for the help with the

quantification of sugars and lipids. We thank Coralie Dal & Aude Balourdet for experimental

support. We thank Morgan David for initial discussion, and Jens Rolff for comments on the

manuscript. This work was funded by ANR-JCJC-0134 & ANR-14-CE02-0009 to YM, and

the DFG grant 5026 FOR InsectInfect.

Authors contributions: Conceived the study and designed the experiments: CZ, YM & MG. Conducted the experiments: CZ & MG. Analyzed the data: MG. Wrote the manuscript: CZ & MG with contributions from YM.

Data archiving statement: the data will be made available on Dryad upon acceptance for publication.

Conflicts of interests statement: The authors declare no conflicting interests.

References

Adámková, A., Mlček, J., Adámek, M., Borkovcová, M., Bednářová, M., Hlobilová, V., Ivana Knížková, Juríková, T. (2020) *Tenebrio molitor* (Coleoptera: Tenebrionidae)—Optimization of Rearing Conditions to Obtain Desired Nutritional Values, *Journal of Insect Science*, *20*(5), 1-10. https://doi.org/10.1093/jisesa/ieaa100

Adamski, Z., Bufo, S. A., Chowański, S., Falabella, P., Lubawy, J., Marciniak, P., Pacholska-Bogalska, J., Salvia, R., Scrano, L., Słocińska, M., Spochacz, M., Szymczak, M., Urbański, A., Walkowiak-Nowicka, K., & Rosiński, G. (2019). Beetles as Model Organisms in Physiological, Biomedical and Environmental Studies - A Review. *Frontiers in physiology*, *10*, 319. https://doi.org/10.3389/fphys.2019.00319

Anderson, J.F. (1974). Responses to starvation in the spiders *Lycosa lenta* (Hentz) and *Filistata hibernalis* (Hentz). *Ecology*, *55*,576-585.

Amarillo-Suárez, A.R., Stillwell, R.C., Fox, C.W. (2011). Natural selection on body size is mediated by multiple interacting factors: a comparison of beetle populations varying naturally and experimentally in body size. *Ecology and Evolution*, *1*, 1-14. https://doi.org/10.1002/ece3.1

Arrese, E.L., Soulages, J.L. (2010). Insect fat body: energy, metabolism, and regulation.
Annual Review of Entomology, 55, 207–225. https://doi.org/10.1146/annurev-ento-112408-
085356

Bargielowski, I., Kaufmann, C., Alphey, L., Reiter, P., Koella, J. (2012). Flight performance
and teneral energy reserves of two genetically-modified and one wild-type strain of the
yellow fever mosquito *Aedes aegypti*. *Vector Borne Zoonotic Diseases, 12,* 1053–1058.
https://doi.org/10.1089/vbz.2012.0994

Barnett, C.A., Suzuki, T.N., Sakaluk, S.K., Thompson, C.F. (2015). Mass-based condition
measures and their relationship with fitness: in what condition is condition? *Journal of*
Zoology, 296, 1–5.

Bar-On, Y.M., Philips, R., Milo, R. (2018). The biomass distribution on Earth. *Proceedings*
of   the   National   Academy   of   Sciences,   115(25),   6506-6511.
https://doi.org/10.1073/pnas.1711842115

Bartoń, K. (2020). MuMIn: Multi-Model Inference. R package version 1.43.17.
https://CRAN.R-project.org/package=MuMIn

Bilton, D.T., Ribera, I., Edward, A.Z. (2019). Water Beetles as Models in Ecology and
Evolution.   *Short   Annual   Review   of   Entomology,   64*(1),   359-377.
https://doi.org/10.1146/annurev-ento-011118-111829

Boggs, C.L. (2009). Understanding insect life histories and senescence through a resource
allocation   lens.   *Functional   Ecology,   23*,   27-37.   https://doi.org/10.1111/j.1365-
2435.2009.01527.x

Berger, D., Walters, R., Gotthard, K. (2008). What limits insect fecundity? Body size- and
temperature-dependent egg maturation and oviposition in a butterfly. *Functional Ecology,*
22(3), 523-529. http://doi.org/10.1111/j.1365-2435.2008.01392.x.

Berger, D., Olofsson, M., Friberg, M., Karlsson, B., Wiklund, C., & Gotthard, K. (2012).
Intraspecific variation in body size and the rate of reproduction in female insects — adaptive

allometry or biophysical constraint? *Journal of Animal Ecology*, *81*(6), 1244–1258.

http://www.jstor.org/stable/23353224

Blanckenhorn, W. (2000). The Evolution of Body Size: What Keeps Organisms Small? *The*

Quarterly Review of Biology, 75(4), 385-407.

Blem, C.R. (1990). Avian energy stores. In: Johnston RF (ed) Current ornithology, vol 7.

Plenums, New York, pp 59–114.

Brown M.E. (1996) Assessing Body Condition in Birds. In: Nolan V., Ketterson E.D. (eds)

Current Ornithology. Current Ornithology, vol 13. Springer, Boston, MA.

https://doi.org/10.1007/978-1-4615-5881-1_3

Burnham, K. P., & Anderson, D. R. (2002). A practical information-theoretic approach.

Model selection and multimodel inference, 2.

Calder, W. A. (1984). Size, function, and life history. - Harvard University Press, Cambridge,

MA.

Calvo, D., Molina, J. M. (2005). Fecundity–Body Size Relationship and Other Reproductive

Aspects of *Streblote panda* (Lepidoptera: Lasiocampidae), *Annals of the Entomological*

Society of America, 98(2), 191–196. https://doi.org/10.1603/0013-

8746(2005)098[0191:FSRAOR]2.0.CO;2

Canavoso, L. E., Jouni, Z. E., Karnas, K. J., Pennington, J. E., & Wells, M. A. (2001). Fat

metabolism in insects. *Annual review of nutrition*, *21*, 23–46.

https://doi.org/10.1146/annurev.nutr.21.1.23

Candy, D.J., Becker, A., Wegener, G. (1997) Coordination and integration of metabolism in

insect flight. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular

Biology, *117*(4), 497-512.

Canty, A., Ripley, B. (2021). boot: Bootstrap R (S-Plus) Functions. R package version 1.3-

27.

Chafino, S., Ureña, E., Casanova, J., Casacuberta, E., Franch-Marro, X., & Martín, D. (2019).
Upregulation of E93 gene expression acts as the trigger for metamorphosis independently of
the threshold size in the beetle *Tribolium castaneum*. *Cell reports, 27*(4), 1039–1049.e2.
https://doi.org/10.1016/j.celrep.2019.03.094

Connat, J.L., Delbecque, J.P.D., Glitho, I., Delachambre, J. (1991). The onset of
metamorphosis in *Tenebrio molitor* larvae (Insecta, Coleoptera) under grouped, isolated and
starved conditions. *Journal of Insect Physiology, 37*(9), 653-657, 659-662.
https://doi.org/10.1016/0022-1910(91)90042-X

Costa, S., Pedro, S., Lourenço, H., Batista, I., Teixeira, B., Bandarra, N. M., Murta, D.,
Nunes, R., Pires, C. (2020). Evaluation of Tenebrio molitor larvae as an alternative food
source. *NFS journal, 21*, 57-64. https://doi.org/10.1016/j.nfs.2020.10.001

Cotton R.T., St. George, R.A. (1929) The mealworms. [U.S. Department of Agriculture.
Technical Bulletin. no. 95.]U.S. Government Printing Office, 38 pages.

Day, T., & Rowe, L. (2002). Developmental thresholds and the evolution of reaction norms
for age and size at life-history transitions. *The American naturalist, 159*(4), 338–350.
https://doi.org/10.1086/338989

DeBano, S. (2008). Morphometric condition as a measure of energetic content and
reproductive potential in *Dactylotum variegatum*. *Journal of Orthoptera Research, 17*, 293-
300. https://doi.org/10.1665/1082-6467-17.2.293

Dixon, A.F.G, Kindlmann, P. (1999). Cost of Flight Apparatus and Optimum Body Size of
Aphid Migrants. *Ecology, 80*(5), 1678–1690. https://doi.org/10.2307/176556

Dmitriew, C., Carroll, J., Rowe, L. (2009). Effects of early growth conditions on body
composition, allometry, and survival in the ladybird beetle Harmonia axyridis. *Canadian*
Journal of Zoology. **87**(2): 175-182. https://doi.org/10.1139/Z09-001

Dnervich, J.M., Papker, R.S., Rauser, C.L., Rutowski, R.L. (2001). Material benefits from
multiple mating in female mealworm beetles (*Tenebrio molitor* L.) *14*, 215–230.

Dubois, G.F., Le Gouar, P.J., Delettre, Y.R., Brustel, H., Vernon, P. (2010). Sex-biased and
body condition dependent dispersal capacity in the endangered saproxylic beetle *Osmoderma*
eremita (Coleoptera: Cetoniidae). *Journal of Insect Conservation, 14,* 679–687.
https://doi.org/10.1007/s10841-010-9296-0

Evenden, M.L., Whitehouse, C.M., Sykes, J. (2014). Factors influencing flight capacity of the
mountain pine beetle (Coleoptera: Curculionidae: Scolytinae). *Environmental Entomology,*
43, 187-96. https://doi.org/10.1603/EN13244

Fligner, M.A., Killeen, T.J. (1976). Distribution-Free Two-Sample Tests for Scale. *Journal*
of the American Statistical Association, 71(353), 210-213.

Funk, D.J., Filchak, K.E. & Feder, J.L. (2002) Herbivorous Insects: Model Systems for the
Comparative Study of Speciation Ecology. *Genetica, 116,* 251–267.
https://doi.org/10.1023/A:1021236510453

Gligorescu, A., Fischer, C. H., Larsen, P. F., Nørgaard, J. V., & Heckman, L.-H. L. (2020).
Production and Optimization of *Hermetia illucens* (L.) Larvae Reared on Food Waste and
Utilized as Feed Ingredient. *Sustainability, 12*(23), 9864. https://doi.org/10.3390/su12239864

Green, A.J. (2001). Mass/length residuals: measures of body condition or generators of
spurious results? *Ecology, 85,* 1473–1483. https://doi.org/10.1890/0012-
9658(2001)082[1473:MLRMOB]2.0.CO;2

Hill, G. E. (2011). Condition-dependent traits as signals of the functionality of vital cellular
processes. *Ecology letters, 14*(7), 625–634. https://doi.org/10.1111/j.1461-0248.2011.01622.x

Hayes, J.P., Shonkwiler, J.S. (2001) Morphometric indicators of body condition: worthwhile
or wishful thinking? In: Spearman JR (ed) Body composition analysis of animals: a handbook
of non-destructive methods. Cambridge University Press, Cambridge, pp 8–38

Honěk, A. (1993). Intraspecific variation in body size and fecundity in insects: a general
relationship. *Oikos, 66,* 483-492. https://doi.org/10.2307/3544943

Ichikawa, T., & Kurauchi, T. (2009). Larval cannibalism and pupal defense against
cannibalism in two species of tenebrionid beetles. *Zoological science, 26*(8), 525–529.
https://doi.org/10.2108/zsj.26.525

Jakob, E.M., Marshall, S.D., Uetz, G.W. (1996). Estimating fitness: A comparison of body
condition indices. *Oikos, 77*, 61-67. https://doi.org/10.2307/3545585

Kasumovic, M. M., & Andrade, M. C. (2006). Male development tracks rapidly shifting
sexual versus natural selection pressures. *Current Biology, 16*(7), R242-R243.
https://doi.org/10.1016/j.cub.2006.03.017

Kasumovic, M.M., Brooks R.C., Andrade, M.C.B. (2009). Body condition but not dietary
restriction prolongs lifespan in a semelparous capital breeder. *Biology Letters,* 5636–638.
http://doi.org/ 10.1098/rsbl.2009.0335

Keil, T. A., & Steinbrecht, R. A. (2010). Insects as model systems in cell biology. *Methods in*
cell biology, 96, 363–394. https://doi.org/10.1016/S0091-679X(10)96016-0

Kleiber, M. (1932). Body size and metabolism. *Hilgardia, 6,* 315–353.
https://doi.org/10.3733/hilg.v06n11p315

Kim, T. K., Yong, H. I., Kim, Y. B., Kim, H. W., & Choi, Y. S. (2019). Edible Insects as a
Protein Source: A Review of Public Perception, Processing Technology, and Research
Trends. *Food science of animal resources, 39*(4), 521–540.
https://doi.org/10.5851/kosfa.2019.e53

Knapp, M., Knappová, J. (2013). Measurement of body condition in a common carabid
beetle, *Poecilus cupreus*: a comparison of fresh weight, dry weight, and fat content. *Journal*
of insect science, 13, 6. https://doi.org/10.1673/031.013.0601

Kotiaho, J.S. (1999). Estimating fitness, comparison of body condition indices revisited.
Oikos, 87, 399–400.

Labocha, M.K., Schutz, H. & Hayes, J.P. (2014). Which body condition index is best? *Oikos,*
123, 111–119. https://doi.org/10.1111/j.1600-0706.2013.00755.x

Locke, M. (1991). Insect epidermal cells. In Physiology of the Insect Epidermis (ed. K.

Binnington and A. Retnakaran), pp. 1-22. Melbourne: CRISCO Publ.

Lowe, W.H., Martin, T.E., Skelly, D.K., Woods, H.A. (2021). Metamorphosis in an Era of

Increasing Climate Variability. *Trends in Ecology & Evolution, 36*(4), 360-375.

https://doi.org/10.1016/j.tree.2020.11.012

McGuire, L.P., Kelly, L.A., Baloun, D.E., Boyle, A.W., Cheng, T.L., Clerc, J., Fuller, N.W.,

Gerson, A.R., Jonasson, K.A., Rogers, E.J., Sommers, A.S, Guglielmo, C.G. (2018).

Common condition indices are no more effective than body mass for estimating fat stores in

insectivorous bats. *Journal of Mammalogy, 99*(5), 1065–1071.

https://doi.org/10.1093/jmammal/gyy103

Moya-Laraño, J., Macías-Ordóñez, R., Blanckenhorn, W. U., & Fernández-Montraveta, C.

(2008). Analysing body condition: mass, volume or density? *Journal of Animal Ecology,*

77(6) 1099-1108. https://doi.org/10.1111/j.1365-2656.2008.01433.x

Morales-Ramos, J. A., Kelstrup, H. C., Rojas, M. G., & Emery, V. (2019). Body Mass

Increase Induced by Eight Years of Artificial Selection in the Yellow Mealworm (Coleoptera:

Tenebrionidae) and Life History Trade-offs. *Journal of Insect Science, 19*(2), 4.

https://doi.org/10.1093/jisesa/iey110

Nandy, B., Joshi, A., Ali, Z.S., Sen, S., Prasad, N.G. (2012). Degree of adpative male mate

choice is positively correlated with female quality variance. *Scientific Reports,* 2, 447.

https://doi.org/10.1038/srep00447

Nakagawa, S., Cuthill, I. C. (2007). Effect size, confidence interval and statistical

significance: a practical guide for biologists. *Biological reviews, 82*(4), 591-605.

https://doi.org/10.1111/j.1469-185x.2007.00027.x

Peig, J. and Green, A. J. (2009). New perspectives for estimating body condition from

mass/length data: the scaled mass index as an alternative method. *Oikos, 118,* 1883-1891.

https://doi.org/10.1111/j.1600-0706.2009.17643.x

Peig, J., Green, A.J. (2010). The paradigm of body condition: a critical reappraisal of current
methods based on mass and length. *Functional Ecology, 24,* 1323–1332.
https://doi.org/10.1111/j.1365-2435.2010.01751.x

R Core Team (2021). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reid, M.L., Purcell, J.R.C. (2011) Condition-dependent tolerance of monoterpenes in an
insect herbivore. *Arthropod-Plant Interactions, 5,* 331-337. https://doi.org/10.1007/s11829-
011-9137-4

Reim, C., Kaufmann, C., Blanckenhorn, W.U. (2009). Size-dependent energetics of
metamorphosis in the yellow dung fly, *Scathophaga stercoraria. Evolutionary Ecology*
Research, 11, 1111-1130. http://dx.doi.org/10.5167/uzh-30455

Reim, C., Teuschl, Y., and Blanckenhorn, W.U. (2006) Size-dependent effects of temperature
and food stress on energy reserves and starvation resistance in yellow dung flies.
Evolutionary Ecology Research, 8, 1215–1234. https://doi.org/10.1111/j.1365-
3032.2007.00589.x

Rivero, A., & Ferguson, H. M. (2003). The energetic budget of Anopheles stephensi infected
with Plasmodium chabaudi: is energy depletion a mechanism for virulence?. *Proceedings.*
Biological sciences, 270(1522), 1365–1371. https://doi.org/10.1098/rspb.2003.2389

Rolff, J., Joop, G. (2002) Estimating condition: pitfalls of using weight as a fitness correlate.
Evolutionary Ecology Research, 4, 931-935.

Rowe, L., Houle, D. (1996). The lek paradox and the capture of genetic variance by condition
dependent traits. *Proceedings of the Royal Society B: Biological Sciences, 263,* 1415-1421.
https://doi.org/10.1098/rspb.1996.0207

Saleh, N.W., Larson, E.L. & Harrison, R.G. (2014). Reproductive Success and Body Size in
the Cricket *Gryllus firmus* . *Journal of Insect Behavior, 27,* 346–356.
https://doi.org/10.1007/s10905-013-9425-1

Schneider, D.S., & Chambers, M.C. (2008). Rogue Insect Immunity. Science, 322, 1199 –
1200. https://doi.org/10.1126/science.1167450

Schmid-Hempel, R., Schmid-Hempel, P. (1998), Colony performance and
immunocompetence of a social insect, *Bombus terrestris*, in poor and variable environments.
Functional Ecology, 12, 22-30. https://doi.org/10.1046/j.1365-2435.1998.00153.x

Scully, L.R., Bidochka, M.J., (2006). Developing insect models for the study of current and
emerging human pathogens. *FEMS Microbiology Letters*, *263*(1), 1–9.
https://doi.org/10.1111/j.1574-6968.2006.00388.x

Simpson, S.J., Raubenheimer, D. (2012). The nature of nutrition: A unifying framework from
animal adaptation to human obesity. Princeton University Press, Princeton, NJ.

Schulte-Hostedde, A.I., Millar, J.S., Hickling, G.J. (2001). Evaluating body condition in
small mammals. *Canadian Journal of Zoology*. *79*(6), 1021-1029.
https://doi.org/10.1139/z01-073

Schulte-Hostedde, A.I., Zinner, B., Millar, J.S., Hickling, G.J. (2005). Restitution of mass-
size residuals: validating body condition indices. *Ecology*, *86*, 155-163.
https://doi.org/10.1890/04-0232

Sinclair, B. J. (2015). Linking energetics and overwintering in temperate insects. *Journal of*
Thermal Biology, *54*, 5-11. https://doi.org/10.1016/j.jtherbio.2014.07.007

Stevenson, R.D, Woods, W.A.Jr (2006). Condition indices for conservation: new uses for
evolving tools. *Integrative and Comparative Biology*, *46*, (6), 1169–1190.
https://doi.org/10.1093/icb/icl052

Stoffel, M.A., Nakagawa, S. and Schielzeth, H. (2017). rptR: repeatability estimation and
variance decomposition by generalized linear mixed effects models. *Methods in Ecology and*
Evolution, 8, 1639-1644. https://doi.org/10.1111/2041-210X.12797

Sturm, R. (2016). Relationship between body size and reproductive capacity in females of the
black field cricket (Orthoptera, Gryllidae). *Austrian Journal of Forest Science*, 48(2).

Tammaru, T., Esperk, T., Castellanos, I. (2002). No evidence for costs of being large in
females of *Orgyia* spp. (Lepidoptera, Lymantriidae): larger is always better. *Oecologia, 133,*
430–438. https://doi.org/10.1007/s00442-002-1057-7

Teuschl, Y., Reim, C. and Blanckenhorn, W.U. (2007). Correlated responses to artificial
body size selection in growth, development, phenotypic plasticity and juvenile viability in
yellow dung flies. *Journal of Evolutionary Biology,   20,   87-103.*
https://doi.org/10.1111/j.1420-9101.2006.01225.x

Thommen, A., Werner, S., Frank, O., Philipp, J., Knitterfeld, O., Quek, Y., Fahmy, K.,
Shevchenko, A., Friedrich, B.M., Jülicher, F., Rink, J.C. (2019). Body size-dependent energy
storage causes Kleiber's law scaling of the metabolic rate in planarians. *Elife,* 2019;8:e38187.
http://doi.org/10.7554/eLife.38187

Thompson, S.N. (2003). Trehalose – The Insect 'Blood' Sugar. *Advances in Insect*
Physiology, *31*, 205–285. http://doi.org/10.1016/S0065-2806(03)31004-5

Townsend, A.K., Clark, A.B., McGowan, K.J., Miller, A.D., Buckles, E.L. (2010). Condition,
innate immunity and disease mortality of inbred crows. *Proceedings of the Royal Society B:*
Biological Sciences, 277, 2875–2883. https://doi.org/10.1098/rspb.2010.0480

van Huis, A. (2021) Prospects of insects as food and feed. *Organic Agriculture, 11,* 301–308.
https://doi.org/10.1007/s13165-020-00290-7

Warner, D. A., Johnson, M. S., & Nagy, T. R. (2016). Validation of Body Condition Indices
and Quantitative Magnetic Resonance in Estimating Body Composition in a Small Lizard.
Journal of experimental zoology. Part A, Ecological genetics and physiology, *325*(9), 588–
597. https://doi.org/10.1002/jez.2053

Warton, D.I., Wright, I.J., Falster, D.S. and Westoby, M. (2006). Bivariate line-fitting
methods   for   allometry.   *Biological   Reviews,   81,   259-291.*
https://doi.org/10.1017/S1464793106007007

Whitman, D.W. (2008). The significance of body size in the Orthoptera: a review. *Journal of*
Orthoptera Research, 7(2): 117-134. https://doi.org/10.1665/1082-6467-17.2.117

Wilder, S.M., Raubenheimer, D., Simpson, S.J. (2016). Moving beyond body condition
indices as an estimate of fitness in ecological and evolutionary studies. *Functional Ecology*,
30, 108-115. https://doi.org/10.1111/1365-2435.12460
971
972 Wilson, A.J., Nussey, D.H. (2010). What is individual quality? An evolutionary perspective.
973 *Trends in Ecology and Evolution, 25,* 207–214. https://doi.org/10.1016/j.tree.2009.10.002

975 Williams, W. I., & Robertson, I. C. (2008). Using automated flight mills to manipulate fat
976 reserves in Douglas-fir beetles (Coleoptera: Curculionidae). *Environmental entomology*,
977 *37*(4), 850–856. https://doi.org/10.1603/0046-225x(2008)37[850:uafmtm]2.0.co;2
978
Yoder, J.A., Christensen, B.S., Keeney, G.D. (2010). Enhanced tolerance to water stress in
adults of the South India strain of the seed beetle, *Callosobruchus maculatus* (Coleoptera:
Bruchidae), as a product of large body size. *European Journal of Entomology*, *107*(2), 271-
275. http://doi.org/10.14411/eje.2010.035

Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects*
models and extensions in ecology with R. Springer Science & Business Media.