Steryl ester formation and accumulation in steroid-degrading bacteria

Johannes Holert[1], Kirstin Brown[1], Ameena Hashimi[1], Lindsay D. Eltis[1], and William W. Mohn[1]

[1]Department of Microbiology & Immunology, Life Sciences Institute, University of British Columbia, Vancouver, British Columbia, Canada

Running title: Bacterial steryl ester synthesis

Correspondence: William W. Mohn, Department of Microbiology & Immunology, Life Sciences Institute, University of British Columbia, 2350 Health Sciences Mall, Vancouver, BC, Canada, V6T 1Z3. Email: wmohn@mail.ubc.ca

ABSTRACT

Steryl esters (SEs) are important storage compounds in many eukaryotes and are often

prominent components of intracellular lipid droplets. Here we demonstrate that selected

Actino- and Proteobacteria growing on sterols are also able to synthesize SEs and to

sequester them in cytoplasmic lipid droplets. We found cholesteryl ester (CE) formation in

members of the actinobacterial genera *Rhodococcus*, *Mycobacterium*, and *Amycolatopsis* as

well as several members of the proteobacterial Cellvibrionales order. CEs maximally

accumulated under nitrogen-limiting conditions, suggesting that steryl ester formation plays a

crucial role for storing excess energy and carbon under adverse conditions. *Rhodococcus jostii*

RHA1 was able to synthesize phytosteryl and cholesteryl esters, the latter reaching up to 7%

of its cellular dry weight and 69% of its lipid droplets. Purified lipid droplets from RHA1

contained CEs, free cholesterol and triacylglycerols (TAGs). In addition, we found formation of

CEs in *Mycobacterium tuberculosis* when grown with cholesterol plus an additional fatty acid

substrate. This study provides the first evidence for steryl ester accumulation in free-living and

pathogenic bacteria and provides a basis for the application of non-pathogenic bacterial whole

cell systems in the biotechnological production of SEs for use in functional foods and

cosmetics.

IMPORTANCE

Oleaginous bacteria exhibit great potential for the production of high-value neutral lipids, such

as triacylglycerols and wax esters. This study describes the formation of steryl esters (SEs) as

neutral lipid storage compounds in sterol-degrading oleaginous bacteria, providing a basis for

biotechnological production of SEs using bacterial systems with potential applications in the

functional food, nutraceutical, and cosmetic industries. We found cholesteryl ester (CE)

formation in several sterol-degrading Actino- and Proteobacteria under nitrogen-limiting

conditions, suggesting an important role of this process in storing energy and carbon under

adverse conditions. In addition, *Mycobacterium tuberculosis* grown on cholesterol

accumulated CEs in the presence of an additional fatty acid substrate.

INTRODUCTION

Many bacteria synthesize and accumulate carbon and energy storage compounds under environmental stress conditions to increase their survival in environments with fluctuating nutrient conditions (1). A much smaller subset of bacteria are oleaginous and accumulate neutral lipids, namely triacylglycerols (TAGs) and wax esters (WEs), as a prominent class of storage compounds. Although TAG and WE synthesis is rarely described in other bacteria, their accumulation is widespread within the actinobacterial genera *Streptomyces*, *Nocardia*, *Rhodococcus*, *Mycobacterium*, *Dietzia*, and *Gordonia*, and has also been reported in the proteobacterial genera *Acinetobacter*, *Alcanivorax* and *Marinobacter* (2-7). TAGs and WEs are highly hydrophobic, non-polar lipids, which are formed by esterification of a glycerol molecule to three fatty acids or by esterification of a long chain fatty alcohol to a fatty acid, respectively (**Fig. 1**). Potential biotechnological applications of TAG and WE producing bacteria gained increasing attention over the last years due to the considerable industrial relevance of these compounds as biofuels and other products such as feed additives, cosmetics, and lubricants (8). Especially oleaginous *Rhodococcus* strains have been suggested as promising microbial cell factories for the large scale production of neutral lipids due to their ability to convert agroindustrial wastes into valuable lipids and their accessibility to cell engineering for increased lipid production (9).

Accumulation of neutral lipids in bacteria is typically induced by environmental stress conditions, in which a suitable carbon source is in excess while additional growth is restricted by a deficiency of other nutrients or environmental conditions. The most prominent example is nitrogen-limitation, which induces TAG and WE synthesis in a wide range of neutral lipid-accumulating Actino- and Proteobacteria. This has been most extensively studied in oleaginous *Rhodococcus* strains, which can accumulate up to 70% of their cellular dry weight as TAGs upon nitrogen depletion, depending on the available carbon source (10-12). In *Mycobacterium tuberculosis*, the causative agent of tuberculosis, dormancy inducing conditions such as nitric oxide (NO) stress or iron-limitation have been shown to initiate accumulation of TAGs and WEs (13, 14). These intracellular carbon and energy storage compounds have been suggested to be important for the pathogenicity of *M. tuberculosis* (15, 16).

TAGs and WEs are sequestered into cytoplasmic lipid droplets, which are multifunctional organelles encapsulated by a phospholipid monolayer and specific proteins (17). Analogous to oleaginous bacteria, many eukaryotes also sequester neutral lipids as carbon and electron donor reservoirs into cytoplasmic lipid droplets. In contrast to bacterial lipid droplets, many eukaryotic lipid droplets contain sterols in addition to TAGs and WEs (18). Sterols constitute a key class of lipids in most animals, fungi, and plants, where they play

essential roles as membrane constituents and precursors for vitamins and steroid hormones.
The dominant sterol in animals is cholesterol, while plants have β-sitosterol and other
phytosterols, and fungi have ergosterol (**Fig. 1**). Within eukaryotic lipid droplets, the sterol 3-
hydroxyl group is typically esterified with fatty acids forming so-called steryl esters (SEs) (**Fig.
1**), which act as a sterol storage pool. Synthesis and breakdown of SEs plays a vital role in
sterol homeostasis and imbalances between free and esterified cholesterol pools are related
to several human diseases, such as Wolman disease, cholesteryl ester storage disease,
atherosclerosis, and Alzheimer's disease (19). In contrast to eukaryotes, sterol synthesis is
rare in bacteria and mainly restricted to *Myxococcales* and *Methylococcales* (20). These
bacterial sterols have structural features that distinguish them from eukaryotic sterols (21).
Further, it is not known whether these bacteria are able to esterify and store their sterols in
lipid droplets.

Sterols are abundant growth substrates for bacteria in the environment, and sterol
degradation has been characterized in many actinobacterial isolates and is highly conserved
in several genera of the Corynebacterineae suborder, namely the genera *Amycolicicoccus,*
Dietzia, Gordonia, Mycobacterium, Nocardia, Rhodococcus, and *Tsukamurella* (22). In
addition, we recently isolated several sterol-degrading strains belonging to the
gammaproteobacterial order Cellvibrionales (23), which are also able to degrade sterols such
as cholesterol. Under aerobic conditions, sterols are degraded by a conserved pathway into
central metabolic moieties such as acetyl-CoA and propionyl-CoA (24), which are
subsequently channeled into central metabolic pathways for energy conservation and
production of biomass. Among those sterol-degrading bacteria are many that are also well
known for their ability to produce TAGs and WEs as neutral lipid storage compounds, such as
the Actinobacteria, *Rhodococcus jostii* RHA1 (RHA1) (25) and *Mycobacterium tuberculosis*
(*Mtb*) (26). Key enzymes in bacterial TAG and WE synthesis are CoA-dependent, bifunctional
WS/DGAT enzymes encoded by *atf* genes. These enzymes catalyze both the acetylation of
DAGs to TAGs and of fatty alcohols to WEs (**Fig. 1**) (27). Most WS/DGATs are promiscuous
enzymes accepting a variety of acyl-CoAs and alcohols as substrates (5, 28). Interestingly, the
WS/DGAT enzyme AtfA from the oleaginous, non-sterol degrading Proteobacterium
Acinetobacter calcoaceticus ADP1 has been shown to have SE-synthase activity when
heterologously expressed in *E. coli* or the yeast *Saccharomyces cerevisiae* (29). Over 50 years
ago, two studies reported that different *Mycobacterium* strains produced cholesteryl fatty acid
esters when growning with cholesterol (30) and that *Mycobacterium smegmatis* also produced
esters of other sterol substrates (31). However, these studies did not provide further insight
into the triggering mechanism or cellular localization of SE accumulation and no further reports
have been published on bacterial steryl ester synthesis since then.

Here we investigate the ability of sterol-degrading bacteria to synthesize lipid storage
compounds from sterol substrates under adverse growth conditions and report formation of
SEs as lipid storage compounds in bacteria. This significantly extents the knowledge about
bacterial SE formation and provides a basis for biotechnological production of SEs using
bacterial systems with potential applications in the functional food, nutraceutical, and cosmetic
industries.
124

RESULTS

Accumulation of SEs in RHA1 under nitrogen-limiting conditions

To investigate formation of lipid-storage compounds from sterol substrates, *Rhodococcus jostii* RHA1 was grown in mineral medium with 1 mM cholesterol or 1 mM of a phytosterol mix, containing 76.4% β-sitosterol, 10.4% β-sitostanol, 7.3% campesterol and 1% campestanol, under nitrogen-limiting and nitrogen-excess conditions. In cholesterol-grown cultures, RHA1 reached a biomass in early stationary phase that was approximately four-fold higher under nitrogen-excess than under nitrogen-limiting conditions (**Fig. 2A**). In contrast to the N-excess cultures, the N-limited cultures had no residual ammonium but did have residual cholesterol. Similar results were obtained during growth on the phytosterol substrate mix (not shown). Thin layer chromatography (TLC) analysis of neutral lipid extracts revealed compound spots with the same R_f-value as an authentic cholesterol-palmitate standard (R_f = 0.78) in N-limited cholesterol and phytosterol cultures, but not in the respective N-excess cultures or in N-limited or N-excess cultures grown on non-sterol substrates (**Fig. 2B)**. In addition, compounds with similar Rf-values as an authentic tripalmitin TAG standard (R_f = 0.24) were present in extracts of all N-limited cultures. Time-dependent sampling of RHA1 grown on cholesterol under N-excess and N-limiting conditions revealed that increased formation of the presumptive cholesteryl esters coincided with the depletion of ammonium from the medium (**Fig. 3A**). Under N-excess conditions only minimal amounts of these compounds accumulated during exponential growth and stationary phase (**Fig. 3B**). When nitrogen was replenished in an N-limited, stationary-phase, cholesterol-grown culture, the accumulated lipids including the presumptive SEs disappeared within 72 h (**Fig. 2B**).

To confirm the formation of CEs, we analyzed native neutral lipid extracts from strain RHA1 grown with cholesterol or succinate under N-limiting conditions in early stationary phase using a gas chromatography (GC)-coupled flame ionization detector (FID). Comparison to a cholesteryl-palmitate (C16:0) and a cholesteryl-heptadecanoate standard (C17:0) confirmed that these CEs accumulated in cholesterol-grown RHA1 cells (**Fig. 4A**), but were absent in succinate-grown cells. In addition, five other compounds from cholesterol-grown cells eluted close to these CEs, which were absent in succinate-grown cells, suggesting that seven CE species, with different fatty acid chain lengths, are formed with cholesterol as substrate. Comparison to a tripalmitate ((C16:0)$_3$) TAG standard suggested the accumulation of up to nine different TAG species in strain RHA1 grown with cholesterol and eleven different TAG species in cells grown with succinate (**Fig. 4A**). Time-dependent sampling starting at the onset of nitrogen depletion from the media followed by GC-FID analysis showed that the CE:TAG ratio shifted from approximately 60% to 72% within the first 24 h of incubation and to 69% after 48 h.

To gain further insight into the nature of SEs accumulating in strain RHA1, neutral lipid

extracts of cholesterol- and phytosterol-grown cultures were fractionated on silica columns

producing five lipid fractions (**supplemental Fig. S1**). SEs eluted in fraction 2 and were well

separated from other lipids, like TAGs (Fractions 3 and 4) and free sterols (fractions 5). Aliquots

of fraction 2 were subjected to alkaline transesterification with methanol. GC-MS analysis

showed that free cholesterol or the free phytosterols (β-sitosterol, β-sitostanol, and

campesterol) were formed from transesterification, which were not present in untransesterified

extracts (**Fig. 5A-B**). Only the respective sterols provided as a growth substrate were detected

in this analysis. In addition, the lack of other free alcohols after transesterification confirmed

the absence of wax esters in neutral lipid extracts. In addition, a range of seven fatty acid

methyl esters (FAMEs) were detected after transesterification of cholesteryl esters, supporting

the results from the GC-FID analysis. Those were identified as methyl esters of C15:0, C16:1,

C16:0, C17:1, C17:0, C18:1 and C18:0 (**Fig. 5A**). The most abundant fatty acids were the odd-

numbered fatty acids, C17:0, C17:1, and C15:0, which constituted more than 75% of produced

FAMEs (**Fig. 5C**). GC-MS analysis of transesterified fraction 2 confirmed that no detectable

amounts of other free alcohols indicative of wax esters were present (not shown). Analysis of

all neutral lipid extracts using a GC-MS method for wax-ester analysis (32) showed that no

detectable amounts of wax esters were present in any sample (not shown), which is in

agreement with earlier reports that wax esters are only formed in minimal amounts in RHA1

(32).

This confirmed that the previously unidentified lipids accumulating in N-limited RHA1

cultures grown with cholesterol or phytosterols are fatty acid steryl esters. In the late stationary

phase (100 h) CEs accounted for 6.8 +/- 1.4% of the cellular dry weight in N-limited cultures

(n = 5 from two independent experiments) based on GC-MS quantification.

Localization of accumulated cholesterol and CEs in RHA1

To confirm that accumulating neutral lipids are localized intracellularly, we stained stationary

phase, cholesterol-grown, washed, N-excess, and N-limited RHA1 cells with the lipophilic

fluorescence stain Nile Red. Microscopic analysis showed that stained lipids accumulated as

lipid droplets in the cell interior (**supplemental Fig. S2**) of N-limited cells but were absent from

N-excess cells. In addition, we incubated RHA1 with fluorescently labelled cholesterol

(BODIPY-cholesterol) and sodium palmitate for 48 h without a nitrogen source. Microscopic

analysis showed that fluorescently labelled cholesterol was mainly present within lipid droplets

inside the cells (**supplemental Fig. S3**) but mostly absent from cell membranes. TLC analysis

of neutral lipid extracts of cells with fluorescently labelled cholesterol showed that no CEs were

formed (not shown), which is most likely because the altered chemical structure of the labelled

cholesterol side chain inhibits esterification. These results confirm that cholesterol and CEs
are mainly stored in intracellular lipid droplets.

To gain further insight into the location of CEs within RHA1 cells, lipid droplets were
isolated from RHA1 cultures grown with cholesterol or sodium palmitate or a mixture of both
substrates under N-limiting conditions. With all three substrate combinations, gradient
centrifugation of total cell extracts produced visible bands of lipid droplet on top of the sucrose
gradient (**supplemental Fig. S4**). TLC analysis revealed that CEs and free cholesterol were
present in neutral lipid extracts of lipid droplets derived from cells grown on cholesterol but not
in lipid droplets derived from cells grown on palmitate alone, while TAGs were present under
all three conditions (**Fig. 6A**). This was confirmed by fractionation and subsequent
transesterification of neutral lipid extracts (not shown). In addition, the lack of other free
alcohols after transesterification confirmed the absence of wax esters in lipid droplets. In two
independent experiments 52% and 58% of lipids in lipid droplets derived from cells grown on
cholesterol were identified as CEs, 31% and 15% as TAGs and 17% and 27% as free
cholesterol. Similar results were obtained for lipid droplets derived from cells grown on
cholesterol plus palmitate, with 50% and 69% CEs, 35% and 23% TAGs and 15% and 8%
cholesterol, while TAGs were the only neutral lipids in lipid droplets derived from cells grown
on palmitate (**Fig. 6B**). In lipid droplets from cells grown on cholesterol, between 60% to 70%
of fatty acids in CEs and TAGs were odd-numbered (C15 and C17), with C17:0 and C17:1
making up around 90% of these fatty acids (**Fig. 6C**). In lipid droplets from cells grown with
cholesterol plus palmitate, 23% to 55% of fatty acids were odd-numbered. TAGs of cells grown
on palmitate contained almost exclusively (~95%) even-numbered fatty acids. (**Fig. 6C**).

Formation of SEs in other sterol-degrading bacteria
To test whether steryl ester formation also occurs in other sterol-degrading bacteria, we
extracted neutral lipids from the Actinobacteria, *Mycobacterium smegmatis* strain mc²155,
Mycobacterium sp. strain BC8-1, *Rhodococcus opacus* strain Chol-4, *Rhodococcus ruber*
strain Chol-4, and *Amycolatopsis* sp. strain ATCC 39116, as well as the
Gammaproteobacteria, *Zhongshania* sp. strain SB11-1A, *Haliaceae* sp. strain BC5-1, and the
BD1-7 clade bacterium strain SB11-3 grown on cholesterol under nitrogen-limiting and
nitrogen-excess conditions. The latter strains were isolated in our lab (23) and are among the
first Proteobacteria reported to be able to degrade sterols. TLC analysis indicated these strains
produced SEs in N-limited cultures in the stationary phase in differing amounts, with highest
yields in the *Rhodococcus*, *Mycobacterium*, and *Zhongshania* strains and lower yields in
Amycolatopsis, *Haliaceae*, and the BD1-7 clade strain (**supplemental Fig. S5A**). Subsequent
fractionation of neutral lipid extracts followed by transesterification and GS-MS analysis
confirmed the formation of CEs in all strains (**supplemental Fig. S5B**).

In addition, we tested steryl ester formation in *Mycobacterium tuberculosis* (*Mtb*)
Erdmann growing on cholesterol and on cholesterol plus sodium palmitate. Because *Mtb* did
not grow when sodium palmitate was added to the medium prior to inoculation we spiked
palmitate into cultures growing on cholesterol after 5 days. TLC analysis of neutral lipid extracts
of *Mtb* indicated the formation of small amounts of CEs in cells growing on cholesterol after 7
days and in cells growing on cholesterol plus palmitate after 12 and 13 days, when cells were
in stationary phase (**supplemental Fig. S6A-B**). Subsequent fractionation of neutral lipid
extracts followed by transesterification and GS-MS analysis confirmed the formation of CEs
(**supplemental Fig. S6C**). The only detectable FAME in this fraction was C16:0. Formation of
free hexacosanol after transesterification suggested that a C_{26}-OH wax ester was also formed.
In addition, we tested whether nitric oxide (NO) stress or iron-limitation, which are known
inducers of neutral lipid accumulation in *Mtb* (13, 14), increase CE formation from cholesterol
or cholesterol plus palmitate. For this, cholesterol-grown cell suspensions of *Mtb* were
incubated with cholesterol alone or cholesterol plus palmitate. To induce NO stress, two pulses
of 0.5 mM spermine-NONOate were added at the beginning of the experiment and again after
16 h of incubation and samples were taken after 16 h and after 2 days of incubation. Spermine
(0.5 mM) was added to control cultures. Deferoxamine (0.5 mM) was added as an iron chelator
and samples were taken after 2 and 4 days of incubation. Confirming the results from the initial
growth experiment, all extracts from *Mtb* cells incubated with cholesterol plus palmitate
contained CEs with C16:0 as only detectable FAME. However, addition of NO or deferoxamine
did not increase the amount of accumulating CEs (**supplemental Fig. S6D**). No significant
accumulation of CEs occurred in treated cells incubated with only cholesterol. Small amounts
of free cholesterol were detected after transesterification of extracts of cell suspensions treated
with NO and incubated for 2 days with cholesterol (not shown).

DISCUSSION

SEs are important sterol storage compounds in many eukaryotic organisms and are prominent components of some eukaryotic lipid droplets. Here we demonstrate that selected sterol-degrading Actino- and Proteobacteria are also able to synthesize SEs and to sequester them in cytoplasmic lipid droplets. We found steryl ester formation in all sterol-degrading strains investigated in this study, including members of the actinobacterial genera, *Rhodococcus*, *Mycobacterium*, and *Amycolatopsis,* as well as members of the proteobacterial Cellvibrionales order, suggesting that the ability to accumulate excess sterols as an electron donor and carbon source under adverse conditions plays a crucial role in sterol-degrading bacteria. This is supported by findings from the 1960s, which showed that different *Mycobacterium* strains produced steryl esters during growth with sterol substrates (30, 31). The ability of other oleaginous bacteria to synthesize SEs has been overlooked in other studies, which exclusively used non-steroidal substrates, such as glucose, gluconate, or benzoate, when investigating neutral lipid accumulation.

Using *Rhodococcus jostii* RHA1 as an oleaginous, sterol-degrading model organism, we found accumulation of minor amounts of CEs during exponential growth on cholesterol when ammonium was available, and a significant increase in CE and TAG accumulation, when ammonium was limiting, indicating that nitrogen-limitation stimulates SE formation. This extends previous reports that nitrogen-limitation is a major inducer for the accumulation of TAGs and WEs in RHA1 with non-steroidal substrates (12, 33). While we found significant accumulation of CEs in the two free-living *Mycobacterium* strains mc^{2}115 and BC8-1 under nitrogen-limiting conditions with cholesterol as the only substrate, the pathogenic strain *Mycobacterium tuberculosis* (*Mtb*) Erdmann did not grow on cholesterol in nitrogen-limited medium. However, small amounts of cholesterol-palmitate accumulated in this strain when growing on cholesterol plus palmitate as substrates without any stress conditions, demonstrating that pathogenic *Mtb* strains are able to accumulate CEs when a sterol and a fatty acid are available. During infection, *Mtb* triggers formation of foamy macrophages (34, 35), which provide a cholesterol and fatty acid-rich microenvironment, and *Mtb* metabolizes these host-derived lipids during infection and persistence (36-38). Interestingly, cholesterol and fatty acid uptake are co-regulated in *Mtb* (39), suggesting that *Mtb* has simultaneous access to both lipid classes *in vivo*. SE synthesis and accumulation could act as a buffer system for excess lipids in *Mtb* providing a way to regulate intracellular lipid metabolism. In addition, SEs could serve as a carbon and electron donor reservoir in *Mtb* under adverse conditions. In a similar manner, TAG accumulation and sequestration in cytoplasmic lipid droplets was shown to occur *in vivo* in pathogenic *Mtb* (16, 40) and was proposed to be important for persistence and reactivation (41). Accumulation of TAGs and WEs was shown to

be stimulated by various dormancy-inducing stress conditions, such as hypoxia, NO stress,
iron-limitation, and increased acidity, which induce expression of several WS/DGAT encoding
genes in *Mtb* (13, 42-44). Although we did not detect increased formation of CEs in *Mtb* under
NO stress or iron-limiting conditions, a combination of stress factors as described in (42) might
provide deeper insight into the regulation of SE synthesis in *Mtb*.

Lipid droplets have been identified in most domains of life, including many eukaryotes,
as well as archaeal and bacterial prokaryotes (45). While the general structure of lipid droplets
is conserved in all these organisms, comprising a neutral lipid core coated by a phospholipid
monolayer and various peripheral proteins, the neutral lipid composition varies significantly
between organisms (17, 46). Our results show that CEs, in addition to TAGs and free
cholesterol, are sequestered in intracellular lipid droplets in RHA1. The lipid droplet
composition in RHA1 grown on cholesterol is very similar to that of lipid droplets found in
selected mammalian lipid droplet-producing cell lines (chinese hamster ovaries CHO K2, rat
kidney normal NKR and immortalized human fibroblasts SV589) (47), with around 50% CEs,
30% TAGs and 20% free cholesterol. The sequestration of cholesterol in lipid droplets is in
agreement with an earlier report showing that fluorescently labeled cholesterol predominantly
localizes in lipid droplets in mycobacteria (48).

Interestingly, the majority of fatty acids stored in TAGs and CEs in cholesterol-grown
RHA1 cells were odd-numbered fatty acids, with C17:0 and C17:1 as main fatty acids, which
is in stark contrast to the mostly even-numbered fatty acids typically found in eukaryotic neutral
lipids. Complete metabolism of one cholesterol molecule through the canonical bacterial 9,10-
seco steroid-degradation pathway is proposed to yield four propionyl-CoAs, four acetyl-CoAs,
one pyruvate and one succinyl-CoA (24). The preponderance of odd-chain fatty acids in neutral
lipids accumulating in RHA1 with cholesterol as carbon source suggests that propionyl-CoA
was a significant building block for neutral-lipid fatty acids. Elevated ratios of odd-numbered to
even-numbered fatty acids have also been found in TAGs accumulating in RHA1 growing with
benzoate under nitrogen-limiting conditions, where propionyl-CoA was presumably generated
through the methylmalonyl-CoA pathway from succinate (12). Cholesterol-derived propionyl-
CoA has been shown to be toxic to *Mtb* if not metabolized (49), and incorporation of propionyl-
CoA into cell wall lipids (50) and TAGs (51) has been suggested as a mechanism to detoxify
propionate in *Mtb*. Cholesterol-degrading bacteria, such as RHA1 and *Mtb*, might encounter
increased cellular propionyl-CoA levels under adverse growth conditions and the formation of
odd-numbered fatty acid CEs might provide an additional sink for this toxic end-product.
However, we did not detect odd-numbered fatty acid CEs in *Mtb*, which accumulated CEs in
significant amounts only in the presence of an additional fatty acid substrate under the tested
conditions.

Sterols are abundant growth substrates for bacteria in the environment and sterol
degradation is conserved in many actinobacterial genera, particularly in members of the
Corynebacterineae (22). Recently, we have also identified sterol-degradation in some isolates
of the oligotrophic marine Gammaproteobacteria (OMG) group belonging to the Cellvibrionales
order (23). Steroid-degrading bacteria are globally distributed and actinobacterial ones are
particularly abundant within soil and rhizosphere as well as in deep ocean habitats. Our results
indicate that SE formation is conserved in sterol-degrading bacteria, suggesting that this trait
might play important roles in these environments. Supporting this hypothesis, most neutral-
lipid accumulating Actinobacteria isolated from soil samples (52) belong to taxa in which the
ability to degrade sterols is conserved (22) and are thus likely to produce SEs. *De novo*
synthesis of sterols is rare in bacteria, being largely restricted to *Myxococcales* and
Methylococcales (20). Further, it is not known whether these bacteria are able to store these
sterols as esters.

We searched the genomes of strain RHA1 and of *Mtb* Erdmann for homologs of one
prokaryotic and fifty eukaryotic proteins with characterized steryl ester synthase activity
available in the Swiss-Prot database, including phosphatidylcholine-sterol *O*-acyltransferases
(EC 2.3.1.43), diacylglycerol *O*-acyltransferases (EC 2.3.1.20), sterol *O*-acyltransferase (EC
2.3.1.26), and phospholipid:diacylglycerol acyltransferases (EC 2.3.1.158). None of these
proteins had any significant similarity (accepted E-values < 0.01) to RHA1 or *Mtb* proteins.
Based on the multifunctionality and substrate promiscuity of TAG and WE producing
WS/DGAT enzymes, it is feasible that these enzymes might carry out an additional function as
steryl ester synthases. Out of the 16 predicted WS/DGATs in RHA1, Atf6 (WP_011594556.1)
and Atf8 (WP_011597548.1) have the highest amino acid sequence identity (40% and 39%,
respectively) to AtfA from the oleaginous, non-sterol degrading Proteobacterium *Acinetobacter*
calcoaceticus ADP1, which was shown to have SE-synthase activity when heterologously
expressed in *E. coli* or *S. cerevisiae* (29). TAG and WE synthase activities have been
confirmed for Atf8 and Atf6, respectively (12, 28) as well as for an Atf6 ortholog in *Rhodococcus*
opacus PD630 (53), and deletions of the respective genes significantly decreased neutral lipid
accumulation. In addition, transcription of *atf8* was strongly induced upon nitrogen depletion in
RHA1 growing on benzoate, exhibiting the highest transcript levels of all 16 *atf* genes under
nitrogen-limiting conditions (12). In contrast, *atf6* transcripts were most abundant under
nitrogen-excess conditions with no apparent transcriptional induction upon nitrogen depletion.
WS/DGAT genes identified in soil communities were exclusively from Actinomycetales (54),
which includes the steroid-degrading suborder *Corynebacterineae*. In contrast, WS/DGAT
genes from marine sediments were exclusively of gammaproteobacterial origin, predominantly
from members of the oligotrophic marine Gammaproteobacteria (OMG) group, which have
been identified as sterol-degrading (22, 23). BLASTp analysis (max E-value 10^{-15}) of AtfA from

Acinetobacter calcoaceticus ADP1 against the proteomes of the cholesteryl ester
accumulating OMG bacteria *Zhongshania* sp. strain SB11-1A, *Haliaceae* sp. strain BC5-1 and
the BD1-7 clade bacterium SB11-3 suggested that these strains encode between four and six
WS/DGAT paralogs, further supporting our hypothesis that these proteins might function as
steryl ester synthases.

Our results indicate that accumulated SEs can be reactivated and degraded in RHA1
upon relief of growth-limiting conditions, suggesting that SEs might play a vital role as a carbon
and energy reservoir in these environments where fluctuating nutrient availability is common.
We searched the genomes of strain RHA1 and of *Mtb* Erdmann for homologs of three
prokaryotic and fifteen eukaryotic proteins with characterized or predicted sterol esterase
activity available in the TrEMBL and Swiss-Prot databases (EC 3.1.1.13). However, none of
the query proteins had significant similarity (accepted coverage > 85%, percent identity > 30%,
E-values < 0.01) to RHA1 or *Mtb* proteins. Strain RHA1 encodes up to 54 putative
lipases/esterase enzymes (33), which were suggested to contribute to neutral lipid degradation,
but no distinct degradation system for endogenous TAGs or WEs has been identified so far.
In contrast, the lipase family protein LipY in *M. tuberculosis* has been shown to have TAG
degradation activity (55).

Interestingly, several animal lineages such as insects and soil-dwelling nematodes
have lost the ability to synthesize sterols *de novo* (56) and thus rely on uptake of sterols with
their diet (57). It is tempting to speculate that sterol and steryl ester accumulating bacteria in
oligotrophic habitats such as soils or freshwater systems might constitute an important source
for sterols for bacteriovorous, sterol-auxotrophic eukaryotes such as *Caenorhabditis elegans*
or *Daphnia magna*.

RHA1 was also able to form esters of the phytosterols β-sitosterol, β-sitostanol, and
campesterol under nitrogen-limiting conditions. In recent years, plant-derived phytosterols and
their esters were recognized as potential agents to lower serum cholesterol levels in
hypercholestemic patients (58, 59) by inhibiting absorption of endogenous and dietary
cholesterol in the intestine. Today phytosterols and phytosteryl esters are approved in many
countries as safe food additives, and several so-called functional foods are marketed
worldwide, which are supplemented with phytosterol and their esters (60). Industrial production
of phytosteryl esters is mainly accomplished by chemical esterification or transesterification of
plant-derived sterols, which requires high temperatures and the use of strong acids or bases
(61). Our study provides a basis for biotechnological production of phytosteryl esters using
non-pathogenic whole-cell bacterial systems, thus eliminating the requirement for elevated
temperature and acid or base catalysts. Similar to RHA1, many other sterol-degrading
Actinobacteria are also able to use phytosterols as growth substrates, providing an excellent
screening resource for efficient phytosteryl ester accumulation.

 MATERIAL AND METHODS

Bacteria and culture conditions

Rhodococcus jostii strain RHA1, as well as *Mycobacterium smegmatis* strain mc^{2}155,

Rhodococcus opacus strain PD630, *Rhodococcus ruber* strain Chol-4, and *Amycolatopsis* sp.

strain ATCC 39116 were cultivated in M9 minimal medium supplemented with trace elements

and vitamin B1 at 30 °C. The marine bacteria *Mycobacterium* strain BC8-1, *Zhongshania* sp.

strain SB11-1A, *Haliaceae* sp. strain BC5-1, and the BD1-7 clade bacterium strain SB11-3

were cultivated in artificial marine medium at 21 °C as described earlier (23). Substrates were

added at the following concentrations: cholesterol 1 mM, sodium palmitate 1.0 mM or 1.7 mM,

succinate 7.0 mM, glucose 5 mM. Starter cultures were grown on 10 mM succinate or 1 mM

cholesterol. Cholesterol and other sterol substrates were added as solids and 1% (w/v) methyl-

β-cyclodextrin was added to the medium, when required, prior to autoclaving. Substrate

removal was monitored by gas chromatography-coupled mass spectrometry (GC-MS) of

organic extracts of acidified culture supernatants as described earlier (62). Nitrogen-limiting

conditions were achieved by reducing the concentration of ammonium chloride in the medium

from 18.70 mM to 0.93 mM. Under N-excess conditions the ratio of carbon to nitrogen in the

growth medium was around 1.5:1, while it was 29:1 under N-limiting conditions. To limit

nitrogen transfer between cultures, starter cultures were washed twice with phosphate-

buffered saline (PBS) without nitrogen. Growth was measured as protein, using the

bicinchoninic acid assay (Thermo Scientific) after hot alkaline cell lysis.

Mycobacterium tuberculosis strain Erdmann (*Mtb*) was grown in 7H9 medium

supplemented with 0.5% tyloxapol and 0.5 mM cholesterol at 37 °C. If required 0.5 mM sodium

palmitate were spiked into growing cultures after 5 days. Growth was measured as optical

density at 600 nm. To test effects of nitric oxide (NO) stress and iron-limitation, cholesterol-

grown *Mtb* was harvested and washed in 7H9 medium without carbon source, and cells were

suspended in the same medium containing 0.5 mM cholesterol and/or 0.5 mM palmitate. 0.5

mM of the NO releasing agent spermine-NONOate was added from a 10 mM stock in 0.01 M

NaOH at the beginning of the experiment and again after 16 h of incubation. Spermine was

added at the same concentration to control cultures. The iron chelating agent deferoxamine

was added at 0.5 mM from a 10 mM stock at the beginning of the experiment.

Biosample IDs and references to cholesterol degradation capability of all strains used

in this study are summarized in the **supplemental Tab. S1**.

Ammonium quantification

Ammonium concentrations were quantified using indophenol blue. Culture supernatant

samples were diluted 25- to 500-fold in distilled water and 200 μl aliquots were distributed into

96-well plates in triplicates. 8 μl of reagent (10% w/v phenol in ethanol), 8 μl catalyst (0.5% w/v

nitroprusside in water) and 20 µl oxidizing solution (20% w/v trisodium citrate and 1% w/v
sodium hydroxide solution, mixed with 5% sodium hypochlorite 4:1 v/v) were added and plates
were incubated at 37 °C for 1-2 h in the dark. Absorbance was measured at 630 nm.
Ammonium concentrations were calculated using a standard curve with 0-2 mg ml^{-1} ammonium
chloride. Alternatively, semi-quantitative ammonium test strips (Quantofix®, Macherey-Nagel,
Germany) were used for instant ammonium quantification.

Lipid extraction and analysis

For lipid extraction, cells were harvested, washed twice with distilled water and frozen at -80 °C.
Mtb cells were autoclaved before freezing. Cell pellets were lyophilized and their dry weight
was determined by weighing the pellets. The pellets were then suspended in 1 ml distilled
water and broken by bead beating (four rounds of 60 sec at 4.5, 5.0, 5.5, and 6.0 speed setting,
respectively, in a FastPrep-24 bead beater; MP Biomedicals, Solon, OH). Between rounds
samples were incubated on ice for 5 min. Neutral lipids were extracted with chloroform-
methanol (2:1, v/v, 1% acetic acid), dried under nitrogen and suspended in chloroform with 1%
acetic acid. If required cholestane was added as internal standard from a 10 mM stock solution
in chloroform prior to organic extraction for normalization and quantification of cholesterol and
FAMEs. Aliquots of neutral lipid extracts were analyzed by thin layer chromatography (TLC)
on 60 F$_{254}$ silica gel plates (Merck) using hexane/diethyl-ether/acetic-acid (90:8:1, v/v/v) as a
solvent system. Cholesterol palmitate, cholesterol, tripalmitin, and stearyl-stearate dissolved
in chloroform were used as standards. The plates were dried, dabbed with 10% cupric sulfate
in 8% phosphoric acid solution and developed at 200 °C for 5-10 min. Native neutral lipid
extracts from non-lyophilized cell pellets of strain RHA1 grown under nitrogen-limiting
conditions with cholesterol or succinate were also analyzed using a GC-coupled flame
ionization detector (FID). For this, 20 ml culture aliquots were harvested, washed twice and
frozen at -80°C at different time points after nitrogen was depleted from the medium. Neutral
lipids were extracted with tert-butyl ether:methanol (10:3, v/v), followed by two extractions with
hexane. Combined organic extracts were dried under nitrogen and suspended in chloroform.
Neutral lipid extracts were analyzed by gas chromatography on an Agilent 8890 series GC
system equipped with a J&W Select Biodiesel GC Column (15 m, 0.32 mm, 0.10 µm), coupled
to a flame ionization detector (FID). GC-injector and detector temperatures were set to 350°C,
and 400°C, respectively. 1 µl sample volumes were injected with an initial purge flow rate of
50 ml min^{-1}, which subsequently decreased to 20 ml min^{-1} after 2 min, and a septum purge
flow rate of 3 ml min^{-1}. The helium flow rate was 8.0972 ml min^{-1}. The oven was set to an initial
temperature of 40°C for 2 min with an equilibration time of 0.5 min. The temperature increased
by 25°C min^{-1} and was held at 170°C for 2 min, then increased by 15°C min^{-1} to 210°C, then
by 5°C min^{-1} to 250°C and held for 5 min. The temperature finally increased by 10°C min^{-1} to
400°C and held for 2 min. The FID gas flow settings were set to 400 ml min^{-1} air flow, 30 ml

min^{-1} H$_2$ fuel flow and 25 ml min^{-1} makeup flow (N$_2$). Cholesteryl palmitate, cholesteryl

heptadecanoate, and tripalmitin TAG dissolved in chloroform were used as standards. CE and

TAG peaks were normalized with the internal standard cholestane for quantification.

Isolation of lipid droplets

Lipid droplets were isolated as previously described (63, 64). Briefly, RHA1 cultures were

harvested by centrifugation (4,000 x g for 10 min), washed three times with 40 ml Buffer A (25

mM tricine, 250 mM sucrose, pH 7.8) and suspended in 8 ml buffer A. After incubation on ice

for 20 min, cells were disrupted by passing them three times through a cooled French pressure

cell at 100 MPa. Cell debris was removed by centrifugation at 4,000 × g for 10 min.

Supernatants were loaded into SW40 tubes (Beckman), layered with 2 ml Buffer B (20 mM

HEPES, 100 mM KCl, 2 mM MgCl2, pH 7.4), and centrifuged at 180,000 x g for 1 h at 4 °C.

The lipid droplet fraction on top of the sucrose gradient was transferred to a 1.5 ml Eppendorf

tube and was washed twice with 200 µl Buffer B and finally suspended in 1 ml buffer B. No

pellets were formed during washing of the lipid droplets suggesting that lipid droplets were not

contaminated by membrane particles. Total lipids were extracted with 5 ml chloroform-acetone

(1:1, v/v). The organic phase was dried under nitrogen and lipids were suspended in 1 ml

chloroform with 1% acetic acid. Aliquots were analyzed by TLC as described above.

Lipid fractionation

For analyzing the fatty acid profile of cellular lipids and lipid droplets, lipid extracts were

fractionated stepwise. Total lipid extracts derived from lipid droplets were first run over a silica

column equilibrated with chloroform and neutral lipid fractions were obtained by eluting with 6

column volumes chloroform with 1% acetic acid. Glycolipid fractions were subsequently eluted

with 6 column volumes acetone (not analyzed further) followed by phospholipid elution with

7.5 column volumes methanol. Neutral lipids from lipid droplets and cellular lipid extractions

were further fractionated by silica column chromatography. Neutral lipid samples were dried,

resuspended in hexane and poured on a silica column equilibrated with the same solvent. 5

different fractions were obtained by eluting sequentially with 5 column volumes hexane, 6

column volumes hexane/diethyl ether (99:1, v/v), 5 column volumes hexane/diethyl ether (95:5,

v/v), 5 column volumes hexane/diethyl ether (92:8, v/v), and 6 column volumes chloroform with

1% acetic acid. Fractions were dried under nitrogen and lipids were suspended in 1 ml

chloroform with 1% acetic acid. If required, aliquots were analyzed by TLC as described above.

Alkaline transesterification with methanol and lipid quantification

Dried cellular lipid extracts or individual lipid fractions were subjected to alkaline

transesterification with methanol modified after (65). For this, dried lipids were resuspended in

methanol/toluene (1:1, v/v), mixed with an equal volume of methanolic KOH and incubated for

1 h at 50 °C. Phospholipid fractions were incubated for 15 min at 30 °C. Free sterols and fatty

acid methyl-esters were extracted twice with 4 volumes hexane/chloroform (4:1, v/v) from 0.15

volumes acetic acid (1 M) and 1 volume water. The organic phase was dried under nitrogen

and lipids were suspended in 1 ml chloroform with 1% acetic acid. Aliquots were analyzed by

GC-MS as described earlier for FAMEs and free cholesterol (62) and for wax esters (32).

Cholesterol and fatty acid methyl-esters (FAMEs) were identified by comparison to authentic

standards as well as matching against the NIST database (National Institute of Standards and

Technology, v11). To quantify total neutral lipids, unfractioned neutral lipid extracts were

subjected to transesterification and FAMEs were normalized on cholestane. To quantify SEs

in cellular lipid extracts, fractions 1 (containing the internal standard cholestane) and 2

(containing steryl esters) were combined prior to transesterification. This allowed normalisation

and quantification of free cholesterol derived from transesterification. The percentage of CEs

per extracted cellular dry weight was estimated by calculating the total weight of CEs from CE

containing fractions and dividing it by the dry weight of the respective extracted cell pellet. For

this, the cholesterol concentration in transesterified fractions was calculated using a

cholesterol-cholestane standard curve. Subsequently, this was transformed into the total

weight of CEs using the molecular masses of all detectable CEs weighted on the peak area

ratio of the respective detected FAMEs (C15:0, C16:1, C16:0, C17:1, C17:0, C18:1, C18:0).

To estimate ratios of neutral lipid classes in lipid droplets, cholestane and methyl-

nonadecanoate were added as internal standards to each fraction prior to transesterification

from 10 mM stock solutions in chloroform. Free cholesterol (in fraction 5) and cholesterol

released from transesterification of fraction 2 were normalized on cholestane and FAMEs

derived from transesterification of CEs and TAGs were normalized on methyl-nonadecanoate

before calculating their molar ratios.

Fluorescence microscopy

To localize accumulating neutral lipids in strain RHA1, we added the fluorescent dye Nile Red

to growing cholesterol cultures under N-limiting or N-excess conditions without cyclodextrin.

Nile red was added to the medium after autoclaving at a final concentration of 0.5 µg ml^{-1} from

a sterile stock solution in DMSO (0.3 mg ml^{-1}). Nitrogen and cholesterol depletion were

confirmed as described above. Cells from both cultures were examined four days after

reaching stationary phase. Cells were washed twice and resuspended in sterile PBS before

microscopy. Slides were examined on a Zeiss Axio Imager M1 fluorescence microscope

equipped with a Zeiss HBO 103 W/2 high-pressure mercury lamp and an AxioCam MRm

camera in phase contrast, differential interference contrast or fluorescence mode.

Fluorescence images were shot using excitation and emission wavelengths of 550 nm and

605 nm, respectively. All Images were processed using the Zeiss AxioVision software (40 V

4.6.3.0).

To further analyze the localization of accumulating cholesterol within cells of RHA1

under nitrogen-limiting conditions, we performed a cell suspension experiment with

fluorescently labeled cholesterol (cholesterol-BODIPY) as substrate under N-limiting
conditions followed by laser scanning confocal microscopy. For that, cholesterol-grown cells
were washed two times with PBS without nitrogen. Depletion of cholesterol in the starter
culture was confirmed by organic extraction and GC-MS analysis. Cells were suspended in
1/5 of the original volume of M9 medium without nitrogen containing 0.5 mM palmitate, 0.5 mM
fluorescently labeled cholesterol (cholesterol-BODIPY) and 1% (w/v) methyl-β-cyclodextrin
and incubated for 48 h at 200 rpm and 30 °C. Cells were washed twice with PBS without
nitrogen containing 0.02% (w/v) cholate and 0.02% (v/v) tylaxopol and suspended in PBS.
Cells were analyzed using a laser scanning confocal microscope (Olympus Fluoview FV1000-
Inverted). A small aliquot of cells was broken by ultrasonication (four rounds of 20 sec using a
Microsom cell disruptor XL set to 5; Mandel scientific Co. Ltd., Guelph, ON, Canada) and
neutral lipid extracts were analyzed by TLC. Undeveloped plates were analyzed using a GE
Typhoon trio + imager with excitation wavelength of 488 nm and an emission filter of 520 nm.
Pre-tests with unlabelled cholesterol and palmitate showed that CE accumulation in RHA1 cell
suspensions started within 22 h of incubation and continued at least until 95 h of incubation
(data not shown).
571

Acknowledgments

We thank Fanrui Meng and Carl Strittmatter for help with fluorescence microscopy. This study

was funded by a Discovery Grant from the Natural Sciences and Engineering Research

Council of Canada.

Data availability

Draft genomes of *Mycobacterium* strain BC8-1, *Zhongshania* sp. strain SB11-1A, *Haliaceae*

sp. strain BC5-1, and the BD1-7 clade bacterium strain SB11-3 have been deposited in the

European Nucleotide Archive (ENA) under the BioProject number PRJEB30766.

References

1. **J Brigham C**, **Kurosawa K**, **Rha C**, **Sinskey AJ**. 2011. Bacterial carbon storage to value added products. J Microbial Biochem Technol **s3**:1–13.

2. **Alvarez HM**, **Steinbüchel A**. 2002. Triacylglycerols in prokaryotic microorganisms. Appl Microbiol Biotechnol **60**:367–376.

3. **Russell NJ**, **Volkman JK**. 1980. The effect of growth temperature on wax ester composition in the psychrophilic bacterium *Micrococcus cryophilus* ATCC 15174. J Gen Microbiol **118**:131–141.

4. **Kalscheuer R**, **Stöveken T**, **Malkus U**, **Reichelt R**, **Golyshin PN**, **Sabirova JS**, **Ferrer M**, **Timmis KN**, **Steinbüchel A**. 2007. Analysis of storage lipid accumulation in *Alcanivorax borkumensis*: Evidence for alternative triacylglycerol biosynthesis routes in bacteria. J Bacteriol **189**:918–928.

5. **Kalscheuer R**, **Steinbüchel A**. 2003. A novel bifunctional wax ester synthase/acyl-CoA:diacylglycerol acyltransferase mediates wax ester and triacylglycerol biosynthesis in *Acinetobacter calcoaceticus* ADP1. J Biol Chem **278**:8075–8082.

6. **Garay LA**, **Boundy-Mills KL**, **German JB**. 2014. Accumulation of high-value lipids in single-cell microorganisms: a mechanistic approach and future perspectives. J Agric Food Chem **62**:2709–2727.

7. **Röttig A**, **Strittmatter CS**, **Schauer J**, **Hiessl S**, **Poehlein A**, **Daniel R**, **Steinbüchel A**. 2016. Role of wax ester synthase/acyl coenzyme a:diacylglycerol acyltransferase in oleaginous *Streptomyces* sp. strain G25. Appl Environ Microbiol **82**:5969–5981.

8. **Röttig A**, **Steinbüchel A**. 2013. Acyltransferases in Bacteria. Microbiol Mol Biol Rev **77**:277–321.

9. **Alvarez HM**, **Herrero OM**, **Silva RA**, **Hernández MA**, **Lanfranconi MP**, **Villalba MS**. 2019. Insights into the metabolism of oleaginous *Rhodococcus* spp. Appl Environ Microbiol **85**:8075–12.

10. **Alvarez HM**, **Mayer F**, **Fabritius D**, **Steinbüchel A**. 1996. Formation of intracytoplasmic lipid inclusions by *Rhodococcus opacus* strain PD630. Arch Microbiol **165**:377–386.

11. **Waltermann M**, **Luftmann H**, **Baumeister D**, **Kalscheuer R**, **Steinbüchel A**. 2000. *Rhodococcus opacus* strain PD630 as a new source of high-value single-cell oil? Isolation and characterization of triacylglycerols and other storage lipids. Microbiology **146** :1143–1149.

12. **Amara S**, **Seghezzi N**, **Otani H**, **Diaz-Salazar C**, **Liu J**, **Eltis LD**. 2016. Characterization of key triacylglycerol biosynthesis processes in rhodococci. Sci Rep **6**:24985.

13. **Daniel J**, **Deb C**, **Dubey VS**, **Sirakova TD**, **Abomoelak B**, **Morbidoni HR**, **Kolattukudy PE**. 2004. Induction of a novel class of diacylglycerol acyltransferases and triacylglycerol accumulation in *Mycobacterium tuberculosis* as it goes into a dormancy-like state in culture. J Bacteriol **186**:5017–5030.

14. **Bacon J**, **Dover LG**, **Hatch KA**, **Zhang Y**, **Gomes JM**, **Kendall S**, **Wernisch L**, **Stoker NG**, **Butcher PD**, **Besra GS**, **Marsh PD**. 2007. Lipid composition and

transcriptional response of *Mycobacterium tuberculosis* grown under iron-limitation in
continuous culture: identification of a novel wax ester. Microbiology **153**:1435–1444.

15. **Stehr M**, **Elamin AA**, **Singh M**. 2013. Lipid inclusions in mycobacterial infections.
Tuberculosis - Current Issues in Diagnosis and Management, Bassam H. Mahboub
and Mayank G. Vats, IntechOpen, doi: 10.5772/54526

16. **Garton NJ**, **Christensen H**, **Minnikin DE**, **Adegbola RA**, **Barer MR**. 2002.
Intracellular lipophilic inclusions of mycobacteria in vitro and in sputum. Microbiology
148:2951–2958.

17. **Zhang C**, **Liu P**. 2017. The lipid droplet: A conserved cellular organelle. Protein &
Cell **8**:796–800.

18. **Ruggles KV**, **Turkish A**, **Sturley SL**. 2013. Making, baking, and breaking: The
synthesis, storage, and hydrolysis of neutral lipids. Annu Rev Nutr **33**:413–451.

19. **Korber M**, **Klein I**, **Daum G**. 2017. Steryl ester synthesis, storage and hydrolysis: A
contribution to sterol homeostasis. Biochim Biophys Acta **1862**:1534–1545.

20. **Wei JH**, **Yin X**, **Welander PV**. 2016. Sterol synthesis in diverse bacteria. Front
Microbiol **7**:990.

21. **Lee AK**, **Banta AB**, **Wei JH**, **Kiemle DJ**, **Feng J**, **Giner J-L**, **Welander PV**. 2018. C-
4 sterol demethylation enzymes distinguish bacterial and eukaryotic sterol synthesis.
Proc Natl Acad Sci USA **115**:5884–5889.

22. **Bergstrand LH**, **Cardenas E**, **Holert J**, **Van Hamme JD**, **Mohn WW**. 2016.
Delineation of steroid-degrading microorganisms through comparative genomic
analysis. MBio **7**:e00166–16.

23. **Holert J**, **Cardenas E**, **Bergstrand LH**, **Zaikova E**, **Hahn AS**, **Hallam SJ**, **Mohn
WW**. 2018. Metagenomes reveal global distribution of bacterial steroid catabolism in
natural, engineered, and host environments. MBio **9**: e02345-17.

24. **Crowe AM**, **Casabon I**, **Brown KL**, **Liu J**, **Lian J**, **Rogalski JC**, **Hurst TE**, **Snieckus
V**, **Foster LJ**, **Eltis LD**. 2017. Catabolism of the last two steroid rings in
Mycobacterium tuberculosis and other Bbacteria. MBio **8**:e00321–17.

25. **van der Geize R**, **Yam K**, **Heuser T**, **Wilbrink MH**, **Hara H**, **Anderton MC**, **Sim E**,
Dijkhuizen L, **Davies JE**, **Mohn WW**, **Eltis LD**. 2007. A gene cluster encoding
cholesterol catabolism in a soil actinomycete provides insight into *Mycobacterium
tuberculosis* survival in macrophages. Proc Natl Acad Sci USA **104**:1947–1952.

26. **Wipperman MF**, **Sampson NS**, **Thomas ST**. 2014. Pathogen roid rage: cholesterol
utilization by *Mycobacterium tuberculosis*. Crit Rev Biochem Mol Biol **49**:269–293.

27. **Alvarez HM**. 2016. Triacylglycerol and wax ester-accumulating machinery in
prokaryotes. Biochimie **120**:28–39.

28. **Barney BM**, **Wahlen BD**, **Garner E**, **Wei J**, **Seefeldt LC**. 2012. Differences in
substrate specificities of five bacterial wax ester synthases. Appl Environ Microbiol
78:5734–5745.

29. **Kalscheuer R**, **Luftmann H**, **Steinbüchel A**. 2004. Synthesis of novel lipids in
Saccharomyces cerevisiae by heterologous expression of an unspecific bacterial
acyltransferase. Appl Environ Microbiol **70**:7119–7125.

30. **Schubert K**, **Kaufmann G**. 1965. Bildung von Sterinestern in der Bakterienzelle.
Biochim Biophys Acta **106**:592–597.

31. **Schubert K**, **Kaufmann G**, **Hörhold C**. 1969. Veresterung von Sterinen mit
Fettsäuren und mit Bernsteinsäure in *Mykobakterien*. Biochim Biophys Acta **176**:163–
169.

32. **Round J**, **Roccor R**, **Li S-N**, **Eltis LD**. 2017. A fatty acyl-CoA reductase promotes
wax ester accumulation in *Rhodococcus jostii* RHA1. Appl Environ Microbiol
83:e00902–17–12.

33. **Hernández MA**, **Mohn WW**, **Martínez E**, **Rost E**, **Alvarez AF**, **Alvarez HM**. 2008.
Biosynthesis of storage compounds by *Rhodococcus jostii* RHA1 and global
identification of genes involved in their metabolism. BMC Genomics **9**:600.

34. **Peyron P**, **Vaubourgeix J**, **Poquet Y**, **Levillain F**, **Botanch C**, **Bardou F**, **Daffé M**,
Emile J-F, **Marchou B**, **Cardona P-J**, **de Chastellier C**, **Altare F**. 2008. Foamy
macrophages from tuberculous patients' granulomas constitute a nutrient-rich
reservoir for *M. tuberculosis* persistence. PLoS Pathog **4**:e1000204.

35. **Russell DG**, **Cardona P-J**, **Kim M-J**, **Allain S**, **Altare F**. 2009. Foamy macrophages
and the progression of the human tuberculosis granuloma. Nat Immunol **10**:943–948.

36. **VanderVen BC**, **Fahey RJ**, **Lee W**, **Liu Y**, **Abramovitch RB**, **Memmott C**, **Crowe
AM**, **Eltis LD**, **Perola E**, **Deininger DD**, **Wang T**, **Locher CP**, **Russell DG**. 2015.
Novel inhibitors of cholesterol degradation in *Mycobacterium tuberculosis* reveal how
the bacterium's metabolism is constrained by the intracellular environment. PLoS
Pathog **11**:e1004679.

37. **Pandey AK**, **Sassetti CM**. 2008. Mycobacterial persistence requires the utilization of
host cholesterol. Proc Natl Acad Sci USA **105**:4376–4380.

38. **Muñoz Elías E**, **Upton A**, **Cherian J**, **McKinney JD**. 2006. Role of the methylcitrate
cycle in *Mycobacterium tuberculosis* metabolism, intracellular growth, and virulence.
Mol Microbiol **60**:1109–1122.

39. **Nazarova EV**, **Montague CR**, **La T**, **Wilburn KM**, **Sukumar N**, **Lee W**, **Caldwell S**,
Russell DG, **VanderVen BC**. 2017. Rv3723/LucA coordinates fatty acid and
cholesterol uptake in *Mycobacterium tuberculosis*. Elife **6**:678.

40. **Daniel J**, **Maamar H**, **Deb C**, **Sirakova TD**, **Kolattukudy PE**. 2011. *Mycobacterium
tuberculosis* uses host triacylglycerol to accumulate lipid droplets and acquires a
dormancy-like phenotype in lipid-loaded macrophages. PLoS Pathog **7**:e1002093.

41. **Low KL**, **Rao PSS**, **Shui G**, **Bendt AK**, **Pethe K**, **Dick T**, **Wenk MR**. 2009.
Triacylglycerol utilization is required for regrowth of in vitro hypoxic nonreplicating
Mycobacterium bovis bacillus Calmette-Guerin. J Bacteriol **191**:5037–5043.

42. **Deb C**, **Lee C-M**, **Dubey VS**, **Daniel J**, **Abomoelak B**, **Sirakova TD**, **Pawar S**,
Rogers L, **Kolattukudy PE**. 2009. A novel in vitro multiple-stress dormancy model
for *Mycobacterium tuberculosis* generates a lipid-loaded, drug-tolerant, dormant
pathogen. PLoS ONE **4**:e6077.

43. **Sirakova TD**, **Deb C**, **Daniel J**, **Singh HD**, **Maamar H**, **Dubey VS**, **Kolattukudy PE**.
2012. Wax ester synthesis is required for *Mycobacterium tuberculosis* to enter in vitro
dormancy. PLoS ONE **7**:e51641.

44.   **Sirakova TD**, **Dubey VS**, **Deb C**, **Daniel J**, **Korotkova TA**, **Abomoelak B**,
Kolattukudy PE. 2006. Identification of a diacylglycerol acyltransferase gene
involved in accumulation of triacylglycerol in *Mycobacterium tuberculosis* under
stress. Microbiology **152**:2717–2725.

45.   **Murphy DJ**. 2011. The dynamic roles of intracellular lipid droplets: from archaea to
mammals. Protoplasma **249**:541–585.

46.   **Waltermann M**, **Steinbüchel A**. 2005. Neutral lipid bodies in prokaryotes: Recent
insights into structure, formation, and relationship to eukaryotic lipid depots. J
Bacteriol **187**:3607–3619.

47.   **Bartz R**, **Li W-H**, **Venables B**, **Zehmer JK**, **Roth MR**, **Welti R**, **Anderson RGW**, **Liu**
P, **Chapman KD**. 2007. Lipidomics reveals that adiposomes store ether lipids and
mediate phospholipid traffic. J Lipid Res **48**:837–847.

48.   **Faletrov Y**, **Brzostek A**, **Plocinska R**, **Dziadek J**, **Rudaya E**, **Edimecheva I**,
Shkumatov V. 2016. Uptake and metabolism of fluorescent steroids by
mycobacterial cells. Steroids **117**:29–37.

49.   **Griffin JE**, **Pandey AK**, **Gilmore SA**, **Mizrahi V**, **McKinney JD**, **Bertozzi CR**,
Sassetti CM. 2012. Cholesterol catabolism by *Mycobacterium tuberculosis* requires
transcriptional and metabolic adaptations. Chem Biol **19**:218–227.

50.   **Russell DG**, **VanderVen BC**, **Lee W**, **Abramovitch RB**, **Kim M-J**, **Homolka S**,
Niemann S, **Rohde KH**. 2010. *Mycobacterium tuberculosis* wears what it eats. Cell
Host Microbe **8**:68–76.

51.   **Singh A**, **Crossman DK**, **Mai D**, **Guidry L**, **Voskuil MI**, **Renfrow MB**, **Steyn AJC**.
2009. *Mycobacterium tuberculosis* WhiB3 maintains redox homeostasis by regulating
virulence lipid anabolism to modulate macrophage response. PLoS Pathog
5:e1000545–16.

52.   **Hauschild P**, **Röttig A**, **Madkour MH**, **Al-Ansari AM**, **Almakishah NH**, **Steinbüchel**
A. 2017. Lipid accumulation in prokaryotic microorganisms from arid habitats. Appl
Microbiol Biotechnol **101**: 2203–2216.

53.   **Alvarez AF**, **Alvarez HM**, **Kalscheuer R**, **Waltermann M**, **Steinbüchel A**. 2008.
Cloning and characterization of a gene involved in triacylglycerol biosynthesis and
identification of additional homologous genes in the oleaginous bacterium
Rhodococcus opacus PD630. Microbiology **154**:2327–2335.

54.   **Lanfranconi MP**, **Alvarez AF**, **Alvarez HM**. 2015. Identification of genes coding for
putative wax ester synthase/diacylglycerol acyltransferase enzymes in terrestrial and
marine environments. AMB Express **5**:42.

55.   **Deb C**, **Daniel J**, **Sirakova TD**, **Abomoelak B**, **Dubey VS**, **Kolattukudy PE**. 2006. A
novel lipase belonging to the hormone-sensitive lipase family induced under
starvation to utilize stored triacylglycerol in *Mycobacterium tuberculosis*. J Biol Chem
281:3866–3875.

56.   **Desmond E**, **Gribaldo S**. 2009. Phylogenomics of sterol synthesis: insights into the
origin, evolution, and diversity of a key eukaryotic feature. Genome Biol Evol **1**:364–
381.

57.   **Wollam J**, **Antebi A**. 2011. Sterol regulation of metabolism, homeostasis, and
development. Annu Rev Microbiol **80**:885–916.

58. **Weingärtner O**, **Böhm M**, **Laufs U**. 2008. Controversial role of plant sterol esters in
the management of hypercholesterolaemia. Eur Heart J **30**:404–409.

59. **Lichtenstein AH**, **Deckelbaum RJ**. 2001. AHA Science Advisory. Stanol/sterol ester-
containing foods and blood cholesterol levels. A statement for healthcare
professionals from the Nutrition Committee of the Council on Nutrition, Physical
Activity, and Metabolism of the American Heart Association. Circulation **103**:1177–
1179.

60. **Bacchetti T**, **Masciangelo S**, **Bicchiega V**, **Bertoli E**, **Ferretti G**. 2011. Phytosterols,
phytostanols and their esters: from natural to functional foods. Mediterr J Nutr Metab
4:165–172.

61. **He W-S**, **Zhu H**, **Chen Z-Y**. 2018. Plant Sterols: Chemical and Enzymatic Structural
Modifications and Effects on Their Cholesterol-Lowering Activity. J Agric Food Chem
66:3047–3062.

62. **Casabon I**, **Crowe AM**, **Liu J**, **Eltis LD**. 2013. FadD3 is an acyl-CoA synthetase that
initiates catabolism of cholesterol rings C and D in actinobacteria. Mol Microbiol
87:269–283.

63. **Ding Y**, **Yang L**, **Zhang S**, **Wang Y**, **Du Y**, **Pu J**, **Peng G**, **Chen Y**, **Zhang H**, **Yu J**,
Hang H, **Wu P**, **Yang F**, **Yang H**, **Steinbüchel A**, **Liu P**. 2012. Identification of the
major functional proteins of prokaryotic lipid droplets. J Lipid Res **53**:399–411.

64. **Ding Y**, **Zhang S**, **Yang L**, **Na H**, **Zhang P**, **Zhang H**, **Wang Y**, **Chen Y**, **Yu J**, **Huo**
C, **Xu S**, **Garaiova M**, **Cong Y**, **Liu P**. 2013. Isolating lipid droplets from multiple
species. Nat Protoc **8**:43–51.

65. **Ichihara K**, **Yamaguchi C**, **Nishijima H**, **Saito K**. 2003. Preparation of FAME from
sterol esters. J Amer Oil Chem Soc **80**:833–834.

  Figure legends

  Fig. 1: Schematic representation of triacylglycerols (TAGs) and wax esters (WEs) as neutral

  lipid storage compounds in bacteria and of a cholesterol-ester. Structures of the major animal,

  plant, and fungal sterols cholesterol, β-sitosterol, and ergosterol.

  Fig. 2: Growth of strain RHA1 under nitrogen-limiting and nitrogen-excess conditions. **A)** at

  early stationary phase (around 100 h). **B)** TLC analysis of neutral lipid extracts from early

  stationary phase N-limited (-N) and N-excess (+N) cultures grown with different carbon

  sources. Abbreviations: CP, cholesterol palmitate; C, cholesterol; TAG, triacylglycerols; Ph,

  phytosterol mix; S, succinate; P, pyruvate; G, glucose; NH_4^+, N replenished in stationary-phase

  C-N culture. The volume of neutral lipid extracts applied to TLC plates was normalized on the

  extracted cellular dry weight of each culture. Red arrows indicate standard spots. Results were

  obtained on two TLC plates which were treated exactly the same way throughout the

  experiment. The results of one representative experiment out of several biological and

  technical replicates are shown for each substrate.

  Fig. 3: Growth of RHA1 on 1 mM cholesterol under nitrogen-limiting conditions **(A)** and

  nitrogen-excess conditions **(B)**. The volume of neutral lipid extracts applied to TLC plates was

  normalized on the basis of culture biomass as protein. One of two reproducible experiments is

  shown.

  Fig. 4: GC-FID analysis of accumulating neutral lipids in nitrogen-limited RHA1 cultures grown

  with cholesterol or succinate. **(A)** GC-FID chromatograms of authentic cholesteryl palmitate

  (C16:0) and cholesteryl heptadecanoate (C17:0) standards (blue line) and of a trimpalmitin

  TAG (C16:0$_3$) standard (orange line) and of neutral lipid extracts of cholesterol- and succinate-

  grown cells harvested 24 h after nitrogen depletion. Seven accumulating CE species were

  identified either by comparison to the standards or based on the proximity of their respective

  peaks to the standard peaks and their absence in succinate grown cells. Identified and

  predicted CEs are labelled with blue arrows, identified and predicted TAG species are labelled

  with orange arrows. **(B)** Ratio of accumulating CEs (blue bars) and TAGs (orange bars) in

  cholesterol-grown nitrogen-limited RHA1 cultures over time starting at nitrogen depletion (0 h).

  Peak areas of the seven CE- and nine TAG-peaks shown in Fig. 4A were normalized on the

  peak area of the internal standard cholestane (not shown) before calculating their respective

  sums. Average values of three biological replicates and their standard deviations are shown.

  Fig. 5: Identification of SEs accumulating in N-limited RHA1 cultures. GC-MS analysis of

  transesterified fraction 2 (black peaks) and untreated fraction 2 (grey peaks) of neutral lipid

extracts of nitrogen-limited cultures grown with **(A)** cholesterol and **(B)** phytosterols.

Transesterification of SEs produced free cholesterol and the free phytosterols β-sitosterol, β-

sitostanol, and campesterol, as well as seven fatty acid methyl esters of C15:0, C16:1, C16:0,

C17:1, C17:0, C18:0 and C18:1. One representative GC-MS chromatogram is shown. **(C)** Fatty

acid profile of transesterified fraction 2 derived from nine cholesterol-grown cultures grown

under nitrogen-limiting conditions from five independent experiments harvested approximately

24h after nitrogen depletion. Odd-numbered fatty acids are coloured blue and even-numbered

fatty acids are coloured red

Fig. 6: Isolation of lipid droplets from strain RHA1 grown with cholesterol (**C**) or palmitate (**P**)

or a mixture of cholesterol plus palmitate (**C+P**) under nitrogen-limiting conditions. **A)** TLC

analysis of neutral lipids derived from total lipid extracts from isolated lipid droplets.

Abbreviations as in previous figures. **B)** Molar ratio of TAGs, CEs and free cholesterol in lipid

droplets derived from cholesterol, cholesterol plus palmitate and palmitate grown cells. **C)**

Ratio of even to odd-numbered fatty acids in CEs and TAGs in lipid droplets and the respective

contributing fatty acid chain lengths. The results of two independent experiments (I and II) are

shown in **B)** and **C)** and of experiment II in **A)**.

Fig. 1: Schematic representation of triacylglycerols (TAGs) and wax esters (WEs) as neutral

lipid storage compounds in bacteria and of a cholesterol-ester. Structures of the major animal,

plant, and fungal sterols cholesterol, β-sitosterol, and ergosterol.

Fig. 2: Growth of strain RHA1 under nitrogen-limiting and nitrogen-excess conditions. **A)** at

early stationary phase (around 100 h). **B)** TLC analysis of neutral lipid extracts from early

stationary phase N-limited (-N) and N-excess (+N) cultures grown with different carbon

sources. Abbreviations: CP, cholesterol palmitate; C, cholesterol; TAG, triacylglycerols; Ph,

phytosterol mix; S, succinate; P, pyruvate; G, glucose; NH_4^+, N replenished in stationary-phase

C-N culture. The volume of neutral lipid extracts applied to TLC plates was normalized on the

extracted cellular dry weight of each culture. Red arrows indicate standard spots. Results were

obtained on two TLC plates which were treated exactly the same way throughout the

experiment. The results of one representative experiment out of several biological and

technical replicates are shown for each substrate.

Fig. 3: Growth of RHA1 on 1 mM cholesterol under nitrogen-limiting conditions **(A)** and

nitrogen-excess conditions **(B)**. The volume of neutral lipid extracts applied to TLC plates was

normalized on the basis of culture biomass as protein. One of two reproducible experiments is

shown

Fig. 4: GC-FID analysis of accumulating neutral lipids in nitrogen-limited RHA1 cultures grown

with cholesterol or succinate. **(A)** GC-FID chromatograms of authentic cholesteryl palmitate

(C16:0) and cholesteryl heptadecanoate (C17:0) standards (blue line) and of a tripalmitin TAG

(C16:0$_3$) standard (orange line) and of neutral lipid extracts of cholesterol- and succinate-

grown cells harvested 24 h after nitrogen depletion. Seven accumulating CE species were

identified either by comparison to the standards or based on the proximity of their respective

peaks to the standard peaks and their absence in succinate grown cells. Identified and

predicted CEs are labelled with blue arrows, identified and predicted TAG species are labelled

with orange arrows. **(B)** Ratio of accumulating CEs (blue bars) and TAGs (orange bars) in

cholesterol-grown nitrogen-limited RHA1 cultures over time starting at nitrogen depletion (0 h).

Peak areas of the seven CE- and nine TAG-peaks shown in Fig. 4A were normalized on the

peak area of the internal standard cholestane (not shown) before calculating their respective

sums. Average values of three biological replicates and their standard deviations are shown.

Fig. 5: Identification of SEs accumulating in N-limited RHA1 cultures. GC-MS analysis of
transesterified fraction 2 (black peaks) and untreated fraction 2 (grey peaks) of neutral lipid
extracts of nitrogen-limited cultures grown with **(A)** cholesterol and **(B)** phytosterols.
Transesterification of SEs produced free cholesterol and the free phytosterols β-sitosterol, β-
sitostanol, and campesterol, as well as seven fatty acid methyl esters of C15:0, C16:1, C16:0,
C17:1, C17:0, C18:0 and C18:1. One representative GC-MS chromatogram is shown. **(C)** Fatty
acid profile of transesterified fraction 2 derived from nine cholesterol-grown cultures grown
under nitrogen-limiting conditions from five independent experiments harvested approximately
24h after nitrogen depletion. Odd-numbered fatty acids are coloured blue and even-numbered
fatty acids are coloured red

Fig. 6: Isolation of lipid droplets from strain RHA1 grown with cholesterol (**C**) or palmitate (**P**) or a mixture of cholesterol plus palmitate (**C+P**) under nitrogen-limiting conditions. **A)** TLC analysis of neutral lipids derived from total lipid extracts from isolated lipid droplets. Abbreviations as in previous figures. **B)** Molar ratio of TAGs, CEs and free cholesterol in lipid droplets derived from cholesterol, cholesterol plus palmitate and palmitate grown cells. **C)** Ratio of even to odd-numbered fatty acids in CEs and TAGs in lipid droplets and the respective contributing fatty acid chain lengths. The results of two independent experiments (I and II) are shown in **B)** and **C)** and of experiment II in **A)**.

901