Steryl ester formation and accumulation in steroid-degrading bacteria

Johannes Holert[1], Kirstin Brown[1], Ameena Hashimi[1], Lindsay D. Eltis[1], and William W. Mohn[1]

[1]Department of Microbiology & Immunology, Life Sciences Institute, University of British Columbia, Vancouver, British Columbia, Canada

Running title: Bacterial steryl ester synthesis

Correspondence: William W. Mohn, Department of Microbiology & Immunology, Life Sciences Institute, University of British Columbia, 2350 Health Sciences Mall, Vancouver, BC, Canada, V6T 1Z3. Email: wmohn@mail.ubc.ca

ABSTRACT

Steryl esters (SEs) are important storage compounds in many eukaryotes and are often
prominent components of intracellular lipid droplets. Here we demonstrate that selected
Actino- and Proteobacteria growing on sterols are also able to synthesize SEs and to
sequester them in cytoplasmic lipid droplets. We found cholesteryl ester (CE) formation in
members of the actinobacterial genera *Rhodococcus*, *Mycobacterium*, and *Amycolatopsis* as
well as several members of the proteobacterial Cellvibrionales order. CEs maximally
accumulated under nitrogen-limiting conditions, suggesting that steryl ester formation plays a
crucial role for storing excess energy and carbon under adverse conditions. *Rhodococcus*
jostii RHA1 was able to synthesize phytosteryl- and cholesteryl esters, the latter reaching up
to 7% of its cellular dry weight and 69% of its lipid droplets. Purified lipid droplets from RHA1
contained CEs, free cholesterol and triacylglycerols. In addition, we found formation of CEs
in *Mycobacterium tuberculosis* when grown with cholesterol plus an additional fatty acid
substrate. This study provides a basis for the application of bacterial whole cell systems in
the biotechnological production of SEs for use in functional foods and cosmetics.

IMPORTANCE

Oleaginous bacteria exhibit great potential for the production of high-value neutral lipids,
such as triacylglycerols and wax esters. This study describes the formation of steryl esters
(SEs) as neutral lipid storage compounds in sterol-degrading oleaginous bacteria, providing
a basis for biotechnological production of SEs using bacterial systems with potential
applications in the functional food, nutraceutical, and cosmetic industries. We found
cholesteryl ester (CE) formation in several sterol-degrading Actino- and Proteobacteria under
nitrogen limiting conditions, suggesting an important role of this process in storing energy
and carbon under adverse conditions. In addition, *Mycobacterium tuberculosis* grown on
cholesterol accumulated CEs in the presence of an additional fatty acid substrate.

INTRODUCTION

Many bacteria synthesize and accumulate carbon and energy storage compounds under
environmental stress conditions to increase their survival in environments with fluctuating
nutrient conditions (1). A much smaller subset of bacteria are oleaginous and accumulate
neutral lipids, namely triacylglycerols (TAGs) and wax esters (WEs), as storage compounds.
Although TAG and WE synthesis is rarely described in other bacteria, their accumulation is
widespread within the actinobacterial genera *Streptomyces*, *Nocardia*, *Rhodococcus*,
Mycobacterium, *Dietzia*, and *Gordonia*, and has also been reported in the proteobacterial
genera *Acinetobacter*, *Alcanivorax* and *Marinobacter* (2-7). TAGs and WEs are highly
hydrophobic, non-polar lipids, which are formed by esterification of a glycerol molecule to
three fatty acids or by esterification of a long chain fatty alcohol to a fatty acid, respectively
(**Fig. 1**). Potential biotechnological applications of TAG and WE producing bacteria gained
increasing attention over the last years due to the considerable industrial relevance of these
compounds as biofuels and other products such as feed additives, cosmetics, and lubricants
(8). Especially oleaginous *Rhodococcus* strains have been suggested as promising microbial
cell factories for the large scale production of neutral lipids due to their ability to convert
agroindustrial wastes into valuable lipids and their accessibility to cell engineering for
increased lipid production (9).

Accumulation of neutral lipids in bacteria is typically induced by environmental stress
conditions, in which a suitable carbon source is in excess while additional growth is restricted
by a deficiency of other nutrients or environmental conditions. The most prominent example
is nitrogen limitation, which induces TAG and WE synthesis in a wide range of neutral lipid-
accumulating Actino- and Proteobacteria. This has been most extensively studied in
oleaginous *Rhodococcus* strains, which can accumulate up to 70% of their cellular dry weight
as TAGs upon nitrogen depletion, depending on the available carbon source (10-12). In
Mycobacterium tuberculosis, the causative agent of tuberculosis, dormancy inducing
conditions such as nitric oxide (NO) stress or iron limitation have been shown to initiate
accumulation of TAGs and WEs (13, 14). These intracellular carbon and energy storage
compounds have been suggested to be important for the pathogenicity of *M. tuberculosis* (15,
16).

TAGs and WEs are sequestered into cytoplasmic lipid droplets, which are
multifunctional organelles encapsulated by a phospholipid monolayer and specific proteins
(17). Analogous to oleaginous bacteria, many eukaryotes also sequester neutral lipids as
carbon and energy reservoirs into cytoplasmic lipid droplets. In contrast to bacterial lipid
droplets, many eukaryotic lipid droplets contain sterols in addition to TAGs and WEs (18).
Sterols constitute a key class of lipids in most animals, fungi, and plants, where they play

essential roles as membrane constituents and precursors for vitamins and steroid hormones. The dominant sterol in animals is cholesterol, while plants have β-sitosterol and other phytosterols, and fungi have ergosterol (**Fig. 1**). Within eukaryotic lipid droplets, the sterol 3-hydroxyl group is typically esterified with fatty acids forming so-called steryl esters (SEs) (**Fig. 1**), which act as a sterol storage pool. Synthesis and breakdown of SEs plays a vital role in sterol homeostasis and imbalances between free and esterified cholesterol pools are related to several human diseases, such as Wolman disease, cholesteryl ester storage disease, atherosclerosis, and Alzheimer's disease (19). In contrast to eukaryotes, sterol synthesis is rare in bacteria and mainly restricted to *Myxococcales* and *Methylococcales* (20). These bacterial sterols have structural features that distinguish them from eukaryotic sterols (21). Further, it is not known whether these bacteria are able to esterify and store their sterols in lipid droplets.

Sterols are abundant growth substrates for bacteria in the environment, and sterol degradation has been characterized in many actinobacterial isolates and is highly conserved in several genera of the Corynebacterineae suborder, namely the genera *Amycolicicoccus, Dietzia, Gordonia, Mycobacterium, Nocardia, Rhodococcus,* and *Tsukamurella* (22). In addition, we recently isolated several sterol-degrading strains belonging to the gammaproteobacterial order Cellvibrionales (23), which are also able to degrade sterols such as cholesterol. Under aerobic conditions, sterols are degraded by a conserved pathway into central metabolic moieties such as acetyl-CoA and propionyl-CoA (24), which are subsequently channeled into central metabolic pathways for energy conservation and production of biomass. Among those sterol-degrading bacteria are many that are also well known for their ability to produce TAGs and WEs as neutral lipid storage compounds, such as the Actinobacteria, *Rhodococcus jostii* RHA1 (RHA1) (25) and *Mycobacterium tuberculosis* (*Mtb*) (26). Key enzymes in bacterial TAG and WE synthesis are CoA-dependent, bifunctional WS/DGAT enzymes encoded by *atf* genes. These enzymes catalyze both the acetylation of DAGs to TAGs and of fatty alcohols to WEs (27). Most WS/DGATs are promiscuous enzymes accepting a variety of acyl-CoAs and alcohols as substrates (5, 28). Interestingly, the WS/DGAT enzyme AtfA from the oleaginous, non-sterol degrading Proteobacterium *Acinetobacter calcoaceticus* ADP1 has been shown to have SE-synthase activity when heterologously expressed in *E. coli* or the yeast *Saccharomyces cerevisiae* (29). Over 50 years ago, two studies reported that different *Mycobacterium* strains produced small amounts of cholesteryl fatty acid esters when incubated with cholesterol in cell suspensions (30) and that *Mycobacterium smegmatis* also produced esters of other sterols (31). However, these studies did not provide further insight into the cellular localization of SE accumulation or which incubation conditions elicit SE formation, and no further reports have been published on bacterial SE synthesis since then.

Here we investigate the ability of sterol-degrading bacteria to synthesize lipid storage
compounds from sterol substrates under adverse growth conditions and report formation of
SEs as lipid storage compounds in bacteria. This significantly extents the knowledge about
bacterial SE formation and provides a basis for biotechnological production of SEs using
bacterial systems with potential applications in the functional food, nutraceutical, and
cosmetic industries.
123

RESULTS

Accumulation of SEs in RHA1 under nitrogen limiting conditions

To investigate formation of lipid-storage compounds from sterol substrates, *Rhodococcus*
jostii RHA1 was grown in mineral medium with 1 mM cholesterol or 1 mM of a phytosterol
mix, containing 76.4% β-sitosterol, 10.4% β-sitostanol, 7.3% campesterol and 1%
campestanol, under nitrogen-limiting and nitrogen-excess conditions. In cholesterol-grown
cultures, RHA1 reached a biomass in early stationary phase that was approximately four-fold
higher under nitrogen excess than under nitrogen-limiting conditions (**Fig. 2A**). In contrast to
the N-excess cultures, the N-limited cultures had no residual ammonium but did have
residual cholesterol. Similar results were obtained during growth on the phytosterol substrate
mix (not shown). Thin layer chromatography (TLC) analysis of neutral lipid extracts revealed
compound spots with the same R_f-value as an authentic cholesterol-palmitate standard ($R_f =$
0.78) in N-limited cholesterol and phytosterol cultures, but not in the respective N-excess
cultures or in N-limited or N-excess cultures grown on non-sterol substrates (**Fig. 2B)**. In
addition, compounds with similar Rf-values as an authentic tripalmitin TAG standard ($R_f =$
0.24) were present in extracts of all N-limited cultures. Time-dependent sampling of RHA1
grown on cholesterol under N-excess and N-limiting conditions revealed that increased
formation of the presumptive cholesteryl esters coincided with the depletion of ammonium
from the medium (**Fig. 3A**). Under N-excess conditions only minimal amounts of these
compounds accumulated during exponential growth and stationary phase (**Fig. 3B**). When
nitrogen was replenished in an N-limited, stationary-phase, cholesterol-grown culture, the
accumulated lipids including the presumptive SEs disappeared within 72 h (**Fig. 2B**).

To confirm the formation of CEs, we analyzed native neutral lipid extracts from strain
RHA1 grown with cholesterol or succinate under N-limiting conditions in early stationary
phase using a gas chromatography (GC)-coupled flame ionization detector (FID).
Comparison to a cholesteryl-palmitate (C16:0) and a cholesteryl-heptadecanoate standard
(C17:0) confirmed that these CEs accumulated in cholesterol-grown RHA1 cells (**Fig. 4A**),
but were absent in succinate-grown cells. In addition, five other compounds from cholesterol-
grown cells eluted close to these CEs, which were absent in succinate-grown cells,
suggesting that seven CE species, with different fatty acid chain lengths, are formed with
cholesterol as substrate. Comparison to a tripalmitate ((C16:0)$_3$) TAG standard suggested
the accumulation of up to nine different TAG species in strain RHA1 grown with cholesterol
and eleven different TAG species in cells grown with succinate (**Fig. 4A**). Time-dependent
sampling starting at the onset of nitrogen depletion from the media followed by GC-FID
analysis showed that the CE:TAG ratio shifted from approximately 60% to 72% within the
first 24 h of incubation and to 69% after 48 h.

To gain further insight into the nature of SEs accumulating in strain RHA1, neutral
lipid extracts of cholesterol- and phytosterol-grown cultures were fractionated on silica
columns producing five lipid fractions (**supplemental Fig. S1**). SEs eluted in fraction 2 and
were well separated from other lipids, like TAGs (Fractions 3 and 4) and free sterols
(fractions 5). Aliquots of fraction 2 were subjected to alkaline transesterification with
methanol. GC-MS analysis showed that free cholesterol or the free phytosterols (β-sitosterol,
β-sitostanol, and campesterol) were formed from transesterification, which were not present
in untransesterified extracts (**Fig. 5A-B**). Only the respective sterols provided as a growth
substrate were detected in this analysis. In addition, the lack of other free alcohols after
transesterification confirmed the absence of wax esters in neutral lipid extracts. In addition, a
range of seven fatty acid methyl esters (FAMEs) were detected after transesterification of
cholesteryl esters, supporting the results from the GC-FID analysis. Those were identified as
methyl esters of C15:0, C16:1, C16:0, C17:1, C17:0, C18:1 and C18:0 (**Fig. 5A**). The most
abundant fatty acids were the odd-numbered fatty acids, C17:0, C17:1, and C15:0, which
constituted more than 75% of produced FAMEs (**Fig. 5C**). GC-MS analysis of transesterified
fraction 2 confirmed that no detectable amounts of other free alcohols indicative of wax
esters were present (not shown). Analysis of all neutral lipid extracts using a GC-MS method
for wax-ester analysis (32) showed that no detectable amounts of wax esters were present in
any sample (not shown), which is in agreement with earlier reports that wax esters are only
formed in minimal amounts in RHA1 (32).
This confirmed that the previously unidentified lipids accumulating in N-limited RHA1
cultures grown with cholesterol or phytosterols are fatty acid steryl esters. In the late
stationary phase (100 h) CEs accounted for 6.8 +/- 1.4% of the cellular dry weight in N-
limited cultures (n = 5 from two independent experiments) based on GC-MS quantification.
185

Localization of accumulated cholesterol and CEs in RHA1
To confirm that accumulating neutral lipids are localized intracellularly, we stained stationary
phase, cholesterol-grown, washed, N-excess, and N-limited RHA1 cells with the lipophilic
fluorescence stain Nile Red. Microscopic analysis showed that stained lipids accumulated as
lipid droplets in the cell interior (**supplemental Fig. S2**) of N-limited cells but were absent
from N-excess cells. In addition, we incubated RHA1 with fluorescently labelled cholesterol
(BODIPY-cholesterol) and sodium palmitate for 48 h without a nitrogen source. Microscopic
analysis showed that fluorescently labelled cholesterol was mainly present within lipid
droplets inside the cells (**supplemental Fig. S3**) but mostly absent from cell membranes.
TLC analysis of neutral lipid extracts of cells with fluorescently labelled cholesterol showed
that no CEs were formed (not shown), which is most likely because the altered chemical

structure of the labelled cholesterol side chain inhibits esterification. These results confirm
that cholesterol and CEs are mainly stored in intracellular lipid droplets.

To gain further insight into the location of CEs within RHA1 cells, lipid droplets were
isolated from RHA1 cultures grown with cholesterol or sodium palmitate or a mixture of both
substrates under N-limiting conditions. With all three substrate combinations, gradient
centrifugation of total cell extracts produced visible bands of lipid droplet on top of the
sucrose gradient (**supplemental Fig. S4**). TLC analysis revealed that CEs and free
cholesterol were present in neutral lipid extracts of lipid droplets derived from cells grown on
cholesterol but not in lipid droplets derived from cells grown on palmitate alone, while TAGs
were present under all three conditions (**Fig. 6A**). This was confirmed by fractionation and
subsequent transesterification of neutral lipid extracts (not shown). In addition, the lack of
other free alcohols after transesterification confirmed the absence of wax esters in lipid
droplets. In two independent experiments 52% and 58% of lipids in lipid droplets derived
from cells grown on cholesterol were identified as CEs, 31% and 15% as TAGs and 17% and
27% as free cholesterol. Similar results were obtained for lipid droplets derived from cells
grown on cholesterol plus palmitate, with 50% and 69% CEs, 35% and 23% TAGs and 15%
and 8% cholesterol, while TAGs were the only neutral lipids in lipid droplets derived from
cells grown on palmitate (**Fig. 6B**). In lipid droplets from cells grown on cholesterol, between
60% to 70% of fatty acids in CEs and TAGs were odd-numbered (C15 and C17), with C17:0
and C17:1 making up around 90% of these fatty acids (**Fig. 6C**). In lipid droplets from cells
grown with cholesterol plus palmitate, 23% to 55% of fatty acids were odd-numbered. TAGs
of cells grown on palmitate contained almost exclusively (~95%) even-numbered fatty acids.
(**Fig. 6C**).
220

Formation of SEs in other sterol-degrading bacteria
To test whether steryl ester formation also occurs in other sterol-degrading bacteria, we
extracted neutral lipids from the Actinobacteria, *Mycobacterium smegmatis* strain mc[2]155,
Mycobacterium sp. strain BC8-1, *Rhodococcus opacus* strain Chol-4, *Rhodococcus ruber*
strain Chol-4, and *Amycolatopsis* sp. strain ATCC 39116, as well as the
Gammaproteobacteria, *Zhongshania* sp. strain SB11-1A, *Haliaceae* sp. strain BC5-1, and
the BD1-7 clade bacterium strain SB11-3 grown on cholesterol under nitrogen-limiting and
nitrogen-excess conditions. The latter strains were isolated in our lab (23) and are among the
first Proteobacteria reported to be able to degrade sterols. TLC analysis indicated these
strains produced SEs in N-limited cultures in the stationary phase in differing amounts, with
highest yields in the *Rhodococcus*, *Mycobacterium*, and *Zhongshania* strains and lower
yields in *Amycolatopsis*, *Haliaceae*, and the BD1-7 clade strain (**supplemental Fig. S5A**).

Subsequent fractionation of neutral lipid extracts followed by transesterification and GS-MS
analysis confirmed the formation of CEs in all strains (**supplemental Fig. S5B**).

In addition, we tested steryl ester formation in *Mycobacterium tuberculosis* (*Mtb*)
Erdmann growing on cholesterol and on cholesterol plus sodium palmitate. Because *Mtb* did
not grow when sodium palmitate was added to the medium prior to inoculation we spiked
palmitate into cultures growing on cholesterol after 5 days. TLC analysis of neutral lipid
extracts of *Mtb* indicated the formation of small amounts of CEs in cells growing on
cholesterol after 7 days and in cells growing on cholesterol plus palmitate after 12 and 13
days, when cells were in stationary phase (**supplemental Fig. S6A-B**). Subsequent
fractionation of neutral lipid extracts followed by transesterification and GS-MS analysis
confirmed the formation of CEs (**supplemental Fig. S6C**). The only detectable FAME in this
fraction was C16:0. Formation of free hexacosanol after transesterification suggested that a
C_{26}-OH wax ester was also formed. In addition, we tested whether nitric oxide (NO) stress or
iron limitation, which are known inducers of neutral lipid accumulation in *Mtb* (13, 14),
increase CE formation from cholesterol or cholesterol plus palmitate. For this, cholesterol-
grown cell suspensions of *Mtb* were incubated with cholesterol alone or cholesterol plus
palmitate. To induce NO stress, two pulses of 0.5 mM spermine-NONOate were added at the
beginning of the experiment and again after 16 h of incubation and samples were taken after
16 h and after 2 days of incubation. Spermine (0.5 mM) was added to control cultures.
Deferoxamine (0.5 mM) was added as an iron chelator and samples were taken after 2 and 4
days of incubation. Confirming the results from the initial growth experiment, all extracts from
Mtb cells incubated with cholesterol plus palmitate contained CEs with C16:0 as only
detectable FAME. However, addition of NO or deferoxamine did not increase the amount of
accumulating CEs (**supplemental Fig. S6D**). No significant accumulation of CEs occurred in
treated cells incubated with only cholesterol. Small amounts of free cholesterol were
detected after transesterification of extracts of cell suspensions treated with NO and
incubated for 2 days with cholesterol (not shown).

DISCUSSION

Steryl esters (SEs) are important storage compounds in many eukaryotes and are often prominent components of intracellular lipid droplets . Here we demonstrate that selected sterol-degrading Actino- and Proteobacteria are also able to synthesize SEs and to sequester them in cytoplasmic lipid droplets. We found SE formation in all sterol-degrading strains investigated in this study, including members of the actinobacterial genera, *Rhodococcus*, *Mycobacterium*, and *Amycolatopsis,* as well as members of the proteobacterial Cellvibrionales order. This is supported by findings from the 1960s, which showed that some *Mycobacterium* strains were able to produce different SE in small amounts (up to 0.5 % of the cellular dry weight for cholesterol) (30, 31). The ability of other oleaginous bacteria to synthesize SEs has been overlooked in other studies, which exclusively used non-steroidal substrates, such as glucose, gluconate, or benzoate, when investigating neutral lipid accumulation. Our results suggest that the ability to accumulate excess sterols as an energy and carbon source under adverse conditions plays a crucial role in sterol-degrading bacteria.

Using *Rhodococcus jostii* RHA1 as an oleaginous, sterol-degrading model organism, we found accumulation of minor amounts of CEs during exponential growth on cholesterol when ammonium was available, and a significant increase in CE and TAG accumulation, when ammonium was limiting, indicating that nitrogen limitation stimulates SE formation. This extends previous reports that nitrogen limitation is a major inducer for the accumulation of TAGs and WEs in RHA1 with non-steroidal substrates (12, 33). While we found significant accumulation of CEs in the two free-living *Mycobacterium* strains mc^{2}115 and BC8-1 under nitrogen limiting conditions with cholesterol as the only substrate, the pathogenic strain *Mycobacterium tuberculosis* (*Mtb*) Erdmann did not grow on cholesterol in nitrogen-limited medium. However, small amounts of cholesterol-palmitate accumulated in this strain when growing on cholesterol plus palmitate as substrates without any stress conditions, demonstrating that pathogenic *Mtb* strains are able to accumulate CEs when a sterol and a fatty acid are available. During infection, *Mtb* triggers formation of foamy macrophages (34, 35), which provide a cholesterol and fatty acid-rich microenvironment, and *Mtb* metabolizes these host-derived lipids during infection and persistence (36-38). Interestingly, cholesterol and fatty acid uptake are co-regulated in *Mtb* (39), suggesting that *Mtb* has simultaneous access to both lipid classes *in vivo*. SE synthesis and accumulation could act as a buffer system for excess lipids in *Mtb* providing a way to regulate intracellular lipid metabolism. In addition, SEs could serve as a carbon and energy reservoir in *Mtb* under adverse conditions. In a similar manner, TAG accumulation and sequestration in cytoplasmic lipid droplets was shown to occur *in vivo* in pathogenic *Mtb* (16, 40) and was proposed to be important for

persistence and reactivation (41). Accumulation of TAGs and WEs was shown to be
stimulated by various dormancy-inducing stress conditions, such as hypoxia, NO stress, iron
limitation, and increased acidity, which induce expression of several WS/DGAT encoding
genes in *Mtb* (13, 42-44). Although we did not detect increased formation of CEs in *Mtb*
under NO stress or iron limiting conditions, a combination of stress factors as described in
(42) might provide deeper insight into the regulation of SE synthesis in *Mtb*.

Lipid droplets have been identified in most domains of life, including many
eukaryotes, as well as archaeal and bacterial prokaryotes (45). While the general structure of
lipid droplets is conserved in all these organisms, comprising a neutral lipid core coated by a
phospholipid monolayer and various peripheral proteins, the neutral lipid composition varies
significantly between organisms (17, 46). Our results show that CEs, in addition to TAGs and
free cholesterol, are sequestered in intracellular lipid droplets in RHA1. The lipid droplet
composition in RHA1 grown on cholesterol is very similar to that of lipid droplets found in
selected mammalian lipid droplet-producing cell lines (chinese hamster ovaries CHO K2, rat
kidney normal NKR and immortalized human fibroblasts SV589) (47), with around 50% CEs,
30% TAGs and 20% free cholesterol. The sequestration of cholesterol in lipid droplets is in
agreement with an earlier report showing that fluorescently labeled cholesterol predominantly
localizes in lipid droplets in mycobacteria (48).

Interestingly, the majority of fatty acids stored in TAGs and CEs in cholesterol-grown
RHA1 cells were odd-numbered fatty acids, with C17:0 and C17:1 as main fatty acids, which
is in stark contrast to the mostly even-numbered fatty acids typically found in eukaryotic
neutral lipids. Complete metabolism of one cholesterol molecule through the canonical
bacterial 9,10-seco steroid-degradation pathway is proposed to yield four propionyl-CoAs,
four acetyl-CoAs, one pyruvate and one succinyl-CoA (24). The preponderance of odd-chain
fatty acids in neutral lipids accumulating in RHA1 with cholesterol as carbon source suggests
that propionyl-CoA was a significant building block for neutral-lipid fatty acids. Elevated ratios
of odd-numbered to even-numbered fatty acids have also been found in TAGs accumulating
in RHA1 growing with benzoate under nitrogen limiting conditions, where propionyl-CoA was
presumably generated through the methylmalonyl-CoA pathway from succinate (12).
Cholesterol-derived propionyl-CoA has been shown to be toxic to *Mtb* if not metabolized (49),
and incorporation of propionyl-CoA into cell wall lipids (50) and TAGs (51) has been
suggested as a mechanism to detoxify propionate in *Mtb*. Cholesterol-degrading bacteria,
such as RHA1 and *Mtb*, might encounter increased cellular propionyl-CoA levels under
adverse growth conditions and the formation of odd-numbered fatty acid CEs might provide
an additional sink for this toxic end-product. However, we did not detect odd-numbered fatty
acid CEs in *Mtb*, which accumulated CEs in significant amounts only in the presence of an
additional fatty acid substrate under the tested conditions.

Sterols are abundant growth substrates for bacteria in the environment and sterol
degradation is conserved in many actinobacterial genera, particularly in members of the
Corynebacterineae (22). Recently, we have also identified sterol-degradation in some
isolates of the oligotrophic marine Gammaproteobacteria (OMG) group belonging to the
Cellvibrionales order (23). Steroid-degrading bacteria are globally distributed and
actinobacterial ones are particularly abundant within soil and rhizosphere as well as in deep
ocean habitats. Our results indicate that SE formation is conserved in sterol-degrading
bacteria, suggesting that this trait might play important roles in these environments.
Supporting this hypothesis, most neutral-lipid accumulating Actinobacteria isolated from soil
samples (52) belong to taxa in which the ability to degrade sterols is conserved (22) and are
thus likely to produce SEs. *De novo* synthesis of sterols is rare in bacteria, being largely
restricted to *Myxococcales* and *Methylococcales* (20) and it is not known whether these
bacteria are able to store these sterols as esters.

We searched the genomes of strain RHA1 and of *Mtb* Erdmann for homologs of one
prokaryotic and fifty eukaryotic proteins with characterized steryl ester synthase activity
available in the Swiss-Prot database, including phosphatidylcholine-sterol *O*-acyltransferases
(EC 2.3.1.43), diacylglycerol *O*-acyltransferases (EC 2.3.1.20), sterol *O*-acyltransferase (EC
2.3.1.26), and phospholipid:diacylglycerol acyltransferases (EC 2.3.1.158). None of these
proteins had any significant similarity (accepted coverage > 85%, percent identity > 30%, E-
values < 0.01) to RHA1 or *Mtb* proteins. Based on the multifunctionality and substrate
promiscuity of TAG- and WE-producing WS/DGAT enzymes, it is feasible that these
enzymes might carry out an additional function as steryl ester synthases. Out of the 16
predicted WS/DGATs in RHA1, Atf6 (WP_011594556.1) and Atf8 (WP_011597548.1) have
the highest amino acid sequence identity (40% and 39%, respectively) to AtfA from the
oleaginous, non-sterol degrading Proteobacterium *Acinetobacter calcoaceticus* ADP1, which
was shown to have SE-synthase activity when heterologously expressed in *E. coli* or *S.*
cerevisiae (29). TAG and WE synthase activities have been confirmed for Atf8 and Atf6,
respectively (12, 28) as well as for an Atf6 ortholog in *Rhodococcus opacus* PD630 (53), and
deletions of the respective genes significantly decreased neutral lipid accumulation. In
addition, transcription of *atf8* was strongly induced upon nitrogen depletion in RHA1 growing
on benzoate, exhibiting the highest transcript levels of all 16 *atf* genes under nitrogen limiting
conditions (12). In contrast, *atf6* transcripts were most abundant under nitrogen excess
conditions with no apparent transcriptional induction upon nitrogen depletion. WS/DGAT
genes identified in soil communities were exclusively from Actinomycetales (54), which
includes the steroid-degrading suborder *Corynebacterineae*. In contrast, WS/DGAT genes
from marine sediments were exclusively of gammaproteobacterial origin, predominantly from
members of the oligotrophic marine Gammaproteobacteria (OMG) group, of which we

recently identified several sterol-degrading strains (22, 23). BLASTp analysis (max E-value

10^{-15}) of AtfA from *Acinetobacter calcoaceticus* ADP1 against the proteomes of the

cholesteryl ester accumulating OMG bacteria *Zhongshania* sp. strain SB11-1A, *Haliaceae*

sp. strain BC5-1 and the BD1-7 clade bacterium SB11-3 suggested that these strains encode

between four and six WS/DGAT paralogs, further supporting our hypothesis that these

proteins might function as steryl ester synthases.

Our results indicate that accumulated SEs can be reactivated and degraded in RHA1

upon relief of growth-limiting conditions, suggesting that SEs might play a vital role as a

carbon and energy reservoir in these environments where fluctuating nutrient availability is

common. We searched the genomes of strain RHA1 and of *Mtb* Erdmann for homologs of

three prokaryotic and fifteen eukaryotic proteins with characterized or predicted sterol

esterase activity available in the TrEMBL and Swiss-Prot databases (EC 3.1.1.13). However,

none of the query proteins had significant similarity (accepted coverage > 85%, percent

identity > 30%, E-values < 0.01) to RHA1 or *Mtb* proteins. Strain RHA1 encodes up to 54

putative lipases/esterase enzymes (33), which were suggested to contribute to neutral lipid

degradation, but no distinct degradation system for endogenous TAGs or WEs has been

identified so far. In contrast, the lipase family protein LipY in *M. tuberculosis* has been shown

to have TAG degradation activity (55).

Interestingly, several animal lineages such as insects and soil-dwelling nematodes

have lost the ability to synthesize sterols *de novo* (56) and thus rely on uptake of sterols with

their diet (57). It is tempting to speculate that sterol and steryl ester accumulating bacteria in

oligotrophic habitats such as soils or freshwater systems might constitute an important

source for sterols for bacteriovorous, sterol-auxotrophic eukaryotes such as *Caenorhabditis*

elegans or *Daphnia magna.*

RHA1 was also able to form esters of the phytosterols β-sitosterol, β-sitostanol, and

campesterol under nitrogen-limiting conditions. In recent years, plant-derived phytosterols

and their esters were recognized as potential agents to lower serum cholesterol levels in

hypercholestemic patients (58, 59) by inhibiting absorption of endogenous, dietary

cholesterol in the intestine. Today phytosterols and phytosteryl esters are approved in many

countries as safe food additives, and several so-called functional foods are marketed

worldwide, which are supplemented with phytosterol and their esters (60). Industrial

production of phytosteryl esters is mainly accomplished by chemical esterification or

transesterification of plant-derived sterols, which requires high temperatures and the use of

strong acids or bases (61). Our study provides a basis for biotechnological production of

phytosteryl esters using non-pathogenic whole-cell bacterial systems, thus eliminating the

requirement for elevated temperature and acid or base catalysts. Similar to RHA1, many

other sterol-degrading Actinobacteria are also able to use phytosterols as growth substrates,

providing an excellent screening resource for efficient phytosteryl ester accumulation.

MATERIAL AND METHODS

Bacteria and culture conditions

Rhodococcus jostii strain RHA1, as well as *Mycobacterium smegmatis* strain mc^{2}155,

Rhodococcus opacus strain PD630, *Rhodococcus ruber* strain Chol-4, and *Amycolatopsis*

sp. strain ATCC 39116 were cultivated in M9 minimal medium supplemented with trace

elements and vitamin B1 at 30 °C. The marine bacteria *Mycobacterium* strain BC8-1,

Zhongshania sp. strain SB11-1A, *Haliaceae* sp. strain BC5-1, and the BD1-7 clade bacterium

strain SB11-3 were cultivated in artificial marine medium at 21 °C as described earlier (23).

Substrates were added at the following concentrations: cholesterol 1 mM, sodium palmitate

1.0 mM or 1.7 mM, succinate 7.0 mM, glucose 5 mM. Starter cultures were grown on 10 mM

succinate or 1 mM cholesterol. Cholesterol and other sterol substrates were added as solids

and 1% (w/v) methyl-β-cyclodextrin was added to the medium, when required, prior to

autoclaving. Substrate removal was monitored by gas chromatography-coupled mass

spectrometry (GC-MS) of organic extracts of acidified culture supernatants as described

earlier (62). Nitrogen-limiting conditions were achieved by reducing the concentration of

ammonium chloride in the medium from 18.70 mM to 0.93 mM. Under N-excess conditions

the ratio of carbon to nitrogen in the growth medium was around 1.5:1, while it was 29:1

under N-limiting conditions. To limit nitrogen transfer between cultures, starter cultures were

washed twice with phosphate-buffered saline (PBS) without nitrogen. Growth was measured

as protein, using the bicinchoninic acid assay (Thermo Scientific) after hot alkaline cell lysis.

Mycobacterium tuberculosis strain Erdmann (*Mtb*) was grown in 7H9 medium

supplemented with 0.5% tyloxapol and 0.5 mM cholesterol at 37 °C. If required 0.5 mM

sodium palmitate were spiked into growing cultures after 5 days. Growth was measured as

optical density at 600 nm. To test effects of nitric oxide (NO) stress and iron limitation,

cholesterol-grown *Mtb* was harvested and washed in 7H9 medium without carbon source,

and cells were suspended in the same medium containing 0.5 mM cholesterol and/or 0.5 mM

palmitate. 0.5 mM of the NO releasing agent spermine-NONOate was added from a 10 mM

stock in 0.01 M NaOH at the beginning of the experiment and again after 16 h of incubation.

Spermine was added at the same concentration to control cultures. The iron chelating agent

deferoxamine was added at 0.5 mM from a 10 mM stock at the beginning of the experiment.

Biosample IDs and references to cholesterol degradation capability of all strains used

in this study are summarized in **supplemental Tab. S1**.

Ammonium quantification

Ammonium concentrations were quantified using indophenol blue. Culture supernatant

samples were diluted 25- to 500-fold in distilled water and 200 µl aliquots were distributed

into 96-well plates in triplicates. 8 µl of reagent (10% w/v phenol in ethanol), 8 µl catalyst

(0.5% w/v nitroprusside in water) and 20 µl oxidizing solution (20% w/v trisodium citrate and

1% w/v sodium hydroxide solution, mixed with 5% sodium hypochlorite 4:1 v/v) were added

and plates were incubated at 37 °C for 1-2 h in the dark. Absorbance was measured at 630

nm. Ammonium concentrations were calculated using a standard curve with 0-2 mg ml^{-1}

ammonium chloride. Alternatively, semi-quantitative ammonium test strips (Quantofix®,

Macherey-Nagel, Germany) were used for instant ammonium quantification.

Lipid extraction and analysis

For lipid extraction, cells were harvested, washed twice with distilled water and frozen at -

80 °C. *Mtb* cells were autoclaved before freezing. Cell pellets were lyophilized and their dry

weight was determined by weighing the pellets. The pellets were then suspended in 1 ml

distilled water and broken by bead beating (four rounds of 60 sec at 4.5, 5.0, 5.5, and 6.0

speed setting, respectively, in a FastPrep-24 bead beater; MP Biomedicals, Solon, OH).

Between rounds samples were incubated on ice for 5 min. Neutral lipids were extracted with

chloroform-methanol (2:1, v/v, 1% acetic acid), dried under nitrogen and suspended in

chloroform with 1% acetic acid. If required cholestane was added as internal standard from a

10 mM stock solution in chloroform prior to organic extraction for normalization and

quantification of cholesterol and FAMEs. Aliquots of neutral lipid extracts were analyzed by

thin layer chromatography (TLC) on 60 F$_{254}$ silica gel plates (Merck) using hexane/diethyl-

ether/acetic-acid (90:8:1, v/v/v) as a solvent system. Cholesterol palmitate, cholesterol, and

tripalmitin dissolved in chloroform were used as standards. The plates were dried, dabbed

with 10% cupric sulfate in 8% phosphoric acid solution and developed at 200 °C for 5-10 min.

Native neutral lipid extracts from non-lyophilized cell pellets of strain RHA1 grown under

nitrogen limiting conditions with cholesterol or succinate were also analyzed using a GC-

coupled flame ionization detector (FID). For this, 20 ml culture aliquots were harvested,

washed twice and frozen at -80 °C at different time points after nitrogen was depleted from

the medium. Neutral lipids were extracted with tert-butyl ether:methanol (10:3, v/v), followed

by two extractions with hexane. Combined organic extracts were dried under nitrogen and

suspended in chloroform. Neutral lipid extracts were analyzed by gas chromatography on an

Agilent 8890 series GC system equipped with a J&W Select Biodiesel GC Column (15 m,

0.32 mm, 0.10 µm), coupled to a flame ionization detector (FID). GC-injector and detector

temperatures were set to 350 °C, and 400 °C, respectively. 1 µl sample volumes were

injected with an initial purge flow rate of 50 ml min^{-1}, which subsequently decreased to 20 ml

min^{-1} after 2 min, and a septum purge flow rate of 3 ml min^{-1}. The helium flow rate was 8.09

ml min^{-1}. The oven was set to an initial temperature of 40 °C for 2 min with an equilibration

time of 0.5 min. The temperature increased by 25 °C min^{-1} and was held at 170 °C for 2 min,

then increased by 15 °C min^{-1} to 210 °C, then by 5 °C min^{-1} to 250 °C and held for 5 min. The

temperature finally increased by 10 °C min^{-1} to 400 °C and held for 2 min. The FID gas flow

settings were set to 400 ml min^{-1} air flow, 30 ml min^{-1} H$_2$ fuel flow and 25 ml min^{-1} makeup

flow (N$_2$). Cholesteryl palmitate, cholesteryl heptadecanoate, and tripalmitin TAG dissolved in

chloroform were used as standards. CE and TAG peaks were normalized with the internal

standard cholestane for quantification.

Isolation of lipid droplets

Lipid droplets were isolated as previously described (63, 64). Briefly, RHA1 cultures were

harvested by centrifugation (4,000 x g for 10 min), washed three times with 40 ml Buffer A

(25 mM tricine, 250 mM sucrose, pH 7.8) and suspended in 8 ml buffer A. After incubation on

ice for 20 min, cells were disrupted by passing them three times through a cooled French

pressure cell at 100 MPa. Cell debris was removed by centrifugation at 4,000 × g for 10 min.

Supernatants were loaded into SW40 tubes (Beckman), layered with 2 ml Buffer B (20 mM

HEPES, 100 mM KCl, 2 mM MgCl2, pH 7.4), and centrifuged at 180,000 x g for 1 h at 4 °C.

The lipid droplet fraction on top of the sucrose gradient was transferred to a 1.5 ml Eppendorf

tube and was washed twice with 200 µl Buffer B and finally suspended in 1 ml buffer B. No

pellets were formed during washing of the lipid droplets suggesting that lipid droplets were

not contaminated by membrane particles. Total lipids were extracted with 5 ml chloroform-

acetone (1:1, v/v). The organic phase was dried under nitrogen and lipids were suspended in

1 ml chloroform with 1% acetic acid. Aliquots were analyzed by TLC as described above.

Lipid fractionation

For analyzing the fatty acid profile of cellular lipids and lipid droplets, lipid extracts were

fractionated stepwise. Total lipid extracts derived from lipid droplets were first run over a

silica column equilibrated with chloroform and neutral lipid fractions were obtained by eluting

with 6 column volumes chloroform with 1% acetic acid. Glycolipid fractions were

subsequently eluted with 6 column volumes acetone (not analyzed further) followed by

phospholipid elution with 7.5 column volumes methanol. Neutral lipids from lipid droplets and

cellular lipid extractions were further fractionated by silica column chromatography. Neutral

lipid samples were dried, resuspended in hexane and poured on a silica column equilibrated

with the same solvent. 5 different fractions were obtained by eluting sequentially with 5

column volumes hexane, 6 column volumes hexane/diethyl ether (99:1, v/v), 5 column

volumes hexane/diethyl ether (95:5, v/v), 5 column volumes hexane/diethyl ether (92:8, v/v),

and 6 column volumes chloroform with 1% acetic acid. Fractions were dried under nitrogen

and lipids were suspended in 1 ml chloroform with 1% acetic acid. If required, aliquots were

analyzed by TLC as described above.

Alkaline transesterification with methanol and lipid quantification

Dried cellular lipid extracts or individual lipid fractions were subjected to alkaline

transesterification with methanol modified after (65). For this, dried lipids were resuspended

in methanol/toluene (1:1, v/v), mixed with an equal volume of methanolic KOH and incubated

for 1 h at 50 °C. Phospholipid fractions were incubated for 15 min at 30 °C. Free sterols and

fatty acid methyl-esters were extracted twice with 4 volumes hexane/chloroform (4:1, v/v)

from 0.15 volumes acetic acid (1 M) and 1 volume water. The organic phase was dried under

nitrogen and lipids were suspended in 1 ml chloroform with 1% acetic acid. Aliquots were

analyzed by GC-MS as described earlier for FAMEs and free cholesterol (62) and for wax

esters (32). Cholesterol and fatty acid methyl-esters (FAMEs) were identified by comparison

to authentic standards as well as matching against the NIST database (National Institute of

Standards and Technology, v11). To quantify total neutral lipids, unfractionated neutral lipid

extracts were subjected to transesterification and FAMEs were normalized on cholestane. To

quantify SEs in cellular lipid extracts, fractions 1 (containing the internal standard cholestane)

and 2 (containing steryl esters) were combined prior to transesterification. This allowed

normalisation and quantification of free cholesterol derived from transesterification. The

percentage of CEs per extracted cellular dry weight was estimated by calculating the total

weight of CEs from CE containing fractions and dividing it by the dry weight of the respective

extracted cell pellet. For this, the cholesterol concentration in transesterified fractions was

calculated using a cholesterol-cholestane standard curve. Subsequently, this was

transformed into the total weight of CEs using the molecular masses of all detectable CEs

weighted on the peak area ratio of the respective detected FAMEs (C15:0, C16:1, C16:0,

C17:1, C17:0, C18:1, C18:0). To estimate ratios of neutral lipid classes in lipid droplets,

cholestane and methyl-nonadecanoate were added as internal standards to each fraction

prior to transesterification from 10 mM stock solutions in chloroform. Free cholesterol (in

fraction 5) and cholesterol released from transesterification of fraction 2 were normalized on

cholestane and FAMEs derived from transesterification of CEs and TAGs were normalized

on methyl-nonadecanoate before calculating their molar ratios.

Fluorescence microscopy

To localize accumulating neutral lipids in strain RHA1, we added the fluorescent dye Nile

Red to growing cholesterol cultures under N-limiting or N-excess conditions without

cyclodextrin. Nile red was added to the medium after autoclaving at a final concentration of

0.5 µg ml^{-1} from a sterile stock solution in DMSO (0.3 mg ml^{-1}). Nitrogen and cholesterol

depletion were confirmed as described above. Cells from both cultures were examined four

days after reaching stationary phase. Cells were washed twice and resuspended in sterile

PBS before microscopy. Slides were examined on a Zeiss Axio Imager M1 fluorescence

microscope equipped with a Zeiss HBO 103 W/2 high-pressure mercury lamp and an

AxioCam MRm camera in phase contrast, differential interference contrast or fluorescence

mode. Fluorescence images were shot using excitation and emission wavelengths of 550 nm

and 605 nm, respectively. All Images were processed using the Zeiss AxioVision software

(40 V 4.6.3.0).

To further analyze the localization of accumulating cholesterol within cells of RHA1

under nitrogen-limiting conditions, we performed a cell suspension experiment with

fluorescently labeled cholesterol (cholesterol-BODIPY) as substrate under N-limiting

conditions followed by laser scanning confocal microscopy. For that, cholesterol-grown cells

were washed two times with PBS without nitrogen. Depletion of cholesterol in the starter

culture was confirmed by organic extraction and GC-MS analysis. Cells were suspended in

1/5 of the original volume of M9 medium without nitrogen containing 0.5 mM palmitate, 0.5

mM fluorescently labeled cholesterol (cholesterol-BODIPY) and 1% (w/v) methyl-β-

cyclodextrin and incubated for 48 h at 200 rpm and 30 °C. Cells were washed twice with PBS

without nitrogen containing 0.02% (w/v) cholate and 0.02% (v/v) tylaxopol and suspended in

PBS. Cells were analyzed using a laser scanning confocal microscope (Olympus Fluoview

FV1000-Inverted). A small aliquot of cells was broken by ultrasonication (four rounds of 20

sec using a Microsom cell disruptor XL set to 5; Mandel scientific Co. Ltd., Guelph, ON,

Canada) and neutral lipid extracts were analyzed by TLC. Undeveloped plates were

analyzed using a GE Typhoon trio + imager with excitation wavelength of 488 nm and an

emission filter of 520 nm. Pre-tests with unlabelled cholesterol and palmitate showed that CE

accumulation in RHA1 cell suspensions started within 22 h of incubation and continued at

least until 95 h of incubation (data not shown).

Acknowledgments

We thank Fanrui Meng and Carl Strittmatter for help with fluorescence microscopy. This study was funded by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada.

Data availability

Draft genomes of *Mycobacterium* strain BC8-1, *Zhongshania* sp. strain SB11-1A, *Haliaceae* sp. strain BC5-1, and the BD1-7 clade bacterium strain SB11-3 have been deposited in the European Nucleotide Archive (ENA) under the BioProject number PRJEB30766.

References

1. **J Brigham C**, **Kurosawa K**, **Rha C**, **Sinskey AJ**. 2011. Bacterial carbon storage to value added products. J Microbial Biochem Technol **s3**:1–13.

2. **Alvarez HM**, **Steinbüchel A**. 2002. Triacylglycerols in prokaryotic microorganisms. Appl Microbiol Biotechnol **60**:367–376.

3. **Russell NJ**, **Volkman JK**. 1980. The effect of growth temperature on wax ester composition in the psychrophilic bacterium *Micrococcus cryophilus* ATCC 15174. J Gen Microbiol **118**:131–141.

4. **Kalscheuer R**, **Stöveken T**, **Malkus U**, **Reichelt R**, **Golyshin PN**, **Sabirova JS**, **Ferrer M**, **Timmis KN**, **Steinbüchel A**. 2007. Analysis of storage lipid accumulation in *Alcanivorax borkumensis*: Evidence for alternative triacylglycerol biosynthesis routes in bacteria. J Bacteriol **189**:918–928.

5. **Kalscheuer R**, **Steinbüchel A**. 2003. A novel bifunctional wax ester synthase/acyl-CoA:diacylglycerol acyltransferase mediates wax ester and triacylglycerol biosynthesis in *Acinetobacter calcoaceticus* ADP1. J Biol Chem **278**:8075–8082.

6. **Garay LA**, **Boundy-Mills KL**, **German JB**. 2014. Accumulation of high-value lipids in single-cell microorganisms: a mechanistic approach and future perspectives. J Agric Food Chem **62**:2709–2727.

7. **Röttig A**, **Strittmatter CS**, **Schauer J**, **Hiessl S**, **Poehlein A**, **Daniel R**, **Steinbüchel A**. 2016. Role of wax ester synthase/acyl coenzyme a:diacylglycerol acyltransferase in oleaginous *Streptomyces* sp. strain G25. Appl Environ Microbiol **82**:5969–5981.

8. **Röttig A**, **Steinbüchel A**. 2013. Acyltransferases in Bacteria. Microbiol Mol Biol Rev **77**:277–321.

9. **Alvarez HM**, **Herrero OM**, **Silva RA**, **Hernández MA**, **Lanfranconi MP**, **Villalba MS**. 2019. Insights into the metabolism of oleaginous *Rhodococcus* spp. Appl Environ Microbiol **85**:8075–12.

10. **Alvarez HM**, **Mayer F**, **Fabritius D**, **Steinbüchel A**. 1996. Formation of intracytoplasmic lipid inclusions by *Rhodococcus opacus* strain PD630. Arch Microbiol **165**:377–386.

11. **Waltermann M**, **Luftmann H**, **Baumeister D**, **Kalscheuer R**, **Steinbüchel A**. 2000. *Rhodococcus opacus* strain PD630 as a new source of high-value single-cell oil? Isolation and characterization of triacylglycerols and other storage lipids. Microbiology **146** :1143–1149.

12. **Amara S**, **Seghezzi N**, **Otani H**, **Diaz-Salazar C**, **Liu J**, **Eltis LD**. 2016. Characterization of key triacylglycerol biosynthesis processes in rhodococci. Sci Rep **6**:24985.

13. **Daniel J**, **Deb C**, **Dubey VS**, **Sirakova TD**, **Abomoelak B**, **Morbidoni HR**, **Kolattukudy PE**. 2004. Induction of a novel class of diacylglycerol acyltransferases and triacylglycerol accumulation in *Mycobacterium tuberculosis* as it goes into a dormancy-like state in culture. J Bacteriol **186**:5017–5030.

14. **Bacon J**, **Dover LG**, **Hatch KA**, **Zhang Y**, **Gomes JM**, **Kendall S**, **Wernisch L**, **Stoker NG**, **Butcher PD**, **Besra GS**, **Marsh PD**. 2007. Lipid composition and

633   transcriptional response of *Mycobacterium tuberculosis* grown under iron-limitation in
634   continuous culture: identification of a novel wax ester. Microbiology **153**:1435–1444.

635   15.   **Stehr M**, **Elamin AA**, **Singh M**. 2013. Lipid inclusions in mycobacterial infections.
636   Tuberculosis - Current Issues in Diagnosis and Management, Bassam H. Mahboub
637   and Mayank G. Vats, IntechOpen, doi: 10.5772/54526

638   16.   **Garton NJ**, **Christensen H**, **Minnikin DE**, **Adegbola RA**, **Barer MR**. 2002.
639   Intracellular lipophilic inclusions of mycobacteria in vitro and in sputum. Microbiology
640   **148**:2951–2958.

641   17.   **Zhang C**, **Liu P**. 2017. The lipid droplet: A conserved cellular organelle. Protein &
642   Cell **8**:796–800.

643   18.   **Ruggles KV**, **Turkish A**, **Sturley SL**. 2013. Making, baking, and breaking: The
644   synthesis, storage, and hydrolysis of neutral lipids. Annu Rev Nutr **33**:413–451.

645   19.   **Korber M**, **Klein I**, **Daum G**. 2017. Steryl ester synthesis, storage and hydrolysis: A
646   contribution to sterol homeostasis. Biochim Biophys Acta **1862**:1534–1545.

647   20.   **Wei JH**, **Yin X**, **Welander PV**. 2016. Sterol synthesis in diverse bacteria. Front
648   Microbiol **7**:990.

649   21.   **Lee AK**, **Banta AB**, **Wei JH**, **Kiemle DJ**, **Feng J**, **Giner J-L**, **Welander PV**. 2018. C-
650   4 sterol demethylation enzymes distinguish bacterial and eukaryotic sterol synthesis.
651   Proc Natl Acad Sci USA **115**:5884–5889.

652   22.   **Bergstrand LH**, **Cardenas E**, **Holert J**, **Van Hamme JD**, **Mohn WW**. 2016.
653   Delineation of steroid-degrading microorganisms through comparative genomic
654   analysis. MBio **7**:e00166–16.

655   23.   **Holert J**, **Cardenas E**, **Bergstrand LH**, **Zaikova E**, **Hahn AS**, **Hallam SJ**, **Mohn WW**.
656   2018. Metagenomes reveal global distribution of bacterial steroid catabolism in
657   natural, engineered, and host environments. MBio **9**: e02345-17.

658   24.   **Crowe AM**, **Casabon I**, **Brown KL**, **Liu J**, **Lian J**, **Rogalski JC**, **Hurst TE**, **Snieckus
659   V**, **Foster LJ**, **Eltis LD**. 2017. Catabolism of the last two steroid rings in
660   *Mycobacterium tuberculosis* and other bacteria. MBio **8**:e00321–17.

661   25.   **van der Geize R**, **Yam K**, **Heuser T**, **Wilbrink MH**, **Hara H**, **Anderton MC**, **Sim E**,
662   **Dijkhuizen L**, **Davies JE**, **Mohn WW**, **Eltis LD**. 2007. A gene cluster encoding
663   cholesterol catabolism in a soil actinomycete provides insight into *Mycobacterium
664   tuberculosis* survival in macrophages. Proc Natl Acad Sci USA **104**:1947–1952.

665   26.   **Wipperman MF**, **Sampson NS**, **Thomas ST**. 2014. Pathogen roid rage: cholesterol
666   utilization by *Mycobacterium tuberculosis*. Crit Rev Biochem Mol Biol **49**:269–293.

667   27.   **Alvarez HM**. 2016. Triacylglycerol and wax ester-accumulating machinery in
668   prokaryotes. Biochimie **120**:28–39.

669   28.   **Barney BM**, **Wahlen BD**, **Garner E**, **Wei J**, **Seefeldt LC**. 2012. Differences in
670   substrate specificities of five bacterial wax ester synthases. Appl Environ Microbiol
671   **78**:5734–5745.

672   29.   **Kalscheuer R**, **Luftmann H**, **Steinbüchel A**. 2004. Synthesis of novel lipids in
673   *Saccharomyces cerevisiae* by heterologous expression of an unspecific bacterial
674   acyltransferase. Appl Environ Microbiol **70**:7119–7125.

30.   **Schubert K**, **Kaufmann G**. 1965. Bildung von Sterinestern in der Bakterienzelle.
Biochim Biophys Acta **106**:592–597.

31.   **Schubert K**, **Kaufmann G**, **Hörhold C**. 1969. Veresterung von Sterinen mit
Fettsäuren und mit Bernsteinsäure in *Mykobakterien*. Biochim Biophys Acta **176**:163–
169.

32.   **Round J**, **Roccor R**, **Li S-N**, **Eltis LD**. 2017. A fatty acyl-CoA reductase promotes
wax ester accumulation in *Rhodococcus jostii* RHA1. Appl Environ Microbiol
83:e00902–17–12.

33.   **Hernández MA**, **Mohn WW**, **Martínez E**, **Rost E**, **Alvarez AF**, **Alvarez HM**. 2008.
Biosynthesis of storage compounds by *Rhodococcus jostii* RHA1 and global
identification of genes involved in their metabolism. BMC Genomics **9**:600.

34.   **Peyron P**, **Vaubourgeix J**, **Poquet Y**, **Levillain F**, **Botanch C**, **Bardou F**, **Daffé M**,
Emile J-F, **Marchou B**, **Cardona P-J**, **de Chastellier C**, **Altare F**. 2008. Foamy
macrophages from tuberculous patients' granulomas constitute a nutrient-rich
reservoir for *M. tuberculosis* persistence. PLoS Pathog **4**:e1000204.

35.   **Russell DG**, **Cardona P-J**, **Kim M-J**, **Allain S**, **Altare F**. 2009. Foamy macrophages
and the progression of the human tuberculosis granuloma. Nat Immunol **10**:943–948.

36.   **VanderVen BC**, **Fahey RJ**, **Lee W**, **Liu Y**, **Abramovitch RB**, **Memmott C**, **Crowe
AM**, **Eltis LD**, **Perola E**, **Deininger DD**, **Wang T**, **Locher CP**, **Russell DG**. 2015.
Novel inhibitors of cholesterol degradation in *Mycobacterium tuberculosis* reveal how
the bacterium's metabolism is constrained by the intracellular environment. PLoS
Pathog **11**:e1004679.

37.   **Pandey AK**, **Sassetti CM**. 2008. Mycobacterial persistence requires the utilization of
host cholesterol. Proc Natl Acad Sci USA **105**:4376–4380.

38.   **Muñoz Elías E**, **Upton A**, **Cherian J**, **McKinney JD**. 2006. Role of the methylcitrate
cycle in *Mycobacterium tuberculosis* metabolism, intracellular growth, and virulence.
Mol Microbiol **60**:1109–1122.

39.   **Nazarova EV**, **Montague CR**, **La T**, **Wilburn KM**, **Sukumar N**, **Lee W**, **Caldwell S**,
Russell DG, **VanderVen BC**. 2017. Rv3723/LucA coordinates fatty acid and
cholesterol uptake in *Mycobacterium tuberculosis*. Elife **6**:678.

40.   **Daniel J**, **Maamar H**, **Deb C**, **Sirakova TD**, **Kolattukudy PE**. 2011. *Mycobacterium
tuberculosis* uses host triacylglycerol to accumulate lipid droplets and acquires a
dormancy-like phenotype in lipid-loaded macrophages. PLoS Pathog **7**:e1002093.

41.   **Low KL**, **Rao PSS**, **Shui G**, **Bendt AK**, **Pethe K**, **Dick T**, **Wenk MR**. 2009.
Triacylglycerol utilization is required for regrowth of in vitro hypoxic nonreplicating
Mycobacterium bovis bacillus Calmette-Guerin. J Bacteriol **191**:5037–5043.

42.   **Deb C**, **Lee C-M**, **Dubey VS**, **Daniel J**, **Abomoelak B**, **Sirakova TD**, **Pawar S**,
Rogers L, **Kolattukudy PE**. 2009. A novel in vitro multiple-stress dormancy model
for *Mycobacterium tuberculosis* generates a lipid-loaded, drug-tolerant, dormant
pathogen. PLoS ONE **4**:e6077.

43.   **Sirakova TD**, **Deb C**, **Daniel J**, **Singh HD**, **Maamar H**, **Dubey VS**, **Kolattukudy PE**.
2012. Wax ester synthesis is required for *Mycobacterium tuberculosis* to enter in vitro
dormancy. PLoS ONE **7**:e51641.

44.   **Sirakova TD**, **Dubey VS**, **Deb C**, **Daniel J**, **Korotkova TA**, **Abomoelak B**,
719   **Kolattukudy PE**. 2006. Identification of a diacylglycerol acyltransferase gene
720   involved in accumulation of triacylglycerol in *Mycobacterium tuberculosis* under stress.
721   Microbiology **152**:2717–2725.

722   45.   **Murphy DJ**. 2011. The dynamic roles of intracellular lipid droplets: from archaea to
723   mammals. Protoplasma **249**:541–585.

724   46.   **Waltermann M**, **Steinbüchel A**. 2005. Neutral lipid bodies in prokaryotes: Recent
725   insights into structure, formation, and relationship to eukaryotic lipid depots. J
726   Bacteriol **187**:3607–3619.

727   47.   **Bartz R**, **Li W-H**, **Venables B**, **Zehmer JK**, **Roth MR**, **Welti R**, **Anderson RGW**, **Liu**
728   **P**, **Chapman KD**. 2007. Lipidomics reveals that adiposomes store ether lipids and
729   mediate phospholipid traffic. J Lipid Res **48**:837–847.

730   48.   **Faletrov Y**, **Brzostek A**, **Plocinska R**, **Dziadek J**, **Rudaya E**, **Edimecheva I**,
731   **Shkumatov V**. 2016. Uptake and metabolism of fluorescent steroids by
732   mycobacterial cells. Steroids **117**:29–37.

733   49.   **Griffin JE**, **Pandey AK**, **Gilmore SA**, **Mizrahi V**, **McKinney JD**, **Bertozzi CR**,
734   **Sassetti CM**. 2012. Cholesterol catabolism by *Mycobacterium tuberculosis* requires
735   transcriptional and metabolic adaptations. Chem Biol **19**:218–227.

736   50.   **Russell DG**, **VanderVen BC**, **Lee W**, **Abramovitch RB**, **Kim M-J**, **Homolka S**,
737   **Niemann S**, **Rohde KH**. 2010. *Mycobacterium tuberculosis* wears what it eats. Cell
738   Host Microbe **8**:68–76.

739   51.   **Singh A**, **Crossman DK**, **Mai D**, **Guidry L**, **Voskuil MI**, **Renfrow MB**, **Steyn AJC**.
740   2009. *Mycobacterium tuberculosis* WhiB3 maintains redox homeostasis by regulating
741   virulence lipid anabolism to modulate macrophage response. PLoS Pathog
742   **5**:e1000545–16.

743   52.   **Hauschild P**, **Röttig A**, **Madkour MH**, **Al-Ansari AM**, **Almakishah NH**, **Steinbüchel**
744   **A**. 2017. Lipid accumulation in prokaryotic microorganisms from arid habitats. Appl
745   Microbiol Biotechnol **101**: 2203–2216.

746   53.   **Alvarez AF**, **Alvarez HM**, **Kalscheuer R**, **Waltermann M**, **Steinbüchel A**. 2008.
747   Cloning and characterization of a gene involved in triacylglycerol biosynthesis and
748   identification of additional homologous genes in the oleaginous bacterium
749   *Rhodococcus opacus* PD630. Microbiology **154**:2327–2335.

750   54.   **Lanfranconi MP**, **Alvarez AF**, **Alvarez HM**. 2015. Identification of genes coding for
751   putative wax ester synthase/diacylglycerol acyltransferase enzymes in terrestrial and
752   marine environments. AMB Express **5**:42.

753   55.   **Deb C**, **Daniel J**, **Sirakova TD**, **Abomoelak B**, **Dubey VS**, **Kolattukudy PE**. 2006. A
754   novel lipase belonging to the hormone-sensitive lipase family induced under
755   starvation to utilize stored triacylglycerol in *Mycobacterium tuberculosis*. J Biol Chem
756   **281**:3866–3875.

757   56.   **Desmond E**, **Gribaldo S**. 2009. Phylogenomics of sterol synthesis: insights into the
758   origin, evolution, and diversity of a key eukaryotic feature. Genome Biol Evol **1**:364–
759   381.

760   57.   **Wollam J**, **Antebi A**. 2011. Sterol regulation of metabolism, homeostasis, and
761   development. Annu Rev Microbiol **80**:885–916.

58. **Weingärtner O**, **Böhm M**, **Laufs U**. 2008. Controversial role of plant sterol esters in
the management of hypercholesterolaemia. Eur Heart J **30**:404–409.

59. **Lichtenstein AH**, **Deckelbaum RJ**. 2001. AHA Science Advisory. Stanol/sterol ester-
containing foods and blood cholesterol levels. A statement for healthcare
professionals from the Nutrition Committee of the Council on Nutrition, Physical
Activity, and Metabolism of the American Heart Association. Circulation **103**:1177–
1179.

60. **Bacchetti T**, **Masciangelo S**, **Bicchiega V**, **Bertoli E**, **Ferretti G**. 2011. Phytosterols,
phytostanols and their esters: from natural to functional foods. Mediterr J Nutr Metab
4:165–172.

61. **He W-S**, **Zhu H**, **Chen Z-Y**. 2018. Plant sterols: Chemical and enzymatic structural
modifications and effects on their cholesterol-lowering activity. J Agric Food Chem
66:3047–3062.

62. **Casabon I**, **Crowe AM**, **Liu J**, **Eltis LD**. 2013. FadD3 is an acyl-CoA synthetase that
initiates catabolism of cholesterol rings C and D in actinobacteria. Mol Microbiol
87:269–283.

63. **Ding Y**, **Yang L**, **Zhang S**, **Wang Y**, **Du Y**, **Pu J**, **Peng G**, **Chen Y**, **Zhang H**, **Yu J**,
Hang H, **Wu P**, **Yang F**, **Yang H**, **Steinbüchel A**, **Liu P**. 2012. Identification of the
major functional proteins of prokaryotic lipid droplets. J Lipid Res **53**:399–411.

64. **Ding Y**, **Zhang S**, **Yang L**, **Na H**, **Zhang P**, **Zhang H**, **Wang Y**, **Chen Y**, **Yu J**, **Huo C**,
Xu S, **Garaiova M**, **Cong Y**, **Liu P**. 2013. Isolating lipid droplets from multiple species.
Nat Protoc **8**:43–51.

65. **Ichihara K**, **Yamaguchi C**, **Nishijima H**, **Saito K**. 2003. Preparation of FAME from
sterol esters. J Amer Oil Chem Soc **80**:833–834.

Figure legends

Fig. 1: Schematic representation of triacylglycerols (TAGs) and wax esters (WEs) as neutral lipid storage compounds in bacteria and of a cholesterol-ester. Structures of the major animal, plant, and fungal sterols cholesterol, β-sitosterol, and ergosterol.

Fig. 2: Growth of strain RHA1 under nitrogen-limiting and nitrogen-excess conditions. **A)** at early stationary phase (around 100 h). **B)** TLC analysis of neutral lipid extracts from early stationary phase N-limited (-N) and N-excess (+N) cultures grown with different carbon sources. Abbreviations: CP, cholesterol palmitate; C, cholesterol; TAG, triacylglycerols; Ph, phytosterol mix; S, succinate; P, pyruvate; G, glucose; NH_4^+, N replenished in stationary-phase C-N culture. The volume of neutral lipid extracts applied to TLC plates was normalized on the extracted cellular dry weight of each culture. Red arrows indicate standard spots. Results were obtained on two TLC plates which were treated exactly the same way throughout the experiment. The results of one representative experiment out of several biological and technical replicates are shown for each substrate.

Fig. 3: Growth of RHA1 on 1 mM cholesterol under nitrogen-limiting conditions **(A)** and nitrogen-excess conditions **(B)**. The volume of neutral lipid extracts applied to TLC plates was normalized on the basis of culture biomass as protein. One of two reproducible experiments is shown.

Fig. 4: GC-FID analysis of accumulating neutral lipids in nitrogen-limited RHA1 cultures grown with cholesterol or succinate. **(A)** GC-FID chromatograms of authentic cholesteryl palmitate (C16:0) and cholesteryl heptadecanoate (C17:0) standards (blue line) and of a trimpalmitin TAG (C16:0$_3$) standard (orange line) and of neutral lipid extracts of cholesterol- and succinate-grown cells harvested 24 h after nitrogen depletion. Seven accumulating CE species were identified either by comparison to the standards or based on the proximity of their respective peaks to the standard peaks and their absence in succinate grown cells. Identified and predicted CEs are labelled with blue arrows, identified and predicted TAG species are labelled with orange arrows. **(B)** Ratio of accumulating CEs (blue bars) and TAGs (orange bars) in cholesterol-grown nitrogen-limited RHA1 cultures over time starting at nitrogen depletion (0 h). Peak areas of the seven CE- and nine TAG-peaks shown in Fig. 4A were normalized on the peak area of the internal standard cholestane (not shown) before calculating their respective sums. Average values of three biological replicates and their standard deviations are shown.

Fig. 5: Identification of SEs accumulating in N-limited RHA1 cultures. GC-MS analysis of
transesterified fraction 2 (black peaks) and untreated fraction 2 (grey peaks) of neutral lipid
extracts of nitrogen-limited cultures grown with **(A)** cholesterol and **(B)** phytosterols.
Transesterification of SEs produced free cholesterol and the free phytosterols β-sitosterol, β-
sitostanol, and campesterol, as well as seven fatty acid methyl esters of C15:0, C16:1,
C16:0, C17:1, C17:0, C18:0 and C18:1. One representative GC-MS chromatogram is shown.
(C) Fatty acid profile of transesterified fraction 2 derived from nine cholesterol-grown cultures
grown under nitrogen-limiting conditions from five independent experiments harvested
approximately 24h after nitrogen depletion. Odd-numbered fatty acids are coloured blue and
even-numbered fatty acids are coloured red

Fig. 6: Isolation of lipid droplets from strain RHA1 grown with cholesterol (**C**) or palmitate (**P**)
or a mixture of cholesterol plus palmitate (**C+P**) under nitrogen-limiting conditions. **A)** TLC
analysis of neutral lipids derived from total lipid extracts from isolated lipid droplets.
Abbreviations as in previous figures. **B)** Molar ratio of TAGs, CEs and free cholesterol in lipid
droplets derived from cholesterol, cholesterol plus palmitate and palmitate grown cells. **C)**
Ratio of even to odd-numbered fatty acids in CEs and TAGs in lipid droplets and the
respective contributing fatty acid chain lengths. The results of two independent experiments
(I and II) are shown in **B)** and **C)** and of experiment II in **A)**.

Fig. 1: Schematic representation of triacylglycerols (TAGs) and wax esters (WEs) as neutral

lipid storage compounds in bacteria and of a cholesterol-ester. Structures of the major

animal, plant, and fungal sterols cholesterol, β-sitosterol, and ergosterol.

Fig. 2: Growth of strain RHA1 under nitrogen-limiting and nitrogen-excess conditions. **A)** at

early stationary phase (around 100 h). **B)** TLC analysis of neutral lipid extracts from early

stationary phase N-limited (-N) and N-excess (+N) cultures grown with different carbon

sources. Abbreviations: CP, cholesterol palmitate; C, cholesterol; TAG, triacylglycerols; Ph,

phytosterol mix; S, succinate; P, pyruvate; G, glucose; NH_4^+, N replenished in stationary-

phase C-N culture. The volume of neutral lipid extracts applied to TLC plates was normalized

on the extracted cellular dry weight of each culture. Red arrows indicate standard spots.

Results were obtained on two TLC plates which were treated exactly the same way

throughout the experiment. The results of one representative experiment out of several

biological and technical replicates are shown for each substrate.

863
864 **Fig. 3:** Growth of RHA1 on 1 mM cholesterol under nitrogen-limiting conditions **(A)** and
865 nitrogen-excess conditions **(B)**. The volume of neutral lipid extracts applied to TLC plates
866 was normalized on the basis of culture biomass as protein. One of two reproducible
867 experiments is shown
868
869

Fig. 4: GC-FID analysis of accumulating neutral lipids in nitrogen-limited RHA1 cultures

grown with cholesterol or succinate. **(A)** GC-FID chromatograms of authentic cholesteryl

palmitate (C16:0) and cholesteryl heptadecanoate (C17:0) standards (blue line) and of a

tripalmitin TAG (C16:0$_3$) standard (orange line) and of neutral lipid extracts of cholesterol-

and succinate-grown cells harvested 24 h after nitrogen depletion. Seven accumulating CE

species were identified either by comparison to the standards or based on the proximity of

their respective peaks to the standard peaks and their absence in succinate grown cells.

Identified and predicted CEs are labelled with blue arrows, identified and predicted TAG

species are labelled with orange arrows. **(B)** Ratio of accumulating CEs (blue bars) and

TAGs (orange bars) in cholesterol-grown nitrogen-limited RHA1 cultures over time starting at

nitrogen depletion (0 h). Peak areas of the seven CE- and nine TAG-peaks shown in Fig. 4A

were normalized on the peak area of the internal standard cholestane (not shown) before

calculating their respective sums. Average values of three biological replicates and their

standard deviations are shown.

Fig. 5: Identification of SEs accumulating in N-limited RHA1 cultures. GC-MS analysis of

transesterified fraction 2 (black peaks) and untreated fraction 2 (grey peaks) of neutral lipid

extracts of nitrogen-limited cultures grown with **(A)** cholesterol and **(B)** phytosterols.

Transesterification of SEs produced free cholesterol and the free phytosterols β-sitosterol, β-

sitostanol, and campesterol, as well as seven fatty acid methyl esters of C15:0, C16:1,

C16:0, C17:1, C17:0, C18:0 and C18:1. One representative GC-MS chromatogram is shown.

(C) Fatty acid profile of transesterified fraction 2 derived from nine cholesterol-grown cultures

grown under nitrogen-limiting conditions from five independent experiments harvested

approximately 24h after nitrogen depletion. Odd-numbered fatty acids are coloured blue and

even-numbered fatty acids are coloured red

Fig. 6: Isolation of lipid droplets from strain RHA1 grown with cholesterol (**C**) or palmitate (**P**)

or a mixture of cholesterol plus palmitate (**C+P**) under nitrogen-limiting conditions. **A)** TLC

analysis of neutral lipids derived from total lipid extracts from isolated lipid droplets.

Abbreviations as in previous figures. **B)** Molar ratio of TAGs, CEs and free cholesterol in lipid

droplets derived from cholesterol, cholesterol plus palmitate and palmitate grown cells. **C)**

Ratio of even to odd-numbered fatty acids in CEs and TAGs in lipid droplets and the

respective contributing fatty acid chain lengths. The results of two independent experiments

(I and II) are shown in **B)** and **C)** and of experiment II in **A)**.

907