Inter-specific variation of maternal thyroid hormones associate with migration, mode of

development, and captivity: a phylogenetic comparative analysis

Bin-Yan Hsu[1], Veli-Matti Pakanen[2,3], Winnie Boner[4], Tapio Eeva[1], Blandine Doligez[5], Ton G.G.

Groothuis[6], Erkki Korpimäki[1], Toni Laaksonen[1], Asmoro Lelono[6], Pat Monaghan[4], Tom Sarraude[1,6],

Barbara Tschirren[7], Robert L. Thomson[8], Jere Tolvanen[2], Rodrigo A. Vásquez[9], Suvi Ruuskanen[1]

[1] Department of Biology, University of Turku, Finland

[2] Ecology and Genetics Research Unit, University of Oulu, Finland

[3] Department of Biological and Environmental Sciences, University of Gothenburg, Sweden

[4] Institute of Biodiversity, Animal Healthy and Comparative Medicine, University of Glasgow,

Scotland, United Kingdom

[5] Department of Biometry and Evolutionary Biology, CNRS UMR 5558, Université de Lyon 1, France

[6] Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, the

Netherlands

[7] Centre for Ecology and Conservation, University of Exeter, United Kingdom

[8] Fitzpatrick Institute of African Ornithology, University of Cape Town, South Africa

[9] Instituto de Ecologíay Biodiversidad, Departamento de Ciencias Ecológicas, Facultad de Ciencias,

Universidad de Chile, Santiago, Chile

Abstract

Animals' life-history traits vary largely along many continuums across species and several

physiological parameters have been proposed as possible mediators that drive the life-history

variation, such as metabolic rates, glucocorticoids, and oxidative stress. Interestingly, thyroid

hormones (THs), despite closely interacting with these physiological parameters, have never been

considered as such mediators. Because of the crucial effects on embryonic development, THs can

also induce transgenerational plasticity when transferred into developing offspring, for instance, via

egg yolks in birds. In this study, we compiled a unique data set of maternal yolk THs in 34 bird

species across 17 families and 6 orders and analysed them with data of life-history traits along

several life-history continuums. Our phylogenetic mixed models estimated higher deposition of the

two most important forms of THs – T3 and T4 -- in both concentrations and total amounts in

migratory species than resident species, and larger total amounts of T3 in precocial species than in

altricial species. However, maternal THs did not show clear associations with any traits of the pace-

of-life syndrome. When taking environmental factors into account, we found that captive species

also deposited higher TH concentrations and larger amounts in the egg yolks than wild species.

These findings suggest that THs are likely involved in the evolution of life-history variation or vice

versa. Accordingly we proposed many angles for further studies.

Key words: life histories, birds, maternal thyroid hormones, phylogenetic comparative analysis

Introduction

Life-history variation constitutes a large part of biodiversity. Across species, life-history traits are often inter-correlated and present along several continuums. For example, many life-history traits, such as lifespan, co-vary with one another and constitute a continuum representing the 'pace of life' (Ricklefs and Wikelski 2002; Dammhahn et al. 2018). On the one end of this continuum, species exhibit faster growth and development, earlier maturation, but faster senescence and shorter lifespan ("fast" species). On the other end, species have slower growth and development, later maturation and first reproduction, but also live longer ("slow" species). Other life-history continuums, such as the nutrient source for developing embryos (from yolk or via placenta), income versus capital breeders, precocial to altricial developmental modes, and various degrees of migratory tendencies, also exist across various taxa in the animal kingdom.

Identifying the mechanisms underlying the evolution of life history differences has fascinated evolutionary biologists for decades. For example, for the pace-of-life (POL) continuum, metabolic rate has been proposed as a mediator as it exhibits a parallel pattern with fast species usually having higher basal and resting metabolic rates (BMR and RMR), while slow species having lower metabolic rates (e.g. Wikelski et al. 2003; Auer et al. 2018). Some studies also proposed other possible mediators, such as glucocorticoids (Hau et al. 2010), oxidative stress (Vágási et al. 2019), and blood glucose level (Tomasek et al. 2019). We propose a novel prediction that thyroid hormones (THs) could be one of the underlying mechanisms explaining the life-history variation, such as the POL continuum. This prediction is likely because THs have regulatory functions on metabolism (Mullur et al. 2014) and interactions with glucocorticoids (Darras et al. 1996; Kulkami and Buchholz 2012; Cortés et al. 2014), yet it is surprising that no studies have yet explored the possibility.

Not only regulating metabolisms in adult animals, THs are also transferred from mothers to developing offspring in all vertebrates (Ruuskanen and Hsu 2018). Such maternally-derived THs (hereafter "maternal THs") play an indispensable role for early embryonic development (Moog et al. 2017; Darras 2019). In humans, the levels of maternal THs are associated with new-born body mass

and children's IQ (Medici et al. 2013; Korevaar et al. 2016). This suggests the variation in maternal

THs can induce transgenerational phenotypic plasticity, which can constitute an important part of

phenotypic variation also in wild animals (Mousseau and Fox 1998; Moore et al. 2019). Nevertheless,

the ecological and evolutionary significance of maternal THs in wild animals was almost totally

neglected (Ruuskanen and Hsu 2018). Recently, several experimental studies manipulating maternal

THs reported inconsistent effects of yolk TH on nestling traits across species (Ruuskanen et al. 2016a;

Hsu et al. 2017, 2019a). Such species differences intrigue us to consider the possibility that maternal

yolk THs might have co-evolved with life-history differences and therefore species respond to

elevated yolk THs differently.

In this study, we explored the potential role of maternal yolk THs underlying the continuums

of life-history strategies in birds by collecting eggs from 34 species of birds and analysing THs in the

egg yolks. In birds, two maternally derived THs – triiodothyronine (T3) and thyroxine (T4) are

transferred and stored in the egg yolk (Sechman and Bobek 1988; Prati et al. 1992). We asked

whether the inter-specific variation of yolk THs would associate with life-history traits across species,

in the meanwhile taking into account some environmental factors that may potentially contribute to

the variation in yolk THs. The associations between maternal THs and life-history traits, if exist,

probably reflect the involvement of maternal THs in the evolution of life-history differences or vice

versa.

Based on the known functions of THs, we considered three life-history continuums, along

which we hypothesized that maternal yolk THs may be associated with the related life-history traits.

The first is the altricial-precocial developmental spectrum. At the two ends of this life-history

continuum, precocial species and altricial species show distinctive ontogenetic patterns of visual and

locomotory ability (Starck and Ricklefs 1998c), hypothalamic-pituitary-adrenal (HPA) axis (Wada

2008), hypothalamic-pituitary-thyroid (HPT) axis (McNabb 2006; de Groef et al. 2013) and

thermoregulation (Price and Dzialowski 2018). Specifically, in precocial species, the thyroid function

already starts in the middle of incubation and the TH levels in thyroid gland and blood circulation

both peak around hatching. By contrast, in altricial species, the thyroid function only starts days

after hatching, and the blood circulating TH levels slowly increase towards the adult level (McNabb

2006; de Groef et al. 2013). Therefore, we hypothesized that such differences in the mode of

development may be associated with yolk TH levels, either because higher maternal THs are

required for a comparatively advanced development at hatching or because the developmental

modes at least partly determines the transferring process of maternal THs, or both .

Second, we considered the life-history differences between migratory and resident species.

THs have been known to regulate seasonal physiology and reproduction (Dardente et al. 2014;

McNabb and Darras 2015; Nishiwaki-Ohkawa and Yoshimura 2016) and also moulting and feather

regeneration (Vézina et al. 2009; Pérez et al. 2018). Several studies have also considered THs as part

of the hormonal mechanism underlying migration (Pérez et al. 2016). We thus hypothesized that

maternal TH levels may be associated with species' migratory status.

Third, we considered the slow-fast pace-of-life (POL) continuum (Ricklefs and Wikelski 2002;

Dammhahn et al. 2018). As introduced above, metabolic rates (Wikelski et al. 2003; Auer et al.

2018), glucocorticoids (Hau et al. 2010), and oxidative stress (Vágási et al. 2018) have been proposed

to mediate the POL-related life-history variation. Interestingly, THs appear to have associations with

all of them. It is well known that THs have regulatory function on metabolism (Mullur et al. 2014),

and in wild birds, within-species studies have confirmed positive correlations between blood THs

and BMR and RMR (e.g. Chastel et al. 2003; Elliott et al 2013; Welcker et al. 2013). THs also have

cross-talks with the HPA axis (Cortés et al. 2014) and may induce higher oxidative stress due to the

metabolic effects (Villanueva et al. 2013). Therefore, it seems possible that THs may also be a

mediator of pace of life.

In addition to life-history differences, some environmental factors may also contribute to the

inter-specific variation of maternal yolk THs and should be taken into account. First, we expected

that the foraging environment, through its direct influence on food composition, may influence the

inter-specific variation of maternal yolk THs. In a previous study with captive wild-type rock pigeons

(*Columba livia livia*), food conditions before and during egg-laying were indeed found affecting

maternal yolk TH levels (Hsu et al. 2016). One possible explanation to this finding probably lies on

the iodine content in the diet, as iodine is an essential ingredient for TH production (Bizhanova and

Kopp 2009). In humans, even mild iodine deficiency during pregnancy has been shown to lead to

deficiency of maternal T4 and hence negative effects on foetal development (Bath et al. 2013;

Trumpff et al. 2013). In wild animals, previous studies have also found that environmental iodine

availability is associated with antler weights of roe deer (*Capreolus capreolus*) via its influence on TH

production (Lehoczki et al. 2011), and freshwater alligators (*Alligator mississippiensis*) have on

average lower plasma T3 and T4 concentrations than the estuarine conspecifics (Boggs et al. 2011).

Therefore, we hypothesized that iodine availability from the environment and diet (e.g. Eckhoff and

Maage 1997) could contribute to part of the variation in maternal yolk THs. Moreover, captive

environment differs in many aspects from a wild environment, such as food availability and safety

from predators. The confinement and regular human disturbance might also pose as chronic

stressors, which may influence THs (e.g. Angelier et al. 2016) and therefore maternal TH deposition.

We therefore hypothesized that captivity may also have influence on maternal yolk THs.

To test these hypotheses, we collected related life-history traits pertaining to each

hypothesis from the literature. The environmental variables were assigned based on the sampled

population. Using birds as a model system, we measured maternal THs in egg yolks from 34 species

across 17 families and 6 order, using a nano-LC-MS/MS protocol (Ruuskanen et al. 2018),

constructing the first and unique interspecific maternal TH data set. Combining the data of life-

history traits and maternal THs, we were able to run phylogenetic mixed-effects models using a

Bayesian framework to test our hypotheses. Potentially,

Materials and Methods

Sample collection

We collected unincubated eggs from 34 species of birds across 17 families and 6 orders (n =

1-21 per species, median=7, Table S1). Nests of wild species were located by extensive nest searches

in the known breeding habitats of each species or from nest-box populations during 2016-2017. We

aimed at collecting eggs from species with varying body sizes and life-history traits and included

both precocial and altricial species. All eggs were collected before clutch completion (i.e. when

found, the eggs were cold and clutch was not completed) and one, randomly sampled egg per clutch

was collected. Therefore, in most of the cases, the laying order of the sampled egg was unknown.

Most of the wild species were sampled from Finnish populations near Turku (60°27'N, 22°16'E), Oulu

(65°01'N, 25°28'E) and Kauhava (63°06'N, 23°04'E) regions, except thorn-tailed rayaditos

(*Aphrastura spinicauda*, sampled mostly from Chiloé island, Chile, 42°40'S, 73°59'W) and collared

flycatchers (*Ficedula albicollis*, from Burgsvik, Gotland, Sweden, 57°02'N, 18°16'E). The eggs of

captive homing pigeons (*C. l. domesticus*) and domesticated chicken (*Gallus gallus domesticus*) and

eggs of semi-captive pheasants (*Phasianus colcichus*) and grey partridges (*Perdix perdix*) were

sampled from local breeders in Finland. The eggs of red jungle fowl (*G. g. gallus*) and rock pigeons (*C.

l. livia*) were sampled from the captive colonies at the University of Groningen, the Netherlands, and

the eggs of captive zebra finches (*Taeniopygia guttata*) were from the University of Glasgow, UK. All

collected eggs were first frozen and stored at -20°C until further analysis. For most of the species,

the frozen eggs were shipped to the University of Turku, where we separated the yolk for hormone

analysis (see *Yolk TH analysis*). Two exceptions are the collared flycatchers and the rock pigeons, for

which the eggs had been dissected and the yolk had been homogenized and mixed in known amount

of milli-Q (MQ) water before shipped to University of Turku and were already ready for extraction.

Eggs of wild species were collected under licenses from the environmental authorities in each

country: Finland, VARELY/665/2016, POPELY/61/2016, VARELY/63/2016, VARELY/412/2016,

VARELY/6085/2016, and Finnish Wildlife Agency 84/2016; Chile, Agricultural Ministry 404/2017,

Sweden, National Board for Laboratory Animals. Captive species were housed and eggs were

collected following all national and international guidelines

<u>Yolk TH analysis</u>

We extracted THs from the egg yolks and measured two THs (T3 and T4) at the University of

Turku and Turku Centre of Biotechnology. Yolk was separated from albumen after a short period of

thawing. Few eggs showed signs of very early development and all such eggs were discarded from

the analysis. After dissection, yolks were weighed (~0.01 g), and a half (for passerines) or a quarter

(for other species) of the yolk was weighed for extraction and we added an equal amount of MQ to

facilitate homogenizing.

We used the extraction protocol described previously (Ruuskanen et al. 2016b). In brief, we

first added 2 ml methanol to the yolk-MQ mixture (ca. 300 mg) and homogenized the sample. A

known amount of $^{13}C_{12}$-T4 (Larodan) was added to each sample as an internal tracer, which allowed

us to correct for extraction efficiency (i.e. recovery). Four ml of chloroform was then added and

samples were centrifuged at +4°C 1900g for 15 min. The supernatant was then collected and the

pellet was re-extracted in 2:1 chloroform-methanol mixture, back-extracted with 0.05% $CaCl_2$ into

aqueous phase, and re-extracted with chloroform:methanol:0.05% $CaCl_2$ mixture (3:49:48). The

aqueous phase was further purified on Bio-Rad AG 1-X2 resin columns, eluted with 70% acetic acid,

and then vacuum-evaporated overnight. Samples from different species were spread over extraction

batches and the extraction batch ID was included as a random intercept in the statistical models.

We used a validated nano-flow liquid chromatography-mass spectrometry (LC-MS) protocol

to measure yolk T3 and T4 simultaneously (Ruuskanen et al. 2018). In brief, the dry TH extracts were

re-suspended in 150 µl 0.1% NH_3 and further diluted with 0.01% NH_3. The dilution factor depended

on the expected amount of T3 and T4 in each sample. Internal standards $^{13}C_6$-T3 and $^{13}C_6$-T4 were

added to identify and also quantify T3 and T4 in the samples. A triple quadruple mass spectrometer

(TSQ Vantage, Thermo Scientific, San Jose, CA) and nano-flow HPLC system Easy-nLC (Thermo

Scientific) were used for the measurements. The on-column quantification limits were 10.6 amol and

17.9 amol for T4 and T3, respectively (Ruuskanen et al. 2018). The data was acquired automatically

using Thermo Xcalibur software (Thermo Fisher Scientific) and subsequently analysed by Skyline

(MacLean et al. 2010). TH concentrations were calculated using the peak area ratios of sample to

internal standard, corrected for recovery, and expressed as pg/mg yolk. We further calculated the

total amount of THs per yolk and expressed it as ng/yolk.

<u>Life-history traits and environmental factors</u>

I. Altricial-Precocial continuum

We specified all our species' **developmental mode** as either precocial or altricial, following Starck

and Ricklefs (1998a). We only had four semi-precocial species in our data set, which all belong to the

family Laridae (gulls and terns) and only one semi-altricial species (the kestrel, *Falco tinnunculus*).

Therefore, semi-precocial and semi-altricial species were pooled with precocial or altricial species,

respectively.

II. Migratory versus resident species

Species' **migratory status** was categorized as either migratory or resident, determined mainly

according to the data compiled in Lehikoinen et al. (2003), McNab (2009) and Pap et al. (2015).

When in contradiction, the assignment was made based on the population from which we collected

eggs.

III. Pace-of-life continuum

The pace-of-life history continuum incorporates many related life-history traits, including a species'

developmental duration, fecundity, metabolic rate, growth rate, and the rate of maturation and

senescence.

(i) The data on **developmental duration** was collected mainly from the amniote life-history

database compiled by Myhrvold et al. (2015) with supplementation from other literature

(see *supplementary materials*). We collected the **incubation duration** and **the time to**

fledging (from hatching) to represent the prenatal and postnatal part of the developmental

period. We additionally calculated two more variables: **the total length of development**

(incubation duration + time to fledging) and **the proportion of prenatal development**

(incubation duration / total developmental length).

(ii) The **growth rate** data originated from the data set in Starck and Ricklefs (1998b) and

supplemented with other literature (see *supplementary materials*). For comparability, the

logistic growth rate constants were used to represent the growth rate of each species.

Whenever multiple sources of the growth rates are available for a species, an average was

calculated and used.

(iii) **Species' body mass, clutch size** and **number of clutches per year** were obtained from

Myhrvold et al. (2015).

(iv) The data of **basal metabolic rates** (BMR) was mainly obtained from McKechnie et al. (2006)

and McNab (2009), supplemented by other literature (see *supplementary materials*).

(v) **Maximum lifespan** and **ages at first reproduction** originated from the database AnAge

(Tacutu et al. 2013), except for the thorn-tailed rayaditos (*Aphrastura spinicauda*), for which

the data was derived from the unpublished data by R.A. Vásquez et al.

IV. Environmental factors

For environmental factors, we created two variables to represent species' **foraging environment** and

captivity. Respectively, species were categorized as marine-bound (foraging on marine food

resources) or terrestrial (foraging on non-marine bound resources) based on the breeding territory

and diet, and wild or captive. Semi-captive species (the pheasant and the grey partridge) were

categorized as captive.

All life-history and environmental data we compiled for this study, as well as the sources we

consulted, the final values used in all our models, will be provided in the supplementary materials or

deposited in a public repository.

Phylogenetic mixed models

Model specifications

Following Hadfield and Nakagawa (2010), we ran phylogenetic mixed models with the R package

MCMCglmm (Hadfield 2010). T3 and T4 concentrations (pg/mg yolk) as well as total contents

(ng/yolk) were ln-transformed and standardized and used as dependent variables in separated

univariate models. Species body mass and other life-history and environmental variables (see *Life-*

history traits and environmental factors) were treated as fixed factors and covariates. In order to

facilitate model fitting, body mass was also ln-transformed and standardized. Before fitting models,

all two-level categorical variables were first dummy-coded as -0.5 and 0.5 (Table S2), to facilitate

model mixing/convergence. The life-history traits that are known to be associated with body mass

(BMR, growth rate, maximum lifespan, and age at first reproduction) were first corrected for body

mass and phylogenetic relatedness among species using the R package *phytools* (Revell 2009, 2012).

BMR data, in particular, was first converted to mass-specific BMR and ln-transformed and corrected

for size and phylogeny.

The four variables of developmental duration were strongly correlated with one another.

Therefore, we did a phylogeny-corrected principal component analysis (PCA) using R package

phytools (Revell 2009, 2012). The first two PCs explained 99.74% of the variance (Table S3). The

loadings suggested that PC1 is highly positively correlated with the time to fledging and the total

length of developmental period, and negatively correlated with the proportion of prenatal

development, while PC2 is negatively correlated with the duration of prenatal development (Table

S3). In subsequent phylogenetic mixed models, we therefore used the scores of PC1 and PC2 for

each species to represent the developmental duration.

Because the data of life-history traits was not available for all species, we defined several

sets of models depending on data availability, listed in Table 1. Each model and data set was thus

pruned for the specific set of species where data was available. Each model set consists of separate

models to analyse the concentrations as well as total contents of T3 and T4, respectively. This is

because in oviparous species, the total contents of maternal hormones in the yolk is fixed after yolk

formation. Therefore, the total content of hormones represents an initial maximum hormone

availability and may represent some different biological meanings from the concentration.

In all models, the phylogenetic relatedness among all species included was treated as

random effects, following Hadfield and Nakagawa (2010). Additionally, species name was also

included as a random factor to control for within-species non-independence among the multiple

eggs from the same species. For this random factor, homing pigeons and domesticated chickens

were coded separately from their wild-type ancestor rock pigeons and red jungle fowl, despite that

they were not distinguished in the phylogeny (i.e. treated as they same species, *Columba livia* and

Gallus gallus). The batch of hormone extraction was also included as an additional random factor.

Response variables were all assumed to follow Gaussian distributions. All models were run for

750000 iterations with a 350 thinning interval and burn-in of 50000, aiming for an approximately

2000 effective sample size for all parameters. Following Houslay and Wilson (2017), we used an

uninformative parameter-expanded prior following a Cauchy distribution (V=1, nu=1, alpha.mu=0,

alpha.V=25^2) for all random factors. Tests using an uninformative invert-Wishart prior or improper

flat prior gave qualitatively same results, suggesting the results are robust against the choice of

priors.

Phylogenetic trees and phylogeny uncertainty

Each phylogeny set (Table 1) was correspondingly pruned and obtained from BirdTree.org (Jetz et al.

2012) based on two phylogenetic backbones (Ericson et al. 2006; Hackett et al. 2008). For each

backbone, 100 possible trees were generated and downloaded. All models were tested repeatedly

across the 100 possible trees in each backbone and the posterior means of parameter estimates and

their 95% credible intervals (CIs) were stored. The average posterior mean and the average

boundaries of the 95% CIs were calculated to account for phylogenetic uncertainty (Rubolini et al.

2015) and reported. The results are highly similar across all possible trees (see Fig. S1 for examples)

and between the two backbones. Therefore we only reported the results based on the Hackett

backbone.

Model diagnostics

As models across all possible phylogenetic trees produced highly similar results, we examined model

performance by re-fitting all models with a consensus tree. The consensus tree was derived using

the R package *phytools* (Revell 2012) for each phylogeny set. Model diagnostics were conducted by

visual inspection on the trace plots for proper mixing and on autocorrelations. Model convergence

was also checked using Gelman diagnostics provided in the R package *coda* (Plummer et al. 2006).

For all models, visual inspection did not raise any red flags for poor mixing or substantial

autocorrelation and all models passed the Gelman diagnostics with all potential scale reduction

factors < 1.05.

Phylogenetic signals

Phylogenetic signal, a.k.a. phylogenetic heritability (H^2), was calculated as

$$H^2 = \frac{\sigma_a^2}{\sigma_a^2 + \sigma_s^2 + \sigma_b^2 + \sigma_e^2}$$

where σ_a^2 represents the variance of the phylogeny; σ_s^2 represents the variance accounted by

individual species; σ_b^2 represents the variance of hormone extraction batches; σ_e^2 represents the

residual variance. The H^2 calculated this way is equivalent to the Pagel's λ (Hadfield and Nakagawa

2010).

Because phylogenetic signal is in fact the proportion of the phylogenetic variance over the

total variance of all random effects, the fixed factors included in the model will influence the value of

the phylogenetic signal by changing the residual variance. Therefore we calculated the phylogenetic

signals for yolk T4 and T3 based on the phylogenetic mixed model that only included the life-history

traits whose 95% CIs did not encompass 0 (i.e. having credible associations). These are the

developmental mode, migratory status, captivity, and body mass (see *Results*). In these models, all

34 species and the full phylogeny set was included and we used a consensus tree from the Hackett

backbone, derived by using the *phytools* package (Revell 2012).

Table 1. Model, phylogeny and data sets defined in this study

Model set	Phylogeny	Data set	Model structure (fixed factors)	Remark
1	All 34 species	Full data set	~developmental mode + migration + body mass + maximum lifespan + foraging environment + captivity	
2a	Excluding *Coturnix japonica**	Excluding domesticated Japanese quails and domesticated chicken	~developmental mode + migration + body mass + clutch size + clutches per year + captivity	Domesticated quails and chicken do not lay clear "clutches" of eggs.
2b	Excluding *Coturnix japonica*	Excluding Japanese quails	~developmental mode + migration + body mass + PC1 and PC2 of developmental period + captivity	Time to fledging data was missing for Japanese quails
3	Excluding *Gallus gallus*	Excluding red jungle fowl and domesticated chicken	~developmental mode + migration + body mass + age at first reproduction + captivity	Age at first reproduction data was missing for both jungle fowl and chicken
4	28 species with BMR data	Species with BMR data	~developmental mode + migration + body mass + BMR + captivity	
5	29 species with growth rate data	Species with growth rate data	~developmental mode + migration + body mass + growth rate + captivity	

* In Model set 2a, both Japanese quails and domesticated chicken were removed from the "data

set" but only *Coturnix japonica* was removed from the phylogeny because *Gallus gallus* was still

needed in the phylogeny for the data from red jungle fowl.

Results

<u>Inter-specific variation in yolk THs and phylogenetic signals</u>

Among the 34 species, we observed a substantial inter-specific variation in both T3 and T4

concentrations (Fig. 1, Table S1). Great tits (*Parus major*) and redshanks (*Tringa totanus*) had the

lowest and highest concentrations of yolk T3 and T4 among the 34 species, respectively. Between

the two species, the difference in average yolk THs reached 100-fold for T3 (great tit, mean±SD =

0.112±0.032 pg/mg yolk, n=11; redshank, mean±SD = 11.242±5.067 pg/mg yolk, n=4), and also

reached 18-fold for T4 (great tit, mean±SD = 0.989±0.292 pg/mg yolk, n=12; redshank, mean±SD =

18.101±4.125 pg/mg yolk, n=4).

The moderate to strong phylogenetic signals indicated that the phylogenetic relatedness

among species accounted for 50%-75% of the inter-specific variation in yolk THs (Table 2). On

average, T3 has stronger phylogenetic signals although their 95% CIs overlapped.

Table 2. Phylogenetic signals (i.e. phylogenetic heritability) for maternal yolk T3 and T4.

Hormone	Parameter	Posterior mean [95% CI]
T3	concentration	0.76 [0.56, 0.93]
	total content	0.77 [0.57, 0.90]
T4	concentration	0.49 [0.16, 0.79]
	total content	0.65 [0.42, 0.87]

<u>Associations between yolk THs and life-history variables</u>

The estimated posterior means and 95% CIs from all models are collected in Fig. 2. Precocial

species clearly deposited larger total amounts of THs in the egg yolks than altricial species (Fig. 3).

The model set 1 supported this difference with positive posterior means and 95% CIs, when

controlling for species body mass and other covariates, especially for T3 contents (Fig. 2D). Also,

migratory species generally deposited higher concentrations and also larger amounts of both THs in

the egg yolks than resident species (Fig. 4), which were supported by the model set 1 (Fig. 2). Species

body mass also strongly correlated with yolk TH content, but not with yolk TH concentration (Fig. 2).

Along the slow-fast pace-of-life continuum, however, no traits were found to be credibly associated

with maternal yolk THs (Fig. 2)

<u>Associations between yolk THs and environmental variables</u>

Between terrestrial and marine-bound species, although the raw data showed that marine-bound

species apparently deposited more yolk THs (Fig. 5), this was not supported by the phylogenetic

mixed models (Fig. 2). Between captive and wild species, credible differences in both yolk T3 and T4,

and in both concentrations and total contents (Fig. 2) were found. This suggested that captive

species deposit higher concentrations and larger amounts of yolk THs than wild species (Fig. 6).

Discussion

<u>Maternal THs and life-history variation</u>

Among the three life-history continuums, our phylogenetic mixed models estimated credible differences of maternal yolk THs between the migratory and resident life histories, and between precocial and altricial developmental modes. However, contradicting to our expectation, none of the pace-of-life (POL) related traits were found credibly associated with maternal yolk THs, after controlling for body mass. The moderate to strong phylogenetic signals also indicate that a substantial proportion of variation in maternal THs is accounted by the phylogeny itself. This suggests a substantial extent of phylogenetic inertia, which may present some evolutionary constraints in maternal TH deposition. Despite the constraint, the credible differences of maternal TH between precocial and altricial species, and between migratory and resident species still suggest that maternal THs are involved in the evolution of these two life-history continuums or vice versa.

THs and migration

Migration is a critical and demanding life-history stage. Many physiological changes are occurring both before and after migration, such as fat deposition, gonadal regression and recrudescence. The temporal overlap with other seasonal activities, such as breeding and moult, also tends to be minimized over evolutionary history in migratory species (Hahn et al. 2015). By contrast, resident species usually have a relatively longer time window to coordinate the physiological transition before and after reproduction compared with migratory species. Therefore, migratory species may require a higher sensitivity or accuracy to fine-tune the transition. Our result of higher yolk TH levels, not only in terms of higher concentrations but also larger amounts, in migratory than resident species probably suggest prenatal programming effects on the coordination of such annual schedule. THs play a central role in the molecular mechanisms of photoperiodic response. The photoperiodic change influences the expression levels of the thyroid-stimulating hormones (TSH) in the hypothalamus, which subsequently induces reciprocal switching of two deiodinases (DIO2,

converting T4 to active T3 and DIO3, converting T4 and T3 to inactive THs) that activate and

inactivate T3, respectively (Nakao et al. 2008; Dardente et al. 2014). This photoperiodic DIO2/DIO3

expression pattern further regulates the seasonal maturation of gonads. Interestingly, in a study of

great tits, the photoperiodic DIO2/DIO3 gene expression pattern was only evident in a Swedish

population but not in a southern German population (Perfito et al. 2012). This difference therefore

seems to suggest a higher sensitivity to photoperiodic change in a northern population, where the

appropriate time window to breed is shorter. Difference in maternal THs could have some

organizational effects on TH responsiveness later in life. There may be still species differences in the

seasonal regulation of DIO2/DIO3 expression, as a similar pattern was not found in European

starlings (*Sturnus vulgaris*, Bentley et al. 2013). Comparison of TH-related gene expression between

conspecific populations that possess different migratory propensity, potentially in combination with

TH manipulation, will probably help identify the elements responsible for such differences.

Precocial development requires more maternal THs?

The result that precocial species deposited higher amounts of THs, particularly T3, in the egg

yolks than altricial species fits well in line with their distinct developmental trajectories (Starck and

Ricklefs 1998c). Compared with precocial species, altricial species seem to temporally shift their

physiological maturity to a later life stage in visual and locomotory ability (Starck and Ricklefs 1998c),

HPA and HPT axis (McNabb 2006; Wada 2008; de Groef et al. 2013) and thermoregulation (Price and

Dzialowski 2018). Our result appears to suggest the requirement of a larger amount of maternal THs

for such an advanced ontogenetic trajectory in precocial birds.

An alternative way to look at the contrasting ontogenetic differences between precocial and

altricial development may be the placement of hatching on the ontogenetic timeline. Regardless of

the developmental modes, the event of hatching requires a lot of physiological changes to prepare

the young for the "outside" world and therefore marks a major life-stage transition. As the altricial

developmental mode is assumed to evolve later than the precocial mode (Starck and Ricklefs 1998c;

de Groef et al. 2013), one way to look at the shift is that the event of hatching is moved "leftwards"

along the ontogenetic timeline in the altricial development so that hatching occurs earlier when the

young is still poorly developed. Whether maternal THs mediates such a shift during the evolution

from precocial to altricial development requires further studies, but according to our data such a

change may coincide with a reduction in TH deposition over the course. We therefore could expect

intermediate levels of maternal THs in semi-precocial and semi-altricial species. Interestingly, the

only semi-altricial species we have measured, the Eurasian kestrel, indeed has higher maternal TH

content than most other altricial species, but lower than most semi-precocial and precocial species

(Fig. 3).

Unlike the differences between migratory and resident species, only the total content of THs

was found associated with the developmental mode. We reiterate here that our model has

accounted for species body mass, so the result is not due to the generally larger body size in

precocial species. This therefore suggests that for some traits it may be the total content that

matters, instead of the *per capita* amount (i.e. concentration).

No size-independent associations between maternal THs and pace of life (POL)

Given the known function of THs on metabolism and interactions with glucocorticoids, it is

surprising that we did not detect any obvious associations between maternal THs and all tested POL-

related traits (Fig. 2). Although yolk TH contents show strong correlations with species body mass,

this is because large species produce large eggs and large yolks, hence higher total amounts of yolk

THs. After controlling for body mass and phylogenetic relatedness, only clutch size showed a

potential trend with T4 contents. The model estimated a negative posterior mean and the 95% CI

was slightly overlapped with 0 (Fig. 2B), implying that species with larger clutch sizes tended to

deposit less T4 in the egg yolks. Such a trend is probably driven by the difference between

passerines and shorebirds (Fig. S2). Confirming such a relationship perhaps would require a larger

sampling scheme that includes more species.

Our hypothesis that maternal THs may be associated with POL is based on the relationships

of blood THs with glucocorticoids and with metabolic rates. The lack of associations between

maternal THs and POL-related traits therefore probably implies that the effects of maternal THs

cannot be translated directly to blood THs. This could be due to limited organizational effects of

maternal THs on the HPT axis or more complex manifestation, such as sex-specific effects. To our

knowledge, there is currently only a couple of experimental studies that have examined how

maternal THs influenced blood THs and metabolic rates during the nestling phase. In the great tits,

yolk TH elevation did not influence nestling RMR before fledging (Ruuskanen et al. 2016a). In the

rock pigeons, similar yolk TH elevation raised the RMR in female nestlings but reduced it in males on

the hatching day, a pattern that was consistent with the effects on plasma T4 concentrations (but

not T3) at day 14 after hatching (in the middle of the nestling phase, Hsu et al. 2017). Although in the

pigeon study, elevated yolk THs showed consistent effects on nestling RMR and plasma T4, the sex-

specific pattern remains unexplained. Nevertheless, a previous study in domesticated chicken

embryos does show that the injection of THs into the yolk influenced the expression of the TH-

transporters in the brain (van Herck et al. 2012). This suggests that differential exposure to maternal

THs may influence brain TH availability. Such changes might have a potential to alter the sensitivity

and responsiveness of the HPT axis for a long term. Therefore, so far the notion that that maternal

THs may organizationally shape the HPT axis have not received substantial support but for the

moment cannot yet be ruled out. Another possibility to explain the lack of associations between

maternal THs and POL-related traits is that blood THs do not have such associations. This would be

quite surprising considering the close associations between THs with metabolic rates,

glucocorticoids, and oxidative stress, but after all such a possibility has not yet been confirmed or

falsified by far.

Maternal THs and life-history variation: cause or consequence?

We have discussed our results as maternal THs could be a mediator over the evolutionary course of

the life-history differences. However, it could be the physiological differences between migratory

and resident life histories, and between precocial and altricial developments that result in

differential maternal TH deposition during egg formation across species. So far, the physiological and

molecular mechanism of yolk hormone deposition in birds is still unclear. However, some

circumstantial evidence have suggested that yolk TH deposition does not reflect blood TH variation

(Hsu et al. 2016) and some regulatory mechanism of yolk TH deposition possibly exists. Thus

comparison in blood TH levels across species with their life histories can provide valuable

information and to our knowledge, no such studies have been done. Interestingly, one study in two

closely-related skylark species, for example, did show that the migratory Eurasian skylarks (*Alauda*

arvensis) have higher pre-breeding blood T3 levels than the resident Asian short-toed larks

(*Calandrella cheleensis*, Zhao et al. 2017). If in general migratory species keep comparatively higher

circulating TH levels than resident species, the higher maternal TH deposition could be a reflection.

A meta-analysis making use of the available data from the literature should be promising to test the

associations between life-history variation and TH physiology.

480   Even if maternal THs are a physiological consequence brought by different life histories,

differential yolk TH deposition may still reinforce the evolution of the life-history variation. Due to

the indispensability for embryonic development, maternal THs likely have short-term and long-term

effects on offspring phenotype, such as nestling development, growth, and physiology (Ruuskanen

et al. 2016a; Hsu et al. 2017, 2019a). If higher maternal THs would lend beneficial effects for a

certain life-history strategy, and physiologically such a life history results in a higher maternal TH

transfer, maternal THs and life-history differences will influence each other and undertake a co-

evolution process. At the moment, this is a bold speculation. However, considering that yolk T3

deposition has a moderate heritability (h^2 = 0.25, Ruuskanen et al. 2016c) and clear between-

individual difference (Hsu et al. 2019b), there might be evolutionary potential for this to happen.

Future studies may examine the temporal and quantitative differences in the expression patterns of

TH-related genes in embryonic brains between migratory and resident species, and between

precocial and altricial species, to know if there has been fundamental differences during the early

developmental stage. Another angle would be examining whether birds hatching from eggs with

elevated yolk THs may deposit higher yolk THs than control birds to test if maternal TH can induce

any non-genetic inheritance.

<u>Maternal THs and environment</u>

In terms of environmental factors, our models suggested differences in maternal yolk THs between

captive and wild species, but not between terrestrial and marine-bound species.

Captive life have favoured higher maternal TH deposition

Life in captivity brings about a lot of differences, such as space confinement, (usually) unlimited food

availability and free of predation (Mason 2010). Such differences from the life in the wild can

influence phenotypic expression (i.e. phenotypic plasticity) and even cause selection for certain

traits (Driscoll et al. 2009). For example, captive zebra finches, domesticated or not, started

incubating earlier and therefore induced a stronger degree of hatching asynchrony than wild zebra

finches (Gilby et al. 2013). In our study, all the captive species have acclimated to captive lives for

generations. Therefore the result that captive species deposit higher levels of THs in the egg yolks

likely reflects some selection under captivity. The actual selective force is elusive at the moment, but

probably related to traits that are favoured in captivity, e.g. faster acclimation/habituation for

anthropogenic diet or perhaps dampened stress response to confined space and constant human

disturbance.

Lack of evidence for the potential link with proximity to the sea

The larger yolk TH contents in marine-bound species compared to terrestrial species, apparent from

the raw data, were not supported by the phylogenetic mixed models. The most likely explanation to

this discrepancy is probably that all sampled marine-bound species are from the order

Charadriiformes, hence we lack of power to statistically partition the effects of environment and

phylogeny. Future studies should therefore broaden the sampling scheme to overcome this

problem. An alternative to test the relationship between foraging environment and maternal THs is

to focus on a smaller clade, such as a single order or a single species inhabiting both inland and

marine-bound populations. In the present study, the eggs of the little ringed plover (*Charadrius*

dubius) was sampled from an inland population. The species, interestingly, had the lowest T4

content among all Charadriiform species, which implies possible influence from the environment.

Experimentally manipulating diet iodine content in a captive model system or supplementing iodine-

rich food to a wild population would be also an effective approach in this respect.

<u>Conclusion</u>

In our unique data set, we measured maternal yolk T3 and T4 from 34 bird species across 6 orders

and analysed their associations with a large set of life-history variables when taking some

environmental factors into account. Our phylogenetic mixed models suggested that migratory

species deposited higher levels of both THs in the egg yolks than resident species, and precocial

species deposited higher total content of T3 in the egg yolks than altricial species. In addition,

captive species also consistently deposited higher levels of yolk THs, suggesting some selection

process under captivity. These identified patterns suggest that maternal THs may be involved in the

process of life-history evolution and can respond to on-going selective forces. Further effort should

be invested to uncover the underlying evolutionary mechanisms.

Acknowledgement

We thank Jon Brommer for his kind and helpful statistical advice in the beginning stage of data

analysis. We thank Jorma Nurmi for excellent nest-finding skills and effort in the field. This study was

funded by a grant from the Academy of Finland to SR (grant no. XXXXXX).

Figure 1. Phylogenetic tree and yolk TH concentrations and total contents of the avian species included in this study. The phylogenetic tree (A) is one

possible tree derived from the Hackett backbone (see text). Different colours in species names represent different orders they belong to. Yolk TH

concentrations (B, D) and total contents (C, E) exhibited substantial inter-specific variation (mean±SD, see Table S1 for exact values and sample sizes). Red

symbols represent resident species, while blue symbols migratory species. Circles represent altricial species, and squares precocial species.

Figure 2. Posterior means (± 95% credible intervals) between yolk THs and all life-history variables

tested in this study. White area presents the estimates from model set 1 (see text and Table 1). The

shaded areas present the variables tested in model set 2 (the light and medium grey areas) and

model set 3-5 (the darkest grey area). In model 2-5, the estimates of the variables that had been

tested in model set 1 were not redundantly presented. For developmental model, the estimate

represents the difference of yolk THs in precocial species from altricial species. For migration, the

554   estimate represents the difference in migratory species from resident species. For captivity, the

555   estimate represents the difference in captive species from wild species. For foraging environment,

556   the estimate represents the difference in marine-bound species from terrestrial species.

Figure 3. Boxplots and species-specific averages of yolk T4 (A) and T3 (B) contents (ng/yolk, ln-

transformed raw data) according to developmental mode. Boxplots represent the median (the

middle line) and the first and the third quartiles (the box), and the whiskers extend to 1.5 times of

the interquartile range. Colored dots represent species-specific means (±SE).

Figure 4. Boxplots and species-specific averages of yolk TH concentrations (A, B, pg/mg, ln-transformed raw data) and total contents (C, D, ng/yolk, ln-transformed raw data) between migratory and resident species. Boxplots represent the median (the middle line) and the first and the third quartiles (the box), and the whiskers extend to 1.5 times of the interquartile range. Colored dots represent species-specific means (±SE).

Figure 5. Boxplots of yolk T4 (A) and T3 (B) contents (ng/yolk, ln-transformed raw data) between

marine-bound and terrestrial species. Boxplots represent the median (the middle line) and the first

and the third quartiles (the box), and the whiskers extend to 1.5 times of the interquartile range.

Colored dots represent species-specific means (±SE).

Figure 6. Boxplots of yolk TH concentrations (A, B, pg/mg, ln-transformed raw data) and total

contents (ng/yolk, ln-transformed raw data) in between captive and wild species. Boxplots represent

the median (the middle line) and the first and the third quartiles (the box), and the whiskers extend

to 1.5 times of the interquartile range. Colored dots represent species-specific means (±SE).

References

Angelier, F., Parenteau, C., Ruault, S., Angelier, N. (2016). Endocrine consequences of an acute stress

under different thermal conditions: a study of corticosterone, prolactin, and thyroid hormones in the

pigeon (*Columba livia*). *Comp. Biochem. Physiol. A,* 196, 38-45.

Auer, S.K., Dick, C.A., Metcalfe, N.B., Reznick, D.N. (2018). Metabolic rate evolves rapidly and in

parallel with the pace of life history. *Nat. Commun.,* 9, 14

Bath, S.C, Steer, C.D., Golding, J., Emmett, P., Rayman, M.P. (2013). Effects of inadequate iodine

status in UK pregnant women on cognitive outcomes in their children: results from the Avon

Longitudinal Study of Parents and Children (ALSPAC). *Lancet,* 382, 331-337.

Bentley, G.E., Tucker, S., Chou, H., Hau, M., Perfito, N. (2013). Testicular growth and regression are

not correlated with Dio2 expression in a wild male songbird, *Sturnus vulgaris*, exposed to natural

changes in photoperiod. *Endocrinology*, 154, 1813-1819.

Bizhanova, A., Kopp, P. (2009). Minireview: the sodium-iodide symporter NIS and pendrin in iodide

homeostasis of the thyroid. *Endocrinology*, 150, 1084-1090.

Boggs, A.S.P., Hamlin, H.J., Lowers, R.H., Guillette, L.J. Jr. (2011). Seasonal variation in plasma thyroid

hormone concentration in coastal versus inland populations of juvenile American alligators (*Alligator*

mississippiensis): influence of plasma iodide concentrations. *Gen. Comp. Endocrinol.,* 174: 362-369.

Chastel, O., Lacroix, A., Kersten, M. (2003). Pre-breeding energy requirements: thyroid hormone,

metabolism and the timing of reproduction in house sparrows *Passer domesticus*. *J. Avian Biol.,* 34,

298-306.

Clements, J.F., Schulenberg, T.S., Iliff, M.J., Roberson, D., Fredericks, T.A., Sullivan, B.L. et al. (2018).

The eBird/Clements checklist of birds of the world: v2018. Downloaded from

http://www.birds.cornell.edu/clementschecklist/download/

Cortés, D.C.C., Langlois, V.S., Fernandino, J.I. (2014). Crossover of the hypothalamic pituitary-

adrenal/interrenal-thyroid, and –gonadal axes in testicular development. *Front. Endocrinol.,* 5, 139.

Dammhahn, M., Dingemanse, N.J., Niemelä, P.T., Réale, D. (2018). Pace-of-life syndormes: a

framework for the adaptive integration of behaviour, physiology and life history. *Behav. Ecol.*

Sociobiol., 72: 62.

Dardente, H., Hazlerigg, D.G., Ebling, F.J.P. (2014). Thyroid hormone and seasonal rhythmicity. *Front.*

Endocrinol., 5, 19.

Darras, V.M. (2019). The role of maternal thyroid hormones in avian embryonic development. *Front.*

Endocrinol., 10, 66.

Darras, V.M., Kotanen, S.P., Geris, K.L., Berghman, L.R., Kühn, E.R. (1996). Plasma thyroid hormone

levels and iodothyronine deiodinase activity following an acute glucocorticoid challenge in

embryonic compared with posthatch chickens. *Gen. Comp. Endocrinol.,* 104, 203-212.

de Groef, B., Grommer, S.V.H., Darras, V.M. (2013). Hatching the cleidoic egg: the role of thyroid

hormone. *Front. Endocrinol.*, 4, 63.

Driscoll, C.A., Macdonald, D.W., O'Brien, S.J. (2009). From wild animals to domesticated pets, an

evolutionary view of domestication. *Proc. Natl. Acad. Sci. U.S.A.*, 106, 9971-9978.

Eckhoff, K.M., Maage, A. (1997). Iodine content in fish and other food products from east Africa

analyzed by ICP-MS. *J. Food Compost. Anal.*, 10, 270-282.

Elliott, K.H., Welcker, J., Gaston, A.J., Hatch, S.A., Palace, V., Hare, J.F. *et al.* (2013). Thyroid

hormones correlate with resting metabolic rate, not daily energy expenditure, in two charadriiform

seabirds. *Biol. Open,* 2, 580-586.

Ericson, P.G.P., Zuccon, D., Ohlson, J.I., Johansson, U.S., Alvarenga, H., Prum, R.O. (2006). Higher-

level phylogeny and morphological evolution of tyrant flycatchers, cotingas, manakins, and their

allies (Aves: Tyrannida). *Mol. Phylogenet. Evol.,* 40, 471-483.

Gilby, A.J., Mainwaring, M.C., Griffith, S.C. (2013). Incubation behaviour and hatching synchrony

differ in wild and captive populations of the zebra finch. *Anim. Behav.*, 85, 1329-1334.

Hackett, S.J., Kimball, R.T., Reddy, S., Bowie, R.C.K., Braun, E.L., Braun, M.J. *et al.* (2008). A

phylogenomic study of birds reveals their evolutionary history. *Science,* 320, 1763-1768.

Hadfield, J.D. (2010). MCMC methods for multi-response generalized linear mixed models: the

MCMCglmm R package. *J. Stat. Softw.,* 33, 1-22.

Hadfield, J.D., Nakagawa, S. (2010). General quantitative genetic methods for comparative biology:

phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *J. Evol.*

Biol., 23, 494-508.

Hahn, T.P., Brazeal, K.R., Schultz, E.M., Chmura, H.E., Cornelius, J.M., Watts, H.E. *et al.* (2015). Annual

schedules. In: *Sturkie's Avian Physiology*, 6th edition. (ed. C.G. Scanes). Elsevier Inc., pp. 847-867.

Hau, M., Ricklefs, R.E., Wikelski, M., Lee, K.A., Brawn, J.D. (2010). Corticosterone, testosterone and

life-history strategies of birds. *Proc. R. Soc. B,* 277, 3203-3212.

Houslay, T.M., Wilson, A.J. (2017). Avoiding the misuse of BLUP in behavioural ecology. *Behav. Ecol.,*

28, 948-952.

Hsu, B.-Y., Dijkstra, C., Darras, V.M., de Vries, B., Groothuis, T.G.G. (2016). Maternal adjustment of

constraint: differential effects of food availability on maternal deposition of macro-nutrients,

steroids and thyroid hormones in rock pigeon eggs. *Ecol. Evol.,* 6, 397-411.

Hsu, B.-Y., Dijkstra, C., Darras, V.M., de Vries, B., Groothuis, T.G.G. (2017). Maternal thyroid

hormones enhance hatching success but decrease nestling body mass in the rock pigeon (*Columba*

livia). *Gen. Comp. Endocrinol.* 240: 174-181.

Hsu, B.-Y., Doligez, B., Gustafsson, L., Ruuskanen, S. (2019a). Transient growth-enhancing effects of

elevated maternal thyroid hormones at no apparent oxidative cost during early postnatal period. *J.*

Avian Biol., 50: e01919.

Hsu, B.-Y., Verhagen, I., Gienapp, P., Darras, V.M., Visser, M.E., Ruuskanen, S. (2019b). Between- and

within-individual variation of maternal thyroid hormone deposition in wild great tits (*Parus major*).

Am. Nat., 194, E96-E108.

Jetz, W., Thomas, G.H.,Joy, J.B., Hartmann, K., Mooers, A.O. (2012). The global diversity of birds in

space and time. *Nature,* 491, 444-448.

Korevaar, T.I., Muetzel, R., Medici, M., Chaker, L., Jaddoe, V.W.V., de Rijke, Y.B. *et al.* (2016).

Association of maternal thyroid function during early pregnancy with offspring IQ and brain

morphology in childhood: a population-based prospective cohort study. *Lacet Diabetes Endocrinol.,*

4. 35-43.

Kulkami, S.S., Buchholz, D.R. (2012). Beyond synergy: corticosterone and thyroid hormone have

numerous interaction effects on gene regulation in *Xenopus tropicalis* tadpoles. *Endocrinology,* 153,

5309-5324.

Lehikoinen, E., Gustafsson, E., Aalto, T., Alho, P., Klemola, H., Laine, J. *et al.* (2003). *Vasinais-Suomen*

Linnut. Turun Lintutieteellinen Yhdistys, Turku. (In Finnish)

Lehoczki, R., Erdélyi, K., Sonkoly, K., Szemethy, L., Csányi. (2011). Iodine distribution in the

environment as a limiting factor for roe deer antler development. *Biol. Trace Elem. Res.*, 139, 168-

176.

MacLean, B., Tomazela, D.M., Shulman, N., Chambers, M., Finney, G.L., Frewen, B. *et al.* (2010).

Skyline: an open source document editor for creating and analyzing targeted proteomics

experiments. *Bioinformatics,* 26, 966-968.

Mason, G.J. (2010). Species differences in responses to captivity: stress, welfare and the comparative

method. *Trends Ecol. Evol.*, 25, 713-721.

McKechnie, A.E., Freckleton, R.P., Jetz, W. (2006). Phenotypic plasticity in the scaling of avian basal

metabolic rate. *Proc. R. Soc. B,* 273, 931-937.

McNab, B.K. (2009). Ecological factors affect the level and scaling of avian BMR. *Comp. Biochem.*

Physiol. A, 152, 22-45.

McNabb, F.M.A. (2006). Avian thyroid development and adaptive plasticity. *Gen. Comp. Endocrinol.,*

147, 93-101.

McNabb, F.M.A., Darras, V.M. (2015). Thyroids. In: *Sturkie's Avian Physiology*, 6[th] edition. (ed. C.G.

Scanes). Elsevier Inc., pp. 535-547.

Medici, M., Timmermans, S., Visser, W., de Muinck Keizer-Schrama, S.M.P.F., Jaddoe, V.W.W.,

Hofman, A. *et al.* (2013). Maternal thyroid hormone parameters during early pregnancy and birth

weight: the generation R study. *J. Clin. Endocrinol. Metab.,* 98, 59-66.

Moog, N.K., Entringer, S., Heim, C., Wadhwa, P.D., Kathmann, N., Buss, C. (2017). Influence of

maternal thyroid hormones during gestation on fetal brain development. *Neuroscience, 342,* 68-100.

Moore, M.P., Whiteman, H.H., Martin, R.A. (2019). A mother's legacy: the strength of maternal

effects in animal populations. *Ecol. Lett.*, 22, 1620-1628.

Mousseau, T.A., Fox, C.W. (1998). *Maternal effects as adaptations.* Oxford University Press, New

York, NY.

Mullur, R., Liu, Y.-Y., Brent, G.A. (2014). Thyroid hormone regulation of metabolism. *Physiol. Rev.,* 94,

355-382.

Myhrvold, N.P., Baldridge, E., Chan, B., Sivam, D., Freeman, D.L., Ernest, S.K.M. (2015). An amniote

life-history database to perform comparative analyses with birds, mammals, and reptiles. *Ecology,*

96, 3109.

Nakao, N., Ono, H., Yamamura, T., Anraku, T., Takagi, T., Higashi, K. *et al.* (2008). Thyrotrophin in the

pars tuberalis triggers photoperiodic response. *Nature*, 452, 317-323.

Nishiwaki-Ohkawa, T., Yoshimura, T. (2016). Molecular basis for regulating seasonal reproduction in

vertebrates. *J. Endocrinol.,* 229, R117-R127.

Pap, P.L., Vágási, C.I., Vincze, O., Osváth, G., Veres-Szászka, J., Czirják, G.Á. (2015). Physiological pace

of life: the link between constitutive immunity, developmental period, and metabolic rate in

European birds. *Oecologia,* 177, 147-158.

Perfito, N., Jeong, S.Y., Silverin, B., Calisi, R.M., Bentley, G.E., Hau, M. (2012). Anticipating spring: wild

populations of great tits (*Parus major*) differ in expression of key genes for photoperiodic time

measurement. *PLoS ONE*, 7, e34997.

Pérez, J.H., Furlow, J.D., Wingfield, J.C., Ramenofsky, M. (2016). Regulation of vernal migration in

Gambel's white-crowned sparrows: role of thyroxine and triiodothyronine. *Horm. Behav.,* 84, 50-56.

Pérez, J.H., Meddle, S.L., Wingfield, J.C., Ramenofsky, M. (2018). Effects of thyroid hormone

manipulation on pre-nuptial molt, luteninzing hormone and testicular growth in male white-crowned

sparrows (*Zonotrichia leuchophrys gambelii*). *Gen. Comp. Endocrinol.,* 255: 12-18.

Plummer, M., Best, N., Cowles, K., Vines, K. (2006). CODA: convergence diagnosis and output analysis

for MCMC. *R News,* 6, 7-11.

Prati, M., Calvo, R., Morreale de Escobar, G. (1992). L-thyroxine and 3,5,3'-triiodothyronine

concentrations in the chicken egg and in the embryo before and after the onset of thyroid function.

Endocrinology, 130, 2651-2659.

Price, E.R., Dzialowski, E.M. (2018). Development of endothermy in birds: patterns and mechanisms.

J. Comp. Physiol. B, 188, 373-391.

Revell, L.J. (2009). Size-correction and principal components for interspecific comparative studies.

Evolution, 63, 3258-3268.

Revell, L.J. (2012). phytools: an R package for phylogenetic comparative biology (and other things).

Methods Ecol. Evol., 3, 217-223.

Ricklefs, R.E., Wikelski, M. (2002). The physiology/life-history nexus. *Trends Ecol. Evol.,* 17, 462-468.

Rubolini, D., Liker, A., Garamszegi, L.Z., Møller, A.P., Saino, N. (2015). Using the BirdTree.org website

to obtain robust phylogenies for avian comparative studies: a primer. *Curr. Zool.,* 61, 959-965.

Ruuskanen, S., Hsu, B.-Y. (2018). Maternal thyroid hormones: an unexplored mechanism underlyinbg

maternal effects in an ecological framework. *Physiol. Biochem. Zool.,* 91, 904-916.

Ruuskanen, S., Darras, V.M., Visser, M.E., Groothuis, T.G.G. (2016a). Effects of experimentally

manipulated yolk thyroid hormone levels on offspring development in a wild bird species. *Horm.*

Behav., 81: 38-44.

Ruuskanen, S., Darras, V.M., de Vries, B., Visser, M.E., Groothuis, T.G.G. (2016b). Experimental

manipulation of food availability leads to short-term intra-clutch adjustment in egg mass but not in

yolk androgen or thyroid hormones. *J. Avian Biol.,* 47: 36-46.

Ruuskanen, S., Gienapp, P., Groothuis, T.G.G., Schaper, S.V., Darras, V.M., Pereira, C. *et al.* (2016c).

Heritable variation in maternally derived yolk androgens, thyroid hormones, and immune factors.

Heredity, 117, 184-190.

Ruuskanen, S., Hsu, B.-Y., Heinonen, A., Vainio, M., Darras, V.M., Sarraude, T. *et al.* (2018). A new

method for measuring thyroid hormones using nano-LC-MS/MS. *J. Chromatogr. B,* 1093-1094, 24-30.

Sechman, A., Bobek, S. (1988). Presence of iodothyronines in the yolk of the hen's egg. *Gen. Comp.*

Endoccrinol., 69, 99-105.

Starck, J.M., Ricklefs, R.E. (1998a). Patterns of development: the altricial-precocial spectrum. In:

Avian Growth and Development: Evolution within the Altricial-Precocial Spectcrum. (eds. Starck, J.M.,

Ricklefs, R.E.) Oxford University Press, New York, NY, pp. 3-30.

Starck, J.M., Ricklefs, R.E. (1998b). Avian growth rate data set. In: *Avian Growth and Development:*

Evolution within the Altricial-Precocial Spectcrum. (eds. Starck, J.M., Ricklefs, R.E.) Oxford University

Press, New York, NY, pp. 381-415.

Starck, J.M., Ricklefs, R.E. (1998c). *Avian Growth and Development: Evolution within the Altricial-*

Precocial Spectrum. Oxford University Press, New York, NY.

Tacutu, R., Craig, T., Budovsky, A., Wuttke, D., Lehmann, G., Taranukha, D. *et al.* (2013). Human

ageing genomic resources: Integrated databases and tools for the biology and genetics of ageing.

Nucleic Acids Res., 41, D1027-D1033.

Tomasek, O., Bobek, L., Kralova, T., Adamkova, M., Albrecht, T. (2019). Fuel for the pace of life:

baseline blood glucose concentration co-evolves with life-history traits in songbirds. *Funct. Ecol.,* 33,

239-249.

Trumpff, C., de Schepper, J., Tafforeau, J., van Oyen, H., Vanderfaeillie, J., Vandervijvere, S. (2013).

Mild iodine deficiency in pregnancy in Europé and its consequences for cognitive and psychomotor

development of children: a review. *J. Trace Elem. Med. Biol.,* 27, 174-183.

Vágási, C.I., Vincze, O., Pătraş, L., Osváth, G., Pénzes, J., Haussmann, M.F. *et al.* (2019). Longevity and

life history coevolve with oxidative stress in birds. *Funct. Ecol.,* 33, 152-161.

van Herck, S.L.J., Geysens, S., Delbaere, J., Tylzanowski, P. Darras, V.M. (2012). Expression profile and

thyroid hormone responsiveness of transporters and deiodinases in early embryonic chicken brain

development. *Mol. Cell. Endocrinol.*, 349, 289-297.

Vézina, F., Gustowska, A., Jalvingh, K.M., Chastel, O., Piersma, T. (2009). Hormonal correlates and

thermoregulatory consequences of molting on metabolic rate in a northerly wintering shorebird.

Physiol. Biochem. Zool., 82, 129-142.

Villanueva, I., Alva-Sánchez, C., Pacheco-Rosado, J. (2013). The role of thyroid hormones as inductors

of oxidative stress and neurodegeneration. *Oxid. Med. Cell. Longev.* 2013, 218145.

Wada, H. (2008). Glucocorticoids: mediators of vertebrate ontogenetic transistions. *Gen. Comp.*

Endocrinol., 156, 441-453.

Welcker, J., Chastel, O., Gabrielsen, G.W., Guillaumin, J., Kitaysky, A.S., Speakman, J.R. *et al.* (2013).

Thyroid hormones correlate with basal metabolic rate but not field metabolic rate in a wild bird

species. *PLoS ONE,* 8, e56229.

Wikelski, M., Spinney, L., Schelsky, W., Scheuerlein, A., Gwinner, E. (2003). Slow pace of life in

tropical sedentary birds: a common-garden experiment on four stonechat populations from different

latitudes. *Proc. R Soc. Lond. B,* 270, 2383-2388.

Zhao, L., Gao, L., Yang, W., Xu, X., Wang, W., Liang, W. *et al.* (2017). Do migrant and resident species

differ in the timing of increases in reproductive and thyroid hormone secretion and body mass? A

case study in the comparison of pre-breeding physiology rhythms in the Eurasian skylark and Asian

short-toed lark. *Avian Res.*, 8, 10.

 Supplementary materials

 The compiled data set and literature sources we consulted will be provided as additional

 supplementary materials or deposited in a public repository upon final publication.

780 **Table S1.** Mean±SD of yolk T3 and T4 concentrations and total contents of all 34 species in this

781 study.

Species	N of T3	Yolk T3 concentration (pg/mg yolk)	Yolk T3 content (ng/yolk)	N of T4	Yolk T4 concentration (pg/mg yolk)	Yolk T4 content (ng/yolk)
Japanese quail, *Coturnix japonica*	21	3.690±1.009	13.035±3.787	21	7.802±1.661	27.514±6.846
Red jungle fowl, *Gallus gallus gallus*	10	6.163±1.852	78.122±25.792	10	3.633±1.215	45.991±16.501
Domesticated chicken, *Gallus gallus domesticus*	5	4.151±1.315	59.809±16.053	5	9.488±1.621	137.149±16.649
Grey partridge, *Perdix perdix*	7	6.803±2.765	31.229±12.491	7	7.633±2.159	34.930±9.072
Ringed-necked pheasant, *Phasianus colchicus*	9	6.743±2.010	74.055±22.609	9	11.060±5.022	120.725±52.997
Common cuckoo, *Cuculus canorus*	8	4.322±1.065	2.998±0.788	8	7.821±2.112	5.397±1.462
Rock pigeon, *Columba livia livia*	9	3.532±0.785	12.685±2.351	9	8.303±2.982	29.978±10.289
Homing pigeon, *Columba livia domesticus*	4	3.103±0.463	12.575±2.084	4	13.339±0.687	53.981±3.341
Eurasian oystercatcher, *Haematopus ostralegus*	1	3.285	40.782	1	8.946	111.068
Northern Lapwing, *Vanellus vanellus*	5	6.051±0.942	46.195±6.027	5	18.050±4.497	139.166±39.438
Little ringed plover, *Charadrius dubius*	1	9.098	22.972	1	17.984	45.409
Common ringed plover, *Charadrius hiaticula*	3	8.021±1.630	28.325±5.727	3	16.965±3.871	59.980±14.134
Redshank, *Tringa tetanus*	4	11.242±5.067	76.245±32.722	4	18.101±4.125	123.911±25.503
Common tern, *Sterna hirundo*	8	3.883±1.250	21.563±7.313	8	12.920±1.844	71.466±11.647
Arctic tern, *Sterna paradisaea*	6	2.681±0.429	13.602±2.926	6	12.229±3.479	61.544±17.678
Common gull, *Larus canus*	3	3.723±3.521	67.738±75.481	3	10.929±3.444	182.853±99.314
Black-headed gull, *Larus ribidundus**	9	2.363±0.883	22.773±9.292	9	12.789±4.801	121.907±43.035
Kestrel, *Falco tinnunculus*	7	3.000±0.356	12.001±2.698	7	7.992±2.238	31.472±9.262
Thorn-tailed rayadito, *Aphrastura spinicauda*	11	1.375±0.503	0.656±0.278	11	4.564±1.647	2.192±0.996
Jackdaw, *Corvus monedula*	5	0.552±0.390	1.079±0.703	6	2.529±1.248	4.982±2.464

Eurasian jay, *Garrulus glandarius*	1	0.485	0.739	1	6.287	9.568
Blue tit, *Parus caeruleus**	9	0.217±0.104	0.051±0.024	9	1.647±0.659	0.385±0.134
Great tit, *Parus major*	11	0.112±0.032	0.053±0.020	12	0.989±0.292	0.458±0.162
Coal tit, *Parus ater**	7	0.186±0.106	0.045±0.027	7	1.301±0.619	0.312±0.157
Zebra finch, *Taeniopygia guttata*	9	2.729±1.084	0.767±0.372	9	8.681±2.733	2.462±1.028
Tree sparrow, *Passer montanus*	7	0.415±0.232	0.187±0.127	7	2.294±1.046	1.015±0.510
Chaffinch, *Fringilla coelebs*	2	2.016±0.896	0.853±0.391	2	8.927±1.798	3.768±0.815
Greenfinch, *Carduelis chloris**	3	1.603±0.285	0.826±0.244	3	4.366±1.276	2.160±0.273
European starling, *Sturnus vulgaris*	7	0.273±0.131	0.334±0.179	7	1.537±0.695	1.794±0.670
Common redstart, *Phoenicurus phoenicurus*	8	2.805±0.850	0.998±0.367	8	5.073±1.151	1.774±0.424
Collared flycatcher, *Ficedula albicollis*	15	1.978±0.489	0.753±0.201	15	7.209±0.998	2.737±0.418
Pied flycatcher, *Ficedula hypoleuca*	15	1.862±0.570	0.740±0.238	15	5.763±1.421	2.307±0.654
Song thrush, *Turdus philomelos*	4	1.222±0.586	1.260±0.678	4	7.157±2.045	7.389±2.607
Blackbird, *Turdus merula*	6	0.939±0.307	1.353±0.525	6	4.881±1.233	7.013±2.147
Fieldfare, *Turdus pilaris*	3	1.587±0.111	1.904±0.257	3	7.521±3.852	9.200±5.259
Redwing, *Turdus iliacus*	4	1.168±0.376	1.009±0.276	4	5.509±1.543	4.745±0.996

782 * In order to match the phylogenetic trees from BirdTree.org, "old" scientific names were still used in this
783 study. Their updated scientific names according to The Clements checklist of birds of the world v2018
784 (Clements et al. 2018) are: Chroicocephalus *ridibundus*, *Cyanistes caeruleus, Periparus ater, Chloris chloris*

785   **Table S2.** Dummy code for the two-level categorical variables included in the phylogenetic mixed

786   models.

	-0.5	0.5
Mode of development	altricial	precocial
Migratory status	resident	migratory
Foraging environment	terrestrial	marine-bound
Captivity	wild	captive

788   **Table S3.** Proportion of variance explained and loadings of the phylogeny-corrected PCA among the

789   four variables of developmental duration.

	PC1	PC2
Proportion of variance	83.26	16.48
Cumulative proportion	83.26	99.74
Loadings		
Incubation duration	0.264	-0.965
Time to fledging	0.998	0.055
Total developmental length	0.942	-0.329
Prenatal proportion	-0.869	-0.489

Figure S1. Histograms of posterior means for each life-history traits tested in model and data set 1

(see Table 1) across 100 possible phylogenetic trees from the Hackett backbone. All results were

highly similar across different trees, indicated by the narrow range of the posterior means.

806   **Figure S2.** Scatterplot of species clutch size and yolk T4 content on a natural-log scale. Dots

807   represent species-specific means and colors represent the orders they belong to.