nkx3.2 mutant zebrafish accommodate the jaw joint loss through a phenocopy of the head shapes of Paleozoic agnathans

Tetsuto Miyashita[1, 2], Pranidhi Baddam[3], Joanna Smeeton[4], A. Phil Oel[1, 5], Natasha Natarajan[4], Brogan Gordon[1], A. Richard Palmer[1], J. Gage Crump[4], Daniel Graf[3, 6], and W. Ted Allison[1, 6]*

[1] Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2E9

[2] Department of Organismal Biology and Anatomy, University of Chicago, Chicago, IL 60637, USA

[3] Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada T6G 2R3

[4] Department of Stem Cell Biology and Regenerative Medicine, W.M. Keck School of Medicine, University of Southern California, Los Angeles, CA 90033, USA

[5] Current Address: Developmental Biology Unit, European Molecular Biology Laboratory, Heidelberg, 69117, Germany

[6] Department of Medical Genetics, University of Alberta, Edmonton, Alberta, Canada T6G 2R7

* Corresponding author. ted.allison@ualberta.ca; 780-492-4430.

Abstract

The vertebrate jaw is a versatile feeding apparatus that facilitated explosive diversification. Its

functions require a joint between the upper and lower jaws, so jaw joint defects — such as

osteoarthritis or even ankylosis — are often highly disruptive and difficult to study. To describe

consequences of jaw joint dysfunction, we engineered two independent null alleles of a single jaw-joint

marker gene, *nkx3.2*, in zebrafish. The mutations caused the fish to become functionally jawless via a

fusion between the upper and lower jaw cartilages (ankylosis). Surviving to adulthood despite lacking

jaw joints, *nkx3.2* mutants accommodate this defect by: a) remodeling their skulls drastically from

juvenile to adult stages; and b) altering their feeding behaviors from suction to ram feeding. As a result

of the remodeling, *nkx3.2* mutants developed superficial similarities to the skull shapes observed in two

lineages of ancient jawless vertebrates (anaspids and furcacaudiform thelodonts), including: a fixed

open gape, reduced snout, and enlarged branchial region. However, no homology exists in individual

skull elements between these taxa, and most of the modified elements in the mutant zebrafish occur

outside known expression domains of *nkx3.2*. Therefore, we interpret the adult *nkx3.2* phenotype not as

a reversal to an ancestral state, but as convergence due to similar functional requirement of feeding

without mobile jaws. This remarkable convergence underscores jaw joint functions as a powerful

constraint during the development of jawed vertebrate skulls, which implies that functionally viable

morphological patterns are finite, with or without functional jaws. Because *nkx3.2* null zebrafish

display prominent joint ankylosis, drastically modified skull shape, and altered feeding behaviors, these

mutants provide a unique model with which to investigate mechanisms of skeletal remodeling and joint

diseases.

Introduction

The jaw is a functionally versatile innovation that facilitated explosive diversification of gnathostomes

(a clade containing jawed vertebrates), but its basic structure is surprisingly simple and highly

conservative (Miyashita, 2016). A jaw consists of 'a hinge and caps': an upper and lower skeletal

levers hinged at a jaw joint (Depew and Simpson, 2006). As the joint enables biting motions, its origin

is considered the final step in the evolutionary assembly of the vertebrate jaws (Cerny et al., 2010;

Kuratani, 2012; Miyashita, 2016). Across jawed vertebrates, the presumptive jaw joint is marked by the

expression of *nkx3.2*, an NK2 class homeobox gene (a.k.a. *bapx*), at the midheight of the embryonic

mandibular arch (Gillis et al., 2013; Lukas and Olsson, 2018a; Miller et al., 2003; Tucker et al., 2004).

Chondrogenesis dorsal to this expression domain gives rise to a palatoquadrate (upper jaw), whereas

chondrogenesis ventral to it forms Meckel's cartilage (lower jaw) (Medeiros and Crump, 2012). This

basic pattern remains conserved among jawed vertebrates, but later development varies. Marginal

bones arise intramembranously around the often endo-/perichondrally ossified jaw cartilages except in

chondrichthyans (sharks, rays, and skates) (Hall, 2015). In mammals, the jaw joint instead forms

between two such intramembranous bones (temporal and dentary), whereas the proximal jaw joint

becomes the malleus-incus interface that is, in mice, no longer affected by *Nkx3.2* knockout (Tucker et

al., 2004). Despite these variations after pharyngeal chondrogenesis, no gnathostome lineage

secondarily lost functional jaws.

By studying functional jaw loss in our new mutant zebrafish, we asked whether — and how —

jaw functions affect the vertebrate skull shape during development. Clinically documented agnathia in

humans typically accompanies severe congenital disorders such as holoprosencephaly and otocephaly,

but the jaw loss is clearly a secondary effect and not a cause in these cases (Bixler et al., 1985; Brown

and Marsh, 1990; Gekas et al., 2010; Schiffer et al., 2002). Instances of temporomandibular joint

ankylosis may result from trauma or infection, or may be congenital (Adekeye, 1983; Chidzonga, 1999;

Manganello-Souza and Mariani, 2003). If untreated, the ankylosis can lead to the 'bird face' deformity

(El-Sheikh et al., 1996). However, these cases do not fully document the effects of functional jaw loss.

In mammalian models, various jaw/skull deformations have been induced by surgical resection,

detachment, or repositioning of the jaw muscles and/or bones (Bayram et al., 2010; Gomes et al., 2012;

Horowitz and Shapiro, 1955; Lifshitz, 1976; Miyazaki et al., 2016; Rodrigues et al., 2009; Sarnat,

1970; Sarnat and Muchnic, 1971; Toledo et al., 2014). These manipulations occurred well after the

formation of the jaw skeleton and muscles, and the jaws remained partially functional because of

unilateral operations or non-comprehensive disruption. The defects and deformities reported in these

studies imply: a) jaw functions are potentially an important factor in shaping the skulls; and b) any

allele disrupting jaw functions would be generally maladaptive. Nevertheless, these implications are

difficult to explore in the absence of an accessible experimental model.

To fill in this gap, we engineered two distinct null alleles of *nkx3.2* in zebrafish. Previously,

transient knockdown of *nkx3.2* during early development (using morpholinos) had shown fusion of the

nascent jaw cartilages in both zebrafish and frogs (Lukas and Olsson, 2018a; Miller et al., 2003). We

confirmed in zebrafish that the mutants recapitulate this phenotype. Surprisingly, mutant zebrafish are

viable despite the loss of the jaw joint persisting through adulthood. Thus functionally jawless, *nkx3.2*$^{-/-}$

zebrafish dramatically alter the skull shapes late in ontogeny to facilitate feeding, with the mouth fixed

open, the snout reduced, and the branchial region expanded. This open-mouth phenotype, previously

unknown in zebrafish or in other jawed vertebrates, also occurred in two extinct lineages of 400-

million-plus year-old jawless vertebrates, anaspids and thelodonts. Even though they share no

homology in individual facial bones, the *nkx3.2*$^{-/-}$ fish accommodate functional jaw loss by converging

onto these ancient, distantly related agnathans in overall head shapes. Thus, *nkx3.2* mutant zebrafish

provide a unique model for both skeletal remodeling and joint diseases such as osteoarthritis, and to

reevaluate evolutionary implications of phenocopies in general.

Methods

Full methodological details are available in Supplementary Information.

Animal Ethics. Zebrafish maintenance and experiments were approved as protocol number

AUP00000077 by the Animal Care and Use Committee: Biosciences at the University of Alberta as

dictated by the Canadian Council on Animal Care. Other zebrafish work was approved by the

University of Southern California Institutional Animal Care and Use Committee.

Molecular genetics. Two disparate regions of the gene were targeted by CRISPR guide RNA (gRNA)

or TALENs, producing two disparate null alleles that produced similar phenotypes (Fig. 1a). One allele

(ua5011) was engineered with CRISPR/Cas9 (Gagnon et al., 2014) targeted at the beginning of the

homeodomain, and it harbors a 20 bp deletion resulting in a frameshift (Fig. 1a; Data Supplement 1).

The disrupted translation of codons is predicted to abrogate production of the critical homeobox

domain, and instead produce random amino acids. This is predicted to produce a non-functional

Nkx3.2 and a null allele. A disparate allele (el802) was generated using TALENs (Barske et al., 2016)

targeted at the start of the gene. This produced a stably inherited gene with 20 bp deletion, removing

the translation start codon (Fig. 1a). The allele $nkx3.2^{el802}$ is predicted to not produce Nkx3.2 protein.

Morphologically, these two alleles are not readily distinguishable from each other (see Results).

Morphometrics. We compared gapes in zebrafish and anaspids (a lineage of extinct jawless

vertebrates) (Blom et al., 2001) by taking the angle between skeletal elements of the upper and lower

lips (Data Supplement 2). In zebrafish, the angle was taken by extrapolating the axis of the premaxilla

until it meets the axis of the dentary. At younger stages (4 and 14 dpfs), the angle was measured

between the axes of the palatal process and Meckel's cartilage. In anaspids, the upper and lower lips

are demarcated by a series of relatively larger scales, which allowed delineation of the gape angle

(measured at where the extrapolated upper and lower lip margins meet). For geometric morphometric

comparison, eight landmarks were assigned to both zebrafish (2 mpf) and anaspid samples to capture

overall head shapes (see Supplementary Information for rationales of landmark assignment). From the

pool of nearly a thousand catalogued specimens of anaspids, we selected a total of 70 specimens that

show lateral compression during the fossilization process, with the least taphonomic artifact, to reflect

lateral view of the heads. Furcacaudiform thelodonts were excluded because only a handful of

exceptionally preserved specimens are available. The sample size is small for this latter group, and all

such specimens were collected from a single locality, making it difficult to control for taphonomic

artifacts. The landmark data (Data Supplement 2) were transformed using the Procrustes

superimposition method (Rohlf, 1999), and the resulting coordinates were compared using Principal

Component Analysis (PCA).

Results

Jaw joint is ankylosed in *nkx3.2* null alleles

At 4 days post fertilization (dpf), *nkx3.2*$^{-/-}$ zebrafish replicate the *nkx3.2* morpholino knockdown

phenotype: the absence of a jaw joint (Miller et al., 2003). The palatoquadrate and Meckel's cartilage

are fused to each other, and the retroarticular process is absent (Fig. 1d, g, i). Apart from the joint

ankylosis, no marked differences exist in body profile, overall size, or survival rates between *nkx3.2*$^{-/-}$

mutants and wildtype. Heterozygotes are morphologically indistinguishable from wildtype (Fig. 1c, f),

and the *nkx3.2* alleles display recessive Mendelian inheritance. Remarkably, these functionally jawless

homozygous mutants survive beyond early larval stages.

Functionally jawless *nkx3.2*$^{-/-}$ zebrafish modify skull shapes late in ontogeny

Counterintuitive to the maladaptive nature of jaw dysfunctions in general, *nkx3.2*$^{-/-}$ zebrafish continue

to grow without a jaw joint. Marked phenotypic differences against wildtype begin to emerge between

the 2nd and 3rd weeks post fertilization (Fig. 2). The lower jaw becomes downturned in *nkx3.2*$^{-/-}$ fish

resulting in a rigidly fixed open mouth, whereas both upper and lower jaws are upturned in wildtype

zebrafish (Fig. 2a, b). This coincides with the onset of ossification and the period of active feeding in

normal juveniles (Cubbage and Mabee, 1996; Kimmel et al., 1995). Although the palatoquadrate and

Meckel's cartilage remain fused in *nkx3.2*[-/-] mutants, skeletal staining reveals that the upper and lower

jaw elements ossify independently of each other — still without a ball-and-socket joint structure (Fig.

2k, l). All skull elements ossify in *nkx3.2*[-/-] without apparent delay.

The lower jaws are increasingly turned downward in *nkx3.2*[-/-] by the end of the first month,

resulting in a greater gape (Fig. 2d, e). The jaw joint is still absent in *nkx3.2*[-/-] mutants: a sheet of

perichondrium lies between the ossifying quadrate and articular — so the two bones remain distinct

elements — but this interface has none of those essential components of synovial diarthrosis (Fig. 2n).

Many other osteological differences emerge by this stage. Normally, the premaxilla and the maxilla

swing forward to sit nearly vertical and are hinged by the kinethmoid for suction feeding (Hernandez,

2000; Hernandez et al., 2007) (Fig. 2c). In *nkx3.2*[-/-] mutants, however, the premaxilla and the maxilla

are oriented posteroventrally and abutted against the anterior margin of the orbit (Fig. 2d). The

kinethmoid is reduced into a fused bony process, unlike a rod-like hinge element in wildtype

(Hernandez et al., 2007). The downturned lower jaws of the *nkx3.2*[-/-] mutants are relatively shorter than

the normal lower jaws of wildtypes. The basihyal protrudes anteroventrally as much as the lower jaw,

implying that the muscle connecting those two elements (m. intermandibularis posterior) (Schilling and

Kimmel, 1997) may be responsible for the lower jaw orientation. As a result of these modifications,

nkx3.2[-/-] mutants have a shorter snout, a fixed open gape, and a dorsoventrally tall profile.

Morphological variations within a cohort of *nkx3.2*[-/-] mutants is greatest at this stage than in any other,

as indicated by the range of variation in lower jaw depression (Fig. 2o).

The adult *nkx3.2*[-/-] phenotype accommodates functional jaw loss

At 2 months of age and approaching sexual maturity nearly all *nkx3.2*[-/-] mutants have a gape angle

greater than 90 degrees (Fig. 2o). These *nkx3.2*[-/-] adults continue to be characterized by the

morphological differences identified at 1 mpf. The nostrils sit between the eyes as the snout is reduced

in length relative to wildtype zebrafish. The skulls appear to be more highly ossified in $nkx3.2^{-/-}$

mutants than in wildtype, where massive bones and cartilages are identified around the interface of

quadrate and articular, in the lower branchial region, and in the laterally expanded operculum (Figs. 2j,

g, 3b; Movie 1). All homozygous mutants showed the descriptive skeletal traits identified here.

Morphologically, the two alleles ($nkx3.2^{ua5011}$ and $nkx3.2^{el802}$) are virtually indistinguishable from each

other at respective ages (Figs. 2j, k, g, 3b), given variations within a strain (for examples, see variation

in gape angles or sexual dimorphism in $nkx3.2^{ua5011}$: Figs. 2o, 3g).

159   With the modified skulls, adult $nkx3.2^{-/-}$ mutants exhibit feeding behaviors markedly different

160   from wildtype (Movie 1). Wildtype zebrafish feed by suction using rapid lower jaw depression and

161   forward swing of the mobile premaxilla-maxilla complex, consistent with general teleost feeding

162   mechanics. In contrast, adult $nkx3.2^{-/-}$ mutants are constrained by the fixed upper jaw unit and open

163   gape. They instead show ram feeding behaviors (swimming through food).

164   This behavior is correlated with the skull remodeling in $nkx3.2^{-/-}$ mutants. In zebrafish skulls,

the bones form endochondrally (quadrate, anguloarticular, basihyal) or intramembranously (premaxilla,

maxilla, dentary, jugal, opercular, preopercular) (Cubbage and Mabee, 1996; Schilling and Kimmel,

1997). Much of the skeletal remodeling observed in adult $nkx3.2^{-/-}$ mutants occurs in the

intramembranous bones — spatially and temporarily well outside the known expression domain of

$nkx3.2$ (Askary et al., 2017; Miller et al., 2003). Until past 1 mpf, the fusion between jaw cartilages is

not completely ossified in these mutants, potentially allowing plastic remodeling (Fig. 2e, l). These

observations support that $nkx3.2^{-/-}$ zebrafish accommodate functional jawlessness through remodeling

of the skull and changes to feeding behavior.

$nkx3.2^{-/-}$ zebrafish converge onto some ancient agnathans in overall head shapes

Through this dramatic remodeling of the skull, $nkx3.2^{-/-}$ zebrafish assume a head shape reminiscent of

two lineages of extinct jawless vertebrates that have laterally compressed body profiles: birkeniiform

anaspids (Fig. 3c), stem cyclostomes known mostly from the Silurian period (Blom et al., 2001;

Miyashita et al., 2019); and furcacaudiid thelodonts (Fig. 3d), much more elusive stem gnathostomes

known from the Silurian and Devonian periods (excluded from quantitative comparison; see Methods

for rationales) (Märss et al., 2007; Wilson and Caldwell, 1998). Qualitatively, the resemblance is

particularly striking in overall head shape characters, including: fixed gape (depressed lower lips),

shortened snout, interorbital position of nostril, proportionally large branchial region, and massive

parietal region behind the occiput.

In linear morphometrics, the gape angle between upper and lower lips reveals that $nkx3.2^{-/-}$

mutants assimilates the anaspid-like condition late in ontogeny (Fig. 2o). The trait in $nkx3.2^{-/-}$ mutants

departs from that in wildtype after the onset of skull ossification and active feeding, and overlaps with

the range occupied by anaspids (grey zone in Fig. 2o). When the jaws are at rest, the gape angle in

wildtype zebrafish is consistently around 30 degrees regardless of ontogenetic stages. This is also the

case for the $nkx3.2^{-/-}$ mutants at 4 and 14 dpf when they still rely on yolk as main intake. The gape

angle increases steadily thereafter.

In landmark-based geometric morphometrics, $nkx3.2^{-/-}$ mutants aligned more closely with

anaspids than wildtype along the principal components (PCs) that explain the adult phenotype (PCs 1

and 5) (Fig. 3e). Nearly one third of the overall shape variation loads on PC1 and primarily concerns

anteroposterior length and dorsoventral height of the entire head. Compared to wildtype zebrafish,

$nkx3.2^{-/-}$ mutants and most anaspids have a much shorter snout (with the lips and the nostrils shifting

posteriorly toward the eye) and dorsoventrally deeper lower lips. PC 5 explains 7.0 % of the overall

shape variation, and the traits that vary along it are orientation of lower lip and relative height of

nostrils, which clearly set wildtype zebrafish apart from $nkx3.2^{-/-}$ mutants and anaspids. On the

Cartesian grid of PCs 1 and 5, $nkx3.2^{-/-}$ mutants overlap with anaspids in morphospace occupation and

apart from wildtype zebrafish (Fig. 3f). The area of overlap also indicates that $nkx3.2^{-/-}$ mutants are

broadly similar to anaspids in these PCs, not just to one or a few individual anaspid taxa. PCs 2-4 (not

shown in Fig. 3) largely explain variations among anaspids or within wildtype/mutant zebrafish

samples and therefore uninformative for comparison between the groups (datasets containing landmark

coordinates and Procrustes transformation are provided in Data Supplement 2).

Discussion

Mutants corroborate the function of *nkx3.2* in joint development

Our *nkx3.2*$^{-/-}$ zebrafish reinforce the morphant-based insight that this transcription factor is essential to

the development of jaw joint in non-mammalian jawed vertebrates (Miller et al., 2003). This is likely

the widespread condition among jawed vertebrates. *Nkx3.2* knockdown also results in fusion between

the palatoquadrate and Meckel's cartilage in amphibians (Lukas and Olsson, 2018a), and similar

phenotype is observed in chicks with ectopic expression of BMP4 and FGF8 that repressed *Nkx3.2*

(Wilson and Tucker, 2004). In mammals, the palatoquadrate (malleus) and Meckel's cartilage (incus)

migrate to form middle ear ossicles. The malleus-incus joint is not affected in *Nkx3.2*$^{-/-}$ mice, although

the malleus becomes narrower in the mutants, and although *Nkx3.2* plays a role in specification of the

gonium and the anterior tympanic ring (Tucker et al., 2004). Aside from the head, *Nkx3.2* plays a role

in various structures in mice and chicks, including axial column (Herbrand et al., 2002; Lettice et al.,

2001; Murtaugh et al., 2001) and visceral lateralities of spleen and pancreas (Hecksher-Sørensen et al.,

2004; Schneider et al., 1999). We will test whether or not *nkx3.2*$^{-/-}$ zebrafish have parallel phenotypes

to these amniote mutants in a forthcoming work.

During development of the jaw joint, *nkx3.2* has been posited to specify the joint interzone by

inhibiting maturation or hypertrophy of the chondrocytes (Miller et al., 2003; Smeeton et al., 2016).

Similar function has been described for *irx7* and *irx5a* in the hyoid joint (Askary et al., 2015). This is

consistent with the predicted regulatory function of *Nkx3.2* in vertebral development, intervertebral and

interphalangeal joint formation, or somatic chondrogenesis in general (Herbrand et al., 2002; Lettice et

al., 2001; Murtaugh et al., 2001) — via repression of *Runx2*, by upregulating *Col2α1*, and/or through a

positive feedback loop with *Sox9* (Smeeton et al., 2016). These insights are based on: a) experimental

results using amniote embryos or somitic mesodermal cell cultures (Cairns et al., 2008; Kawato et al.,

2012; Lengner et al., 2005; Murtaugh et al., 2001; Provot et al., 2006; Yamashita et al., 2009; Zeng et

al., 2002); and b) clinical and genetic studies of human pathologies, including osteoarthritis and

spondylo-megaepiphyseal-metaphyseal dysplasia (including pseudoepiphyses) (Caron et al., 2015;

Hellemans et al., 2009). Before this study, however, no mutants were available to specifically address

these potential mechanisms of *nkx3.2* functions in the mandibular arch.

The open-mouth phenotype results from plastic remodeling

The late onset and topology of skull/jaw remodeling suggests that *nkx3.2*$^{-/-}$ zebrafish accommodate the

loss of the jaw joint in a plastic response. Other than the absence of jaw joint, *nkx3.2*$^{-/-}$ zebrafish appear

otherwise normal until metamorphosis (14–21 dpf) (Fig. 1d, g). As the skull ossifies, however, the

observed phenotype becomes increasingly prominent (Fig. 2b, d, g, j). Although the fused jaw

cartilages clearly result from *nkx3.2* mutation, it seems unlikely that *nkx3.2* loss-of-function, by itself,

yields all the rest of phenotypic effects described here. A genome-wide analysis suggests that *nkx3.2*

patterning effects in the skull are restricted to the mid-portion of the mandibular arch (Askary et al.,

2017). Nor does *nkx3.2* have known expression or function in the intramembranously ossified skull

elements (Miller et al., 2003; Tucker et al., 2004), which are dramatically modified in our *nkx3.2*$^{-/-}$ fish.

The hypertrophied mandibular cartilage may result from *nkx3.2* loss-of-function in regulating local

chondrogenesis (Smeeton et al., 2016). Still, further investigation is warranted because in amphibians

an ectopic cartilage formed in the mandibular arch with *Nkx3.2* overexpression, not repression (Lukas

and Olsson, 2018b).

Comparative survey across vertebrates supports our interpretation of the adult *nkx3.2*$^{-/-}$

phenotype as skeletal remodeling to accommodate the jaw joint loss. Discrete variation in cichlid jaw

morphology has an epigenetic basis in behaviorally mediated skeletal remodeling, where gaping

frequencies in juveniles correlate with dimensions of the retroarticular process (Hu and Albertson,

2017). The magnitude of morphological changes in the adult $nkx3.2^{-/-}$ phenotype also appears

consistent with a series of surgical experiments in mammalian jaw skeletons (Bayram et al., 2010;

Gomes et al., 2012; Horowitz and Shapiro, 1955; Lifshitz, 1976; Miyazaki et al., 2016; Rodrigues et

al., 2009; Sarnat, 1970; Sarnat and Muchnic, 1971; Toledo et al., 2014) or with the 'bird face'

deformity observed in clinical cases of the temporomandibular joint ankylosis in humans (El-Sheikh et

al., 1996). Collectively, these studies show that latent potentials of development allow a plastic trait to

become expressed. Variations resulting from developmental plasticities — whether induced by

environmental cue, developmental perturbation, or mutation — are often non-random and adaptive

(West-Eberhard, 2005a, 2005b, 2003), as documented across wide ranges of taxa and structures

including cichlid jaw skeletons (Hu and Albertson, 2017; Stewart and Albertson, 2010). Ram feeding

facilitated by the fixed open gape in the adult $nkx3.2^{-/-}$ zebrafish corroborates non-random, adaptive

accommodation of the jaw joint defect.

Jaw joint functions constrain skull morphology in vertebrates

The drastically remodeled skull of adult $nkx3.2^{-/-}$ zebrafish highlights jaw joint functions as an

important factor in morphological diversity of jawed vertebrate skulls (Depew et al., 2005; Depew and

Compagnucci, 2008; Depew and Simpson, 2006). Simultaneously, the absence of jaw joint (or jaw

apparatus altogether) also limits functionally viable forms. This is implied by the superficial

convergence in head shapes between $nkx3.2^{-/-}$ zebrafish and two Paleozoic agnathan lineages (anaspids

and furcacaudiform thelodonts). The $nkx3.2^{-/-}$ zebrafish and these agnathans share a functional

requirement — the absence of a hinge joint between upper and lower lips — and transversely

compressed body profile. No evidence suggests any more similarities in the otherwise wildtype-like

young mutants (Fig. 1d, g) until the lower jaw skeletons begin rotating posteroventrally post 14 dpf

(Fig. 2b, d). Therefore, we interpret the anaspid/thelodont-like traits in the adult *nkx3.2*$^{-/-}$ zebrafish not

as recapitulations of conserved early vertebrate development (atavism), but instead as assimilations of

the patterns previously exploited by early vertebrates in a zebrafish skull (convergence).

Thus, the jaw joint loss (*nkx3.2* loss-of-function phenotype) released *nkx3.2*$^{-/-}$ zebrafish from

forms that can facilitate jaw movement, and exposed them to a different functional requirement of

feeding without mobile jaws. A fixed open gape — accompanied by modification of the skull elements

— is one morphological solution to the functional loss of jaw joint, which is corroborated by ram

feeding exhibited by *nkx3.2*$^{-/-}$ mutants (Movie 1). The magnitude of modification documented post-

hatching (Figs. 1, 2) is a testament to significant functional optimization within the bounds of the

gnathostome bauplan. Simultaneously, well-constrained occupancy by the *nkx3.2*$^{-/-}$ in the PCA plot

(Fig. 3f) implies that alternative morphological patterns are either functionally non-viable (e.g., fixed

closed gape) or developmentally non-accesible (e.g., ectopic formation of a mouth). Thus, the *nkx3.2*$^{-/-}$

mutants provide a rare case study. The results suggest that the vertebrate bauplan allows a limited

repertoire of functionally viable morphological patterns, onto which forms may converge under a given

functional constraint.

In support of our interpretation, remarkable convergences arise via developmental plasticity

under similar functional requirements — even in the absence of homology in individual skeletal

elements — between taxa widely separated chronologically and phylogenetically. Bichirs (*Polypterus*)

routinely trained under terrestrial conditions modify the pectoral fin skeleton through plasticity and

develop morphological conditions observed in stem tetrapods (Standen et al., 2014). In another, *vav2*

and *waslb* mutant zebrafish develop limb-like long bones connected by joints within the pectoral fins

(Hawkins et al., 2018). As shown in *nkx3.2*$^{-/-}$ zebrafish, these experimental manipulations illustrate

through phenotypic accommodation that functions are a strong constraint for adaptive forms, even after

development has laid out phylogenetically conserved patterns (such as homologies of the individual

bones).

nkx3.2 mutants as a unique model for skeletal development and disease

Mechanisms of developmentally plastic remodeling, whether at the level of whole-animal behavior or

gene transcription, remain an elusive component of skeletal development that is challenging to test

experimentally. Therefore, $nkx3.2^{-/-}$ zebrafish provide a unique experimental system in which to test the

role of a joint in skeletal development further. For example, further investigations are warranted to

determine precisely: a) what genetic mechanisms regulate the remodeling of intramembranous

ossifications; b) what components of the remodeling process respond to the jaw joint dysfunction; and

c) whether any feedback exists to coordinate remodeling between developmentally independent, but

functionally connected units (e.g., premaxilla-maxilla complex and kinethmoid). Such insights would

begin to fill in knowledge gaps regarding skeletal and joint diseases including osteoarthritis and joint

ankylosis.

The non-atavistic nature of the $nkx3.2^{-/-}$ phenotype is potentially useful to reevaluate

evolutionarily inspired interpretations of phenocopies in general. There are no evolutionary

relationships in the similarities between $nkx3.2^{-/-}$ zebrafish and anaspids or thelodonts. Anaspids are a

stem cyclostome lineage (Miyashita et al., 2019) (Fig. 3g), and thus a poor surrogate for an ancestral

state. The lineages between anaspids and thelodonts, or those between thelodonts and the gnathostome

crown, do not share a similar combination of morphological traits, but instead are dorsoventrally

depressed forms (Janvier, 1996; Miyashita, 2016; Miyashita et al., 2019) (Fig. 3g). Finally, head

similarities between the mutants and the stem taxa do not extend beyond overall configuration. No

skull elements are lost or replaced in $nkx3.2^{-/-}$ zebrafish to achieve the anaspid/thelodont-like

micromery (Blom et al., 2001; Janvier, 1996; Märss et al., 2007). Therefore, in no way does the adult

$nkx3.2^{-/-}$ phenotype represent a reversal to an ancestral state. Although experimental phenocopies are

often interpreted as atavistic (ancestral reversal), alternative interpretations are seldom tested (Smith

and Schneider, 1998). Here, our $nkx3.2^{-/-}$ zebrafish present developmental plasticity as a testable

alternative to atavistic reversal to explain a phenocopy. By showing that a gnathostome can survive

without a jaw, the anaspid/thelodont-like *nkx3.2*$^{-/-}$ zebrafish also offer a comparative model to make

inferences about the functional morphology of these long-extinct agnathans.

Data availability. Sequence information, measurements, and landmark coordinates are available as

data supplements to this paper.

Author contributions. T.M. conceived project; T.M. and A.P.O. performed experiments resulting in

nkx3.2^{ua5011}; J.S. and N.N. performed experiments resulting in *nkx3.2*^{el802}; P.B. and B.G. provided μCT

scanning and histological sampling; T.M. and P.B. conducted morphometric analyses; T.M., P.B.,

A.R.P., J.G.C., D.G., and W.T.A. analyzed data; T.M. wrote the paper with inputs from P.B., A.P.O.,

J.S., A.R.P., J.G.C., D.G., and W.T.A.

Competing interests. We have no competing interests.

Funding. This work was supported by Natural Science and Engineering Research Council grants

RGPIN-2014-04863, RGPIN-2014-06311, and RGPIN-2015-06006 (to A.R.P., D.G., and W.T.A.,

respectively) and National Institute of Health grants R35DE027550 and K99 DE027218 (to J.G.C. and

J.S., respectively).

Acknowledgements. We thank B.F. Eames (University of Saskatchewan) for facilitating our

collaboration.

References

Adekeye EO. 1983. Ankylosis of the mandible: Analysis of 76 cases. *Journal of Oral and Maxillofacial Surgery* **41**:442–449. doi:10.1016/0278-2391(83)90129-5

Askary A, Mork L, Paul S, He X, Izuhara AK, Gopalakrishnan S, Ichida JK, McMahon AP, Dabizljevic S, Dale R, Mariani FV, Crump JG. 2015. Iroquois proteins promote skeletal joint formation by maintaining chondrocytes in an immature state. *Developmental Cell* **35**:358–365. doi:10.1016/j.devcel.2015.10.004

Askary A, Xu P, Barske L, Bay M, Bump P, Balczerski B, Bonaguidi MA, Crump JG. 2017. Genome-wide analysis of facial skeletal regionalization in zebrafish. *Development* **144**:2994–3005. doi:10.1242/dev.151712

Barske L, Askary A, Zuniga E, Balczerski B, Bump P, Nichols JT, Crump JG. 2016. Competition between Jagged-Notch and Endothelin1 signaling selectively restricts cartilage formation in the zebrafish upper face. *PLoS Genet* **12**:e1005967. doi:10.1371/journal.pgen.1005967

Bayram B, Uckan S, Cetinsahin A, Arman Ozcirpici A, Ozdemir H, Yazici C. 2010. Repositioning of the masseter muscle and its effect on skeletal growth. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology* **109**:e1–e5. doi:10.1016/j.tripleo.2009.12.041

Bixler D, Ward R, Gale DD. 1985. Agnathia-holoprosencephaly: a developmental field complex involving face and brain. Report of 3 cases. *J Craniofac Genet Dev Biol Suppl* **1**:241–249.

Blom H, Märss T. 2010. The interrelationships and evolutionary history of anaspid fishes In: Elliott DK, Maisey JG, Yu X, Miao D, editors. Morphology, Phylogeny and Palaeogeography of Fossil Fishes – Honoring Meemann Chang. Munich: Verlag Dr. Fredrich Pfeil.

Blom H, Märss T, Miller CG. 2001. Silurian and earliest Devonian birkeniid anaspids from the Northern Hemisphere. *Earth and Environmental Science Transactions of The Royal Society of Edinburgh* **92**:263–323. doi:10.1017/S0263593300000250

Brown DM, Marsh JL. 1990. Agnathia and Associated Malformations: A Case Report. *Cleft Palate Journal* **27**:415–418. doi:10.1597/1545-1569_1990_027_0415_aaamac_2.3.co_2

Cairns DM, Sato ME, Lee PG, Lassar AB, Zeng L. 2008. A gradient of Shh establishes mutually repressing somitic cell fates induced by Nkx3.2 and Pax3. *Developmental Biology* **323**:152–165. doi:10.1016/j.ydbio.2008.08.024

Caron MMJ, Emans PJ, Surtel DAM, van der Kraan PM, van Rhijn LW, Welting TJM. 2015. BAPX-1/NKX-3.2 acts as a chondrocyte hypertrophy molecular switch in osteoarthritis. *Arthritis & Rheumatology* **67**:2944–2956. doi:10.1002/art.39293

Cerny R, Cattell M, Sauka-Spengler T, Bronner-Fraser M, Yu F, Medeiros DM. 2010. Evidence for the prepattern/cooption model of vertebrate jaw evolution. *PNAS* **107**:17262–17267. doi:10.1073/pnas.1009304107

Chidzonga MM. 1999. Temporomandibular joint ankylosis: review of thirty-two cases. *British Journal of Oral and Maxillofacial Surgery* **37**:123–126. doi:10.1054/bjom.1997.0089

Cubbage CC, Mabee PM. 1996. Development of the cranium and paired fins in the zebrafish *Danio*
rerio (Ostariophysi, Cyprinidae). *Journal of Morphology* **229**:121–160.
doi:10.1002/(SICI)1097-4687(199608)229:2<121::AID-JMOR1>3.0.CO;2-4

Depew MJ, Compagnucci C. 2008. Tweaking the hinge and caps: testing a model of the organization of
jaws. *J Exp Zool* **310B**:315–335. doi:10.1002/jez.b.21205

Depew MJ, Simpson CA. 2006. 21st Century neontology and the comparative development of the
vertebrate skull. *Dev Dyn* **235**:1256–1291. doi:10.1002/dvdy.20796

Depew MJ, Simpson CA, Morasso M, Rubenstein JLR. 2005. Reassessing the Dlx code: the genetic
regulation of branchial arch skeletal pattern and development. *Journal of Anatomy* **207**:501–
561. doi:10.1111/j.1469-7580.2005.00487.x

Donoghue PCJ, Forey PL, Aldridge RJ. 2000. Conodont affinity and chordate phylogeny. *Biological*
Reviews **75**:191–251. doi:null

El-Sheikh MM, Medra AM, Warda MH. 1996. Bird face deformity secondary to bilateral
temporomandibular joint ankylosis. *Journal of Cranio-Maxillofacial Surgery* **24**:96–103.
doi:10.1016/S1010-5182(96)80020-5

Gagnon JA, Valen E, Thyme SB, Huang P, Ahkmetova L, Pauli A, Montague TG, Zimmerman S,
Richter C, Schier AF. 2014. Efficient mutagenesis by Cas9 protein-mediated oligonucleotide
insertion and large-scale assessment of single-guide RNAs. *PLOS ONE* **9**:e98186.
doi:10.1371/journal.pone.0098186

Gekas J, Li B, Kamnasaran D. 2010. Current perspectives on the etiology of agnathia-otocephaly.
European Journal of Medical Genetics **53**:358–366. doi:10.1016/j.ejmg.2010.09.002

Gillis JA, Modrell MS, Baker CVH. 2013. Developmental evidence for serial homology of the
vertebrate jaw and gill arch skeleton. *Nat Commun* **4**:1436. doi:10.1038/ncomms2429

Gomes FEF, Moraes RB, Luz JG de C. 2012. Effects of temporal muscle detachment and
coronoidotomy on facial growth in young rats. *Brazilian Oral Research* **26**:348–354.
doi:10.1590/S1806-83242012000400011

Hall BK. 2015. Bones and Cartilage: Developmental and Evolutionary Skeletal Biology. 2nd edition.
London: Academic Press.

Hawkins MB, Henke K, Harris MP. 2018. Latent developmental potential to form limb-like skeletal
structures in zebrafish. *bioRxiv* 450619. doi:10.1101/450619

Hecksher-Sørensen J, Watson RP, Lettice LA, Serup P, Eley L, Angelis CD, Ahlgren U, Hill RE. 2004.
The splanchnic mesodermal plate directs spleen and pancreatic laterality, and is regulated by
Bapx1/Nkx3.2. *Development* **131**:4665–4675. doi:10.1242/dev.01364

Hellemans J, Simon M, Dheedene A, Alanay Y, Mihci E, Rifai L, Sefiani A, van Bever Y, Meradji M,
Superti-Furga A, Mortier G. 2009. Homozygous inactivating mutations in the NKX3-2 gene
result in spondylo-megaepiphyseal-metaphyseal dysplasia. *The American Journal of Human*
Genetics **85**:916–922. doi:10.1016/j.ajhg.2009.11.005

Herbrand H, Pabst O, Hill R, Arnold H-H. 2002. Transcription factors Nkx3.1 and Nkx3.2 (Bapx1)
play an overlapping role in sclerotomal development of the mouse. *Mechanisms of*
Development **117**:217–224. doi:10.1016/S0925-4773(02)00207-1

Hernandez LP. 2000. Intraspecific scaling of feeding mechanics in an ontogenetic series of zebrafish,
Danio rerio. *Journal of Experimental Biology* **203**:3033–3043.

Hernandez LP, Bird NC, Staab KL. 2007. Using zebrafish to investigate cypriniform evolutionary
novelties: Functional development and evolutionary diversification of the kinethmoid. *Journal*
of Experimental Zoology Part B: Molecular and Developmental Evolution **308B**:625–641.
doi:10.1002/jez.b.21166

Horowitz SL, Shapiro HH. 1955. Modification of skull and jaw architecture following removal of the
masseter muscle in the rat. *American Journal of Physical Anthropology* **13**:301–308.
doi:10.1002/ajpa.1330130208

Howland HC, Merola S, Basarab JR. 2004. The allometry and scaling of the size of vertebrate eyes.
Vision Research **44**:2043–2065. doi:10.1016/j.visres.2004.03.023

Hu Y, Albertson RC. 2017. Baby fish working out: an epigenetic source of adaptive variation in the
cichlid jaw. *Proc R Soc B* **284**:20171018. doi:10.1098/rspb.2017.1018

Janvier P. 2007. Homologies and evolutionary transitions in early vertebrate history In: Anderson JS,
Sues H-D, editors. Major Transitions in Vertebrate Evolution. Bloomington: Indiana University
Press. pp. 57–121.

Janvier P. 1996. Early Vertebrates, Oxford Monographs on Geology and Geophysics. Oxford:
Clarendon Press.

Kawato Y, Hirao M, Ebina K, Shi K, Hashimoto J, Honjo Y, Yoshikawa H, Myoui A. 2012. Nkx3.2
promotes primary chondrogenic differentiation by upregulating Col2a1 transcription. *PLOS*
ONE **7**:e34703. doi:10.1371/journal.pone.0034703

Keating JN, Donoghue PCJ. 2016. Histology and affinity of anaspids, and the early evolution of the
vertebrate dermal skeleton. *Proc R Soc B* **283**:20152917. doi:10.1098/rspb.2015.2917

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of embryonic
development of the zebrafish. *Dev Dyn* **203**:253–310. doi:10.1002/aja.1002030302

Klingenberg CP. 2011. MorphoJ: an integrated software package for geometric morphometrics.
Molecular Ecology Resources **11**:353–357. doi:10.1111/j.1755-0998.2010.02924.x

Kuratani S. 2012. Evolution of the vertebrate jaw from developmental perspectives. *Evolution &*
Development **14**:76–92. doi:10.1111/j.1525-142X.2011.00523.x

Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. 2016. CHOPCHOP v2: a web tool for the
next generation of CRISPR genome engineering. *Nucleic Acids Res* **44**:W272–W276.
doi:10.1093/nar/gkw398

Lengner CJ, Hassan MQ, Serra RW, Lepper C, Wijnen AJ van, Stein JL, Lian JB, Stein GS. 2005.
Nkx3.2-mediated repression of Runx2 promotes chondrogenic differentiation. *J Biol Chem*
280:15872–15879. doi:10.1074/jbc.M411144200

Lettice L, Hecksher-Sørensen J, Hill R. 2001. The role of Bapx1 (Nkx3.2) in the development and
evolution of the axial skeleton. *Journal of Anatomy* **199**:181–187. doi:10.1046/j.1469-
7580.2001.19910181.x

Lifshitz J. 1976. Comparative anatomic study of mandibular growth in rats after bilateral resections of
superficial masseter, posterior temporal, and anterior digastric muscles. *J Dent Res* **55**:854–858.
doi:10.1177/00220345760550052301

Lukas P, Olsson L. 2018a. Bapx1 is required for jaw joint development in amphibians. *Evolution &*
Development **20**:192–206. doi:10.1111/ede.12267

Lukas P, Olsson L. 2018b. Bapx1 upregulation is associated with ectopic mandibular cartilage
development in amphibians. *Zoological Letters* **4**:16. doi:10.1186/s40851-018-0101-3

Manganello-Souza LC, Mariani PB. 2003. Temporomandibular joint ankylosis: Report of 14 cases.
International Journal of Oral and Maxillofacial Surgery **32**:24–29.
doi:10.1054/ijom.2002.0308

Märss T, Turner S, Karatajūtė-Talimaa V. 2007. Handbook of Paleoichthyology / "Agnatha" II.
Thelodonti. München: Verlag Dr. Friedrich Pfeil.

Medeiros DM, Crump JG. 2012. New perspectives on pharyngeal dorsoventral patterning in
development and evolution of the vertebrate jaw. *Developmental Biology* **371**:121–135.
doi:10.1016/j.ydbio.2012.08.026

Miller CT, Yelon D, Stainier DYR, Kimmel CB. 2003. Two endothelin 1 effectors, hand2 and bapx1,
pattern ventral pharyngeal cartilage and the jaw joint. *Development* **130**:1353–1365.
doi:10.1242/dev.00339

Miyashita T. 2016. Fishing for jaws in early vertebrate evolution: A novel hypothesis of mandibular
confinement. *Biological Reviews* **91**:611–657. doi:10.1111/brv.12187

Miyashita T. 2012. Comparative Analysis of the Anatomy of the Myxinoidea and the Ancestry of Early
Vertebrate Lineages (Unpublished M.Sc. thesis). Edmonton: University of Alberta.

Miyashita T, Coates MI, Farrar R, Larson P, Manning PL, Wogelius RA, Edwards NP, Anné J,
Bergmann U, Palmer AR, Currie PJ. 2019. Hagfish from the Cretaceous Tethys Sea and a
reconciliation of the morphological–molecular conflict in early vertebrate phylogeny. *PNAS*
116:2146–2151. doi:10.1073/pnas.1814794116

Miyazaki M, Yonemitsu I, Takei M, Kure-Hattori I, Ono T. 2016. The imbalance of masticatory
muscle activity affects the asymmetric growth of condylar cartilage and subchondral bone in
rats. *Archives of Oral Biology* **63**:22–31. doi:10.1016/j.archoralbio.2015.11.020

Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. 2014. CHOPCHOP: a CRISPR/Cas9 and
TALEN web tool for genome editing. *Nucleic Acids Res* **42**:W401–W407.
doi:10.1093/nar/gku410

Murtaugh LC, Zeng L, Chyung JH, Lassar AB. 2001. The chick transcriptional repressor Nkx3.2 acts
downstream of Shh to promote BMP-dependent axial chondrogenesis. *Developmental Cell*
1:411–422. doi:10.1016/S1534-5807(01)00039-9

Provot S, Kempf H, Murtaugh LC, Chung U, Kim D-W, Chyung J, Kronenberg HM, Lassar AB. 2006.
Nkx3.2/Bapx1 acts as a negative regulator of chondrocyte maturation. *Development* **133**:651–
662. doi:10.1242/dev.02258

Rodrigues L, Traina AA, Nakamai LF, Luz JG de C. 2009. Effects of the unilateral removal and
dissection of the masseter muscle on the facial growth of young rats. *Brazilian Oral Research*
23:89–95. doi:10.1590/S1806-83242009000100015

Rohlf FJ. 2018. tpsDIG. New York.

Rohlf FJ. 1999. Shape Statistics: Procrustes Superimpositions and Tangent Spaces. *J of Classification*
16:197–223. doi:10.1007/s003579900054

Sansom RS, Freedman K, Gabbott SE, Aldridge RJ, Purnell MA. 2010. Taphonomy and affinity of an
enigmatic Silurian vertebrate, *Jamoytius kerwoodi* White. *Palaeontology* **53**:1393–1409.
doi:10.1111/j.1475-4983.2010.01019.x

Sarnat BG. 1970. The face and jaws after surgical experimentation with the septovomeral region in
growing and adult rabbits. *Acta Oto-Laryngologica* **69**:1–30. doi:10.3109/00016487009131762

Sarnat BG, Muchnic H. 1971. Facial skeletal changes after mandibular condylectomy in the adult
monkey. *J Anat* **108**:323–338.

Schiffer C, Tariverdian G, Schiesser M, Thomas MC, Sergi C. 2002. Agnathia-otocephaly complex:
Report of three cases with involvement of two different Carnegie stages. *American Journal of
Medical Genetics* **112**:203–208. doi:10.1002/ajmg.10672

Schilling TF, Kimmel CB. 1997. Musculoskeletal patterning in the pharyngeal segments of the
zebrafish embryo. *Development* **124**:2945–2960.

Schneider A, Mijalski T, Schlange T, Dai W, Overbeek P, Arnold H-H, Brand T. 1999. The homeobox
gene it NKX3.2 is a target of left–right signalling and is expressed on opposite sides in chick
and mouse embryos. *Current Biology* **9**:911-S1. doi:10.1016/S0960-9822(99)80397-2

Smeeton J, Askary A, Gage Crump J. 2016. Building and maintaining joints by exquisite local control
of cell fate. *WIREs Dev Biol* **6**:e245. doi:10.1002/wdev.245

Smith KK, Schneider RA. 1998. Have gene knockouts caused evolutionary reversals in the mammalian
first arch? *BioEssays* **20**:245–255. doi:10.1002/(SICI)1521-1878(199803)20:3<245::AID-
BIES8>3.0.CO;2-Q

Standen EM, Du TY, Larsson HCE. 2014. Developmental plasticity and the origin of tetrapods. *Nature*
513:54–58. doi:10.1038/nature13708

Stewart TA, Albertson RC. 2010. Evolution of a unique predatory feeding apparatus: Functional
anatomy, development and a genetic locus for jaw laterality in Lake Tanganyika scale-eating
cichlids. *BMC Biology* **8**:8. doi:10.1186/1741-7007-8-8

Toledo LG, Cavalcanti SCXB, Corrêa L, Luz JGC. 2014. Effects of injury or removal of the articular
disc on maxillomandibular growth in young rats. *Journal of Oral and Maxillofacial Surgery*
72:2140–2147. doi:10.1016/j.joms.2014.06.445

Tucker AS, Watson RP, Lettice LA, Yamada G, Hill RE. 2004. Bapx1 regulates patterning in the
middle ear: Altered regulatory role in the transition from the proximal jaw during vertebrate
evolution. *Development* **131**:1235–1245. doi:10.1242/dev.01017

West-Eberhard MJ. 2005a. Developmental plasticity and the origin of species differences. *PNAS*
102:6543–6549. doi:10.1073/pnas.0501844102

West-Eberhard MJ. 2005b. Phenotypic accommodation: adaptive innovation due to developmental
plasticity. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*
304B:610–618. doi:10.1002/jez.b.21071

West-Eberhard MJ. 2003. Developmental Plasticity and Evolution, 1 edition. ed. Oxford ; New York:
Oxford University Press.

Wilson J, Tucker AS. 2004. Fgf and Bmp signals repress the expression of Bapx1 in the mandibular
mesenchyme and control the position of the developing jaw joint. *Developmental Biology*
266:138–150. doi:10.1016/j.ydbio.2003.10.012

Wilson MVH, Caldwell MW. 1998. The Furcacaudiformes: A new order of jawless vertebrates with
thelodont scales, based on articulated Silurian and Devonian fossils from northern Canada.
Journal of Vertebrate Paleontology **18**:10–29. doi:10.1080/02724634.1998.10011031

Wilson MVH, Caldwell MW. 1993. New Silurian and Devonian fork-tailed "thelodonts" are jawless
vertebrates with stomachs and deep bodies. *Nature* **361**:442–444. doi:10.1038/361442a0

Yamashita S, Andoh M, Ueno-Kudoh H, Sato T, Miyaki S, Asahara H. 2009. Sox9 directly promotes
Bapx1 gene expression to repress Runx2 in chondrocytes. *Experimental Cell Research*
315:2231–2240. doi:10.1016/j.yexcr.2009.03.008

Zeng L, Kempf H, Murtaugh LC, Sato ME, Lassar AB. 2002. Shh establishes an Nkx3.2/Sox9
autoregulatory loop that is maintained by BMP signals to induce somitic chondrogenesis. *Genes*
Dev **16**:1990–2005. doi:10.1101/gad.1008002

Figures

Figure 1. Null alleles of *nkx3.2* in zebrafish result in the jaw joint ankylosis.

(**a**) Mutations engineered with CRISPR/Cas9 or TALEN in the zebrafish gene *nkx3.2* that

encodes the homeobox transcription factor protein Nkx3.2 (a.k.a. Bapx1). Top: The gene *nkx3.2*

contains two exons (purple), the second of which encodes the homeobox domain (DNA binding

domain, blue). The diagram is approximately to scale (exon 1 = 310 bp), except the 3' untranslated

region (3'UTR). Allele *nkx3.2^{el802}* was generated with TALEN technology that deleted 20 bp from the

start of the gene (orange dashed line), eliminating the start codon. Allele *nkx3.2^{ua5011}* was engineered

with CRISPR/Cas9 to generate a 20 bp deletion (red box) and frameshift that is predicted to eliminate

the homeodomain. Bottom: Schematic of predicted protein following CRISPR mutagenesis. In the

allele *nkx3.2^{ua5011}*, the frameshift (20 bp deletion) is predicted to disrupt the translation and abrogate

production of the critical homeobox domain. This is predicted to produce random amino acids (red

hatching). (**a'**) Sequencing results from allele *nkx3.2^{ua5011}* (lower) compared to a wildtype zebrafish

(upper).

At 4 dpf, the chondrocrania are compared by *sox10*:eGFP expression in chondrocytes (**b–d**, **h**,

i) or by alcian blue staining of cartilages (**e–g**) between age-matched specimens. (**b**, **e**) The

chondrocranial morphology of wildtypes (AB background; *sox10*:eGFP) at 4 dpf in left lateral view, in

sox10:eGFP expression within chondrocytes (b) and alcian blue staining of cartilages (e). (**c**, **f**) The

chondrocranial morphology of *nkx3.2* heterozygous mutants (*nkx3.2^{+/ua5011}*; *sox10*:eGFP) at 4 dpf in

left lateral view, in *sox10*:eGFP expression within chondrocytes (c) and alcian blue staining of

cartilages (f). (**d**, **g**) The chondrocranial morphology of *nkx3.2* homozygous mutants (*nkx3.2^{ua5011}*;

sox10:eGFP) at 4 dpf in left lateral view, in *sox10*:eGFP expression within chondrocytes (d) and alcian

blue staining of cartilages (g). (**h**, **i**) Comparison of jaw morphology between age-matched wildtype

(AB background; *sox10*:eGFP) (h) and *nkx3.2* homozygous mutants (*nkx3.2^{ua5011}*; *sox10*:eGFP) (i) at 4

dpf in left lateral view, using *sox10*:eGFP expression within chondrocytes. White broken lines

delineate the mandibular cartilages. (**j**) Schematic drawing of the mandibular cartilages in zebrafish at 4

dpf in left lateral view, showing wildtype morphology (see panel e).

Abbreviations: asterisk (*), jaw joint; **aa**, anguloarticular; **ar**, articular; **bh**, basihyal; **ch**,

ceratohyal; **d**, dentary; **dpf**, days post fertilization; **hm**, hyomandibula; **ih**, interhyal; **j**, jugal; **ke**,

kinethmoid; **lp**, lower lip; **mc**, Meckel's cartilage; **m**, maxilla; **n**, nostril or aperture of nasohypophyseal

system; **o**, operculum; **pm**, premaxilla; **po**, preopercular; **pq**, palatoquadrate; **q**, quadrate; **ra**,

retroarticular process; **sm**, symplectic; **up**, upper lip.

Figure 2. Ontogeny of *nkx3.2*⁻/⁻ zebrafish documents drastic remodeling of the skull.

(**a**, **b**) Comparison of the skull morphology between age-matched wildtype (AB strain) (a) and

nkx3.2 homozygous mutant (*nkx3.2ua5011*) (b) at 21 dpf in right lateral view (inverted for consistency

with other panels), using alcian blue and alizarin red staining. (**c**, **d**) Comparison of skull morphology

between age-matched wildtype (AB) (c) and *nkx3.2* homozygous mutant (*nkx3.2ua5011*) (d) at 1mpf in

left lateral view, using alcian blue and alizarin red staining. (**e**) Detailed morphology of jaw skeleton in

nkx3.2 homozygous mutant (*nkx3.2ua5011*) at 1 mpf in right lateral view (inverted for consistency with

other panels), using alcian blue and alizarin red staining.

(**f**, **g**) Comparison of skull morphology between age-matched wildtype (AB) (f) and *nkx3.2*

homozygous mutant (*nkx3.2^{ua5011}*) (g) at 2 mpf in left lateral view, using alcian blue and alizarin red

staining. (**h**) Interpretive drawing of the specimen in panel g.

(**i**, **j**) Comparison of skull morphology between age-matched wildtype (AB) (i) and *nkx3.2*

homozygous mutant (*nkx3.2^{el802}*) (j) at 2 mpf in left lateral view, using alcian blue and alizarin red

staining. This second independent null allele of *nkx3.2* confirms the phenotypes reported herein.

(**k**–**n**) Detailed morphology of jaw joint in wildtype, or the joint-less interface between the

upper and lower jaws in *nkx3.2$^{-/-}$* mutants. (**k**, **l**) Detailed morphology of junction between upper and

lower jaws in age-matched wildtype (k) and *nkx3.2* homozygous mutant (*nkx3.2^{ua5011}*) (l) at 21 dpf in

left lateral view, using alcian blue and alizarin red staining. (**m**, **n**) Sagittal section of junction between

upper and lower jaws in age-matched wildtype (AB) (m) and *nkx3.2* homozygous mutant (*nkx3.2^{ua5011}*)

(n) at 1 mpf, stained with eosin and hematoxylin. (**n'**) Detailed view of section in panel n.

(**o**) Ontogeny of lower jaws in wildtype and *nkx3.2$^{-/-}$* zebrafish measured here as gape (angle

between upper and lower lips). See Data Supplement 2 for original measurements. Values from

wildtype and mutant (blue and red) are plotted as boxes of first and third quartile, with middle line

displaying mean, and whiskers communicating maximum and minimum values (*n* = sample size).

Photographs are male representative specimens at 60 dpf. Grey area indicates range of gape observed in

fossil anaspids (see Supplementary Information for details and Data Supplement 2 for original

measurements).

Figure 3. Adult *nkx3.2⁻/⁻* zebrafish converge onto overall head shapes of anaspids and thelodonts.

(**a**, **b**) Skulls of adult zebrafish in left lateral view via micro-computed tomography (μCT)

showing wildtype and *nkx3.2⁻/⁻* null mutant zebrafish (allele *ua5011*; 20 bp deletion in homeodomain).

The jaw joint is indicated by an asterisk (*) in wildtype (a) and absent in mutant (b). Adult mutants

display dramatic phenotypes in the jaw, snout, lips, and orobranchial regions. See also 3D rendering

(Movie 1). (**c**, **d**) Skulls of extinct jawless vertebrates, showing general resemblance to the skull shape

of *nkx3.2⁻/⁻* zebrafish. Morphometric analysis corroborates this similarity. (c) Skull of an anaspid

(birkeniid birkeniiform) from the Upper Silurian Cape Philips Formation of Cornwallis Island, Canada

(Geological Survey of Canada C-26661-005) in left lateral view. (d) Skull of *Sphenonectris turnerae*, a

thelodont (furcacaudiid furcacaudiform) from the Lower Devonian Road River Formation of Northwest

Territories, Canada (University of Alberta Laboratory for Vertebrate Palaeontology, specimen number

42212).

(**e**, **f**) Landmark-based geometric morphometric comparison of *nkx3.2* phenotype using the thin-

plate-spline Procrustes superimposition and principal component (PC) analysis. (**e**) Histogram showing

loading on each principal component in eigenvalue (%). PCs 1 and 5 are highlighted in blue. These two

PCs were chosen for comparison between groups, as these are the two largest components that set apart

nkx3.2 homozygous mutants and wildtype zebrafish from each other at adult stage. PCs 2-4, though

accounting for greater variation than PC5, primarily distinguish among anaspids, or within wildtype

and mutant zebrafish groups. The original dataset is available (Data Supplement 2). (**f**) A Cartesian plot

of PCs 1 and 5, comparing morphospace occupation between wildtypes (2 mpf) (blue), *nkx3.2*

homozygous mutants (2 mpf) (red), and anaspids (green), each with 90% ellipse. End of each axis is

labeled with a thin-plate-spline shape at that position (dark outline) against mean shape (grey

silhouette). A representative specimen is shown with landmarks labeled (yellow circles) for each group

in left lateral view. Each data point is a unique biological replicate (a specimen). The anaspid example

is *Birkenia elegans*, National Museum of Scotland specimen number 1929.5.6 from a Silurian locality

of Scotland. For anaspids, each data point is labeled with taxonomic identifications: B, *Birkenia*

elegans; C, Birkeniidae indet. from the Cornwallis Island; H, *Pharyngolepis oblongus*; L, *Lasanius*

problematicus; R, *Ryncholepis parvulus*; T, *Pterygolepis nitidus*.

(**g**) Simplified phylogenetic tree of vertebrates to illustrate distant relationships among the taxa

compared in this paper. A 60 dpf gravid female specimen is shown for *nkx3.2*$^{-/-}$ zebrafish to capture

variations among the mutants (compare with the male in panel e). Photographs of an anaspid and a

thelodont are the same individuals in the respective panels showing cranial morphology (d, e). Grey

triangle indicates a crown group (not to scale). Dark silhouettes and long branches indicate crown

groups, whereas grey silhouettes and short branches represent extinct lineages.

Movie

Movie 1. Feeding behaviour and 3D μCT scans of jawless zebrafish. Phenotypic comparison of *nkx3.2$^{-/-}$* and wildtype zebrafish. Three-dimensional rendering of μCT scan of the age-matched *nkx3.2$^{-/-}$* (ua5011) and wildtype (AB strain) zebrafish at 2 mpf, each followed by filming of feeding behavior on brine shrimp played at 1/10 original speed.

Supplementary Information

Methods

Zebrafish lines, husbandry, and ethical approvals

Zebrafish maintenance and experiments were approved as protocol number AUP00000077 by the

Animal Care and Use Committee: Biosciences at the University of Alberta as dictated by the Canadian

Council on Animal Care. Other zebrafish work was approved by the University of Southern California

Institutional Animal Care and Use Committee.

For *nkx3.2*^{ua5011}, we generated this mutant lineage against AB background. Cas 9 and sgRNAs

targeted for *nkx3.2* were coinjected with sgRNA that disrupts GFP GA5'

(CTCGGGCTGAAGCTCGGCG), at stages between fertilization and first cleavage, to fertilized eggs

collected from the crossing of ua3140 ubi:switch/AB+RH and the background AB line. At 3 dpf,

injected larvae were sorted for reduced expression of ubiquitous GFP and the presence of dsRed

fluorescence in heart. These larvae provided the P_0 population. Sequencing of genomic DNA extracted

from fin clips of the P_0 adults identified a female with a 20 bp deletion to the homeodomain-coding

region of *nkx3.2* (ua5011). The P_0 female carrying this mutation was crossed to the *sox10*:GFP

transgenic line, and progenies were sorted at 3 dpf for the presence of *sox10*:GFP expression and the

absence of the two markers (ubiquitous GFP expression and red fluorescent heart). ua5011

heterozygotes were identified by both sequencing of extracted gDNA and RFLP analysis using one

XmaI (NEB: R01805) restriction site within the deleted region (primers for genotyping: 5'–

GGACGAGACGGATCAGGAATC–3'; 5'–CACTCGGCGTGTTCGGTAAA–3'). These F_1

heterozygotes were incrossed for F_2 embryos, which were genotyped by RFLP analysis and phenotyped

at 4 dpf by identifying *sox10*:GFP-positive chondrocytes and staining cartilages using alcian blue.

Homozygotes were reared with a strictly small-grained diet to the adult stage. In this study,

nkx3.2^{ua5011/ua5011} represent F_2 generation derived from the P_0 mutant female and a wildtype male (AB;

sox10:GFP), whereas comparative wildtype (AB; *sox10*:GFP) come from incrossing of half-siblings of

the P_0 male.

Embryos were incubated at 28 ˚C, and treated with 0.003% PTU (1-phenyl-2-thiourea) in 10%

Hank's saline starting at 24 hpf. Larvae were introduced to the nursery at 1 week to 10 dpf. Genomic

DNA was extracted from clipped fins of 3-5 dpf larvae or from adults. Preserved tissues, embryos,

larvae, and adults were all fixed in 4% PFA, and stored in 100% EtOH or MeOH at -20 ˚C except for

adults (preserved in 70% EtOH at 4˚C).

For *nkx3.2*^{el802}, TALEN mutagenesis and husbandry protocols followed the previously

published description (Barske et al., 2016).

Molecular genetics

Nkx3.2 protein is a transcription factor with a homeobox DNA binding domain that is 100% conserved

in amino acid sequence among zebrafish, mouse, and human homologs. In zebrafish, a single *nkx3.2*

gene is apparent in the genome, and its homology to mammalian *NKX3.2* is strongly supported by gene

synteny: e.g. the neighbor genes flanking *nkx3.2* on zebrafish Chromosome 14 (*wdr1* and *bod1l1*) are

positioned coordinately in mouse, human and spotted gar.

To generate *nkx3.2*^{ua5011}, we designed sgRNA to five different targets near or within the *nkx3.2*

homeodomain, which were all injected:

GGCGGCCATCTGACGTCGCT

GGCTGACGCCAGCAGATCGG

AAGCAGCGGAAGAAGCGCTC

GAGCGCTTCTTCCGCTGCTT

GGCCGCGTTCTCCCACGCGC

These targets were selected using the web resource CHOPCHOP (Labun et al., 2016; Montague et al.,

2014). Following the protocol developed by Gagnon and colleagues (Gagnon et al., 2014), two

different oligonucleotides were ordered: one containing a target sequence led by the SP6 promoter

(ATTTAGGTGACACTATA) and followed by the overlapping region

(GTTTTAGAGCTAGAAATAGCAAG) of the reverse oligonucleotide; and the reverse containing the

constant, Cas9-binding domain of sgRNA. These oliogonucleotides were annealed after 5-minute

incubation at 95˚C, through graded cooling (-2˚C s^{-1} to 85˚C; -0.1˚C s^{-1} to 25˚C), and filled in for the

non-overlapping regions using T4 DNA polymerase (NEB: M0203S). To synthesize sgRNAs using

these templates, MegaScriptTM SP6 Transcription Kit (Ambion: AM1330) was used. The RNAs were

precipitated in ammonium acetate solution, suspended in UltraPure^{TM} H$_2$O, and stored in 2-3 µl

aliquots at -80 ℃. For injection, sgRNA(s) were diluted to 400-600 ng µl^{-1}, with 1 µl mixed with 1 µl

aliquot of Cas9 nuclease from Streptococcus pyogenes (NEB: #M0646) at 1 µg ml^{-1}. This solution was

mixed with 3 µl of 0.2M KCl, 0.2 % phenol red, and ddH$_2$O. The final injection volume per embryo

was approximately 5 nl, with 400-600 pg sgRNA and 1 ng Cas9 nuclease. For control, GFP 5'GA was

used at the stage P$_0$ when ubi:switch/RH+AB was crossed, which can be phenotyped by reduction of

ubiquitous GFP in the progenies with dsRed expression in heart.

Morphometrics

Rationales for the selection of metric traits. The purpose of our quantitative comparison is to test

phenotypic similarities and differences qualitatively identified in *nkx3.2$^{-/-}$* mutants with respect to

wildtype zebrafish and anaspids. On the one hand, the skulls of *nkx3.2$^{-/-}$* mutants clearly depart from

wildtype morphologically at adult stage (Fig. 2). To describe this morphological departure in greater

details, we will present comparison of skeletal growth between *nkx3.2$^{-/-}$* and wildtype elsewhere. On

the other hand, it is difficult to assess observed similarities between *nkx3.2$^{-/-}$* phenotype and the general

head configuration in anaspids. Zebrafish and anaspids are distant to each other phylogenetically: the

former is nested deep within, in the ascending order, cypriniforms, teleosts, neopterygians,

actinopterygians, osteichthyans, and gnathostomes, whereas anaspids represent either a stem

gnathostome or even a stem cyclostome lineage (Donoghue et al., 2000; Janvier, 2007, 1996; Keating

and Donoghue, 2016; Miyashita et al., 2019). The dermatocranium of a zebrafish is macromeric,

although that of an anaspid is largely micromeric with scales of acellular bone (Blom et al., 2001).

There is no morphological correspondence in individual elements of the skull roof between zebrafish

and anaspids. The parabranchial cavity is closed by the operculum in zebrafish, whereas each branchial

pouch had its own outlet in the series of external pores in anaspids (Blom et al., 2001). A single gene

mutation in *nkx3.2* did not reverse these, and other morphological differences accumulated after the last

common ancestor of zebrafish and anaspids. For morphometric comparison, we chose metric traits that

can be identified in both zebrafish and anaspids.

Linear morphometrics. The fixed open gape is similar between *nkx3.2*[-/-] mutants and anaspids. We

compared this qualitative similarity by taking the angle between supporting skeletal elements of the

upper and lower lips in both taxa. In zebrafish, the angle was taken by extrapolating the axis of the

premaxilla until it meets the axis of the dentary. At stages younger than the onset of dermal ossification

(4 and 14 dpfs), the angle was measured between the axes of the palatal process (palatoquadrate) and

Meckel's cartilage. In anaspids, the upper and lower lips are demarcated by a series of plates or

relatively larger scales, which allowed delineation of the gape angle.

In *nkx3.2*[-/-] zebrafish, the gape angle increases progressively with age. The timing of lower-lip

depression roughly coincides with the onset of dermal ossification. Variation of the gape angle among

individuals largely overlaps with that of anaspids (Fig. 2o). However, the increase in gape angle also

correlates with an increase in body size, and anaspids have much greater range of body size than

zebrafish. Unlike *nkx3.2*[-/-] zebrafish, the gape angle appears to vary independently of body size in

anaspids. No correlation exists between gape angle and orbit diameter in anaspids, regardless of

whether among those specimens in the size range of zebrafish or across the entire clade ($r = -0.114$; $P = $

0.306). Although eye size is generally negatively allometric in vertebrates (Howland et al., 2004), the

orbit diameter is one reliable, structurally intact metric trait in this clade, because most specimens are

not preserved in entire body length, and because other reference measurements (e.g., body height) are

affected by taphonomic deformation or simply not preserved in most specimens (Blom et al., 2001;

Blom and Märss, 2010; Janvier, 1996; Sansom et al., 2010).

Gape angle exhibits a roughly normal distribution within each of the age class of both *nkx3.2*[-/-]

and wildtype zebrafish and among anaspids (see Data Supplement 2). For anaspids, 95% of the

variation was calculated. This range is shown in Fig. 2o, independent of age (unknown in anaspid

samples) or size (unrelated to variation of gape angles among anaspids). Measurement values are

available in Data Supplement 2.

Geometric morphometrics. Eight landmarks were assigned to both zebrafish (2 mpf) and anaspid

samples for geometric morphometric comparison (Fig. 3f). These landmarks capture general

configuration of the heads (1: anterior tip of upper lip; 2: junction between upper and lower lips; 3:

anterior tip of lower lip; 4: nostril, or nasohypophyseal opening; 5: anterior extremity of orbit; 6:

posterior extremity of orbit; 7: trunk-head boundary at dorsal outline; 8: ventral point of hypobranchial

region). These landmarks describe structures homologous across vertebrates (landmarks 4, 5, 6) or

geometrically determined positions comparable across vertebrates. They are free of morphological

discontinuity between anaspids and gnathostomes (none of the landmarks represents anaspid- or

gnathostome-specific morphology). TpsDig (Rohlf, 2018) was used to place landmarks on two-

dimensional images of anaspid and zebrafish. The digitized file was entered into MorphoJ software

(Klingenberg, 2011) and all images were aligned using the anterior and posterior extremity of the

orbits. These landmark data were transformed using the procrustes superimposition method (Rohlf,

1999), and the resulting coordinates were compared using Principal Component Analysis (PCA). In

PCA, PC scores from anapsid, *nkx3.2*$^{+/+}$ and *nkx3.2*$^{-/-}$ zebrafish were grouped by equal frequency

ellipses with a P value of 0.95.

Rationales for selection of comparative taxa. Only anaspids were included as a comparative outgroup.

There are many other lineages of jawless vertebrates, including living cyclostomes, heterostracans,

thelodonts, galeaspids, pituriaspids, and osteostracans, in the order of nested hierarchy toward the

crown-group gnathostomes (Janvier, 2007, 1996). Living cyclostomes are difficult to compare as they

have an anguilliform profile and lack ossified skeletons, or cartilages unambiguously comparable to the

jaw/lip elements of the zebrafish skull (Miyashita, 2016, 2012; Miyashita et al., 2019). Extinct jawless

vertebrates generally have a depressiform profile(Janvier, 1996). Therefore, direct shape comparison is

difficult with the compressiform zebrafish and anaspids. Among non-depressiform jawless stem

gnathostomes, the lips are typically preserved poorly. Furcacaudiform thelodonts have a lateromedially

compressed body profile, and have the lip morphology consistent with that of anaspids (Wilson and

Caldwell, 1998, 1993). These similarities suggest that depressed lower lips are a general condition

along the gnathostome stem, as reconstructed conventionally across the stem group. However, we did

not include furcacaudiforms in our morphometric comparison. In furcacaudiforms, the lips are

demarcated by numerous scales, which make gape angle difficult to determine (Märss et al., 2007).

Landmarks cannot be assigned confidently either. The nasohypophyseal opening cannot be located

precisely because of the micromeric nature of the integument on the dorsal side of the head (Wilson

and Caldwell, 1998, 1993). The transition from head to trunk is ambiguous along the dorsal outline

because there is no apparent change in morphology of the scales (Wilson and Caldwell, 1998, 1993).

As such, anaspids remain as the only comparable outgroup taxon to *nkx3.2*$^{-/-}$ mutant zebrafish.

Tissue Preparation and Histology

Two-month-old *nkx3.2*$^{+/+}$ and *nkx3.2*$^{-/-}$ zebrafish were fixed in 4% paraformaldehyde for 24hrs.

Zebrafish were eviscerated prior to decalcification with 0.5M ethylenediaminetetraacetic acid (EDTA)

solution for 4 weeks. Samples were dehydrated post decalcification in a series of graded ethanol and

embedded in paraffin. Tissue blocks were embedded in a sagittal orientation and sections were cut at

7μm using a 820 Spencer microtome. Hematoxylin and eosin staining was performed on zebrafish

sections by firstly placing sections in an oven at 60 °C for 10 min. The deparaffinized sections were

rehydrated using xylene and graded ethanol (100%, 95%, 70%), followed by staining with hematoxylin

and eosin. The slides were then dehydrated and mounted using Permount.

Skeletal preparation

Alcian blue staining of cartilages partly followed the protocol provided by Michael Shapiro (University

of Utah). Specimens fixed in 4% PFA were rinsed with ddH$_2$O and transferred to 70% EtOH. Once

equilibrated, larvae were immersed in alcian blue solution (0.167 mg/ml alcian blue; 15% acetic acid;

70% EtOH), rinsed through EtOH/ddH$_2$O series, and washed in a saturated sodium borate solution.

Specimens were immersed in trypsin solution (0.125% trypsin; 30% sodium borate) overnight, washed

in 1% KOH solution, bleached in 0.15% H$_2$O$_2$ 0.1% KOH, 25% glycerol solution, immersed through a

1% KOH/glycerol graded series into 100% glycerol for storage. Specimens older than 21 dpf were

immersed in 0.005% alizarin red solution in 1% KOH overnight, after the first 1% KOH wash and

before bleaching in 0.15% H$_2$O$_2$ 0.1% KOH.

Filming

Wildtype (AB; *sox10*:GFP) and *nkx3.2*$^{ua5011/ua5011}$ were filmed at 2 months post fertilization to record

feeding behavior (Movie S1). A fish was placed in a 1.4L tank with dark background and given brine

shrimp larvae. Feeding was filmed using Canon EOS 760D at 30 frames s^{-1} in dimensions 1280×720

pixels. The films were cropped and assembled using iMovie (ver. 10.1.9, © 2001-2018 Apple Inc.) and

slowed to 1/10 original speed.

Imaging

Micro- Computed Tomography (µCT). 2-month-old zebrafish were scanned using MILabs µCT

scanner. Scans were reconstructed at a voxel size of 25µm. Images were analyzed using AVIZO 3-

dimensional software (Milabs, Utrecht, Netherlands). 2-Dimensional images used for linear and

geometric morphometrics were obtained from AVIZO.

Microscopy. Fluorescent images were acquired on a Zeiss Axio Observer.Z1 with LSM 700 confocal

microscope via ZEN 2010 software (version 6.0, Carl Zeiss MicroImaging, Oberkochen). Brightfield

imaging of stained preps was performed on a Leica MZ16F dissection microscope (Concord ON,

Canada) with 12.8 megapixel digital camera (DP72, Olympus; Richmond Hill ON, Canada).

DATA SUPPLEMENTS

Data supplement 1. Sequence information for the *nkx3.2* allele ua5011.

Data supplement 2. Gape angles in zebrafish; gape angles and orbit diameter in anaspids; thin-plate-

spline landmark data in zebrafish (wildtype and *nkx3.2*$^{-/-}$) and anaspids used for Procrustes

transformation, which were subjected to principal component analysis.

863