nkx3.2 mutant zebrafish accommodate jaw joint loss through a phenocopy of the head shapes of Paleozoic jawless fish

Tetsuto Miyashita[1, 2], Pranidhi Baddam[3], Joanna Smeeton[4], A. Phil Oel[1, 5], Natasha Natarajan[4], Brogan Gordon[1], A. Richard Palmer[1], J. Gage Crump[4], Daniel Graf[3, 6], and W. Ted Allison[1, 6]*

[1] Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2E9

[2] Department of Organismal Biology and Anatomy, University of Chicago, Chicago, IL 60637, USA

[3] Department of Dentistry, University of Alberta, Edmonton, Alberta, Canada T6G 2R3

[4] Department of Stem Cell Biology and Regenerative Medicine, W.M. Keck School of Medicine, University of Southern California, Los Angeles, CA 90033, USA

[5] Current Address: Developmental Biology Unit, European Molecular Biology Laboratory, Heidelberg, 69117, Germany

[6] Department of Medical Genetics, University of Alberta, Edmonton, Alberta, Canada T6G 2R7

* Corresponding author. ted.allison@ualberta.ca; 780-492-4430.

KEY WORDS: Skeletal remodeling, Developmental plasticity, Jaw, Joint, Zebrafish, Agnatha, *Nkx3.2*, *Bapx*

RUNNING TITLE: Functionally jawless zebrafish

ABSTRACT

The vertebrate jaw is a versatile feeding apparatus that facilitated explosive diversification. To

function, it requires a joint between the upper and lower jaws, so jaw joint defects — such as

osteoarthritis or even ankylosis — are often highly disruptive and difficult to study. To describe

consequences of jaw-joint dysfunction, we engineered two independent null alleles of a single jaw-joint

marker gene, *nkx3.2*, in zebrafish. These mutations caused zebrafish to become functionally jawless via

fusion of the upper and lower jaw cartilages (ankylosis). Despite lacking jaw joints, *nkx3.2* mutants

survive to adulthood and accommodate this defect by: a) remodeling their skulls; and b) altering their

behavior from suction feeding to ram feeding. As a result of remodeling, *nkx3.2* mutants developed

superficial similarities to the skull shapes observed in two lineages of ancient jawless vertebrates

(anaspids and furcacaudiid thelodonts), including: a fixed open gape, reduced snout, and enlarged

branchial region. However, no homology exists in individual skull elements between these taxa, and

most of the modified elements in the mutant zebrafish occur outside known expression domains of

nkx3.2. Therefore, we interpret the adult *nkx3.2* phenotype not as a reversal to an ancestral state, but as

convergence due to similar functional requirement of feeding without moveable jaws. This remarkable

convergence strongly suggests that jaw movements themselves dramatically influence the development

of jawed vertebrate skulls, which implies that functionally viable skull morphologies are finite, with or

without functional jaws. Because *nkx3.2* null zebrafish display prominent joint ankylosis, drastically

modified skull shape, and altered feeding behaviors, these mutants provide a unique model with which

to investigate mechanisms of skeletal remodeling and joint diseases.

INTRODUCTION

The jaw is a functionally versatile innovation that facilitated explosive diversification of gnathostomes

(a clade containing jawed vertebrates), but its basic structure is surprisingly simple and highly

conserved (Miyashita, 2016). A jaw consists of 'a hinge and caps': upper and lower skeletal levers

hinged at a jaw joint (Depew and Simpson, 2006). As the joint enables biting motions, its origin is

considered the final step in the evolutionary assembly of the vertebrate jaw (Cerny et al., 2010;

Kuratani, 2012; Miyashita, 2016). Across jawed vertebrates, the presumptive jaw joint is marked by the

expression of *nkx3.2*, an NK2 class homeobox gene (a.k.a. *bapx*), at the midheight of the embryonic

mandibular arch (Gillis et al., 2013; Lukas and Olsson, 2018a; Miller et al., 2003; Tucker et al., 2004).

Chondrogenesis dorsal to this expression domain gives rise to a palatoquadrate (upper jaw), whereas

chondrogenesis ventral to it forms Meckel's cartilage (lower jaw) (Medeiros and Crump, 2012). This

basic pattern remains conserved among jawed vertebrates, but later development varies. Marginal

bones arise intramembranously around the often endo-/peri-chondrally ossified jaw cartilages except in

chondrichthyans (sharks, rays, and skates) (Hall, 2015). In mammals, the jaw joint instead forms

between two such intramembranous bones (temporal and dentary), whereas the proximal jaw joint

becomes the malleus-incus interface that is, in mice, no longer affected by *Nkx3.2* knockout (Tucker et

al., 2004). Despite these variations after pharyngeal chondrogenesis, no gnathostome lineage

secondarily lost functional jaws.

By studying functional jaw loss in our new mutant zebrafish, we asked whether — and how —

jaw functions affect vertebrate skull shape during development. Clinically documented agnathia in

humans typically accompanies severe congenital disorders such as holoprosencephaly and otocephaly,

but jaw loss is clearly a secondary effect and not a cause in these cases (Bixler et al., 1985; Brown and

Marsh, 1990; Gekas et al., 2010; Schiffer et al., 2002). Instances of temporomandibular joint ankylosis

(stiffening due to bone fusion) may result from trauma or infection, or may be congenital (Adekeye,

1983; Chidzonga, 1999; Manganello-Souza and Mariani, 2003). If untreated, the ankylosis can lead to

the 'bird face' deformity (El-Sheikh et al., 1996). However, these cases do not fully document the

effects of functional jaw loss. In mammalian models, various jaw/skull deformations have been

induced by surgical resection, detachment, or repositioning of the jaw muscles and/or bones (Bayram et

al., 2010; Gomes et al., 2012; Horowitz and Shapiro, 1955; Lifshitz, 1976; Miyazaki et al., 2016;

Rodrigues et al., 2009; Sarnat, 1970; Sarnat and Muchnic, 1971; Toledo et al., 2014). These

manipulations occurred well after formation of the jaw skeleton and muscles, and the jaws remained

partially functional because of unilateral operations or non-comprehensive disruption. The defects and

deformities reported in these studies imply: a) jaw movements are potentially an important factor in

shaping the skulls; and b) any allele disrupting jaw movements would be generally maladaptive.

Nevertheless, these implications are difficult to explore without an accessible experimental model.

To fill this gap, we engineered two distinct null alleles of *nkx3.2* in zebrafish. Previously,

transient knockdown of *nkx3.2* during early development (using morpholinos) had shown fusion of the

nascent jaw cartilages in both zebrafish and frogs (Lukas and Olsson, 2018a; Miller et al., 2003). We

confirmed in zebrafish that the mutants reproduce this phenotype. Surprisingly, mutant zebrafish are

viable — despite loss of the jaw joint — and grow through to adulthood. Functionally jawless as a

result, *nkx3.2*$^{-/-}$ zebrafish dramatically alter skull shape late in ontogeny to facilitate feeding, with the

mouth fixed open, the snout reduced, and the branchial region expanded. This open-mouth phenotype,

previously unknown in zebrafish or any other jawed vertebrates, also occurred in two extinct lineages

of 400-million-plus year-old jawless vertebrates, anaspids and thelodonts. Even though they share no

homology in individual facial bones, *nkx3.2*$^{-/-}$ fish accommodate loss of a functional jaw by converging

onto these ancient, distantly related agnathan head shapes. Thus, *nkx3.2* mutant zebrafish provide a

unique model for both skeletal remodeling and joint diseases such as osteoarthritis, and to reevaluate

evolutionary implications of phenocopies in general.

MATERIALS AND METHODS

Animal Ethics

Zebrafish maintenance and experiments were approved as protocol number AUP00000077 by the

Animal Care and Use Committee: Biosciences at the University of Alberta as dictated by the Canadian

Council on Animal Care. Other zebrafish work was approved by the University of Southern California

Institutional Animal Care and Use Committee.

Animal husbandry

Embryos were incubated at 28 ˚C, and treated with 0.003% PTU (1-phenyl-2-thiourea) in 10% Hank's

saline starting at 24 hpf. Larvae were introduced to the nursery at 1 week to 10 dpf. Genomic DNA was

extracted from clipped fins of 3-5 dpf larvae or from adults. Preserved tissues, embryos, larvae, and

adults were all fixed in 4% PFA, and stored in 100% EtOH or MeOH at -20 ˚C except for adults

(preserved in 70% EtOH at 4˚C).

Molecular genetics

Nkx3.2 protein is a transcription factor with a homeobox DNA binding domain that is 100% conserved

in amino acid sequence among zebrafish, mouse, and human homologs. In zebrafish, a single *nkx3.2*

gene is apparent in the genome, and its homology to mammalian *NKX3.2* is strongly supported by gene

synteny: e.g. the neighbor genes flanking *nkx3.2* on zebrafish Chromosome 14 (*wdr1* and *bod1l1*) are

positioned coordinately in mouse, human and spotted gar.

Two disparate regions of the gene were targeted by CRISPR guide RNA (gRNA) or TALENs,

producing two disparate null alleles that produced similar phenotypes (Fig. 1A). One allele (ua5011)

was engineered with CRISPR/Cas9 (Gagnon et al., 2014) targeted at the beginning of the

homeodomain, and it harbors a 20 bp deletion resulting in a frameshift (Fig. 1A; Data Supplement 1).

The disrupted translation of codons is predicted to abrogate production of the critical homeobox

domain, and instead produce random amino acids. This is predicted to produce a non-functional

Nkx3.2 and a null allele. A disparate allele (el802) was generated using TALENs (Barske et al., 2016)

targeted at the start of the gene. This produced a stably inherited gene with 20 bp deletion, removing

the translation start codon (Fig. 1A). The allele $nkx3.2^{el802}$ is predicted to not produce Nkx3.2 protein.

Morphologically, these two alleles are not readily distinguishable from each other (see Results).

CRISPR

To generate $nkx3.2^{ua5011}$, we designed sgRNA to five different targets near or within the $nkx3.2$

homeodomain, which were all injected:

GGCGGCCATCTGACGTCGCT

GGCTGACGCCAGCAGATCGG

AAGCAGCGGAAGAAGCGCTC

GAGCGCTTCTTCCGCTGCTT

GGCCGCGTTCTCCCACGCGC

These targets were selected using the web resource CHOPCHOP (Labun et al., 2016; Montague et al.,

2014). Following the protocol developed by Gagnon and colleagues (2014), two different

oligonucleotides were ordered: one containing a target sequence led by the SP6 promoter

(ATTTAGGTGACACTATA) and followed by the overlapping region

(GTTTTAGAGCTAGAAATAGCAAG) of the reverse oligonucleotide; and the reverse containing the

constant, Cas9-binding domain of sgRNA. These oliogonucleotides were annealed after 5-minute

incubation at 95°C, through graded cooling (-2°C s^{-1} to 85°C; -0.1°C s^{-1} to 25°C), and filled in for the

non-overlapping regions using T4 DNA polymerase (NEB: M0203S). To synthesize sgRNAs using

these templates, MegaScript^{TM} SP6 Transcription Kit (Ambion: AM1330) was used. The RNAs were

precipitated in ammonium acetate solution, suspended in UltraPure^{TM} H$_2$O, and stored in 2-3 μl

aliquots at -80 ℃. For injection, sgRNA(s) were diluted to 400-600 ng μl^{-1}, with 1 μl mixed with 1 μl

aliquot of Cas9 nuclease from *Streptococcus pyogenes* (NEB: #M0646) at 1 μg ml^{-1}. This solution was

mixed with 3 μl of 0.2M KCl, 0.2 % phenol red, and ddH$_2$O. The final injection volume per embryo

was approximately 5 nl, with 400-600 pg sgRNA and 1 ng Cas9 nuclease. For control, GFP 5'GA was

used at the stage P$_0$ when ubi:switch/RH+AB was crossed, which can be phenotyped by reduction of

ubiquitous GFP in the progenies with dsRed expression in heart.

We generated $nkx3.2^{ua5011}$ against AB background. Cas 9 and sgRNAs targeted for $nkx3.2$ were

coinjected with sgRNA that disrupts GFP GA5' (CTCGGGCTGAAGCTCGGCG), at stages between

fertilization and first cleavage, to fertilized eggs collected from the crossing of ua3140

ubi:switch/AB+RH and the background AB line. At 3 dpf, injected larvae were sorted for reduced

expression of ubiquitous GFP and the presence of dsRed fluorescence in heart. These larvae provided

the P$_0$ population. Sequencing of genomic DNA extracted from fin clips of the P$_0$ adults identified a

female with a 20 bp deletion to the homeodomain-coding region of *nkx3.2* (ua5011). The P_0 female

carrying this mutation was crossed to the *sox10*:GFP transgenic line, and progenies were sorted at 3 dpf

for the presence of *sox10*:GFP expression and the absence of the two markers (ubiquitous GFP

expression and red fluorescent heart). ua5011 heterozygotes were identified by both sequencing of

extracted gDNA and Restriction Fragment Length Polymorphism (RFLP) analysis using one XmaI

(NEB: R01805) restriction site within the deleted region (primers for genotyping: 5'–

GGACGAGACGGATCAGGAATC–3'; 5'–CACTCGGCGTGTTCGGTAAA–3'). These F_1

heterozygotes were incrossed for F_2 embryos, which were genotyped by RFLP analysis and phenotyped

at 4 dpf by identifying *sox10*:GFP-positive chondrocytes and staining cartilages using alcian blue.

Homozygotes were reared with a strictly small-grained diet to the adult stage. In this study,

nkx3.2$^{ua5011/ua5011}$ represent F_2 generation derived from the P_0 mutant female and a wildtype male (AB;

sox10:GFP), whereas comparative wildtype (AB; *sox10*:GFP) come from incrossing of half-siblings of

the P_0 male.

Tissue preparation and histology

One- and two-month-old *nkx3.2*$^{+/+}$ and *nkx3.2*$^{-/-}$ zebrafish were fixed in 4% paraformaldehyde for

24hrs. Zebrafish were eviscerated prior to decalcification with 0.5M ethylenediaminetetraacetic acid

(EDTA) solution for 4 weeks. Samples were dehydrated post decalcification in a series of graded

ethanol and embedded in paraffin. Tissue blocks were embedded in a sagittal orientation and sections

were cut at 7μm using a 820 Spencer microtome. Hematoxylin and eosin staining was performed on

zebrafish sections by firstly placing sections in an oven at 60 °C for 10 min. The deparaffinized

sections were rehydrated using xylene and graded ethanol (100%, 95%, 70%), followed by staining

with hematoxylin and eosin. The slides were then dehydrated and mounted using Permount.

Skeletal preparation

Alcian blue staining of cartilages partly followed the protocol provided by Michael Shapiro (University

of Utah). Specimens fixed in 4% PFA were rinsed with ddH$_2$O and transferred to 70% EtOH. Once

equilibrated, larvae were immersed in alcian blue solution (0.167 mg/ml alcian blue; 15% acetic acid;

70% EtOH), rinsed through EtOH/ddH$_2$O series, and washed in a saturated sodium borate solution.

Specimens were immersed in trypsin solution (0.125% trypsin; 30% sodium borate) overnight, washed

in 1% KOH solution, bleached in 0.15% H$_2$O$_2$ 0.1% KOH, 25% glycerol solution, immersed through a

1% KOH/glycerol graded series into 100% glycerol for storage. Specimens older than 21 dpf were

immersed in 0.005% alizarin red solution in 1% KOH overnight, after the first 1% KOH wash and

before bleaching in 0.15% H_2O_2 0.1% KOH.

Filming

Wildtype (AB; *sox10*:GFP) and *nkx3.2*[ua5011/ua5011] were filmed at 2 months post fertilization to record

feeding behavior (Movie S1). A fish was placed in a 1.4L tank with dark background and given brine

shrimp larvae. Feeding was filmed using Canon EOS 760D at 30 frames s^{-1} in dimensions 1280×720

pixels. The films were cropped and assembled using iMovie (ver. 10.1.9, © 2001-2018 Apple Inc.) and

slowed to 1/10 original speed.

Imaging

Micro-computed tomography (µCT). Two-month-old zebrafish were scanned using MILabs µCT

scanner. Scans were reconstructed at a voxel size of 25µm. Images were analyzed using AVIZO 3-

dimensional software (Milabs, Utrecht, Netherlands). 2-Dimensional images used for linear and

geometric morphometrics were obtained from AVIZO.

Microscopy. Fluorescent images were acquired on a Zeiss Axio Observer.Z1 with LSM 700 confocal

microscope via ZEN 2010 software (version 6.0, Carl Zeiss MicroImaging, Oberkochen). Brightfield

imaging of stained preps was performed on a Leica MZ16F dissection microscope (Concord ON,

Canada) with 12.8 megapixel digital camera (DP72, Olympus; Richmond Hill ON, Canada).

Morphometrics

Rationales for morphometric comparison. The purpose of our quantitative comparison is to test

phenotypic similarities and differences qualitatively identified in *nkx3.2*[-/-] mutants with respect to

wildtype zebrafish and anaspids (and thelodonts for gape angles). On the one hand, the skulls of *nkx3.2*[-/-]

/- mutants clearly depart from wildtype morphologically at adult stage (Fig. 2). To describe this

morphological departure in greater details, we will present comparison of skeletal growth between

nkx3.2[-/-] and wildtype elsewhere. On the other hand, it is difficult to assess observed similarities

between *nkx3.2*[-/-] phenotype and the general head configuration in anaspids. Zebrafish and anaspids are

distant to each other phylogenetically: the former is nested deep within, in the ascending order,

cypriniforms, teleosts, neopterygians, actinopterygians, osteichthyans, and gnathostomes, whereas

anaspids represent either a stem gnathostome or even a stem cyclostome lineage (Donoghue et al.,

2000; Janvier, 2007, 1996; Keating and Donoghue, 2016; Miyashita et al., 2019). The dermatocranium

of a zebrafish is macromeric, although that of an anaspid is largely micromeric with scales of acellular

bone (Blom et al., 2001). There is no morphological correspondence in individual elements of the skull

roof between zebrafish and anaspids. The parabranchial cavity is closed by the operculum in zebrafish,

whereas each branchial pouch had its own outlet in the series of external pores in anaspids (Blom et al.,

2001). A single gene mutation in *nkx3.2* did not reverse these, and other morphological differences

accumulated after the last common ancestor of zebrafish and anaspids. For morphometric comparison,

we chose metric traits that can be identified in both zebrafish and anaspids.

Linear morphometrics. The fixed open gape is similar between $nkx3.2^{-/-}$ mutants and anaspids. We

compared this trait by taking the angle between supporting skeletal elements of the upper and lower

lips in both taxa. In zebrafish, the angle was taken by extrapolating the axis of the premaxilla until it

meets the axis of the dentary. At stages younger than the onset of dermal ossification (4 and 14 dpfs),

the angle was measured between the axes of the palatal process (palatoquadrate) and Meckel's

cartilage. In anaspids, the upper and lower lips are demarcated by a series of plates or relatively larger

scales, which allowed delineation of the gape angle (measured at where the extrapolated upper and

lower lip margins meet). The angle was measured similarly in thelodonts, except that the lip margins

were identified along the series of small marginal scales. Gape angle exhibits a roughly normal

distribution within each of the age class of both $nkx3.2^{-/-}$ and wildtype zebrafish and among anaspids

(see Data Supplement 2). In addition to gape angles, we measured lengths of skulls and lower jaws in

zebrafish, and orbit diameter in anaspids and thelodonts. Original measurements are available in Data

Supplement 2.

In $nkx3.2^{-/-}$ zebrafish, the gape angle increases progressively with age, and thus with increasing

body size. Anaspids and thelodonts overall have much greater range of body size than zebrafish. Unlike

$nkx3.2^{-/-}$ zebrafish, however, the gape angle appears to vary independently of body size in anaspids. No

correlation exists between gape angle and orbit diameter in anaspids, regardless of whether among

those specimens falling in the size range of zebrafish or across the entire clade ($r = -0.114$; $P = 0.306$).

Although eye size is generally negatively allometric in vertebrates (Howland et al., 2004), the orbit

diameter is one reliable, structurally intact metric trait in this clade, because most specimens are not

preserved in entire body length, and because other reference measurements (e.g., body height) are

affected by taphonomic deformation or simply not preserved in most specimens (Blom et al., 2001;

Blom and Märss, 2010; Janvier, 1996; Sansom et al., 2010). In thelodonts, the relationship remains to

be tested between body size and gape angles because of small sample size ($n = 5$).

Geometric morphometics. Eight landmarks were assigned to both zebrafish (2 mpf) and anaspid

samples for geometric morphometric comparison. These landmarks capture general configuration of

the heads (1: anterior tip of upper lip; 2: junction between upper and lower lips; 3: anterior tip of lower

lip; 4: nostril, or nasohypophyseal opening; 5: anterior extremity of orbit; 6: posterior extremity of

orbit; 7: trunk-head boundary at dorsal outline; 8: ventral point of hypobranchial region). These

landmarks describe structures homologous across vertebrates (landmarks 4, 5, 6) or geometrically

determined positions comparable across vertebrates. They are free of morphological discontinuity

between anaspids and gnathostomes (none of the landmarks represents anaspid- or gnathostome-

specific morphology). TpsDig (Rohlf, 2018) was used to place landmarks on two-dimensional images

of anaspid and zebrafish. The digitized file was entered into MorphoJ software (Klingenberg, 2011)

and all images were aligned using the anterior and posterior extremity of the orbits. These landmark

data were transformed using the procrustes superimposition method (Rohlf, 1999), and the resulting

coordinates were compared using Principal Component Analysis (PCA). In PCA, PC scores from

anapsid, *nkx3.2*$^{+/+}$ and *nkx3.2*$^{-/-}$ zebrafish were grouped by equal frequency ellipses with a P value of

0.95.

Rationales for selection of comparative taxa in geometric morphometrics. From the pool of nearly a

thousand catalogued specimens of anaspids, we selected a total of 70 specimens that show lateral

compression during the fossilization process, with the least taphonomic artifact, to reflect lateral view

of the heads. Furcacaudiid thelodonts were excluded from geometric morphometrics because only a

handful of exceptionally preserved specimens are available. The sample size is small for this latter

group ($n < 5$), and all such specimens were collected from a single locality, making it difficult to

identify (and thus control for) taphonomic artifacts. In addition, landmarks cannot be assigned

confidently in this group. The nasohypophyseal opening cannot be located precisely because of the

micromeric nature of the integument on the dorsal side of the head (Wilson and Caldwell, 1998, 1993).

The transition from head to trunk is ambiguous along the dorsal outline because there is no apparent

change in morphology of the scales (Wilson and Caldwell, 1998, 1993).

There are many other lineages of jawless vertebrates, including living cyclostomes,

heterostracans, thelodonts, galeaspids, pituriaspids, and osteostracans, in the order of nested hierarchy

toward the crown-group gnathostomes (Janvier, 2007, 1996). Living cyclostomes are difficult to

compare as they have an anguilliform profile and lack ossified skeletons, or cartilages unambiguously

comparable to the jaw/lip elements of the zebrafish skull (Miyashita, 2016, 2012; Miyashita et al.,

2019). Extinct jawless vertebrates generally have a depressiform profile (Janvier, 1996). Therefore,

direct shape comparison is difficult with the compressiform zebrafish and anaspids.

Among non-depressiform jawless stem gnathostomes, the lips are typically preserved poorly.

Furcacaudiid thelodonts have a lateromedially compressed body profile, and have the lip morphology

consistent with that of anaspids (Wilson and Caldwell, 1998, 1993). These similarities suggest that

depressed lower lips are a general condition along the gnathostome stem, as reconstructed

conventionally across the stem group.

RESULTS

Jaw joint is ankylosed in *nkx3.2* null alleles

At 4 days post fertilization (dpf), $nkx3.2^{-/-}$ zebrafish replicated the *nkx3.2* morpholino knockdown

phenotype: the absence of a jaw joint (Miller et al., 2003). The palatoquadrate and Meckel's cartilage

fused together, and the retroarticular process was absent (Fig. 1D, G, I). Apart from joint ankylosis, no

marked differences were apparent in overall morphology or survival rates between $nkx3.2^{-/-}$ mutants

and wildtype (some minor difference in skull size and lower jaw proportions are discussed in the next

section). Heterozygotes were morphologically indistinguishable from wildtype (Fig. 1C, F), and the

nkx3.2 alleles displayed recessive Mendelian inheritance (F_2 genotypes followed Mendelian ratio;

heterozygotes developed wildtype morphology). Remarkably, these functionally jawless homozygous

mutants survived beyond early larval stages.

Functionally jawless $nkx3.2^{-/-}$ zebrafish modify skull shapes late in ontogeny

Contrary to the maladaptive nature of jaw dysfunctions in general, $nkx3.2^{-/-}$ zebrafish continued to grow

without a jaw joint. Marked phenotypic differences against wildtype began to emerge between the 2^{nd}

and 3^{rd} weeks post fertilization (Fig. 2). The lower jaw became downturned in $nkx3.2^{-/-}$ fish resulting in

a rigidly fixed open mouth, whereas both upper and lower jaws were upturned in wildtype zebrafish

(Fig. 2A, B). This timing coincides with the onset of ossification and the period of active feeding in

normal juveniles (Cubbage and Mabee, 1996; Kimmel et al., 1995). Although the palatoquadrate and

Meckel's cartilage remained fused in $nkx3.2^{-/-}$ mutants, skeletal staining reveals that the upper and

lower jaw elements ossified independently of each other — still without a ball-and-socket joint

structure (Fig. 3A, B). All skull elements in $nkx3.2^{-/-}$ mutants ossified without apparent delay.

The lower jaws were increasingly turned downward in $nkx3.2^{-/-}$ by the end of the first month,

resulting in a greater gape (Fig. 2D, E). The jaw joint was still absent in $nkx3.2^{-/-}$ mutants: a sheet of

perichondrium lies between the ossifying quadrate and articular — so the two bones remained distinct

elements — but this interface had none of those essential components of synovial diarthrosis (Fig. 3D,

E). Many other osteological differences emerged by this stage. Normally, the premaxilla and the

maxilla swing forward to sit nearly vertical and are hinged by the kinethmoid for suction feeding

(Hernandez, 2000; Hernandez et al., 2007) (Fig. 2C). In *nkx3.2*$^{-/-}$ mutants, however, the premaxilla and

the maxilla became oriented posteroventrally and abutted against the anterior margin of the orbit (Fig.

2D). The kinethmoid was reduced into a fused bony process, unlike a rod-like hinge element in

wildtype (Hernandez et al., 2007). The downturned lower jaws of the *nkx3.2*$^{-/-}$ mutants were relatively

shorter than the normal lower jaws of wildtypes. The basihyal protruded anteroventrally as much as the

lower jaw, implying that the muscle connecting those two elements (m. intermandibularis posterior)

(Schilling and Kimmel, 1997) may be responsible for the lower jaw orientation. As a result of these

modifications, *nkx3.2*$^{-/-}$ mutants had a shorter snout, a fixed open gape, and a dorsoventrally tall profile.

Linear morphometrics corroborated the departure from normal morphology in *nkx3.2*$^{-/-}$ mutants

in the latter half of the first month (14 dpf onward). For absolute size, no significant difference (*P* >

0.05) in skull length emerged between wildtype and *nkx3.2*$^{-/-}$ mutants except at 4 dpf (*t* = 2.202; *P* =

0.338) (Fig. 4B). Also at this stage, the lower jaws appear to be shorter relative to skull length in the

mutants than in the wildtypes (*t* = 2.809; *P* = 0.0078) (Fig. 4C). These minor but statistically significant

differences at 4 dpf may be a direct consequence of the ankylosis between palatoquadrate and Meckel's

cartilage. By the second week, however, differences between the mutants and wildtypes became non-

significant in these metric traits. The lower jaw depression (gape angle > 45º) in the mutants was

pronounced at 21 dpf (*t* = -11.834; *P* << 0.01) (Fig. 4A), but proportional changes to lower jaw lengths

in the same mutants were only expressed in significant magnitude at 1 month (21 dpf: *t* = 1.7225; *P* =

0.091559; 30 dpf: *t* = 10.56; *P* << 0.01) (Fig. 4C). This lag between the two traits indicates that the rate

of skeletal growth in the jaws followed changes in their orientation for a greater gape (and thus

resulting in functional shift). Morphological variations within a cohort of *nkx3.2*$^{-/-}$ mutants were greater

at this stage than in any other, as indicated by the range of variation in orientations and relative lengths

of the lower jaws (Fig. 4A, C).

The adult *nkx3.2*$^{-/-}$ phenotype accommodates functional jaw loss

At 2 months of age and approaching sexual maturity nearly all *nkx3.2*$^{-/-}$ mutants had a gape angle

greater than 90 degrees (Fig. 4A). These *nkx3.2*$^{-/-}$ adults continued to be characterized by the

morphological differences identified at 1 mpf. The nostrils sat between the eyes because the snout was

reduced in length relative to wildtype zebrafish. The skulls appeared to be more highly ossified in

nkx3.2$^{-/-}$ mutants than in wildtype, where massive bones and cartilages were identified around the

interface of quadrate and articular, in the lower branchial region, and in the laterally expanded

operculum (Figs. 2J, G, M, N, 5B; Movie S1). All homozygous mutants showed the descriptive skeletal

traits identified here. Morphologically, the phenotypic effects of the two alleles (*nkx3.2^{ua5011}* and

nkx3.2^{el802}) were virtually indistinguishable from each other at respective ages (Fig. 2G, J), given

variations within a strain (for examples, see variation in gape angles, relative lengths of lower jaws, or

sexual dimorphism in *nkx3.2^{ua5011}*: Figs.2M, N, 4A, C).

337   With their modified skulls, adult *nkx3.2$^{-/-}$* mutants exhibited feeding behaviors markedly

different from wildtype (Fig. 5; Movie S1). Wildtype zebrafish feed by suction using rapid lower jaw

depression and a forward swing of the mobile premaxilla-maxilla complex shortly followed by opening

of the operculum (Fig. 5A), consistent with general teleost feeding mechanics (Alexander, 1970, 1969;

Lauder, 1980, 1979; Westneat, 2005, 2004). In contrast, adult *nkx3.2$^{-/-}$* mutants were constrained by the

fixed upper jaw unit and open gape. They instead showed ram feeding behaviors (swimming through

food) (Fig. 5B). The jaws remained fixed, and no significant dorsoventral movement was observed.

Whereas in wildtype one complete cycle of the jaw opening and closing took approximately 80

milliseconds (and approximately a tenth of a second to the closure of the operculum), *nkx3.2$^{-/-}$* mutants

required double that time from changing direction of swimming toward food (0 s) to doing so again

away from the food (0.2 s) in this particular feeding episode in Fig. 5B. A detailed analysis of the

feeding mechanics is beyond the scope of this paper and is currently the focus of our study, with

filming at higher speed and resolution.

350   This ram-feeding behavior was correlated with skull remodeling in *nkx3.2$^{-/-}$* mutants. In

zebrafish skulls, the bones form endochondrally (quadrate, anguloarticular, basihyal) or

intramembranously (premaxilla, maxilla, dentary, jugal, opercular, preopercular) (Cubbage and Mabee,

1996; Schilling and Kimmel, 1997). Much of the skeletal remodeling observed in adult *nkx3.2$^{-/-}$*

mutants occurred in the intramembranous bones — spatially and temporarily well outside the known

expression domain of *nkx3.2* (Askary et al., 2017; Miller et al., 2003). Until past 1 mpf, the fusion

between jaw cartilages was not completely ossified in these mutants, potentially allowing plastic

remodeling (Fig. 2e, l). These observations suggest that *nkx3.2$^{-/-}$* zebrafish accommodate functional

jawlessness through remodeling of the skull and changes to feeding behavior.

nkx3.2$^{-/-}$ zebrafish converge onto agnathans in overall head shapes

Through this dramatic remodeling of the skull, *nkx3.2$^{-/-}$* zebrafish assumed a head shape reminiscent of

two lineages of extinct jawless vertebrates that have laterally compressed body profiles: 1) birkeniiform

anaspids (Fig. 6C), stem cyclostomes known mostly from the Silurian period (Blom et al., 2001;

Miyashita et al., 2019); and 2) furcacaudiid thelodonts (Fig. 6D), much more elusive stem

gnathostomes known from the Silurian and Devonian periods (Märss et al., 2007; Wilson and Caldwell,

1998). Qualitatively, the resemblance is particularly striking in overall head shape characters,

including: fixed gape (depressed lower lips), shortened snout, interorbital position of nostril,

proportionally large branchial region, and massive parietal region behind the occiput.

In linear morphometrics, the gape angle between upper and lower lips reveals that $nkx3.2^{-/-}$

mutants closely resembles the anaspid- and thelodont-like conditions late in ontogeny (Fig. 4A). The

trait in $nkx3.2^{-/-}$ mutants departed from that in wildtype after the onset of skull ossification and active

feeding (14–21 dpf), and overlaps with the range occupied by anaspids and thelodonts in later stages (1

and 2 mpf). When the jaws were at rest, the gape angle in wildtype zebrafish was consistently around

30 degrees regardless of ontogenetic stages. This was also the case for the $nkx3.2^{-/-}$ mutants at 4 and 14

dpf when they still rely on yolk as main or partial intake. The gape angle increased steadily thereafter

in the mutants.

In landmark-based geometric morphometrics, $nkx3.2^{-/-}$ mutants aligned more closely with

anaspids than wildtype along the principal components (PCs) that explain the adult phenotype (PCs 1

and 5) (Fig. 6E, F). Nearly one third of the overall shape variation loaded on PC1 and primarily

concerned anteroposterior length and dorsoventral height of the entire head. Compared to wildtype

zebrafish, $nkx3.2^{-/-}$ mutants and most anaspids had a much shorter snout (with the lips and the nostrils

shifting posteriorly toward the eye) and dorsoventrally deeper lower lips. PC 5 explained 7.0 % of

overall shape variation, and the traits that vary along it were orientation of lower lip and relative height

of nostrils, which clearly set wildtype zebrafish apart from $nkx3.2^{-/-}$ mutants and anaspids. On the

Cartesian grid of PCs 1 and 5, $nkx3.2^{-/-}$ mutants overlapped with anaspids in morphospace occupation

and apart from wildtype zebrafish (Fig. 6F). The area of overlap also indicated that $nkx3.2^{-/-}$ mutants

are broadly similar to anaspids in these PCs, not just to one or a few individual anaspid taxa. PCs 2-4

(not shown in Fig. 6) largely explain variation among anaspids or within wildtype/mutant zebrafish

samples and were therefore uninformative for comparison between the groups (datasets containing

landmark coordinates and Procrustes transformation are provided in Data Supplement 3).

DISCUSSION

Mutants corroborate the function of *nkx3.2* in joint development

Our *nkx3.2*[-/-] zebrafish reinforce the morphant-based insight that this transcription factor is essential to

the development of jaw joint in non-mammalian vertebrates (Miller et al., 2003). This is likely the

widespread condition among jawed vertebrates. *Nkx3.2* knockdown also results in fusion between the

palatoquadrate and Meckel's cartilage in amphibians (Lukas and Olsson, 2018a), and a similar

phenotype is observed in chicks with ectopic expression of BMP4 and FGF8 that repressed *Nkx3.2*

(Wilson and Tucker, 2004). In mammals, the palatoquadrate (malleus) and Meckel's cartilage (incus)

migrate to form middle ear ossicles. The malleus-incus joint is not affected in *Nkx3.2*[-/-] mice, even

though the malleus becomes narrower in the mutants, and even though *Nkx3.2* plays a role in

specification of the gonium and the anterior tympanic ring (Tucker et al., 2004). Aside from the head,

Nkx3.2 plays a role in various structures in mice and chicks, including the axial column (Herbrand et

al., 2002; Lettice et al., 2001; Murtaugh et al., 2001) and visceral lateralities of the spleen and pancreas

(Hecksher-Sørensen et al., 2004; Schneider et al., 1999). We will test in a forthcoming work whether or

not *nkx3.2*[-/-] zebrafish have parallel phenotypes to these amniote mutants in the axial skeletons or

visceral literalities.

During development of the jaw joint, *nkx3.2* is thought to specify the joint interzone by

inhibiting maturation or hypertrophy of the chondrocytes (Miller et al., 2003; Smeeton et al., 2016).

Similar functions have been ascribed to *irx7* and *irx5a* in the hyoid joint (Askary et al., 2015). This is

consistent with the predicted regulatory function of *Nkx3.2* in vertebral development, intervertebral and

interphalangeal joint formation, or somatic chondrogenesis in general (Herbrand et al., 2002; Lettice et

al., 2001; Murtaugh et al., 2001) — via repression of *Runx2*, by upregulating *Col2α1*, and/or through a

positive feedback loop with *Sox9* (Smeeton et al., 2016). These insights are based on: a) experimental

results using amniote embryos or somitic mesodermal cell cultures (Cairns et al., 2008; Kawato et al.,

2012; Lengner et al., 2005; Murtaugh et al., 2001; Provot et al., 2006; Yamashita et al., 2009; Zeng et

al., 2002); and b) clinical and genetic studies of human pathologies, including osteoarthritis and

spondylo-megaepiphyseal-metaphyseal dysplasia (including pseudoepiphyses) (Caron et al., 2015;

Hellemans et al., 2009). Before this study, however, no mutants were available to specifically address

these potential mechanisms of *nkx3.2* functions in the mandibular arch.

The open-mouth phenotype results from plastic remodeling

The late onset and topology of skull/jaw remodeling suggests that *nkx3.2*[-/-] zebrafish accommodate the

loss of the jaw joint via a plastic response. Other than the absence of jaw joint, *nkx3.2*[-/-] zebrafish

appear normal until metamorphosis (14–21 dpf) (Fig. 1D, G). As the skull ossifies, however, the

observed phenotype becomes increasingly prominent (Fig. 2B, D, G, J, M, N). Although the fused jaw

cartilages clearly result from *nkx3.2* mutation, *nkx3.2* loss-of-function, by itself, seems unlikely to yield

all the rest of phenotypic effects described here. A genome-wide analysis suggests that *nkx3.2*

patterning effects in the skull are restricted to the mid-portion of the mandibular arch (Askary et al.,

2017). Nor does *nkx3.2* have known expression or function in the intramembranously ossified skull

elements (Miller et al., 2003; Tucker et al., 2004), which are dramatically modified in our $nkx3.2^{-/-}$ fish.

The hypertrophied mandibular cartilage may result from *nkx3.2* loss-of-function in regulating local

chondrogenesis (Smeeton et al., 2016). Still, further investigation is warranted because in amphibians

an ectopic cartilage formed in the mandibular arch with *Nkx3.2* overexpression, not repression (Lukas

and Olsson, 2018b).

A comparative survey across vertebrates supports our interpretation of the adult $nkx3.2^{-/-}$

phenotype as skeletal remodeling to accommodate the jaw joint loss. Discrete variation in cichlid jaw

morphology has an epigenetic basis in behaviorally mediated skeletal remodeling, where gaping

frequencies in juveniles correlate with dimensions of the retroarticular process (Hu and Albertson,

2017). The magnitude of morphological changes in the adult $nkx3.2^{-/-}$ phenotype also appears

consistent with a series of surgical experiments in mammalian jaw skeletons (Bayram et al., 2010;

Gomes et al., 2012; Horowitz and Shapiro, 1955; Lifshitz, 1976; Miyazaki et al., 2016; Rodrigues et

al., 2009; Sarnat, 1970; Sarnat and Muchnic, 1971; Toledo et al., 2014) or with the 'bird face'

deformity observed in clinical cases of the temporomandibular joint ankylosis in humans (El-Sheikh et

al., 1996). Collectively, these studies show that latent potentials of development allow a plastic trait to

become expressed in jaw skeletons.

Variation resulting from developmental plasticities — whether induced by environmental cues,

developmental perturbation, or mutation — are often non-random and adaptive (Palmer, 2012; West-

Eberhard, 2005a, 2005b, 2003). Such non-random, adaptive responses are documented across wide

ranges of taxa and structures, including: more robust claws in crabs fed with harder food items (Smith

and Palmer, 1994); longer or shorter appendages of barnacles transplanted between wave-exposed and

protected shores (Kaji and Palmer, 2017; Neufeld and Palmer, 2008); and thickened shells of

gastropods exposed to predator cues (Appleton and Palmer, 1988; Edgell and Neufeld, 2008).

Phylogenetically closer to zebrafish, cichlids have been extensively studied for developmental

plasticity in jaw skeletons, such as: the relationship with gape frequencies and retroarticular processes,

mentioned above (Hu and Albertson, 2017); antisymmetric development of the jaws in scale-eating

specialists (Stewart and Albertson, 2010); and various dietary effects on feeding morphology (Galis,

1993; Greenwood, 1965; Liem and Osse, 1975; Meyer, 1987; Wimberger, 1992, 1991). Ram feeding

facilitated by the fixed open gape in the adult $nkx3.2^{-/-}$ zebrafish corroborates non-random, adaptive

accommodation of the jaw joint defect.

Jaw joint functions constrain skull morphology in vertebrates

The drastically remodeled skull of adult $nkx3.2^{-/-}$ zebrafish highlights jaw movements as an important

factor in the development — and thus morphological diversity — of jawed vertebrate skulls (Depew et

al., 2005; Depew and Compagnucci, 2008; Depew and Simpson, 2006). Functional jaw loss resulting

from $nkx3.2$-null mutations allowed the mutants to depart so markedly in morphology from their

wildtype cousins, despite the nearly identical genetic backgrounds. This departure implies that

movements at the jaw joint limit skull forms. The specific combination of traits observed in adult

$nkx3.2^{-/-}$ zebrafish — e.g., nostrils in interorbital position, premaxilla and maxilla abutted against

antorbital wall, kinethmoid reduced, basihyal protrusion — is likely maladaptive and unavailable to

zebrafish when the jaws function properly. Simultaneously, the absence of a jaw joint (or jaw apparatus

altogether) also limits functionally viable forms. This interpretation is bolstered by the superficial

convergence in head shapes between $nkx3.2^{-/-}$ zebrafish and two Paleozoic agnathan lineages (anaspids

and furcacaudiform thelodonts). The $nkx3.2^{-/-}$ zebrafish and these agnathans share a functional

requirement — the absence of a hinge joint between upper and lower lips — and transversely

compressed body profile. No evidence suggests any more similarities in the otherwise wildtype-like

young mutants (Fig. 1D, G) until the lower jaw skeletons begin rotating posteroventrally post 14 dpf

(Fig. 2B, D). Therefore, we interpret the anaspid/thelodont-like traits in the adult $nkx3.2^{-/-}$ zebrafish not

as recapitulations of conserved, genetically hard-wired early vertebrate development (atavism), but

rather as parallel developmentally plastic responses to shared growth conditions that were experienced

by early vertebrates in a zebrafish skull (convergence).

Thus, jaw-joint loss ($nkx3.2$ loss-of-function phenotype) released $nkx3.2^{-/-}$ zebrafish from

developmental constraints to facilitate jaw movement, and exposed them to a different functional

requirement: feeding without mobile jaws. A fixed open gape — accompanied by modification of the

skull elements — is one morphological solution to the functional loss of the jaw joint, which is

corroborated by ram feeding exhibited by $nkx3.2^{-/-}$ mutants (Fig. 5B; Movie S1). The magnitude of

modification documented post-hatching (Figs. 1–5) is a testament to significant functional optimization

within the bounds of the gnathostome bauplan. Simultaneously, well-constrained occupancy by the

$nkx3.2^{-/-}$ in the PCA plot (Fig. 6F) implies that alternative morphological patterns are either

functionally non-viable (e.g., fixed closed gape) or developmentally non-accessible (e.g., ectopic

formation of a mouth). Thus, the $nkx3.2^{-/-}$ mutants provide a rare case study. The results suggest that

the vertebrate bauplan allows a limited repertoire of functionally viable morphological patterns, onto

which forms may converge under a given functional constraint.

In support of our interpretation, remarkable convergences arise via developmental plasticity

under similar functional requirements — even in the absence of homology in individual skeletal

elements — between taxa widely separated chronologically and phylogenetically. Bichirs (*Polypterus*)

routinely trained under terrestrial conditions modify the pectoral fin skeleton through plasticity and

develop morphological conditions observed in stem tetrapods (Standen et al., 2014). In another, *vav2*

and *waslb* mutant zebrafish develop limb-like long bones connected by joints within the pectoral fins

(Hawkins et al., 2018, preprint). West-Eberhard (2005b) reviewed a goat born with congenital paralysis

of forelimbs, which performed bipedal locomotion. This behavioral accommodation led to highly

modified musculoskeletal anatomy of the axial column, pelvis, and hindlimbs (Slijper, 1942a, 1942b).

As shown in *nkx3.2$^{-/-}$* zebrafish, these experimental manipulations and clinical reports illustrate that

adaptive forms emerge from developmentally plastic responses to functional constraints — and

therefore may converge onto phenotype independently exploited by a distant lineage — even after

development has laid out lineage-specific patterns (such as homologies of individual bones).

***nkx3.2* mutants as a unique model for skeletal development and disease**

Mechanisms of developmentally plastic remodeling, whether at the level of whole-animal behavior or

gene transcription, remain an elusive component of skeletal development that is challenging to test

experimentally. Therefore, *nkx3.2$^{-/-}$* zebrafish provide a unique experimental system in which to test

further the role of a joint in skeletal development. For example: a) What genetic mechanisms regulate

the remodeling of intramembranous ossifications? b) What components of the remodeling process

respond to jaw-joint dysfunction? and c) Does any feedback exists to coordinate remodeling between

developmentally independent, but functionally connected units (e.g., premaxilla-maxilla complex and

kinethmoid)? Such insights would begin to fill in knowledge gaps regarding skeletal and joint diseases

including osteoarthritis and joint ankylosis.

The non-atavistic nature of the *nkx3.2$^{-/-}$* phenotype is potentially useful to reevaluate

evolutionarily inspired interpretations of phenocopies in general. There are no evolutionary

relationships in the similarities between *nkx3.2$^{-/-}$* zebrafish and anaspids or thelodonts. Anaspids are a

stem cyclostome lineage (Miyashita et al., 2019) (Fig. 6G), and thus a poor surrogate for an ancestral

state. The lineages between anaspids and thelodonts, or those between thelodonts and the gnathostome

crown, do not share a similar combination of morphological traits, but instead are dorsoventrally

depressed forms (Janvier, 1996; Miyashita, 2016; Miyashita et al., 2019) (Fig. 6G). Finally, head

similarities between the mutants and the stem taxa do not extend beyond overall configuration. No

skull elements are lost or replaced in *nkx3.2*$^{-/-}$ zebrafish to achieve the anaspid/thelodont-like

micromery (Blom et al., 2001; Janvier, 1996; Märss et al., 2007). Therefore, the adult *nkx3.2*$^{-/-}$

phenotype clearly does not represent a reversal to an ancestral state. Although experimental

phenocopies are often interpreted as atavistic (ancestral reversal), alternative interpretations are seldom

tested (Smith and Schneider, 1998). Here, our *nkx3.2*$^{-/-}$ zebrafish present developmental plasticity as a

testable alternative to atavistic reversal to explain a phenocopy. By showing that a gnathostome can

survive without a jaw, the anaspid/thelodont-like *nkx3.2*$^{-/-}$ zebrafish also offer a comparative model to

make inferences about the functional morphology of these long-extinct agnathans.

Acknowledgements

We thank B.F. Eames (University of Saskatchewan) for facilitating our collaboration; M.E. Bronner

(California Institute of Technology), S.J. Childs (University of Calgary), M.I. Coates, V.E. Prince, and

M.W. Westneat (University of Chicago) for discussion; and members of the Allison and Prince labs for

maintenance of *nkx3.2*^{ua5011}.

Competing interests

The authors declare no competing or financial interests.

Author contributions

T.M. conceived project; T.M. and A.P.O. performed experiments resulting in *nkx3.2*^{ua5011}; J.S. and

N.N. performed experiments resulting in *nkx3.2*^{el802}; P.B. and B.G. provided μCT scanning and

histological sampling; T.M. and P.B. conducted morphometric analyses; T.M., P.B., A.R.P., J.G.C.,

D.G., and W.T.A. analyzed data; T.M. wrote the paper with inputs from P.B., A.P.O., J.S., A.R.P.,

J.G.C., D.G., and W.T.A.

Funding. This work was supported by Natural Science and Engineering Research Council grants

RGPIN-2014-04863, RGPIN-2014-06311, and RGPIN-2015-06006 (to A.R.P., D.G., and W.T.A.,

respectively) and National Institute of Health grants R35DE027550 and K99 DE027218 (to J.G.C. and

J.S., respectively).

Supplementary information

Sequence information, measurements, and landmark coordinates are available as data supplements to

this paper.

Movie S1. Feeding behaviour and 3D μCT scans of jawless zebrafish. Phenotypic comparison of

nkx3.2$^{-/-}$ and wildtype zebrafish. Three-dimensional rendering of μCT scan of the age-matched *nkx3.2*$^{-/-}$

(ua5011) and wildtype (AB strain) zebrafish at 2 mpf, each followed by filming of feeding behavior on

brine shrimp played at 1/10 original speed.

Data supplements

Data supplement 1. Sequence information for the *nkx3.2* allele ua5011.

Data supplement 2. Gape angles and other linear measurements in zebrafish; gape angles and orbit

diameter in anaspids.

Data supplement 3. Thin-plate-spline landmark data in zebrafish (wildtype and *nkx3.2*$^{-/-}$) and anaspids

used for Procrustes transformation, which were subjected to principal component analysis.

REFERENCES

Adekeye EO. 1983. Ankylosis of the mandible: Analysis of 76 cases. *J. Oral Maxillofac. Surg.* **41**:442–

449. doi:10.1016/0278-2391(83)90129-5

Alexander RM. 1970. Mechanics of the feeding action of various teleost fishes. *J. Zool.* **162**:145–156.

doi:10.1111/j.1469-7998.1970.tb01261.x

Alexander RMcN. 1969. Mechanics of the feeding action of a cyprinid fish. *J. Zool.* **159**:1–15.

doi:10.1111/j.1469-7998.1969.tb03067.x

Appleton RD, Palmer AR. 1988. Water-borne stimuli released by predatory crabs and damaged prey

induce more predator-resistant shells in a marine gastropod. *PNAS* **85**:4387–4391.

doi:10.1073/pnas.85.12.4387

Askary A, Mork L, Paul S, He X, Izuhara AK, Gopalakrishnan S, Ichida JK, McMahon AP,

Dabizljevic S, Dale R, Mariani FV, Crump JG. 2015. Iroquois proteins promote skeletal joint

formation by maintaining chondrocytes in an immature state. *Dev. Cell* **35**:358–365.

doi:10.1016/j.devcel.2015.10.004

Askary A, Xu P, Barske L, Bay M, Bump P, Balczerski B, Bonaguidi MA, Crump JG. 2017. Genome-

wide analysis of facial skeletal regionalization in zebrafish. *Development* **144**:2994–3005.

doi:10.1242/dev.151712

Barske L, Askary A, Zuniga E, Balczerski B, Bump P, Nichols JT, Crump JG. 2016. Competition

between Jagged-Notch and Endothelin1 signaling selectively restricts cartilage formation in the

zebrafish upper face. *PLoS Genet.* **12**:e1005967. doi:10.1371/journal.pgen.1005967

Bayram B, Uckan S, Cetinsahin A, Arman Ozcirpici A, Ozdemir H, Yazici C. 2010. Repositioning of

the masseter muscle and its effect on skeletal growth. *Oral Surg. Med. Pathol. Radiol.*

Endodontol. **109**:e1–e5. doi:10.1016/j.tripleo.2009.12.041

Bixler D, Ward R, Gale DD. 1985. Agnathia-holoprosencephaly: A developmental field complex

involving face and brain. Report of 3 cases. *J. Craniofac. Genet. Dev. Biol. Suppl* **1**:241–249.

Blom H, Märss T. 2010. The interrelationships and evolutionary history of anaspid fishes In: Elliott

DK, Maisey JG, Yu X, Miao D, editors. Morphology, Phylogeny and Palaeogeography of

Fossil Fishes – Honoring Meemann Chang. Munich: Verlag Dr. Fredrich Pfeil.

Blom H, Märss T, Miller CG. 2001. Silurian and earliest Devonian birkeniid anaspids from the

Northern Hemisphere. *Eart. Envir. Sci. Trans. R. Soc. Edinbur.* **92**:263–323.

doi:10.1017/S0263593300000250

Brown DM, Marsh JL. 1990. Agnathia and associated malformations: A case report. *Clef. Palat. J.*

27:415–418. doi:10.1597/1545-1569_1990_027_0415_aaamac_2.3.co_2

Cairns DM, Sato ME, Lee PG, Lassar AB, Zeng L. 2008. A gradient of Shh establishes mutually
repressing somitic cell fates induced by Nkx3.2 and Pax3. *Dev. Biol.* **323**:152–165.
doi:10.1016/j.ydbio.2008.08.024

Caron MMJ, Emans PJ, Surtel DAM, van der Kraan PM, van Rhijn LW, Welting TJM. 2015. BAPX-
1/NKX-3.2 acts as a chondrocyte hypertrophy molecular switch in osteoarthritis. *Arthrit.*
Rheumatol. **67**:2944–2956. doi:10.1002/art.39293

Cerny R, Cattell M, Sauka-Spengler T, Bronner-Fraser M, Yu F, Medeiros DM. 2010. Evidence for the
prepattern/cooption model of vertebrate jaw evolution. *Proc. Nat. Acad. Sci.* **107**:17262–17267.
doi:10.1073/pnas.1009304107

Chidzonga MM. 1999. Temporomandibular joint ankylosis: Review of thirty-two cases. *Brit. J. Oral*
Maxillofac. Surg. **37**:123–126. doi:10.1054/bjom.1997.0089

Cubbage CC, Mabee PM. 1996. Development of the cranium and paired fins in the zebrafish *Danio*
rerio (Ostariophysi, Cyprinidae). *J. Morphol.* **229**:121–160. doi:10.1002/(SICI)1097-
4687(199608)229:2<121::AID-JMOR1>3.0.CO;2-4

Depew MJ, Compagnucci C. 2008. Tweaking the hinge and caps: Testing a model of the organization
of jaws. *J. Exp. Zool.* **310B**:315–335. doi:10.1002/jez.b.21205

Depew MJ, Simpson CA. 2006. 21st Century neontology and the comparative development of the
vertebrate skull. *Dev. Dyn.* **235**:1256–1291. doi:10.1002/dvdy.20796

Depew MJ, Simpson CA, Morasso M, Rubenstein JLR. 2005. Reassessing the Dlx code: The genetic
regulation of branchial arch skeletal pattern and development. *J. Anat.* **207**:501–561.
doi:10.1111/j.1469-7580.2005.00487.x

Donoghue PCJ, Forey PL, Aldridge RJ. 2000. Conodont affinity and chordate phylogeny. *Biol. Rev.*
75:191–251.

Edgell TC, Neufeld CJ. 2008. Experimental evidence for latent developmental plasticity: Intertidal
whelks respond to a native but not an introduced predator. *Biol. Let.* **4**:385–387.
doi:10.1098/rsbl.2008.0204

El-Sheikh MM, Medra AM, Warda MH. 1996. Bird face deformity secondary to bilateral
temporomandibular joint ankylosis. *J. Cranio-Maxillofac. Surg.* **24**:96–103.
doi:10.1016/S1010-5182(96)80020-5

Gagnon JA, Valen E, Thyme SB, Huang P, Ahkmetova L, Pauli A, Montague TG, Zimmerman S,
Richter C, Schier AF. 2014. Efficient mutagenesis by Cas9 protein-mediated oligonucleotide
insertion and large-scale assessment of single-guide RNAs. *PLoS One* **9**:e98186.
doi:10.1371/journal.pone.0098186

Galis F. 1993. Interactions between the pharyngeal jaw apparatus, feeding behaviour, and ontogeny in
the cichlid fish, Haplochromis piceatus: A study of morphological constraints in evolutionary
ecology. *J. Exp. Zool.* **267**:137–154. doi:10.1002/jez.1402670207

Gekas J, Li B, Kamnasaran D. 2010. Current perspectives on the etiology of agnathia-otocephaly.
European J. Med. Genet. **53**:358–366. doi:10.1016/j.ejmg.2010.09.002

Gillis JA, Modrell MS, Baker CVH. 2013. Developmental evidence for serial homology of the
vertebrate jaw and gill arch skeleton. *Nat. Commun.* **4**:1436. doi:10.1038/ncomms2429

Gomes FEF, Moraes RB, Luz JG de C. 2012. Effects of temporal muscle detachment and
coronoidotomy on facial growth in young rats. *Braz. Oral Res.* **26**:348–354.
doi:10.1590/S1806-83242012000400011

Greenwood PH. 1965. Environmental effects on the phar-yngeal mill of a cichlid fish,Astatoreochromis
alluaudi,and their taxonomic implications. *Proc. Linn. Soc. Lond.* **176**:1–10.

Hall BK. 2015. Bones and Cartilage: Developmental and Evolutionary Skeletal Biology. 2nd edition.
London: Academic Press.

Hawkins MB, Henke K, Harris MP. 2018. Latent developmental potential to form limb-like skeletal
structures in zebrafish. *bioRxiv* 450619. doi:10.1101/450619

Hecksher-Sørensen J, Watson RP, Lettice LA, Serup P, Eley L, Angelis CD, Ahlgren U, Hill RE. 2004.
The splanchnic mesodermal plate directs spleen and pancreatic laterality, and is regulated by
Bapx1/Nkx3.2. *Development* **131**:4665–4675. doi:10.1242/dev.01364

Hellemans J, Simon M, Dheedene A, Alanay Y, Mihci E, Rifai L, Sefiani A, van Bever Y, Meradji M,
Superti-Furga A, Mortier G. 2009. Homozygous inactivating mutations in the NKX3-2 gene
result in spondylo-megaepiphyseal-metaphyseal dysplasia. *Am. J. Hum. Genet.* **85**:916–922.
doi:10.1016/j.ajhg.2009.11.005

Herbrand H, Pabst O, Hill R, Arnold H-H. 2002. Transcription factors Nkx3.1 and Nkx3.2 (Bapx1)
play an overlapping role in sclerotomal development of the mouse. *Mec. Dev.* **117**:217–224.
doi:10.1016/S0925-4773(02)00207-1

Hernandez LP. 2000. Intraspecific scaling of feeding mechanics in an ontogenetic series of zebrafish,
Danio rerio. *J. Exp. Biol.* **203**:3033–3043.

Hernandez LP, Bird NC, Staab KL. 2007. Using zebrafish to investigate cypriniform evolutionary
novelties: Functional development and evolutionary diversification of the kinethmoid. *J. Exp.
Zool.* **308B**:625–641. doi:10.1002/jez.b.21166

Horowitz SL, Shapiro HH. 1955. Modification of skull and jaw architecture following removal of the
masseter muscle in the rat. *Am. J. Phys. Anthropol.* **13**:301–308. doi:10.1002/ajpa.1330130208

Howland HC, Merola S, Basarab JR. 2004. The allometry and scaling of the size of vertebrate eyes.
Vis. Res. **44**:2043–2065. doi:10.1016/j.visres.2004.03.023

Hu Y, Albertson RC. 2017. Baby fish working out: An epigenetic source of adaptive variation in the
cichlid jaw. *Proc R Soc B* **284**:20171018. doi:10.1098/rspb.2017.1018

Janvier P. 2007. Homologies and evolutionary transitions in early vertebrate history In: Anderson JS,
Sues H-D, editors. Major Transitions in Vertebrate Evolution. Bloomington: Indiana University
Press. pp. 57–121.

Janvier P. 1996. Early Vertebrates, Oxford Monographs on Geology and Geophysics. Oxford:
Clarendon Press.

Kaji T, Palmer AR. 2017. How reversible is development? Contrast between developmentally plastic
gain and loss of segments in barnacle feeding legs. *Evolution* **71**:756–765.
doi:10.1111/evo.13152

Kawato Y, Hirao M, Ebina K, Shi K, Hashimoto J, Honjo Y, Yoshikawa H, Myoui A. 2012. Nkx3.2
promotes primary chondrogenic differentiation by upregulating Col2a1 transcription. *PLoS One*
7:e34703. doi:10.1371/journal.pone.0034703

Keating JN, Donoghue PCJ. 2016. Histology and affinity of anaspids, and the early evolution of the
vertebrate dermal skeleton. *Proc. R. Soc. B* **283**:20152917. doi:10.1098/rspb.2015.2917

Kiaer J. 1924. The Downtonian fauna of Norway. I. Anaspida with a geological introduction. *Skrifter*
6:1–139.

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of embryonic
development of the zebrafish. *Dev. Dyn.* **203**:253–310. doi:10.1002/aja.1002030302

Klingenberg CP. 2011. MorphoJ: An integrated software package for geometric morphometrics. *Mol.*
Ecol. Res. **11**:353–357. doi:10.1111/j.1755-0998.2010.02924.x

Kuratani S. 2012. Evolution of the vertebrate jaw from developmental perspectives. *Evol. Dev.* **14**:76–
92. doi:10.1111/j.1525-142X.2011.00523.x

Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. 2016. CHOPCHOP v2: A web tool for the
next generation of CRISPR genome engineering. *Nuc. Acids Res* **44**:W272–W276.
doi:10.1093/nar/gkw398

Lauder GV. 1980. Evolution of the feeding mechanism in primitive actionopterygian fishes: A
functional anatomical analysis of *Polypterus*, *Lepisosteus*, and *Amia*. *J. Morphol.* **163**:283–317.
doi:10.1002/jmor.1051630305

Lauder GV. 1979. Feeding mechanics in primitive teleosts and in the halecomorph fish *Amia calva*. *J.*
Zool. **187**:543–578. doi:10.1111/j.1469-7998.1979.tb03386.x

Lengner CJ, Hassan MQ, Serra RW, Lepper C, Wijnen AJ van, Stein JL, Lian JB, Stein GS. 2005.
Nkx3.2-mediated repression of Runx2 promotes chondrogenic differentiation. *J. Biol. Chem.*
280:15872–15879. doi:10.1074/jbc.M411144200

Lettice L, Hecksher-Sørensen J, Hill R. 2001. The role of Bapx1 (Nkx3.2) in the development and
evolution of the axial skeleton. *J. Anat.* **199**:181–187. doi:10.1046/j.1469-
7580.2001.19910181.x

Liem KF, Osse JWM. 1975. Biological versatility, evolution, and food resource exploitation in African
cichlid fishes. *Integr. Comp. Biol.* **15**:427–454. doi:10.1093/icb/15.2.427

Lifshitz J. 1976. Comparative anatomic study of mandibular growth in rats after bilateral resections of
superficial masseter, posterior temporal, and anterior digastric muscles. *J. Dent. Res.* **55**:854–
858. doi:10.1177/00220345760550052301

Lukas P, Olsson L. 2018a. Bapx1 is required for jaw joint development in amphibians. *Evol. Dev.*
20:192–206. doi:10.1111/ede.12267

Lukas P, Olsson L. 2018b. Bapx1 upregulation is associated with ectopic mandibular cartilage
development in amphibians. *Zool. Let.* **4**:16. doi:10.1186/s40851-018-0101-3

Manganello-Souza LC, Mariani PB. 2003. Temporomandibular joint ankylosis: Report of 14 cases.
International J. Oral Maxillofac. Surg. **32**:24–29. doi:10.1054/ijom.2002.0308

Märss T, Turner S, Karatajūtė-Talimaa V. 2007. Handbook of Paleoichthyology / "Agnatha" II.
Thelodonti. München: Verlag Dr. Friedrich Pfeil.

Medeiros DM, Crump JG. 2012. New perspectives on pharyngeal dorsoventral patterning in
development and evolution of the vertebrate jaw. *Dev. Biol.* **371**:121–135.
doi:10.1016/j.ydbio.2012.08.026

Meyer A. 1987. Phenotypic plasticity and heterochrony in *Cichlasoma managuense* (Pisces, Cichlidae)
and their implications for speciation in cichlid fishes. *Evolution* **41**:1357–1369.
doi:10.1111/j.1558-5646.1987.tb02473.x

Miller CT, Yelon D, Stainier DYR, Kimmel CB. 2003. Two endothelin 1 effectors, hand2 and bapx1,
pattern ventral pharyngeal cartilage and the jaw joint. *Development* **130**:1353–1365.
doi:10.1242/dev.00339

Miyashita T. 2016. Fishing for jaws in early vertebrate evolution: A novel hypothesis of mandibular
confinement. *Biol. Rev.* **91**:611–657. doi:10.1111/brv.12187

Miyashita T. 2012. Comparative Analysis of the Anatomy of the Myxinoidea and the Ancestry of Early
Vertebrate Lineages (Unpublished M.Sc. thesis). Edmonton: University of Alberta.

Miyashita T, Coates MI, Farrar R, Larson P, Manning PL, Wogelius RA, Edwards NP, Anné J,
Bergmann U, Palmer AR, Currie PJ. 2019. Hagfish from the Cretaceous Tethys Sea and a
reconciliation of the morphological–molecular conflict in early vertebrate phylogeny. *Proc.*
Nat. Acad. Sci. **116**:2146–2151. doi:10.1073/pnas.1814794116

Miyazaki M, Yonemitsu I, Takei M, Kure-Hattori I, Ono T. 2016. The imbalance of masticatory
muscle activity affects the asymmetric growth of condylar cartilage and subchondral bone in
rats. *Arc. Oral Biol.* **63**:22–31. doi:10.1016/j.archoralbio.2015.11.020

Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. 2014. CHOPCHOP: A CRISPR/Cas9 and
TALEN web tool for genome editing. *Nuc. Acids Res.* **42**:W401–W407.
doi:10.1093/nar/gku410

Murtaugh LC, Zeng L, Chyung JH, Lassar AB. 2001. The chick transcriptional repressor Nkx3.2 acts
downstream of Shh to promote BMP-dependent axial chondrogenesis. *Dev. Cell* **1**:411–422.
doi:10.1016/S1534-5807(01)00039-9

Neufeld CJ, Palmer AR. 2008. Precisely proportioned: Intertidal barnacles alter penis form to suit
coastal wave action. *Proc. R. Soc. B* **275**:1081–1087. doi:10.1098/rspb.2007.1760

Palmer AR. 2012. Developmental plasticity and the origin of novel forms: Unveiling cryptic genetic
variation via "use and disuse." *J. Exp. Zool. Part* **318B**:466–479. doi:10.1002/jez.b.21447

Provot S, Kempf H, Murtaugh LC, Chung U, Kim D-W, Chyung J, Kronenberg HM, Lassar AB. 2006.
Nkx3.2/Bapx1 acts as a negative regulator of chondrocyte maturation. *Development* **133**:651–
662. doi:10.1242/dev.02258

Rodrigues L, Traina AA, Nakamai LF, Luz JG de C. 2009. Effects of the unilateral removal and
dissection of the masseter muscle on the facial growth of young rats. *Braz. Oral Res.* **23**:89–95.
doi:10.1590/S1806-83242009000100015

Rohlf FJ. 2018. tpsDIG. New York.

Rohlf FJ. 1999. Shape statistics: Procrustes superimpositions and tangent spaces. *J. Classif.* **16**:197–
223. doi:10.1007/s003579900054

Sansom RS, Freedman K, Gabbott SE, Aldridge RJ, Purnell MA. 2010. Taphonomy and affinity of an
enigmatic Silurian vertebrate, *Jamoytius kerwoodi* White. *Palaeontology* **53**:1393–1409.
doi:10.1111/j.1475-4983.2010.01019.x

Sarnat BG. 1970. The face and jaws after surgical experimentation with the septovomeral region in
growing and adult rabbits. *Acta Oto-Laryngol.* **69**:1–30. doi:10.3109/00016487009131762

Sarnat BG, Muchnic H. 1971. Facial skeletal changes after mandibular condylectomy in the adult
monkey. *J. Anat.* **108**:323–338.

Schiffer C, Tariverdian G, Schiesser M, Thomas MC, Sergi C. 2002. Agnathia-otocephaly complex:
Report of three cases with involvement of two different Carnegie stages. *Am. J. Med. Genet.*
112:203–208. doi:10.1002/ajmg.10672

Schilling TF, Kimmel CB. 1997. Musculoskeletal patterning in the pharyngeal segments of the
zebrafish embryo. *Development* **124**:2945–2960.

Schneider A, Mijalski T, Schlange T, Dai W, Overbeek P, Arnold H-H, Brand T. 1999. The homeobox
gene it NKX3.2 is a target of left–right signalling and is expressed on opposite sides in chick
and mouse embryos. *Curr. Biol.* **9**:911-S1. doi:10.1016/S0960-9822(99)80397-2

Slijper EJ. 1942a. Biologic-anatomical investigations on the bipedal gait and upright posture in
mammals, with special reference to a little goat, born without forelegs. I. *Proc. Konink. Ned.*
Akad. Wet. **45**:288–295.

Slijper EJ. 1942b. Biologic-anatomical investigations on the bipedal gait and upright posture in
mammals, with special reference to a little goat, born without forelegs. II. *Proc. Konink. Ned.*
Akad. Wet. **45**:407–415.

Smeeton J, Askary A, Gage Crump J. 2016. Building and maintaining joints by exquisite local control
of cell fate. *WIREs Dev. Biol.* **6**:e245. doi:10.1002/wdev.245

Smith KK, Schneider RA. 1998. Have gene knockouts caused evolutionary reversals in the mammalian
first arch? *BioEssays* **20**:245–255. doi:10.1002/(SICI)1521-1878(199803)20:3<245::AID-
BIES8>3.0.CO;2-Q

Smith LD, Palmer AR. 1994. Effects of manipulated diet on size and performance of brachyuran crab
claws. *Science* **264**:710–712. doi:10.1126/science.264.5159.710

Standen EM, Du TY, Larsson HCE. 2014. Developmental plasticity and the origin of tetrapods. *Nature*
513:54–58. doi:10.1038/nature13708

Stewart TA, Albertson RC. 2010. Evolution of a unique predatory feeding apparatus: Functional
anatomy, development and a genetic locus for jaw laterality in Lake Tanganyika scale-eating
cichlids. *BMC Biol.* **8**:8. doi:10.1186/1741-7007-8-8

Toledo LG, Cavalcanti SCXB, Corrêa L, Luz JGC. 2014. Effects of injury or removal of the articular
disc on maxillomandibular growth in young rats. *J. Oral Maxillofac. Surg.* **72**:2140–2147.
doi:10.1016/j.joms.2014.06.445

Tucker AS, Watson RP, Lettice LA, Yamada G, Hill RE. 2004. Bapx1 regulates patterning in the
middle ear: Altered regulatory role in the transition from the proximal jaw during vertebrate
evolution. *Development* **131**:1235–1245. doi:10.1242/dev.01017

West-Eberhard MJ. 2005a. Developmental plasticity and the origin of species differences. *Proc. Nat.*
Acad. Sci. **102**:6543–6549. doi:10.1073/pnas.0501844102

West-Eberhard MJ. 2005b. Phenotypic accommodation: Adaptive innovation due to developmental
plasticity. *J. Exp. Zool.* **304B**:610–618. doi:10.1002/jez.b.21071

West-Eberhard MJ. 2003. Developmental Plasticity and Evolution, 1 edition. ed. Oxford ; New York:
Oxford University Press.

Westneat MW. 2004. Evolution of levers and linkages in the feeding mechanisms of fishes. *Integr*
Comp Biol **44**:378–389. doi:10.1093/icb/44.5.378

Westneat MW. 2005. Skull biomechanics and suction feeding in fishes. Fish Physiology, Fish
Biomechanics. Academic Press. pp. 29–75. doi:10.1016/S1546-5098(05)23002-9

Wilson J, Tucker AS. 2004. Fgf and Bmp signals repress the expression of Bapx1 in the mandibular
mesenchyme and control the position of the developing jaw joint. *Dev. Biol.* **266**:138–150.
doi:10.1016/j.ydbio.2003.10.012

Wilson MVH, Caldwell MW. 1998. The Furcacaudiformes: A new order of jawless vertebrates with
thelodont scales, based on articulated Silurian and Devonian fossils from northern Canada. *J.*
Vert. Paleontol. **18**:10–29. doi:10.1080/02724634.1998.10011031

Wilson MVH, Caldwell MW. 1993. New Silurian and Devonian fork-tailed "thelodonts" are jawless
vertebrates with stomachs and deep bodies. *Nature* **361**:442–444. doi:10.1038/361442a0

Wimberger PH. 1992. Plasticity of fish body shape. The effects of diet, development, family and age in
two species of *Geophagus* (Pisces: Cichlidae). *Biol. J. Linn. Soc.* **45**:197–218.
doi:10.1111/j.1095-8312.1992.tb00640.x

Wimberger PH. 1991. Plasticity of jaw and skull morphology in the Neotropical cichlids *Geophagus*
Brasiliensis and *G. Steindachneri*. *Evolution* **45**:1545–1563. doi:10.1111/j.1558-
5646.1991.tb02662.x

Yamashita S, Andoh M, Ueno-Kudoh H, Sato T, Miyaki S, Asahara H. 2009. Sox9 directly promotes
Bapx1 gene expression to repress Runx2 in chondrocytes. *Exp. Cell Res.* **315**:2231–2240.
doi:10.1016/j.yexcr.2009.03.008

Zeng L, Kempf H, Murtaugh LC, Sato ME, Lassar AB. 2002. Shh establishes an Nkx3.2/Sox9
autoregulatory loop that is maintained by BMP signals to induce somitic chondrogenesis. *Genes*
Dev. **16**:1990–2005. doi:10.1101/gad.1008002

FIGURES

Fig. 1. Null alleles of *nkx3.2* in zebrafish result in the jaw joint ankylosis.

(**A**) Mutations engineered with CRISPR/Cas9 or TALEN in the zebrafish gene *nkx3.2* that

encodes the homeobox transcription factor protein Nkx3.2 (a.k.a. Bapx1). <u>Top:</u> The gene *nkx3.2*

contains two exons (purple), the second of which encodes the homeobox domain (DNA binding

domain, blue). The diagram is approximately to scale (exon 1 = 310 bp), except the 3' untranslated

region (3'UTR). Allele *nkx3.2^{el802}* was generated with TALEN technology that deleted 20 bp from the

start of the gene (orange dashed line), eliminating the start codon. Allele *nkx3.2^{ua5011}* was engineered

with CRISPR/Cas9 to generate a 20 bp deletion (red box) and frameshift that is predicted to eliminate

the homeodomain. <u>Bottom:</u> Schematic of predicted protein following CRISPR mutagenesis. In the

allele *nkx3.2^{ua5011}*, the frameshift (20 bp deletion) is predicted to disrupt the translation and abrogate

production of the critical homeobox domain. This is predicted to produce random amino acids (red

hatching). (**A'**) Sequencing results from allele *nkx3.2^{ua5011}* (lower) compared to a wildtype zebrafish

(upper).

At 4 dpf, the chondrocrania are compared by *sox10*:eGFP expression in chondrocytes (**B–D**, **H**,

I) or by alcian blue staining of cartilages (**E–G**) between age-matched specimens. (**B**, **E**) The

chondrocranial morphology of wildtypes (AB background; *sox10*:eGFP) at 4 dpf in left lateral view, in

sox10:eGFP expression within chondrocytes (B) and alcian blue staining of cartilages (E). (**C**, **F**) The

chondrocranial morphology of *nkx3.2* heterozygous mutants (*nkx3.2^{+/ua5011}*; *sox10*:eGFP) at 4 dpf in

left lateral view, in *sox10*:eGFP expression within chondrocytes (C) and alcian blue staining of

cartilages (F). (**D**, **G**) The chondrocranial morphology of *nkx3.2* homozygous mutants (*nkx3.2^ua5011^*;

sox10:eGFP) at 4 dpf in left lateral view, in *sox10*:eGFP expression within chondrocytes (D) and alcian

blue staining of cartilages (G). (**H**, **I**) Comparison of jaw morphology between age-matched wildtype

(AB background; *sox10*:eGFP) (H) and *nkx3.2* homozygous mutants (*nkx3.2^ua5011^*; *sox10*:eGFP) (I) at 4

dpf in left lateral view, using *sox10*:eGFP expression within chondrocytes. White broken lines

delineate the mandibular cartilages. (**J**) Schematic drawing of the mandibular cartilages in zebrafish at

4 dpf in left lateral view, showing wildtype morphology (see panel E).

Abbreviations: asterisk (*), jaw joint; **aa**, anguloarticular; **ar**, articular; **bh**, basihyal; **ch**,

ceratohyal; **d**, dentary; **dpf**, days post fertilization; **hm**, hyomandibula; **ih**, interhyal; **j**, jugal; **ke**,

kinethmoid; **lp**, lower lip; **mc**, Meckel's cartilage; **m**, maxilla; **n**, nostril or aperture of nasohypophyseal

system; **o**, operculum; **pm**, premaxilla; **po**, preopercular; **pq**, palatoquadrate; **q**, quadrate; **ra**,

retroarticular process; **sm**, symplectic; **up**, upper lip.

Fig. 2. Ontogeny of *nkx3.2⁻/⁻* zebrafish documents drastic remodeling of the skull.

(**A**, **B**) Comparison of the skull morphology between age-matched wildtype (AB strain) (A) and *nkx3.2* homozygous mutant (*nkx3.2ua5011*) (B) at 21 dpf in right lateral view (inverted for consistency with other panels), using alcian blue and alizarin red staining. (**C**, **D**) Comparison of skull morphology between age-matched wildtype (AB) (C) and *nkx3.2* homozygous mutant (*nkx3.2ua5011*) (D) at 1mpf in left lateral view, using alcian blue and alizarin red staining. (**E**) Detailed morphology of jaw skeleton in *nkx3.2* homozygous mutant (*nkx3.2ua5011*) at 1 mpf in right lateral view (inverted for consistency with other panels), using alcian blue and alizarin red staining.

(**F**, **G**) Comparison of skull morphology between age-matched wildtype (AB) (F) and *nkx3.2* homozygous mutant (*nkx3.2ua5011*) (G) at 2 mpf in left lateral view, using alcian blue and alizarin red staining. (**H**) Interpretive drawing of the specimen in panel g.

(**I**, **J**) Comparison of skull morphology between age-matched wildtype (AB) (I) and *nkx3.2* homozygous mutant (*nkx3.2el802*) (J) at 2 mpf in left lateral view, using alcian blue and alizarin red staining. This second independent null allele of *nkx3.2* confirms the phenotypes reported herein.

884   (**K–N**) Comparison of overall morphology at 2 mpf among age-matched wildtype (AB)

885   female (K) and male (L) and *nkx3.2* homozygous mutant (*nkx3.2^{el802}*) female (M) and male (N) in left

886   lateral view. Scale bars = 5 mm. Abbreviations follow Fig. 1.

Fig. 3. Detailed morphology of jaw joint in wildtype, or the joint-less interface between the upper and lower jaws in *nkx3.2*$^{-/-}$ mutants.

(**A**, **B**) Detailed morphology of junction between upper and lower jaws in age-matched wildtype (A) and *nkx3.2* homozygous mutant (*nkx3.2*^{ua5011}) (B) at 21 dpf in left lateral view, using alcian blue and alizarin red staining. (**C**, **D**) Sagittal section of junction between upper and lower jaws in age-matched wildtype (AB) (C) and *nkx3.2* homozygous mutant (*nkx3.2*^{ua5011}) (D) at 1 mpf, stained with eosin and hematoxylin. (**E**) Detailed view of section in panel D.

A. Ontogeny of gape angles in wildtype and mutant zebrafish

B. Overall skull growth in wildtype and mutant zebrafish

C. Ontogeny of lower jaws in wildtype and mutant zebrafish

Fig. 4. Growth of *nkx3.2*$^{-/-}$ mutants in linear and proportional traits.

(**A**) A marked departure from normal morphology (wildtype = blue) occurs in lower jaw

orientations of *nkx3.2*$^{-/-}$ zebrafish (= red) past 14 dpf, coinciding in timing with metamorphosis (onset

of intramembranous ossification in the skulls and active feeding). Fixed open gapes in the mutants at 1

to 2 mpf are comparable to those of Paleozoic agnathan lineages, birkeniiform anaspids and

furcacaudiid thelodonts (green). The orientations were measured here as gape, an angle between upper

and lower lips at natural, resting position.

(**B**) Wildtype and *nkx3.2*$^{-/-}$ zebrafish do not differ significantly from each other in absolute

sizes, except at 4 dpf. Here, skull length is selected to illustrate this general observation. (**C**)

Proportional changes appear to follow shape changes in skeletal remodeling. The box plot shows

phenotypic separation in relative lower jaw length between wildtype and *nkx3.2*$^{-/-}$ zebrafish at 1 and 2

mpf, even though a significant difference developed in lower jaw orientation by 21 dpf.

Values are plotted as boxes of first and third quartile, with middle line displaying mean, and

whiskers communicating maximum and minimum values (n = sample size, same across A–C). Asterisk

indicates level of statistically significant difference in means (*t*-test): *, $P < 0.05$; **, $P < 0.01$.

Photographs of zebrafish are male representative specimens at 60 dpf (Fig. 2L, N). Drawings are:

Pharyngolepis oblonga as a general representative of anaspids (after Blom et al., 2001; Kiaer, 1924);

and *Furcacauda fredholmae* as a general representative of furcacaudiid thelodonts (after Wilson and

Caldwell, 1993). See Data Supplement 2 for all original measurements.

Fig. 5. Functionally jawless *nkx3.2^{-/-}* mutants perform ram feeding.

(**A**) Wildtype zebrafish (6 mpf) use suction feeding, following the general principles of suction feeding mechanics of actiopterygians: lower jaw depression, forward swing of premaxilla and maxilla, expansion of parabranchial cavity, and recoiling motions in that order. An entire cycle takes approximately 0.1 s. (**B**) No suction feeding was observed in *nkx3.2^{-/-}* zebrafish (6 mpf); instead, they perform ram feeding (swim through food) with the fixed open gape. In this particular feeding episode, the mutant initiated a cycle with detection of food (change in swimming orientation) and turned laterally to exit that swimming trajectory in approximately 0.2 s. Frame by frame still images from a film captured at 60 frames per second. The movie is available as Supplementary Information (Movie S1).

Fig. 6. Adult *nkx3.2^{-/-}* zebrafish converge onto overall head shapes of anaspids and thelodonts.

(**A**, **B**) Skulls of adult zebrafish in left lateral view via micro-computed tomography (μCT) showing wildtype and *nkx3.2^{-/-}* null mutant zebrafish (allele *ua5011*; 20 bp deletion in homeodomain). The jaw joint is indicated by an asterisk (*) in wildtype (A) and absent in mutant (B). Adult mutants display dramatic phenotypes in the jaw, snout, lips, and orobranchial regions. See also 3D rendering in multiple angles (Movie S1). (**C**, **D**) Skulls of extinct jawless vertebrates, showing general resemblance to the skull shape of *nkx3.2^{-/-}* zebrafish. Morphometric analysis corroborates this similarity. (C) Skull of an anaspid (birkeniid birkeniiform) from the Upper Silurian Cape Philips Formation of Cornwallis Island, Canada (Geological Survey of Canada C-26661-005) in left lateral view. (D) Skull of *Sphenonectris turnerae*, a thelodont (furcacaudiid furcacaudiform) from the Lower Devonian Road River Formation of Northwest Territories, Canada (University of Alberta Laboratory for Vertebrate Palaeontology, specimen number 42212).

(**E**, **F**) Landmark-based geometric morphometric comparison of *nkx3.2* phenotype using the thin-plate-spline Procrustes superimposition and principal component (PC) analysis. (**E**) Histogram

showing loading on each principal component in eigenvalue (%). PCs 1 and 5 are highlighted in blue.

These two PCs were chosen for comparison between groups, as these are the two largest components

that set apart *nkx3.2* homozygous mutants and wildtype zebrafish from each other at adult stage. PCs 2-

4, though accounting for greater variation than PC5, primarily distinguish among anaspids, or within

wildtype and mutant zebrafish groups. The original dataset is available (Data Supplement 3). (**F**) A

Cartesian plot of PCs 1 and 5, comparing morphospace occupation between wildtypes (2 mpf) (blue),

nkx3.2 homozygous mutants (2 mpf) (red), and anaspids (green), each with 90% ellipse. End of each

axis is labeled with a thin-plate-spline shape at that position (dark outline) against mean shape (grey

silhouette). A representative specimen is shown with landmarks labeled (yellow circles) for each group

in left lateral view. Each data point is a unique biological replicate (a specimen). The anaspid example

is *Birkenia elegans* (National Museum of Scotland specimen number 1929.5.6 from a Silurian locality

of Scotland). For anaspids, each data point is labeled with taxonomic identifications: B, *Birkenia*

elegans; C, Birkeniidae indet. from the Cornwallis Island; H, *Pharyngolepis oblongus*; L, *Lasanius*

problematicus; R, *Ryncholepis parvulus*; T, *Pterygolepis nitidus*.

(**G**) Simplified phylogenetic tree of vertebrates to illustrate distant relationships among the taxa

compared in this paper. A 60 dpf gravid female specimen is shown for *nkx3.2*$^{-/-}$ zebrafish (Fig. 3M).

Photographs of an anaspid and a thelodont are the same individuals in the respective panels showing

cranial morphology (D, E). Grey triangle indicates a crown group (not to scale). Dark silhouettes and

long branches indicate crown groups, whereas grey silhouettes and short branches represent extinct

lineages.