A multivariate phylogenetic comparative method incorporating a flexible function between discrete and continuous traits

Yuki Haba[1] & Nobuyuki Kutsukake[2]

1. Department of Ecology and Evolutionary Biology, Princeton University (corresponding author: yhaba@princeton.edu)
2. Department of Evolutionary Studies of Biosystems, Sokendai, The Graduate University for Advanced Studies, Hayama, Kanagawa 240-0193, Japan, (kutsu@soken.ac.jp)

Abstract

One major challenge of using the phylogenetic comparative method (PCM) is the analysis of the

evolution of interrelated continuous and discrete traits in a single multivariate statistical

framework. In addition, more intricate parameters such as branch-specific directional selection

have rarely been integrated into such multivariate PCM frameworks. Here, originally motivated

to analyze the complex evolutionary trajectories of group size (continuous variable) and social

systems (discrete variable) in African subterranean rodents, we develop a flexible approach using

approximate Bayesian computation (ABC). Specifically, our multivariate ABC-PCM method

allows the user to flexibly model an underlying latent evolutionary function between continuous

and discrete traits. The ABC-PCM also simultaneously incorporates complex evolutionary

parameters such as branch-specific selection. This study highlights the flexibility of ABC-PCMs

in analyzing the evolution of phenotypic traits interrelated in a complex manner.

Key words: Phylogenetic comparative method (PCM), approximate Bayesian computation

(ABC), multivariate analysis, social evolution, African mole rats

Introduction

Phylogenetic comparative methods (PCMs) provide a powerful statistical framework for investigating the patterns and processes of trait evolution (Felsenstein 1985; Harvey & Pagel 1991; Nunn 2011; Garamszegi 2014a, 2014b). The recent development of PCMs permits analyses of biologically interrelated discrete and continuous variables in a single multivariate statistical framework (Table 1). The development of such multivariate PCMs is crucial for two reasons. First, conducting two separate univariate analyses for a continuous trait and a discrete trait is redundant when the two variables are interrelated. Second, and more importantly, separate univariate analyses will miss the opportunity to consider important biological links between these traits.

The threshold model (Felsenstein 2005, 2012) was among the first PCMs to fully combine both discrete and continuous traits. The idea of the threshold model was originally developed in quantitative genetics by Wright (1934) to understand how multiple underlying genetic loci contribute to categorical traits such as the number of digits in guinea pigs. The threshold model assumes an unobservable continuous trait called 'liability' that underlies a discrete trait of interest. Because the liability is a continuous trait, Brownian motion has been conventionally used to model its evolution. Then, the state of the discrete trait of interest is determined by whether the liability trait value is below or above a particular threshold. This model allows users to incorporate both continuous and discrete traits in a straightforward way, as well as to estimate the covariance between the liability trait and other continuous traits of interest. However, because liability is unobservable, it is impossible to directly infer a latent function between the discrete trait and other observable continuous traits of interest. Moreover, although it is convenient to assume that the discrete trait is determined by a simple threshold (which can be treated as a probit link function in a framework of a phylogenetic generalized linear mixed model, PGLMM; Hadfield 2015; also see Ives and Garland 2014), it is unclear if the assumption is always biologically valid. At the very least, it is desirable for researchers to be able to assume other forms of latent functions (Fig. 1).

In addition to Felsenstein's threshold model, other methods that can link discrete traits and continuous traits have been proposed (Ives & Garland 2010, 2014; Hadfield and Nakagawa 2009; Hadfield 2015; see Table 1). For example, Ives and Garland (2010) developed a phylogenetic logistic regression to test the effects of observable continuous independent traits on a discrete dependent trait (phylogenetic logistic regression, Ives & Garland 2010). Hadfield and Nakagawa proposed a phylogenetic generalized linear mixed model employing a Bayesian approach (MCMCglmm, Hadfield and Nakagawa 2009; Hadfield 2015; also see Ives and

Garland 2014 for other approaches and a comparison of their performance). These approaches

successfully extended traditional linear models to enable nonlinear link functions (logit function

in a phylogenetic logistic regression; logit or probit functions in MCMCglmm) between discrete

and continuous traits. Notably, a probit-GLM can be mathematically equivalent to the threshold

model (see Hadfield 2015). However, limitations to these models still exist. For example, the

model by Ives and Garland (2010) assumes that the continuous traits are known values measured

empirically and does not allow the continuous traits to evolve along the tree (see Felsenstein

2012 and Hadfield 2015 for models that do not have such assumption; but also see Ives and

Garland 2014 for analyses of the relatively small effects of phylogenetic signal on continuous

traits). Moreover, the link function in the models is predetermined. In most cases, equipped

functions (e.g., logit or probit functions) are those that have useful mathematical properties for

analyzing a relationship between discrete and continuous traits. Still, cases may exist in which it

is biologically valid to establish a more complicated function between discrete and continuous

traits. Therefore, it is ideal to prepare a framework that enables researchers to examine a flexible

function to test specific hypotheses of interest.

Furthermore, analyses will be even more complex if additional variables of interest are

included, such as the presence of branch-specific directional selection (Kutsukake and Innan

2013, 2014). This complication often prevents the description of a likelihood function of the

model, which most conventional PCMs require (e.g., the maximum likelihood, Bayesian

approach). The aforementioned PCMs (Felsenstein 2005, 2012; Ives and Garland 2010, 2014;

Hadfield and Nakagawa 2009; Hadfield 2015) cannot incorporate branch-specific directional

selection into their models.

Here, we propose a PCM using approximate Bayesian computation (ABC) to overcome the

difficulties discussed above (Fig. 2). ABCs have been shown to facilitate flexible analyses in a

comparative framework and therefore have increasingly been applied to PCMs with intricate

evolutionary scenarios (Bokma 2010; Slater *et al*., 2012; Kutsukake and Innan 2013, 2014; Janzen

et al. 2015; Harano & Kutsukake 2018). Briefly, an ABC-PCM estimates parameters of interest by

simulating phenotypic evolution without a likelihood function (Beaumont 2010; Bertorelle *et al*.

2010; Csillery *et al*. 2010). The proposed parameters are accepted only when the simulated data

and real data are similar, and the accepted data comprise posterior distributions of parameters.

Thanks to this flexibility, ABC-PCMs enable researchers to test evolutionary models whose

likelihood function is mathematically intractable.

Our initial motivation for extending ABC-PCMs was to analyze a heterogeneous evolutionary

pattern between group size and social system in African subterranean mole rats (family

Bathyergidae; Table 2). In this clade, species have varied social structures that span solitary,

social, and eusocial states. Eusociality has independently evolved twice: in naked mole rats

(*Heterocephalus glaber*) and in Damaraland mole rats (*Fukomys damarensis*) (Jarvis 1981;

Sherman *et al*. 1991; Bennett & Faulkes 2000; Faulkes & Bennett 2013). We hypothesized that

the probabilities of evolutionary transitions among eu/social and solitary states (discrete trait)

change depending on group size (continuous trait). To capture all of these complex evolutionary

features, we are required to incorporate (1) the evolutionary trajectory of sociality and group

size, (2) the parameters for the trait-dependent latent functions, and (3) the presence of branch-

specific directional selection on group size in the two eusocial species. Certain aspects of these

features can be analyzed using previous methods; however, it is not possible to incorporate all

factors in a single model with those methods. Then, based on the example data, we discuss how

this ABC-PCM can be used for inferring other similar complex evolutionary processes.

Methods

ABC-PCM

Our ABC-PCM extends a previously developed framework (Fig. 2; Kutsukake & Innan 2013,

2014) designed to analyze heterogeneous evolutionary modes whose likelihood is not

straightforward to describe. We assume knowledge of the species tree Ψ, which consists of

information on tree shape and the lengths of all branches in the topology. We also assume that

trait data at the tips of the tree **D** have been observed.

In bivariate analyses of continuous and discrete traits, several causational patterns are

possible. For example, a discrete trait can be determined by a continuous trait that evolves on its

own, or vice versa. Alternatively, it is also possible to assume no *a priori* causality between the

continuous or discrete traits (Hadfield and Nakagawa 2009; Ives and Garland 2010). Any of

these cases can be modeled by the ABC-PCM framework proposed in this paper. Moreover, this

framework can be used regardless of whether a given value is latent (e.g., liability in the

threshold model) or measurable.

We hereafter consider a simple case of interrelated evolution in which a continuous trait

determines a discrete trait. In our example study of African subterranean mole rats, both traits

were measurable.

Briefly, our ABC-PCM process is implemented as follows (see Fig. 2 for a visual

schematic).

Let Λ be the parameter set to be estimated based on a hypothesis.

126    (Step 1) Determine prior distributions for all parameters in Λ. If prior biological

127    knowledge is available, it can be used to set informative (i.e., strong) prior distributions.

128    (Step 2) Parameters used in the simulation (Λ') are randomly generated from the prior

129    distributions.

130    (Step 3) Using Λ', the trait evolution is simulated on the phylogeny Ψ. In the current

131    model, the trait simulation has two parts corresponding to continuous and discrete trait

132    evolution (Fig. 1).

133      (Step 3a) The continuous trait evolves via Brownian motion (Felsenstein 1985; note

134      that other evolutionary models can be used; see Kutsukake and Innan 2013).

135      (Step 3b) Then the discrete trait is determined according to the probability of

136      transition between two states (A and B) as a function of the continuous trait. Note

137      again that this setting can be relaxed such that causation between continuous and

138      discrete traits can be flexibly changed according to a hypothesis of interest. Here,

139      $p(x)$ is the probability function that state A changes to B given the continuous trait x.

140      Similarly, $q(x)$ is the probability that state B changes to A given the continuous trait

141      x (Fig. 1).

142    (Step 4) Calculate the likelihood by comparing simulated data Θ with the real data $\boldsymbol{D}$ and

143    determine whether the parameter set is accepted or rejected. A joint probability (full

144    likelihood) for the comparison of n species can be calculated. In most cases, this

145    probability is difficult to obtain. In such cases, a composite likelihood that is proportional

146    to the full likelihood can be used as an approximate proxy. Intraspecific variation in the

147    trait data can be considered by assuming a certain distribution for the trait when

148    calculating the likelihood.

149    Then the acceptance or rejection of the parameters can be determined based on the

150    likelihood . Several methods of judgment exist (Marjoram *et al*. 2003; Marjoram and

151    Tavare 2006).

152    (Step 5) Repeat Steps 2–4 until a sufficient number of parameters Λ' is accepted. Then,

153    posterior distributions and credible intervals can be estimated.

155 Although the fundamental structure of ABC is straightforward, the number/choice of summary

156 statistics and the width of tolerance for judging the acceptance or rejection of simulated data at

157 Step 4 are controversial, and there is no general consensus on the choice of summary statistics

158 and tolerance (Beaumont *et al*. 2002; Csillery *et al*. 2010; Leuenberger & Wegmann 2010). In

this study, we use a combination of perfect match and joint probability as summary statistics (see

Acceptance and summary statistics below for details).

Application—social evolution in African mole rats

One example of complex evolution where continuous and discrete traits are interrelated is

social evolution. Sociality in animals can be characterized using a discrete classification based on

mating and/or social systems (e.g., Shultz *et al*. 2011). Average group size of a species, a

continuous trait, is also an important variable in characterizing sociality in animals (e.g.,

comparative analyses: Faulkes *et al*. 1997; Sheehan *et al*. 2015). Loss of sociality is correlated

with a decrease in group size (secondary loss of sociality; Wcislo & Danforth 1997; Beauchamp

1999; Sheehan *et al*. 2015). Likewise, cooperative social systems are likely to appear as the

group size increases (e.g., limited dispersal by ecological constraints: Emlen 1982; Duffy &

Macdonald 2010). Thus, the complex social evolution across African mole rats is an ideal system

to test our framework. Here, species-specific sociality and the mean group size for each species

(x) are the target traits.

Dataset

We surveyed the literature for field data on the sociality (solitary, social, or eusocial), mean

group size, and/or distribution of group sizes in each species (Table 2). When the distribution of

group size was available, we calculated its mean and standard deviation for each species. For

solitary species, we regarded the mean group size as one. In reality, however, their group sizes

can deviate from this value, because females may have dependent, pre-dispersal offspring; thus,

the group size of solitary species can also have a distribution. Therefore, we incorporated a

realistic value for the variance of solitary species (Table 2). We used the phylogeny presented in

Faulkes *et al*. (2004), who used mitochondrial genes 12s rRNA and cyt *b*. The mean value of the

estimated divergent interval in millions of years was used as the length of each branch.

Parameter set and trait simulation

The trait simulation included six parameters to be estimated: the group size of the most

recent common ancestor (MRCA) (θ), the rate of evolution (μ), the parameters of directional

selection (k_n and k_d), and the parameters for latent functions (a and b, or c and d, depending on

the model used; see next section). Table 3 shows the notation and settings of the prior

distributions.

The evolutionary process of the continuous trait was as follows. First, we generated the

values of six parameters from the prior distributions. Then, based on the proposed value of x at

the root (i.e., the group size of the MRCA, θ), we randomly determined the state (i.e., sociality)

of the MRCA by either $p(x)$ or $q(x)$. The number of evolutionary events that change traits (i.e.,

either an increase or decrease in group size) on each branch was respectively modeled as a

random number from a Poisson distribution with a mean equaling the product of the evolutionary

rate, μ (trait change per million years), and branch length τ (millions of years). The degree of

trait change by one evolutionary event is a random value from an exponential distribution, with

its mean being an arbitrary value φ (set to 0.01). φ corresponds to the average effect of one

change on the trait value. We used an exponential distribution because a change in trait due to

one evolutionary event would have small effects on traits in most cases, and large effects less

frequently. Thus,

$\qquad \Delta^+ = \mathrm{Pois}(\mu \times \tau_i) \times \exp(\varphi) \dots$ trait increase in a branch

$\qquad \Delta^- = \mathrm{Pois}(\mu \times \tau_i) \times \exp(\varphi) \dots$ trait decrease in a branch

where τ_i is the length of the ith branch. The total trait change on the branch is calculated as $\Delta^+ -$

Δ^-.

$\qquad$ Changes in group size caused by one evolutionary event would depend on the initial group

size. For example, an increase in the group size by one individual will have different biological

meanings in solitary and eusocial species. Therefore, we transformed the value of group size to a

natural logarithmic scale during trait simulation. Because the mean group size x cannot be less

than one, we enforced a lower bound of one. That is, when a trait change resulted in a value less

than one on a branch, the change was not implemented.

$\qquad$ We also tested whether there were selective pressures for increasing group size in the two

branches leading to eusocial species. In our ABC-PCM model, the parameter k represents

directional selection (Kutsukake & Innan 2013, 2014). The selection parameter k biases the

number of positive or negative trait changes such that the number of trait changes $\mathrm{Pois}(\mu \times \tau_i) \times$

$\exp(\varphi)$ is multiplied or divided by k, respectively (see Kutsukake & Innan 2013 for more details).

Thus,

$\qquad \Delta^+ = \mathrm{Pois}(\mu \times \tau_i \times k) \times \exp(\varphi) \dots$ trait increase in a branch

$\qquad \Delta^- = \mathrm{Pois}(\mu \times \tau_i / k) \times \exp(\varphi) \dots$ trait decrease in a branch

and again, the total trait change on the branch is calculated as $\Delta^+ - \Delta^-$.

$\qquad$ When $k = 1$, the evolutionary mode of trait evolution is asymptotically identical to the

Brownian motion. On the other hand, a significant departure of k from 1 is used as a signature of

directional selection.

In this analysis, we set selection parameters for the branches for both naked mole rats and

Damaraland mole rats as k_n and k_d, respectively (Table 3), to test our hypothesis that branch-

specific selective pressure has increased group size in eusocial species (Jarvis & Bennett 1993;

Young *et al.* 2015). When the 95% CI of the posterior distribution of k was larger than 1, it was

considered a signature of directional selection for larger group size.

Models of latent evolutionary functions

At each evolutionary step, the discrete trait value was determined by the continuous trait value

according to the latent functions (see Step 3b of *ABC-PCM*). Because the shape of this function

between sociality and group size was unknown, we took advantage of the flexibility of our ABC-

PCM framework and tested two examples of evolutionary models with different transition

functions, $p(x)$ and $q(x)$. We note that this framework is not limited to these two models but can

use any function, depending on the focal biological traits and hypotheses of interest. For

simplicity, we set the latent evolutionary functions such that $p(x) + q(x) = 1$, but this is not

required if one desires otherwise.

Model 1: Logistic function

In this model, we used a logistic function as the latent evolutionary function, which was similar

to a previous model using a generalized linear model for a binary dependent term (Ives and

Garland 2010). The transition functions $p(x)$ and $q(x)$ were defined as follows:

$$p(\mathrm{x}) = 1 - 1/(1 + \exp(-a \times (x-b)))$$

$$q(\mathrm{x}) = 1/(1 + \exp(-a \times (x-b)))$$

The parameter a determined the curvature of $p(x)$ and $q(x)$, i.e., the effects of the group size on

sociality (Fig. 1). When a is equal to 0, $p(x)$ and $q(x)$ are flat and the transition between social and

solitary is independent of group size (Fig. 1, right). When a is positive, a species with large group

size is likely to be social; when a is negative, a species with large group size is likely to be solitary.

Importantly, when $|a|$ is large enough, the transition between social and solitary is determined by

a certain group size, which virtually behaves like a step function (Fig. 1, left). The parameter b is

the x value at which $p(x)$ and $q(x)$ equal 0.5, and determines the group size at which the probability

of transitioning from solitary to social becomes larger than the probability of the reverse. Because

the curvature and midpoint of $p(x)$ and $q(x)$ (i.e., a and b) were unknown *a priori*, we set broad

prior distributions of a and b (Table 3).

Model 2: Exponential function

In the second model, we used an exponential function as the latent evolutionary function. We set

the transition functions $p(x)$ and $q(x)$ as follows:

When $d > 0$
$$\begin{cases} p(x) = \begin{cases} 1 - \exp\left(-c \times (x - d)\right), & x > d \\ 0, & x \le d \end{cases} \\ q(x) = \begin{cases} \exp\left(-c \times (x - b)\right), & x > d \\ 1, & x \le d \end{cases} \end{cases}$$

When $d \le 0$
$$\begin{cases} p(x) = \begin{cases} 1 - \exp\left(c \times (x - d)\right), & x > d \\ 0, & x \le d \end{cases} \\ q(x) = \begin{cases} \exp\left(c \times (x - d)\right), & x > d \\ 1, & x \le d \end{cases} \end{cases}$$

These functions describe an exponential decrease or diminishing increase in the transition

probability, and a given species with a mean group size of one is always a solitary species. Similar

to a and b in model 1, c and d determine the curvature of the exponential function and the point at

which the exponential decrease/increase begins, respectively. Note that one side of the exponential

function used here is asymptotically equal to either 0 or 1 when x is sufficiently large, but that of

the other side is not; therefore, we set the probability of state change as 0 or 1 when x is smaller

than d, at which the functions reach 0 or 1, respectively.

Acceptance and summary statistics

Among ABC-PCMs, no clear consensus has been reached concerning the number/choice of

summary statistics or the width of tolerance for judging the acceptance or rejection of simulated

data. In the present analyses, we used two summary statistics to assess the fit of the simulated

data to actual data. First, for the discrete variable (i.e., sociality), we only accepted simulations in

which the simulated data were a perfect match to the real data. For the continuous variable, we

used a conventional method that uses a joint probability (Kutsukake & Innan 2013, 2014); we

first calculated the probability that the real trait value is gained under a simulated trait value for

all 10 species. When calculating the probability, group size (back-transformed to an arithmetic

scale from a log-transformed value) was assumed to be normally distributed with a mean and

standard deviation equal to those of the real data. Then we calculated the product of those 10

probabilities and used the joint probability as a summary statistic. Simulated parameter sets were

accepted in proportion to the joint probability. For example, if the joint probability was 0.8 for a

simulation, the parameter set of the simulation has an 80% chance of being accepted. Because

the joint probability can be considered a direct likelihood, this method is superior to the standard

ABCs, which depend on the choice of summary statistics and arbitrary tolerance (discussed in Kutsukake & Innan 2013, 2014).

In total, 500 parameter sets were collected for estimating posterior distributions. All results were visualised using R version 3.5.3 (R Core Team 2017). The simulation code was written in C and is available in a public repository (https://github.com/YukiHaba/ABC-PCM).

Results

Model 1: Logistic functions

When the latent evolutionary function was logistic, the social system of the MRCA was estimated to be solitary in 61.4% of the cases (Fig. 2a, node A). The accepted functions of the transition probabilities varied widely (Fig. 3a), but showed several consistent patterns. First, the curvature parameter of functions, a, was positive in all cases (Table 3, Fig. 3b; see Supplementary material for a low risk of type I error), indicating that the probability of transition from a solitary to social species increased as group size increased. In addition, $p(x)$ typically increased rapidly around $x = 2$ to 4, which corresponded to the peak of the posterior distribution of b (Table 3, Fig. 3c).

At each internal node before reaching the common ancestor of social species, the inferred social system was predominantly solitary (Fig. 2a). The common ancestor of solitary species was likely to be solitary (Fig. 2a, node C), and a similar pattern was evident at the node of the common ancestor of *B. janetta* and *B. suillus*. By contrast, at the nodes leading to the clade of *Fukomis*, predominantly social states were inferred (Fig. 2a, node B). The estimated group sizes were consistent with the inferred social systems at internal nodes (Fig. 2b–d); the more social the system was likely to be, the larger the group size.

We detected marginal directional selection for increasing group size in the branch that led to naked mole rats (the proportion of $k_n < 1$ was 6.4%, Fig. 2f). By contrast, we did not detect significant directional selection in the branch leading to another eusocial species, Damaraland mole rats (the proportion of $k_d < 1$ was 15.8%, Fig. 2g). Based upon this non-significant result for k_d, we repeated the estimation after excluding k_d (i.e., directional selection was assumed only for the branch leading to naked mole rats), but the results did not change qualitatively (data not shown).

Model 2: Exponential functions

With the exponential latent evolutionary functions, the MRCA was inferred to be solitary in 60.4% of the cases (Fig. 4a, node A), similar to model 1. Again, although the accepted functions

of the transition probabilities between social and solitary varied (Fig. 5a), the curvature of $p(x)$,

c, was always positive (Table 3, Fig. 5b; see Supplementary material for a low risk of type I

error).

Group size and sociality at each internal node were almost identical to those in model 1.

Within the clade of *Fukomis*, we inferred predominantly social states (Fig. 4a, node B) and

relatively large group sizes (Fig. 4c). Within the clade of *Bathyergus* and *Georychus*, in contrast,

the social system at each node was consistently solitary (Fig. 4a, node C) and the group size was

around 2 to 4 (Fig. 4d).

We detected significant directional selection for increasing group size in the branch

leading to naked mole rats (the proportion of $k_n < 1$ was 3.4%; Fig. 4f). However, we did not

detect significant directional selection in the branch to eusocial Damaraland mole rats (the

proportion of $k_d < 1$ was 16.2%; Fig. 4g). Again, based on this non-significant result for k_d, we

repeated the estimation after excluding k_d (i.e., directional selection was assumed only for the

branch leading to naked mole rats); however, the result did not change qualitatively (data not

shown).

Discussion

This study proposes an extension of a previously described ABC-PCM (Kutsukake & Innan

2013, 2014) to analyze complex evolutionary scenarios in which discrete and continuous traits

are biologically interwoven. This is the first ABC-PCM framework that allows simultaneous

analyses of the interdependent evolution of discrete and continuous traits. In addition, our model

offers at least two advantages over other existing methods.

First, this study incorporated a feature that has rarely been included in a PCM framework:

branch-specific directional selection (Kutsukake & Innan 2013, 2014; Harano & Kutsukake

2018). This inclusion was possible thanks to the flexibility of the ABC-PCM, which does not

require the mathematical expression or analytic solution of a likelihood function. Second, our

multivariate model can incorporate flexible user-defined functions that describe the evolutionary

relationship between continuous and categorical traits. Previous methods can incorporate

nonlinear functions such as logit and probit functions (Ives and Garland 2010; Hadfield &

Nakagawa 2009; Hadfield 2015), but the choice of a latent function is less flexible than in our

present method. Another well-established method for simultaneously analyzing both continuous

and categorical traits is the threshold model (Felsenstein 2005, 2012). The threshold model is

equivalent to a phylogenetically controlled linear model with a probit latent function (see

Supplementary Material in Hadfield 2015 for a useful graphical representation). Although the

assumption is mathematically convenient for analyses, it may be an oversimplification of the

evolutionary relationship between the continuous and categorical traits of interest.

Applying this new approach to data on African subterranean rodents, we estimated the

intricate evolutionary trajectory of sociality and group size. Two models with different latent

evolutionary functions were tested: logistic and exponential. Although the exponential function

has not been used in previous PCMs, we believe that this function is suitable for our study

species for two reasons. First, group size should have a minimum value of one, and second, a

species whose group size is one must be a solitary species. Although these functions have

different mathematical characteristics and are not nested, they have common features such as a

monotonic increase and an asymptotical approach from one to zero. Potentially due to these

common features, the estimated parameters showed similar posterior distributions. In both

models the social state of the MRCA was not decisively solitary or social, and its estimated

group size varied (node A in Figs. 2a and 4a).

The latent evolutionary functions $p(x)$ and $q(x)$ had a consistent pattern in both models

(Figs. 3 and 5). Namely, most of the accepted functions had a steep transition around a group

size of 2 to 4. This indicates that this particular window of group size was an evolutionary

tipping point of the transition between solitary and social states. This study is the first to

quantitatively infer the range of group size that is crucial to the evolution of sociality in this

clade.

We tested differential selective pressure on group size in the branches leading to the two

eusocial species in this clade (k_n and k_d); we detected directional selection (k_n) for larger group

size in the branch leading to naked mole rats in model 2 (Fig. 4f) but not in model 1 (Fig. 2f). In

both models, a selective pressure was not detected in the branch leading to Damaraland mole rats

(k_d, Figs. 2g and 4g). Thus, although both species are deemed eusocial, the two species may have

undergone qualitatively different evolutionary paths (Burda *et al*. 2000). Future studies should

explore the differences in the ecological and evolutionary causes of the evolution of these two

eusocial species.

One critical limitation of this case study was the relatively low sample size (10 species;

see Supplementary material). This sample size is caused by three inevitable limitations. First, the

family Bathyergidae is a monotypic group composed of less than 20 OTUs, which is an

inevitable constraint on sample size. Second, detailed data for group size are not available for all

species of this family, which further limits the available data. Finally, and most importantly,

expanding the number of study species to include closely related non-subterranean species is

questionable, as it is highly likely that the underground ecological niche has affected predation

pressure and consequently sociality. More broadly, it remains unclear whether heterogeneity of

traits other than the traits of interest would affect biological interpretation in PCM studies based

on a large sample size. It is generally believed that larger sample sizes (a larger number of

species, or more precisely, a larger number of evolutionary transitions) would produce a more

powerful test in PCMs. While this is true, it remains largely undiscussed how the inclusion of

species that have fundamentally different ecological or behavioral traits would affect the results.

In this sense, in addition to a large-scale comparison, it is important to conduct PCMs that focus

on a taxon that shares fundamental traits, even if the sample size is not very large.

One major challenge in ABC-PCMs is that they are computationally intensive. In our

study, for example, it took several days to accept one parameter set (MacPro, OS 10.6.7, 2 x 2.93

GHz Quad-Core Intel Xeon; also see Kutsukake & Innan 2013). To compare the performance of

our model to an available method, we ran a bivariate model on our dataset and phylogeny using

MCMCglmm (Hadfield & Nakagawa 2009; Hadfield 2015) assuming that group size and

sociality follow Gaussian and probit ("threshold") families, respectively. For simplicity, we ran

the program without incorporating intraspecific variation, branch-specific selection, or the

bounding of group size. The MCMCglmm results qualitatively agreed with our results (e.g., 95%

CI of the covariance between group size and sociality > 0), yet the process took only a few

minutes to an hour on a standard laptop, depending on the parameters (results not shown). Thus,

the scope of the present ABC-PCM algorithm is currently limited to analyses of relatively small

to moderate numbers of species (29 species in Harano & Kutsukake 2018), but it is not suitable

for a large number of species. Improvements in both efficient simulation algorithms and

computational power will enable analyses based on a larger number of species.

In summary, we have developed a flexible multivariate ABC-PCM that has great potential

for testing biologically intricate scenarios of trait evolution. Although our analyses considered the

evolutionary transition of sociality in a relatively small number of species, our method can be

applied to other topics and to larger datasets. Despite the fact that our analyses focused on a simple

case in which a continuous trait affects the state of a discrete trait, other causational patterns can

be dealt with by a flexible setting of an evolutionary simulation. Furthermore, our ABC-PCM

framework can also be extended to model more complex evolutionary trajectories, such as

asymmetric transitions between states and/or more than two states of discrete traits with different

transition functions. Thus, our method allows evolutionary biologists to explore various

hypotheses of interest concerning the evolution of interrelated traits.

Acknowledgements

We would like to thank Dustin R. Rubenstein, Rafael Maia, and Margaret E. O'Brien at Columbia

University and Jessica Zung at Princeton University for useful comments and advice. Masahito

Tsuboi at University of Oslo also provided useful insights on the earlier version of manuscript.

Two reviewers gave detailed and helpful comments on the manuscript. This study was financially

supported by MEXT (No. 25711025) to K. N.

Author contributions

434   Y. H. and K. N. conceived, designed, and performed the analysis. Both discussed the result and

435   wrote the final manuscript.

Conflict of interest

438   None declared.

Literature cited

Bennett, N.C., Jarvis, J.U.M. & Cotterill, F. P. D. 1994. The colony structure and reproductive biology of the afrotropical Mashona mole-rat, Cryptomys darlingi. *J. Zool*. **234**: 477-487.

Beaumont, M.A. 2010. Approximate Bayesian computation in evolution and ecology. *Ann. Rev. Ecol. Evol. Syst.* **41**:379–406.

Beaumont, M. A., Zhang, W. & Balding, D. J. 2002. Approximate Bayesian computation in population genetics. *Genetics* **162**:2025–2035.

Beauchamp, G. 1999. The evolution of communal roosting in birds: origin and secondary losses. *Behav. Ecol.* **10**: 675-687.

Bennett, N.C. & Faulkes, C.G. 2000. African mole-rats: ecology and eusociality. Cambridge, UK: Cambridge University Press.

Bokma, F., 2010. Time, species, and separating their effects on trait variance in clades. *Syst. Biol.* **59**:602–607.

Burda, H., Honeycutt, R.L., Begall, S., Locker-Grütjen, O. & Scharff, A. 2000. Are naked and common mole-rats eusocial and if so, why? *Behav. Ecol. Sociobiol.* **47**: 293-303.

Burda, H. & Kawalika M. 1993. Evolution of eusociality in the Bathyergidae: the case of the giant mole-rat (*Cryptomys mechowi*). *Naturwissenschaften* **80**:235-237.

Csillery, K., Blum, M.G., Gaggiotti, O.E. & François, O. 2010. Approximate Bayesian computation (ABC) in practice. *Trends. Ecol. Evol.* **25**:410–418.

Duffy, J.M. & Macdonald, K.S. 2010. Kin structure, ecology and the evolution of social organization in shrimp: a comparative analysis. *P. R. Soc. B.* **277**: 575-584.

Emlen, S.T. 1982. The evolution of helping. I. An ecological constraints model. *Am. Nat.* **119**: 29–39.

Faulkes, C.G., Bennett, N.C., Bruford, M.W., O'brien, H.P., Aguilar, G.H. & Jarvis, J.U.M. 1997. Ecological constraints drive social evolution in the African mole–rats. *P. R. Soc. B.* **264**: 1619-1627.

Faulkes, C.G., Verheyen, E., Verheyen, W., Jarvis, J.U.M. & Bennett, N.C. 2004.

Phylogeographical patterns of genetic divergence and speciation in African mole-rats

(Family: Bathyergidae). *Mol. Ecol.* **13**: 613-629.

Faulkes, C.G. & Bennett, N.C. 2013. Plasticity and constraints on social evolution in African

mole-rats : ultimate and proximate factors. *Philos. T. R. Soc. B.* **368**:20120347.

Felsenstein, J. 1985. Phylogenies and the comparative method. *Am. Nat.* **125**: 1–15.

Felsenstein, J. 2005. Using the quantitative genetic threshold model for inferences between and

within species. *Philos. T. R. Soc. B.* **360**: 1427-1434.

Felsenstein, J. 2012. A comparative method for both discrete and continuous characters using the

threshold model. *Am. Nat.* **179**: 145-156.

Garamszegi, L.Z. 2014a. *Modern Phylogenetic Comparative Methods and their Application in*

Evolutionary Biology - Concepts and Practice, (L.Z. Garamszegi, ed.). Springer,

Heidelberg.

Garamszegi, L.Z. 2014b. Uncertainties due to within-species variation in comparative studies:

measurement errors and statistical weights. In: *Modern Phylogenetic Comparative*

Methods and their Application in Evolutionary Biology - Concepts and Practice, (L.Z.

Garamszegi, ed.) pp. 157-199. Springer, Heidelberg.

Hadfield, J.D. & Nakagawa, S. 2009. General quantitative methods for comparative biology:

phylogenies, taxonomies and multi-trait models for continuous and categorical characters.

J. Evol. Biol. **23**: 494-508.

Hadfield, J. D. 2015. Increasing the efficiency of MCMC for hierarchical phylogenetic models of

categorical traits using reduced mixed models. *Methods in Ecology and Evolution*

6.6:706-714.

Harano, T. & Kutsukake, N. 2018. Directional selection in the evolution of elongated upper

canines in clouded leopards and sabre-toothed cats. *J. Evol. Biol.*, **31**: 1268-1283.

Harvey P.H. & Pagel, M.D. 1991. The comparative method in evolutionary biology. Oxford

University Press, Oxford

Ives, A. R. & Garland, T. 2010. Phylogenetic logistic regression for binary dependent variables.

Syst. Biol. **59**: 9-26.

Ives, A. R. & Garland, T. 2014. Phylogenetic Regression for Binary Dependent Variables.

In: *Modern Phylogenetic Comparative Methods and their Application in Evolutionary*

Biology - Concepts and Practice (L.Z. Garamszegi, ed.), pp. 231-261. Springer,

Heidelberg.

Janzen, T., Hoehna, S. & Etienne, R. S. 2015. Approximate Bayesian Computation of

diversification rates from molecular phylogenies: introducing a new efficient summary

statistic, the nLTT. Methods Ecol. Evol. **6**: 566-575.

Jarvis, J.U.M. & Bennett, N.C. 1993. Eusociality has evolved independently in two genera of

bathyergid mole-rats—but occurs in no other subterranean mammal. *Behav. Ecol.*

Sociobiol. **33**: 253-260.

Jarvis, J.U.M. 1981. Eusociality in a mammal: cooperative breeding in naked mole-rat colonies.

Science **212**: 571–573.

Jones, K., Bielby, J., Cardillo, M. & Fritz, S. 2009. PanTHERIA: a species-level database of life

history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**:

2648.

Kutsukake, N. & Innan, H. 2013. Simulation-based likelihood approach for evolutionary models

of phenotypic traits on phylogeny. *Evolution* **67**: 355-367.

Kutsukake, N. & Innan, H. 2014. Detecting phenotypic selection by approximate Bayesian

computation (ABC) in phylogenetic comparative methods. In: *Modern Phylogenetic*

Comparative Methods and their Application in Evolutionary Biology - Concepts and

Practice (L.Z. Garamszegi, ed.), pp. 409-424. Springer, Heidelberg

515 Leuenberger, C. & Wegmann, D. 2010. Bayesian computation and model selection without

516   likelihoods. *Genetics* **184**: 243-252.

517 Lewis, P. O. 2001. A likelihood approach to estimating phylogeny from discrete morphological

518   character data. Syst. Biol. **50**: 913–925.

519 Marjoram P, Tavare S. 2006. Modern computational approaches for analysing molecular genetic

520   variation data. Nat. Genet. Rev. **7**:759–770

521 Marjoram P, Molitor J, Plagnol V, Tavare S. 2003. Markov chain Monte Carlo without

522   likelihoods. Proc. Natl. Acad. Sci. USA **100**:15324–15328

523 Nunn, C.L. 2011. The comparative approach in evolutionary anthropology and biology.

524   University of Chicago Press, Chicago

525 Pagel, M. 1994. Detecting correlated evolution on phylogenies: a general method for the

526   comparative analysis of discrete characters. *P. R. Soc. B.* **255**: 37-45.

527 R core team. 2017. R: A language and environment for statistical computing. *R Found. Stat.*

528   *Comput. Vienna, Austria.*

529 Sheehan, M.J., Botero, C.A., Hendry, T.A., Sedio, B.E., Jandt, J.M., Weiner, S., Toth, A.L. &

530   Tibbetts, E.A. 2015. Different axes of environmental variation explain the presence vs.

531   extent of cooperative nest founding associations in *Polistes* paper wasps. *Ecol. Lett.* **18**:

532   1057-67.

533 Sherman, P.W., Jarvis, J.U.M. & Alexander, R.D. 1991. The biology of the naked mole-rat.

534   Princeton, NJ: Princeton University Press.

535 Shultz, S., Opie, C. & Atkinson, Q.D. 2011. Stepwise evolution of stable sociality in primates.

536   *Nature* **479**: 219-222.

537 Sichilima, A.M., Faulkes, C.G. & Bennett, N.C. 2008. Field evidence for aseasonality of

538   reproduction and colony size in the Afrotropical giant mole-rat *Fukomys mechowii*

539   (Rodentia: Bathyergidae). *Afr. Zool*. **43**: 144-149.

Sichilima, A.M., Bennett, N.C., Faulkes, C.G. & Bronner, G.N. 2011. Field evidence for colony

size and aseasonality of breeding and in Ansell's mole-rat, *Fukomys anselli* (Rodentia:

Bathyergidae). *Afr. Zool.* **46**: 334-339.

Slater, G. J., Harmon, L.J., Wegmann, D., Joyce, P., Revell, L.J. & Alfaro, M.E. 2012. Fitting

models of continuous trait evolution to incompletely sampled comparative data using

approximate Bayesian computation. *Evolution* **66**:752–762 .

Van Daele, P.A., Blonde, P., Stjernstedt, R. & Adriaens, D. 2013. A new species of African

mole-rat (Fukomys, Bathyergidae, Rodentia) from the Zaire-Zambezi watershed. *Zootaxa*

3636: 171-189.

Wcislo, W.T. & Danforth, B.N. 1997. Secondarily solitary: the evolutionary loss of social

behavior. *Trends Ecol. Evol.* **12**: 468-474.

Wright, S. 1934. An analysis of variablity in number of digits in an inbred strain of guniea pigs.

Genetics. **19**(6): 506-536.

Young, A.J., Jarvis J.U.M., Barnaville, J. & Bennett, N.C. 2015. Workforce effects and the

evolution of complex society in wild Damaraland mole rats. *Am. Nat.* **186**: 302-311.

Table 1. **Summary of previous methods and the originality of this study.**

| | Approach | |
Trait	Univariate models	Multivariate models (link function)
Discrete	Mk model[1] Threshold model[2,*] PLogReg[3] (w/o independent variables)	Mk model[1] Threshold model[2,*]
Continuous	PLogReg[3] MCMCglmm[4,5]	MCMCglmm[4,5]
Both discrete and continuous	-	Threshold model[2] (determined by a threshold)[*] PLogReg[3] (logit link) MCMCglmm[4,5] (logit and probit link) This study (researcher-defined function)

[1]Pagel 1994; Lewis 2001
[2]Felsenstein 2012
[3]Phylogenetic logistic regression, Ives and Garland 2010
[4]Hadfield and Nakagawa 2009
[5]Hadfield 2015
[*]continuous trait (liability) that is unobservable
563

564 **Table 2**. **Data used in this study.**

Species	Social system	Group size, mean (SD)
Naked mole rat, *Heterocephalus glaber*[1]	eusocial	75 (48.65)[A]
Damaraland mole rat, *Fukomys damarensis*[1]	eusocial	11 (6.26)
Whyte's mole rat, *Fukomys hottentotus*[1]	social	5.16 (2.62)[B]
Mechow's mole rat, *Fukomys mechowii*[2]	social	9.91 (2.49)
Mashona mole rat, *Fukomys darlingi*[3]	social	7 (not available[C])
Ansell's mole rat, *Fukomys anselli*[4]	social	8.72 (2.15)
Cape mole rat, *Georychus capensis*[5]	solitary	1[D]
Cape dune mole rat, *Bathyergus suillus*[5]	solitary	1[D]
Namaqua dune mole rat, *Bathyergus janetta*[5]	solitary	1[D]
Silvery mole rat, *Heliophobius argenteocinereus*[5]	solitary	1[D]

565
[1]Bennett & Faulkes 2000; [2]Sichilima *et al*. 2008; [3]Bennett *et al*. 1994; [4]Sichilima *et al*. 2011; [5]Burda & Kawalika
1993; see Van Daele *et al*. (2013) for a recent classification.
[A]Because group sizes were categorized at an interval of 25 individuals, we used the mean value of each range for
calculating the mean and SD (e.g., 37.5 for groups of 25 to 50 individuals).
[B]When calculating the mean group size of this species, we used a group size of 11 when it exceeded 10, because the
original data for group size pooled group sizes larger than 10 into ">10" (p. 92 in Bennett & Faulkes 2000).
[C]Because no data on intraspecific variation were available, we used the mean value of the SD, 2.42, of the other three
social species.
[D]The mean litter size ranges from 2.46 to 5.94 in solitary species (Jones *et al*. 2009). To account for this, we set a
value of 2.56 as their SD to cover the range of possible group sizes.

Table 3. Parameters estimated in this study.

Parameter	Notation	Prior distribution	Posterior distribution (95% CI)	
			Model 1 (logistic)	Model 2 (exponential)
Most recent common ancestor (MRCA)	θ	U(1.001, 10)	1.1–5.22	2.45–9.15
Baseline evolutionary rate (per million years)	μ	U(0.001, 30)	6.4–29.5	3.62–28.3
Directional selection (naked mole rats)	k_n	U(0.001, 30)	0.99–3.57	1.01–1.93
Directional selection (Damaraland mole rats)	k_d	U(0.001, 5)	0.82–4.22	0.82–4.64
Curvature of transition functions (solitary $\rightleftarrows$ social)	*a or c*	U(-15,15)	*a*: 1.52–14.50	*c*: 1.28–14.60
Position of transition functions	*b or d*	U(0,10)	*b*: 1.52–4.94	*d*: 1.28–5.07

Fig. **1. Variation in multivariate PCMs with a latent function between discrete and**
continuous traits. Conventional PCMs can explore various latent evolutionary functions
between continuous and discrete traits (Table 1). Here, we consider a simple example of the
evolutionary link between a continuous and a discrete trait. The continuous trait (x) evolves by a
process similar to Brownian motion. The discrete trait, on the other hand, evolves according to
the transition probabilities $p(x)$ and $q(x)$, i.e., the latent evolutionary functions. The state of the
discrete trait at each time step is shown as a white (A) or black (B) square. At one extreme, the
transition of discrete states is determined by a certain "threshold" value of the continuous trait.
At the other extreme, the discrete trait and the continuous trait are independent. In such a case,
the transition probabilities do not vary with the continuous traits (right). In our framework, any
functions between the two extreme cases can be incorporated based on biological hypotheses
(middle). Furthermore, x can be either a measurable trait or a latent value. Here, logistic and
exponential functions are shown as examples.

Fig. **2. Our ABC-PCM with latent evolutionary functions.** Schematic of our ABC-PCM
approach. First, determine parameters of interest and set prior distributions (Step 1). Example
parameters include phenotype of the MRCA (θ_0), evolutionary rate (μ), directional selection
parameter (k), latent function parameter (a), and so on. Next, generate random values for
parameters from the prior distributions (Step 2). Trait simulations on a phylogeny Ψ are then
conducted (Step 3), and the simulated data Θ are compared to real data D to determine whether
the data are accepted or rejected (Step 4). A number (N) of simulations are conducted until
enough samples are collected to infer the posterior distribution (Step 5).

Fig. **3. The estimated evolutionary trajectory with the latent function being logistic (model 1)**. (a) The estimated parameters on the phylogeny. The social state of each tip species is shown as black (solitary) or white (social) squares. Asterisks indicate eusocial species. The proportion of solitary/social states and the 95% CI of the posterior distribution of group size (within brackets) are shown at each internal node. (b–d) The distributions of simulated group size at nodes A (MRCA), B, and C in the phylogeny. (e) The posterior distribution of the baseline evolutionary rate. (f, g) The posterior distributions of the selection coefficients k_n and k_d. The dashed line at $k = 1$ indicates the proportion of simulations in which directional selection for larger group size was detected (black histograms).

618 **Fig**. **4. Accepted latent functions in model 1**. (a) All 500 accepted latent functions $p(x)$, i.e., the

619 probability of transition from a solitary state to a social state as a function of group size. Most

620 accepted functions increased steeply at group sizes of 2 to 4. (b, c) The posterior distributions of a

621 and b, the curvature of the function and the group size at which $p(x)$ equals 0.5.

Fig. **5. The estimated evolutionary trajectory with the latent function being exponential (model 2)**. (a) The estimated parameters on the phylogeny. The social state of each tip species is shown as black (solitary) or white (social) squares. Asterisks indicate eusocial species. The proportion of solitary/social states and the 95% CI of the posterior distribution of group size (within brackets) are shown at each internal node. (b–d) The distributions of simulated group size at nodes A (MRCA), B, and C in the phylogeny. (e) The posterior distribution of baseline evolutionary rate. (f, g) The posterior distributions of the selection coefficients k_n and k_d. The dashed line at $k = 1$ indicates the proportion of simulations in which directional selection for larger group size was detected (black histograms).

637 **Fig**. **6. Accepted latent functions in model 2.** (a) All 500 accepted latent functions corresponding

638 to $p(x)$. Similar to the case in model 1, most of the accepted functions increased at a group size of

639 2 to 4. (b, c) The posterior distributions of c and d, the curvature of the function and the group size

640 at which $p(x)$ equals 0.