A Structurally-Validated Multiple Sequence Alignment

of 497 Human Protein Kinase Domains

Vivek Modi

Roland L. Dunbrack, Jr.*

Fox Chase Cancer Center

333 Cottman Avenue

Philadelphia PA 19111

* Correspondence: Roland.Dunbrack@fccc.edu

Abstract

Protein kinases are important in a large number of signaling pathways. Their dysregulation is

involved in a number of human diseases, especially cancer. Studies on the structures of

individual kinases have been used to understand the functions and phenotypes of mutations in

other kinases that do not yet have experimental structures. The key factor in accurate inference

by homology is an accurate sequence alignment. We present a parsimonious structure-based

sequence alignment of 497 human protein kinase domains excluding atypical kinases, even

those with related but somewhat different folds. Starting with a computed multiple sequence

alignment, the alignment was manually refined in Jalview based on pairwise structural

superposition onto a single kinase (Aurora A), followed by sequence alignment of the remaining

kinases to their closest relatives with known structures. The alignment is arranged in 17 blocks

of conserved regions and unaligned blocks in between that contain insertions of varying lengths

present in only a subset of kinases. The aligned blocks contain well-conserved elements of

secondary structure and well-known functional motifs, such as the DFG and HRD motifs. We

validated the multiple sequence alignment by a pairwise, all-against-all alignment of 272 human

kinases with known crystal structures. Our alignment has true-positive rate (TPR) and positive

predictive value (PPV) accuracies of 97%. The remaining inaccuracy in our alignment comes

from a few structures with shifted elements of secondary structure, and from the boundaries of

aligned and unaligned regions, where compromises need to be made to encompass the

majority of kinases. A new phylogeny of the protein kinase domains in the human genome

based on our alignment indicates that 14 kinases previously labeled as "OTHER" can be

confidently placed into the CAMK group. These kinases comprise the Aurora kinases, Polo

kinases, ULK kinases, Calcium/calmodulin-dependent kinase kinases, and STK36.

Introduction

Protein kinases catalyze the transfer of a phosphoryl group from an ATP molecule to substrate proteins[1], and are crucial for cellular signaling pathways[2]. Mutations in kinases that lead to gain of function are frequently observed in many cancer types[3,4], while mutations may also result in drug resistance rendering existing drugs inefficient[3]. Humans have over 500 genes that catalyze the phosphorylation of proteins, collectively called the 'kinome'[5].

Protein kinase activity is found in a number of protein families and superfamilies in the human proteome. The vast majority of human kinases come from one very large, diverse family that share a common fold consisting of an N-terminal lobe, composed of 5 β-sheet strands and an α-helix called the C-helix, and a C-terminal lobe comprising 6 α-helices[6]. The active site is located between the two lobes where the activation and catalytic loops form the ATP and substrate binding sites.

In 2002, Manning and coworkers identified 518 kinase genes in the human kinome[5] which they divided into 478 typical kinase genes (13 of them containing two kinase domains, for a total of 491) and 40 atypical kinase genes. The annotation of the human genome has improved since the Manning paper; currently Uniprot identifies 483 human proteins containing 496 typical kinase domains (https://www.uniprot.org/docs/pkinfam). Uniprot identifies 29 atypical human kinases. Some of these are distantly related to the typical kinase domain, thus making them a superfamily, including Alpha kinases[7], ADCK kinases[8], RIO kinases[9], FAM20C kinases[10], and the PI3-PI4 kinase family, which contains the protein kinases ATM, ATR, and MTOR[11]. In addition, there are proteins that do not appear to share an evolutionary relationship with typical kinases that also phosphorylate proteins, such as pyruvate dehydrogenase kinases[12].

For any large protein family, an accurate multiple sequence alignment is the basis of an accurate phylogeny[13] and structural and functional inferences[14]. In 2002, Manning et al. built a phylogenetic tree of 491 typical kinase domains from a multiple sequence alignment created

without using any kind of structural information[5]. The accompanying poster and image of this

tree is still widely used in scientific papers and presentations on kinases[15]. Multiple sequence

alignments of kinases have been used to extend structural and functional information from the

kinases with known structures to those without known structures. This includes the

conformations of active and inactive kinases[16-18], predictions of substrate specificity[19], and

analysis of kinase-drug interactions[20,21].

A common problem in multiple sequence alignments of large, diverse protein families is

that they are very 'gappy,' i.e., containing many gap characters in every sequence in order to

align inserted segments of different lengths that may be present in only a small subset of the

sequences. This is true of the alignment used to produce the kinome tree of Manning et al.[5] The

gappy regions are usually present between major elements of secondary structure, where the

family members may have widely divergent sequence loop lengths due to numerous insertions

that occurred in different lineages of family members. The gappiness makes alignments difficult

to visualize and produces errors in phylogenetic inference[22].

In this paper, we present a parsimonious, structure-based multiple sequence alignment

(MSA) of 497 human typical kinase domains from 484 human genes. We developed the MSA

from pairwise structure alignment to a reference kinase (Aurora A kinase) and sequence

alignment for kinases of unknown structure to their closest homologues of known structure. One

of our central goals was to develop a parsimonious alignment of kinases containing as few gap

regions as possible. Our alignment therefore contains aligned blocks (in upper case letters) that

represent common structural elements, usually of secondary structure elements and important

motifs, present in most or all of the kinases. These regions are separated by left-justified,

unaligned sequence regions in lower case letters containing insertions of different lengths

present in only subsets of the kinases. Our alignment contains 17 aligned blocks and 16

unaligned regions between them.

We validated this alignment with an all-against-all pairwise structure alignment of 272

protein kinases, and we show that the alignment is significantly more accurate than the Manning

alignment and other recent alignments for classifying kinases[16,23]. Finally, we used our

alignment to produce an accurate phylogeny of the kinase domains. Guided by the phylogeny

and HMMs for each of group we assign 14 kinases previously categorized as "OTHER" by

Manning et al. to the CAMK group, including Aurora kinases, Polo-like kinases,

Calcium/calmodulin-dependent kinase kinases, and Unc51-like kinases (ULKs).

RESULTS

Typical and atypical kinases

Kinase sequences were identified from the list of human kinases and pseudokinases

provided by Uniprot[24] (https://www.uniprot.org/docs/pkinfam), which divides 'typical' kinases into

nine groups (AGC, CAMK, CK1, CMGC, NEK, RGC, STE, TKL, TYR) and a group of OTHER

kinases. It also contains a list of 'atypical' kinases divided into six families (ADCK, Alpha-type,

PI3-PI4-related, RIO, PDK/BCKDK, and FASTK). To identify any kinases which are not included

in the Uniprot list, we searched all human Uniprot sequences with PSI-BLAST and each human

kinase domain as query. As a result, we were able to identify Uniprot entry PEAK3_HUMAN as

an additional human kinase. It was identified in searches starting with PRAG1 and PEAK1. The

relationship with protein kinases was confirmed with hhpred[25].

Manning et al. did not include any atypical kinase in their sequence alignment and

phylogenetic tree. However, before discarding them from our dataset we wanted to examine the

atypical kinase structures available from the PDB. Since Aurora A is a good representative of

typical protein kinase domains, we have used its structure as a reference to identify structural

similarities and differences between typical kinases and atypical kinases. The kinase domain of

Aurora A has 251 amino acid residues with 8 helices, 7 β strands, and all the known conserved

motifs without any unusually long insertions (Figure 1).

Figure 1: Structure of Aurora A (PDB: 3E5A_A) representing typical protein kinase domain. For the
description of the labeled structural elements refer to Table 2.

Structure alignment with Aurora A indicated that four of the atypical kinase families are

homologous to typical kinases (Figure 2), containing some elements of the typical kinase fold

but containing changes and additions in elements of secondary structure. These include ADCK,

Alpha-type, PI3-PI4-related, and RIO kinases. The ADCK (aarF-domain containing) kinases

consist of 5 proteins: ADCK1, ADCK2, COQ8A (ADCK3), COQ8B (ADCK4), and ADCK5. Only

the structure of COQ8A is available (PDB:4PED[8]). The structure consists of 384 residues, 13

helices, and 8 beta sheet strands. Structure alignment with FATCAT aligned 192 residues with

an RMSD of 3.92 Å, covering the N-terminal domain, the HRD and DFG motifs, and the E and F

helices of the C-terminal domain. COQ8A's N-terminal domain contains an additional

subdomain of five alpha helices, three of which precede the typical kinase domain and two of

which are inserted between beta strand B3 and the C-helix. Instead of the activation loop

leading into the F-helix, the DFG motif leads into a bundle of four alpha helices that precede

COQ8A's F-helix, which is followed by one additional helix.

Figure 2: Representative structures from four different families of atypical kinases showing differences
from typical kinase domains. **a)** ADCK protein kinase family - ADCK3 (4PED_A); **b)** Alpha-type - TRPM7
(1IAJ_A); **c)** PI3/PI4 - PIK3CA (4L2Y_A) and **d)** RIO-type Ser/Thr kinase family - RIOK1 (4OTP_A). The
regions without any structural similarity to the typical kinase domain are colored in yellow, as identified
by FATCAT and CEalign (in Pymol) after aligning to Aurora A (3E5A_A).

There are six kinases in the Alpha-type kinase family: ALPK1, ALPK2, ALPK3, EEF2K,

TRPM6, and TRPM7. The structure of mouse TRPM7 (PDB:1IAJ[26]) has been determined; only

the N-terminal domain and the E helix could be aligned to AURKA with an RMSD of 5.8 Å over

120 residues. The remainder of the C-terminal domain of TRMP7 consists of two beta sheet

strands, large coil regions, and a short helix. The human PI3/PI4 kinases consist of 7 genes:

ATM, ATR, MTOR, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PRKDC, and SMG1. All of these

except SMG1 have known structures (PIK3CB is represented by a structure of mouse PIK3CB).

The structure of PIK3CA (PDB: 4L2Y[27]) aligns with Aurora A with RMSD 6.0 Å over 168

residues. The structures of two of the three RIO kinases (RIOK1, RIOK2, and RIOK3) are

known. The structures of RIOK1 (PDB: 4OTP[28]) and RIOK2 (PDB: 6FDN[29]) can be aligned with

AURKA with 5.1 and 4.6 Å RMSD over 176 amino acids, because the orientation of the HG, HH,

and HI helices are entirely different.

Two of the atypical protein kinase families listed by Uniprot do not appear to be

homologous to typical kinases. The PDK/PCKDK protein kinase family consists of BCKDK,

PDK1, PDK2, PDK3, and PDK4. The structures of all of these proteins are known (with BCKDK

represented by rat BCKDK in the PDB by entry 3TZ5[30] and PDK1 by 2Q8H[31]), and none of them

resemble typical protein kinases. They consist of an N-terminal domain in the form of a bundle

of 4 long α helices and a C-terminal domain of a 5-stranded β sheet and 3 α helices. ECOD

(Evolutionary Classification of Protein Domains) also does not classify these structures in the

same homology group as the typical kinases[32]. While there is no structure of FAST kinase

(Uniprot FASTK_HUMAN), the program hhpred[25] found that the closest homologues in the PDB

are restriction endonucleases (e.g., PDB:1CW0[33]), which do not appear to be homologous to

typical kinases.

Overall, our examination indicated that every atypical family has significant differences

from the typical kinase domain in the arrangement or presence of secondary structural

elements. These differences make any an alignment with the sequences of the typical kinase

domain approximate and partial. Therefore, we did not include any atypical kinase sequence in

our multiple sequence alignment of human protein kinases.

Table 1: Number of kinase domains in each family in the multiple sequence alignment

	AGC	CAMK	CK1	CMGC	NEK	RGC	STE	TKL	TYR	OTHER	Total
Uniprots	62	94	12	65	11	5	47	42	90	62	484*
Domains	62	96	12	65	11	5	47	42	94	63	497
Domains with PDB	29	52	10	39	3	0	27	25	64	25	274
Included in validation	29	52	10	39	3	0	27	25	64	23	272

* 6 kinase genes have one AGC and one CAMK domain each (RPS6KA1, RPS6KA2, RPS6KA3, RPS6KA4, RPS6KA5 and RPS6KA6),
and are counted twice in this table; 2 kinases have 2 CAMK domains (OBSCN, SPEG); 4 kinases have 2 TYR domains (JAK1, JAK2,
JAK3, TYK2); 1 kinase has 2 domain in the OTHER category (EIF2AK4) (484+6+2+4+1=497 domains). Two kinases AGC_PDPK2P
and AGC_PRKY could be pseudogenes according to their Uniprot annotations.

A summary of the 497 typical kinase domains from 484 Uniprot sequences included in

our dataset and the available structures in the PDB are provided in Table 1. Thirteen kinases

have two kinase domains in the sequence (see caption to Table 1), and some kinases were

reassigned to different groups than the Uniprot or Manning designations (discussed below). The

domain boundaries were initially identified using the domain annotations in Uniprot, while some

of them were updated during the process of alignment. We labeled sequences by their HGNC

gene names[34] with a group name identifier from Uniprot. For example, the sequence for the

gene AKT1 is labeled as AGC_AKT1 and EGFR is labeled TYR_EGFR. Some of the common

kinases have gene names which are not easily recognized; for example VEGFR1, VEGFR2,

and VEGFR3 are labeled TYR_FLT1, TYR_KDR, and TYR_FLT4 respectively. A table of

Uniprot accession ids, Uniprot entry names, HGNC gene names, HGNC accessions, gene

name synonyms, kinase group names, and domain boundaries is provided in Table S1. These

sequences were used to create the MSA utilizing a variety of alignment programs and structural

information.

Forming a multiple sequence alignment of 497 human protein kinase domains

        To examine the boundaries of conserved segments and common insertions across
kinases, we first aligned the structure of Aurora A kinase N- and C-terminal domains separately
to one representative structure of each of 271 other human kinases in the PDB using the
program FATCAT[35]. We then used the SE program[36] to read the structural alignments and print
the corresponding sequence alignments. SE prints regions of low structural similarity (usually
because of insertions or deletions) in lower case letters, and left-justifies them. An example is
shown in Figure 3. To identify regions of structural variability in kinases, we counted the number
of times a residue in the Aurora A sequence was structurally aligned across all the pairwise
alignments output by SE (i.e. printed in upper case). This provides a count of how often each
residue is structurally conserved, indicating the locations of insertions, deletions, and structural
variations across the kinases (Figure 4). The tallest bars in the plot display the conserved
regions in the alignments while the shorter bars represent low similarity regions or segments
abutting common insertions and deletions. We used these blocks of conserved regions and
intermittent low similarity regions to guide the formation of the MSA.

Figure 3: Structural alignment of the activation loop region of CAMK_AURKA and CAMK_PRKAA1 by **a)**
FATCAT and **b)** the SE program. Structure alignment programs like FATCAT often introduce gaps in low
similarity loop regions to align segments that are not necessarily homologous. The SE program takes
coordinates of the superposed structures and produces an optimized sequence alignment with
structurally similar regions in upper case and low similarity residues in lower case letters to distinguish
them from aligned residues.

        The creation of the MSA was a multi-step process. The initial alignment of all the kinase
domain sequences was done using ClustalOmega[37] which aligned the main conserved regions

211   in a majority of the sequences up to the beginning of the activation loop. Because of very large

212   insertions in the activation loop and in the C-terminal domain in some kinases, the C-terminal

213   domain was aligned only within some families. For example, the AGC-family Great wall kinase

214   (AGC_MASTL) has a 548 amino acid insertion in the activation loop that caused the entire AGC

215   family to be misaligned in the C-terminal domain with respect to the other families.

Figure 4: Pairwise structural alignments of Aurora A to 271 human protein kinases using FATCAT. The x-
axis represents Aurora A residue numbers; the y-axis displays the number of times each Aurora A
residue position is aligned to other kinases in the pairwise alignments. For a description and location of
the labeled aligned blocks in Aurora A structure refer to Table 2 and Figure 1. The list of 272 aligned
structures is provided in Table S1.

This alignment was manually edited in Jalview[38]. As shown in Figure 5a, coloring the

sequence in "Clustal" format in Jalview greatly aided in adjusting the alignment since it

highlighted both level of sequence conservation and physical characteristics of amino acids in

each column (hydrophobicity, positive charge, negative charge, etc.). In addition, a file with the

secondary structure element boundaries of the kinases of known structure was used to highlight

conserved alpha helices and beta sheet strands. Secondary structure predictions performed

with our own secondary structure prediction program based on a deep convolutional neural

network and visualized in Jalview were added (Figure 5b). The alignment was improved in

cycles of several steps:

Figure 5: Snapshot of section of N-terminal region of the multiple sequence alignment of human protein kinases displayed using **a)** Clustal coloring scheme in Jalview which colors the residues by their chemical nature if they are conserved or similar to each other; **b)** secondary structure features of proteins. The secondary structure information for proteins with known structures was obtained from PDB files. For the proteins where a crystal structure is not available, secondary structure predictions were performed with our program SecNet (unpublished). The experimentally known and predicted regions are shown in darker and lighter shades of the same color, respectively, with beta sheets in red and helices in blue. Loops are displayed in gray and the residues which were not resolved in crystal structures are colored in black.

1) The main conserved regions identified in Figure 4 were edited to form aligned blocks in the

MSA, according to the structure alignments of AURKA N and C terminal domains to the

kinases with known structures. These blocks were made as long as possible while

respecting the structure alignments.

2) Regions between the aligned blocks that contain variable sequence lengths were left

unaligned, and edited in Jalview so that there were left-justified and denoted in lower-case

letters. They were separated from the aligned blocks by two empty columns on each end.

These blocks were made as short as possible while respecting the structure alignments.

3) The sequences of kinases in each kinase group without known structure were aligned to

their closest homologues of known structure, such that all sequence motifs were identified in

each kinase. This was greatly facilitated by sorting the alignment by family (AGC, CAMK,

etc.) in Jalview, which was possible because of the naming convention described above

indicating group membership (AGC_AKT1, etc.) and sorting the alignment by these

identifiers.

4) Some kinases without close homologues of known structure required pairwise sequence

alignment to other individual kinases with PSI-BLAST and/or hhpred in order to identify

regions of the kinase domain that occur after very long insertions or that are highly divergent

from other kinases. A small number of kinases were not closely related to any mammalian

kinase of known structure, and instead required a sequence alignment to a yeast or insect

kinase of known structure, which could then be structurally aligned to AURKA. Examples

include: OTHER_TP53RK (33% identity with yeast BUD32, PDB:4WWA[39]); the second

kinase domain of OTHER_EIF2AK4 (47% identity to yeast GCN2, PDB:1ZY5[40]);

OTHER_PINK1 (47% identity with *Triboleum castaneum* PINK1, PDB:5OAT[41]); and

OTHER_PAN3 (49% identical to *Drosophila* PAN3, PDB:4BWP[42]).

5) The G-helix required structure alignments of a region consisting of the G helix itself and the

HF~HG and HG~HH loops, since the position of the G helix is highly variable within the C-

terminal domain of kinases (Figure 1). A small number of kinases do not contain a G-helix

as indicated by their experimental structures or those of close homologues (OTHER_BUB1,

OTHER_HASPIN) and in a few cases by secondary structure prediction and alignment to

more distant homologues (OTHER_RPS6KC1, OTHER_SCYL1, OTHER_SCYL3,

OTHER_PEAK1, OTHER_PEAK3, OTHER_PKDCC, OTHER_PRAG1, OTHER_TP53RK,

OTHER_PXK, OTHER_RNASEL, OTHER_RPS6KL1, OTHER_POMK, OTHER_STK31,

STE_EIF2AK1).

Our final alignment of 497 kinase sequences consists of 2229 columns including all the

residue positions and gaps. It has 17 aligned blocks and 16 unaligned low similarity regions.

The 17 aligned blocks consist of 8 segments from the N-terminal domain (B1N, B1C, B2, B3,

HC, B4, B5, HD) and 9 segments for the C-terminal domain (HE, CL, ALN, ALC, HF, FL, HG,

HH, HI). Each block is named for the main element of secondary structure that it contains,

although they each contain adjacent loop regions. We name the lower-case insertion regions by

the pieces of structure that they connect separated by a tilde, e.g. B2~B3 is the segment

between B2 and B3. B1N and B1C are the N and C terminal segments of beta strand 1; some

kinases such as STE_MAP3K8 have an insertion in the strand that produces an extrusion of the

chain. CL is the catalytic loop that contains the HRD motif. This segment also contains B6 and

B7, which are short beta strands in most kinases. It also contains a 5-residue insertion in one

kinase, OTHER_POMK, which is evident in the structure of mouse POMK (PDB:5GZ8[43]). This

segment remains part of the aligned region CL, since it occurs in only one kinase. The activation

loop is divided into N and C terminal aligned blocks, named ALN and ALC, of 21 and 17

residues respectively. These are long enough to include the common phosphorylation sites of

tyrosine kinases (at positions 13 and 14 from the activation loop N terminus) and

serine/threonine kinases (at position 12 from the activation loop C-terminus)[44]. Many kinases

have long insertions between ALN and ALC. ALN contains the B8 strand prior to the DFG motif

at the N-terminus of the activation loop. ALC contains the short EF helix, which extends several

residues beyond the APE motif at the C-terminus of the activation loop. FL ("F loop") is a short

beta turn motif with consensus sequence PP[FY] between HF and HG that is conserved in most

kinases both in sequence and in structure.

Table 2: Summary of multiple sequence alignment

#	Aligned Region Name	Hanks Name	Common Length of Aligned Region	Aurora A Res. of Aligned Region	Column Number of Aligned Region[a]	Unaligned Region Name	Median Length of Unaligned Region	Aurora A Res. [Pos] of Unaligned Region[b]	Max Length	Kinase with longest insertion
1	B1N	I	4	133-136	1-4	B1N~B1C	0	[136-137]	12	IRAK4
2	B1C	I	7	137-143	21-27	B1~B2	0	[143-144]	12	PAN3
3	B2	I	9	144-152	44-52	B2~B3	2	153-154	36	PINK1
4	B3	II	11	155-165	93-103	B3~HC	7	166-174	32	KKCC1/2
5	HC	III	14	175-188	140-153	HC~B4	0	[188-189]	22	PINK1
6	B4	IV	16	189-204	180-195	B4~B5	0	[204-205]	220	E2AK3
7	B5	IV	10	205-215	420-430	B5~HD	0	[215-216]	10	STK40
8	HD	V	9	216-224	445-453	HD~HE	2	225-226	480	KS6C1
9	HE	VIa	25	227-250	939-962	HE~B6	0	[250-251]	41	MKNK1
10	CL	VIb	16	251-266	1008-1028	CL~ALN	0	[266-267]	298	SRPK2
11	ALN	VII	21	267-286	1331-1351	ALN~ALC	1	[286-287]	548	GWL
12	ALC	VIII	17	287-303	1904-1920	ALC~HF	0	[303-304]	28	PI3R4
13	HF	IX	22	304-325	1953-1975	HF~FL	0	[325-326]	13	BMPR2
14	FL	IX	6	326-331	1993-1998	FL~HG	0	[331-332]	46	MP2K2
15	HG	X	13	332-344	2049-2061	HG~HH	11	345-353	109	CDC7
16	HH	XI	20	354-373	2175-2194	HH~HI	0	[373-374]	10	IKKA
17	HI	XI	10	374-383	2209-2218					

[a] The length of aligned regions by Aurora residue and alignment column numbers differ at CL because
OTHER_STK16 has a five residue insertion; and HF because 30 CAMK kinases have a one residue
insertion in HF.
[b]Aurora A residues in square brackets indicate that Aurora has an unaligned region of length zero.

A list of aligned and unaligned blocks is provided in Table 2, including their positions in

Aurora A, the column numbers in the MSA, and their length(s). Their positions within the SE

alignments are notated in Figure 4. Five of the 16 unaligned regions have nonzero median

lengths. The longest of these are the B3~HC and HG~HH regions. We left the B3~HC region

unaligned because in most of the 62 AGC and 96 CAMK kinases, the region is in the form of a

helix called the B helix while in the other families it takes on a coil form. The HG~HH unaligned

region is highly divergent in structure because of the variation in the position of the G helix. We

have created sequence logos with the program Weblogo[45] to visualize the conservation of

residues in all the aligned blocks of our MSA (Figure 6). The logos show the well-known

conserved motifs including the HRD motif in the catalytic look and the DFG and APE motifs in

the activation loop, as well as hydrophobic positions in the beta sheet strands and alpha helices.

For instance, positions 6, 7, and 10 in the G helix contain predominantly hydrophobic amino

acids.

Figure 6: Sequence logos displaying conservation of residues created for all the aligned blocks of the
MSA using the webserver Weblogo (http://weblogo.threeplusone.com/). The y-axis represents the
column numbers from the MSA. For a description of aligned blocks see Table 2. CL is split into two parts
for better visualization.

Structural validation of the MSA

As described above, the MSA was guided by pairwise alignments of kinase structures to a single kinase (AURKA). However, to determine the accuracy of our MSA we have compared it with the sequence alignments derived from pairwise structure alignments of 272 human kinases in the PDB. Because changes in conformation of the activation loop or movement of the C-helix may affect the corresponding alignment, we used structures that carry an inward disposition of the C-helix as often as possible, as determined by our recent classification of the active and inactive states of kinases[46]. The resulting structure alignments from FATCAT were read by SE to print the unaligned blocks in lower case letters. A residue pair in any two kinases is assumed to be correctly aligned in the MSA if it is also aligned in the pairwise structural alignment of the two kinases. To do the validation we have computed three quantities as described in Methods: 1) True positive rate (TPR): the number of residue pairs which are aligned in the MSA and also in pairwise structure alignments divided by the total number of residue pairs aligned in the structure alignments; 2) Positive predictive value (PPV): the number of residue pairs which are aligned in the MSA and also in the structure alignments divided by the total number of residue pairs aligned in the MSA; 3) The Jaccard similarity index[47]: the sum of the number of residue pairs that are identically aligned in both the MSA and the structure alignments divided by the total number of unique residue pairs aligned in the MSA or the structure alignments or both (counting each pair only once). The Jaccard index shows the overlap between MSA and structural alignments, and penalizes both under- and over-prediction of aligned residues in the MSA.

The average values and distributions of these quantities are presented in Table 3 for our MSA. We have also compared the quality of our alignment with the previously published alignments by Manning et al.[5], Möbitz[16], Kwon et al.[23], a hidden Markov model (HMM) derived from our MSA, and the initial ClustalOmega alignment. The average TPR for our MSA is 0.97, which is significantly better than the Möbitz (0.88), Kwon (0.90), and Manning (0.80) MSAs. The

initial ClustalOmega alignment had a TPR of only 0.74.. Similarly, our MSA also has the highest

PPV (0.97) and the highest value for the Jaccard index of 0.94. The HMM derived from our MSA

has TPR of 0.92, which is less accurate than the MSA itself but more accurate than the Möbitz,

Manning, and Kwon alignments. It may therefore be of use in aligning kinases from other

species to our MSA of human kinases.

Table 3. Average values for TPR, PPV and Jaccard similarity

	Kinases in alignment	Kinases in validation	TPR	PPV	Jaccard similarity	Gap regions per kinase
FoxChase*	497	272	0.969	0.967	0.938	18.7
HMM**	497	272	0.922	0.919	0.857	-
Manning	491	272	0.799	0.809	0.694	44.6
Möbitz	489	271	0.882	0.880	0.799	19.9
Kwon	494	271	0.901	0.908	0.835	139.6
Clustal	497	272	0.739	0.754	0.612	22.3

* Our alignment
** HMM derived from our alignment
TPR = true positive rate
PPV = positive predictive value

We calculated the 'gappiness' of each element, which we identify as the average

number of gap regions in each sequence in the MSA. These are also contained in Table 3. Our

alignment is the least gappy, with average number of gap regions of 19. While we have 16

unaligned regions, three of the aligned regions contain short gapped regions internally to

accommodate one or more kinases with an unusual insertion in the aligned region. The Möbitz

and ClustalOmega alignments are slightly more gappy than ours, while the Manning and Kwon

alignment are substantially gappier with 45 and 140 gap regions per sequence respectively.

Figure 7: Histogram displaying the discrepancy between benchmark pairwise structural alignments of kinases and **a)** FoxChase and **b)** Manning's multiple sequence alignment. The y-axis represents the number of pairwise alignments when a residue pair is aligned in the benchmark but is not aligned in the MSA. The values are plotted with Aurora A residue numbering as a reference on the x-axis.

We examined the positions where the structure alignments and our MSA are discrepant (Figure 7a). In some cases, the discrepancies occur in positions within the aligned blocks that are immediately adjacent to the unaligned segments. This is because the positions of the aligned blocks are not ideal for every single kinase but are a form of consensus position. For example, 30% of the alignments have a discrepancy near the B4/B5 boundary. In the remaining cases, they represent structural shifts of elements of secondary structure in some kinases relative to the kinase domain. This is particularly true of the C-helix, which shows a discrepancy between the sequence alignment and the structural alignment at a rate of about 5%.

Examination of cases where the C-helix is misaligned indicates that the sequence alignments

are probably correct, and the differences with the structure alignment are because of shifts in

the position or orientation of the C-helix relative to the rest of the N-terminal domain. For

example, the C-helix is misaligned in 50% of the structure alignments of MAP2K1 and MAP2K2,

despite the fact that the conserved C-helix glutamic acid that forms a salt bridge with a lysine

residue in the B3 strand in most kinases is correctly aligned in our MSA. Another example is the

C-helix of STK38 (PDB: 6BXI[48]), which is rotated by 100°, and thus the conserved glutamic acid

residue and the entire C-helix do not align with the homologous residues in closely related

kinases. An active form structure of STK38 is not available, but is predicted to have the C-helix

salt bridge to the lysine residue in the β3 strand[48].

We performed the same analysis of the alignment errors for the Manning et al. alignment

(Figure 7b). While there are more misalignments throughout the kinase domain in the Manning

alignment than in ours, the G-helix region is misaligned in about 50% of the pairwise alignments

in comparison to the structure alignments. This is likely due to lower sequence conservation in

this region and the absence of readily identified sequence motif,like the HRD and DFG motifs.

The segment before the C-helix is also poorly aligned compared to structure alignments

because of the presence of the B helix in AGC and CAMK kinases and coil region elsewhere, as

noted above.

Some kinases have very long insertions that have confounded some previous multiple

sequence alignments. AGC_MASTL (Great wall kinase) contains a 548-residue long insertion

within the activation loop (residues 188-735) and is the longest kinase domain sequence at 801

amino acids. In the Manning and Möbitz alignments, the kinase domain is defined as residues

35-310, and both miss the entirety of the C-terminal domain that follows the activation loop.

OTHER_RPS6KC1 (Ribosomal protein S6 kinase delta-1) has a 480-residue long insertion

(residues 419-898) in the HD~HE unaligned region. In the Manning alignment, the kinase

domain begins on residue 822; in our alignment it begins at residue 337. OTHER_CDC7 has a

98-residue long insertion (residues 441-539) between HG and HH, while the kinase domain is

defined as residues 58-472 in the Kwon alignment, thus missing the HH and HI helices.

Figure 8: Unrooted phylogenetic tree of human protein kinases created by using our MSA. The nine
kinase groups are displayed in different colors with OTHER in gray. 13 of the 14 OTHER kinases assigned
to CAMK group by our analysis are shown as a dendrogram on the right side (PLK5 is a truncated kinase
domain and is not shown in the tree). The figure was created using the iTOL webserver and can be
accessed at https://itol.embl.de/shared/foxchase

Phylogenetic trees and group membership

In their paper on the human kinome, Manning et. al provided a phylogenetic tree and

classified the human protein kinases into nine groups extending the early Hanks[49] and Hunter[50]

schemes. These groups consisted of AGC, CAMK, CK1, CMGC, NEK, RGC, STE, TKL and

TYR. A total of 83 protein kinases were placed in an OTHER category because no significant

relationship to any of the nine groups was recognized. However, this classification was done

with only a limited amount of structural information, and as shown above, the Manning multiple

sequence alignment was only 80% correct. We have revisited the phylogeny and classification

of kinases to see if we can assign groups to some of the OTHER kinases by benefiting from our

structure guided multiple sequence alignment.

Because using aligned blocks tends to result in better phylogenies[51], we built the tree

with the 17 conserved blocks from the alignment (Figure 8). The tree was created using the

neighbor joining algorithm in the software MegaX[52] and was visualized using the webserver

iTOL[53]. The resulting tree clusters most of the kinases into the previously recognized nine

groups. Uniprot includes the NEK kinases as a separate group, which also appears in our tree.

In our tree, the RGC kinases form a small sub-branch within the TKL group, but we have

retained the designation.

Among the kinases which are assigned to a group by Uniprot, we observed that eight

STE kinases, MAP3K7 (TAK1 in Manning), MAP3K9 (MLK1), MAP3K10 (MLK2), MAP3K11

(MLK3), MAP3K12 (ZPK), MAP3K13 (LZK), MAP3K20 (ZAK), and MAP3K21 (MLK4) form a

tight cluster in the TKL group branch. These were also in the TKL branch of the Manning tree.

Similarly, six sequences consisting of the second domains of RPS6KA1, RPS6KA2, RPS6KA3,

RPS6KA4, RPS6KA5, and RPS6KA6 which were previously annotated to be in the AGC group

by Uniprot also cluster in CAMKs (as they are in the Manning tree). The first domains of these

kinases are AGC members.

Because of their remote homology, most of the OTHER kinases branch out into separate

clades in between the major groups. Due to the smaller size of these clades and relatively low

similarity between the members we have not classified them as individual kinase groups. In our

tree there are seven OTHER kinases in Uniprot that are correctly assigned to groups by

Manning. Four kinases -- STK32A, STK32B, STK32C, and RSKR, form a branch within the

AGC group. These kinases were also classified as AGCs by Manning et. al. (labeled YANK1,

YANK2, YANK3, SgK494 respectively). Three kinases, CSNK2A1, CSNK2A2, and CSNK2A3,

form a tight cluster within the CMGC group. These are listed as OTHER by Uniprot and while

CSNK2A1 and CSNK2A2 were designated CMGCs by Manning. One kinase, PBK (also called

TOPK) is assigned to STE in Uniprot, but to OTHER by Manning. In our tree, it sits just outside

the TKL group but we would still classify it as OTHER.

Figure 9: HMM scores of CAMK and AGC kinases. HMMs were built on our alignment of AGC and CAMK
kinases according to the original Manning assignments. Several subfamilies of kinases have CAMK scores
in the same range as other bona fide CAMK kinases (magenta). These scores are much higher than their
scores with the AGC HMM (blue). These kinases subfamilies consist of AURK (AURKA, AURKB, AURKC),
CAMKK (CAMKK1 and CAMKK2), PLK (PLK1, PLK2, PLK3, PLK4), ULK (ULK1, ULK2, ULK3), and STK36.
458

We have identified a set of 14 kinases from the OTHER category in both Manning and

Uniprot that can be appropriately assigned to the CAMK group. One branch is formed in the

middle of the CAMK group by nine kinases consisting of AURKA, AURKB, AURKC, CAMKK1,

CAMKK2, PLK1, PLK2, PLK3, and PLK4 (Figure 8). PLK5 is a pseudokinase consisting only of

the C-terminal domain, although mouse PLK5 is full-length[54]. We have included it in the CAMK

group because of its close sequence relationship with the other PLKs. The second branch,

which is on the periphery of the CAMK group, comprises four kinases, ULK1, ULK2, ULK3, and

STK36 (labeled Fused in Manning). ULK4 is more distantly related, and we have left it in the

OTHER group.

To confirm the changes in group membership, we created HMM profiles for each of the

nine groups of kinases, omitting the kinases potentially misclassified by Uniprot (MAPKs in STE

and second domain of RPS6KA1-6 in AGC). We then scanned each of the 497 kinase

sequences against the nine group HMM profiles. A cutoff score of 200 was consistent with the

assignments by Uniprot except for the changes described above. The novel assignments are

the 14 kinases that we can confidently move from OTHER to CAMK described above. The HMM

scores clearly assign them to CAMK rather than AGC or OTHER, since the new CAMKs cluster

with the other CAMK kinases (Figure 9), well above the diagonal line where CAMK=AGC.

Using the new assignments in the AGC, CAMK, CMGC and TKL groups, we created

new HMM profiles to identify if any OTHER sequences could also be reassigned. However, in

the second iteration none of the OTHER category kinases exhibited high scores against any

group HMM profile.

DISCUSSION

Typical kinase domains possess a well-defined fold that is similar across the available

structures. Many kinases are not well studied – the so-called dark kinome[55], and it is possible to

generate hypotheses about their sequence-structure-function relationships by examining their

phylogenetic and structural relationships to well studied kinases. To enable this and for many

other purposes, we have created a structurally-validated, multiple sequence alignment of 497

human protein kinase domains - fully annotated with gene, protein, group name, UniProt

accession identifiers, and residue numbers. The MSA contains 17 aligned blocks of conserved

elements of typical kinase domain and 16 intermittent low similarity regions with varying length

insertions. Our aim was to create a parsimonious alignment without unnecessary gaps; the

residues in low similarity regions were therefore not aligned but formatted as left-justified blocks

of lower case letters to distinguish them from aligned regions. It is reminiscent of the first

multiple sequence alignment of kinases produced by Hanks et al. in 1988[49], which the authors

also described as "parsimonious."

Alignments are only useful if they are accurate. While several multiple sequence

alignments of human kinases have been published and are available online[5,16,19,23], none of

them has been structurally validated. We assessed the accuracy of our alignment with a set of

all-against-all pairwise structural alignments of 272 human kinases, and calculated true positive

rates (TPR), positive predictive values (PPV), and the Jaccard similarity index. In a large-scale

benchmark of sequence alignment methods, we referred to them as f_D (for developer) and f_M

(for modeler) for TPR and PPV respectively[56]. Yona and Levitt subsequently used these values

(renamed Q_D and Q_M) to benchmark profile-profile sequence alignments, and added Q_C or

$Q_{Combined}$[57], which is simply the Jaccard index.. The Jaccard index penalizes both overprediction

and underprediction in our sequence alignments. We used all three values for our alignment

(0.97, 0.97, and 0.94 respectively) to demonstrate that our alignment is more accurate than the

others available.

The errors in our MSA of kinases are mostly limited to the boundaries of conserved

blocks where the variability of residue positions across kinases make their unambiguous

placement in aligned blocks difficult. However, structure alignments do not always align every

homologous pair of residues in two proteins. This occurs when residues are disordered in one of

the structures or where there is significant conformational change. In a small number of kinases

the only structure available has a significantly rotated C-helix. Structure alignment therefore

sometimes does not align the homologous residues of the C helix in two kinases. The same is

true for the G-helix in some kinases, which may be positioned in different locations within the C-

terminal domain, but retains a homologous sequence and structure, and thus is aligned

differently in our MSA than in the structure alignments.

With a more accurate MSA in hand, we have revisited the phylogenetic tree of human

kinases. The widely used phylogenetic tree of Manning et al is based on an alignment that has

TPR and PPV values of only 0.80 and 0.81. Uniprot also provides a classification of kinases into

the same groups as Manning et al (with the addition of NEK kinases). The Uniprot annotations

that differ from Manning et al. are all incorrect. Fourteen kinases are placed in the wrong groups

by Uniprot, and another seven are placed in OTHER but can easily be placed within one of the

defined groups, in agreement with the Manning annotations. One kinase (PBK or TOPK) is

labeled as STE by Uniprot but we agree with Manning's annotation of OTHER.

Of greater interest, our phylogenetic tree and hidden Markov models for each group can

help us confidently assign 14 kinases to the CAMK group; these are listed as OTHER by

Manning et al. and Uniprot: AURKA, AURKB, AURKC, CAMKK1, CAMKK2, PLK1, PLK2, PLK3,

PLK4, PLK5, ULK1, ULK2, ULK3, and STK36. Manning et al. put the CAMKK, ULK, PLK

kinases, and STK36 (called Fused in the kinome poster) adjacent to the CAMK group. From our

hidden Markov model of CAMK kinases and a phylogenetic tree based on our MSA, these

kinases fit clearly into the CAMK group. Experimental data confirm that these assignments are

correct. CAMKK1 and CAMKK2 (Ca^{2+}/calmodulin-dependent kinase kinase 1 and 2) both

phosphorylate Ca^{2+}/calmodulin-dependent kinases and bind calmodulin[58-60]. There is also direct

evidence of calmodulin binding to PLK1[61] and of a calmodulin homologue, calcium-and-integrin-

binding protein (CIB), to both PLK2 (Snk in the kinome poster) and PLK3 (Fnk)[62]. There is

evidence suggesting the role of calcium ions in the activity of ULK1 through its phosphorylation

by the CAMK kinase AMPK[63].

Manning et al. put the Aurora kinases on the same branch but not in the AGC group,

which is closely related to the CAMK group. Both groups possess a B helix that is not present in

the other families. The HMM and the phylogenetic tree show that the three Aurora kinases fit

better into the CAMK group than the AGC group. In an earlier study with colleagues, we have

shown experimental evidence that Aurora A binds calmodulin[64], supporting its assignment to the

CAMK group. Calmodulin also binds to Aurora B kinase (AURKB), preventing its degradation

via the E3 ligase FBXL2 subunit[65].

Our MSA provides the benefit of a common numbering scheme using the columns of the

alignment facilitating comparison across all the kinase sequences. The identification of

equivalent residue positions helps in generalizing experimental data from one kinase to another.

For example, substrate specificity is highly correlated with the amino acid type at a small

number of positions within the substrate binding site[66]. Creixell found that specificity could also

be modulated by more remote sites[19], based on a multiple sequence alignment of kinases

derived with Clustal W. It is likely that our more accurate alignment would facilitate this analysis

and produce more reliable predictions. Other areas where an accurate alignment and phylogeny

may be useful are in predicting inhibitor specificity[67], regulatory mechanisms through protein-

protein interactions, and computational protein design of kinases with altered functionality[68].

Our alignment is included as supplemental data and on our website, and will be updated

as new structures are determined. We hope that it will be of use in kinase biology and

therapeutic development.

METHODS

Identification of human protein kinases

The list of human typical and atypical protein kinases was obtained from Uniprot website

(https://www.uniprot.org/docs/pkinfam). To identify any unlisted kinases we searched human

sequences in Uniprot with PSI-BLAST using the typical and atypical protein kinase on the

Uniprot page as queries. PSI-BLAST was also used to identify structures of human kinases or

their closest homologues in the PDB. The structures of atypical kinases (or homologues thereof)

were examined structurally using FATCAT pairwise alignments and structural superposition by

CEalign in Pymol to Aurora A. Four of the atypical kinase families are visibly related to typical

kinases, but contain significant fold differences. The other two families are not homologous to

typical kinases.

A total of 497 typical kinase domain sequences from 484 kinase genes (13 genes have

two kinase domains each) were used to create the MSA. These sequences were initially divided

into 9 phylogenetic groups as per the Uniprot nomenclature: AGC, CAMK, CK1, CMGC, NEK,

RGC, TKL, TYR, and STE, and a tenth group of diverse kinases designated OTHER. Gene

names   were   retrieved   from   the   Human   Gene   Nomenclature   Committee   website

(http://genenames.org)[34]. Each kinase sequence was labeled by group name underscore HGNC

gene name, for example AGC_PRKACA for KAPCA_HUMAN. The 13 kinases that have two

kinase domains in the polypeptide chain were labeled with an underscore, for example

TYR_JAK1_1 and TYR_JAK1_2. The boundaries were determined with PSI-BLAST of the full-

length Uniprot sequence against the PDB.

Multiple Sequence Alignment

The kinase sequences (except some with very long insertions like GWL) were aligned

using ClustalOmega[69] to prepare an initial alignment. This was manually edited using Jalview to

make sure that conserved motifs such as the DFG and HRD motif were aligned across most of

the sequences[70]. The sequences with low sequence similarity to most of the other kinases and

those containing long insertions were difficult to align. To improve the accuracy of the alignment,

pairwise structural alignment of the kinases which have a crystal structure was performed using

with the structure of Aurora kinase (3E5A_A) and the program FATCAT[71]. However, for the

kinases where a structure was not known, alignment of the closest known structures to Aurora A

were used to edit the alignment with Jalview[72]. In a few cases, the most closely related

structures were not human or even mammalian kinases. In these cases, the non-human kinase

was structurally aligned to Aurora A and the target kinase was added to the MSA by transitive

alignment. For a few distant kinases where a closely related structure or sequence was not

known, HHPred was used to identify similarity to another kinase[73].

Structural validation of the MSA

The MSA was structurally validated using a set of pairwise structural alignments as a

benchmark. The benchmark consists of all vs all pairwise structural alignments of 272 kinases

with known structures in the PDB. Two kinases were excluded: the BUB1B kinase domain in the

cryo-EM structure of the anaphase promoting complex (PDB: 5KHU, chain Q[74]) is completely

disordered; the structure of PBK (TOPK_HUMAN, PDB: 5J0A[75]) contains two monomers with

half of the N-terminal domains of each chain swapped with the other monomer, making it

difficult to align to other kinase domain structures. The structure-based sequence alignments

were created using FATCAT in rigid mode and optimized using SE[36]. For kinases with multiple

structures known the structure for validation was selected based on their conformational states

using our previously published nomenclature[46]. The active state BLAminus conformation was

preferred over others, followed by different kinds of DFGin inactive states - ABAminus, BLAplus,

BLBminus, BLBplus, BLBtrans and DFGout-BBAminus.

A residue pair between two kinases in the MSA was considered to be aligned if it was

also aligned in the benchmark pairwise structural alignment of the same kinases. Using this

information the accuracy of the MSA was assessed by computing three quantities TPR, PPV

and the Jaccard similarity index. For each pair of sequences, we first calculate the number of

aligned residue pairs that are present in both the sequence alignment and the structure

alignment ($N_{correct}$). The TPR is the ratio of $N_{correct}$ and the number of residue pairs aligned in the

structure alignment (N_{struct}). For computation of the TPR, residue pairs in the structure alignment

are skipped if either or both residues are contained in the unaligned (lower-case) blocks of the

sequence alignment. This takes care of situations that occur when two kinases have identical

length segments between two of our aligned blocks; the structure alignment program would

align them but they would be indicated as unaligned in our sequence alignment. The alignment

of Kwon et al. also includes unaligned regions in lowercase and is treated in the same way.

PPV is the ratio of $N_{correct}$ and the number of aligned residue pairs in the sequence

alignment (N_{seq}). For the PPV, residue pairs in the aligned blocks of the sequence alignment are

skipped if one or both residues are aligned to gap characters in the structure alignment. This is

usually either because the residues are disordered (no coordinates) in one of the structures or

because there is a significant conformational change of a loop and the residues are aligned to

gaps. The Jaccard similarity index is the ratio of $N_{correct}$ and the number of unique aligned pairs

in either the structure alignment or the sequence alignment (counting each only once). For the

Jaccard index, all the pairs skipped in TPR and PPV are also skipped. A script for calculating

these values is available on https://github.com/DunbrackLab/Kinases. The aligned residues

from the pairwise structure alignments are available on https://zenodo.org/record/3445533

(DOI 10.5281/zenodo.3445533).

We also compared our MSA accuracy with the previously published alignments. These

alignments did not contain residue ranges in the Uniprot sequences, and used different

nomenclature for the protein names. To identify a correspondence between the sequences in

previously published alignments and our MSA, we performed PSI-BLAST searches of each of

their sequences against Uniprot and renamed them according to our scheme

(groupname_genename).

Phylogenetic tree

The phylogenetic tree of human protein kinases was created from an MSA obtained by

deleting the unaligned regions from our MSA. A distance matrix using the p-distance in the

program MegaX[52] was created which was used to create a phylogenetic tree with the neighbor-

joining algorithm. The tree was saved in Newick format and uploaded to iTOL webserver for

visualization where each clade was colored according to its kinase group

(https://itol.embl.de/shared/foxchase).

HMM profiles of kinase groups

We used the hmmbuild program of the HMMER3 package[76] to create HMM profiles for

each of the nine kinase groups. The input MSA for each group was extracted from the main

MSA and any empty columns were deleted. Before creating the HMMs the following sequences

were removed from the STE group - MAP3K7, MAP3K9, MAP3K10, MAP3K11, MAP3K12,

MAP3K13, MAP3K20, MAP3K21 and PBK, since PSI-BLAST searches indicated that Uniprot

had misclassified these. For the same reason, RPS6KA1_2, RPS6KA2_2, RPS6KA3_2,

RPS6KA4_2, RPS6KA5_2 and RPS6KA6_2 were removed from the AGC group. All 497 kinase

sequences were run against the nine HMM profiles using the program hmmsearch[76]. The

scores of each sequence against nine HMMs were sorted and the group with highest score

against each sequence was identified.

Acknowledgments

662 This work was supported by NIH grant R35 GM122517 to RLD. We thank Maxim Shapovalov

663 for the program *SecNet* for prediction of protein secondary structure.

Supplemental Files

TableS1.txt – data on each of 497 kinases in the alignment in text format

TableS1.xlsx -- data on each of 497 kinases in the alignment in Excel format

Human-PK-v1.aln – our alignment in Clustal format (version 1)

Human-PK-alignment-v1.mfa – our alignment in FASTA format, suitable for visualization in

Jalview

Human-PK-sequences-v1.fasta – the domain sequences of 497 kinases (unaligned)

References

1.	Adams JA. Kinetic and catalytic mechanisms of protein kinases. *Chem Rev* **101**:2271-2290 (2001).

2.	Blume-Jensen P, Hunter T. Oncogenic kinase signalling. *Nature* **411**:355-365 (2001).

3.	Lahiry P, Torkamani A, Schork NJ, Hegele RA. Kinase mutations in human disease: interpreting genotype-phenotype relationships. *Nat Rev Genet* **11**:60-74 (2010). 10.1038/nrg2707.

4.	Zhang J, Yang PL, Gray NS. Targeting cancer with small molecule kinase inhibitors. *Nat Rev Cancer* **9**:28-39 (2009). 10.1038/nrc2559.

5.	Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. The protein kinase complement of the human genome. *Science* **298**:1912-1934 (2002).

6.	Ten Eyck LF, Taylor SS, Kornev AP. Conserved spatial patterns across the protein kinase family. *Biochim Biophys Acta* **1784**:238-243 (2008). 10.1016/j.bbapap.2007.11.002.

7.	Middelbeek J, Clark K, Venselaar H, Huynen MA, Van Leeuwen FN. The alpha-kinase family: an exceptional branch on the protein kinase tree. *Cell Mol Life Sci* **67**:875-890 (2010).

8.	Stefely JA, Reidenbach AG, Ulbrich A, Oruganty K, Floyd BJ, Jochem A, Saunders JM, Johnson IE, Minogue CE, Wrobel RL, Barber GE, Lee D, Li S, Kannan N, Coon JJ, Bingman CA, Pagliarini DJ. Mitochondrial ADCK3 employs an atypical protein kinase-like fold to enable coenzyme Q biosynthesis. *Mol Cell* **57**:83-94 (2015). 10.1016/j.molcel.2014.11.002.

9.	LaRonde-LeBlanc N, Wlodawer A. A family portrait of the RIO kinases. *J Biol Chem* **280**:37297-37300 (2005).

10.	Xiao J, Tagliabracci VS, Wen J, Kim S-A, Dixon JE. Crystal structure of the Golgi casein kinase. *Proceedings of the National Academy of Sciences* **110**:10574-10579 (2013).

11.	Blackford AN, Jackson SP. ATM, ATR, and DNA-PK: the trinity at the heart of the DNA damage response. *Mol Cell* **66**:801-817 (2017).

12.	Steussy CN, Popov KM, Bowker-Kinley MM, Sloan RB, Harris RA, Hamilton JA. Structure of pyruvate dehydrogenase kinase: Novel folding pattern for a serine protein kinase. *J Biol Chem* **276**:37443-37450 (2001).

13.	Ogden TH, Rosenberg MS. Multiple sequence alignment accuracy and phylogenetic inference. *Syst Biol* **55**:314-328 (2006).

14. Jiang Y, Oron TR, Clark WT, Bankapur AR, D'Andrea D, Lepore R, Funk CS, Kahanda I, Verspoor
KM, Ben-Hur A. An expanded evaluation of protein function prediction methods shows an
improvement in accuracy. *Genome biology* **17**:184 (2016).
15. Chartier M, Chenard T, Barker J, Najmanovich R. Kinome Render: a stand-alone and web-
accessible tool to annotate the human protein kinome tree. *PeerJ* **1**:e126 (2013).
10.7717/peerj.126.
16. Möbitz H. The ABC of protein kinase conformations. *Biochimica et Biophysica Acta (BBA)-
Proteins and Proteomics* **1854**:1555-1566 (2015).
17. Brooijmans N, Chang YW, Mobilio D, Denny RA, Humblet C. An enriched structural kinase
database to enable kinome-wide structure-based analyses and drug discovery. *Protein Sci*
19:763-774 (2010).
18. McSkimming DI, Rasheed K, Kannan N. Classifying kinase conformations using a machine
learning approach. *BMC Bioinformatics* **18**:86 (2017).
19. Creixell P, Palmeri A, Miller CJ, Lou HJ, Santini CC, Nielsen M, Turk BE, Linding R. Unmasking
determinants of specificity in the human kinome. *Cell* **163**:187-201 (2015).
20. Rahman R, Ung PM-U, Schlessinger A. KinaMetrix: a web resource to investigate kinase
conformations and inhibitor space. *Nucleic Acids Res* **47**:D361-D366 (2018).
21. van Linden OP, Kooistra AJ, Leurs R, de Esch IJ, de Graaf C. KLIFS: A knowledge-based structural
database to navigate kinase-ligand interaction space. *J Med Chem* (2013). 10.1021/jm400378w.
22. Hartmann S, Vision TJ. Using ESTs for phylogenomics: can one accurately infer a phylogenetic
tree from a gappy alignment? *BMC Evol Biol* **8**:95 (2008).
23. Kwon A, Scott S, Taujale R, Yeung W, Kochut KJ, Eyers PA, Kannan N. Tracing the origin and
evolution of pseudokinases across the tree of life. *Sci Signal* **12**:eaav3810 (2019).
24. Magrane M, UniProt Consortium. UniProt Knowledgebase: a hub of integrated protein data.
Database **2011**:bar009 (2011).
25. Hildebrand A, Remmert M, Biegert A, Soding J. Fast and accurate automatic structure prediction
with HHpred. *Proteins* **77 Suppl 9**:128-132 (2009). 10.1002/prot.22499.
26. Yamaguchi H, Matsushita M, Nairn AC, Kuriyan J. Crystal structure of the atypical protein kinase
domain of a TRP channel with phosphotransferase activity. *Mol Cell* **7**:1047-1057 (2001).
27. Zhao Y, Zhang X, Chen Y, Lu S, Peng Y, Wang X, Guo C, Zhou A, Zhang J, Luo Y, Shen Q, Ding J,
Meng L, Zhang J. Crystal Structures of PI3Kalpha Complexed with PI103 and Its Derivatives: New
Directions for Inhibitors Design. *ACS Med Chem Lett* **5**:138-142 (2014). 10.1021/ml400378e.
28. Ferreira-Cerca S, Kiburu I, Thomson E, LaRonde N, Hurt E. Dominant Rio1 kinase/ATPase catalytic
mutant induces trapping of late pre-40S biogenesis factors in 80S-like ribosomes. *Nucleic Acids
Res* **42**:8635-8647 (2014). 10.1093/nar/gku542.
29. Maurice F, Pérébaskine N, Thore S, Fribourg S. In vitro dimerization of human RIO2 kinase. *RNA
Biology* **in press**:1-10 (2019). 10.1080/15476286.2019.1653679.
30. Tso S-C, Qi X, Gui W-J, Chuang JL, Morlock LK, Wallace AL, Ahmed K, Laxman S, Campeau PM,
Lee BH, Hutson SM, Tu BP, Williams NS, Tambar UK, Wynn RM, Chuang DT. Structure-based
design and mechanisms of allosteric inhibitors for mitochondrial branched-chain α-ketoacid
dehydrogenase kinase. *Proceedings of the National Academy of Sciences* **110**:9728-9733 (2013).
31. Kato M, Li J, Chuang JL, Chuang DT. Distinct structural mechanisms for inhibition of pyruvate
dehydrogenase kinase isoforms by AZD7545, dichloroacetate, and radicicol. *Structure* **15**:992-
1004 (2007). 10.1016/j.str.2007.07.001.
32. Cheng H, Schaeffer RD, Liao Y, Kinch LN, Pei J, Shi S, Kim B-H, Grishin NV. ECOD: an evolutionary
classification of protein domains. *PLOS Comput Biol* **10**:e1003926 (2014).
33. Tsutakawa SE, Jingami H, Morikawa K. Recognition of a TG mismatch: the crystal structure of
very short patch repair endonuclease in complex with a DNA duplex. *Cell* **99**:615-623 (1999).

34. Braschi B, Denny P, Gray K, Jones T, Seal R, Tweedie S, Yates B, Bruford E. Genenames. org: the
HGNC and VGNC resources in 2019. *Nucleic Acids Res* **47**:D786-D792 (2018).
35. Ye Y, Godzik A. FATCAT: a web server for flexible structure comparison and structure similarity
searching. *Nucleic Acids Res* **32**:W582-585 (2004).
36. Tai C-H, Vincent JJ, Kim C, Lee B. SE: an algorithm for deriving sequence alignment from a pair of
superimposed structures. *BMC Bioinformatics* **10**:S4 (2009).
37. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M,
Söding J. Fast, scalable generation of high-quality protein multiple sequence alignments using
Clustal Omega. *Mol Syst Biol* **7**:539 (2011).
38. Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2—a multiple
sequence alignment editor and analysis workbench. *Bioinformatics* **25**:1189-1191 (2009).
39. Zhang W, Collinet B, Graille M, Daugeron MC, Lazar N, Libri D, Durand D, van Tilbeurgh H. Crystal
structures of the Gon7/Pcc1 and Bud32/Cgi121 complexes provide a model for the complete
yeast KEOPS complex. *Nucleic Acids Res* **43**:3358-3372 (2015). 10.1093/nar/gkv155.
40. Padyana AK, Qiu H, Roll-Mecak A, Hinnebusch AG, Burley SK. Structural basis for autoinhibition
and mutational activation of eukaryotic initiation factor 2alpha protein kinase GCN2. *J Biol Chem*
280:29289-29299 (2005). 10.1074/jbc.M504096200.
41. Kumar A, Tamjar J, Waddell AD, Woodroof HI, Raimi OG, Shaw AM, Peggie M, Muqit MM, van
Aalten DM. Structure of PINK1 and mechanisms of Parkinson's disease-associated mutations.
eLife **6** (2017). 10.7554/eLife.29985.
42. Christie M, Boland A, Huntzinger E, Weichenrieder O, Izaurralde E. Structure of the PAN3
pseudokinase reveals the basis for interactions with the PAN2 deadenylase and the GW182
proteins. *Mol Cell* **51**:360-373 (2013). 10.1016/j.molcel.2013.07.011.
43. Nagae M, Mishra SK, Neyazaki M, Oi R, Ikeda A, Matsugaki N, Akashi S, Manya H, Mizuno M, Yagi
H, Kato K, Senda T, Endo T, Nogi T, Yamaguchi Y. 3D structural analysis of protein O-mannosyl
kinase, POMK, a causative gene product of dystroglycanopathy. *Genes Cells* **22**:348-359 (2017).
10.1111/gtc.12480.
44. Xu Q, Malecka KL, Fink L, Jordan EJ, Duffy E, Kolander S, Peterson JR, Dunbrack RL, Jr. Identifying
three-dimensional structures of autophosphorylation complexes in crystals of protein kinases.
Sci Signal **8**:rs13 (2015). 10.1126/scisignal.aaa6711.
45. Crooks GE, Hon G, Chandonia J-M, Brenner SE. WebLogo: a sequence logo generator. *Genome
Res* **14**:1188-1190 (2004).
46. Modi V, Dunbrack RL. Defining a new nomenclature for the structures of active and inactive
kinases. *Proceedings of the National Academy of Sciences* **116**:6818-6827 (2019).
47. Jaccard P. La distribution de la flore dans la zone alpine. *Revue générale des sciences pures et
appliqué* **15 Dec.**:961-967 (1907).
48. Xiong S, Lorenzen K, Couzens AL, Templeton CM, Rajendran D, Mao DY, Juang Y-C, Chiovitti D,
Kurinov I, Guettler S. Structural basis for auto-inhibition of the NDR1 kinase domain by an
atypically long activation segment. *Structure* **26**:1101-1115. e1106 (2018).
49. Hanks SK, Quinn AM, Hunter T. The protein kinase family: conserved features and deduced
phylogeny of the catalytic domains. *Science* **241**:42-52 (1988).
50. Hunter T. [1] Protein kinase classification. Methods Enzymol. Volume 200: Elsevier; 1991. p 3-37.
51. Talavera G, Castresana J. Improvement of phylogenies after removing divergent and
ambiguously aligned blocks from protein sequence alignments. *Syst Biol* **56**:564-577 (2007).
52. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis
across computing platforms. *Mol Biol Evol* **35**:1547-1549 (2018).
53. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation
of phylogenetic and other trees. *Nucleic Acids Res* **44**:W242-W245 (2016).

54.   de Cárcer G, Manning G, Malumbres M. From Plk1 to Plk5: functional evolution of polo-like
799   kinases. *Cell cycle* **10**:2255-2262 (2011).

800   55.   Needham EJ, Parker BL, Burykin T, James DE, Humphrey SJ. Illuminating the dark
801   phosphoproteome. *Sci Signal* **12**:eaau8645 (2019).

802   56.   Sauder JM, Arthur JW, Dunbrack RL, Jr. Large-scale comparison of protein sequence alignment
803   algorithms with structure alignments. *Proteins: Structure, Function and Genetics* **40**:6-22 (2000).

804   57.   Yona G, Levitt M. Within the twilight zone: a sensitive profile-profile comparison tool based on
805   information theory. *J Mol Biol* **315**:1257-1275 (2002).

806   58.   Tokumitsu H, Wayman GA, Muramatsu M, Soderling TR. Calcium/calmodulin-dependent protein
807   kinase kinase: identification of regulatory domains. *Biochemistry* **36**:12823-12827 (1997).

808   59.   Osawa M, Tokumitsu H, Swindells MB, Kurihara H, Orita M, Shibanuma T, Furuya T, Ikura M. A
809   novel target recognition revealed by calmodulin in complex with Ca 2+-calmodulin-dependent
810   kinase kinase. *Nat Struct Mol Biol* **6**:819 (1999).

811   60.   Tokumitsu H, Muramatsu M-a, Ikura M, Kobayashi R. Regulatory mechanism of
812   Ca2+/calmodulin-dependent protein kinase kinase. *J Biol Chem* **275**:20090-20095 (2000).

813   61.   Dai G, Qian Y, Chen J, Meng F-L, Pan F-Y, Shen W-G, Zhang S-Z, Xue B, Li C-J. Calmodulin
814   activation of polo-like kinase 1 is required during mitotic entry. *Biochem Cell Biol* **91**:287-294
815   (2013).

816   62.   Kauselmann G, Weiler M, Wulff P, Jessberger S, Konietzko U, Scafidi J, Staubli U, Bereiter-Hahn J,
817   Strebhardt K, Kuhl D. The polo-like protein kinases Fnk and Snk associate with a Ca2+-and
818   integrin-binding protein and are regulated dynamically with synaptic plasticity. *The EMBO*
819   *journal* **18**:5528-5539 (1999).

820   63.   Shang L, Chen S, Du F, Li S, Zhao L, Wang X. Nutrient starvation elicits an acute autophagic
821   response mediated by Ulk1 dephosphorylation and its subsequent dissociation from AMPK.
822   *Proceedings of the National Academy of Sciences* **108**:4788-4793 (2011).

823   64.   Plotnikova OV, Pugacheva EN, Dunbrack RL, Golemis EA. Rapid calcium-dependent activation of
824   Aurora-A kinase. *Nature communications* **1**:64 (2010). 10.1038/ncomms1061

825   ncomms1061 [pii].

826   65.   Mallampalli RK, Glasser JR, Coon TA, Chen BB. Calmodulin protects Aurora B on the midbody to
827   regulate the fidelity of cytokinesis. *Cell Cycle* **12**:663-673 (2013).

828   66.   Brinkworth RI, Breinl RA, Kobe B. Structural basis and prediction of substrate specificity in
829   protein serine/threonine kinases. *Proceedings of the National Academy of Sciences* **100**:74-79
830   (2003).

831   67.   Anastassiadis T, Deacon SW, Devarajan K, Ma H, Peterson JR. Comprehensive assay of kinase
832   catalytic activity reveals features of kinase inhibitor selectivity. *Nat Biotechnol* **29**:1039 (2011).

833   68.   Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J, Shimizu E, Tsien JZ, Schultz
834   PG, Rose MD. A chemical switch for inhibitor-sensitive alleles of any protein kinase. *Nature*
835   **407**:395 (2000).

836   69.   Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F,
837   Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. Clustal W and Clustal X
838   version 2.0. *Bioinformatics* **23**:2947-2948 (2007). btm404 [pii]

839   10.1093/bioinformatics/btm404.

840   70.   Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2--a multiple
841   sequence alignment editor and analysis workbench. *Bioinformatics* **25**:1189-1191 (2009).
842   10.1093/bioinformatics/btp033.

71. Ye Y, Godzik A. Flexible structure alignment by chaining aligned fragment pairs allowing twists.
Bioinformatics **19 Suppl 2**:246-255 (2003).
72. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and
PSI-BLAST: a new generation of database programs. *Nucleic Acids Res* **25**:3389-3402 (1997).
73. Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology detection and
structure prediction. *Nucleic Acids Res* **33**:W244-248 (2005). 33/suppl_2/W244 [pii]

10.1093/nar/gki408.
74. Yamaguchi M, VanderLinden R, Weissmann F, Qiao R, Dube P, Brown NG, Haselbach D, Zhang W,
Sidhu SS, Peters JM, Stark H, Schulman BA. Cryo-EM of Mitotic Checkpoint Complex-Bound
APC/C Reveals Reciprocal and Conformational Regulation of Ubiquitin Ligation. *Mol Cell* **63**:593-
607 (2016). 10.1016/j.molcel.2016.07.003.
75. Dong C, Tang X, Xie Y, Zou Q, Yang X, Zhou H. The crystal structure of an inactive dimer of PDZ-
binding kinase. *Biochem Biophys Res Commun* **476**:586-593 (2016). 10.1016/j.bbrc.2016.05.166.
76. Eddy SR. A new generation of homology search tools based on probabilistic inference. Genome
Informatics 2009: Genome Informatics Series Vol 23: World Scientific; 2009. p 205-211.