Mating-type specific divergent *RPL22* paralogs are essential, differentially regulated, and control aspects of sexual reproduction in *Cryptococcus neoformans*

Giuseppe Ianiri[1], Yufeng "Francis" Fang[1], Tim A. Dahlmann[2], Guilhem Janbon[3], Ulrich Kück[2], and Joseph Heitman[1*]

[1]Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC, 27710, USA. [2]Allgemeine und Molekulare Botanik, Ruhr-Universität Bochum, 44780, Bochum, Germany. [3]Unité Biologie des ARN des Pathogènes Fongiques, Département de Mycologie, Institut Pasteur, Paris, 75015, France.

[*]Address correspondence to Joseph Heitman, Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, United States of America. Phone +1 (919) 684-2824. Fax (919) 684-2790. Email heitm001@duke.edu

Abstract

The *MAT* locus of *Cryptococcus neoformans* has a bipolar organization characterized by an unusually large structure, spanning over 100 kb. *MAT* genes have been characterized by functional genetics as being involved in sexual reproduction and virulence. However, classical gene replacement failed to achieve mutants for five *MAT* genes (*RPL22, RPO41, MYO2, PRT1, RPL39*), indicating that they are likely essential. In the present study, targeted gene replacement was performed in a diploid strain for both the α and **a** alleles of the ribosomal genes *RPL22* and *RPL39*. Mendelian analysis of the progeny confirmed that both *RPL22* and *RPL39* are essential for viability. Ectopic integration of the *RPL22* paralog of opposite *MAT* identity in the heterozygous *RPL22***a**/*rpl22*αΔ or *RPL22*α/*rpl22***a**Δ mutant strains failed to complement their essential phenotype. Evidence suggests that this is due to differential expression of the *RPL22* genes, and an RNAi-dependent mechanism that contributes to control *RPL22***a** expression. Furthermore, via CRISPR/Cas9 technology the *RPL22* alleles were exchanged in haploid *MAT*α and *MAT***a** strains of *C. neoformans*. These *RPL22* exchange strains displayed morphological and genetic defects during bilateral mating. The *C. neoformans MAT***a** strain in which the *RPL22***a** gene was exchanged with *RPL22*α displayed sensitivity to hydroxyurea and high temperature, and had a morphological phenotype consistent with a defective cell cycle. These results contribute to elucidate functions of *C. neoformans* essential mating type genes that may constitute a type of imprinting system to promote inheritance of nuclei of both mating types.

Introduction

Infectious diseases cause significant morbidity and mortality worldwide in both developed and developing countries. Fungal infections are common in humans and impact the majority of the world's population, but are often underestimated (BROWN *et al.* 2012). The *Cryptococcus* species complex includes basidiomycetous fungal pathogens that can cause lung infections and life-threatening meningoencephalitis in both normal and immunocompromised patients, accounting for approximately 1 million annual infections globally and almost 200,000 annual mortalities (RAJASINGHAM *et al.* 2017). The available drugs to treat *Cryptococcus* infections are amphotericin B, 5-flucytosine, and azoles. These drugs are characterized by limited spectrum, toxicity, unavailability in some countries, and emergence of drug resistance (BROWN *et al.* 2012).

In fungi, the mechanisms that govern sexual reproduction are controlled by specialized regions called mating type (*MAT*) loci. The genomic organization of the *MAT* loci can differ among fungi. The tetrapolar mating type includes two *MAT* loci, the P/R locus encoding pheromones and pheromone receptor genes defining sexual identity and mate recognition, and the HD locus encoding homeodomain transcription factors that govern post-mating developmental processes. In the tetrapolar system the P/R and HD loci are located on different chromosomes and segregate independently during meiosis, hence generating recombinant *MAT* systems. Conversely, in the bipolar mating system both the P/R and the HD loci are linked on the same chromosome, and recombination in this region is suppressed. A variant of the bipolar system is a mating conformation called pseudobipolar, in which the P/R and the HD loci are located on the same chromosome but unlinked, thus allowing (limited) recombination (COELHO *et al.* 2017).

C. neoformans has a well-defined sexual cycle that is controlled by a bipolar *MAT* system that is derived from an ancestral tetrapolar state. The *C. neoformans MAT* locus evolved in a unique configuration as it spans over 100 kb and contains more than 20 genes that control cell identity, sexual reproduction, infectious spore production, and virulence. The two opposite *C. neoformans MAT*α and *MAT***a** alleles include divergent sets of the same genes that evolved by extensive remodeling from common ancestral DNA regions. Both the *MAT*α and *MAT***a** allele contain five predicted essential genes: *RPO41, PRT1, MYO2, RPL39,* and *RPL22* (LENGELER *et al.* 2002; FRASER *et al.* 2004). Rpo41 is a mitochondrial RNA polymerase that transcribes

mitochondrial genes and also synthesizes RNA primers for mitochondrial DNA replication

(SANCHEZ-SANDOVAL *et al.* 2015). Prt1 is a subunit of the eukaryotic translation initiation factor

3 (eIF3) that plays a critical role in translation (BEZNOSKOVA *et al.* 2015). Myo2 is a myosin

heavy chain type V that is involved in actin-based transport of cargos and is essential in *S.*

cerevisiae (JOHNSTON *et al.* 1991). Rpl39 and Rpl22 are ribosomal proteins.

This study focused on the *MAT* ribosomal proteins, demonstrating that both *RPL39* and

RPL22 α and **a** alleles are essential in *C. neoformans.* Because Rpl22 in yeast and vertebrates has

been found to play specialized functions and extra-ribosomal roles (GABUNILAS AND CHANFREAU

2016; KIM AND STRICH 2016; ZHANG *et al.* 2017; ABRHAMOVA *et al.* 2018), we aimed to

characterize the functions of the *C. neoformans RPL22*α and *RPL22***a** paralogs. We found that

ectopic integration of an *RPL22* paralog failed to complement the essential phenotype due to the

mutation of the *RPL22* of opposite mating type. We found differential expression of the *C.*

*neoformans RPL22*α and *RPL22***a** paralogs during mating, and discovered an RNAi-mediated

mechanism that contributes to control *RPL22***a** expression. Next, using the CRISPR/Cas9

technology *RPL22* alleles were exchanged in haploid *MAT*α and *MAT***a** strains of *C. neoformans*

and this resulted in morphological and genetic defects during bilateral mating. In summary, these

studies reveal a novel role for diverged essential ribosomal proteins in controlling fungal sexual

reproduction.

Materials and methods

Strains and culture conditions. The strains utilized in the present study are listed in table S1. Heterozygous mutants were generated in the diploid *C. neoformans* strain AI187 (*MATα*/*MAT***a** *ade2*/*ADE2 ura5*/*URA5*) according to a previously developed strategy (IANIRI AND IDNURM 2015). *C. neoformans* strain AI187 was generated through the fusion of strains JF99 (*MAT***a** *ura5*) and M001 (*MATα ade2*) (IDNURM 2010). For transformation of haploid *C. neoformans* strains, we employed H99α and KN99**a** (NIELSEN *et al.* 2003). All of the strains were maintained on yeast extract-peptone dextrose (YPD) agar medium.

Molecular manipulation of *C. neoformans*. For the generation of heterozygous mutants, 1.5 kb regions flanking the genes of interest were amplified by PCR and fused with the *NAT* marker through *in vivo* recombination in *S. cerevisiae* as previously described (IANIRI AND IDNURM 2015). Split-marker gene replacement alleles were amplified from *S. cerevisiae* transformants with primers JOHE43263/ALID1229 and JOHE43264/ALID1230 in combination with ai37 and JOHE44324, respectively. The amplicons were precipitated onto gold beads and transformed into *C. neoformans* with a Bio-Rad particle delivery system (TOFFALETTI *et al.* 1993); W7 hydrochloride was added to YPD + 1 M Sorbitol to increase the efficiency of homologous recombination (ARRAS AND FRASER 2016). Transformants were selected on YPD + NAT and screened for homologous recombination events by PCR with primers external to the replaced regions in combination with primers specific for the *NAT* marker, and with gene-specific internal primers. The primers used are listed in Table S2.

For complementation experiments, a region of 2397 bp including the *RPL22*α gene with its promoter and terminator was amplified by PCR from *C. neoformans* H99α genome and cloned in pCR™2.1 according to the manufacturers' instructions. Similarly, a region of 2648 bp including the *RPL22***a** gene with its promoter and terminator was amplified by PCR from *C. neoformans* KN99**a** genome and cloned in pCR™2.1 according to manufacturers' instruction. Plasmids were recovered from *E. coli* TOP10 and sequenced to identify error-free clones (Table S2). Sequence confirmed plasmids were digested with SpeI-XhoI and SpeI-NotI to obtain regions including the *RPL22*α and *RPL22***a** genes, respectively. These fragments were purified and subcloned within the pSDMA57 plasmid for safe haven complementation (ARRAS *et al.* 2015) digested with the same enzymes. The recombinant plasmids were recovered from *E. coli*

TOP10, linearized with AscI, PacI, or BaeI, and transformed through biolistic in *C. neoformans*

heterozygous mutants GI56 (*RPL22α/rpl22*a∆) and GI81 (*RPL22*a/*rpl22*α∆) as described above.

C. neoformans transformants were selected on YPD + neomycin G418, and subjected to DNA

extraction and PCR analyses to identify transformants having the correct insert within the safe

haven region (ARRAS *et al.* 2015).

The recently-developed TRACE technology (Transient CRISPR/Cas9 coupled with

electroporation) (FAN AND LIN 2018) was utilized for the generation of the 5′∆ *RPL22*a strain

GI228 and the *RPL22* exchange alleles. For 5′∆ *RPL22*a, a homology directed repair (HDR)

template consisting of 1.5 kb sections flanking the region upstream *RPL22*a targeted by sRNA

was fused with the *NAT* marker through *in vivo* recombination in *S. cerevisiae* as described

above. For the generation of the *RPL22* exchange alleles, HDR templates were generating by

fusing ~1.0 kb fragments flanking the *RPL22* genes with the ORF of the opposite *RPL22*

paralog. For the generation of a chimeric c*RPL22*α (c = chimeric), the N-terminal region of

*RPL22*α (from nucleotide 1 to 268) and the C terminal region of *RPL22*a (from nucleotide 253 to

600) were combined together by PCR, fused with ~1.0 kb regions flanking the *RPL22*a gene, and

employed as an HDR template. All HDR templates were assembled using overlap PCR as

described in (DAVIDSON *et al.* 2002).

Specific guide RNAs (gRNA) were designed according to (FANG *et al.* 2017) using

EuPaGDT (http://grna.ctegd.uga.edu/) available on FungiDB (https://fungidb.org/fungidb/).

Complete gRNAs were generated by one-step overlap PCR, in which a bridge primer that

comprises the 20 nucleotide gRNA guide sequences was utilized to integrate the U6 promoters

(amplified from *C. deneoformans* XL280 genomic DNA) and the gRNA scaffold [amplified

from the plasmid pYF515 (FANG *et al.* 2017)]. *CAS9* was amplified from pXL1-Cas9 (FAN AND

LIN 2018). Safe Haven 2 (SH2) sequence was obtained from (UPADHYA *et al.* 2017). All PCR-

amplifications were conducted using Phusion High-Fidelity DNA Polymerase (NEB). *C.*

neoformans was transformed with *CAS9*, gRNAs, and HDR templates in through electroporation

following the previously reported protocol (FAN AND LIN 2018). Transformants were screened

for homologous recombination events by PCR as previously indicated.

Genetic analyses and scanning electron microscopy of reproductive structures. The

heterozygous strains generated were grown on Murashige-Skoog (MS) medium to induce

meiosis and sporulation. Haploid *C. neoformans* mutants were crossed with *C. neoformans* WT

strains of compatible mating type (H99 *MATα* and KN99**a** *MAT***a**) on MS medium and monitored

for the formation of sexual structures. Spores were micromanipulated and allowed to germinate

onto YPD agar for 3 to 4 days at 30°C, and then tested for the segregation of the genetic markers.

For the heterozygous strains the markers were nourseothricin resistance (NAT^R) or sensitivity

(NAT^S), *ura5/URA5, ade2/ADE2, MATα*/*MAT***a**, plus neomycin G418 resistance (NEO^R) or

sensitivity (NEO^S) for the complementing strains. For the haploid strains they were either NAT^R-

NAT^S or NEO^R-NEO^S, and *MATα* or *MAT***a**. The analyses were performed by spotting 2 µl of

cell suspensions onto YPD + nourseothricin (100 µg/ml) or neomycin (100 µg/ml), YNB +

adenine (20 mg/L) or YNB + uracil (40 mg/L). The mating type was scored by crossing haploid

progeny to strains KN99**a** and H99 on MS media supplemented with adenine and uracil, and by

evaluating the formation of sexual structures by microscopy (IDNURM 2010; IANIRI AND IDNURM

2015). For NAT^R colonies the mating type was confirmed by PCR with primers JOHE39201-

JOHE39202 (*MAT***a**) and JOHE39203-JOHE39204 (*MATα*).

Scanning electron microscopy (SEM) was performed at the North Carolina State

University Center for Electron Microscopy, Raleigh, NC, USA. Samples were prepared for SEM

as previously described (FU AND HEITMAN 2017). Briefly, a small MS agar block containing

hyphae was excised and fixed in 0.1 M sodium cacodylate buffer, pH = 6.8, containing 3%

glutaraldehyde at 4°C for several weeks. Before imaging, the agar block was rinsed with cold 0.1

M sodium cacodylate buffer, pH = 6.8 three times and post-fixed in 2% osmium tetroxide in cold

0.1 M cacodylate buffer, pH = 6.8 for 2.5 hours at 4°C. Then the block was critical-point dried

with liquid CO_2 and sputter coated with 50 Å of gold/palladium with a Hummer 6.2 sputter

coater (Anatech). The samples were viewed at 15KV with a JSM 5900LV scanning electron

microscope (JEOL) and captured with a Digital Scan Generator (JEOL) image acquisition

system.

RT-qPCR analysis during mating and statistical analysis. For RT-qPCR analysis of *RPL22*

expression during mating, strains were grown overnight in liquid YPD, and cellular density was

adjusted to 1 x 10^9 CFU/mL. Equal amounts of each cellular suspension of strains to be analyzed

were mixed, and 5 spots of 300 µl were placed onto one plate of MS agar per day of incubation.

Control conditions were the single strains on YPD agar (2 spots of 300 µl per day of incubation).

Every 24 h, cells were scraped off the MS plate, washed once with sterile water, lyophilized, and

kept at -80°C until RNA extraction. RNA extraction was performed with the standard TRIzol

protocol following the manufacturers' instructions (RIO *et al.* 2010). Extracted RNA was treated

with DNase and purified with an RNA clean and concentration kit (Zymo Research). Then, 3 µg

of purified RNA were converted into cDNA via the Affinity Script QPCR cDNA synthesis kit

(Agilent Technologies). cDNA synthesized without the RT/RNase block enzyme mixture was

utilized as a control for genomic DNA contamination. Approximately 500 pg of cDNA were

utilized to measure the relative expression level of target genes through quantitative real-time

PCR (RT-qPCR) using the Brilliant III ultra-fast SYBR green QPCR mix (Agilent Technologies)

in an Applied Biosystems 7500 Real-Time PCR System. A control without template RNA was

included for each target. Technical triplicates and biological triplicates were performed for each

sample. Gene expression levels were normalized using the endogenous reference gene *GDP1*

and determined using the comparative ΔΔCt method.

To determine whether the relative gene expression levels between strains of the same

mating reaction in the same day of incubation (for example, *RPL22*α and *RPL22***a** expression in

WT H99α x KN99**a** cross after 48 h of incubation) exhibited statistically significant differences

($p<0.05$, $p<0.01$, $p<0.001$), the unpaired student's t-test with Welch's correction was applied. To

compare the results of different strains in different mating reactions, ordinary one-way ANOVA

with Tukey's multiple comparison test was applied. Because in these comparisons we were

interested in monitoring the changes in gene expression following genetic manipulation, only

statistically significant differences ($p<0.05$, $p<0.01$, $p<0.001$) were displayed for the expression

levels of the same gene on the same day of incubation in separate mating reactions (for example,

*RPL22***a** expression in WT H99α x KN99**a** cross compared to *RPL22***a** expression in *rdp1*Δ x

*rdp1*Δ bilateral cross after 48 h of incubation). Statistical analyses were performed using the

software PRISM8 (GraphPad, https://www.graphpad.com/scientific-software/prism/).

RNA structure modeling. RNA structure modeling was conducted with RNAfold (LORENZ *et*

al. 2011) with default settings.

sRNA data processing. Small RNA (sRNA) sequencing libraries from *C. neoformans* WT H99

× KN99**a** cross and *rdp1*Δ bilateral cross are as described in (WANG *et al.* 2010). The adapters

sequences (5-prime: GTTCAGAGTTCTACAGTCCGACGATC; 3-prime:

TCGTATGCCGTCTTCTGCTTGT) were removed by using cutadapt v1.9 (MARTIN 2011) and

trimmed reads were mapped with bowtie v1.2.2 (LANGMEAD *et al.* 2009) against the *MAT***a**

(AF542528.2) and *MATα* (AF542529.2) loci from *C. neoformans* strains 125.91 and H99,

respectively (LENGELER *et al.* 2000). Mapping was performed by allowing a single nucleotide

mismatch and up to five alignments within both mating type loci, and the H99α genome.

Furthermore, reads showing a single perfect match were considered in order to identify their

genetic origin. Read counts were calculated with SAMtools' depth function and by using custom

made Perl scripts (LI *et al.* 2009; DAHLMANN AND KÜCK 2015). The read counts were

normalized against tRNA mapping reads (tRNA read counts per 100,000 reads). The

normalization factors for the WT mating and the *rdp1Δ* mating were calculated with 1.386 and

0.728, respectively.

Chemical genetic screen and phenotypic analysis. Phenotypic analysis was performed for all

the strains listed in table S1 with the standard 10-fold serial dilution method. Tested conditions

and stresses included: temperatures of 4°C, 25°C, 30°C, 37°C, 38°C, 39°C; antifungal drugs,

such as amphotericin B (AmB, 1.5 µg/mL), 5-fluorocytosine (5-FC, 100 µg/mL), fluconazole

(FLC, 20 µg/mL), FK506 (1 µg/mL), rapamycin (1 µg/mL); cell wall and plasma membrane

stressors, such as YPD and YP supplemented with NaCl (1.5 M and 1 M, respectively) and

sorbitol (2M and 1.5 M, respectively), caffeine (10 mM), calcofluor white (4 mg/mL), Congo

Red (0.8%); genotoxic, oxidative, nitrosative and other stress-inducing agents, such as ethidium

bromide (10 µg/mL), sodium nitrite (NaNO$_2$, 1.5 mM), UV (150 µJ x cm^2), hydrogen peroxide

(H$_2$O$_2$, 3 mM), cycloeximide (0.15 µg/mL), dithiothreitol (DTT, 15 mM), hydroxyurea (125

mM), tunicamycin (0.7 µg/mL), benomyl (2.5 µg/mL), cadmium sulphate (CdSO$_4$, 30 µM).

Unless indicated, plates were incubated at 30°C for 3 to 6 days and photographed.

Data Availability Statement. Strains and plasmids are available upon request. The authors

affirm that all data necessary for confirming the conclusions of the article are present within the

article, figures, and tables.

Results

The *MAT* ribosomal genes *RPL22* and *RPL39* are essential. Both the α and **a** alleles of the five predicted *MAT* essential genes (*RPL39, RPL22, MYO2, RPO41, PRT1*) were identified in the genomes of *C. neoformans* H99α and KN99**a**, and were subjected to targeted mutagenesis in the *C. neoformans* diploid strain AI187 according to the strategy reported by Ianiri and Idnurm with minor modifications (IANIRI AND IDNURM 2015). Briefly, cassettes for targeted gene replacement were generated by *in vivo* recombination in *S. cerevisiae*, and amplified by PCR to perform targeted mutagenesis via split-marker coupled with the use of the non-homologous end joining (NHEJ) inhibitor W7 hydrochloride (FU *et al.* 2006; ARRAS AND FRASER 2016). These modifications were critical to increase the rate of homologous recombination, and allowed the generation of heterozygous mutants for both the *MAT*α and *MAT*a alleles of the *RPL39*, *RPL22*, and *MYO2* genes. Despite these modifications, heterozygous mutants for *RPO41* and *PRT1* were not obtained. This study focuses on the characterization of the *RPL39* and *RPL22* genes encoding ribosomal proteins.

Heterozygous mutants were confirmed by PCR analyses, and then transferred onto MS medium supplemented with adenine and uracil and allowed to undergo meiosis, sporulation, and basidiospore production. Spores were micromanipulated on YPD agar and subjected to phenotypic analysis to assess the segregation of the four available markers (*URA5, ADE2, MAT, NAT*). Because the predicted essential genes were deleted by insertion of the *NAT* marker, the absence of NAT[R] progeny indicates an essential gene function. The other 3 markers were tested to exclude defects in meiosis: while the *URA5/ura5* and *ADE2/ade2* loci were expected to segregate independently, the segregation of the *MAT* region was expected to be linked to the mutated alleles, with progeny being only *MAT*a when derived from *MAT*α heterozygous deletion mutants, and only *MAT*α when derived from *MAT*a heterozygous deletion mutants.

Heterozygous mutants for the *MAT* ribosomal proteins Rpl39 and Rpl22 produced basidiospores that displayed a rate of germination ranging from 36% to 41% (Table 1). Mendelian analysis of the progeny confirmed that both the *MAT*a and *MAT*α alleles of Rpl39 and Rpl22 encode an essential function (Table 1). Figure 1 shows an example of the genetic analysis performed on progeny derived from strains GI233 (*RPL39*a/*rpl39*αΔ) and GI56 (*RPL22*α/*rpl22*aΔ). As expected, in all cases the progeny inherited only one *MAT* allele, which is the opposite of the mutated gene. One exception is that one NAT[R] progeny was obtained from

sporulation of the heterozygous *RPL22α/rpl22a*Δ; further PCR analysis revealed that this strain

has the mutated *rpl22a*Δ*NAT* and an extra copy of the *RPL22α* gene, suggesting that its *NAT*

resistance is likely due to aneuploidy of chromosome 5 (1n + 1) where the *MAT* locus resides.

The *C. neoformans* RPL22 paralogs are highly similar. The *RPL22***a** and *RPL22α* genes share

83% identity at the DNA level, and the encoded proteins differ in 5 amino acids that are located

in the N-terminal region (Fig. S1 A, B). With the exception of intron 1 that is 132 bp for *RPL22α*

and 116 bp for *RPL22***a**, both of the *C. neoformans RPL22* genes contain 4 exons and 3 introns of

identical length. Intron 1 shares ~75% nucleotide identity, intron 2 shares ~75% nucleotide

identity, and intron 3 shares ~60% nucleotide identity (Fig. 2A). *In silico* analysis of intron

features revealed a canonical NG|GTNNGT motif at the donor sites for both *RPL22α* and

*RPL22***a**, and both canonical and non-canonical acceptor motifs CAG|G/C for *RPL22α* and

YAG|Y/G for *RPL22***a**. Four branch sites were predicted for each *RPL22* gene, with the

canonical motif CTRAY being more represented (Fig. 2B). Note that the vertical bar represents

the exon–intron junction, and Y and R indicate nucleotides with pyrimidine and purine bases,

respectively. The length of the predicted polypyrimidine tracts ranged from 9 to 33 nt, and they

differed only in intron 1 (Fig. 2A).

The *rpl22*a mutant was not complemented by the opposite paralog. We sought to determine

whether the Rpl22 *MAT* proteins play a specialized role in *C. neoformans.* The first approach

was based on the heterologous expression of *RPL22***a** or *RPL22α* in the heterozygous mutants

*RPL22***a**/*rpl22α*Δ (strain GI81) and *RPL22α/rpl22a*Δ (strain GI54). Briefly, the *RPL22* genes

including promoters and terminators were cloned into plasmid pSDMA57 for safe haven

complementation with the *NEO* selectable marker (ARRAS *et al.* 2015); the empty plasmid served

as the control. NAT and NEO double drug resistance coupled with PCR analyses to confirm safe

haven integration and lack of plasmid catenation were the basis to select transformants for

analysis; in some cases, integration at the safe haven was not achieved, and a heterozygous

mutant with ectopic integration of the plasmid was chosen for analysis.

Complementing heterozygous strains were sporulated on MS media, and segregation of

the five markers available (*NEO, NAT, MAT, URA5, ADE2*) was determined in the germinated

progeny. The criteria for the successful complementation of the mutant phenotype were: 1)

presence of both NAT[R] and NEO[R] markers in which the progeny's *MAT* region segregated with

the mutated *rpl22ΔNAT* allele – this indicates that the ectopic integration of the *RPL22-NEO*

allele was able to confer viability in progeny that inherited the essential gene *rpl22* deletion 2)

absence of only NAT[R] progeny, because those that inherit the *rpl22ΔNAT* allele are inviable; 3)

presence of NEO[R] progeny that, together with progeny sensitive to *NAT* and *NEO*, inherited the

MAT allele that contains the non-mutated copy of *RPL22*. Progeny that were both *MAT***a** and

*MAT*α were expected to be NAT[R] and either aneuploid or diploid.

Results from the complementation experiments in the heterozygous mutants are

summarized in Table 2. For the heterozygous *RPL22*α/*rpl22***a**Δ (GI56), ectopic introduction of a

WT copy of *RPL22***a** (strain GI151) was able to restore viability in the progeny, with the

generation of three NAT[R] and NEO[R] progeny that were *MAT***a** (Fig. 3). Several attempts with

both biolistic and electroporation of plasmid pSDMA57 + *RPL22*α in strain GI56 yielded a low

number of transformants, a high percentage of plasmid catenation, and lack of integration at the

safe haven. Nevertheless, in two independent experiments 2 heterozygous *RPL22*α/*rpl22***a**Δ +

*RPL22*α mutants (strains GI102 and GI154) with a single ectopic copy of the plasmid pSDMA57

+ *RPL22*α were isolated. Genetic analysis of basidiospores dissected from strain GI102 revealed

surprising findings, with no NEO[R] progeny and two NAT[R] progeny that had an extra copy of

*MAT*α, which includes the *RPL22*α gene and explains their NAT resistance (Fig. 3). Regarding

transformant GI154, out of 20 spores that germinated, only one was solely NAT[R] and three were

both NAT[R] and NEO[R], and all were both *MAT***a** and *MAT*α. These results indicated that the

*RPL22*α gene did not complement the *rpl22***a** mutation. Finally, sporulation of strain GI104

bearing the empty plasmid pSDMA57 also produced one spore that was both *MAT***a** and *MAT*α

(Fig. 3).

Integration of *RPL22*α at the safe haven (strain GI86) restored viability in progeny

derived from the heterozygous *RPL22***a**/*rpl22*αΔ mutant, with the generation of four NAT[R] and

NEO[R] strains that were *MAT*α. For *RPL22***a**/*rpl22*αΔ + *RPL22***a**, none of the transformants

obtained had the plasmid integrated at the safe haven. Several transformants having a single

ectopic copy of the plasmid were sporulated on MS media, but none was able to form

basidiospores for genetic analysis; a representative strain (GI150) is included in Table 2. These

results indicate that *RPL22***a**/*rpl22*αΔ + *RPL22***a** complementing strains failed to sporulate and

hence we could not assess proper functional complementation through genetic analysis of the

meiotic progeny. Finally, as expected introduction of the empty plasmid in *RPL22***a**/*rpl22*αΔ

(strain GI83) did not restore progeny viability.

***RPL22*a expression is regulated by the RNAi pathway.** Next, we sought to determine the

expression levels of the *RPL22* genes during mating (H99 x KN99**a**) on MS medium in

comparison to vegetative growth on YPD agar. While after 24 h of incubation both *RPL22* genes

were expressed at a similar level, at 48 h *RPL22*α expression drastically decreased and remained

lower than that of *RPL22***a** up to 96 h (Fig. 4A). Expression of *RPL22***a** was also higher than

*RPL22*α during vegetative growth on YPD, reflecting the results obtained during mating (Fig.

S2). These results indicate that the two *RPL22* alleles are differentially expressed, with *RPL22***a**

expression being higher than *RPL22*α during both vegetative growth and mating.

What are the mechanisms that control *RPL22* expression? Our hypothesis was that

inefficiently spliced introns of *RPL22*α could trigger RNA interference (RNAi) through the

SCANR complex with subsequent silencing of the gene (DUMESIC *et al.* 2013). Analysis of

intron retention (IR) and the splicing pattern of *RPL22*α in several conditions revealed that intron

1 of *RPL22*α is subject to IR, whereas there is minimal IR for introns 2 and 3 (Fig. S3). Mapping

small RNA data from an H99α x KN99**a** cross against the *MAT***a** and *MAT*α regions, we found

that no reads mapped against the *RPL22* genes, including intronic regions and intron-exon

junctions. This observation argues against the hypothesis that there is a direct role of RNAi in

governing *RPL22* gene expression.

Interestingly, analysis of the region surrounding the *RPL22* genes revealed that more than

54,989 sRNA reads map to the 2.2 kb region upstream of the *RPL22***a** gene, which includes the

LTR11 and *LTR14* elements, and a candidate long non-coding RNA (lncRNA) predicted based

on BLAST analyses. Several sRNA mapping parameters were tested to evaluate whether the

sRNA reads map to multiple locations in the genome of *C. neoformans*, allowing us to determine

the origin of sRNA reads that were mapped to the 5´ upstream region of *RPL22***a**. We found that

the majority of the reads originated from the 5′ upstream region of the *RPL22***a** gene, while

others originated from regions of the genome distant from *MAT* corresponding to the lncRNAs

CNAG_12037 and CNAG_13142, and the region upstream of the lncRNA CNAG_12435.

Analysis of the region upstream of the *RPL22***a** gene in an *rdp1*Δ *MAT*α x *rdp1*Δ *MAT***a**

bilateral cross revealed a drastic reduction in sRNA reads (Fig. 4B), consistent with a role for

RNAi in governing these sRNA. We then performed RT-qPCR of the *rdp1*Δ x *rdp1*Δ bilateral

cross, and found that *RPL22***a** expression was lower than *RPL22*α expression at 72 h and 96 h of

incubation (Fig. 4C). Compared to a WT H99α x KN99**a** cross, *RPL22***a** expression was overall

lower whereas that of *RPL22*α was higher (Fig. S4A). Increased expression of *RPL22*α in the

*rdp1*Δ mutant bilateral cross corroborates previous findings (WANG *et al.* 2010), although

*RPL22*α expression seems to be indirectly regulated by RNAi because no sRNA map to the

regions surrounding the *RPL22*α gene (Fig S5).

We also performed RT-qPCR during a bilateral cross between mutants for the SCANR

complex component Gwc1, and found strong downregulation [fold change (FC) < 0.5] of both

*RPL22***a** and *RPL22*α (Fig. 4D; Fig. S4B). Lastly, we used the recently-developed CRISPR/Cas9

technology to accurately delete the *MAT* region in *C. neoformans* KN99**a** that is upstream of

*RPL22***a** and that is targeted by the abundant sRNA. The strain generated was named GI288 (5′Δ

*RPL22***a**), and a schematic representation of the strategy is shown in Fig. S6A. RT-qPCR

expression analysis during mating revealed that *RPL22***a** expression remained higher than

*RPL22*α, except at 72 h of incubation (Fig. 4E). As compared to *RPL22* expression during the

WT cross, *RPL22***a** was strongly downregulated in the GI228 (5′Δ *RPL22***a**) x H99α cross at 48 h

and 72 h of incubation, while expression of *RPL22*α remained unchanged (Fig. S6B).

Interestingly, expression of *RPL22***a** during the GI288 (5′Δ *RPL22***a**) x H99α cross mirrors that of

*RPL22***a** during the *rdp1*Δ bilateral cross, in accord with an RNAi-dependent mechanism

regulating *RPL22***a** expression (Fig. S6C). Despite the downregulation of *RPL22***a**, the GI228

mutant strain does not display any morphological defect during vegetative growth, and its ability

to mate with H99 and generate viable progeny was not compromised; as expected, genetic

analysis of progeny derived from the GI228 x H99 cross showed co-segregation of *NAT* with

*MAT***a** (data not shown).

RPL22 exchange allele strains exhibit sexual reproduction defects. We next sought to

determine whether functional complementation of the *RPL22* genes could be achieved by

replacing either of them with the opposite *RPL22* paralog at the native locus within the *MAT*

loci. To this end, we generated exchange alleles of *RPL22*α and *RPL22***a** by means of

CRISPR/Cas9, whose use was critical due to suppressed recombination within *MAT*. Because the

RPL22 genes share a high level of identity (83%) (Fig. S1), specificity of the gene replacement

was achieved by designing two specific guide RNA (gRNA) molecules that determined the sites

for homologous recombination. Transformation was performed by electroporation with the

simultaneous introduction of the three sgRNAs [at the 5′ and 3′ of the *RPL22* gene, and one for

the Safe Haven 2 (SH2)], the homology-directed repair (HDR) *RPL22* gene, and selectable

markers *NAT* or *NEO,* which were introduced in the Safe Haven 2 (SH2) region to avoid

unnecessary ectopic mutations that could interfere with the resulting phenotype. Recipients for

transformations were the most isogenic strains available for *C. neoformans* serotype A, H99

(*MAT*α) and KN99**a** (*MAT***a**) (NIELSEN *et al.* 2003; JANBON *et al.* 2014; FRIEDMAN *et al.* 2018).

In two independent transformation attempts, precise gene replacement of *RPL22*α with

*RPL22***a** at its native location within the *MAT*α locus of H99 and correct integration of *NAT* in

the SH2 were readily obtained. This resulted in the generation of *rpl22*α::*RPL22***a** *SH2*::*NAT*

mutant strains (YFF92) that differ from their parental strain at only the *RPL22* gene. A schematic

representation of the exchange strains generated is shown in Figure S7. Conversely, via the same

strategy (i.e. two gRNA at the 5′ and 3′ of the *RPL22*) an *RPL22***a** exchange allele strain could

not be isolated. Because the *RPL22* genes differ in only 5 amino acids that are located in the N-

terminal region (Fig. S1, S7), to replace *RPL22***a** with the Rpl22α coding gene a different

strategy based on CRISPR/Cas9 was employed. In this approach we generated a chimeric

*RPL22*α (c*RPL22*α) HDR template, which consisted of the N-terminus of *RPL22*α fused with the

C-terminus of *RPL22***a**, and this was introduced into *C. neoformans* strain KN99**a** together with

new gRNAs designed to target the 5′ region of *RPL22***a** (Fig 5A; Fig. S7). Several independent

*rpl22***a**::*RPL22*α^{N}-*RPL22***a**^{C} *SH2*::*NEO* exchange strains were obtained, and one (strain YFF113)

was chosen for further experiments. Allele exchange strains YFF92 and YFF113 were used in

unilateral and bilateral crosses to evaluate both their *MAT*-specificity and the phenotypic

consequence due to the absence of one *RPL22* gene. In the presence of only *RPL22***a** (cross

KN99**a** x YFF92α) or *RPL22*α (cross H99α x YFF113**a**), the strains displayed no altered

morphology and had WT mating ability, spore germination, independent segregation of the

markers, and uniparental inheritance of mitochondria (Fig. 5B; File S2; Table 3). These results

indicate that the absence of one Rpl22 paralog does not affect mating efficiency or meiosis when

the other paralog is present in the native location within *MAT*. Interestingly, RT-qPCR revealed a

higher level of expression of the chimeric c*RPL22*α gene in the KN99**a** background compared to

that of *RPL22***a** in the H99 background at 72 h and 96 h of incubation, hence displaying an

opposite trend compared to the WT cross (Fig. 5D; Fig. S8A).

Because the approach utilized to generate strain YFF113 was successful, we employed
the same gRNA to replace the *RPL22***a** gene with a native copy of the *RPL22*α gene. After
several unsuccessful attempts, we obtained one *rpl22***a**::*RPL22*α strain (YFF116) with the *NEO*
marker integrated ectopically in the genome, and not in the *SH2* locus as planned (Fig. 6A; Fig.
S7). Similar to findings presented above, in the presence of only the *RPL22*α gene (cross H99α x
YFF116**a**) no morphological and genetic defects were observed (Fig. 6C; File S2; Table 3).
Remarkably, during bilateral crosses strains with exchanged *RPL22* genes (YFF92α x YFF116**a**)
exhibited a high percentage of irregular basidia (Fig. 6D). High resolution scanning electron
microscopy revealed that the majority of the basidia had no basidiospores, while others had a
morphology defect (Figure 6E-F; File S3), and a low number had irregular basidiospore chains
collapsed on the basidia (Figure 6G; File S3); the formation of clamp connections was not
affected (Figure 6H; File S3). Basidiospores germinated from cross YFF92α x YFF116**a**
displayed a low germination rate (12%) and irregular segregation of the meiotic markers, with 3
progeny being *MAT***a**-*RPL22*α, no *RPL22***a**-*MAT*α progeny, and 4 progeny being both *MAT*α and
*MAT***a**, hence aneuploid or diploid (Table 3). RT-qPCR analysis during YFF92 x YFF116 mating
revealed low expression (FC < 1) of both *RPL22* genes from 24 h to 96 h (Fig. 6B; Fig. S8B).

Phenotypic analysis of *rpl22* mutant strains. Heterozygous deletion mutants and exchange
strains (Table S1) were tested for altered phenotypic traits (see Materials and Methods for
details). Of the 28 stresses tested, few phenotypic differences among the isolates were observed
on FLC, YP + NaCl, caffeine, 39°C and 4°C (Fig. S8). Remarkably, exchange strain YFF116
(*rpl22***a**::*RPL22*α *NEO*) displayed a marked sensitivity to 39°C and hydroxyurea compared to its
parental strain KN99**a** (Fig. 7A, Fig. S8). Microscopic analysis of strain YFF116 grown on
hydroxyurea revealed severe morphological defects, including larger cells with irregular shapes
and abnormally elongated buds (Fig. 7B). Surprisingly, strain YFF116 also displayed cellular
defects when grown in the absence of stress, although these were less severe and occurred in
fewer cells.

Discussion

A previous study reported that the *MAT* locus of *C. neoformans* contains five genes (*RPL39, RPL22, MYO2, RPO41, PRT1*) that encode proteins required for viability (FRASER *et al.* 2004). The essential nature of these genes was inferred based on the inability to mutate them in a haploid strain of *C. neoformans.* In addition to *C. neoformans*, *Candida albicans* is the only other fungus known to encode essential genes within the *MAT* locus (HULL *et al.* 2000; SRIKANTHA *et al.* 2012). It is likely that one function of *MAT*-essential genes is to constrain recombination and serve as a genetic buffer constraining loss of portions of the *MAT* locus, although they might also serve other functions related to development, sexual reproduction, and virulence.

Here the functions of the α and **a** *MAT* specific alleles encoding the *C. neoformans* ribosomal proteins Rpl22 and Rpl39 were characterized. Given suppressed recombination within the *C. neoformans MAT* loci, mutation of the *RPL22* and *RPL39* genes was challenging. Heterozygous mutants for these genes were successfully generated through the combined use of a biolistic split marker approach and the compound W7 to inhibit non-homologous end joining and enhance homologous recombination (FU *et al.* 2006; ARRAS AND FRASER 2016). Through deletions in the *C. neoformans* **a**/α diploid strain AI187 and Mendelian analysis of recombinant F1 progeny obtained following sexual reproduction and spore dissection, we demonstrated that both the α and **a** paralogs of the *RPL22* and *RPL39* genes are essential (Table 1; Fig. 1). Conversely, in *S. cerevisiae* the *RPL22* and *RPL39* orthologs are not required for viability (STEFFEN *et al.* 2012; KIM AND STRICH 2016), indicating evolutionary divergence of essential ribosomal genes between Ascomycetous and Basidiomycetous yeasts.

The ribosome was thought to be a constant, conserved, uniform protein translation machine. Recent studies have revealed novel and unexpected findings for the ribosome, in particular complex heterogeneity and specialized activity that confers regulatory control in gene expression (WARNER AND MCINTOSH 2009; NARLA AND EBERT 2010; XUE AND BARNA 2012). While in mammals ribosomal proteins are encoded by single genes, in yeasts, plants, and flies, ribosomal proteins are encoded by several genes. A remarkable example is the model yeast *S. cerevisiae*, in which, following a genome duplication event, 59 of the 78 ribosomal proteins are encoded by two retained gene copies and share high sequence similarity, but are in most cases not functionally redundant, and have been found to play specialized functions [(XUE AND BARNA

2012) and references within it]. Specialized ribosomes have been identified also in plants, flies, zebrafish, and mice (KOMILI *et al.* 2007; MCINTOSH AND WARNER 2007; XUE AND BARNA 2012), but it is not known whether they exist in microbial pathogens.

We sought to determine the functions of the Rpl22 *MAT* proteins and whether they play a specialized role in *C. neoformans.* A number of studies have converged to reveal diverse and specialized roles for the Rpl22 ribosomal paralogs. In *S. cerevisiae*, haploid *rpl22*a mutants are cold sensitive, and display reduced invasive growth and a longer doubling time compared to *rpl22*b (STEFFEN *et al.* 2012; KIM AND STRICH 2016). Moreover, *rpl22*a mutations perturb bud site selection and cause random budding, while *rpl22*b mutations do not, and overexpression of *RPL22*B in *rpl22*a mutants fails to restore bud site selection (KOMILI *et al.* 2007). Vertebrates also express Rpl22 paralogs, called Rpl22 and Rpl22-like1 (RPL22-l1). Mice lacking *RPL22* are viable and have specific αβ T-cell developmental defects, likely attributable to compensation by Rpl22-l1 in other tissues (ANDERSON *et al.* 2007). Recent studies in both yeast and mammals suggest an extraribosomal role for Rpl22 paralogs in binding target mRNAs and regulating their expression (GABUNILAS AND CHANFREAU 2016; ZHANG *et al.* 2017; ABRHAMOVA *et al.* 2018).

In *C. neoformans*, we found that *RPL22*α was not able to complement the essential phenotype due to mutations of *RPL22***a**, whereas ectopic introduction of *RPL22***a** in *RPL22***a**/*rpl22*αΔ resulted in sporulation failure. Conversely, viability was restored in progeny derived from heterozygous *RPL22***a**/*rpl22*αΔ + *RPL22*α and *RPL22*α/*rpl22***a**Δ + *RPL22***a** strains (Fig. 3; Table 2). In a parallel approach we ectopically inserted an *RPL22* paralog into a *C. neoformans* haploid strain, and then attempted to mutate the native opposite *MAT* copy. Also in this case deletion mutants could not be recovered, further supporting the observation of a failure of complementation between the two *C. neoformans RPL22* paralogs (data not shown). We further demonstrate that during both mitotic growth and sexual reproduction *RPL22***a** expression is much higher than *RPL22*α (Fig. 4A, Fig. S2). Considering that the heterozygous mutants *RPL22***a**/*rpl22*αΔ and *RPL22*α/*rpl22***a**Δ are viable, we propose two possible models to explain the lack of complementation. The first is a model involving an expression effect in which differential expression levels of the two *RPL22* paralogs at ectopic locations might hamper functional complementation; the second is a model involving position effect in which each *RPL22* paralog has to be in its own *MAT* locus.

We determined that an RNAi-mediated mechanism regulates *RPL22***a** expression and that it involves a region located upstream of *RPL22***a** that includes the *LTR11* and *LTR14* elements and a predicted lncRNA; when this upstream region is silenced by sRNA, expression of *RPL22***a** is enhanced. Conversely, in the absence of sRNA (i.e. in an RNAi mutant background), or when the sRNA-targeted region was deleted (Fig. S6A), *RPL22***a** expression was strongly decreased (Fig. 4 B – E, Fig. S4, Fig. S6). This is a novel and intriguing epigenetic mechanism of gene expression regulation within the *MAT***a** locus of *C. neoformans*. Examples of LTR elements silenced by sRNA have been described also in plants and mammals as a mechanism of genome protection (ŠURBANOVSKI *et al.* 2016; MARTINEZ *et al.* 2017; SCHORN *et al.* 2017; MARTINEZ 2018). There are also other locations within the *MAT***a** locus that are characterized by *LTR* elements that are also robustly targeted by sRNA in an RNAi-dependent manner (Fig. 4B), and future studies will elucidate their impact on the gene expression, mating, and genome stability.

While we found differential expression between the *RPL22***a** and *RPL22*α paralogs, and have identified epigenetic regulation of *RPL22***a** expression, the approaches employed did not enable us to define whether the Rpl22 paralogs play specific cellular roles. We then applied a newly developed CRISPR approach to generate haploid isogenic *C. neoformans* strains exchanging the *MAT RPL22* genes: *MAT*α-*RPL22***a** (strain YFF92α) and *MAT***a**-*RPL22*α (strain YFF116**a**) (Fig. S7). Unilateral crosses involving *MAT*α-*RPL22***a** x KN99**a**, and H99α x *MAT***a**-*RPL22*α, exhibited sexual reproduction features similar to the wild type cross H99α x KN99**a**, including dikaryotic hyphae, clamp connections, basidia, and basidiospore chains (Figs. 5-6, File S2). Conversely, the bilateral cross *MAT*α-*RPL22***a** x *MAT***a**-*RPL22*α (cross YFF92α x YFF116**a**) produced regular hyphae and clamp connections, but with irregular basidia and few or no spores that were characterized by a low germination and viability following microdissection, suggesting a defect in nuclear fusion, meiosis, or sporulation (Fig. 6, File S3). This may be due to the drastic reduction of both *RPL22***a** and *RPL22*α expression (Fig. 6B), combined with a phenotypic and morphological defect of the strain *MAT***a**-*RPL22*α (YFF116). Indeed, this strain displayed increased sensitivity to hydroxyurea (Fig. 7, Fig. S8), a chemical that causes DNA damage, and hinders cell cycle progression and signaling checkpoint protection (SINGH AND XU 2016).

Lastly, we have also generated a chimeric c*RPL22*α-*MAT***a** (YFF113) exchange strain of *C. neoformans* to initially test the phenotypic consequences of exchanging Rpl22, with a focus on the 5 amino acid differences located in the N-terminal region of the protein (Fig. 5; Fig. S7).

While this exchange strain does not display any morphological or phenotypic defects (Fig. 5;

Fig. S9), its analysis turned out to be of interest with respect to the mechanisms of regulation of

RPL22α. The strains *MAT***a**-c*RPL22α* (YFF113) and *MAT***a**-*RPL22α* (YFF116) both encode an

Rpl22α protein, yet they display very different *RPL22* expression patterns and distinct

phenotypes (Fig. S8C, Fig S9). In *S. cerevisiae* introns play a crucial role for *RPL22* expression,

with Rpl22 playing an extraribosomal role in inhibiting the splicing of the *RPL22B* pre-mRNA

transcript through direct binding of its intron (GABUNILAS AND CHANFREAU 2016; ABRHAMOVA

et al. 2018). Moreover, this mechanism of autoregulation seems to be conserved also in

Kluyveromyces lactis, a Saccharomycotina species that did not undergo the whole genome

duplication event and retains only one copy of the *RPL22* gene (SCANNELL et al. 2007). Similar

mechanisms might operate to control expression of *C. neoformans RPL22α*. Considering the

high expression of the chimeric c*RPL22α* but not *RPL22α* (Fig. S8C), one could hypothesize that

their different introns could potentially have a regulatory role in *RPL22α* expression. The two

Rpl22α-coding genes in strains *MAT***a**-c*RPL22α* (YFF113) and *MAT***a**-*RPL22α* (YFF116) differ

only in the 3′ region, which for strain YFF113 is from *RPL22***a** and includes introns 2 and 3.

Intron 1, which is the largest and most divergent between the *RPL22* genes (Fig. 2; Figs. S7,

S10), is the same in the *RPL22α* and c*RPL22α* allele and can therefore be excluded. Introns 2 of

*RPL22***a** and *RPL22α* share high sequence similarities and have the same intronic features (Fig.

2; Fig. S10). Introns 3 share lower sequence similarities, and the main differences are found in

pre-mRNA secondary structure and nucleotide composition in the region between the branch site

location and the 3′ acceptor site (Fig. 2; Fig S10), which is known to affect splicing and gene

expression (GAHURA *et al.* 2011; PLASS *et al.* 2012; ZAFRIR AND TULLER 2015). Furthermore,

another issue could also be that the canonical branch site of intron 3 of *RPL22α* might be too

close to the donor site with inhibition of the lariat formation, while intron 3 of *RPL22***a** has a

possible more distal canonical branch site (Fig. 2). Based on these observations, we speculate

that intron 3 of *RPL22α* might be a candidate for regulatory function.

Our findings, such as the absence of morphological defects of heterozygous *RPL22/rpl22*

mutants, the lack of cross complementation between the *RPL22* paralogs, and the morphological

and genetic defect of exchange strain MAT**a-***RPL22α*, may support a model in which the two

RPL22 MAT essential genes operate as a type of imprinting system to ensure fidelity of sexual

reproduction to enforce coordinate segregation of the opposite *MAT* nuclei in the dikaryotic

hyphae.

ACKNOWLEDGMENTS

We thank Shelly Clancey and Alexander Idnurm for critical comments on the manuscript.

FUNDING INFORMATION

This work was supported by NIH/NIAID R01 grant AI50113-15 and by NIH/NIAID R37 MERIT

award AI39115-21 (to J.H.) and grant KU 517/ 15-1 (U.K.) from the German Research Foundation

(DFG).

References

Abrhamova, K., F. Nemcko, J. Libus, M. Prevorovsky, M. Halova *et al.*, 2018 Introns provide a platform for intergenic regulatory feedback of *RPL22* paralogs in yeast. PLoS One 13**:** e0190685.

Anderson, S. J., J. P. Lauritsen, M. G. Hartman, A. M. Foushee, J. M. Lefebvre *et al.*, 2007 Ablation of ribosomal protein L22 selectively impairs αß T cell development by activation of a p53-dependent checkpoint. Immunity 26**:** 759-772.

Arras, S. D., J. L. Chitty, K. L. Blake, B. L. Schulz and J. A. Fraser, 2015 A genomic safe haven for mutant complementation in *Cryptococcus neoformans*. PLoS One 10**:** e0122916.

Arras, S. D., and J. A. Fraser, 2016 Chemical inhibitors of non-homologous end joining increase targeted construct integration in *Cryptococcus neoformans*. PLoS One 11**:** e0163049.

Beznoskova, P., S. Wagner, M. E. Jansen, T. von der Haar and L. S. Valasek, 2015 Translation initiation factor eIF3 promotes programmed stop codon readthrough. Nucleic Acids Res 43**:** 5099-5111.

Brown, G. D., D. W. Denning, N. A. Gow, S. M. Levitz, M. G. Netea *et al.*, 2012 Hidden killers: human fungal infections. Sci Transl Med 4**:** 165rv113.

Coelho, M. A., G. Bakkeren, S. Sun, M. E. Hood and T. Giraud, 2017 Fungal Sex: The Basidiomycota. Microbiol Spectr 5(3). FUNK-0046-2016..

Dahlmann, T. A., and U. Kück, 2015 Dicer-dependent biogenesis of small RNAs and evidence for microRNA-like RNAs in the penicillin producing fungus *Penicillium chrysogenum*. PLOS One 10**:** e0125989.

Davidson, R. C., J. R. Blankenship, P. R. Kraus, M. de Jesus Berrios, C. M. Hull *et al.*, 2002 A PCR-based strategy to generate integrative targeting alleles with large regions of homology. Microbiology 148**:** 2607-2615.

Dumesic, P. A., P. Natarajan, C. Chen, I. A. Drinnenberg, B. J. Schiller *et al.*, 2013 Stalled spliceosomes are a signal for RNAi-mediated genome defense. Cell 152**:** 957-968.

Fan, Y., and X. Lin, 2018 Multiple applications of a transient CRISPR-Cas9 coupled with electroporation (TRACE) system in the *Cryptococcus neoformans* species complex. Genetics 208**:** 1357-1372.

Fang, Y., L. Cui, B. Gu, F. Arredondo and B. M. Tyler, 2017 Efficient genome editing in the oomycete *Phytophthora sojae* using CRISPR/Cas9. Curr Protoc Microbiol 44**:** 21a.21.21-21a.21.26.

Fraser, J. A., S. Diezmann, R. L. Subaran, A. Allen, K. B. Lengeler *et al.*, 2004 Convergent evolution of chromosomal sex-determining regions in the animal and fungal kingdoms. PLoS Biol 2**:** e384.

Friedman, R. Z., S. R. Gish, H. Brown, L. Brier, N. Howard *et al.*, 2018 Unintended side effects of transformation are very rare in *Cryptococcus neoformans*. G3: Genes, Genomes, Genetics 8**:** 815-822.

Fu, C., and J. Heitman, 2017 *PRM1* and *KAR5* function in cell-cell fusion and karyogamy to drive distinct bisexual and unisexual cycles in the *Cryptococcus* pathogenic species complex. PLoS Genet 13**:** e1007113.

Fu, J., E. Hettler and B. L. Wickes, 2006 Split marker transformation increases homologous integration frequency in *Cryptococcus neoformans*. Fungal Genetics and Biology 43**:** 200-212.

Gabunilas, J., and G. Chanfreau, 2016 Splicing-mediated autoregulation modulates Rpl22p expression in *Saccharomyces cerevisiae*. PLoS Genet 12**:** e1005999.

Gahura, O., C. Hammann, A. Valentova, F. Puta and P. Folk, 2011 Secondary structure is required for 3' splice site recognition in yeast. Nucleic Acids Res 39**:** 9759-9767.

Hull, C. M., R. M. Raisner and A. D. Johnson, 2000 Evidence for mating of the "asexual" yeast *Candida albicans* in a mammalian host. Science 289**:** 307-310.

Ianiri, G., and A. Idnurm, 2015 Essential gene discovery in the basidiomycete *Cryptococcus neoformans* for antifungal drug target prioritization. mBio 6**:** e02334-02314.

Idnurm, A., 2010 A tetrad analysis of the basidiomycete fungus *Cryptococcus neoformans*. Genetics 185**:** 153–163.

Janbon, G., K. L. Ormerod, D. Paulet, E. J. Byrnes, V. Yadav *et al.*, 2014 Analysis of the genome and transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex RNA expression and microevolution leading to virulence attenuation. PLoS Genet 10**:** e1004261.

Johnston, G. C., J. A. Prendergast and R. A. Singer, 1991 The *Saccharomyces cerevisiae MYO2* gene encodes an essential myosin for vectorial transport of vesicles. J Cell Biol 113**:** 539-551.

Kim, S. J., and R. Strich, 2016 Rpl22 is required for *IME1* mRNA translation and meiotic induction in *S. cerevisiae*. Cell Div 11**:** 10.

Kol, G., G. Lev-Maor and G. Ast, 2005 Human-mouse comparative analysis reveals that branch-site plasticity contributes to splicing regulation. Hum Mol Genet 14**:** 1559-1568.

Komili, S., N. G. Farny, F. P. Roth and P. A. Silver, 2007 Functional specificity among ribosomal proteins regulates gene expression. Cell 131**:** 557-571.

Langmead, B., C. Trapnell, M. Pop and S. L. Salzberg, 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biology 10**:** R25.

Lengeler, K. B., D. S. Fox, J. A. Fraser, A. Allen, K. Forrester *et al.*, 2002 Mating-type locus of *Cryptococcus neoformans*: a step in the evolution of sex chromosomes. Eukaryotic Cell 1**:** 704-718.

Lengeler, K. B., P. Wang, G. M. Cox, J. R. Perfect and J. Heitman, 2000 Identification of the *MAT*a mating-type locus of *Cryptococcus neoformans* reveals a serotype A *MAT*a strain thought to have been extinct. Proc Natl Acad Sci U S A 97**:** 14455-14460.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The sequence alignment/map format and SAMtools. Bioinformatics 25**:** 2078-2079.

Lorenz, R., S. H. Bernhart, C. Honer Zu Siederdissen, H. Tafer, C. Flamm *et al.*, 2011 ViennaRNA package 2.0. Algorithms Mol Biol 6**:** 26.

Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads. 2011 17**:** 3.

Martinez, G., 2018 tRNA-derived small RNAs: New players in genome protection against retrotransposons. RNA Biol 15**:** 170-175.

Martinez, G., S. G. Choudury and R. K. Slotkin, 2017 tRNA-derived small RNAs target transposable element transcripts. Nucleic Acids Res 45**:** 5142-5152.

McIntosh, K. B., and J. R. Warner, 2007 Yeast ribosomes: variety is the spice of life. Cell 131**:** 450-451.

Narla, A., and B. L. Ebert, 2010 Ribosomopathies: human disorders of ribosome dysfunction. Blood 115**:** 3196-3205.

Nielsen, K., G. M. Cox, P. Wang, D. L. Toffaletti, J. R. Perfect *et al.*, 2003 Sexual cycle of *Cryptococcus neoformans* var. *grubii* and virulence of congenic **a** and αisolates. Infect Immun 71**:** 4831-4841.

Plass, M., C. Codony-Servat, P. G. Ferreira, J. Vilardell and E. Eyras, 2012 RNA secondary
structure mediates alternative 3'ss selection in *Saccharomyces cerevisiae.* RNA 18**:** 1103-
1115.
Rajasingham, R., R. M. Smith, B. J. Park, J. N. Jarvis, N. P. Govender *et al.*, 2017 Global burden
of disease of HIV-associated cryptococcal meningitis: an updated analysis. Lancet Infect
Dis 17**:** 873-881.
Rio, D. C., M. Ares, Jr., G. J. Hannon and T. W. Nilsen, 2010 Purification of RNA using TRIzol
(TRI reagent). Cold Spring Harb Protoc 2010**:** pdb.prot5439.
Sanchez-Sandoval, E., C. Diaz-Quezada, G. Velazquez, L. F. Arroyo-Navarro, N. Almanza-
Martinez *et al.*, 2015 Yeast mitochondrial RNA polymerase primes mitochondrial DNA
polymerase at origins of replication and promoter sequences. Mitochondrion 24**:** 22-31.
Scannell, D. R., G. Butler and K. H. Wolfe, 2007 Yeast genome evolution--the origin of the
species. Yeast 24**:** 929-942.
Schorn, A. J., M. J. Gutbrod, C. LeBlanc and R. Martienssen, 2017 LTR-retrotransposon control
by tRNA-derived small RNAs. Cell 170**:** 61-71.e11.
Schwartz, S., E. Hall and G. Ast, 2009 SROOGLE: webserver for integrative, user-friendly
visualization of splicing signals. Nucleic Acids Res 37**:** W189-192.
Schwartz, S. H., J. Silva, D. Burstein, T. Pupko, E. Eyras *et al.*, 2008 Large-scale comparative
analysis of splicing signals and their corresponding splicing factors in eukaryotes.
Genome Res 18**:** 88-103.
Singh, A., and Y.-J. Xu, 2016 The cell killing mechanisms of hydroxyurea. Genes 7**:** 99.
Srikantha, T., K. J. Daniels, C. Pujol, N. Sahni, S. Yi *et al.*, 2012 Nonsex genes in the mating
type locus of *Candida albicans* play roles in a/alpha biofilm formation, including
impermeability and fluconazole resistance. PLoS Pathog 8**:** e1002476.
Steffen, K. K., M. A. McCormick, K. M. Pham, V. L. MacKay, J. R. Delaney *et al.*, 2012
Ribosome deficiency protects against ER stress in *Saccharomyces cerevisiae*. Genetics
191**:** 107-118.
Šurbanovski, N., M. Brilli, M. Moser and A. Si-Ammour, 2016 A highly specific microRNA-
mediated mechanism silences LTR retrotransposons of strawberry. The Plant Journal 85**:**
70-82.
Toffaletti, D. L., T. H. Rude, S. A. Johnston, D. T. Durack and J. R. Perfect, 1993 Gene transfer
in *Cryptococcus neoformans* by use of biolistic delivery of DNA. J. Bacteriol. 175**:** 1405-
1411.
Upadhya, R., W. C. Lam, B. T. Maybruck, M. J. Donlin, A. L. Chang *et al.*, 2017 A fluorogenic
C. neoformans reporter strain with a robust expression of m-cherry expressed from a safe
haven site in the genome. Fungal Genet Biol 108**:** 13-25.
Wang, X., Y. P. Hsueh, W. Li, A. Floyd, R. Skalsky *et al.*, 2010 Sex-induced silencing defends
the genome of *Cryptococcus neoformans* via RNAi. Genes Dev 24**:** 2566-2582.
Warner, J. R., and K. B. McIntosh, 2009 How common are extraribosomal functions of
ribosomal proteins? Mol Cell 34**:** 3-11.
Xue, S., and M. Barna, 2012 Specialized ribosomes: a new frontier in gene regulation and
organismal biology. Nat Rev Mol Cell Biol 13**:** 355-369.
Zafir, Z., and T. Tuller, 2015 Nucleotide sequence composition adjacent to intronic splice sites
improves splicing efficiency via its effect on pre-mRNA local folding in fungi. RNA
(New York, N.Y.) 21**:** 1704-1718.

Zhang, Y., M. N. O'Leary, S. Peri, M. Wang, J. Zha *et al.*, 2017 Ribosomal proteins Rpl22 and
Rpl22l1 control morphogenesis by regulating pre-mRNA splicing. Cell Rep 18**:** 545-556.

 Figure legends

Figure 1. The *MAT* locus contains essential genes encoding ribosomal proteins Rpl39 and
Rpl22. Representative example of genetic analysis of two *C. neoformans* heterozygous mutants
[GI233 (*RPL39***a**/*rpl39α*Δ) and GI56 (*RPL22α*/*rpl22***a**Δ)]. The first colony in red box in the top
left corner represents the original heterozygous mutant from which the progeny analyzed
originated. The remaining isolates were haploid germinated spore progeny grown on control
medium YPD, YPD + NAT, SD –uracil, and SD – adenine. The *MAT* type of the progeny is
indicated on the right panel (Δ indicates the heterozygous mutant).

Figure 2. Intron analysis of *C. neoformans RPL22* paralogs. (A) Alignment of introns 1, 2, and 3
of *RPL22α* and *RPL22***a**. Donor and acceptor splice sites are indicated in red. Branch sites were
predicted using a combination of the online software SROOGLE (http://sroogle.tau.ac.il/)
(SCHWARTZ *et al.* 2009) and SVM-BP finder
(http://regulatorygenomics.upf.edu/Software/SVM_BP/): branch sites based on the algorithm of
Kol are represented in blue (KOL *et al.* 2005), those based on the algorithm of Schwarts are in
purple (SCHWARTZ *et al.* 2008). Manual adjustments were performed based on the websites
instructions and score prediction; both algorithms failed to identify a CTRAY canonical branch
site of intron 3 of *RPL22***a** (boxed). The polypyrimidine tracts are underlined. The percentage of
identity is indicated in parentheses, while the length (in bp) of the introns is indicated at the end
on each alignment. (B) Consensus sequences at the 5′ splice site, the branch site, and the 3′ splice
site constructed using WebLogo 3.3. bits, binary digits (http://weblogo.threeplusone.com/).
Underlined are the canonical donor GT and acceptor AG.

Figure 3. *RPL22***a** complements to restore viability of *rpl22***a** mutants, *RPL22α* does not. Genetic
analysis of *C. neoformans* complemented NEO[R] strains GI151, GI102, GI154 and GI104 derived
from heterozygous mutant GI56 (*RPL22α*/*rpl22***a**Δ). The first colony in the top left corner
represents the original heterozygous mutant from which the progeny analyzed originate (red
boxes). The mating type in NAT[R] colonies was scored by genetic crosses and by PCR with
primers JOHE39201-JOHE39204. For PCR, the WT (H99 *MATα* and KN99**a** *MAT***a**) and the
heterozygous strain that was analyzed were used as controls; the NAT[R] progeny is indicated with
"P", and the number that corresponds to the position in the plate. C = negative control.

Figure 4. RNAi contributes to control expression of *RPL22***a**. *RPL22*α and *RPL22***a** expression during *C. neoformans* mating between WT H99α x KN99**a** cross (A), *rdp1*Δ and *gwc1*Δ bilateral crosses (C, D), and GI228 x H99 cross (E) for 24, 48, 72 and 96 h of incubation. Ct values were converted to expression level (fold change) through comparison with the endogenous reference *GDP1* (ΔΔct analysis); asterisk indicates p <0.05 for each *RPL22*α and *RPL22***a** comparison for the same day of incubation. Note the different scales on the y axes of the graphs in A - C - D – E. (B) sRNA obtained during H99α x KN99**a** cross (black) and *rdp1*Δ bilateral cross (red) were mapped to the reference *MAT***a** locus of *C. neoformans* (accession number AF542528.2); genes are represented in grey in the middle panel; in blue the *LTR* and transposable elements. *LTR11, LTR14* and *RPL22***a** of interest in this study are indicated in bold.

Figure 5. Construction, analysis and sexual reproduction of *RPL22* exchange strains. (A) Schematic representation of the generation of the *C. neoformans RPL22* exchange strains YFF92 and YF113. Lightning bolts in different colors denote different gRNA targeting sites. (B) Mating phenotypes during crosses between H99α x KN99**a**, and YFF92 x KN99**a**, H99α x YFF113, and YFF92 x YFF113. The scale bar is 100 μm. (C) Genetic analysis of progeny obtained from the YFF92 x YFF113 bilateral cross; note the expected 1:1 segregation of the *SH2*::*NAT* and *SH2*::*NEO* markers in progeny produced from YFF92 x YFF113 cross, and the independent segregation of the *MAT* loci, with *MAT***a** indicated in grey and *MAT*α indicated in white. (D) RT-qPCR of *RPL22***a** and c*RPL22*α expression during the YFF92 x YFF113 cross. Asterisk indicates p<0.05 for each c*RPL22*α and *RPL22***a** comparison for the same day of incubation.

Figure 6. Reciprocal *RPL22* exchange strains exhibit defects in sexual reproduction. (A) Schematic representation of the generation of the *C. neoformans RPL22* swapped strain YFF116. Lightning bolts in different colors denote different gRNA targeting sites. (B) RT-qPCR of *RPL22***a** and *RPL22*α expression during YFF92 x YFF116 cross; asterisk indicates p<0.05 for each c*RPL22*α and *RPL22***a** comparison for the same day of incubation. (C - D) Mating phenotypes during crosses between the cross H99 x YFF116, and YFF92 x YFF116. The scale bar is 100 μm. (E) Scanning electron microscopy of sexual structures of the YFF92 x YFF116 cross; note that the majority of the basidia are bald with no basidiospore chains. The scale bar is 10 μm. (F) Scanning electron microscopy of an irregular basidium produced by the YFF92 x YFF116 cross. The scale bar is 10 μm. (G) Scanning electron microscopy of a basidium with a

collapsed chain of basidiospores produced by the YFF92 x YFF116 cross. The scale bar is 1 μm.

(H) Scanning electron microscopy of a regular unfused clump connection produced by the

YFF92 x YFF116 cross. The scale bar is 1 μm.

Figure 7. Phenotypic analysis of *RPL22*α *MAT***a** exchange strain. (A) Phenotype of the *C.*

neoformans WT parental strain KN99**a** and the exchange *RPL22* strain YFF116 on YPD at 30°C

(control condition), on hydroxyurea (125 μg/mL), and at 39°C. (B) Microphotographs of *C.*

neoformans KN99**a** and strain YFF116 after growth on YPD at 30°C and on hydroxyurea (125

μg/mL). The scale bar is 10 μm.

Table 1. Genetic analysis of *C. neoformans* heterozygous mutants for the *MAT*α and *MAT***a** ribosomal proteins Rpl39 and Rpl22.

Genotype	Strain	Protein function	Germinated basidiospores/ dissected basidiospores	NAT-R progeny/ basidiospore germinated	Phenotype
*RPL39*α/*rpl39***a**Δ	GI230	Large Subunit Ribosomal Protein L39	31/80 – 38.7%	0/31	Inviable
*RPL39***a**/*rpl39*αΔ	GI233		29/70 – 41.4%	0/29	Inviable
*RPL22*α/*rpl22***a**Δ	GI56	Large Subunit Ribosomal Protein L22	23/63 – 36.5%	1/23*	Inviable
*RPL22***a**/*rpl22*αΔ	GI81		28/72 – 38.8%	0/28	Inviable

* The NAT-R progeny obtained is likely aneuploid for chromosome 5, because PCR confirmed the presence of the *RPL22*α allele and the mutated copy of the *RPL22***a.**

Table 2. Genetic analysis of *C. neoformans* complementing strains of heterozygous mutants *RPL22**a**/rpl22*α∆ and *RPL22*α*/rpl22**a**∆

Strain	Genotype	Germinated basidiospores/ dissected basidiospores	*NAT*-R	*NEO*-R	*NAT*-R + *NEO*-R	*MAT**a***	*MAT*α	*MAT**a*** - *MAT*α	Phenotype
GI86	*RPL22**a**/rpl22*α∆*NAT* + *RPL22*α*-NEO*	19/70	0	8	**4**	15	**4**	0	Viable
GI150	*RPL22**a**/rpl22*α∆*NAT* + *RPL22**a**-NEO*	NA	NA	NA	NA	NA	NA	NA	NA
GI83	*RPL22**a**/rpl22*α∆*NAT* + empty *NEO* vector	11/63	0	7	0	11	0	0	Inviable
GI151	*RPL22*α*/rpl22**a**∆NAT* + *RPL22**a***-NEO	19/63	0	6	**3**	**3**	16	0	Viable
GI102	*RPL22*α*/rpl22**a**∆NAT* + *RPL22*α*-NEO*	17/58	2*	0	0	0	15	2	Inviable
GI154	*RPL22*α*/rpl22**a**∆NAT* + *RPL22*α*-NEO*	20/69	1*	12	3	0	16	4	Inviable
GI104	*RPL22*α*/rpl22**a**∆NAT* + empty *NEO* vector	10/57	0	8	1	0	9	1	Inviable

* NAT-R progeny with 1 extra copy of *RPL22α*

NA: not available (strain GI150 did not produce basidiospores for analysis)

In bold and underlined are the progeny with markers that indicate success of the complementation experiment

Table 3. Genetic analysis of crosses between allele exchange strains

Cross	Spores dissected	Spores germinated	*MATα-RPL22α*	*MATa - RPL22a*	*MATa - MATα*	Mitochondria from *MATa*	Mitochondria from *MATα*
H99 x KN99**a**	60	27	11	14	2	27	0

Cross	Spores dissected	Spores germinated	*MATα-RPL22a*	*MATa - RPL22a*	*MATa - MATα*	NAT-R progeny (*MATa*; *MATα*; *MATa/α*)	Mitochondria from *MATa*	Mitochondria from *MATα*
YFF92 x KN99**a**	60	18	9	6	3	7 (4; 2; 1)	18	0

Cross	Spores dissected	Spores germinated	*MATa - cRPL22α*	*MATα-RPL22α*	*MATa - MATα*	NEO-R progeny (*MATa*; *MATα*; *MATa/α*)	Mitochondria from *MATa*	Mitochondria from *MATα*
YFF113 x H99	62	49	27	19	3	31 (16; 12; 3)	49	0

Cross	Spores dissected	Spores germinated	*MATa-RPL22α*	*MATα-RPL22α*	*MATa - MATα*	NEO-R progeny (*MATa*; *MATα*; *MATa/α*)	Mitochondria from *MATa*	Mitochondria from *MATα*
YFF116 x H99	66	43	23	17	3	23 (10; 10; 3)	43	0

Cross	Spores dissec.	Spores germ.	*MATa-cRPL22α*	*MATα-RPL22a*	*MATa - MATα*	NEO-R (*MATa*; *MATα*; *MATa/α*)	NAT-R (*MATa*; *MATα*; *MATa/α*)	NEO-R NAT-R	Mito. *MATa*	Mito. *MATa*
YFF92 x YFF113	63	28	14	13	1	14 (9; 5; 0)	14 (6; 7; 0)	0	28	0
YFF92 x YFF116	58	7	3	0	4	1 (1; 0; 0)	4 (0; 0; 4)	0	NA	NA

NA: not available

GI233 - *RPL39***a**/*rpl39*αΔ

GI56 - *RPL22*α/*rpl22***a**Δ

YPD   NAT   no adenine   no uracil   *MAT*

A

```
Introns 1 (75.86% identity)
Intron 1_α    gtgcgttttattttttctttttgattttctcgtatcccatgtaatcctgattgcaaacgt
Intron 1_a    gtgcgtttcct--------------tttctcatatcccatgtaaccctcattgcaacttc
              *******  *        ***** ************ *** *******

Intron 1_α    ttgcaacattgggttgactggctcttgtccctaatcttacccatttccgctaatatgactc
Intron 1_a    ttgcaattttgggttg--tggatgctgcgctaaatttagccatttccgctaacgtgtctt
              ******   *******    **   ** *  **** *** **************  ** **

Intron 1_α    gttttcaaacag  132
Intron 1_a    gtgttcaaacag  116
              ** ********

Introns 2 (74.55% identity)
Intron 2_α    gtgagtgtgcccaaaaaatcactatttgaataatgttctgatggtcgttgcacag  55
Intron 2_a    gtcagtacgcccgaaaagttagtagatgagttgtgttctgatggttgtttcacag  55
              ** ***   **** **** * * ** *** *  *********** *** *****

Introns 3 (60.42% identity)
Intron 3_α    gttagtcatgtgtaacacgcctgatattatatattaacatttctacag  48
Intron 3_a    gttagtaacgtgtagatacattgatactatatactgattgcctcatag  48
              ****** * *****   *  ***** ******* * *      * **
```

B

RPL22α

RPL22a

GI151 - *RPL22α/rpl22**a**Δ + RPL22**a***

GI102 - *RPL22α/rpl22**a**Δ + RPL22α*

GI154 - *RPL22α/rpl22**a**Δ + RPL22α*

GI104 - *RPL22α/rpl22**a**Δ + pSDMA57*

| YPD | NAT | NEO | no adenine | no uracil | *MAT* |

A

YFF92 (*MAT*α) - *rpl22α::RPL22**a** SH2::NAT*

H99

ATG   *RPL22α (616 bp)*   TAA

*RPL22**a** (600 bp)*

SH2

NAT

YFF113 (*MAT**a***) - *rpl22a::RPL22α^N-RPL22**a**^C SH2::NEO*

ATG   *RPL22**a** (600 bp)*   TAA

RPL22α   *RPL22**a***
(1-268 bp)   (253-600 bp)

SH2

NEO

KN99**a**

B

H99 x KN99**a**   KN99**a** x YFF92   H99 x YFF113   YFF92 x YFF113

C

YPD   Mating type

a ⍺

NEO   NAT

D

▨ *RPL22**a*** in H99   ▨ c*RPL22α* in KN99**a**

Expression level (ΔΔct) Fold Change

1.5

1.0

0.5

24   48   72   96

Incubation time (hours)

*

A YFF116 (*MAT***a**) - *rpl*22**a**::*RPL*22α *NEO*

KN99**a**

ATG   *RPL22***a** (600 bp)   TAA

*RPL22*α (616 bp)

NEO

B

▨ *RPL22***a** in H99   ▨ *RPL22*α in KN99**a**

Expression level (ΔΔct) Fold Change

*

Incubation time (hours)

C H99 x YFF116

D YFF92 x YFF116

E

F

G

H

Figure S1. Alignment between *C. neoformans RPL22*α and *RPL22***a.** DNA sequence (A), and Rpl22α and Rpl22**a** proteins (B). In A introns are highlighted in red, and nucleotide changes that result in different amino acids between the Rpl22α and Rpl22**a** proteins are color coded with the corresponding amino acids boxed and highlighted in the same color in B. In B only the initial 60 amino acids are shown. Both in A and B, an asterisk (*) indicates positions which have a single, fully conserved residue, a colon (:) indicates conservation between groups of strongly similar properties - scoring > 0.5 in the Gonnet PAM 250 matrix, and a period (.) indicates conservation between groups of weakly similar properties - scoring =< 0.5 in the Gonnet PAM 250 matrix.

Figure S2 Expression of *RPL22*α and *RPL22***a** during vegetative growth of *C. neoformans.* WT strains H99α and KN99**a** were grown in YPD media for 24, 48, 72 and 96 h. Ct values were converted to expression level (fold change) through comparison with the endogenous reference *GDP1* ($^\Delta$ct analysis). The data presented in this supplemental figure were used as comparative condition to calculate the expression levels of *RPL22*α and *RPL22***a** during H99α x KN99**a** mating of Figure 4A. Asterisk indicates p<0.05 for each *RPL22*α and *RPL22***a** comparison for the same day of incubation.

Figure S3. Intron retention and splice junction analysis for *RPL22*α following growth in several conditions. High retention of intron 1 at 30°C static growth. Detected alternative splicing at intron 1 and 2 at 37°C static growth.

Figure S4. Comparison of the expression levels of *RPL22*α and *RPL22***a** in WT and RNAi mutants. RT-qPCR data from Figures 4A, 4C and 4D, were replotted to allow direct comparison of the *RPL22*α and *RPL22***a** expression levels during WT H99α x KN99**a** cross compared to *rdp1*Δ (A) and *gwc1*Δ (B) bilateral crosses for 24, 48, 72 and 96 h of incubation. Statistical analysis was carried out using one-way ANOVA with Tukey's multiple comparison test. Because on these comparisons we were interested in monitoring the changes

in gene expression following genetic manipulation, we display only statistically significant differences (* for $p<0.05$, ** for $p<0.01$, and *** for $p<0.001$) of the expression levels of the same gene for the same day of incubation in separate mating reactions.

Figure S5. sRNA obtained during H99α x KN99**a** cross (black) and *rdp1Δ MAT*α x *rdp1Δ MAT***a** bilateral cross (red) were mapped to the reference *MAT*α locus of *C. neoformans* (accession number AF542529.2). Genes are represented in grey in the middle panel, and the *LTR* and transposable elements are in blue.

A

B

Figure S6. Analysis of 5′ control element of *RPL22***a** gene during sexual reproduction. (A) Schematic representation of the CRISPR/Cas9 strategy used to generate strain *C. neoformans* GI288 (5′ Δ *RPL22***a**) through deletion of the region targeted by sRNA upstream *RPL22***a** during H99α x KN99**a** cross. (B) RT-qPCR data from Figures 4A and 4D were replotted to allow direct comparison of the *RPL22***a** and *RPL22*α expression levels during WT H99α x KN99**a** cross compared to GI288**a** (5′ Δ *RPL22***a**) x H99α cross for 24, 48, 72, and 96 h of incubation. (C) RT-qPCR data from Figures 4C and 4D were replotted to allow direct comparison of the *RPL22***a** and *RPL22*α expression levels during *rdp1*Δ bilateral cross compared to GI288 x H99 cross for 24, 48, 72 and 96 h of incubation. For both B and C, statistical analysis was carried out using one-way ANOVA with Tukey's multiple comparison test. These comparisons were to monitor the changes in gene expression following genetic manipulation, and only statistically significant differences of the expression levels of the same gene in the same day of incubation in separate mating reactions are displayed (* for $p<0.05$, and ** for $p<0.01$).

Figure S7. Features of the *RPL22*α and *RPL22***a** genes in *C. neoformans* WT H99α and KN99**a** (A), respectively, and in the *RPL22* exchange strains YFF92, YFF113, and YFF116 (B). Boxes represent exons. The start and stop codons, the different amino acids between the Rpl22α and Rpl22**a** proteins, and the introns (" i ") are indicated. The chimeric *RPL22*α is indicated as c*RPL22*α (c = chimera).

Figure S8. Comparison of the expression levels of *RPL22α*, c*RPL22α*, and *RPL22***a** during mating in the *RPL22* exchange strains. (A) RT-qPCR data from Figure 4A and Figure 5D

were replotted to allow direct comparison of the *RPL22*α and *RPL22***a** expression levels during WT H99α x KN99**a** cross compared to c*RPL22*α and *RPL22***a** expression levels during YFF92α x YFF113**a** cross for 24, 48, 72, and 96 h of incubation. (B) RT-qPCR data from Figure 4A and Figure 6B were replotted to allow direct comparison of the *RPL22*α and *RPL22***a** expression levels during WT H99α x KN99**a** cross compared to YFF92α x YFF116**a** cross for 24, 48, 72, and 96 h of incubation. (C) RT-qPCR data from Figure 5D and Figure 6B were replotted to allow direct comparison of the *RPL22*α and *RPL22***a** expression levels during YFF92α x YFF116**a** cross compared to c*RPL22*α and *RPL22***a** expression levels during YFF92α x YFF113**a** cross for 24, 48, 72, and 96 h of incubation. Statistical analysis was carried out using one-way ANOVA with Tukey's multiple comparison test. Because in these comparisons we were interested in monitoring the changes in gene expression following genetic manipulation, we display only statistically significant differences (* for $p<0.05$, and ** for $p<0.01$) of the expression levels of the same gene in the same day of incubation in separate mating reactions.

Figure S9. Phenotypic analysis of the *C. neoformans* heterozygous and mutant strains generated in this study (Table S1) on several stressors. For details see materials and methods.

*RPL22*α   *RPL22***a**

Intron 1

Intron 2

Intron 3

Figure S10. RNAfold secondary structure prediction of the regions including the branch sites and the acceptor sites of introns 1, 2, and 3 of the *RPL22*α and *RPL22***a** genes.

File S2

Morphological analysis of WT strains during mating

Morphological analysis of YFF92 x KN99**a** (only *RPL22***a**) strains during mating

Morphological analysis of YFF113 x H99 (only *RPL22α*) strains during mating

Morphological analysis of *RPL22* exchange strains (YFF92 x YFF113) during mating

File S3

Morphological analysis of YFF116 x H99 (only *RPL22α*) strains during mating

Morphological analysis of *RPL22* exchange strains (YFF92 x YFF116) during mating

Morphological analysis of *RPL22* exchange strains (YFF92 x YFF116) during mating

Formation of irregular basidiospores chains during YFF92 x YFF116 mating