Title:

Cis-regulatory differences in isoform expression associate with life history strategy variation in Atlantic salmon

Authors:

Jukka-Pekka Verta [1,2], Paul Vincent Debes [1,2], Nikolai Piavchenko [1], Annukka Ruokolainen [1], Outi Ovaskainen [1], Jacqueline Emmanuel Moustakas-Verho [1,2], Seija Tillanen [1,2], Noora Parre [1], Tutku Aykanat [1,2], Jaakko Erkinaro [3] & Craig Robert Primmer [1,2]

Affiliations:

1. Organismal and Evolutionary Biology Research Programme
   University of Helsinki
   Viikinkaari 9, 00790, Helsinki, Finland
2. Institute of Biotechnology, University of Helsinki, Finland
3. Natural Resources Institute Finland (LUKE),
   Oulu, Finland

Co-corresponding authors: craig.primmer@helsinki.fi, jukka-pekka.verta@helsinki.fi

Keywords:

life history traits, genotype-phenotype association, transcript isoforms, *cis*-regulation, puberty timing, testes differentiation

Abstract:

A major goal in biology is to understand how evolution shapes variation in individual life histories. Genome-wide association studies have been successful in uncovering genome regions linked with traits underlying life history variation in a range of species. However, lack of functional studies of the discovered genotype-phenotype associations severely restrains our understanding how alternative life history traits evolved and are mediated at the molecular level. Here, we report a *cis*-regulatory mechanism whereby expression of alternative isoforms of the transcription co-factor *vestigial-like 3* (*vgll3*) associate with variation in a key life history trait, age at maturity, in Atlantic salmon (*Salmo salar*). Using a common-garden experiment, we first show that *vgll3* genotype associates with puberty timing in one-year-old salmon males. By way of temporal sampling of *vgll3* expression in ten tissues across the first year of salmon development, we identify a pubertal transition in *vgll3* expression where maturation coincided with a 66% reduction in testicular *vgll3* expression. The *late* maturation allele was not only associated with a tendency to delay puberty, but also with expression of a rare transcript isoform of *vgll3* pre-puberty. By comparing absolute *vgll3* mRNA copies in heterozygotes we show that the expression difference between the *early* and *late* maturity alleles is largely *cis*-regulatory. We propose a model whereby expression of a rare isoform from the *late* allele shifts the liability of its carriers towards delaying puberty. These results reveal how regulatory differences can be a central mechanism for the evolution of life history traits.

Author summary:

Alternative life history strategies are an important source of diversity within populations and promote the maintenance of adaptive capacity and population resilience. However, in many cases the molecular basis of different life history strategies remains elusive. Age at maturity is a key adaptive life history trait in Atlantic salmon and has a relatively simple genetic basis. Using salmon age at maturity as a model, we report a mechanism whereby different transcript isoforms of the key age at maturity gene, *vestigial-like 3* (*vgll3*), associate with variation in the timing of male puberty. Our results show how gene regulatory differences in conjunction with variation in gene transcript structure can encode for complex alternative life histories.

Introduction:

Some of the most ecologically and evolutionarily important traits involve alternative life-history strategies and the trade-offs therein [1]. One such example is maturation age, whereby early maturing individuals avoid the increased risk of mortality associated with later reproduction, but often with the trade-off of lower fecundity [2]. Genome-wide association studies have been successful in uncovering genome regions controlling for life-history trait variation such as survival and reproductive success in Soay sheep [3], early versus late age at maturity in Atlantic salmon [4,5], early versus late run timing in Chinook salmon and steelhead trout [6–8], age and size at flowering in monkeyflowers [9] and alternative reproductive strategies in ruffs [10]. However, despite this progress mapping the genetic basis of life history traits, moving beyond genome-level associations to an understanding of the molecular basis of such variation in life history traits and trade-offs has thus far remained elusive. This lack of functional insight into the discovered genotype-phenotype associations severely restrains our understanding of the molecular basis of evolution in life history traits [11].

In Atlantic salmon, age at maturity is a key adaptive life history trait that is evolving under a trade-off between survival and fecundity. Later-maturing, and therefore larger, individuals have higher reproductive fitness at spawning than earlier maturing (smaller) individuals [12,13], but also have a higher risk of mortality prior to first reproduction [2]. Males mature, on average, earlier than females and at smaller size, which is indicative of a potential sexual conflict for age at maturity in the species [14,15]. Genetic variation in a single genome region including the transcription co-factor gene *vestigial-like 3* (*vgll3*) was recently found to explain nearly 40% of variation in sea-age at maturity in both male and female salmon across 57 Northern-European wild populations [4], making salmon age-at maturity a genetically tractable model to study the molecular bases of life-history trait variation. Further, variation in *VGLL3* also associated with age at puberty in humans [16,17], indicating that functional mechanisms of puberty timing may be shared across vertebrates. However, in all cases the functional basis of the association between genetic variation in *vgll3* and age at maturity remains unknown.

The pattern of association observed in natural salmon populations can be used to generate hypotheses pertaining the functional mechanism that translates genetic variation to different maturation phenotypes. The strongest association between age at maturity and *vgll3* genotype maps to a single nucleotide polymorphism (SNP) in a non-coding region approximately 7000 basepairs 3' of *vgll3*, followed in statistical significance by mutations in *vgll3* coding sequence [4,5]. This suggests that genetic variation in linked *cis*-regulatory sequences (e.g. enhancers) may underlie the functional

differences among salmon *vgll3* genotypes by changing their expression level or pattern. In other
species, a similar pattern of linked non-coding changes is often observed in association studies for
life history traits [6,7,10,18,19], suggesting that regulatory changes may be a common driver of
evolution in life history strategies. However, non-coding and coding hypotheses pertaining the control
of maturation age are nuanced and not mutually exclusive; enhancer mutations can have effects
unlinked to their genotype (they can act in *trans*) [20,21] and non-coding mutations can alter transcript
structure, stability and translation via alternative transcription start sites and splicing [22–24].

To uncover the functional genetic basis of age at maturity variation in Atlantic salmon, we first
validated the *vgll3* association with male maturation timing in different genetic backgrounds and in
common garden settings by using 32 controlled-crossed families with known *vgll3* genotypes.
Uncertainty remains regarding the tissues and developmental stages during which *vgll3* expression
potentially influences the maturation phenotype. *Vgll3* expression and function has been linked to
diverse processes such as adipocyte differentiation and myogenesis in mice [25,26], immune function,
adiposity and puberty timing in humans [16,17]and cell fate commitment in salmon tissues including
adipose and gonads [27,28], but genotype comparisons, *vgll3* transcript structure or *cis*-regulation have
not been investigated. We determined *vgll3* temporal expression dynamics during the first year of
salmon development up to male maturation by way of a time-series sampling of ten tissues from a
total of 273 individuals, characterized *vgll3* transcript structure and further tested the importance of
cis-regulatory mechanisms among *vgll3* genotypes by assaying *vgll3* allele-specific expression in
heterozygotes. Our results reveal a potential functional mechanism that links allele-specific isoform
expression differences in *vgll3* to variation in maturation timing and thus mediates variation in life
history strategies.

Results:

vgll3 **genotype predicts male maturation frequency**

To uncover the functional genetic basis of age at maturity variation in Atlantic salmon, we first

validated that the genotype at the *vgll3* locus on chromosome 25 (termed "*E*" for early sea-age at

maturity and "*L*" for late sea-age at maturity after previous association study [4]) is associated with

age at maturity in controlled conditions by rearing 656 individuals with known *vgll3* genotypes from

32 families in common garden conditions (Fig. S1). We assessed all males for maturation status non-

lethally at the start of the breeding season (nine months post-hatching) and modeled maturation

probabilities based on observed frequencies of mature versus non-mature fish as a function of *vgll3*

genotype (see Methods). *Vgll3*EE* individuals showed a 10 times higher frequency of maturation

(20%) compared to individuals with *vgll3*LL* genotypes (2%, Fig. S2). Correspondingly, maturation

probability for *vgll3*EE* males (20.7%) was significantly higher compared to males with *vgll3*LL*

genotypes (3%) (probit 95% credible interval: 0.59 to 3.07, *P*=0.006, *vgll3*EE* n*=93 and *vgll3*LL*

n=92) (Fig. 1A). Maturation probability for males heterozygous for *vgll3* (15.4%) was closer to

*vgll3*EE* males compared to *vgll3*LL* males, and significantly different from *vgll3*LL* males (probit

95% credible interval: 0.32 to 3.88, *P*=0.018, *vgll3*EL* n*=199), but not *vgll3*EE* males.

We additionally modeled maturation probability as a function of *vgll3* genotype and fish length,

hypothesizing that early growth positively affects maturation probability [29]. Indeed, maturation

probability positively associated with fish length before the start of the breeding season (eight months

post-hatching) (probit 95% credible interval: 0.36 to 0.94, *P*=1e-4, Fig. 1B). Irrespective of size

effect, the *vgll3*EE* genotype was still associated with a higher maturation probability compared to

the *vgll3*LL* genotype (probit 95% credible interval: 0.54 to 2.84, *P*=0.005), as well as *vgll3*EL*

compared to *vgll3*LL* (probit 95% credible interval: 0.26 to 3.6, *P*=0.026).

These results demonstrate that the *vgll3*E* allele associated with young sea-age at maturity in natural

populations is associated with earlier maturation also in controlled conditions, and also supports

stronger dominance of *vgll3*E* allele as observed in males from natural populations [4]. Seeing that

concurring results were observed in another wild-derived stock of Atlantic salmon [30], the genotype-

phenotype association between *vgll3* and maturation timing is reproducible in controlled conditions

and likely common in North-European strains of Atlantic salmon.

Figure 1. *Vgll3* **genotype associates with male maturation probability in controlled conditions.** (**A**) Probability of maturation at nine months post-hatching in controlled conditions is significantly higher in *vgll3*EE* individuals compared to *vgll3*LL* individuals. Points represent means and whiskers represent 95% credible intervals from a Bayesian Markov Chain Monte-Carlo animal model (see Methods). N_{EE}=98, N_{EL}=204, N_{LL}=82 (**B**) Maturation probability modeled as a function of fish length one month before the breeding season. Blue, *vgll3*EE*; grey, *vgll3*EL*; red, *vgll3*LL*.

vgll3 expression correlates with maturation inhibition

Until now, it has remained unknown by which functional means, tissues, cell types and developmental time-points genetic variation in *vgll3* influences the maturation phenotype. In order to investigate the functional mechanisms resulting in earlier maturation in males carrying the *vgll3*EE* genotype and to establish the tissues and temporal time-points of relevance for *vgll3* expression, we first characterized *vgll3* transcript expression profiles of ten tissues and whole salmon juveniles in a time-series of samples including 13 time points and all *vgll3* genotypes by using reverse-transcription droplet digital PCR (RT-ddPCR) and a genotype-specific assay on exon 2 of the *vgll3* transcript (total *n*=291, Fig. S3).

Vgll3 expression in the heart was high relative to other tissues (Fig. 2A), but also very variable, similar to results from three-year old juveniles from natural populations [28]. Expression in liver showed a linear downward trend from younger to older individuals, although the overall expression level remained very low relative to other tissues (Fig. S4). *Vgll3* expression in muscle was lowest among tissues, while immature ovaries showed intermediate and relatively stable expression (Fig. 2A). Interestingly, while *vgll3* expression was high in immature testes, expression was reduced by 66% in mature testes compared to immature testes (37 copies/ng versus 107 copies/ng, i.e. 70 copies/ng fewer, 95% CI [47-93 fewer], two-tailed *t*-test, df=25.9, *P*=1e-06, n_{mature}=14, $n_{immature}$=65, Fig. 2B).

Vgll3 expression was similarly lower in stripped mature testes, containing mainly mitotic cell types,

and in ejaculate, containing mainly mature sperm, compared to immature testes matched for sampling

time-point (58 copies/ng (stripped testes) versus 128 copies/ng i.e. 70 copies/ng fewer, 95% CI [30-

111 fewer], two-tailed *t*-test, df=21.8, *P*=0.002, $n_{stripped}$=8; 19 copies/ng (ejaculate) versus 134

copies/ng i.e. 115 copies/ng fewer, 95% CI [50-181 fewer], df=6.6, *P*=0.005, $n_{ejaculate}$=4, respectively,

Fig. 2B). Taken together, these results indicate that in salmon gonad tissues, *vgll3* expression is high

in immature testes and down-regulated upon the onset of male puberty.

To further test whether *vgll3* down-regulation coincides with the activation of the male puberty

pathway, we compared *vgll3* expression relative to the expression of *amh,* which is a repressor of

male puberty and Sertoli cell differentiation, and *igf3*, which promotes spermatogonial germ cell

differentiation [31–34]. Consistent with our visual classification of gonad morphology, immature testes

were characterized by typically higher *amh* expression compared to *igf3* expression, whereas

morphologically mature testes with differentiated (running) milt showed typically equal, or higher

expression of *igf3* compared to *amh* (Fig. S5). Clustering based on *amh/igf3* expression ratio and

vgll3 expression level further revealed that mature testes were separated in gene expression space

from immature testes (bootstrap LRT [35] *P*<0.0001) and characterized by both low *amh/igf3* ratio

and low *vgll3* expression, indicating that down-regulation of both *amh* and *vgll3* expression is

commonly associated with completing puberty (Fig. 2C). In sum, these results point towards a

mechanism by which male puberty is associated with a down-regulation of *vgll3* expression,

corroborating previous studies that suggested *vgll3* influences Sertoli cell differentiation [27].

While the effects of *vgll3* down-regulation on the downstream transcriptional landscape of testes cell

types remain largely unknown, transcriptomic studies indicate that *vgll3* expression correlates with

changes in the Hippo signaling pathway, known to control cell differentiation and organ size [27,28].

To gain additional insight into the expression patterns of key players of Hippo signaling pathway and

vgll3 during male puberty, we re-analyzed RNA-sequencing data from immature, prepubertal and

pubertal testes from an independent dataset [34]. We found that while maturation correlates with a

reduction in *vgll3* expression, the Hippo pathway antagonist for *vgll3*, *yap1* [26,36,37], showed the

opposite expression pattern with increasing expression as maturation proceeded (Fig. S6). *Vgll3* was

co-expressed within a group of 7073 other genes out of which 5670 were assigned function in the

NCBI RefSeq database. Gene Ontology analysis revealed an over-representation in *"transcriptional*

coactivator function" in the *vgll3* co-expressed cluster (GO:0003713, *P*=0.049, Fig. S6), which

included multiple histone acetyltransferases (LOC106607167, LOC106580491, LOC106585100),

nuclear receptor coactivators (LOC106607631, LOC100196047, LOC106571447, LOC106579720,

LOC106590658), CREB-binding proteins (LOC106590442, LOC106564464) and transcriptional adapter proteins (LOC106581187, LOC106577491), suggesting that *vgll3* is embedded in a larger gene regulatory network potentially influencing gene expression on a transcriptomic level.

Figure 2. *Vgll3* **expression across salmon tissues during the first year of development.** (**A**) *Vgll3* reaches highest expression levels in the heart and immature testes. For sample size see Supplemental Fig. 3. Line, loess smoothing; shading, 95% confidence interval. (**B**) Male maturation is associated with a 66% drop in *vgll3* expression compared to immature testes expression levels. Reduction in *vgll3* expression is observed in all major tissues of the mature testes. Stripped testes; mature testes with ejaculate removed manually. Error bars show SE. (**C**) *Vgll3* expression shows a strong

association with relative expression levels of *amh/igf3*. Squares; mature testes, diamonds; pubertal samples in transition to mature morphology, spheres; immature testes. Ellipses delineate estimated 95% confidence intervals for each cluster. Expression is measured using RT-ddPCR.

Complex transcript expression differences between *vgll3* genotypes

The difference in *vgll3* expression level between mature and immature testes led us to hypothesize that the observed higher probability of maturation in *vgll3*EE* males may be linked to differences in *vgll3* expression in immature testes of *vgll3*EE* and *vgll3*LL* males, thus resulting in differing maturation probabilities for males with alternative genotypes. We tested this hypothesis by comparing the testicular expression from immature individuals with different *vgll3* genotypes.

Using three different RT-ddPCR assays including two that tagged transcribed SNPs within the *vgll3* coding sequence, we observed a complex pattern of expression differences where association with *vgll3* genotype varied between different exons of the *vgll3* transcript. Using an assay spanning the boundary of exon 1 and 2, we observed no expression difference between *vgll3* genotypes (P=0.564, Fig. 3), whereas an assay spanning an SNP in exon 2 showed a 14% lower expression in *vgll3*EE* genotypes compared to *vgll3*LL* genotypes (91 versus 106 copies/ng i.e. 15 copies/ng fewer, 95% CI [3-28 fewer], two-sided *t*-test, df=31, P=0.014, n_{EE}=12 n_{LL}=22, Fig. 3). The opposite pattern was observed using an assay spanning an SNP in exon 3, where *vgll3*EE* genotypes showed 18.5% higher expression compared to *vgll3*LL* genotypes (97 versus 79 copies/ng i.e. 18 copies/ng more, 95% CI [0.08-36 more], two-sided *t*-test, df=25, P=0.049, n_{EE}=13 n_{LL}=23, Fig. 3). The other two tissues with highest *vgll3* expression levels, heart and adipose, did not show a clear relationship between *vgll3* expression level and genotype, maturation or age as assayed on the exon 2 assay (Fig. S7 & Fig. S8).

Figure 3. *Vgll3* expression differences in immature testes using three RT-ddPCR assays in different *vgll3* regions. Individuals with *vgll3*EE* (blue) and *vgll3*LL* (red) genotypes show varying levels of expression differences along the

vgll3 transcript. Points represent the mean of two replicates per individual sample (triangles; genotype means, *P*-values;
two-sided *t*-test). Grey lines connect RT-ddPCR assays with the approximate locations of their amplification targets on
the NCBI annotated *vgll3* transcript model with lines representing introns, rectangles representing exons.

Rare *vgll3* transcript isoforms in immature testes

The results from RT-ddPCR assays spanning three exons indicated that expression differences
between *vgll3* genotypes are restricted to specific exons of the *vgll3* transcript. This suggests that
vgll3 genotype may influence the structure of *vgll3* transcript being expressed, because conflicting
expression differences within a gene can be produced by confounding signal from multiple transcript
isoforms. We tested this by characterizing *vgll3* transcript structure in immature testes using 5' and
3' RACE (Rapid Amplification of cDNA Ends). Using pooled samples of five individuals for each
of the *vgll3*EE*, *vgll3*EL* and *vgll3*LL* genotypes, we observed that all genotypes expressed multiple
isoforms that differed in length 5' of exon 3 (Fig. S9). We Sanger sequenced of a total of 278 clones
to determine the structure of the expressed isoforms in *vgll3*EL* genotypes and observed at least six
distinct RACE fragments representing at least two distinct isoforms (from a total of 202 quality
sequences, Fig 4A). The longer isoform aligned with the NCBI annotated transcript with multiple
RACE fragments representing alternative lengths of the 5' UTR or exon 1. The shorter second
isoform was transcribed from an alternative 5' UTR within the NCBI annotated first intron with
translation likely initiated from start codons within exon 2. Further, Sanger sequencing revealed a
previously undetected C/T SNP in the alternative 5' UTR at position ssa25:28,655,795, where the
genotype was linked to the *vgll3*E/vgll3*L* genotype.

To further confirm the expression levels of *vgll3* isoforms, we re-analyzed publicly available RNA-
seq data from male gonads [34] using Cufflinks [38]. Combined RNA-seq data from three immature,
three prepubertal and three pubertal individuals that we identified as *vgll3*EL* heterozygotes
supported our discovery of *vgll3* isoforms with alternative transcription start sites. Twenty-two RNA-
seq reads mapped to an un-annotated region within intron 1 of the NCBI annotated transcript that
spliced into exon 2 with a canonical GT/AG 5'/3' (donor/acceptor) sequence at the splice site (Fig.
4B), and that corresponded to the alternative 5' UTR identified above. The abundance of the shorter
isoform accounted for 18% of the expression of the sum of *vgll3* isoforms (Cufflinks FPKM 0.56 out
of total of FPKM 0.56 + 2.52).

Taken together, based on varying expression across the exons of the *vgll3* transcript in conjunction
with the detection of multiple *vgll3* isoforms, we hypothesize that fish carrying *vgll3*EE* and
*vgll3*LL* genotypes may preferentially express alternative isoforms of *vgll3* that have different

transcription start sites, 5' UTRs and N-terminal protein sequences.

Figure 4. Characterization of a *vgll3* transcript isoform with an alternative 5' UTR. (A) 5' and 3' RACE sequencing

of pooled total RNA from *vgll3*EL* individuals identified an isoform with an alternative 5' UTR. Browser tracks of

Sanger sequencing show (from top) coverage, NCBI annotations, identified UTR, introns and intron junctions, and

representative RACE fragments with their alignment to the reference gene model (in light green). Highlighted region

shows gene annotations and Sanger sequencing tracks of a representative RACE fragment corresponding to the alternative

5' UTR containing isoform. **(B)** RNA-seq sequencing coverage and splice events from combined immature, pre-pubertal

and pubertal testes. Numbers correspond to RNA-seq reads spanning intron splice sites. Gene models depict transcripts

identified using Cufflinks [38].

***Cis*-regulatory differences in *vgll3* track alternative isoforms**

Expression of alternative transcript isoforms can be mediated by *cis* and/or *trans*-regulatory

mechanisms, where expression of isoforms is influenced by regulatory differences linked or unlinked,

respectively, to the genotype at the locus. We hypothesized that if the expression differences observed

between homozygote *vgll3*EE* and *vgll3*LL* individuals were due to *cis*-mediated isoform

expression differences, the differing expression levels in homozygotes would be conserved as allele-

specific expression within *vgll3*EL* heterozygotes. Expression level differences between the alleles

in heterozygotes can be accredited to differences in *cis*-regulation because both alleles are being

expressed in the same cellular *trans*-acting environment [39].

Using allele-specific assays for the newly discovered 5' UTR, exon 2 and exon 3, we observed that

heterozygotes showed allele-specific expression differences consistent with a largely *cis*-regulatory

basis of isoform expression differences. For the alternative 5' UTR, heterozygotes tended to express

the *vgll3*L* allele at a higher level compared to the *vgll3*E* allele, with on average 54% of mRNA

molecules representing from the *vgll3*L* allele. Using Poisson modeling of expression rates, 21 of

the 35 tested heterozygotes (60%) showed significantly higher *vgll3*L* expression at *P*-value

threshold of 0.05, while only five (14%) showed higher *vgll3*E* allele expression at the same *P*-value

threshold (Fig. 5A).

For exon 2, where we observed higher expression in *vgll3*LL* homozygotes compared to *vgll3*EE*

individuals, allele-specific expression showed a tendency towards higher expression of the *vgll3*L*

allele with on average 53% of the *vgll3* mRNA molecules expressed from the *vgll3*L* allele (Fig.

5A), versus 50.4% expected from heterozygous DNA used as control for differences in efficiencies

of allele-specific probes. Thirty-one of the 47 heterozygotes analyzed (66%) showed significantly

higher *vgll3*L* allele expression, while only one individual (2%) showed significantly higher *vgll3*E*

allele expression (*P*<0.05, Poisson rate-ratio test, Fig. 5A). In terms of expressed mRNA molecules,

the average magnitude of the *cis*-regulatory differences observed between *vgll3*E* and *vgll3*L* alleles

(6.8 mRNA molecules per ng of RNA) was large enough to account for 89% of the average expression

differences between *vgll3*EE* and *vgll3*LL* homozygotes (15.3 mRNA molecules, i.e.

$(2*6.8)/15.3*100=89\%$).

For exon 3, where *vgll3*LL* homozygotes showed lower expression compared to *vgll3*EE*, allele-

specific expression differences were also consistent with *cis*-regulation with on average 46.2% of

mRNA molecules expressed from the *vgll3*L* allele in *vgll3*EL* heterozygotes (compared to 49.4%

in heterozygous DNA control). Thirty-one of 35 tested heterozygotes (89%) showed significantly

lower expression of the *vgll3*L* allele (*P*<0.05). The average magnitude of the *cis*-regulatory

difference (7.5 mRNA molecules per ng of RNA) accounted for 83% of the average homozygote

expression difference (18 mRNA molecules, i.e. $(2*7.5)/18*100=83\%$).

For further validation of the *cis*-regulatory differences in *vgll3*, we re-analyzed publicly available

RNA-seq data from immature male gonads [34]. We identified three immature individuals as

heterozygous for the same *vgll3* SNPs we analyzed with our allele-specific RT-ddPCR assays and

quantified *vgll3* allele-specific expression based on RNA-seq reads overlapping the SNP positions.

Inspection of RNA-seq reads revealed the individuals to be heterozygous all of the three linked SNPs

in *vgll3* targeted by our RT-ddPCR assays (Fig. 5B). Consistent with our results from RT-ddPCR,

the newly discovered alternative 5' UTR showed strong allele-specific expression, being expressed

almost solely from the *vgll3*L* allele (8 versus 1 reads, *P*=0.038). The *vgll3*L* allele in exon 2 had

higher expression compared to the *vgll3*E* allele in all three heterozygote individuals (13 versus 8
reads, 10 versus 5 reads, 11 versus 10 reads), and when reads from all three males were combined
(42 reads versus 25 reads, binomial exact test *P*=0.049, Fig. 5B). For exon 3, the difference in *vgll3*L*
and *vgll3*E* alleles was non-significant (*P*=0.44).

These results show that the *vgll3* allele expression differences observed between alternative
homozygotes tend to be conserved in direction and in magnitude when assayed in heterozygotes. Re-
analysis of an independent data-set supports this conclusion. In sum, our results strongly support that
the differences in maturation timing between *vgll3* genotypes being driven by *cis*-regulated
expression differences of alternative *vgll3* isoforms.

Figure 5. Allele-specific expression of transcribed SNPs supports a *cis*-regulatory basis for differences in *vgll3*
isoform expression. (**A**) RT-ddPCR assays of *vgll3* allele-specific expression in heterozygotes show that alternative
homozygote expression differences are conserved within heterozygotes. For the alternative 5' UTR, exon 2 and exon 3,
on average 54%, 53% and 46% of *vgll3* mRNA molecules are derived from the *late* allele, respectively. For exon 2 and
exon 3 assay the null hypothesis for testing of 1:1 allele-specific expression ratio is calculated from heterozygote DNA
(dashed line, *N*=10). The color scale indicates significant (blue; *vgll3*E* allele more expressed, red; *vgll3*L* allele more
expressed) and non-significant (grey) allele-specific expression *P*-values (estimated from Poisson modelling of positive
RT-ddPCR droplets). (**B**) Re-analysis of an independent RNA-seq data set is consistent with the RT-ddPCR results and
shows allele-specific expression of an alternative 5' UTR and exon 2. Tracks represent (from top) RNA-seq coverage,
identified transcript models, and RNA-seq read coverage of SNPs. *P*-value are computed using a binomial test.

Discussion:

The molecular basis of variation in life history traits and their trade-offs have thus far remained
elusive because many life history traits are either polygenic, experimentally intractable, or their
genetic basis has only recently been discovered. Here, we show that variation in age at maturity, a
key life history trait, is associated with *cis*-regulatory differences in isoform expression of the
transcription co-factor gene *vgll3* in immature testes of Atlantic salmon males. Using targeted
expression assays, time series sampling in ten tissues and characterization of transcript structure, we
discovered that genotypes that tend to mature early versus those that tend to mature late express

different mixtures of *vgll3* transcript isoforms pre-puberty. The expression differences were allele-

specific in heterozygotes, suggesting that divergence in *cis*-regulation underlies the associations

between *vgll3* genotype, isoform expression and maturation probability.

We found strong evidence that *vgll3* genotypes associated with maturation age phenotypes do so

through a mechanism of *cis*-regulated differences in transcript isoform expression in immature testes.

The expressed *vgll3* isoforms differed in their transcription start sites, but are predicted to all produce

a functional *vgll3* protein. Alternative transcription start sites are thought to be important for

determining cell-types and their proliferation by influencing not only protein sequence, but also

through post-transcriptional regulation such as mRNA translation and stability [22,40,41]. Validating

the functional differences between *vgll3* isoforms and characterizing isoform expression patterns of

vgll3 in additional tissues will be an important avenue for future research. Expression of different

transcript isoforms has been established as the primary mechanism of important developmental

turning points in several other species. In many insects for example, alternatively spliced isoforms of

the gene *doublesex* act as sex determining factors [42]. Most species-specific differences in transcript

isoform expression have been linked to *cis*-regulatory differences [43,44]. *Cis*-regulatory changes that

affect alternative splicing are thought to reside within the un-spliced pre-mRNA [44]. For expression

from alternative transcription start sites such conservation of *cis*-regulatory placement has not been

shown, as initiation of transcription can be influenced by regulatory elements not only in the core

promoter, but also in distal enhancer sites [45]. Overall, genetic variation influencing *cis*-regulatory

loci is known to be an important source of phenotypic diversity and a common source of evolutionary

change [46–52]. The results reported here support the notion that this is also a strong candidate for the

molecular mechanisms underlying variation in complex life histories as well.

The transcription co-factor function of *vgll3* suggests that isoform changes in *vgll3* expression

transmit downstream to expression changes in gene regulatory networks associated with cell

proliferation and organ development [37]. Like its vertebrate and invertebrate homologs, the salmon

vgll3 has been proposed to control cell differentiation through interacting with the TEAD-family of

transcription factors of the Hippo signaling pathway [27,28]. Vgll3, like related co-factors in mice,

humans and *Drosophila*, is predicted to compete for TEAD accessibility with alternative co-factors

downstream of the Hippo signaling pathway controlling for organ size (Yap/Taz co-factors) [26,36].

Consistent with this, by analyzing RNA-sequencing data from immature, prepubertal and pubertal

testes, we found that *vgll3* and *yap1* showed opposite expression patterns with respect to maturity

status. Salmon *vgll3* expression was also found to negatively correlate with *yap1* expression in early

development [28]. We hypothesize that changes in *vgll3* isoform abundance associated with *vgll3*E*

and *vgll3*L* alleles may influence the tendency of cell differentiation in immature testes by changing interaction between Vgll3 and TEADs and/or Vgll3-Yap (Fig. 6). Clarifying the molecular relationships between *vgll3,* TEADs and *yap1* remain as important topics for future research.

Figure 6. Model for a molecular mechanism of *vgll3* genotype-dependent liability in puberty timing. Hypothetical distributions represent relative abundances of *vgll3* isoforms in individuals with given *vgll3* genotypes (blue, *vgll3*E*; red, *vgll3*L*). The association between *vgll3* genotype and maturation timing arises from *cis*-regulatory differences influencing *vgll3* isoform expression. The *vgll3*L* allele tends express more of the shorter isoform compared to the *vgll3*E* allele, thus changing the balance between *vgll3*-controlled differentiation versus self-renewal. TEAD interaction with *vgll3* isoforms tip the balance between tendency for self-renewal (higher expression of the short isoform) and differentiation (long isoform).

The relatively low expression level of *vgll3* and variation around genotype means also merit careful consideration of the limits of our study. First, some of the variation around the mean expression levels within *vgll3* genotypes can possibly be explained by additional genetic variation segregating in the experimental families and influencing *vgll3* expression in *cis* and in *trans*. Second, as the overall expression level of *vgll3* is very low, an alternative interpretation of the results could be that the small but statistically significant differences in expression across different exons of the *vgll3* transcript are driven by non-biological sources, such as noise in the measurement of low expression levels, or differences in assay efficiency and specificity. However, concurring results from re-analysis of an independent data-set, our careful control experiments, and the fact that the observed expression differences were specific to tissues and not observed in a systematic manner across all samples do not support this interpretation. The possibility exists also that *vgll3* is included in, but not determining the output of, the molecular pathway determining maturation timing, or that variation in the timing of measurement versus initiation of puberty causes variation in the genotype-phenotype association. Further functional validation through manipulation of isoform expression and allele-specific expression is an important future direction for validating the role of *vgll3* isoform-specific *cis*-

regulation in maturation timing.

It is important to note that although we observed a *cis*-regulatory differences associated with isoform

expression and maturation probability, it does not exclude the possibility that protein-coding changes

within the *vgll3* gene [4,5] could play additional functional roles in determining maturation timing.

Further, inferring an expression-genotype-phenotype association in male gonads does not exclude the

possibility that *vgll3* also has functional relevance in other tissues/processes (e.g. adipogenesis). In

fact, pleiotropic effects of single genes that are under the control of tissue and development-specific

regulatory sequences fits well with what we understand from the modularity of *cis*-regulation [46].

The expression pattern of *vgll3* and its paralog indeed support additional functions for the gene pair

in adipogenesis and female gonad development [28]. In addition, we observed relatively high *vgll3*

expression in heart as well as testes, which may indicate a pleiotropic action of *vgll3* in heart

development and function through the Hippo signaling pathway [53]. The role of changes in isoform

expression of *vgll3* in such pleiotropic effects warrants future research.

Although the effects of protein-coding changes on maturation phenotype were not addressed here,

support for the benefits of *cis*-regulatory changes compared to protein-coding changes in the

evolution of phenotypic diversity can be drawn from studies on morphological traits. Regulatory

changes often show higher modularity across tissues and development, additivity, and lower

pleiotropy, which are thought to inflict less evolutionary constraints compared to protein-coding

changes [46,54], especially in cases where proteins modulate the transcription of other genes as is the

case with *vgll3*. These features of evolvability of regulation are also well demonstrated here, where

we observed alternative expression dynamics (e.g. rising versus diminishing expression in heart and

liver, respectively, as individuals age), and genetic effects driving expression differences restricted

to one tissue and developmental stage (immature testes). Overall, these features are expected to lead

to *cis*-regulation being a more likely source of evolution in life history traits compared to coding

changes.

We have shown here that polymorphisms in the *vgll3* region associate significantly with one-year-

old male age at maturity in controlled conditions. The effect of *vgll3* genotype on maturation timing

in our experiment was arguably smaller compared to that observed in wild populations and for sea-

age at maturity [4]. Possible explanations for the divergent finding in the two studies can be related to

the differences in (*i*) traits (age at maturity versus probability of maturation), (*ii*) modeling approaches

(based on logit versus in our case probit distributions), (*iii*) wild versus husbandry environments, (*iv*)

differences in the assessment of maturity status, (*v*) variation in *vgll3* contribution to age at maturity

phenotypes across populations and (*vi*) possible differences in the genetic architecture of the distinct

age at maturity phenotypes (sea-age at maturity versus early male maturity).

Taken together, here we have revealed key functional insights into the temporal expression dynamics

and likely molecular mechanisms that translate genetic variation in a major effect locus to variation

in life history strategies in Atlantic salmon. Many life history traits are encoded by many genes where

it has remained challenging to identify the loci responsible for variation [55]. In the light of the results

presented here, gene regulatory evolution and isoform variation are likely candidates for mechanisms

mediating alternative life history strategies that merits further investigation across species.

Materials and Methods:

Crosses used in the study

We controlled-crossed eight 2x2 factorial matings of unrelated individuals that each included one *vgll3*EE* and one *vgll3*LL* male and female (see ref *13* for genotyping details), thereby resulting in 32 families with reciprocal *vgll3* genotypes *EE*, *EL*, *LE* and *LL* (allele order denotes female and male parents of origin respectively, Supplemental Fig. 1). We used parents from the "Oulujoki" broodstock, which is a mixture of several salmon populations from northern Baltic rivers and is maintained by the Natural Resources Institute Finland (LUKE) [56]. All parents were selected for having a homogenous age-at-maturity-associated haplotype encompassing the *vgll3* (chromosome 25 paralog) coding sequence, the top non-coding association [4], and the coding region of the adjacent *akap11* gene (collectively referred to as "*vgll3* genotype").

Husbandry design and conditions

Following fertilization in October 2017, familial egg batch replicates were raised in two egg incubators until first feeding (~ 2 months post-hatching) with mesh-separators to contain families in individual compartments. Families were assigned to compartments randomly. At first feeding, familial batches were combined and 48 individuals per family were randomly selected and distributed evenly across eight 0.25 m^3 tanks using recirculated water i.e. on average, each tank included six individuals per family. Prior to tank allocation, each individual was tagged with visible elastomer (VIE) at the base of the caudal fin to enable identification of the *vgll3* genotype of an individual by a fluorescent color (initial *n*=1300). Light cycle and water temperature (min=6.3, max=17.7 °C) during husbandry corresponded to the local seasonal cycle at 61°N. Fish were fed first with live *Artemia* for three days, after which we fed commercial aquaculture feed *ad libitum* (Raisio Baltic Blend) with pellet diameter increasing over time according to fish size. Temperature, oxygen and nitrogen-component levels were monitored on a regular basis. Experimentation was conducted under animal experimentation license ESAVI/2778/2018.

Individual tagging, family assignment and morphological measurements

At eight months post-hatching and before the onset of the first breeding season we tagged remaining fish using passive integrated transponder (PIT) tags and collected fin-clips. We extracted DNA using standard chelex or salt extraction methods, genotyped all fish for 141 informative SNPs [57], assigned their genotypic sex based on genotyping coverage of the salmon sex determining locus *sdy* and assigned individuals to families using the likelihood method implemented in SNPPIT [58]. We recorded the length and mass of all fish monthly after PIT-tagging. We non-lethally assessed the maturity status of all males by testing for the presence of running milt at nine months post-hatching.

Morphological measurement and non-lethal maturation assessment were performed in a random order
such that the assessor did not have knowledge of *vgll3* genotype.

Statistical model of male maturation probability

We modeled the probability of males reaching maturity within their first year given their *vgll3*
genotypes by using a GLMM (generalized linear mixed model) approach to implement a genetic
threshold model that takes full account of the relatedness among individuals [59,60]. We estimated
maturation probability using a generalized linear animal model with probit-link function under
Bayesian Markov Chain Monte Carlo simulations with the R-software [61] version 3.6.1 and package
MCMCglmm [62] for the following model: $y = vgll3 + length + a + t + e$, where y is a vector of
maturation binaries, *length* is mean-centered and variance scaled natural logarithm of length (cm),
vgll3 is the genotype, *a* is the additive genetic relationship-matrix predicted animal effect [63], *t* is the
common environmental tank effect and *e* the error term (with fixed variance of 1). We used priors
following a χ_1^2 distribution [64] and based inferences on 10,000 samples collected every 1,000
iteration after burn-in of 10,000. Model convergence and adequate sampling was ensured by visual
trace plot inspection, autocorrelation < 0.1 for lag-2 parameter samples, and using Heidelberger and
Welch's convergence diagnostic [65]. We transformed maturation probabilities from probit-scale to
normal probability scale for ease of interpretation. Heterozygote *vgll3*LE* and *vgll3*EL* (order of
alleles denotes maternal origin) were considered as one group as initial analysis indicated that
maturation probabilities between these genotypes were not significantly different.

Temporal sampling, RNA extraction and RT-ddPCR

We sampled individuals with *vgll3*EE, vgll3*EL, vgll3*LE* and *vgll3*LL* monthly (3-32 in total per
genotype per time point) and assessed sex, gonad size and maturity by dissection. We sacrificed the
animals by MS-222 overdose and dissected ten tissues (brain (timepoints 12-13 months post hatch);
heart, liver, muscle (timepoints 3-13 months post hatch), adipose (timepoints 5-13 months post
hatch); immature ovary (timepoints 6-13 months post hatch); immature testes (timepoints 7-13
months post hatch); mature testes (timepoints 8 and 10-12 months post hatch); stripped mature testes
(timepoints 12-13 months post hatch) and ejaculate (timepoint 13 months post hatch)). As immature
male gonads had no recordable weight on a scale with a detection limit of 10 mg thus preventing the
use of gonadal-somatic index, we classified each sacrificed male fish visually as immature (not
enlarged, transparent color gonads), pubertal (slightly enlarged, transparent or white color gonads) or
mature (enlarged white color gonads). We flash froze the samples on liquid nitrogen and stored the
samples in -80 °C. We homogenized the tissues using an OMNI Bead Ruptor Elite machine, in 2 ml
or 7 ml (for mature gonads) with 2.4 mm steel beads. We used Macherey-Nagel NucleoSpin RNA

kits in single and in 96-well plate formats to extract total RNA and treated the samples with additional

DNase (Invitrogen Turbo Dnase) to remove any residual DNA. We verified total RNA quality using

Agilent BioAnalyzer for a random set of 24 samples and found all samples to have RIN>9. We

measured total RNA yield using ThermoFischer Qubit and Quant-it reagents, diluted the samples to

working concentrations and re-measured RNA concentrations. For samples in Figs. 2 and 3 (all gonad

samples) we repeated the RNA concentration measurement once more and calculated the mean of the

duplicate measurements, which we used in data normalization. We used ~20 ng (2-8 months post-

hatch) or ~50 ng (8-14 months post-hatch) of total RNA as template in one-step reverse transcription

droplet digital PCR (RT-ddPCR) [66] (BioRad) with TaqMan probes specific to *vgll3* alleles to

quantify *vgll3* absolute expression (Supplementary table 1). Measurement of *amh* and *igf3* expression

was performed as above, with the exception of using ~2 ng of total RNA template and TaqMan probes

that were labeled with alternative fluorophores and assayed in the same RT-ddPCR reactions. We

verified the specificity of *vgll3*-genotype specific TaqMan assays with genomic and synthetic DNA

templates (Fig. S10) and optimized melting temperatures for all assays using a gradient. Genomic

DNA could not be used as control for the *vgll3* 5' UTR assay because the assay spans an intron-exon

boundary. Genomic and synthetic DNA controls and null control were included in each RT-ddPCR

vgll3 run, while genomic DNA and null control were included in *Amh* and *Igf3* RT-ddPCR runs.

Expression analyses

We analyzed RT-ddPCR results using BioRad QuantaSoft software. We manually set minimum

signal thresholds for *vgll3*L* and *vgll3*E* channels because visual inspection of RT-ddPCR signal

indicated that, for *vgll3* exon 2 assay, the *vgll3*L* specific allele showed low background signal on

the *vgll3*E* channel (Fig. S10). We excluded samples that failed *vgll3* RT-ddPCR reaction using a

pre-determined minimum threshold of 10,000 quality-filter passed droplets to include only samples

where expression could be estimated with low error. We excluded *amh/igf3* expression data from

samples where RT-ddPCR signal was saturated (over 98% of droplets were positive) because mRNA

abundances cannot be accurately estimated with saturated RT-ddPCR data. We excluded seven

samples where the genotypic sex was inconsistent with morphological sex of the gonads. We

excluded three mis-labeled individuals with inconsistent *vgll3* genotypes and allelic expression (e.g.

vgll3*EE homozygotes with expression signal only from *vgll3*LL* specific probes). We removed two

heterozygote samples that showed extreme monoallelic expression of the *vgll3*L* allele (fractional

abundance over 90%), possibly representing labeling or sampling error. We then used custom R

scripts to normalize the number of *vgll3* mRNA molecules to the amount of input RNA and analyze

the data and plot the results (available upon acceptance).

For investigation of the effect of age/season and *vgll3* genotype to *vgll3* expression in immature testes

and immature ovaries, we used the following regression model: $y = vgll3 + x + e$, where y is *vgll3*

total mRNA copies, x is days post-hatch, *vgll3* is the genotype and e the error term. We tested the

effect of *vgll3* and x (days post-hatch) individually using a corresponding model.

We characterized the relative expression levels of the key male maturity genes *amh* and *igf3* by way

of Poisson modeling of expression rate-ratio for each sample. The number of positive droplets with

expression signal in RT-ddPCR data follows a Poisson distribution with lambda being the rate of

positive   droplets   over   all   droplets   ([66],   https://www.bio-

rad.com/webroot/web/pdf/lsr/literature/Bulletin_6407.pdf). We calculated the Poisson rates of expression from

amh and *igf3* -positive droplets over the total amount of analyzed droplets (the rate ratio of *amh/igf3*

expression) using the R function *poisson.test*.

We tested for presence of distinct clusters of gene expression patterns on *amh/igf3* and *vgll3*

expression space using Gaussian mixture modeling as implemented in the R package *Mclust* version

5 [35]. We chose the best fitting mixture of Gaussian distributions based on Bayesian Information

Criterion using the function *mclustBIC* and further tested the best fit mixture model for the presence

of more than one distributions using a bootstrapping likelihood-ratio test as implemented in the

function *mclustBootstrapLRT* with 1000 bootstrap replications.

RACE

To perform RACE we followed the manufacturer's instructions using Takara SMARTer 5'/3' RACE-

kit with the following modifications. We pooled equal amounts (25 ng) of immature testis total RNA

from six individuals per genotype with either *vgll3*EE, vgll3*EL* or *vgll3*LL* genotypes to create

three genotype specific pools, and performed RACE with primers aligning to exon 3 of the *vgll3*

transcript. After analysis on agarose gel, we purified the RACE PCR products using Macherey-Nagel

NucleoMag DNA binding beads, cloned the PCR products to a pRACE vector using the SMARTer

kit reagents, transformed the plasmids to *E. coli* and grew the transformation at 37 degrees C o/n on

LB-ampicillin (100 mg/ml) plates. Sanger sequencing of transformed PCR products was performed

as described in ref [67]. Briefly, colonies were picked into 96 deep-well plates containing 1 ml growth

media and incubated in a shaker o/n. An aliquot (150 ul) was transfered to a PCR plate and spin down.

The pellet was dissolved in 150 water and heated to 95 C for 10 min and 2 ul was used as template

in PCR with universal and reverse primers (UP-47 and RP-48). The obtained PCR products were

sequenced with UP and RP universal primers using BigDye Chemistry v 3 and analyzed on ABI

3130xl Sequencer (Life Technologies) at the DNA Sequencing and Genomics lab Institute of

Biotechnology University of Helsinki. We used *Geneious Prime 2020.1.2* to automatically detect and

trim the sequences of vector DNA and low-quality sequences above 1% base error rate. We aligned

the sequences to the *vgll3* region downloaded from NCBI using the *Geneious RNA* mapper with the

highest sensitivity settings and allowing for novel intron detection.

Allele-specific expression

Testing of *vgll3* allele-specific expression in heterozygotes was performed in a similar manner as

above for *amh/igf3* expression (test for Poisson rate ratio), with the exception that signal from positive

droplets now represented expression from either the *vgll3*E* or the *vgll3*L* allele. We defined the

expression rate of *vgll3*E* allele as the number of droplets positive for the *vgll3*E* probes over all

probes that passes the QuantaSoft standard quality filter (including empty droplets), and

correspondingly for the *vgll3*L* allele. We then tested for difference in Poisson rates for each

individual heterozygote using the R function *poisson.test* and *P*-value threshold of 0.05. We

calculated the null expectation for equal expression rates based on RT-ddPCR with 10 replicates of

vgll3 heterozygous DNA as template. The assay for the alternative 5' UTR spanned an intron-exon

boundary, and therefore we could not use genomic DNA as control (the assay does not amplify with

DNA as template). We assumed 50% efficiency for this assay.

We estimated that on average the *cis*-regulatory difference in *vgll3* allele-specific expression can

account for 89% of the average expression difference between *vgll3*EE* and *vgll3*LL* homozygotes

in exon 2 and 83% in exon 3. We estimated the 89% for exon 2 as follows (analogous calculation for

exon 3 not shown): On average *vgll3*EL* individuals expressed 113 *vgll3* mRNA molecules (per ng

of input RNA, Fig 3A). Of these, on average, 53.5 molecules were expressed from the *vgll3*E* allele

and 60.3 molecules from the *vgll3*L* allele, giving an average *vgll3*L* relative abundance of 53%

(60.3/(53.5+60.3)*100=53%). From above we can see that average difference between *vgll3*E* and

vgll3*L mRNA counts is 6.8 mRNA molecules (60.3-53.5=6.8). Corresponding, average difference

between *vgll3*EE* and *vgll3*LL* genotypes is 15.3 mRNA molecules (106.35-91.06=15.3). The

percentage of *vgll3*EE*- *vgll3*LL* difference accounted by *vgll3*E*- *vgll3*L cis*-expression difference

is thus (2*6.8)/15.3*100=89%, as *vgll3*EE* and *vgll3*LL* have two of either allele each.

RNA-seq analyses

We re-analyzed RNA-sequencing data from immature, prepubertal and pubertal Atlantic salmon

testis samples [34] in order to investigate the relative expression patterns of *vgll3* and *yap1* genes. We

downloaded raw RNA-seq data for immature (SRR8479243, SRR8479245, SRR8479246),

prepubertal (SRR8479244, SRR8479249, SRR8479250) and pubertal (SRR8479242, SRR8479248,

SRR8479247) testes from the *Sequence Read Archive* and filtered the reads for quality using *Fastp*

and default parameters [68].We created an alignment reference with *STAR* [69] and the parameter –

runMode genomeGenerate, using the Atlantic salmon reference genome and annotation files

downloaded from *NCBI* [70]. We aligned and quantified the quality-filtered reads using *STAR* in

manual   two-pass   mode   and   with   the   following   parameters: *--outFilterIntronMotifs*

RemoveNoncanonicalUnannotated   –chimSegmentMin   10   –outFilterType   BySJout   –

alignSJDBoverhangMin 1 –alignIntronMin 20 –alignIntronMax 1000000 –alignMatesGapMax

1000000 –quantMode GeneCounts –alignEndsProtrude 10 ConcordantPair --limitOutSJcollapsed

5000000. We combined the read counts from *STAR* into a single table using a custom *R* script and

normalized the data using *DESeq2 varianceStabilizingNormalization* [71] and independently with

Clust [72] (automatic normalization method: 101 3 4), so that expression patterns of genes could be

compared independently from their total expression level. Last, we used custom scripts in *R* to

analyze the co-expressed genes and plot the results. *NCBI RefSeq* annotations for genes were fetched

using *annotationhub* package and Gene Ontology over-representation in co-expression groups was

tested using *clusterProfiler* package [73]. Reads from isoform variants were quantified using

ggsashimi [74]

We quantified *vgll3* allele-specific expression in the RNA-seq data by comparing reads overlapping

SNP positions Ssa25: 28655795, Ssa25:28656101 and Ssa25:28658151 (the two latter being same

SNPs used in our crossing scheme and allele-specific TaqMan assays). We changed the SNP

nucleotide to "N" in order to remove reference-biased mapping and re-aligned the RNA-seq reads

using *STAR,* excluding multimapping reads and reads with more than one mis-match by defining

parameters *-outFilterMultimapNmax 1 -outFilterMismatchNmax 1* (in addition to the parameters

defined above). We then counted the number of overlapping *vgll3*L* and *vgll3*E* reads by manual

inspection of RNA-seq read tracks using *IGV* [75] and tested the significance of allele-specific

expression to the null expectation of 1:1 expression with binomial test.

Acknowledgements:

We thank members of Primmer lab for assistance with the common garden experiment, Lars Paulin for assistance in RACE and Sanger sequencing, and LUKE staff for assistance with the initial crosses. Financial support was provided by the University of Helsinki, Academy of Finland (grant number 314254) and the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement number 742312).

Author contributions:

- JPV and CRP designed the time-series sampling and expression experiments.
- JPV implemented and supervised the time-series sampling and expression analyses, designed and optimized the method, analyzed the data and wrote the initial draft.
- PVD and CRP designed the experiments validating *vgll3* association with male maturation probability.
- PVD implemented and supervised validation of *vgll3* association with male maturation probability and developed the associated statistical method.
- AR performed supporting lab analyses and optimized the genotyping method.
- NPi, AR, OO, JMV, ST and NPa assisted with fish rearing, tagging and sampling.
- TA designed the genotyping method.
- JE provided initial material for crosses.
- CRP conceived and supervised the project and acquired the funding.
- JPV and CRP wrote the final manuscript with input from PVD and JMV.
- All authors approved the manuscript.

Data availability:

Data will be made available on Dryad database following manuscript acceptance.

Code availability:

Code will be made available on GitHub following manuscript acceptance.

719 **Supplementary figures.**

722 **Figure S1.** 16 females with 16 males were crossed in a 2x2 factorial design, producing 32 families

723 with *vgll3* genotypes *EE, EL, LE* and *LL* (order denotes maternal and paternal origin of alleles).

726 **Figure S2.** Maturation frequencies at nine months post-hatching. Number above bars indicates the

727 number of mature individuals over all individuals.

732 **Figure S3.** *Vgll3* expression in all investigated tissues and whole fry. Stripped testes; mature testes

733 with ejaculate manually removed.

736 **Figure S4.** *Vgll3* expression in liver shows a linear downward trend, but the overall expression level

737 is low. Green line, loess smoothing; grey shading, 95% confidence interval.

Figure S5. Expression of puberty indicator genes in male testes. Average *amh/igf3* expression ratio
is higher in immature testes and maturation coincides with relative down-regulation of *amh* in relation
to *igf3*. Pubertal samples show large variation in *amh/igf3* ratio. Testes are separated into categories
based on morphology (see methods).

Figure S6. Re-analysis of data from ref [34] Expression patterns of *vgll3* and *yap1* in immature, prepubertal and pubertal testes. (A) Points represent normalized expression levels (variance stabilizing normalization [71]). (B) Points represent the relative normalized expression level means [72] from three biological replicates in each case. (C) Relative expression patterns of 7074 genes in *vgll3* co-expression cluster. Black points; *vgll3*, grey points; *yap1*.

Figure S7. Expression of *vgll3* in the heart. All trendlines non-significant at *P*-value threshold 0.05.

Figure S8. Expression of *vgll3* in adipose tissue. All trendlines non-significant at *P*-value threshold 0.05.

Figure S9. Agarose gel of 5' and 3' RACE PCR reactions. Both 5' and 3' RACE reactions amplify multiple RACE fragments of different lengths. RACE was performed using primers aligning to exon 3 of the *vgll3* transcript.

Figure S10. Genomic DNA and synthetic DNA controls of allele-specific *vgll3* assays.

768 **Supplemental tables:**

770 **Table S1.** TaqMan assays used in the study.

Vgll3 5' UTR Forward Primer	*Vgll3* 5' UTR Reverse Primer	*Vgll3* 5' UTR *Late* probe	*Vgll3* 5' UTR *Early* probe
GTCGTCATACTTTCACCAGAACACA	TGTAAACACCCAGCTTCTTCGAA	AAAAGTTGTAGGACGTTTC	AAAAGTTGTAGAACGTTTC

Vgll3 Exon 2 Forward Primer	*Vgll3* Exon 2 Reverse Primer	*Vgll3* Exon 2 *Early* probe	*Vgll3* Exon 2 *Late* probe
GAAGCTGGGTGTTTACAGTAGGAT	TGTCTCCGCCCTGGAAAC	TGCTGCTCCATGCTGT	TGCTGCTCCGTGCTGT

Vgll3 Exon 3 Forward Primer	*Vgll3* Exon 3 Reverse Primer	*Vgll3* Exon 3 *Early* probe	*Vgll3* Exon 3 *Late* probe
AGCCCAGGGATACACAGTGA	GTGGGCCAGGCTGAGG	CCACCTCTGTGTTCACA	CCACCTCTGTCTTCACA

Amh Forward Primer	*Amh* Reverse Primer	*Amh* probe
ATGAAAGAGTGACAGAAAAGCTGAGT	TGCAGCTCACACAGAAACGT	CCCTCCCCTGCTCCC

Igf3 Forward Primer	*Igf3* Reverse Primer	*Igf3* probe
CAACAAACTGAGATGAACAGTCTGAAC	CAGTCGCACAGCCAAGAATAAAA	TCGGCACAAAATATCT

References:

1. Stearns SC. The evolution of life histories. Oxford University Press; 1992.

2. Stearns SC. Life history evolution: successes, limitations, and prospects. Die Naturwissenschaften. 2000;87: 476–486. doi:10.1007/s001140050763

3. Johnston SE, Gratten J, Berenos C, Pilkington JG, Clutton-Brock TH, Pemberton JM, et al. Life history trade-offs at a single locus maintain sexually selected genetic variation. Nature. 2013;502: 93–95. doi:10.1038/nature12489

4. Barson NJ, Aykanat T, Hindar K, Baranski M, Bolstad GH, Fiske P, et al. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. Nature. 2015;528: 405–408. doi:10.1038/nature16062

5. Ayllon F, Kjaerner-Semb E, Furmanek T, Wennevik V, Solberg MF, Dahle G, et al. The vgll3 locus controls age at maturity in wild and domesticated Atlantic salmon (Salmo salar L.) males. PLoS Genetics. 2015;11. doi:10.1371/journal.pgen.1005628

6. Prince DJ, O'Rourke SM, Thompson TQ, Ali OA, Lyman HS, Saglam IK, et al. The evolutionary basis of premature migration in Pacific salmon highlights the utility of genomics for informing conservation. Science advances. 2017;3: e1603198. doi:10.1126/sciadv.1603198

7. Narum SR, Genova AD, Micheletti SJ, Maass A. Genomic variation underlying complex life-history traits revealed by genome sequencing in Chinook salmon. Proc Biol Sci. 2018;285. doi:10.1098/rspb.2018.0935

8. Hess JE, Zendt JS, Matala AR, Narum SR. Genetic basis of adult migration timing in anadromous steelhead discovered through multivariate association testing. Proc Biol Sci. 2016;283: 20153064. doi:10.1098/rspb.2015.3064

9. Troth A, Puzey JR, Kim RS, Willis JH, Kelly JK. Selective trade-offs maintain alleles underpinning complex trait variation in plants. Science. 2018;361: 475–478. doi:10.1126/science.aat5760

10. Lamichhaney S, Fan G, Widemo F, Gunnarsson U, Thalmann DS, Hoeppner MP, et al. Structural genomic changes underlie alternative reproductive strategies in the ruff (Philomachus pugnax). Nat Genet. 2016;48: 84–88. doi:10.1038/ng.3430

11. Heyland A, Flatt T. Mechanisms of Life History Evolution : The Genetics and Physiology of Life History Traits and Trade-Offs. OUP Oxford; 2011.

12. Fleming IA, Einum S. Reproductive Ecology: A Tale of Two Sexes. Wiley-Blackwell; 2010. pp. 33–65. doi:10.1002/9781444327755.ch2

13. Mobley KB, Granroth-Wilding H, Ellmen M, Vaha J-P, Aykanat T, Johnston SE, et al. Home ground advantage: Local Atlantic salmon have higher reproductive fitness than dispersers in the wild. Science advances. 2019;5: eaav1112. doi:10.1126/sciadv.aav1112

14. Fleming IA. Pattern and variability in the breeding system of Atlantic salmon (Salmo salar), with comparisons to other salmonids. Canadian Journal of Fisheries and Aquatic Sciences. 1998;55: 59–76.

15. Czorlich Y, Aykanat T, Erkinaro J, Orell P, Primmer CR. Rapid sex-specific evolution of age at maturity is shaped by genetic architecture in Atlantic salmon. Nature Ecology & Evolution. 2018;2: 1800–1807. doi:10.1038/s41559-018-0681-5

16. Cousminer DL, Berry DJ, Timpson NJ, Ang W, Thiering E, Byrne EM, et al. Genome-wide association and longitudinal analyses reveal genetic loci linking pubertal height growth, pubertal timing and childhood adiposity. Human Molecular Genetics. 2013;22: 2735–2747. doi:10.1093/hmg/ddt104

17. Perry JRB, Day F, Elks CE, Sulem P, Thompson DJ, Ferreira T, et al. Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. Nature. 2014;514: 92-+. doi:10.1038/nature13545

18. Pearse DE, Barson NJ, Nome T, Gao G, Campbell MA, Abadía-Cardoso A, et al. Sex-dependent dominance maintains migration supergene in rainbow trout. Nature Ecology & Evolution. 2019;69: 1–12. doi:10.1038/s41559-019-1044-6

19. Lamichhaney S, Fuentes-Pardo AP, Rafati N, Ryman N, McCracken GR, Bourne C, et al. Parallel adaptive evolution of geographically distant herring populations on both sides of the North Atlantic Ocean. Proceedings of the National Academy of Sciences. 2017;54: 201617728-E3461. doi:10.1073/pnas.1617728114

20. Bateman JR, Johnson JE, Locke MN. Comparing enhancer action in cis and in trans. Genetics. 2012;191: 1143–1155. doi:10.1534/genetics.112.140954

21. Tian K, Henderson RE, Parker R, Brown A, Johnson JE, Bateman JR. Two modes of transvection at the eyes absent gene of Drosophila demonstrate plasticity in transcriptional regulatory interactions in cis and in trans. Bosco G, editor. PLoS Genetics. 2019;15: e1008152. doi:10.1371/journal.pgen.1008152

22. Reyes A, Huber W. Alternative start and termination sites of transcription drive most transcript isoform differences across human tissues. Nucleic Acids Res. 2018;46: 582–592. doi:10.1093/nar/gkx1165

23. Wang X, Hou J, Quedenau C, Chen W. Pervasive isoform-specific translational regulation via alternative transcription start sites in mammals. Mol Syst Biol. 2016;12: 875. doi:10.15252/msb.20166941

24. Wiestner A, Schlemper RJ, Maas A van der, Skoda RC. An activating splice donor mutation in the thrombopoietin gene causes hereditary thrombocythaemia. Nat Genet. 1998;18: 49–52. doi:10.1038/ng0198-49

25. Halperin DS, Pan C, Lusis AJ, Tontonoz P. Vestigial-like 3 is an inhibitor of adipocyte differentiation. Journal of Lipid Research. 2013;54: 473–481. doi:10.1194/jlr.m032755

26. Figeac N, Mohamed AD, Sun C, Schönfelder M, Matallanas D, Garcia-Munoz A, et al. VGLL3 operates via TEAD1, TEAD3 and TEAD4 to influence myogenesis in skeletal muscle. Journal of cell science. 2019;132: jcs225946. doi:10.1242/jcs.225946

27. Kjaerner-Semb E, Ayllon F, Kleppe L, Sørhus E, Skaftnesmo K, Furmanek T, et al. Vgll3 and the Hippo pathway are regulated in Sertoli cells upon entry and during puberty in Atlantic salmon testis. Scientific Reports. 2018;8: 1912. doi:10.1038/s41598-018-20308-1

28. Kurko J, Debes PV, House A, Aykanat T, Erkinaro J, Primmer CR. Transcription Profiles of Age-at-Maturity-Associated Genes Suggest Cell Fate Commitment Regulation as a Key Factor in the Atlantic Salmon Maturation Process. G3 (Bethesda). 2019; g3.400882.2019. doi:10.1534/g3.119.400882

29. Taranger GL, Carrillo M, Schulz RW, Fontaine P, Zanuy S, Felip A, et al. Control of puberty in farmed fish. General and Comparative Endocrinology. 2010;165: 483–515. doi:10.1016/j.ygcen.2009.05.004

30. Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE, Parre N, et al. Large single-locus effects for maturation timing are mediated via condition variation in Atlantic salmon. bioRxiv. 2019;11: 780437. doi:10.1101/780437

31. Schulz RW, Franca LR, Lareyre J-J, LeGac F, Chiarini-Garcia H, Nobrega RH, et al. Spermatogenesis in fish. General and Comparative Endocrinology. 2010;165: 390–411. doi:10.1016/j.ygcen.2009.02.013

32. Pfennig F, Standke A, Gutzeit HO. The role of Amh signaling in teleost fish--Multiple functions not restricted to the gonads. General and Comparative Endocrinology. 2015;223: 87–107. doi:10.1016/j.ygcen.2015.09.025

33. Morais R, Crespo D, Nobrega RH, Lemos MS, Kant HJG van de, Franca LR, et al. Antagonistic regulation of spermatogonial differentiation in zebrafish (Danio rerio) by Igf3 and Amh. Molecular and Cellular Endocrinology. 2017;454: 112–124. doi:10.1016/j.mce.2017.06.017

34. Skaftnesmo KO, Edvardsen RB, Furmanek T, Crespo D, Andersson E, Kleppe L, et al. Integrative testis transcriptome analysis reveals differentially expressed miRNAs and their mRNA targets during early puberty in Atlantic salmon. BMC Genomics. 2017;18: 801. doi:10.1186/s12864-017-4205-5

35. Scrucca L, Fop M, Murphy TB, Raftery AE. mclust 5: Clustering, classification and density estimation ssing Gaussian finite mixture models. The R journal. 2016;8: 289–317. doi:10.1186/s12942-015-0017-5

36. Meng Z, Moroishi T, Guan K-L. Mechanisms of Hippo pathway regulation. Genes Dev. 2016;30: 1–17. doi:10.1101/gad.274027.115

37. Simon E, Faucheux C, Zider A, Theze N, Thiebaud P. From vestigial to vestigial-like: the Drosophila gene that has taken wing. Development Genes and Evolution. 2016;226: 297–315. doi:10.1007/s00427-016-0546-3

38. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, et al. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nat Protocols. 2012;7: 562–578. doi:10.1038/nprot.2012.016

39. Wittkopp PJ, Haerum BK, Clark AG. Evolutionary changes in cis and trans gene regulation. Nature. 2004;430: 85–88.

40. Sendoel A, Dunn JG, Rodriguez EH, Naik S, Gomez NC, Hurwitz B, et al. Translation from unconventional 5′ start sites drives tumour initiation. Nature. 2017;541: 494–499.

41. (DGT) TFC and the RP and C. A promoter-level mammalian expression atlas. Nature. 2014;507: 462–470. doi:10.1038/nature13182

42. Burtis KC, Baker BS. Drosophila doublesex gene controls somatic sexual differentiation by producing alternatively spliced mRNAs encoding related sex-specific polypeptides. Cell. 1989;56: 997–1010. doi:10.1016/0092-8674(89)90633-8

43. Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, et al. The Evolutionary Landscape of Alternative Splicing in Vertebrate Species. Science. 2012;338: 1587–1593. doi:10.1126/science.1230612

44. Gao Q, Sun W, Ballegeer M, Libert C, Chen W. Predominant contribution of cis-regulatory divergence in the evolution of mouse alternative splicing. Mol Syst Biol. 2015;11: 816. doi:10.15252/msb.20145970

45. Smale ST, Kadonaga JT. The RNA polymerase II core promoter. Annu Rev Biochem. 2003;72: 449–479.

46. Carroll SB. Evo-devo and an expanding evolutionary synthesis: A genetic theory of morphological evolution. Cell. 2008;134: 25–36. doi:10.1016/j.cell.2008.06.030

47. Schwartz C, Balasubramanian S, Warthmann N, Michael TP, Lempe J, Sureshkumar S, et al. Cis-regulatory Changes at FLOWERING LOCUS T Mediate Natural Variation in Flowering Responses of Arabidopsis thaliana. Genetics. 2009;183: 723–732. doi:10.1534/genetics.109.104984

48. Chan YF, Marks ME, Jones FC, Villarreal G, Shapiro MD, Brady SD, et al. Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer. Science. 2010;327: 302–305.

49. Frankel N, Erezyilmaz DF, McGregor AP, Wang S, Payre F, Stern DL. Morphological evolution caused by many subtle-effect substitutions in regulatory DNA. Nature. 2011;474: 598–603. doi:10.1038/nature10200

50. O'Brown NM, Summers BR, Jones FC, Brady SD, Kingsley DM. A recurrent regulatory change underlying altered expression and Wnt response of the stickleback armor plates gene EDA. Elife. 2015;4. doi:10.7554/elife.05290

51. Indjeian VB, Kingman GA, Jones FC, Guenther CA, Grimwood J, Schmutz J, et al. Evolving new skeletal traits by cis-regulatory changes in mone morphogenetic proteins. Cell. 2016;164: 45–56. doi:10.1016/j.cell.2015.12.007

52. Gautier M, Yamaguchi J, Foucaud J, Loiseau A, Ausset A, Facon B, et al. The genomic basis of color pattern polymorphism in the harlequin ladybird. Current Biology. 2018;28: 1–7. doi:10.1016/j.cub.2018.08.023

53. Wang J, Liu S, Heallen T, Martin JF. The Hippo pathway in the heart: pivotal roles in development, disease, and regeneration. Nature Reviews Cardiology. 2018;15: 672–684. doi:10.1038/s41569-018-0063-3

54. Stern DL, Orgogozo V. The loci of evolution: how predictable is genetic evolution? Evolution.
2008;62: 2155–2177. doi:10.1111/j.1558-5646.2008.00450.x

55. Price T, Schluter D. ON THE LOW HERITABILITY OF LIFE-HISTORY TRAITS. Evolution.
1991;45: 853–861. doi:10.1111/j.1558-5646.1991.tb04354.x

56. Erkinaro J, Laine A, Maki-Petays A, Karjalainen TP, Laajala E, Hirvonen A, et al. Restoring
migratory salmonid populations in regulated rivers in the northernmost Baltic Sea area, Northern
Finland - biological, technical and social challenges. Journal of Applied Ichthyology. 2011;27: 45–
52. doi:10.1111/j.1439-0426.2011.01851.x

57. Aykanat T, Lindqvist M, Pritchard VL, Primmer CR. From population genomics to
conservation and management: a workflow for targeted analysis of markers identified using
genome-wide approaches in Atlantic salmon Salmo salar. Journal of fish biology. 2016;89: 2658–
2679. doi:10.1111/jfb.13149

58. Anderson EC. Computational algorithms and user-friendly software for parentage-based tagging
of Pacific salmonids. 2010.

59. Sorensen DA, Genetics SA, 1995. Bayesian inference in threshold models using Gibbs
sampling. Genetics Selection Evolution. 1995;27: 229–249.

60. Hadfield JD. Increasing the efficiency of MCMC for hierarchical phylogenetic models of
categorical traits using reduced mixed models. O'Hara RB, editor. Methods in Ecology and
Evolution. 2015;6: 706–714. doi:10.1111/2041-210x.12354

61. Team RC. R: A Language and Environment for Statistical Computing. 2013;

62. Hadfield J. MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. Journal of Statistical Software. 2010;33: 1–22.

63. Henderson C. Sire evaluation and genetic trends. Journal of Animal Ecology. 1973; 10–43.
doi:10.1093/ansci/1973.symposium.10

64. Villemereuil P de, Gimenez O, Doligez B. Comparing parent–offspring regression with
frequentist and Bayesian animal models to estimate heritability in wild populations: a simulation
study for Gaussian and binary traits. Freckleton R, editor. Methods in Ecology and Evolution.
2013;4: 260–275. doi:10.1111/2041-210x.12011

65. Heidelberger P, Welch PD. A spectral method for confidence interval generation and run length
control in simulations. Communications of the ACM. 1981;24: 233–245.
doi:10.1145/358598.358630

66. Hindson CM, Chevillet JR, Briggs HA, Gallichotte EN, Ruf IK, Hindson BJ, et al. Absolute
quantification by droplet digital PCR versus analog real-time PCR. Nat Meth. 2013;10: 1003–1005.
doi:10.1038/nmeth.2633

67. Laitinen RAE, Immanen J, Auvinen P, Rudd S, Alatalo E, Paulin L, et al. Analysis of the floral
transcriptome uncovers new regulators of organ determination and gene families related to flower
organ differentiation in Gerbera hybrida (Asteraceae). Genome Res. 2005;15: 475–486.
doi:10.1101/gr.3043705

68. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34: i884–i890. doi:10.1093/bioinformatics/bty560

69. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

70. Lien S, Koop BF, Sandve SR, Miller JR, Kent MP, Nome T, et al. The Atlantic salmon genome provides insights into rediploidization. Nature. 2016;533: 200–205. doi:10.1038/nature17164

71. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology. 2014;15: 550. doi:10.1186/s13059-014-0550-8

72. Abu-Jamous B, Kelly S. Clust: automatic extraction of optimal co-expressed gene clusters from gene expression data. Genome biology. 2018;19: 172. doi:10.1186/s13059-018-1536-8

73. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological themes among gene clusters. Omics : a journal of integrative biology. 2012;16: 284–287. doi:10.1089/omi.2011.0118

74. Garrido-Martín D, Palumbo E, Guigó R, Breschi A. ggsashimi: Sashimi plot revised for browser- and annotation-independent splicing visualization. Plos Comput Biol. 2018;14: e1006360. doi:10.1371/journal.pcbi.1006360

75. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative genomics viewer. Nature biotechnology. 2011;29: 24–26. doi:10.1038/nbt.1754