A multifaceted cellular damage repair and prevention pathway promotes high level tolerance to β-lactam antibiotics

Jung-Ho Shin [1, 2], Donghui Choe[4, 5], Brett Ransegnola[1, 2], Hye-Rim Hong [1, 2], Ikenna Onyekwere[1, 2], Byung-Kwan Cho[4, 5, 6], and Tobias Dörr [1, 2, 3a]

[1] Weill Institute for Cell and Molecular Biology, Cornell, University, Ithaca, NY 14853, USA

[2] Department of Microbiology, Cornell University, Ithaca NY 14853, USA

[3] Cornell Institute of Host-Microbe Interactions and Disease, Cornell University, Ithaca NY 14853, USA

[4] Department of Biological Sciences, Korea Advanced Institute of Science and Technology, Daejeon 34141, Republic of Korea.

[5] KI for the BioCentury, Korea Advanced Institute of Science and Technology, Daejeon 34141, Republic of Korea.

[6] Intelligent Synthetic Biology Center, Daejeon 34141, Republic of Korea.

[a] corresponding author: tdoerr@cornell.edu

Abstract

Bactericidal antibiotics are powerful drugs due to their ability to not only inhibit essential bacterial functions, but to convert them into toxic (and potentially lethal) processes. However, many important bacterial pathogens are remarkably tolerant against bactericidal drugs, due to inducible stress responses that repair antibiotic-induced damage. The mechanistic details of how stress responses promote whole population tolerance in important human pathogens are unknown. The two-component system VxrAB of the diarrheal pathogen *Vibrio cholerae,* a model system for high-level β-lactam

tolerance, is induced by exposure to cell wall acting antibiotic and controls a gene

network encoding highly diverse functions, including cell wall synthesis functions and

iron uptake systems. Here, we show that positive control over cell wall synthesis

functions only partially explains high level β-lactam tolerance. We find that in addition to

cell wall damage, β-lactam antibiotics inappropriately induce the Fur-regulated iron

starvation response, causing an increase in intracellular free iron levels and colateral

oxidative damage. We propose that VxrAB reduces antibiotic-induced toxic influx of

Fe^{2+} and concomitant metabolic perturbations by selectively downregulating iron uptake

transporters. Our results suggest that the ability to counteract diverse antibiotic-induced

stresses promotes high-level antibiotic tolerance and highlight the complex responses

elicited by antibiotics in addition to their primary mechanism of action.

Introduction

The discovery and development of antibiotics is one of the most important biomedical

breakthroughs in human history. The introduction of drugs that selectively kill bacteria

(bactericidal antibiotics) or inhibit their growth (bacteriostatic antibiotics) has directly

saved countless lives by enabling treatment of once common causes of death like

pneumonia and sepsis and indirectly enabled other medical breakthroughs, including

transplantation medicine and chemotherapy [1,2]. All this progress is threatened due to

the rapid increase in the occurrence of antibiotic treatment failure. We urgently need to

develop new therapeutics and improve the effectiveness of existing ones; this would be

aided by a precise understanding of antibiotic mechanisms of action, and especially of

bacterial strategies to avoid their noxious effects. In principle, bacteria resist antibiotics

by either developing the ability to grow in their presence (antibiotic resistance, ABR) or

to simply stay alive in their presence for extended time periods (antibiotic

tolerance/persistence) [3-8]. While the mechanisms and consequences of ABR are

relatively well-established, antibiotic tolerance remains poorly understood, limiting our

ability to develop antibiotic adjuvants that increase the efficacy of existing drugs.

The β-lactam antibiotics (penicillins, cephalosporins, carbapenems, cephamycins and

monobactams) are highly potent bactericidal agents. Their typically lethal action results

from their ability to simultaneously inhibit multiple targets (i.e, the transpeptidase

domain of multiple penicillin-binding proteins [PBPs]), which ultimately causes bacterial

cells to deplete essential cell wall precursors and self-destruct through the activity of

endogenous, cell wall lytic enzymes ('autolysins'; endopeptidases, amidases and lytic

transglycosylases) [9-12]. However, we and others have recently shown that many

clinically significant Gram-negative pathogens are remarkably β-lactam tolerant. The

cholera pathogen *Vibrio cholerae,* the opportunistic pathogen *Pseudomonas aeruginosa*

and clinical isolates of *Enterobacteriaceae* all survive treatment with β-lactam antibiotics

(including the "last resort" agent meropenem) by forming non-dividing, cell wall deficient

spheroplasts [11,13,14]. Upon removal of the antibiotic, spheroplasts rapidly revert to wild-

type shape and exponential growth, providing a possible explanation for treatment

failure in the clinic. Spheroplasts are reminiscent of bacteria in the so-called L-form

state [15,16] with the crucial distinction that spheroplasts, unlike L-forms, do not divide in

the presence of antibiotics.

The extent of antibiotic tolerance is a reflection of the cell's ability to mount a stress

response to counteract primary damage resulting from antibiotic-target interactions [17,18].

Numerous bacteria encode cell envelope damage response systems (alternative sigma

factors and two-component response regulators) that sense damage to the cell wall or

membranes and consequently upregulate complex regulons aimed at damage repair [19-

25]. In *V. cholerae*, high-level β-lactam tolerance depends on the cell wall damage

sensing two-component system VxrAB (also known as WigKR) [17]. The vast VxrAB

regulon is induced by cell wall-acting antibiotics and includes cell wall synthesis

functions and those involved in biofilm formation and type VI secretion, *inter alia* [17,26,27].

How VxrAB promotes high level tolerance against bactericidal β-lactams is unknown,

but logically must rely on repair of cellular damage inflicted by β-lactam antibiotics, *i.e.*,

primarily cell envelope damage. However, in addition to their primary mechanism of

action, bactericidal antibiotics have also been proposed to generate toxic reactive

oxygen species (ROS) as a byproduct, which are thought to enhance cell death [28-31].

This idea has been highly controversial and the resulting debate about it is ongoing [32-37].

Here, we present evidence supporting a multifactorial role for VxrAB in promoting high-

level β-lactam tolerance. We have found that the ability to upregulate cell wall synthesis

functions only accounts for part of the high-level tolerance observed in wild-type *V.*

cholerae. In addition, we propose that VxrAB's role in promoting tolerance is to

downregulate iron acquisition systems which are inappropriately induced due to the

accumulation of reactive oxygen species that are produced upon exposure to penicillin.

Our data highlight the pleiotropic nature of antibiotic mechanism of action and provide a

blueprint for the mechanistic underpinnings of the high-level β-lactam tolerance often

observed in significant Gram-negative pathogens.

Results

ChIP-Seq and RNA-Seq reveal a complex cell wall damage response

We previously elucidated the transcriptional network controlled by VxrAB upon

overexpression of a phosphomimetic version of VxrB (VxrB^{D78E}). Here, we directed our

attention to genes directly controlled by the VxrAB response. To this end, we performed

a ChIP-Seq experiment using a strain with functional, His-tagged, chromosomally

encoded VxrB assayed during penicillin (PenG) treatment. After 180 min of PenG

exposure (100 µg/ml, 10x MIC; these are the same conditions used in our previous

study[17]), 424 putative VxrB binding events were identified by comparison with a similarly

treated control strain expressing wild type, untagged VxrB (**Fig. 1A,** see **Table S3** for

complete list). We then validated the ChIP-Seq results via electromobility shift assays

(EMSA) of the 10 DNAs from the strongest VxrB binding events (**Fig. S1**) and, for a

subset, differential regulation via S1 nuclease assays of PenG-treated cells. We then

integrated this ChIP-Seq dataset with our re-analyzed (with more stringent parameters,

see Methods for details) previous RNA-Seq dataset of overexpressed VxrB^{D78E} [17] (**Fig.**

1B, see **Table S4** and **S5** for a complete list) to score VxrB binding events that

coincided with transcriptional up- or downregulation of the associated gene(s) (**Table**

S5). Our analysis resulted in the identification of 102 genes/operons that contained a

putative VxrB binding site and exhibited significantly increased, VxrB-dependent

expression levels, and 38 genes/operons that were scored as candidates for direct

binding and downregulation (**Table S5**). Preliminary functional assignments using the

KEGG database [38] revealed that the VxrB regulon controlled a large number (76.4%

107/140) of genes of unknown function. Among the annotated upregulated genes, many

coded for functions related to cell envelope biology (ABC transporters, peptidoglycan

synthesis, and membrane synthesis) (**Fig. S2**). Among the downregulated genes, we

observed genes encoding proteins involved in iron uptake systems, flagellar

biosynthesis, carbon metabolism, chemotaxis, cell wall lytic enzymes and

lipopolysaccharide biosynthesis (**Fig. S2**). Importantly, genes encoding cell wall

synthesis functions (e.g. class A PBP1a [aPBP1a]), whose Vxr-dependent

transcriptional regulation we have extensively validated previously [17], exhibited strong

VxrB binding sites (**Fig. 1D**) and differential, Vxr-dependent regulation in S1 nuclease

assays, increasing our confidence in the robustness of the ChIP-Seq dataset. Both

ChIP-Seq and RNA-Seq also corroborated autoregulation of the VxrAB operon (**Fig**. **1B,**

Table S1). We noted that while many of the putatively Vxr-controlled genes exhibited

VxrB binding in addition to differential regulation by phosphomimetic VxrB and VxrAB-

dependent regulation upon exposure to PenG (**Fig. 1D,** PBP1a and ZapB), some genes

were apparently indirectly regulated as they clearly exhibited transcriptional

downregulation (both upon overexpression of VxrB^{D78E} and exposure to PenG) without

a clear signal in our VxrB ChIP-Seq experiment (**Fig. 1D,** *flaA*). We speculate that this

may reflect either true indirect regulation or perhaps differential binding of the

phosphorylated versus the non-phosphorylated form of VxrB. Thus, the VxrAB cell wall

stress response controls a gene network composed of diverse functions with an

apparent enrichment for functions required for cell envelope homeostasis and iron

acquisition.

DNA footprinting analysis identifies putative VxrB binding sites

Based on an alignment of putative VxrB binding sites, we used the MEME suite [39] to

identify the canonical VxrB consensus binding site ("Vxr box"). We based our analysis

on the 102 upregulated and the 38 downregulated genes separately, reasoning that

binding site variations may account for up- *versus* downregulation. This resulted in the

putative VxrB binding motif TTGACAAAA-N2-TTGAC (N=any nucleotide) for the

upregulated genes and CGCTCATTTTTTAACCAAGT for the downregulated genes.

However, these putative binding motifs were poorly conserved across all Vxr-controlled

genes. We hypothesize that this may be reflective of different binding characteristics of

the phosphorylated vs. non-phosphorylated forms of VxrB; both forms might be

expected to occur after PenG treatment and could bind to different motifs. To further

characterize the VxrB binding site we conducted DNAse I footprinting assays using

purified VxrB and promoter regions of putative Vxr-controlled genes (**Fig. 2AB, S3**). We

again chose genes with the strongest VxrB binding as indicated in our EMSA and ChIP-

Seq experiments. In 10 out of 15 promoters, VxrB protected a ~ 50 bp sequence from

DNAse I digestion at low µM concentrations (**Fig. S3**), again corroborating direct binding

to these promoters. However, apparent affinity varied between promoters, with the

promoters of *pbp1a*, encoding aPBP1a, *mgtE*, encoding a magnesium transporter and

murJ, encoding the lipid II flippase exhibiting the highest degree of protection (**Fig. 2AB,**

Fig. S3). Surprisingly, we were again unable to construct a well-conserved binding site

for VxrB based on this subset of promoters, despite three independent assays

unequivocally demonstrating binding. This apparent binding promiscuity suggests that

VxrB might have the capacity to bind to a wide range of promoter sequences *in vivo*

and/or that it requires as yet unidentified accessory factors to modulate binding.

Control of cell wall synthesis only partially explains VxrAB's requirement for β-

lactam tolerance

To elucidate the mechanism of β-lactam tolerance in *V. cholerae,* we sought to dissect

the individual contributions of VxrB-regulated genes to resistance to killing. Since the

primary mechanism of action of β-lactam antibiotics is inhibition of cell wall synthesis,

we first investigated the role of Vxr-induced PG biogenesis in antibiotic tolerance. We

took two approaches to eliminate VxrB's positive control over PG synthesis. For

aPBP1a, we engineered a strain with a deletion in the second, functionally semi-

redundant [40,41] aPBP1b, which also contains an IPTG-inducible PBP1a as its sole copy.

As expected, growth of this strain was IPTG concentration-dependent (**Fig. S4**). We

then grew this strain either in high concentrations of IPTG (high PBP1a levels) or very

low concentrations that markedly reduced growth rate (**Fig. S4**). Addition of PenG to low

aPBP cultures resulted in only a modest (5 – 10-fold) decrease in viability (**Fig. 2C**).

This observation suggests that upregulation of PBP1a is indeed critical for survival in

the presence of PenG, but cannot account entirely for the steep, ~10.000-fold decrease

in post-antibiotic plating efficiency (**Fig. 2E** and [17]) of the Δ*vxrAB* mutant.

To corroborate these results, we also created a mutant with a partial in the putative

VxrB binding site of the *murJ* gene as determined from our DNA footprinting experiment.

MurJ-mediated PG precursor translocation is a critical bottleneck for cell wall synthesis

[42,43]. We thus expected that introducing a mutation that removes Vxr's positive control

over this step would allow us to gauge the significance of VxrB control over cell wall

synthesis. S1 nuclease assays confirmed that in the VxrB binding site deletion strain,

murJ expression was not significantly induced by PenG, similar to a Δ*vxrAB* strain (**Fig.**

2D). Similar to our observations with low PBP1a expression, the Vxr-independent *murJ*

strain exhibited modest (10-fold) lethality upon exposure to PenG (**Fig. 2E**). We

conclude that the ability to upregulate cell wall synthesis functions via VxrAB accounts

for only some of its ability to promote tolerance.

The* VxrB *regulon controls iron acquisition systems

Based on our previous RNA-Seq data, VxrB downregulates genes involved in iron

acquisition [17]. Indeed, examination of the RNA-Seq/ChIP-Seq datasets revealed a high

overlap between genes regulated by VxrB with those regulated by the iron starvation

response repressor Fur [44]. 47 out of the putative 63 Fur-regulated genes were also

downregulated by VxrB and 20 out of these 63 genes contained both a putative VxrB

and a Fur binding site (**Fig. 3AB**). Thus, the VxrB system appears to control multiple

functions required for iron homeostasis. Consistent with a role in maintaining proper iron

homeostasis, the Δ*vxrAB* mutant was more sensitive to growth inhibition by Fe^{2+} than

the wild type strain (**Fig. S5**).

Given VxrAB's apparent role in downregulating iron acquisition in response to cell wall damage, we expected that surplus iron accumulates in a PenG-treated $\Delta vxrAB$ mutant and that this might be detrimental to cells under these conditions. To test this, we measured intracellular iron levels using ICP-MS during treatment with PenG. Before treatment, both WT and the $\Delta vxrAB$ mutant strain contained similar iron levels at ~280 ng/mg protein (**Fig. 3C**). After 1 h of PenG exposure, WT iron levels remained relatively constant while the iron content of the $\Delta vxrAB$ strain started to increase significantly by ~60 % compared to baseline. Iron levels increased slightly in the WT between 90 min and 3 h, while levels in the $\Delta vxrAB$ mutant remained significantly higher. Consistent with an increase in free intracellular iron, the WT also became more sensitive to the iron-dependent quinone antibiotic streptonigrin [45,46] upon treatment with PenG (**Fig. 3D).** Thus, PenG exposure causes enhanced iron influx. If this iron influx is detrimental, we would expect iron chelators to alleviate the $\Delta vxrAB$ β-lactam sensitivity defect. Indeed, addition of 250 µM (0.25 x MIC) of the intracellular iron chelator 2,2' bipyridyl reproducibly increased survival of the $\Delta vxrAB$ mutant 50 – 100-fold (**Fig. 3E**); the ability to partially rescue could be reverted by addition of iron to the growth medium. As a caveat, we cannot fully exclude that the iron chelator reduces metabolic activity, and thus the rate of cell wall turnover, in a subpopulation of cells (*e.g.,* those that stochastically have lower iron levels to begin with), which could non-specifically increase tolerance levels, similar to what has been suggested before [33]. However, in conjunction with our ICP-MS/streptonigrin data demonstrating increased iron uptake in

PenG-treated cells, we tentatively interpret these data as evidence of iron-dependent

toxicity in $\Delta vxrAB$ cells.

Penicillin exposure induces ROS defense systems, ROS and oxidative damage

Iron is thought to be toxic to cells due to its ability to convert hydrogen peroxide (a

byproduct of metabolic processes) into oxygen radicals that pleiotropically oxidize and

thus damage macromolecules such as proteins and DNA [47] and generally disturb metal

homeostasis [48]. Indeed, antibiotics have been proposed to induce the increased

production of ROS through metabolic perturbations [29,31,37], however, the extent to which

this contributes to killing by antibiotics remains highly controversial [32-34]. To interrogate

the potential significance of ROS to lethality of the $\Delta vxrAB$ mutant, we first assessed

whether oxidative stress response systems were induced by PenG. We found that

PenG rapidly (within ~20 min) induced members of the H_2O_2 responsive OxyR1 regulon

(*prxA* and *dps*) [49-51], in the WT. Transcript levels of both OxyR1 readouts remained high

initially and then decreased to below baseline after ~ 60 min (*prxA*) or 180 min (*dps*),

suggesting that the wild type strain induces a response that detoxifies hydrogen

peroxide (**Fig. 4A, S6**). Both transcripts are induced by H_2O_2 in an OxyR1-dependent

manner [49,50] and we validated responsiveness to hydrogen peroxide for the *prxA*

transcript in our strain **(Fig. S6A).** Thus, *prxA* and *dps* are *bona fide* readouts for H_2O_2

production. Interestingly, the transcript for the response regulator OxyR1 itself was

upregulated upon PenG exposure, with dynamics similar to those observed for

members of its regulon (**Fig. S6B**). In the model organism *E. coli*, *oxyR* transcript levels

do not change under H_2O_2 exposure [52] and our results suggest that *V. cholerae* might

employ a different mode of regulation of its oxidative stress responses. A return to

baseline coincided with the upregulation of *V. cholerae*'s two catalases (*katB* and *katG*;

only *katG* appears to be under OxyR1 control in the cholera pathogen [50,51]) at later

stages of antibiotic exposure (**Fig. 4A**, **Fig. S6B**), likely revealing the detoxification

mechanism that enables the repression of the OxyR1 regulon. Consistent with a H_2O_2

sensing function, KatG was also induced by hydrogen peroxide (**Fig. S6A**).

In the Δ*vxrAB* strain, *oxyR1* baseline levels were slightly increased compared to the wild

type, while *prxA* and *dps* levels were similar. After exposure to PenG, *oxyR1* levels

decreased at first but the transcript remained detectable and above the wild-type

baseline for longer than the WT background (**Fig. 4A, S6**). The *prxA* and *dps* transcripts

exhibited a dynamic pattern with strong induction (*dps)* or transient downregulation

(*prxA*) in the mutant. We observed a pattern similar to *oxyR1* with a gene coding for

another oxidative stress mitigation factor, the Alkyl hydroperoxide reductase C, *ahpC*

(**Fig. S6B**). Interestingly, the transcript for *katG* remained uninduced in the *vxrAB*

mutant.

We confirmed that transcriptional regulation also resulted in increased catalase protein

activity using a gel-based activity assay for catalases (**Fig. 4B,** see Methods for details)

after exposure to PenG. We also included an assay for SOD (**Fig. 4B**) activity, since

superoxide dismutases are broadly conserved, important ROS scavenger enzymes [53].

Consistent with the observed transcript levels (**Fig. 4A**), catalases were only induced in

the WT, but not the mutant (**Fig. 4C**). In contrast, SOD activity increased similarly in

both the WT and the *vxrAB* mutant (**Fig. 4D**). Taken together, these results suggest that

both the WT and Δ*vxrAB* induce responses upon exposure to PenG that are associated

with oxidative stress, albeit with different, complex dynamics.

Since the transcriptional response to PenG suggested that cells may experience

oxidative damage, we next sought to measure ROS, and the consequences of their

generation, directly. First, we used a highly sensitive, fluorimetric assay (Amplex Red) to

measure intracellular hydrogen peroxide concentrations. Upon exposure to PenG, both

the WT and Δ*vxrAB* mutant experienced a transient increase in hydrogen peroxide

concentrations, with a peak between 20 and 60 minutes, followed by a decrease back to

baseline. However, the H_2O_2 concentration in the *vxrAB* mutant was significantly

elevated and at peak (20 - 60 min) accumulated more than double the concentration

measured in wild type cells (**Fig. 5A**). As expected, and consistent with the accurate

ability of Amplex Red to report H_2O_2, a Δ*oxR1* Δ*katB* Δ*katG* triple mutant, which lacks

the ability to elaborate the primary H_2O_2 detoxification systems, exhibited both higher

baseline and PenG-induced levels of H_2O_2 than the WT and Δ*vxrAB* mutant; additionally,

external spiking of catalase into our assay reactions completely abolished the signal

(**Fig. 5A**). We also measured H_2O_2 in the supernatant of PenG-treated cells, since H_2O_2

should be able to diffuse through membranes to some degree. We did not, however,

note a meaningful accumulation of peroxide in the supernatant of any of our mutants

(**Fig. 5B**), demonstrating that at least for the duration of our measurements, there was

no significant diffusion out of the cell.

We next assessed the cellular consequences of oxidative damage using a protein

oxidation assay via Western Blot. Consistent with our results showing enhanced iron

sensitivity of the mutant, protein oxidation levels were increased in the $\Delta vxrAB$ mutant

vs. WT even prior to addition of antibiotic (**Fig. 5C**). Addition of PenG resulted in a rapid

(within 20 min) increase in protein oxidation levels in the WT, which decreased after 40

min but stayed above baseline throughout β-lactam exposure. In contrast, the *vxrAB*

mutant steadily accumulated oxidized proteins throughout the exposure time,

corroborating that oxidative stress has a more profound effect on the *vxrAB* mutant than

the WT strain. Both, wild type and the *vxrAB* mutant exhibited increased protein

oxidation when exposed to hydrogen peroxide as a positive control (**Fig. 5C, Fig. S7**).

Taken together, these data suggest that both WT and $\Delta vxrAB$ experience oxidative

stress in the presence of PenG and the $\Delta vxrAB$ mutant is unable to mount a sufficient

response.

The electron transport chain is required for both maximal killing and H_2O_2

production

How does PenG treatment result in production of ROS? A possible origin is the electron

transport chain (ETC), which in *V. cholerae* is a major generator of ROS under normal

growth conditions [54]. To test the ETC's involvement in PenG-induced ROS production,

we created mutants in two ETC components, namely *V. cholerae's* principal sodium

translocating NADH dehydrogenase NqrA [55] and *ubiA*, which encodes a key initial step

towards ubiquinone biosynthesis. NqrA has been shown to be a significant source of

ROS under normal growth conditions [54], but for both mutants, we expect a drastic

reduction in ETC electron flow. Crucially, neither mutation resulted in a significantly

decreased growth rate when grown in medium containing the fermentable carbon

source glucose (albeit, both mutations caused a slight increase in lag phase) (**Fig. S8A**).

Reduced growth rates can affect β-lactam sensitivity non-specifically [56] and mutants

conferring slower growth can thus confound the interpretation of experiments conducted

with such strains. Consistent with the expectation of reduced ETC activity in these

mutants, disruption of *nqrA* and *ubiA* in both the WT and the Δ*vxrAB* mutant background

resulted in a marked reduction in PenG-induced accumulation of $H_2O_2,$ as measured by

Amplex Red (**Fig. S8B**). To test the hypothesis that the reduction in H_2O_2 also results in

lack of killing in the Δ*vxrAB* mutant, we conducted time-dependent killing experiments

with these strains. As observed previously, WT viability remained stable for the entire

duration of the experiment while the Δ*vxrAB* mutant displayed a ~10,000-fold decrease

in viability. The Δ*nqrA* and *ubiA* mutants were as tolerant as the WT. However, a

combination of ETC mutations with Δ*vxrAB* reproducibly increased survival by 10- (*nqrA*)

or 100-fold (*ubiA*) (**Fig. 6A)**. Thus, maximal Δ*vxrAB* death after PenG exposure

depends on the presence of the ETC.

ROS detoxification systems promote tolerance at high cell densities

We next asked whether ROS damage mitigation systems were required for WT

tolerance. We created knockout mutants deficient in the iron starvation repressor Fur

(expected to result in an increase in intracellular iron and thus enhanced ROS), SodB

and a triple mutant defective in both catalases and OxyR1. We grew all strains from late

exponential into stationary phase and periodically withdrew aliquots that were then

exposed to PenG for 3 h, and measured survival. As a comparison, we subjected the

Δ*vxrAB* mutant to the same treatment. While all mutants except for Δ*vxrAB* exhibited

wild-type survival patterns in exponential phase, entry into stationary phase coincided

with significant killing in all mutants that are expected to experience enhanced ROS

production (**Fig. 6B**). Consistent with these results, *sodB* had previously answered a

Transposon Insertion Sequencing (TnSeq) screen for factors that are required for PenG

tolerance [57]. We thus propose that ROS contribute significantly to β-lactam-mediated

killing in stationary phase (see Discussion for details).

Penicillin exposure induces the Fur regulon

Thus far, we have shown that VxrAB controls iron acquisition and that a lack in the

ability to do so may result in enhanced PenG-mediated killing. But why do PenG-treated

Δ*vxrAB* cells increase their iron load? According to evidence presented above, H_2O_2

accumulates upon exposure to PenG. In *Bacillus subtilis,* peroxide is sensed by the

transcriptional regulator PerR, a member of the Fur family of repressors [58]. PerR senses

peroxide via oxidation of an iron cofactor [59], which results in induction of the PerR ROS

detoxification regulon [60-62]. *V. cholera*e, like many other Gram-negative bacteria, lacks

PerR; but in other bacteria, peroxide is thought to directly inactivate Fur through

oxidation of its iron cofactor [63]. Derepression of the Fur regulon in turn would result in

the upregulation of iron acquisition systems. To test the hypothesis that PenG induces

the Fur regulon, we used S1 nuclease mapping to monitor transcript levels of the

canonical Fur regulon readout *ryhB*[64], a small RNA [65]. Consistent with our line of

thinking, *ryhB* was highly induced by addition of external H_2O_2 **(Fig. 7A).** In *E. coli, sodB*

transcript levels are negatively controlled by *ryhB,* and consistent with a similar

regulatory mechanism in *V. cholerae, ryhB* induction (**Fig. 7A**) coincided with a

disappearance of the *sodB* transcript during treatment with hydrogen peroxide (**Fig. 7A**)*.*

Consistent with a regulation via iron starvation sensing, *ryhB* transcript also increased

dramatically upon exposure of cells to the cation chelator EDTA (**Fig. 7A**). Upon PenG

exposure, *ryhB* transcript was reproducibly induced ~2-fold in both the WT and Δ*vxrAB*

after 20 min, followed by a return to baseline. After prolonged PenG-treatment (180 min),

ryhB remained uninduced in the WT, but increased dramatically in the Δ*vxrAB* mutant.

Consistent with these transcriptional changes, a P$_{ryhB}$-*msfGFP* transcriptional reporter

construct also increased in fluorescence after exposure to PenG in both, WT and

Δ*vxrAB* (**Fig. 7B**). Other Fur-regulated genes (*hutA, hutB* and *vc2212*), also exhibited a

marked, and in some cases transient, upregulation upon PenG exposure (**Fig. 7B**).

These results suggest that indeed the Fur regulon is induced by exposure to PenG.

Interestingly, we noticed in our ICP-MS dataset that concomitantly with iron uptake,

manganese concentrations also increased in PenG-treated cells (**Fig. 7C**). Manganese

has been proposed to be a cellular response to "rescue" iron-dependent enzymes that

have lost their iron cofactor due to iron starvation or oxidative damage; biologically

appropriate, Mn^{2+} uptake is thus under control of Fur in Gram-negative model

organisms [66-68]. While a specific Mn^{2+} importer has not been identified in *V. cholerae* [69],

our data suggest that the cholera pathogen might possess a similar regulatory

arrangement.

Discussion

Our overall model of VxrAB-mediated tolerance is depicted in **Fig. 8;** based on the evidence presented in this study, we propose the following: PenG induces cell wall degradation, resulting in perturbations of the electron transport chain. Our work does not directly address the nature of this perturbation, but it may simply result from topological alterations of the cytoplasmic membrane brought about by loss of the cell wall. Interestingly, and consistent with this line of thinking, other cell wall deficient cells (L-Forms) also appear to accumulate ROS [70,71]. We hypothesize that partial reduction of molecular oxygen through the ETC (particularly NqrA) results in the production of superoxide O_2^-. O_2^- is removed by SOD, which catalyzes the formation of hydrogen peroxide (H_2O_2). H_2O_2 induces the *oxyR1* regulon, but we propose that it also inappropriately inactivates the Fur repressor via oxidation of its Fe^{2+} cofactor. Inactivation of Fur (i.e., induction of the Fur regulon) results in the upregulation of iron uptake systems, which increases the intracellular iron pool. Intracellular iron reacts with H_2O_2 to form hydroxyl radical HO^-, which damages proteins and other macromolecules. The wild-type strain mounts effective responses to all these types of damage. SOD, AhpC and catalases detoxify superoxide and H_2O_2. VxrAB induces cell wall synthesis functions to prepare cells for speedy recovery once the antibiotic is removed. At the same time VxrAB also downregulates some iron acquisition genes and other Fur functions to dampen detrimental iron influx, and thus decrease oxidative damage. Interestingly, some of our data suggest that ROS production might contribute significantly to killing in later growth phases (early stationary phase), and stationary phase-specific oxidative damage is not without precedent. Periplasmic SODs are under

control of stationary phase regulators in diverse bacteria and yeast [72-74] and SOD is

required for antibiotic tolerance in stationary phase in *P. aeruginosa* as well [75]. We

hypothesize that during rapid growth in exponential phase, the primary mechanism of

action (cell wall degradation) is likely the more important contributor to cell death, while

in stationary phase (slower/no growth, less cell wall damage), the harmful effects of

ROS accumulation may become more visible. This is noteworthy, since in the natural

environment, bacterial growth can be expected to resemble stationary phase more than

rapid growth in exponential phase [76]. On separate note, β-lactam-mediated killing in

stationary phase may seem counterintuitive at first, since at least in the well-

characterized model organism *E. coli*, cessation of growth upon entry into this growth

phase is generally assumed to result in complete protection from β-lactam-induced

damage [56,77]. However, the *vxrAB* mutant still exhibited significant PenG-dependent

mortality even when PenG exposure only occurred several hours into stationary phase,

i.e., far into a period of non-growth, suggesting that this mutant experiences PenG-

induced cell envelope damage even when populations stop growing. We thus propose

that in *V. cholerae,* cell wall turnover (and thus PenG's ability to inflict damage)

continues at least for some time after cells stop growing. A similar phenomenon could

explain the unusual observation that the Lyme disease pathogen *Borrelia burgdorferi*

remains susceptible to killing by amoxicillin in stationary phase [78].

The connection between antibiotic exposure and ROS production and particularly the

extent of the contribution of ROS to killing is still actively debated in the field. We agree

with a previous hypothesis [34] that some of the interpretive difficulties in other studies

may have arisen from the low concentrations of antibiotics used (often around the MIC).

High concentrations of bactericidal antibiotics might kill cells too quickly through their

primary mechanism of action to measure a response to the proposed secondary

mechanism. Lower concentrations around the MIC might result in day-to-day variation

of killing dynamics that are difficult to reproduce under different conditions. In contrast,

bacteria like *V. cholerae* (as well as *P. aeruginosa, Klebsiella pneumoniae, K.*

aerogenes and *Enterobacter cloacae*), exhibiting almost population-wide tolerance to β-

lactams [11,13,14,57], may be more appropriate model organisms to dissect the layers of

physiological changes occurring as a consequence of antibiotic exposure. In these

tolerant bacteria, the primary mechanism of action (degradation of the cell wall) is

recapitulated; however, unlike in other bacteria, this does not result in lysis and cell

death, allowing us to examine a large population of live cells after treatment with high

concentrations (10 x MIC or more) of antibiotic.

One of the main critiques of the idea that ROS contribute to killing was that some

groups failed to measure meaningful ROS levels (and the expected responses to them)

in antibiotic-treated cells in the model organism *E. coli* [33,34]. This has been challenged

by other groups [35,37,79]. Here, we have shown that *V. cholerae* accumulates ROS and

induces defense systems against them after exposure to PenG. We propose that in the

vxrAB mutant, antibiotic-induced lethality is multifactorial. Both, cell wall defects and

ROS may contribute to killing. Importantly, other detrimental consequences of iron influx

may also play a role, for example mismetalation of key enzymes [48]. While ROS

production and iron influx are measurable in the wild type as well (to a lower degree

than the mutant), it is likely that its multiple, highly effective defense systems prevent

antibiotic killing. This thinking is consistent with a previous observation that compared to

E. coli and *P. aeruginosa,* exposure of *V. cholerae* to sub-MIC concentrations of β-

lactam antibiotics only caused a moderate increase in mutagenesis (a possible

consequence of ROS accumulation) [80]. Thus, the cholera pathogen's high tolerance to

β-lactam exposure might be the consequence of an exceptional ability to repair cell

damage.

Most means to directly probe the effect of ROS on cellular physiology under antibiotic

stress are not without caveats. Growth under anaerobic conditions changes growth

rates and thus response to antibiotics non-specifically (and it is for this reason that we

did not consider killing under anaerobic conditions here). Bipyridyl might simply reduce

metabolism in a subset of cells (those at the lower end of the iron content normal

distribution), causing non-specific induction of "tolerance" through metabolic shutdown.

While we cannot entirely rule out this possibility, we demonstrate that PenG-treatment

simultaneously induces iron uptake and H_2O_2 production, raising the strong possibility

that the increase in iron levels has toxic consequences. As a side-note, iron withdrawal

is a common antibacterial strategy in the human host and might thus significantly affect

antibiotic susceptibility *in vivo*, regardless of whether the mechanism is through

metabolic shutdown or iron detoxification [81]. Since iron influx into antibiotic-stressed

cells can also be detrimental, the extent of the host's ability to control iron levels in a

specific infection locale might thus significantly and divergently influence antibiotic

susceptibility in different host tissues.

While it is difficult to conclude beyond doubt that ROS contribute significantly to killing

by antibiotics in the Δ*vxrAB* mutant and we thus hesitate to unequivocally do so here,

we conclude that β-lactams (and likely other antibiotics) elicit more complex cellular

responses than just those associated with their primary mechanism of action. A more

context-, metabolism- and physiology-centered view of antibiotic action [82-85] might

therefore yet yield critical insight to enable the development of novel adjuvants that

increase the efficacy of antibiotics in the complex infection environment. Lastly, with or

without ROS, our data suggest that targeting bacterial stress responses could be a

powerful strategy to enhance the activity of bactericidal drugs against tolerant

pathogens.

Acknowledgments

Research in the Dörr laboratory is supported by NIAID grant 1R01AI143704. This work

was partially supported by the Korea Bio Grand Challenge Project

(2018M3A9H3024759 to B.K.C.) through the National Research Foundation of Korea

(NRF) funded by the Ministry of Science and ICT, Republic of Korea. We thank the

Helmann lab (Cornell University) for sharing equipment and reagents. We further thank

Stavroula Hatzios (Yale University), John Helmann (Cornell University), Joseph Peters

(Cornell University), James Imlay (University of Illinois) and members of the Dörr lab for

helpful comments.

Figure legends

Figure 1 | ChIP-Seq and RNA-Seq analysis reveals a complex VxrB regulon.

(**A**) Overview of VxrB-binding profiles across the *V. cholerae* genome after β-lactam

exposure (PenG 100µg/ml [10x MIC] for 3 hours). Red peaks indicate VxrB binding sites

above the significance cutoff (q-value<10^{-10}; MACS). (**B**) RNA-Seq MA plot is presented

as fold change between VxrB^{D78E} overexpressed sample and control sample. Red dots

indicate differentially expressed genes after VxrB^{D78E} overexpression (P-value < 0.001).

Representative VxrB target genes are marked on both chromosomes and on the MA

plot. **(C)** Venn diagram depicting overlap between RNA-seq and ChIP-Seq datasets **(D)**

Regulatory modes of representative VxrB-controlled genes in response to PenG. Upper

plot shows VxrB binding patterns (assessed via ChIP-Seq), lower panel shows

transcriptional response upon overexpression of VxrB ^{D78E} (assessed via RNA-Seq).

RNA expression was confirmed by S1 nuclease mapping assay in wild type and *Δvxr*AB

mutant strains after exposure to PenG (100 µg/ml for 3 hours).

Figure 2 | Peptidoglycan synthesis control only partially explains *Δvxr*AB lethality.

(A,B) DNAse I footprinting results for VxrB interaction with promoters controlling cell

wall synthesis functions. Red boxes indicate core DNA regions protected by purified

VxrB protein (final concentration 0.17 - 1.38 µM) on the *murJ* (A) or *pbp1A* (B) promoter.

Predicted start codons of each gene are presented in bold letters. VBS, putative **V**xrB

Binding **S**equence **(C- E)** Reduction of cell wall synthesis gene expression causes a

moderate tolerance defect. **(C)** Survival of a Δ*pbp1b* Δ*pbp1a* P$_{iptg}$:*pbp1a* strain after 6

hours of PenG exposure. P_{up} indicates conditions of high IPTG concentrations (100 μM),

P_{down} low IPTG concentration (2 μM). Survival fraction is cfu/ml of PenG-treated cultures

normalized to untreated cultures grown in and plated on medium containing the same

IPTG concentration. **(D)** Deleting the putative VxrB binding site from the *murJ* promoter

abrogates induction of *murJ* by PenG exposure. Transcript levels were measured by S1

nuclease mapping. Numbers represent induction as measured by band intensity,

normalized to pre-exposure WT. n.d., not determined. (**E)** Time-dependent killing

experiment in the presence of PenG (100 μg/ml, 10 x MIC). All data are means (+/-

standard error) of 3 independent biological replicates.

Figure 3 | VxrAB controls genes involved in iron homeostasis. (A) Venn diagram

showing overlap between the VxrB regulon and known *V. cholerae* Fur-controlled genes.

(as per RegPrecise3.0 DB, http://regprecise.lbl.gov/RegPrecise/). 20 out of 47 Fur-

regulated genes also have a VxrB binding site and are differentially regulated by

VxrB^{D78E}. (**B)** Genomic organization of iron uptake systems and their putative Fur/VxrB

binding sites. **(C)** Time-course of iron accumulation in PenG-treated WT and Δ*vxrAB*

cells as measured by ICP-MS. Data are average of 3 independent biological replicates,

error bars represent standard deviation. $*$$*$P=0.001; $*$$*$$*$P<0.0003 (paired t-test) **(D)**

Streptonigrin-sensitivity of PenG-treated cells. Cultures were grown to a density of ~2 x

10^8 cfu/ml. Before ("- PenG") and after addition of PenG (100 μg/ml, 10x MIC), ~ 5 x 10^7

cells were spread on an LB agar plate. In the PenG-treated samples ("+ PenG"), beta

lactamase solution was also added to remove the antibiotic. Streptonigrin (5 mg/ml, in

DMSO) was then added on a filter disk, followed by incubation for 24 h at 37 ˚C. Values

represent average diameter of clearance zones (+/- standard deviation) of six biological

replicates. SNG, streptonigrin. **(E)** Iron chelation reduces $\Delta vxrAB$ death in the presence

of PenG. Time-dependent killing experiment in the absence or presence of 250 μM

bipyridyl (Bip) with or without addition of 250 μM Fe^{2+}. Data are average of 3

independent biological replicates, error bars represent standard deviation.

Figure 4 | PenG exposure induces ROS defense systems (A) RNA expression

profiles of first line ROS defense systems in *V.cholerae* monitored by S1 mapping assay.

Wild type and *Δvxr*AB were grown to exponential phase (OD_{600}~0.5) in liquid LB

medium and treated with 100ug/ml PenG. Samples were collected at different time

points as indicated and S1 nuclease protection assay carried out to quantify the

expression level of each gene. Quantified average values (normalized to WT before

PenG exposure) from three independent experiments are presented under the signals

of each gene. **(B)** Gel based catalase and superoxide dismutase (SOD) activity staining.

Wild type and *Δvxr*AB cells were grown to exponential phase (OD_{600}~0.5) in LB medium

and exposed to PenG. Samples were collected at the indicated time points. 30 μg/well

of crude proteins were separated by 5% (catalases) or 8% (SODs) native PAGE at 4 °C

followed by staining (see Methods for details). To quantify the SOD or catalases activity,

stained gels were scanned and analyzed by Odyssey Imaging Systems (LI-COR

Biosciences) and Image Studio Lite software. Rpol α subunit (RNA polymerase I, α

subunit) was visualized as internal control. Coomassie staining (loading control) was

used to ensure consistent protein loading between samples. **(C)** and **(D)** Quantified

activity values from the gel shown in **(B)**. Data are average values from three

independent experiments normalized to untreated wild type, for **(C)** catalase or **(D)**

superoxide dismutase activity.

Figure 5 | PenG induces hydrogen peroxide production and protein oxidation (A)

(A-B) Total intracellular **(A)** or supernatant **(B)** hydrogen peroxide (H_2O_2) levels were

measured using Amplex^{TM} Red Hydrogen Peroxide/Peroxidase Assay Kit and

normalized to total cellular protein level (measured by Qubit^{TM}). Data are average

values from three independent experiments, error bars represent standard deviation.

Insert represents magnified data to accentuate subtle differences between mutants and

conditions. Cat, catalase. **(C)** Detection of proteins damage following oxidative stress

triggered by PenG treatment. To measure the total cellular protein oxidation, OxyBlot

Protein Oxidation Detection Kit (Sigma-Aldrich) was used. Wild type and *Δvxr*AB mutant

were grown to OD_{600}~0.4 and harvested at different times after PenG treatment. 0.5 mM

H_2O_2 treated cells were used as a positive protein oxidation control with oxidized

standard marker ($_{OX}$M). Both DNP-derivatized or nonderivatized (negative control)

protein samples were separated by polyacrylamide gel electrophoresis followed by

Western blotting with-DNP specific antibody.

Figure 6 | ROS generation and detoxification modulate PenG-induced lethality

(A) Time-dependent killing of respiratory chain mutants (Δ*nqrA,* encoding NADH

dehydrogenase and Δ*ubiA*, encoding an early step of ubiquinone synthesis) in the

presence of PenG (100 µg/ml, 10xMIC). All strains were grown in LB 0.2 % glucose.

Data are mean of three independent biological replicates, error bars represent standard

deviation. **(B)** Growth-phase dependent killing assay. At designated time-points, a 5 ml

aliquot was withdrawn and exposed to 100 µg/ml PenG for 3 h. Cfu/ml for each time

point was assessed before (solid lines) and after addition of PenG (dotted lines).

Figure 7 | PenG and hydrogen peroxide induce the Fur iron starvation response.

(A) Expression levels of the Fur-regulated small RNA *ryhB* and the RyhB target *sodB*

were quantified using S1 nuclease mapping. Cells were grown to mid-exponential

phase followed by exposure to H_2O_2, PenG or the positive control 2 mM EDTA (a strong

chelator of divalent cations) for the indicated duration. Numbers represent expression

levels normalized to untreated control (averages from three independent replicates). **(B)**

To confirm the induction of *ryhB* transcript by PenG treatment, the native *ryhB* promoter

was fused with *msfGFP* and introduced into WT and *ΔvxrA* mutant cells. Each strain

was cultured until mid-exponential phase, when 1mM EDTA, 0.5mM FeCl or 100µg/ml

PenG were added for the indicated duration. PenG-mediated induction of other Fur-

regulated genes (*hutA/B*, coding for heme transport systems and *vc2212*, which

encodes a component of a ferric citrate uptake system) was measured by S1 nuclease

mapping. **(C)** Time-course of manganese accumulation in PenG-treated WT and

Δ*vxrAB* cells as measured by ICP-MS. Data are average of 3 independent biological

replicates, error bars represent standard deviation. ∗∗P=0.001; ∗∗∗P<0.0003 (paired t-

test)

Figure 8 | Proposed model for VxrAB-mediated antibiotic tolerance.

PenG induces ETC dysfunction, resulting in the production of ROS. Hydrogen peroxide

illegitimately induces the Fur regulon, resulting in the upregulation of iron acquisition

systems, resulting in enhanced iron influx and oxygen radical generation. VxrAB

dampens this response by downregulating iron acquisition genes, in addition to its

primary role as a positive regulator of cell wall synthesis genes. SO, superoxide.

MATERIALS AND METHODS

Bacterial strains and culture conditions.

All *V. cholerae* strains used in this study were derivatives of *V. cholerae* WT El Tor strain N16961 (Heidelberg *et al.,* 2000) or E7946 (for chitin-induced transformation) listed in Table S1. *V. cholerae* was grown on Luria-Bertani (LB) medium with or without 0.2% glucose or in M9 medium containing 0.2% glucose at 37°C unless otherwise indicated. All media contained 200 µg/ml streptomycin (to which *V. cholerae* is resistant). Where applicable, growth media were supplemented with 100 µg/ml Benzylpenicillin potassium salt (PenG) (Fisher BioReagents, cat# BP914-100), isopropyl-β-D-1-thiogalactopyranoside (IPTG; final 0.1~1mM) (GOLDBIO, cat# 12481C50), or catalase 1µg/ml (Sigma-Aldrich, Cat# c9322)**.** *Escherichia coli* DH5α and DH5α λpir were used for routine DNA cloning (Sambrook *et al*., 1989 and Platt R *et al.,* 2000). Unless indicated otherwise, LB liquid media were inoculated from an overnight culture and incubated at 37°C with shaking at 200 rpm until reaching mid-exponential phase (OD$_{600}$ ~0.4). Where applicable, trimethoprim (Sigma, cat #T7883, dissolved in DMSO) was used at 15 µg/ml final concentration.

Strain construction

Deletion mutants were generated via replacement of the open reading frame/binding site with the scar sequence TAATGCGGCCGCACTCGAGTAATAATGATGA or, where indicated, a trimethoprim resistance cassette. Knockouts were generated either by homologous recombination using the suicide vector pCVD442 [86] or by chitin-induced

natural transformation, both as described before [87] [17]. The following primer

combinations were used (see supplemental material for sequence information).

murJ Δvbs: TDP1062/1064 + TDP1066/1067, fused via stitching PCR with nesting

primers TDP1078/1079, followed by chitin transformation using *vc1807*::trim (generated

using primers TDP597/598 + TDP599/600 [on trimR from the V. cholerae HAITI strain [88]]

+TDP601/602, fused via stitching PCR with nesting primers TDP603/604) as primary

selection.

Δ*oxyR1*: TDP1057/58 + TDP1059/60, cloned into Xba1-digested pCVD442 using

isothermal assembly [89], followed by knockout via homologous recombination.

Δ*katG:* TDP1100/1101 + TDP1102/1103, cloned into Xba1-digested pCVD442 using

isothermal assembly, followed by knockout via homologous recombination.

Δ*katB*: TDP1103/1104 + TDP1105/1106, cloned into Xba1-digested pCVD442 using

isothermal assembly, followed by knockout via homologous recombination.

lacZ::P$_{ryhB}$:*msfGFP*: TDP1152/1235 + TDP1237/38, cloned into Sma1-digested pTD101,

a suicide vector for chromosomal integration into the *lacZ* locus [90].

Δ*nqrA*: TDP1388/1389 + TDP1390/1391, fused via stitching PCR with nesting primers

TDP1392/1393, followed by chitin transformation using *vc1807*::trim (see above) as

primary selection.

Δ*ubiA*::*trim* TDP1399/1401 + TDP1404/1403 + TDP636/637, fused via stitching PCR

with nesting primers TDP1405/1406, followed by chitin transformation.

Δ*sodB*: TDP828/829 + TDP830/831, cloned into Xba1-digested pCVD442 using

isothermal assembly, followed by knockout via homologous recombination.

Δ*fur*. TDP834/835 + TDP836/837, cloned into Xba1-digested pCVD442 using isothermal

assembly, followed by knockout via homologous recombination.

Pulldown of VxrB binding DNA fragments and ChIP-Seq library construction. Cells

carrying either 6xHis+VxrB or the native gene were grown in LB medium to mid-

logarithmic phase (at an OD_{600} ~0.5). Following treatment with PenG (100 µg/ml, 3

hours), 30 ml aliquots were spun down and the pellets were saved at -80°C. For cross-

linking, pellets were resuspended in 1ml buffer CJ1 [100 mM Na_2HPO_4, 600 mM NaCl,

and 5 mM imidazole;pH 8.0] and 30 µl of 34% formaldehyde was added, followed by

rocking at room temperature for 10 min. To quench cross-linking reaction, 133 µl of 1 M

Glycine pH7.5 (final molar concentration 0.133 M) was added, and the cells were

shaken gently at 4°C for 30 min. Cells were collected by centrifugation, and the pellets

were resuspended in 1 ml buffer CJ2 [100 mM Na_2HPO_4, 600 mM NaCl, 0.05% tween^{TM},

and 30 mM imidazole;pH 8.0] after washing twice with same CJ1 buffer. Cells were then

lysed by sonication, followed by addition of Micrococcal nuclease (100 U ml^{-1}, NEB

cat# M0247S) and RNase A (1 mg ml^{-1}) for fragmentation of genomic DNA and

removal of RNA. 950 µl of supernatant was collected after centrifugation.

For DNA-VxrB complex purification, HisPur^{TM} Ni-NTA magnetic beads (cat#88831-

Thermo Scientific) were equilibrated with six bed volumes of CJ2 buffer [100 mM

Na_2HPO_4, 600 mM NaCl, 0.05% tween^{TM}, and 30 mM imidazole;pH 8.0]. 500 µg of

crude lysate (protein concentration determined by Bradford assay) was mixed with the

Ni-NTA resin and incubated at 4°C for 4 hours on rotation mixer. The bead slurry was

recovered via magnetic stand for 5 min standing and washed twice with 400 µl buffer

CJ3 [100 mM Na_2HPO_4, 600 mM NaCl, 0.05% tween[TM], and 50 mM imidazole; pH 8.0].

The VxrB-DNA complexes were then eluted from the beads by addition of CJ4 elution

buffer [100 mM Na_2HPO_4, 600 mM NaCl, 0.05% tween[TM], and 500 mM imidazole;pH

8.0]. Enriched target DNAs were purified by using PCR Clean-Up Kits (Thermo Fisher

Scientific) after reversal of crosslinking at 65°C overnight. The desired DNA fragment

size (200~300 bps) was then purified by gel extraction. For illumina HiSeq, each DNA

library was constructed with specific barcode sequences, according to the

manufacturer's instruction of TruSeq ChIP Library Preparation Kit (IP-202-1012,

Illumina). The constructed libraries were sequenced using 100-bp paired-end read

protocol on an Illumina HiSeq 2500 platform (Illumina) at the Biotechnology Resource

Center (BRC) at Cornell University.

ChIP-Seq, binding peak detection, and motif analysis. Sequencing data was

processed by CLC Genomics Workbench (CLC Bio). First, raw sequencing reads were

quality trimmed using a Trim Sequence Tool with a quality limit of 0.05. Any reads with

more than two ambiguous bases were discarded at this step. Then, quality trimmed

reads were mapped onto the *V. cholerae* reference genome (NCBI accession

NC_002505.1 and NC_002506.1) with mismatch cost of 2, insertion/deletion cost of 3,

length fraction of 0.9, and similarity fraction of 0.9. Non-specific matches were mapped

randomly. Mapping files (.bam) were submitted to the Model-based Analysis of ChIP-

Seq 2 (MACS2) software with a mapping file of a mock sample as a control (Zhang *et*

al., 2008). Peaks were detected with shift size of 50 and q-value cutoff of 10^{-10}. For motif

analysis, peaks with q-value lower than 10^{-40} were selected and 100 bp DNA sequences

centered at peak summits were submitted to MEME Suite software v4.11.4 (Bailey *et al*.,

2009).

RNA-Seq analysis

Sequencing data (obtained previously) were also processed on a CLC Genomics

Workbench (CLC Bio). Raw sequencing reads were quality trimmed with the same

method with ChIP-Seq reads. Quality trimmed reads were mapped strand specifically

on the reference genome sequence by Transcriptome Analysis Tool with the following

parameters: maximum number of mismatches of 2, length fraction, and similarity

fraction of both 0.9. Number of mapped reads on each gene was normalized by

DESeq2 package in Bioconductor v3.4 (Love *et al*., 2014). Genes with p-value (padj,

DESeq2) lower than 0.001 were scored as differentially expressed genes (DEGs).

Purification of VxrB protein. Wild-type VxrB protein was purified from *E. coli* BL21

(DE3/pLysS) cells containing pET15b based recombinant plasmid. The coding

sequence for WT VxrB was cloned into pET15b (Novagen) into the *Nde*I and *Bam*HI

restriction sites. For purification of VxrB from this construct, an overnight culture from a

single colony was used to inoculate 1 liter of LB medium. Cells were grown with

vigorous shaking at 37°C to an optical density at 600 nm (OD$_{600}$) of 0.5 and were

induced with 1 mM (final concentration) isopropyl-β-D-thiogalactopyranoside (IPTG) for

6 h at 30°C. Harvested cells were re-suspended in binding buffer [20 mM Tris-HCl (pH

7.9), 0.5 M NaCl, 1 mM TCEP (tris(2-carboxyethyl)-phosphine), and 5 mM imidazole]

and cell extracts were prepared by sonication and centrifugation at 16,000 rpm for 45

min. Cell extracts were loaded onto a cobalt-charged NTA column and then washed

with 6 volumes of binding buffer followed by 6 volumes of washing buffer [20 mM Tris-

HCl (pH 8.0), 0.5 M NaCl, and 50 mM imidazole]. VxrB protein was eluted with 10

volumes of elution buffer [20 mM Tris-HCl (8.0) and 0.5 M NaCl] containing linear

imidazole gradients from 100 to 500 mM. Fractions containing VxrB protein were pooled

and dialyzed against buffer J1 [20 mM Tris-HCl (pH 8.0), 250 mM NaCl, 5% (vol/vol)

glycerol, and 4 mM EDTA] to remove imidazole and cobalt and then transferred to two

extra dialysis buffer J2 [20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol, and 0.2

mM DTT] and buffer J3 [20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 30% glycerol, and 1

mM DTT]. The purified VxrB protein was concentrated by centrifugal filter devices

(Millipore, 3,000 MW CO) before injection onto High load TM (16/60) pg Superdex G75

column in FPLC system (Pharmacia). The column was equilibrated with buffer SG [20

mM Tris-HCl (pH 8.0), 150 mM NaCl, and 1 mM DTT]. Eluted fractions were monitored

through UV detector. Concentration of purified VxrB was estimated in triplicate by

Bradford assay (Bio-Rad) using BSA (bovine serum albumin, Sigma-Aldrich) as the

calibration standard at A595. Measurement of UV absorbance at 280 nm, combined

with calculated molar extinction coefficient (ε_{280} = 46470 M^{-1} cm^{-1}, http://expasy.org/cgi-

bin/protparam), gave nearly identical values. The purity of protein was confirmed

through Coomassie blue staining of loaded protein samples in the SDS-PAGE gel. The

protein was stored in final storage buffer JHS [20 mM Tris-HCl (pH 8.0), 150 mM NaCl,

1 mM DTT, 30% Glycerol, 1 mM MgCl$_2$, and 1 mM ATP] at -80°C.

DNase I footprinting assay. The probe DNAs were prepared by using the "crush and

soaking method" (Sambrook *et al.*, 1989) after PCR with the primer pairs shown in

Table S2. The 5′ end of the probes was labeled with 6-FAM (Fluorescein). Binding

reactions were performed as described below for EMSA, except using 250 ng probe

DNA in a 40 μl reaction mixture. DNA probes were incubated with increasing

concentrations of VxrB (0.17, 0.34, 0.68, and 1.38 μM). After 20 min incubation at room

temperature, 40 μl of 5 mM $CaCl_2$ and 10 mM $MgCl_2$ was added, followed by adding

200 U of RQI RNase-free DNase I (Promega) for 1 min. The cleavage reaction was

stopped by adding 90 μl stop solution (200 mM NaCl, 30 mM EDTA, 1% sodium

dodecyl sulfate, 125 μg ml^{-1} of glycogen), followed by DNA extraction and

precipitation. Samples were analyzed by capillary electrophoresis on an ABI 3730x1

DNA analyzer (Life Technologies). The DNA probe only with no added VxrB was

analyzed in parallel. The core protected regions were indicated with red lines and its

corresponding DNA sequences were shown at the bottom of peak profiles.

Preparation of total RNA and S1 nuclease mapping analysis. Wild type and mutant

strains were cultured to mid-logarithmic phase (at an OD_{600} of 0.4 to 0. 5) in LB

medium. Following application of a stress (PenG or H_2O_2), total RNA was extracted

using the "hot phenol method" as described in Shin *et al.*, 2014, albeit without lysozyme

pretreatment. RNA was quantified and quality-controlled by absorbance spectroscopy

and confirmed by resolving RNA samples on 1.3% formaldehyde-agarose gels. Gene-

specific DNA oligonucleotide probes for *vxrA*, *vxrB*, *pbp1A*, *zapB*, *flaA*, *murJ*, *oxyR1*,

sodB, *sodA*, *katG*, *katB*, *prxA*, *oxyR2*, *ahpC*, *dps*, *fur*, and *ryhB* transcripts were

generated by PCR amplification using *V. cholerae* wild-type genomic DNA as template.

The appropriate primer pairs are listed in Table S2. Each specific DNA probe was

radiolabeled with (γ-^{32}P) ATP and T4 polynucleotide kinase and 30,000-40,000 cpm of

labeled probe was used in each reaction. For S1 assays, 100 µg of total RNA was

pelleted and lyophilized. The total RNA pellet was then carefully resuspended in 20 µl

hybridization buffer [40 mM PIPES (pH 6.4), 400 mM NaCl, 1 mM EDTA, 80% (v/v)

formamide]. Individual samples were incubated at 95°C for 25 min and cooled to 42°C.

Following overnight incubation, 300 µl of S1 nuclease mix containing 100 units of S1

nuclease in S1 nuclease buffer [280 mM NaCl, 30 mM NaOAc (pH 4.4), 4.5 mM ZnOAc]

was added and incubated at 37°C for 45 min. The reaction was terminated by addition

of 75 µl of S1 nuclease termination solution (2.5 M NH_4OAc, 0.05 M EDTA). The DNA-

RNA hybrid was precipitated by adding 400 µl of isopropanol (100 % v/v) and the pellet

was washed with 70% (v/v) ethanol, vacuum dried, and re-suspended in 13 µl alkaline

loading dye (Sambrook *et al.*, 1989). The protected DNA fragments were then resolved

by 6% (w/v) polyacrylamide gels containing 7 M urea. Gels were dried using a Hoefer

Scientific Drygel Slab Gel Dryer 01385 and then imaged on a phosphor imaging screen

(Typhoon FLA 7000; GE); bands were quantified using Multi Gauge V3.0 (Fuji).

Quantified average fold intensity (compared to non-treated sample) from three

independent experiments were presented under the each signal with standard deviation

values.

Killing Assay. For most time-dependent killing experiments, overnight cultures of wild

type and each mutant strains were grown at 37°C in LB medium in the presence or

absence of additional supplement such as IPTG or 0.2% glucose. At each time-point,

samples were collected, serially diluted (sterile PBS) and spot-plated for cfu/ml

determination (plates were incubated at 30 °C for 20 – 24 h). Where indicated, PenG

(100 µg/ml), 250 µM Bip, or 250 µM $FeCl_2$ were added. For the growth phase dependent

experiment, wild type, *ΔvxrAB*, *Δfur*, *ΔsodB* and *ΔoxyR1ΔkatGΔkatB* mutants were

cultured in the presence of 1µg/ml catalase for overnight seed (catalase is required for

ΔoxyR1 growth) in 100 mL LB in flasks. At designated time points, 5ml cell cultures

were transferred to new tubes and cultured with or without PenG for 3 hours; cfu/mL

was determined before and after addition of the antibiotic.

Streptonigrin sensitivity assay

Cells were diluted 100fold from overnight cultures and grown to a density of ~2 x 10^8

cfu/ml. At each time point (before addition of 100 µg/ml PenG and 3 h after), 200 µL of

culture were spread on 25 ml LB agar plates using glass beads. At the 3 h time point, 1

mg/ml (final concentration) β-lactamase (Sigma, catalog # P4524) was added prior to

plating. Following plating, a sterile filter disk containing 5 µL of 5 mg/ml streptonigrin

solution (Sigma, catalog # S1014, dissolved in DMSO) was placed in the center of the

plate. Plates were then incubated 24 h at 37 °C. Measurements are averages of six

biological replicates.

ICP-MS analysis for intracellular metal concentration measurement. A single

colony of the indicated *V. cholerae* strains from LB plates was inoculated into 5 mL LB

with streptomycin (200 µg/mL) and grown overnight. The next morning, 0.1% of

overnight cultured seeds were inoculated in 200 ml LB supplemented with antibiotics.

These cells were grown shaking at 37°C for 2~3 hours until they reached an OD_{600} of

~0.5. Samples were collected before and at indicated time points after PenG treatment.

At the indicated time points, cells were pelleted by centrifugation (10 min, 20000 rcf)

and washed three times with phosphate buffered saline (PBS) buffer containing 0.1 M

EDTA followed by two chelex (BioRad, cat # 1422822)-treated PBS buffer washes to

remove all external metal ions. Cells were then resuspended in 400 μl of chelex-treated

PBS buffer with 1% Triton X-100 containing 75 mM NaN3 to induce lysis (Jolliffe et al.,

1981) and incubated at 37°C for 90 min. 10 μl of each supernatant was used to

measure total cell protein concentration by Bradford assay. 600 μl of 5% HNO3 with 0.1%

(v/v) Triton X-100 were then added to the rest of the supernatant, which was boiled at

95°C for 30 min. After centrifugation (10 min, 20000 rcf), the supernatant was then

diluted 1000fold with 1% HNO3. Metal levels were measured by ICP-MS (Perkin-Elmer

ELAN DRC II using ammonia as the reaction gas and gallium as an internal standard)

and normalized against total cell protein concentration. Data represent with average

values from biological triplicates. Significant differences between wild type and *Δvxr*AB

after PenG exposure were determined by paired R t-test as indicated: ∗∗P=0.001;

∗∗∗P<0.0003.

Catalase and SOD activity assay. Catalase activity staining was performed as

described in Cho *et al.,* 1997. Briefly, cell extracts (30 μg per well, obtained by 20 cycles

of sonication on ice at 20 % power, 3 s pulse, followed by 10 s cooling) were separated

on a 5% native polyacrylamide gel and electrophoresed at 4°C with 80 volt. The gel was

washed in distilled water and then treated with 1 mM H_2O_2 for 20 min. After treatment,

the gel was rinsed and transferred to 1% (w/v) ferric chloride and 1% (w/v) potassium

ferricyanide solution. The reaction was stopped by washing twice in distilled water.

Superoxide dismutase activity staining was performed as described in [91]. The cell

extracts (30 µg/well) were electrophoresed on a 8% native polyacrylamide gel under the

same conditions as described above for catalase activity, after which the gel was

incubated in a solution containing 2.5 mM nitroblue tetrazolium (NBT) for 25 min and

then in 50 mM phosphate buffer (pH 7.8) containing 28 µM riboflavin and 28 mM

tetramethyl ethylene diamine for 20 min in the dark. The gel was placed in distilled

water and exposed on a light box for 15 min until the dark blue background color

appeared. To quantify SOD and catalase activity, stained gels were scanned and

analyzed by Odyssey Imaging Systems (LI-COR Biosciences) and Image Studio Lite

software. Quantified average values from three independent experiments are presented

as a fold activity against wild type non-treated sample for SODs and Catalases.

Western blot analysis to detect alpha subunit of RNA polymerase. For the internal

control of SOD and catalase activity, the same crude extracts were resolved on two

independent 12 % polyacrylamide gels for each western blotting and coomassie

staining. After electrophoresis the proteins were transferred to a membrane by using a

semi-drying transfer system (I Blot 2-Invitrogen). The membrane was then blocked with

blocking solution (dry skim milk dissolved in 20 mM Tris-HCl (pH7.8), 150 mM

NaCl, 0.1% Triton X-100) overnight. The membrane was incubated with monoclonal

anti-αRpol antibody (BioLegend-cat#663104) for two hours, washed with TBST (20 mM

Tris-HCl (pH7.8), 150 mM NaCl, 0.1% Triton X-100), and incubated with anti-mouse

secondary antibody (IRDye 800CW, Li-Cor cat# 926-32211). All images were scanned

on an Odyssey CLx imaging device (LI-COR Biosciences) and visualized using image

studio lite software.

Measurement of total cellular H_2O_2 level. Total cellular hydrogen peroxide (H_2O_2)

level was measured using the Amplex^{TM} Red Hydrogen Peroxide/Peroxidase Assay Kit

(Invitrogen™ cat#A22188) and normalized to total cellular proteins. Wild type and

indicated mutant stains were cultured in LB medium (in the absence or presence of

additional 0.2% glucose when it needed) until OD_{600} ~0.5. PenG (100 µg/ml) was then

added and 1ml cell culture was withdrawn and pelleted at each indicated time point.

Pellets were resuspended in 100 µl cold 1xPBS buffer and lysed by sonication, followed

by centrifugation at 4°C (10 min, 20000 rcf). 3 µl of each supernatant was used to

measure total cell protein concentration by Bradford assay. 20µl extracts from cleaned

supernatant were reacted with kit components following manufacturer's instruction. For

H_2O_2 quantification, the reaction product resorufin was read by fluorometer (QUBIT 3,

Invitrogen) at Ex/Em of: ~571/585 nm. Obtained values were normalized to total protein

concentration.

Detection of oxidized proteins. Oxidized proteins containing carbonyl groups were

detected using the OxyBlot Protein Oxidation Detection Kit (Sigma-Aldrich: cat#S7150).

Briefly, wild type and *ΔvxrAB* mutant were grown to exponential growth phase and

harvested at different times after PenG treatment. 0.5 mM H_2O_2 treated cells were used

as a positive protein oxidation control with oxidized standard marker ($_{OX}$M). The protein

sample (10µg/well) was incubated with dinitrophenylhydrazine (DNPH) (derivatization

reaction) for 15 min, followed by neutralization with a solution containing glycerol and β-

mercaptoethanol. Both DNP-derivatized or nonderivatized (for negative control) protein

samples were electrophoresed on a 10% SDS-PAGE and then transferred to a

nitrocellulose membrane. After blocking, the membrane was incubated for 1 h with a

rabbit antibody against DNP (cat# D9656), followed by incubation with secondary

antibody: goat anti-rabbit IgG (Li-COR, IRDye® 800CW cat# 926-32211) for 1 h at room

temperature as recommended by the manufacturer. The signals were scanned on an

Odyssey CLx imaging device (LI-COR Biosciences) and visualized using image studio

lite software.

Microscopy. All images were taken on a Leica MDi8 microscope (Leica Microsystems,

GmbH, Wetzlar, Germany) with a PECON Temp-Controller 2000-1 (Erbach, Germany),

heated stage at 37 °C for growth experiments, or room temperature. Time-lapse

microscopy was performed by imaging frames five minutes apart and data were

analyzed by ImageJ program. The msfGFP expressing cells were imaged at

$485_{Ex}/530_{Em}$ for two second exposure time. Collected images were processed in ImageJ

by subtracting background and adjusting brightness/contrast uniformly across all

fluorescent images. Representative images were taken at least three independent

experiments.

1. Spellberg, B. The future of antibiotics. *Crit Care* **18**, 228 (2014).
2. Fair, R.J. & Tor, Y. Antibiotics and bacterial resistance in the 21st century. *Perspect Medicin Chem* **6**, 25-64 (2014).
3. Lewis, K. Persister cells: molecular mechanisms related to antibiotic tolerance. *Handb Exp Pharmacol*, 121-133 (2012).
4. Lewis, K. Persister cells. *Annu Rev Microbiol* **64**, 357-372 (2010).
5. Lewis, K. Persister cells, dormancy and infectious disease. *Nat Rev Microbiol* **5**, 48-56 (2007).
6. El-Halfawy, O.M. & Valvano, M.A. Antimicrobial heteroresistance: an emerging field in need of clarity. *Clin Microbiol Rev* **28**, 191-207 (2015).
7. Wilmaerts, D., Windels, E.M., Verstraeten, N. & Michiels, J. General Mechanisms Leading to Persister Formation and Awakening. *Trends Genet* **35**, 401-411 (2019).
8. Balaban, N.Q.*, et al.* Definitions and guidelines for research on antibiotic persistence. *Nat Rev Microbiol* (2019).
9. Kitano, K., Tuomanen, E. & Tomasz, A. Transglycosylase and endopeptidase participate in the degradation of murein during autolysis of Escherichia coli. *J Bacteriol* **167**, 759-765 (1986).
10. Flores-Kim, J., Dobihal, G.S., Fenton, A., Rudner, D.Z. & Bernhardt, T.G. A switch in surface polymer biogenesis triggers growth-phase-dependent and antibiotic-induced bacteriolysis. *Elife* **8**(2019).
11. Dorr, T., Davis, B.M. & Waldor, M.K. Endopeptidase-mediated beta lactam tolerance. *PLoS Pathog* **11**, e1004850 (2015).
12. Cho, H., Uehara, T. & Bernhardt, T.G. Beta-lactam antibiotics induce a lethal malfunctioning of the bacterial cell wall synthesis machinery. *Cell* **159**, 1300-1311 (2014).
13. Monahan, L.G.*, et al.* Rapid conversion of Pseudomonas aeruginosa to a spherical cell morphotype facilitates tolerance to carbapenems and penicillins but increases susceptibility to antimicrobial peptides. *Antimicrob Agents Chemother* **58**, 1956-1962 (2014).
14. Cross, T.*, et al.* Spheroplast-mediated carbapenem tolerance in Gram-negative pathogens *Antimicrob Agents Chemother* (2019).
15. Errington, J., Mickiewicz, K., Kawai, Y. & Wu, L.J. L-form bacteria, chronic diseases and the origins of life. *Philos Trans R Soc Lond B Biol Sci* **371**(2016).
16. Errington, J. Cell wall-deficient, L-form bacteria in the 21st century: a personal perspective. *Biochem Soc Trans* **45**, 287-295 (2017).
17. Dorr, T.*, et al.* A cell wall damage response mediated by a sensor kinase/response regulator pair enables beta-lactam tolerance. *Proc Natl Acad Sci U S A* **113**, 404-409 (2016).
18. Dörr, T., Lewis, K. & Vulic, M. SOS response induces persistence to fluoroquinolones in Escherichia coli. *PLoS Genet* **5**, e1000760 (2009).

19.   Raivio, T.L. Everything old is new again: an update on current research on the
Cpx envelope stress response. *Biochim Biophys Acta* **1843**, 1529-1541 (2014).
20.   Helmann, J.D. Bacillus subtilis extracytoplasmic function (ECF) sigma factors
and defense of the cell envelope. *Curr Opin Microbiol* **30**, 122-132 (2016).
21.   Radeck, J., Fritz, G. & Mascher, T. The cell envelope stress response of Bacillus
subtilis: from static signaling devices to dynamic regulatory network. *Curr Genet*
63, 79-90 (2017).
22.   Wall, E., Majdalani, N. & Gottesman, S. The Complex Rcs Regulatory Cascade.
Annu Rev Microbiol **72**, 111-139 (2018).
23.   Guest, R.L. & Raivio, T.L. Role of the Gram-Negative Envelope Stress Response
in the Presence of Antimicrobial Agents. *Trends Microbiol* **24**, 377-390 (2016).
24.   Bhagirath, A.Y.*, et al.* Two Component Regulatory Systems and Antibiotic
Resistance in Gram-Negative Pathogens. *Int J Mol Sci* **20**(2019).
25.   Bury-Mone, S.*, et al.* Global analysis of extracytoplasmic stress signaling in
Escherichia coli. *PLoS Genet* **5**, e1000651 (2009).
26.   Cheng, A.T., Ottemann, K.M. & Yildiz, F.H. Vibrio cholerae Response Regulator
VxrB Controls Colonization and Regulates the Type VI Secretion System. *PLoS*
Pathog **11**, e1004933 (2015).
27.   Teschler, J.K., Cheng, A.T. & Yildiz, F.H. The Two-Component Signal
Transduction System VxrAB Positively Regulates Vibrio cholerae Biofilm
Formation. *J Bacteriol* **199**(2017).
28.   Dwyer, D.J., Kohanski, M.A., Hayete, B. & Collins, J.J. Gyrase inhibitors induce
an oxidative damage cellular death pathway in Escherichia coli. *Mol Syst Biol* **3**,
91 (2007).
29.   Kohanski, M.A., Dwyer, D.J., Hayete, B., Lawrence, C.A. & Collins, J.J. A
common mechanism of cellular death induced by bactericidal antibiotics. *Cell* **130**,
797-810 (2007).
30.   Kohanski, M.A., Dwyer, D.J., Wierzbowski, J., Cottarel, G. & Collins, J.J.
Mistranslation of membrane proteins and two-component system activation
trigger antibiotic-mediated cell death. *Cell* **135**, 679-690 (2008).
31.   Dwyer, D.J., Kohanski, M.A. & Collins, J.J. Role of reactive oxygen species in
antibiotic action and resistance. *Curr Opin Microbiol* **12**, 482-489 (2009).
32.   Imlay, J.A. Diagnosing oxidative stress in bacteria: not as easy as you might
think. *Curr Opin Microbiol* **24**, 124-131 (2015).
33.   Liu, Y. & Imlay, J.A. Cell death from antibiotics without the involvement of
reactive oxygen species. *Science* **339**, 1210-1213 (2013).
34.   Keren, I., Wu, Y., Inocencio, J., Mulcahy, L.R. & Lewis, K. Killing by bactericidal
antibiotics does not depend on reactive oxygen species. *Science* **339**, 1213-1216
(2013).
35.   Hong, Y., Zeng, J., Wang, X., Drlica, K. & Zhao, X. Post-stress bacterial cell
death mediated by reactive oxygen species. *Proc Natl Acad Sci U S A* **116**,
10064-10071 (2019).
36.   Dwyer, D.J., Collins, J.J. & Walker, G.C. Unraveling the physiological
complexities of antibiotic lethality. *Annu Rev Pharmacol Toxicol* **55**, 313-332
(2015).

37. Dwyer, D.J.*, et al.* Antibiotics induce redox-related physiological alterations as
part of their lethality. *Proc Natl Acad Sci U S A* **111**, E2100-2109 (2014).
38. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res **28**, 27-30 (2000).
39. Bailey, T.L.*, et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic*
Acids Res **37**, W202-208 (2009).
40. Dorr, T.*, et al.* Differential requirement for PBP1a and PBP1b in in vivo and in
vitro fitness of Vibrio cholerae. *Infect Immun* **82**, 2115-2124 (2014).
41. Dorr, T.*, et al.* A novel peptidoglycan binding protein crucial for PBP1A-mediated
cell wall biogenesis in Vibrio cholerae. *PLoS Genet* **10**, e1004433 (2014).
42. Ruiz, N. Lipid Flippases for Bacterial Peptidoglycan Biosynthesis. *Lipid Insights* **8**,
21-31 (2015).
43. Teo, A.C. & Roper, D.I. Core Steps of Membrane-Bound Peptidoglycan
Biosynthesis: Recent Advances, Insight and Opportunities. *Antibiotics (Basel)* **4**,
495-520 (2015).
44. Davies, B.W., Bogard, R.W. & Mekalanos, J.J. Mapping the regulon of Vibrio
cholerae ferric uptake regulator expands its known network of gene regulation.
Proc Natl Acad Sci U S A **108**, 12467-12472 (2011).
45. White, J.R. & Yeowell, H.N. Iron enhances the bactericidal action of streptonigrin.
Biochem Biophys Res Commun **106**, 407-411 (1982).
46. Yeowell, H.N. & White, J.R. Iron requirement in the bactericidal mechanism of
streptonigrin. *Antimicrob Agents Chemother* **22**, 961-968 (1982).
47. Imlay, J.A. The molecular mechanisms and physiological consequences of
oxidative stress: lessons from a model bacterium. *Nat Rev Microbiol* **11**, 443-454
(2013).
48. Imlay, J.A. The mismetallation of enzymes during oxidative stress. *J Biol Chem*
289, 28121-28128 (2014).
49. Xia, X.*, et al.* OxyR-activated expression of Dps is important for Vibrio cholerae
oxidative stress resistance and pathogenesis. *PLoS One* **12**, e0171201 (2017).
50. Wang, H.*, et al.* Catalases promote resistance of oxidative stress in Vibrio
cholerae. *PLoS One* **7**, e53383 (2012).
51. Wang, H.*, et al.* OxyR2 Modulates OxyR1 Activity and Vibrio cholerae Oxidative
Stress Response. *Infect Immun* **85**(2017).
52. Storz, G., Tartaglia, L.A. & Ames, B.N. The OxyR regulon. *Antonie Van*
Leeuwenhoek **58**, 157-161 (1990).
53. Johnson, L.A. & Hug, L.A. Distribution of reactive oxygen species defense
mechanisms across domain bacteria. *Free Radic Biol Med* (2019).
54. Muras, V., Dogaru-Kinn, P., Minato, Y., Hase, C.C. & Steuber, J. The Na+-
Translocating NADH:Quinone Oxidoreductase Enhances Oxidative Stress in the
Cytoplasm of Vibrio cholerae. *J Bacteriol* **198**, 2307-2317 (2016).
55. Steuber, J.*, et al.* The structure of Na(+)-translocating of NADH:ubiquinone
oxidoreductase of Vibrio cholerae: implications on coupling between electron
transfer and Na(+) transport. *Biol Chem* **396**, 1015-1030 (2015).
56. Tuomanen, E., Cozens, R., Tosch, W., Zak, O. & Tomasz, A. The rate of killing of
Escherichia coli by beta-lactam antibiotics is strictly proportional to the rate of
bacterial growth. *J Gen Microbiol* **132**, 1297-1304 (1986).

57. Weaver, A.I.*, et al.* Genetic Determinants of Penicillin Tolerance in Vibrio
cholerae. *Antimicrob Agents Chemother* **62**(2018).
58. Fillat, M.F. The FUR (ferric uptake regulator) superfamily: diversity and versatility
of key transcriptional regulators. *Arch Biochem Biophys* **546**, 41-52 (2014).
59. Lee, J.W. & Helmann, J.D. The PerR transcription factor senses H2O2 by metal-
catalysed histidine oxidation. *Nature* **440**, 363-367 (2006).
60. Helmann, J.D.*, et al.* The global transcriptional response of Bacillus subtilis to
peroxide stress is coordinated by three transcription factors. *J Bacteriol* **185**, 243-
253 (2003).
61. Faulkner, M.J. & Helmann, J.D. Peroxide stress elicits adaptive changes in
bacterial metal ion homeostasis. *Antioxid Redox Signal* **15**, 175-189 (2011).
62. Duarte, V. & Latour, J.M. PerR vs OhrR: selective peroxide sensing in Bacillus
subtilis. *Mol Biosyst* **6**, 316-323 (2010).
63. Varghese, S., Wu, A., Park, S., Imlay, K.R. & Imlay, J.A. Submicromolar
hydrogen peroxide disrupts the ability of Fur protein to control free-iron levels in
Escherichia coli. *Mol Microbiol* **64**, 822-830 (2007).
64. Seo, S.W.*, et al.* Deciphering Fur transcriptional regulatory network highlights its
complex role beyond iron metabolism in Escherichia coli. *Nat Commun* **5**, 4910
(2014).
65. Masse, E. & Gottesman, S. A small RNA regulates the expression of genes
involved in iron metabolism in Escherichia coli. *Proc Natl Acad Sci U S A* **99**,
4620-4625 (2002).
66. Kehres, D.G., Janakiraman, A., Slauch, J.M. & Maguire, M.E. Regulation of
Salmonella enterica serovar Typhimurium mntH transcription by H(2)O(2), Fe(2+),
and Mn(2+). *J Bacteriol* **184**, 3151-3158 (2002).
67. Martin, J.E., Waters, L.S., Storz, G. & Imlay, J.A. The Escherichia coli small
protein MntS and exporter MntP optimize the intracellular concentration of
manganese. *PLoS Genet* **11**, e1004977 (2015).
68. Ikeda, J.S., Janakiraman, A., Kehres, D.G., Maguire, M.E. & Slauch, J.M.
Transcriptional regulation of sitABCD of Salmonella enterica serovar
Typhimurium by MntR and Fur. *J Bacteriol* **187**, 912-922 (2005).
69. Fisher, C.R., Wyckoff, E.E., Peng, E.D. & Payne, S.M. Identification and
Characterization of a Putative Manganese Export Protein in Vibrio cholerae. *J*
Bacteriol **198**, 2810-2817 (2016).
70. Kawai, Y.*, et al.* Cell growth of wall-free L-form bacteria is limited by oxidative
damage. *Curr Biol* **25**, 1613-1618 (2015).
71. Kawai, Y.*, et al.* Crucial role for central carbon metabolism in the bacterial L-form
switch and killing by beta-lactam antibiotics. *Nat Microbiol* (2019).
72. Schnell, S. & Steinman, H.M. Function and stationary-phase induction of
periplasmic copper-zinc superoxide dismutase and catalase/peroxidase in
Caulobacter crescentus. *J Bacteriol* **177**, 5924-5929 (1995).
73. Longo, V.D., Gralla, E.B. & Valentine, J.S. Superoxide dismutase activity is
essential for stationary phase survival in Saccharomyces cerevisiae.
Mitochondrial production of toxic oxygen species in vivo. *J Biol Chem* **271**,
12275-12280 (1996).

74. Gort, A.S., Ferber, D.M. & Imlay, J.A. The regulation and role of the periplasmic copper, zinc superoxide dismutase of Escherichia coli. *Mol Microbiol* **32**, 179-191 (1999).

75. Martins, D.*, et al.* Superoxide dismutase activity confers (p)ppGpp-mediated antibiotic tolerance to stationary-phase Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **115**, 9797-9802 (2018).

76. Haugan, M.S., Charbon, G., Frimodt-Moller, N. & Lobner-Olesen, A. Chromosome replication as a measure of bacterial growth rate during Escherichia coli infection in the mouse peritonitis model. *Sci Rep* **8**, 14961 (2018).

77. Keren, I., Kaldalu, N., Spoering, A., Wang, Y. & Lewis, K. Persister cells and tolerance to antimicrobials. *FEMS Microbiol Lett* **230**, 13-18 (2004).

78. Sharma, B., Brown, A.V., Matluck, N.E., Hu, L.T. & Lewis, K. Borrelia burgdorferi, the Causative Agent of Lyme Disease, Forms Drug-Tolerant Persister Cells. *Antimicrob Agents Chemother* **59**, 4616-4624 (2015).

79. Brynildsen, M.P., Winkler, J.A., Spina, C.S., MacDonald, I.C. & Collins, J.J. Potentiating antibacterial activity by predictably enhancing endogenous microbial ROS production. *Nat Biotechnol* **31**, 160-165 (2013).

80. Gutierrez, A.*, et al.* beta-Lactam antibiotics promote bacterial mutagenesis via an RpoS-mediated reduction in replication fidelity. *Nat Commun* **4**, 1610 (2013).

81. Sheldon, J.R. & Skaar, E.P. Metals as phagocyte antimicrobial effectors. *Curr Opin Immunol* **60**, 1-9 (2019).

82. Yang, J.H.*, et al.* A White-Box Machine Learning Approach for Revealing Antibiotic Mechanisms of Action. *Cell* **177**, 1649-1661 e1649 (2019).

83. Radlinski, L. & Conlon, B.P. Antibiotic efficacy in the complex infection environment. *Curr Opin Microbiol* **42**, 19-24 (2018).

84. Yang, J.H., Bening, S.C. & Collins, J.J. Antibiotic efficacy-context matters. *Curr Opin Microbiol* **39**, 73-80 (2017).

85. Stokes, J.M., Lopatkin, A.J., Lobritz, M.A. & Collins, J.J. Bacterial Metabolism and Antibiotic Efficacy. *Cell Metab* (2019).

86. Donnenberg, M.S. & Kaper, J.B. Construction of an eae deletion mutant of enteropathogenic Escherichia coli by using a positive-selection suicide vector. *Infect Immun* **59**, 4310-4317 (1991).

87. Dalia, A.B. Natural Cotransformation and Multiplex Genome Editing by Natural Transformation (MuGENT) of Vibrio cholerae. *Methods Mol Biol* **1839**, 53-64 (2018).

88. Hubbard, T.P.*, et al.* A live vaccine rapidly protects against cholera in an infant rabbit model. *Sci Transl Med* **10**(2018).

89. Gibson, D.G.*, et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**, 343-345 (2009).

90. Dorr, T., Cava, F., Lam, H., Davis, B.M. & Waldor, M.K. Substrate specificity of an elongation-specific peptidoglycan endopeptidase and its implications for cell wall architecture and growth of Vibrio cholerae. *Mol Microbiol* **89**, 949-962 (2013).

91. Beauchamp, C. & Fridovich, I. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal Biochem* **44**, 276-287 (1971).

A

VxrB, 100 ug/ml PenG 3H — Chromosome I

ChIP-Seq (x10²)

(LPS) VC0215 (PBP1B) VC0602 (PBP5,6/RlpA) VC0947/8 (Flagella assembly) VC2188-93 (PBP1A) VC2635

ChIP-Seq (x10²)

VxrB+His, 100 ug/ml PenG 3H

VxrB, 100 ug/ml PenG 3H — Chromosome II

ChIP-Seq (x10²)

(ShyA) VCA0079 (FhuD) VCA0227 (VxrAB) VCA0565/6 (DcuC) VCA0665

ChIP-Seq (x10²)

VxrB+His, 100 ug/ml PenG 3H

B

Log_2(Fold-change; over exp./control) vs Log_2 (Mean Expression)

DEGs (n= 818)
non-DEG (n=2870)

VC0157
vxrB
pbp1b
vxrA
pbp1a
hutA

C

RNA-Seq (818 genes)   ChIP-Seq (424 peaks)
p-value < 0.001 q-value $<10^{-10}$

678 genes   140 genes (127 peaks)   297 peaks

Up (102 genes)
Down (38 genes)

D

0.5 kb — VC2634 Penicillin-binding protein 1A VC2635 VC2636

ChIP-Seq (x10³) — VxrB, 100PG 3H

RNA-Seq normalized mapped reads (x10³) — WT control — VxrB O/E

Transcripts (S1 nuclease mapping) — Wild Type / ΔvxrAB — min (100 μg/ml) 0 20 180 / 0 20 180 — pbp1A — rRNAs

Directly activated gene

0.25 kb — VC2685 ZapB VC2686 VC2687

ChIP-Seq (x10³) — VxrB, 100PG 3H

RNA-Seq normalized mapped reads (x10²) — WT control — VxrB O/E

Transcripts (S1 nuclease mapping) — Wild Type / ΔvxrAB — min (100 μg/ml) 0 20 180 / 0 20 180 — zapB — rRNAs

Directly repressed gene

0.3kb — VC2187 Flagellin VC2188 VC2189

ChIP-Seq (x10³) — VxrB, 100PG 3H

RNA-Seq normalized mapped reads (x10²) — WT control — VxrB O/E

Transcripts (S1 nuclease mapping) — Wild Type / ΔvxrAB — min (100 μg/ml PenG) 0 20 60 180 / 0 20 60 180 — flaA — rRNAs

Poor binding of VxrB or indirectly regulated gene

Figure 2

A *murJ* promoter

No His-VxrB

His-VxrB (µM): 0.17, 0.34, 0.68, 1.38

ATTCTATCAGCATTTTTTACCAATGCTAATACTTATCTGATCTCACTATGACATT--38N--**TTG**

VBS

B *pbp1A* promoter

No His-VxrB

His-VxrB (µM): 0.17, 0.34, 0.68, 1.38

Cooperative binding region

AGGCAAACCCTTTTTTAGTTCTTTTAGTTTCTTTACAATTTTCTGATAATCGAG--328N--**GTG**

VBS

C

P_IPTG → *pbp1a*

fraction survival: *pbp1a* (P_up), *pbp1a* (P_down)

E

ΔVBS → *murJ*

wt   ΔvxrA   murJ_ΔVBS

cfu/ml (log scale) vs PenG Exposure (hours)

D

	Wild Type						ΔvxrAB						ΔVBS on P_murJ						min (100µg/ml PenG)
	0	20	40	60	90	180	0	20	40	60	90	180	0	20	40	60	90	180	

murJ

1   1.3±0.2   4±0.5   15±3.1   16±1.7   6±1.8   n.d   n.d

rRNAs

A

B

C

D

A

B

C

A

B

A

B

C