A multifaceted cellular damage repair and prevention pathway promotes high level tolerance to β-lactam antibiotics

Jung-Ho Shin[1, 2], Donghui Choe[4, 5], Brett Ransegnola[1, 2], Hye-Rim Hong[1, 2], Ikenna Onyekwere[1, 2], Trevor Cross[1,2], Qiaojuan Shi[9], Byung-Kwan Cho[4, 5, 6], Lars F. Westblade[7,8], Ilana L. Brito[9], and Tobias Dörr[1, 2, 3a]

[1] Weill Institute for Cell and Molecular Biology, Cornell, University, Ithaca, NY 14853, USA

[2] Department of Microbiology, Cornell University, Ithaca, NY 14853, USA

[3] Cornell Institute of Host-Microbe Interactions and Disease, Cornell University, Ithaca, NY 14853, USA

[4] Department of Biological Sciences, Korea Advanced Institute of Science and Technology, Daejeon 34141, Republic of Korea.

[5] KI for the BioCentury, Korea Advanced Institute of Science and Technology, Daejeon 34141, Republic of Korea.

[6] Intelligent Synthetic Biology Center, Daejeon 34141, Republic of Korea.

[7] Department of Pathology and Laboratory Medicine, Weill Cornell Medicine, New York, NY, USA.

[8] Division of Infectious Diseases, Department of Medicine, Weill Cornell Medicine, New York, NY, USA

[9] Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, 14850, USA

[a] corresponding author: tdoerr@cornell.edu

Abstract

Bactericidal antibiotics are powerful agents due to their ability to convert essential bacterial functions into lethal processes. However, many important bacterial pathogens are remarkably tolerant against bactericidal antibiotics due to inducible damage repair

responses. The cell wall damage response two-component system VxrAB of the

gastrointestinal pathogen *Vibrio cholerae* promotes high-level β-lactam tolerance and

controls a gene network encoding highly diverse functions, including negative control

over multiple iron uptake systems. How this system contributes to tolerance is poorly

understood. Here, we show that β-lactam antibiotics cause an increase in intracellular

free iron levels and collateral oxidative damage, which is exacerbated in the Δ*vxrAB*

mutant. Mutating major iron uptake systems drastically increased Δ*vxrAB* tolerance to β-

lactams. We propose that VxrAB reduces antibiotic-induced toxic iron and concomitant

metabolic perturbations by downregulating iron uptake transporters and show that iron

sequestration enhances tolerance against β-lactam therapy in a mouse model of

cholera infection. Our results suggest that a microorganism's ability to counteract

diverse antibiotic-induced stresses promotes high-level antibiotic tolerance, and

highlights the complex secondary responses elicited by antibiotics.

Introduction

The discovery and development of antibiotics is one of the most important biomedical

breakthroughs in human history. The introduction of agents that selectively kill bacteria

(bactericidal antibiotics) or inhibit their growth (bacteriostatic antibiotics) has directly

saved countless lives by enabling treatment of once common causes of death (*e.g.,*

pneumonia and sepsis) and indirectly facilitated other medical breakthroughs, including

transplantation medicine and chemotherapy [1,2]. Unfortunately, these significant

advances in medicine are threatened due to the rapid increase in the occurrence of

antibiotic treatment failure. We urgently need to develop new therapeutics and improve

the effectiveness of existing ones; this requires a precise understanding of antibiotic

mechanisms of action, and especially of bacterial strategies to avoid their noxious

effects. In principle, bacteria resist antibiotics by either developing the ability to grow in

their presence (antibiotic resistance, ABR) or to simply stay alive in their presence for

extended time periods (antibiotic tolerance/persistence)[3-8]. While the mechanisms and

consequences of ABR are relatively well-established, antibiotic tolerance remains poorly

understood, limiting our ability to effectively treat bacterial infections.

The β-lactam antibiotics (penicillins, cephalosporins, carbapenems, cephamycins and

monobactams) are highly potent bactericidal agents. Typically, their lethal action results

from their ability to simultaneously inhibit multiple targets (i.e., the transpeptidase

domain of multiple penicillin-binding proteins [PBPs]), which ultimately causes bacterial

cells to deplete essential cell wall precursors and self-destruct through the activity of

endogenous, cell wall lytic enzymes ('autolysins'; endopeptidases, amidases and lytic

transglycosylases) [9-12]. However, we and others have recently shown that many

clinically significant Gram-negative pathogens are remarkably tolerant to β-lactams. The

agent of cholera, *Vibrio cholerae,* the opportunistic pathogen *Pseudomonas aeruginosa*

and members of the *Enterobacterales* all survive treatment with β-lactams (including the

potent so-called "last resort" agent meropenem [a carbapenem]) by forming non-

dividing, cell wall deficient spheroplasts [11,13,14]. Upon removal of the β-lactam,

spheroplasts rapidly revert to wild-type shape and growth, providing a possible

explanation for treatment failure in the clinic. Spheroplasts are reminiscent of bacteria in

the so-called "L-form" state [15,16] with the crucial distinction that spheroplasts, unlike L-

forms, do not divide in the presence of antibiotics.

The extent of antibiotic tolerance is a reflection of the cell's ability to mount a stress

response to counteract primary damage resulting from antibiotic-target interactions [17,18].

Numerous bacteria encode cell envelope damage response systems (alternative sigma

factors and two-component response regulators) that sense damage to the cell wall or

membranes and consequently upregulate complex regulons aimed at damage repair [19-

25]. In *V. cholerae*, high-level β-lactam tolerance depends on the cell wall damage

sensing two-component system VxrAB (also known as WigKR)[17], which is broadly

conserved among *Vibrionaceae*. The vast VxrAB regulon is induced by cell wall-acting

antibiotics and includes cell wall synthesis functions and those involved in biofilm

formation and type VI secretion, *inter alia* [17,26,27]. How VxrAB promotes high level

tolerance against bactericidal β-lactams is unknown, but presumably relies on repair of

cellular damage inflicted by β-lactam antibiotics, *i.e.*, primarily cell envelope damage.

However, in addition to their primary mechanism of action, bactericidal antibiotics have

also been proposed to generate toxic reactive oxygen species (ROS) as a byproduct,

which are thought to enhance cell death [28-31]. This idea has been highly controversial

and the resulting debate is ongoing [32-37].

Here, we present evidence supporting a multifactorial role for VxrAB in promoting high-

level β-lactam tolerance. We have found that the ability to upregulate cell wall synthesis

only accounts for part of the high-level tolerance observed in wild-type *V. cholerae.* In

addition, we propose that VxrAB's role in promoting tolerance is to downregulate iron

acquisition systems which are inappropriately induced due to the accumulation of ROS

that are produced upon exposure to penicillin. Our data highlight the pleiotropic nature

of antibiotic mechanism of action and provide a blueprint for the mechanistic

underpinnings of the high-level β-lactam tolerance often observed in significant Gram-

negative pathogens.

Results

Spheroplast-mediated β-lactam tolerance is widespread in human pathogens

Recently, we and others have reported that clinically significant bacterial pathogens

exhibit high tolerance to β-lactam antibiotics via formation of cell wall deficient

spheroplasts when exposed to antibiotics in various growth media, including human

serum[11,13,14]. Here, we asked if spheroplast formation could also be observed in the

standard bacterial growth medium LB (Lysogeny Broth). We tested a well-

characterized[14] panel of clinical isolates of *Enterobacterales* as well as known

spheroplast-formers, *Pseudomonas aeruginosa* and *V. cholerae*, for their ability to

survive exposure to the carbapenem antibiotic meropenem. Meropenem is not

susceptible to hydrolysis by the AmpC β-lactamase which some members of the

Enterobacterales encode[38] and was thus appropriate for these experiments. Tolerance

levels for almost all isolates (with the exception of *E. coli* lab strain MG1655), were very

high (comparable to or higher than our previous observations in human serum and

Brain-Heart Infusion broth[14]) (**Fig. 1A**). Crucially, tolerance in all cases coincided with

the formation of membranous (membranes visualized by FM4-64 stain, **Fig. 1B**)

spheroplasts after 3h of incubation. Thus, β-lactam tolerance phenotypes are

recapitulated in LB, underscoring the usefulness of this growth medium to probe β-

lactam tolerance.

ChIP-Seq and RNA-Seq reveal a complex cell wall damage response

We previously elucidated the transcriptional network controlled by VxrAB upon

overexpression of a phosphomimetic version of VxrB (VxrB^{D78E}). Here, we directed our

attention to genes directly controlled by the VxrAB response. To this end, we performed

a ChIP-Seq experiment using a strain with functional, polyhistidine-tagged,

chromosomally encoded VxrB assayed during penicillin (PenG) treatment. After 180 min

of PenG exposure (100 µg/ml, 10 × MIC; same conditions used in our previous study[17]),

424 putative VxrB binding events were identified by comparison with a similarly treated

control strain expressing wild-type, untagged VxrB (**Fig. 2A,** see **Table S3** for complete

list). We then validated the ChIP-Seq results via electrophoretic mobility shift assays

(EMSA) of the 10 DNAs from the strongest VxrB binding events (**Fig. S1**) and, for a

subset, differential regulation via S1 nuclease assays of PenG-treated cells. We then

integrated this ChIP-Seq dataset with our re-analyzed (with more stringent parameters,

see Methods for details) previous RNA-Seq dataset of overexpressed VxrB^{D78E} [17] (**Fig.**

2B, see **Table S4** and **S5** for a complete list) to score VxrB binding events that

coincided with transcriptional up- or downregulation of the associated gene(s) (**Table**

S5). Our analysis resulted in the identification of 102 genes/operons that contained a

putative VxrB binding site and exhibited significantly increased, VxrB-dependent

expression levels, and 38 genes/operons that were scored as candidates for direct

binding and downregulation (**Fig. 2C**). Preliminary functional assignments using the

KEGG database [39] revealed that the VxrB regulon controlled a large number (76.4%

107/140) of genes of unknown function. Among the annotated upregulated genes, many

coded for functions related to cell envelope biology (ABC transporters, peptidoglycan

synthesis, and membrane synthesis) (**Fig. S2**). Among the downregulated genes, we

observed genes encoding proteins involved in iron uptake systems, flagella

biosynthesis, carbon metabolism, chemotaxis, cell wall lytic enzymes and

lipopolysaccharide biosynthesis (**Fig. S2**). Importantly, genes encoding cell wall

synthesis functions [*e.g.*, class A PBP1a (aPBP1a)], whose Vxr-dependent

transcriptional regulation we have extensively validated[17], exhibited strong VxrB binding

sites (**Fig. 2D**) and differential, Vxr-dependent regulation in S1 nuclease assays,

increasing our confidence in the robustness of the ChIP-Seq dataset. Both ChIP-Seq

and RNA-Seq also corroborated autoregulation of the VxrAB operon (**Fig**. **2B, Table**

S1). We noted that while many of the putative Vxr-controlled genes exhibited VxrB

binding in addition to differential regulation by phosphomimetic VxrB and VxrAB-

dependent regulation upon exposure to PenG (**Fig. 2D,** *pbp1a* and *zapB*), some genes

were apparently indirectly regulated as they clearly exhibited transcriptional

downregulation (both upon overexpression of VxrB^{D78E} and exposure to PenG) without

a clear signal in our VxrB ChIP-Seq experiment (**Fig. 2D,** *flaA*). We speculate that this

may reflect indirect regulation as a consequence of large scale metabolic alterations

that might be expected from induction of the VxrAB regulon. Thus, the VxrAB cell wall

stress response controls a gene network composed of diverse functions with an

apparent enrichment for functions required for cell envelope homeostasis and iron

acquisition.

DNA footprinting analysis identifies putative VxrB binding sites

Based on an alignment of putative VxrB binding sites, we used the MEME suite [40] to

identify the canonical VxrB consensus binding site ("Vxr box"). We based our analysis

on the 102 upregulated and the 38 downregulated genes separately, reasoning that

binding site variations may account for up- *versus* downregulation. This resulted in the

putative VxrB binding motif TTGACAAAA-N2-TTGAC (N = any nucleotide) for the

upregulated genes and CGCTCATTTTTTAACCAAGT for the downregulated genes.

However, these putative binding motifs were poorly conserved across all Vxr-controlled

genes. We hypothesize that this may be reflective of different binding characteristics of

the phosphorylated vs. non-phosphorylated forms of VxrB; both forms might be

expected to occur after PenG treatment and could bind to different motifs. To further

characterize the VxrB binding site we conducted DNase I footprinting assays using

purified VxrB and promoter regions of putative Vxr-controlled genes (**Fig. 3AB, Fig. S3**).

We again chose genes with the strongest VxrB binding as indicated in our EMSA and

ChIP-Seq experiments. We identified 10 promoters where VxrB protected a ~ 50 bp

sequence from DNase I digestion at low µM concentrations (**Fig. 3AB, Fig. S3**), again

corroborating direct binding to these promoters. However, apparent affinity varied

between promoters, with the promoters of *pbp1a*, encoding aPBP1a, *mgtE*, encoding a

magnesium transporter and *murJ*, encoding the lipid II flippase exhibiting the highest

degree of protection (**Fig. 3AB, Fig. S3)**. Surprisingly, we were again unable to

construct a well-conserved binding site for VxrB based on this subset of promoters,

despite three independent assays unequivocally demonstrating binding. This apparent

binding promiscuity suggests that VxrB might have the capacity to bind to a wide range

of promoter sequences *in vivo* and/or that it requires as yet unidentified accessory

factors to modulate binding.

Control of cell wall synthesis only partially explains VxrAB's requirement for β-

lactam tolerance

To elucidate the mechanism of β-lactam tolerance in *V. cholerae,* we sought to dissect

the individual contributions of VxrB-regulated genes to resistance to killing. Since the

primary mechanism of action of β-lactam antibiotics is inhibition of cell wall synthesis,

we first investigated the role of Vxr-induced PG biogenesis in antibiotic tolerance. We

took two approaches to eliminate VxrB's positive control over PG synthesis. For

aPBP1a, we engineered a strain with a deletion in the second, functionally semi-

redundant [41,42] aPBP1b, which also contains an IPTG-inducible PBP1a as its sole copy.

As expected, growth of this strain was IPTG concentration-dependent (**Fig. S4**). We

then grew this strain either in high concentrations of IPTG (high PBP1a levels) or very

low concentrations that markedly reduced growth rate (**Fig. S4**). Addition of PenG to low

aPBP concentration cultures resulted in only a modest (5 – 10-fold) decrease in viability

(**Fig. 3C**). This observation suggests that upregulation of PBP1a is indeed critical for

survival in the presence of PenG, but cannot account entirely for the steep, ~10,000-fold

decrease in post-antibiotic plating efficiency (**Fig. 3D** and [17]) of the Δ*vxrAB* mutant.

To corroborate these results, we also created a mutant with a partial deletion in the

putative VxrB binding site of the *murJ* gene as determined from our DNA footprinting

experiment. MurJ-mediated PG precursor translocation is a critical bottleneck for cell

wall synthesis and upstream of all periplasmic cell wall assembly functions [43,44]. We

thus expected that introducing a mutation that removes VxrB's positive control over this

step would allow us to gauge the significance of VxrB control over cell wall synthesis as

a whole. S1 nuclease assays confirmed that in the VxrB binding site deletion strain,

murJ expression was not significantly induced by PenG, similar to a Δ*vxrAB* strain (**Fig.**

3E). Similar to our observations with low PBP1a expression, the Vxr-independent *murJ*

strain exhibited 10-fold loss in viability upon exposure to PenG (**Fig. 3D**). We conclude

that the ability to upregulate cell wall synthesis functions via VxrAB accounts for only

some of its ability to promote tolerance.

The* VxrB *regulon controls iron acquisition systems

Based on our previous RNA-Seq data, VxrB downregulates genes involved in iron

acquisition [17]. Indeed, examination of the RNA-Seq/ChIP-Seq datasets revealed a high

overlap between genes regulated by VxrB with those regulated by the iron starvation

response repressor Fur [45]. 47 out of the putative 63 Fur-regulated genes were also

downregulated by VxrB and 20 out of these 63 genes contained both a putative VxrB

and a Fur binding site (**Fig. 4AB**). Thus, the VxrB system appears to control multiple

functions required for iron homeostasis.

Given VxrAB's apparent role in downregulating iron acquisition in response to cell wall

damage, we expected that surplus iron accumulates in a PenG-treated $\Delta vxrAB$ mutant

and that this might be detrimental to cells under these conditions. To test this, we

measured intracellular iron levels using ICP-MS during treatment with PenG. Before

treatment, both the wild-type and $\Delta vxrAB$ mutant strains contained similar iron levels at

~280 ng/mg protein (**Fig. 4C**). After 1 h of PenG exposure, wild-type iron levels

remained relatively constant while the iron content of the $\Delta vxrAB$ strain started to

increase significantly by ~60% compared to baseline. Iron levels increased slightly in

the wild-type strain between 90 min and 3 h, while levels in the $\Delta vxrAB$ mutant

remained significantly higher. Consistent with an increase in free intracellular iron, the

wild-type strain also became more sensitive to the iron-dependent quinone antibiotic

streptonigrin [46,47] upon treatment with PenG (**Fig. 4D).** Thus, PenG exposure causes

enhanced iron influx.

Penicillin exposure induces the Fur regulon

Why do PenG-treated $\Delta vxrAB$ cells increase their iron load? Exposure to β-lactam

antibiotics has been shown to result in the accumulation of H_2O_2 in other bacteria [31,48,49]

and we hypothesized that hydrogen peroxide might directly activate Fur via oxidation of

its iron cofactor. Derepression of the Fur regulon in turn would result in the upregulation

of iron acquisition systems. We first confirmed that PenG exposure resulted in H_2O_2

production in *V. cholerae* as well. We used a highly sensitive, fluorimetric assay

(Amplex Red) to measure intracellular hydrogen peroxide concentrations. Upon

exposure to PenG, both the wild-type strain and $\Delta vxrAB$ mutant experienced a transient

increase in hydrogen peroxide concentrations, with a peak between 20 and 60 minutes,

followed by a decrease back to baseline. However, the hydrogen peroxide concentration

in the *vxrAB* mutant was significantly elevated and at peak (20 - 60 min) accumulated

more than double the concentration measured in wild-type cells (**Fig. 5A**). As expected,

and consistent with the accurate ability of Amplex Red to report hydrogen peroxide, a

Δ*oxyR1* Δ*katB* Δ*katG* triple mutant, which lacks the ability to elaborate the primary

hydrogen peroxide detoxification systems, exhibited both higher baseline and PenG-

induced levels of hydrogen peroxide than the wild-type and Δ*vxrAB* mutant; additionally,

external spiking of catalase into our assay reactions completely abolished the signal

(**Fig. 5A**)*.* We also measured hydrogen peroxide in the supernatant of PenG-treated

cells, since hydrogen peroxide might diffuse through membranes to some degree. We

did not, however, note a meaningful accumulation of peroxide in the supernatant of any

of our mutants (**Fig. S5A**), demonstrating that at least for the duration of our

measurements, there was no significant diffusion out of the cell.

Next, we tested whether the observed accumulation of peroxide resulted in upregulation

of the *fur* regulon. We used S1 nuclease mapping to monitor transcript levels of the

canonical Fur regulon readout *ryhB*[50], a small RNA [51]. Consistent with our line of

thinking, *ryhB* was highly induced by addition of external hydrogen peroxide **(Fig. 5B).** In

E. coli, sodB transcript levels are negatively controlled by *ryhB,* and consistent with a

similar regulatory mechanism in *V. cholerae, ryhB* induction (**Fig. 5B**) coincided with a

disappearance of the *sodB* transcript during treatment with hydrogen peroxide (**Fig.**

5B)*.* Consistent with a regulation via iron starvation sensing, the *ryhB* transcript also

increased dramatically upon exposure of cells to the cation chelator EDTA (**Fig. S5B**).

Upon PenG exposure, *ryhB* transcript was reproducibly induced ~2-fold in both the wild-

type and ∆*vxrAB* strains after 20 min, followed by a return to baseline. Other Fur-

regulated genes (*hutA, hutB* and *vc2212*), also exhibited a marked, and in some cases

transient, upregulation upon PenG exposure (**Fig. 5B**). After prolonged PenG-treatment

(180 min), *ryhB* remained uninduced in the wild-type, but increased dramatically in the

∆*vxrAB* mutant (**Fig. S5B**). Consistent with these transcriptional changes, a P$_{ryhB}$-

msfGFP transcriptional reporter construct also increased in fluorescence after exposure

to PenG in both wild-type and ∆*vxrAB* strains (**Fig. 5C**), to similar degree as in the

positive control, the metal chelator EDTA. These results suggest that the Fur regulon is

indeed induced by exposure to PenG. Interestingly, we noticed in our ICP-MS dataset

that concomitantly with iron uptake, manganese concentrations also increased in PenG-

treated cells (**Fig. S5C**). Manganese has been proposed to be a cellular response to

"rescue" iron-dependent enzymes that have lost their iron cofactor due to iron starvation

or oxidative damage; biologically appropriately, manganese ion uptake is thus under

control of Fur in Gram-negative model organisms [52-54]. While a specific manganese ion

importer has not been identified in *V. cholerae* [55], our data suggest that the cholera

pathogen might possess a similar regulatory arrangement.

Interfering with iron uptake partially restores tolerance to the ∆vxrAB mutant

If PenG-induced iron influx is detrimental, we would expect iron chelators to alleviate the

∆*vxrAB* β-lactam sensitivity defect. Indeed, addition of 250 µM (0.25 x MIC) of the

intracellular iron chelator 2,2' bipyridyl reproducibly increased survival of the ∆*vxrAB*

mutant 50 – 100-fold (**Fig. S6**); the ability to partially rescue could be reverted by

addition of iron to the growth medium. To probe the involvement of iron uptake systems

even further, we sequentially deleted known iron uptake systems in the Δ*vxrAB*

background and measured survival in the resulting mutants in the presence of PenG.

Several combinations of Δ*vxrAB* with iron uptake system mutants reduced killing,

particularly at early timepoints (**Fig. S7A-C**). Crucially, the Δ*vxrAB* Δ*vctPDGC* Δ*feoABC*

Δ*fbpABC* mutant rescued the tolerance defect by ~100-fold (**Fig. 6A**). The Vct, Feo and

Fbp systems encode uptake systems for the siderophore vibriobactin, ferrous and ferric

iron, respectively [56]; their importance is consistent with them likely being the main

drivers of iron uptake under our experimental conditions (LB broth). In addition, we

serendipitously found that the Δ*vxrAB* mutant is growth-defective on 10% sucrose

plates, and this was exacerbated by addition of iron to the plate (**Fig. 6B**). Strikingly,

deleting Δ*vctPDGC* Δ*feoABC* Δ*fbpABC* in the Δ*vxrAB* background completely restored

growth on sucrose with or without 100 μM $FeSO_4$ (but did not substantially increase

plating efficiency at 1 mM $FeSO_4$) (**Fig. 6BC**). Similar to our observations in PenG,

combinations of Δ*vxrAB* with different combinations of iron uptake mutants had varying

effects on restoration of growth on iron-supplemented sucrose (**Fig. S7 D-F**); most

notably a mutation in the siderophore uptake system VctP itself was able to significantly

improve plating efficiency of the Δ*vxrAB* mutant at lower iron concentrations (**Fig. S7D**).

Thus, under cell envelope stress conditions, iron uptake significantly contributes to

Δ*vxrAB* cell death or growth inhibition.

The iron chelator deferoxamine promotes β-lactam tolerance in a mouse model of

cholera infection

Since our data thus far suggest iron influx is detrimental to tolerance in the absence of

mitigating factors, we reasoned that the clinical use of iron chelators like

deferoxamine[57-61] may increase bacterial antibiotic tolerance as a collateral effect. We

employed a mouse model of cholera infection (see Methods for details) to test this

hypothesis *in vivo*. Streptomycin-treated mice were infected with *V. cholerae* and then

treated with either deferoxamine, the oral β-lactam antibiotic amoxicillin, or both (**Fig.**

7A). *In vivo* pathogen survival under antibiotic pressure was quantified by assessing

time-resolved viable cell counts from stool samples. As expected, viable counts from

amoxicillin-treated mice drastically decreased over time (**Fig. 7B**), while deferoxamine

alone did not affect pathogen survival compared to vehicle control. Strikingly, however,

addition of deferoxamine to amoxicillin treatment completely undermined the latter

antibiotic's lethal effect and induced high-level amoxicillin tolerance (**Fig. 7B**). Thus,

limiting available iron reduces antibiotic effectiveness *in vivo*; these results are also

consistent with a model where detrimental iron accumulation results from β-lactam

exposure.

Penicillin exposure induces ROS defense systems and oxidative damage

Iron is thought to be toxic to cells due to its ability to convert hydrogen peroxide into

oxygen radicals that pleiotropically oxidize and damage macromolecules such as

proteins and DNA [62] and generally disturb metal homeostasis [63]. Indeed, antibiotics

have been proposed to induce the increased production of ROS through metabolic

perturbations [29,31,37], however, the extent to which this contributes to killing by antibiotics

remains highly controversial [32-34]. To interrogate the potential significance of ROS to

lethality of the Δ*vxrAB* mutant, we first assessed whether oxidative stress response

systems were induced by PenG. We found that PenG rapidly (within ~20 min) induced

members of the hydrogen peroxide responsive OxyR1 regulon (*prxA* and *dps*) [64-66] in

the wild-type strain (**Fig. 8A**). Transcript levels of both OxyR1 readouts remained high

initially and then decreased to below baseline after ~ 60 min (*prxA*) or 180 min (*dps)*,

suggesting that the wild-type strain induces a response that detoxifies (*prxA*) or protects

against (*dps*) hydrogen peroxide (**Fig. 8A, S8A**). Both transcripts are induced by

hydrogen peroxide in an OxyR1-dependent manner [64,65] and we validated

responsiveness to hydrogen peroxide for the *prxA* transcript in our *V. cholerae* wild-type

strain **(Fig. S8B).** Thus, *prxA* and *dps* are *bona fide* readouts for hydrogen peroxide

production. Interestingly, the transcript for the response regulator OxyR1 itself was

upregulated upon PenG exposure, with dynamics similar to those observed for

members of its regulon (**Fig. 8A**). In the model organism *E. coli*, *oxyR* transcript levels

do not change under hydrogen peroxide exposure [67] and our results suggest that *V.*

cholerae might employ a different mode of regulation of its oxidative stress responses.

A return to baseline coincided with the upregulation of *V. cholerae*'s two catalases (*katB*

and *katG*; only *katG* appears to be under OxyR1 control in the cholera pathogen [65,66]) at

later stages of antibiotic exposure (**Fig. 8A**, **Fig. S8B**), likely revealing the detoxification

mechanism that enables the repression of the OxyR1 regulon. Consistent with its

hydrogen peroxide sensing function, KatG was also induced by hydrogen peroxide (**Fig.**

S8A).

In the ∆*vxrAB* strain, *oxyR1* baseline levels were slightly increased compared to the

wild-type, while *prxA* and *dps* levels were similar. After exposure to PenG, *oxyR1* levels

decreased at first but the transcript remained detectable and above the wild-type

baseline for longer than the wild-type background (**Fig. 8A, S8**). The *prxA* and *dps*

transcripts exhibited a dynamic pattern with strong induction (*dps)* or transient

downregulation (*prxA*) in the mutant. We observed a pattern similar to *oxyR1* with a

gene coding for another oxidative stress mitigation factor, the alkyl hydroperoxide

reductase C, *ahpC* (**Fig. S8**). Interestingly, the transcript for *katG* remained uninduced

in the *vxrAB* mutant.

We confirmed that transcriptional regulation also resulted in increased catalase protein

activity using a gel-based activity assay for catalases (**Fig. S9A,** see Methods for

details) after exposure to PenG. We also included an assay for superoxide dismutase

(SOD) activity, since SODs are broadly conserved, important ROS scavenging

enzymes[68]. Consistent with the observed transcript levels (**Fig. 8B**), catalases were only

induced in the wild-type, but not the mutant (**Fig. 8B**). In contrast, SOD activity

increased similarly in both the wild-type and the *vxrAB* mutant (**Fig. 8C**). Taken

together, these results suggest that both the wild type and ∆*vxrAB* induce responses

upon exposure to PenG that are associated with oxidative stress, albeit with different,

complex dynamics.

We next assessed the cellular consequences of oxidative damage using a protein

oxidation assay via Western Blot. Consistent with our results showing enhanced iron

sensitivity of the mutant, protein oxidation levels were increased in the $\Delta vxrAB$ mutant

vs. wild-type even prior to addition of antibiotic (**Fig. 8D**). Addition of PenG resulted in a

rapid (within 20 min) increase in protein oxidation levels in the wild-type, which

decreased after 40 min but stayed above baseline throughout β-lactam exposure. In

contrast, the *vxrAB* mutant steadily accumulated oxidized proteins throughout the

exposure time, corroborating that oxidative stress has a more profound effect on the

vxrAB mutant than the wild-type strain. Both wild-type and the *vxrAB* mutant exhibited

increased protein oxidation when exposed to hydrogen peroxide as a positive control

(**Fig. 8D, Fig. S9B**). Taken together, these data suggest that both wild-type and $\Delta vxrAB$

experience oxidative stress in the presence of PenG, but the $\Delta vxrAB$ mutant is unable

to mount a mitigating response.

The electron transport chain is required for both maximal killing and hydrogen

peroxide production

How does PenG treatment result in production of ROS? A possible origin is the electron

transport chain (ETC), which in *V. cholerae* is a major generator of ROS under normal

growth conditions [69]. To test the ETC's involvement in PenG-induced ROS production,

we created mutants in two ETC components, namely *V. cholerae's* principal sodium

translocating NADH dehydrogenase NqrA [70] and *ubiA*, which encodes a key initial step

towards ubiquinone biosynthesis. NqrA has been shown to be a significant source of

ROS under normal growth conditions [69], but for both mutants, we expect a drastic

reduction in ETC electron flow. Crucially, neither mutation resulted in a significantly

decreased growth rate when grown in medium containing the fermentable carbon

source glucose (albeit, both mutations caused a slight increase in lag phase) (**Fig.**

S10A). Reduced growth rates can affect β-lactam susceptibility non-specifically [71] and

mutants conferring slower growth can thus confound the interpretation of experiments

conducted with such strains. Consistent with the expectation of reduced ETC activity in

these mutants, disruption of *nqrA* and *ubiA* in both the WT and the ∆*vxrAB* mutant

background resulted in a marked reduction in PenG-induced accumulation of hydrogen

peroxide, as measured by Amplex Red (**Fig. S10B**). To test the hypothesis that the

reduction in hydrogen peroxide also results in lack of killing in the ∆*vxrAB* mutant, we

conducted time-dependent killing experiments with these strains. As observed

previously, wild-type viability remained stable for the entire duration of the experiment

while the ∆*vxrAB* mutant displayed a ~10,000-fold decrease in viability. The ∆*nqrA* and

∆*ubiA* mutants were as tolerant as the wild-type. However, a combination of ETC

mutations with ∆*vxrAB* reproducibly increased survival by 10- (∆*nqrA*) or 100-fold

(∆*ubiA*) (**Fig. 8E**). Thus, ∆*vxrAB* lethality after PenG exposure partially depends on the

presence of the ETC.

ROS detoxification systems promote tolerance at high cell densities

We next asked whether ROS damage mitigation systems were required for wild-type

tolerance. We created knockout mutants deficient in the iron starvation repressor Fur

(expected to result in an increase in intracellular iron and thus enhanced ROS), SodB

and a triple mutant defective in both catalases and OxyR1. We grew all strains from late

exponential into stationary phase and periodically withdrew aliquots that were then

exposed to PenG for 3 h, and measured survival. As a comparison, we subjected the

$\Delta vxrAB$ mutant to the same treatment. While all mutants except for $\Delta vxrAB$ exhibited

wild-type survival patterns in exponential phase, entry into stationary phase coincided

with significant killing in all mutants that are expected to experience enhanced ROS

production (**Fig. 8F**). Consistent with these results, *sodB* had previously answered a

Transposon Insertion Sequencing (TnSeq) screen for factors that are required for PenG

tolerance [72]. We thus propose that ROS contribute significantly to β-lactam-mediated

killing in stationary phase (see Discussion for details).

Discussion

Our overall model of VxrAB-mediated tolerance is depicted in **Fig. 9;** based on the

evidence presented in this study, we propose the following: PenG induces cell wall

degradation, resulting in perturbations of the ETC. Our work does not directly address

the nature of this perturbation, but it may simply result from topological alterations of the

cytoplasmic membrane brought about by loss of the cell wall. Interestingly, and

consistent with this line of thinking, other cell wall deficient cells (L-Forms) also appear

to accumulate ROS[73,74]. We hypothesize that partial reduction of molecular oxygen

through the ETC (particularly NqrA) results in the production of superoxide ion.

Superoxide ions are removed by SOD, which catalyzes the formation of hydrogen

peroxide. Hydrogen peroxide induces the *oxyR1* regulon, but we propose that it also

inappropriately inactivates the Fur repressor via oxidation of its ferrous ion cofactor.

Inactivation of Fur (i.e., induction of the Fur regulon) results in the upregulation of iron

uptake systems, which increases the intracellular iron pool. Intracellular iron reacts with

hydrogen peroxide to form hydroxyl radical, which damages proteins and other

macromolecules. The wild-type strain mounts effective responses to all these types of

damage. SOD, AhpC and catalases detoxify superoxide ions and hydrogen peroxide.

VxrAB induces cell wall synthesis functions to prepare cells for speedy recovery once

the antibiotic is removed. At the same time VxrAB also downregulates some iron

acquisition genes and other Fur functions to dampen detrimental iron influx and

potentially concomitant oxidative damage. Interestingly, some of our data suggest that

ROS production might contribute significantly to killing in later growth phases (early

stationary phase), and stationary phase-specific oxidative damage is not without

precedent. Periplasmic SODs are under control of stationary phase regulators in diverse

bacteria and yeast [75-77] and SOD is required for antibiotic tolerance in stationary phase

in *P. aeruginosa* as well [78]. We hypothesize that during rapid growth in exponential

phase, the primary mechanism of action (cell wall degradation) is likely the more

important contributor to cell death, while in stationary phase (slower/no growth, less cell

wall damage), the harmful effects of ROS accumulation may become more visible. This

is noteworthy, since in the natural environment, bacterial growth can be expected to

resemble stationary phase more than rapid growth in exponential phase [79]. On a

separate note, β-lactam-mediated killing in stationary phase may seem counterintuitive

at first, since at least in the well-characterized model organism *E. coli*, cessation of

growth upon entry into this growth phase is generally assumed to result in complete

protection from β-lactam-induced damage [71,80]. However, the *vxrAB* mutant still

exhibited significant PenG-dependent mortality even when PenG exposure only

occurred several hours into stationary phase, i.e., far into a period of non-growth,

suggesting that this mutant experiences PenG-induced cell envelope damage even

when populations stop growing. We thus propose that in *V. cholerae,* cell wall turnover

(and thus PenG's ability to inflict damage) continues at least for some time after cells

stop growing. A similar phenomenon could explain the unusual observation that the

Lyme disease pathogen *Borrelia burgdorferi* remains susceptible to killing by cell wall

acting antibiotics in stationary phase [81].

The connection between antibiotic exposure and ROS production and particularly the

extent of the contribution of ROS to killing is still actively debated in the field. We agree

with a previous hypothesis[34] that some of the interpretive difficulties in other studies

may have arisen from the low concentrations of antibiotics used (often around the MIC).

High concentrations of bactericidal antibiotics might kill cells too quickly through their

primary mechanism of action to measure a response to the proposed secondary

mechanism. Lower concentrations around the MIC might result in day-to-day variation

of killing dynamics that are difficult to reproduce under different conditions. In contrast,

bacteria like *V. cholerae* (as well as *P. aeruginosa, Klebsiella pneumoniae, Klebsiella*

(formerly *Enterobacter*) *aerogenes* and *Enterobacter cloacae),* exhibiting almost

population-wide tolerance to high concentrations of β-lactams [11,13,14,72], may be more

appropriate model organisms to dissect the layers of physiological changes occurring as

a consequence of antibiotic exposure. In these tolerant bacteria, the primary

mechanism of action (degradation of the cell wall) is recapitulated; however, unlike in

other bacteria, this does not result in lysis and cell death, allowing us to examine a large

population of live cells after treatment with high concentrations (10 × MIC or more) of

antibiotic.

One of the main critiques of the idea that ROS contribute to killing was that some

groups failed to measure meaningful ROS levels (and the expected responses to them)

in antibiotic-treated cells in the model organism *E. coli* [33,34]. This has been challenged

by other groups[35,37,82]. Here, we have shown that *V. cholerae* accumulates ROS and

induces defense systems against them after exposure to PenG. We propose that in the

vxrAB mutant, antibiotic-induced lethality is multifactorial. Both, cell wall defects and

ROS may contribute to killing. Importantly, other detrimental consequences of iron influx

may also play a role, for example mismetalation of key enzymes [63]. While ROS

production and iron influx are measurable in the wild-type as well, albeit to a lower

degree than the mutant, it is likely that its multiple, highly effective defense systems

prevent antibiotic killing. This thinking is consistent with a previous observation that

compared to *E. coli* and *P. aeruginosa,* exposure of *V. cholerae* to sub-MIC of β-lactam

antibiotics only caused a moderate increase in mutagenesis (a possible consequence of

ROS accumulation) [83]. Thus, the cholera pathogen's high tolerance to β-lactam

exposure might be the consequence of an exceptional ability to repair cell damage.

Since we demonstrate here that PenG-treatment induces toxic iron uptake, it is

important to note that iron withdrawal is a common antibacterial strategy in the human

host (and deferoxamine is clinically used to treat iron homeostasis disorders), which

might significantly affect antibiotic susceptibility *in vivo*[84]. The extent of the host's ability

to control iron levels in a specific infection locale might significantly and divergently

influence antibiotic susceptibility in different host tissues. In summary, β-lactams (and

likely other antibiotics) elicit more complex cellular responses than just those associated

with their primary mechanism of action. A more context-, metabolism- and physiology-

centered view of antibiotic action [85-88] might therefore yet yield critical insight to enable

the development of novel adjuvants that increase the efficacy of antibiotics in the

complex infection environment. Lastly, with or without ROS, our data suggest that

targeting bacterial stress responses could be a powerful strategy to enhance the activity

of bactericidal drugs against tolerant pathogens.

Acknowledgments

Research in the Dörr laboratory is supported by NIH grants R01AI143704 and

R01GM130971. This work was partially supported by the Korea Bio Grand Challenge

Project (2018M3A9H3024759 to B.K.C.) through the National Research Foundation of

Korea (NRF) funded by the Ministry of Science and ICT, Republic of Korea. Research in

the Brito Laboratory is funded by the Packard Foundation, Sloan Foundation, Pew

Foundation and NIH grant 1DP2HL141007. We thank the Helmann lab (Cornell

University) for sharing equipment and reagents. We further thank Stavroula Hatzios

(Yale University), John Helmann (Cornell University), Joseph Peters (Cornell

University), James Imlay (University of Illinois) and members of the Dörr lab for helpful

comments.

Figure legends

Figure 1 | Spheroplast formation in diverse bacterial pathogens. (A) Time-dependent killing of a panel of bacterial isolates in the presence of meropenem (MEM10, 10 µg/ml). Cells were grown overnight and sub-cultured for 30 minutes at 37°C before being exposed to meropenem at 10 µg/mL and moved to a 37°C stationary incubator. Samples were taken at each indicated time point and spot-titered onto LB medium to quantify colony forming units (CFU/mL) throughout meropenem treatment. **(B)** images of cells corresponding to the time points in **(A)** were taken after labeling with membrane stain FM4-64. *E. cloacae, Enterobacter cloacae; K. aerogenes, Klebsiella aerogenes; K. pneumoniae, Klebsiella pneumoniae; E. coli, Escherichia coli*

Figure 2 | ChIP-Seq and RNA-Seq analysis reveals a complex VxrB regulon. **(A)** Overview of VxrB-binding profiles across the *V. cholerae* genome after β-lactam exposure (PenG 100µg/ml [10x MIC] for 3 hours). Red peaks indicate VxrB binding sites above the significance cutoff (q-value<10^{-10}; MACS). **(B)** RNA-Seq MA plot is presented as fold change between VxrB^{D78E} overexpressed sample and control sample. Red dots indicate differentially expressed genes after VxrB^{D78E} overexpression (P-value < 0.001). Representative VxrB target genes are marked on both chromosomes and on the MA plot. **(C)** Venn diagram depicting overlap between RNA-seq and ChIP-Seq datasets **(D)** Regulatory modes of representative VxrB-controlled genes in response to PenG. Upper plot shows VxrB binding patterns (assessed via ChIP-Seq), lower panel shows transcriptional response upon overexpression of VxrB ^{D78E} (assessed via RNA-Seq). RNA expression was confirmed by S1 nuclease mapping assay in wild type and *Δvxr*AB mutant strains after exposure to PenG (100 µg/ml for 3 hours).

Figure 3 | Peptidoglycan synthesis control only partially explains *Δvxr*AB lethality.

(A and B) DNase I footprinting results for VxrB interaction with promoters controlling cell wall synthesis functions. Red boxes indicate core DNA regions protected by purified VxrB protein (final concentration 0.17 - 1.38 µM) on the *murJ* (A) or *pbp1A* (B) promoter. Identified transcriptional start site (TSS; +1) and promoter elements (-10 and -35) were derived from published datasets[89,90], and are presented with the VxrB binding site. VBS, putative **V**xrB **B**inding **S**equence. **(C-E)** Reduction of cell wall synthesis gene expression causes a tolerance defect. **(C)** Survival of a $\Delta pbp1b$ $\Delta pbp1a$ P$_{iptg}$:*pbp1a* strain after 6 hours of PenG exposure. P$_{up}$ indicates conditions of high IPTG concentrations (100 µM), P$_{down}$ low IPTG concentration (2 µM). Survival fraction is cfu/ml of PenG-treated cultures normalized to untreated cultures grown in and plated on medium containing the same IPTG concentration. (**D**) Time-dependent killing experiment in the presence of PenG (100 µg/ml, 10 x MIC). All data are means (+/- standard error) of 3 independent biological replicates. **(E)** Deleting the putative VxrB binding site from the *murJ* promoter abrogates induction of *murJ* by PenG exposure. Transcript levels were measured by S1 nuclease mapping. Numbers represent induction as measured by band intensity, normalized to pre-exposure WT. n.d., not determined.

Figure 4 | VxrAB controls genes involved in iron homeostasis. (A) Venn diagram showing overlap between the VxrB regulon and known *V. cholerae* Fur-controlled genes. (as per RegPrecise3.0 DB, http://regprecise.lbl.gov/RegPrecise/). 20 out of 47 Fur-

regulated genes also have a VxrB binding site and are differentially regulated by

VxrB[D78E]. **(B)** Genomic organization of iron uptake systems and their putative Fur/VxrB

binding sites. **(C)** Time-course of iron accumulation in PenG-treated WT and Δ*vxrAB*

cells as measured by ICP-MS. Data are average of 3 independent biological replicates,

error bars represent standard deviation. ∗∗P=0.001; ∗∗∗P<0.0003 (paired t-test). **(D)**

Streptonigrin-sensitivity of PenG-treated cells. Cultures were grown to a density of ~2 x

10^8 cfu/ml. Before ("- PenG") and after addition of PenG (100 µg/ml, 10x MIC), ~ 5 x 10^7

cells were spread on an LB agar plate. In the PenG-treated samples ("+ PenG"), beta

lactamase solution was also added to remove the antibiotic. Streptonigrin (5 µL of 5

mg/ml, in DMSO) was then added on a filter disk, followed by incubation for 24 h at 37

˚C. Values represent average diameter of clearance zones (+/- standard deviation) of

six biological replicates. SNG, streptonigrin.

Figure 5 | PenG induces hydrogen peroxide production and the Fur iron

starvation response. (A) Total intracellular hydrogen peroxide (H_2O_2) levels were

measured using Amplex[TM] Red Hydrogen Peroxide/Peroxidase Assay Kit and

normalized to total cellular protein level (measured by Qubit[TM]). Data are average

values from three independent experiments, error bars represent standard deviation.

Cat, catalase (1µg/ml). **(B)**. To test PenG-mediated induction of Fur-regulated genes,

S1 nuclease mapping assays were performed on the small RNA *ryhB,* the RyhB target

sodB, *hutA/B*, coding for heme transport systems and *vc2212*, which encodes a

component of a ferric citrate uptake system). Cells were grown to mid-exponential

phase followed by exposure to H_2O_2 or PenG for the indicated duration. Numbers

represent expression levels normalized to untreated control (averages from three

independent replicates). **(C)** To confirm the induction of *ryhB* transcript by PenG

treatment, the native *ryhB* promoter was fused with *msfGFP* and introduced into WT

and *ΔvxrA* mutant cells. Each strain was cultured until mid-exponential phase, when

1mM EDTA or 100µg/ml PenG were added for the indicated duration. Scale bar (5 µM).

Quantified GFP fluorescence intensities are presented with average values from 3

independent biological replicates (cell, n>500), error bars represent standard deviation.

∗∗∗∗ P<0.0001 (paired t-test).

Figure 6 | Disruption of iron influx systems partially restores tolerance of the

ΔvxrAB mutant against PenG exposure. (A) Time-dependent killing experiment in the

presence of PenG. **(B)** and **(C)** Iron uptake mutants restore *ΔvxrAB* growth on 10%

sucrose. The indicated strains were grown to exponential phase (OD$_{600}$~0.5) in LB

medium and spot-plated on 10% Sucrose plates with or without additional iron sulfate.

An example is shown in **(B),** quantification of cfu is shown in **(C).** Data are average of 3

independent biological replicates, error bars represent standard deviation. ∗∗∗ P<0.0003;

∗∗∗∗ P<0.0001 (paired t-test).

Figure 7 | The iron chelator deferoxamine (DFOA) promotes amoxicillin (AMX)

tolerance *in vivo*. (A) Schema of experimental design. Briefly, mice were pretreated

with streptomycin (5 mg/mL) for 2 days and then intragastrically inoculated with *V.*

cholerae (~1.97x10^{10} cells in 200 µl PBS). After 24 h, mice in each group received water

(H$_2$O), amoxicillin (50 mg/kg), deferoxamine (200 µl of a 100 µM solution), or both

amoxicillin and deferoxamine. **(B)** Excretion of *V. cholerae*. Fecal samples were

collected every two hours and assessed for their viable cell content (Colony Forming

Units, CFU) to monitor the level of *V. cholerae* excretion. Each symbol represents fecal

samples from an individual mouse (n=4). The number of bacterial CFU was normalized

per milligram of feces (CFU/mg). ∗∗ P<0.005; ∗∗∗ P<0.001 (two-way ANOVA).

Figure 8 | PenG exposure is associated with ROS production (A) RNA expression

profiles of first line ROS defense systems in *V. cholerae* monitored by S1 nuclease

mapping assay. Quantified average values (normalized to WT before PenG exposure)

from three independent experiments are presented under the signals of each gene. **(B**

and C) Quantification of the SOD and catalase activity values from the gel shown in **Fig.**

S9. Data are average values from three independent experiments normalized to

untreated wild type. **(D)** Detection of protein oxidation after PenG treatment. To

measure total cellular protein oxidation, OxyBlot Protein Oxidation Detection Kit (Sigma-

Aldrich) was used. For positive protein oxidation control, oxidized standard marker ($_{OX}$M)

was used. Both DNP-derivatized or nonderivatized (negative control) protein samples

were separated by polyacrylamide gel electrophoresis followed by Western blotting

with-DNP specific antibody. **(E)** Time-dependent killing of respiratory chain mutants

($\Delta nqrA$, encoding NADH dehydrogenase and $\Delta ubiA$, encoding an early step of

ubiquinone synthesis) in the presence of PenG (100 µg/ml, 10xMIC). All strains were

grown in LB 0.2 % glucose. **(F)** Growth-phase dependent killing assay of ROS

detoxification systems (Fur, SodB, OxyR1, KatG, and KatB). At designated time-points,

a 5 ml aliquot was withdrawn and exposed to 100 µg/ml PenG for 3 h. CFU/mL for each

time point was assessed before (solid lines) and after addition of PenG (dotted lines).

Data are mean of three independent biological replicates and error bars represent

standard deviation.

Figure 9 | Proposed model for VxrAB-mediated antibiotic tolerance. PenG induces

ETC dysfunction during transition of cell shape from rod to spheroplast, resulting in the

production of ROS. Hydrogen peroxide illegitimately induces the Fur regulon, resulting

in the upregulation of iron acquisition systems, resulting in enhanced iron influx and

lethal oxygen radical generation. VxrAB dampens this response by downregulating iron

acquisition genes, in addition to its primary role as a positive regulator of cell wall

synthesis genes. SO, superoxide.

MATERIALS AND METHODS

Bacterial strains and culture conditions.

All *V. cholerae* strains used in this study were derivatives of *V. cholerae* WT El Tor

strain N16961 or E7946 (for chitin-induced transformation) listed in **Table S1**. *V.*

cholerae was grown on Luria-Bertani (LB) medium with or without 0.2% glucose or in

M9 medium containing 0.2% glucose at 37°C unless otherwise indicated. All media

contained 200 µg/ml streptomycin (to which *V. cholerae* is resistant). Where applicable,

growth media were supplemented with 100 µg/ml Benzylpenicillin potassium salt (PenG)

(Fisher BioReagents, cat# BP914-100), isopropyl-β-D-1-thiogalactopyranoside (IPTG;

final 0.1~1mM) (GOLDBIO, cat# 12481C50), or catalase 1µg/ml (Sigma-Aldrich, Cat#

c9322)**.** *Escherichia coli* DH5α and DH5α λpir were used for routine DNA

cloning. Unless indicated otherwise, LB liquid media were inoculated from an overnight

culture and incubated at 37°C with shaking at 200 rpm until reaching mid-exponential

phase (OD_{600} ~0.4). Where applicable, trimethoprim (Sigma, cat #T7883, dissolved in

DMSO) was used at 15 µg/ml final concentration.

Strain construction

Deletion mutants were generated via replacement of the open reading frame/binding

site with the scar sequence TAATGCGGCCGCACTCGAGTAATAATGATGA or, where

indicated, a trimethoprim resistance cassette. Knockouts were generated either by

homologous recombination using the suicide vector pCVD442 [91] or by chitin-induced

natural transformation, both as described before [92] [17]. The following primer

combinations were used (see supplemental material for sequence information).

murJ Δ*vbs*: TDP1062/1064 + TDP1066/1067, fused via stitching PCR with nesting

primers TDP1078/1079, followed by chitin transformation using *vc1807*::trim [generated

using primers TDP597/598 + TDP599/600 (on trimR from the *V. cholerae* HAITI strain

[93])+TDP601/602, fused via stitching PCR with nesting primers TDP603/604] as primary

selection. lacZ::P$_{ryhB}$:*msfGFP*: TDP1152/1235 + TDP1237/38, cloned into *SmaI*-

digested pTD101, a suicide vector for chromosomal integration into the *lacZ* locus [95].

Δ*nqrA*: TDP1388/1389 + TDP1390/1391, fused via stitching PCR with nesting primers

TDP1392/1393, followed by chitin transformation using *vc1807*::trim (see above) as

primary selection. Δ*ubiA::trim* TDP1399/1401 + TDP1404/1403 + TDP636/637, fused

via stitching PCR with nesting primers TDP1405/1406, followed by chitin transformation.

Δ*oxyR1*: TDP1057/58 + TDP1059/60, Δ*katG:* TDP1100/1101 + TDP1102/1103, Δ*katB*:

TDP1103/1104 + TDP1105/1106, Δ*sodB*: TDP828/829 + TDP830/831, Δ*fur*:

TDP834/835 + TDP836/837 cloned into *XbaI*-digested pCVD442 using isothermal

assembly [94] and Δ*vctPDGC:* TD-JHS394/395 (by amplification from gene block;

vctPDGC deletion), Δ*feoABC:* TD-JHS397/398 (*feoABC* deletion), Δ*fbpABC:* TD-

JHS400/401(*fbpABC* deletion), Δ*vc2211:* TD-JHS403/404 (*vc2211* deletion), Δ*vc2212:*

TD-JHS406/407 (*vc2212* deletion), Δ*tonB2exbB2:* TD-JHS409/410 (*tonB2exbB2*

deletion), and Δ*hutA:* TD-JHS412/413 (*hutA* deletion) cloned into *SphI/XmaI*-digested

pCVD442 followed by knockout via homologous recombination.

Pulldown of VxrB binding DNA fragments and ChIP-Seq library construction. Cells

carrying either 6xHis+VxrB or the native gene were grown in LB medium to mid-

logarithmic phase (at an OD_{600} ~0.5). Following treatment with PenG (100 µg/ml, 3

hours), 30 ml aliquots were spun down and the pellets were saved at -80°C. For cross-

linking, pellets were resuspended in 1ml buffer CJ1 [100 mM Na_2HPO_4, 600 mM NaCl,

and 5 mM imidazole (pH 8.0)] and 30 µl of 34% formaldehyde was added, followed by

rocking at room temperature for 10 min. To quench cross-linking reaction, 133 µl of 1 M

Glycine pH7.5 (final molar concentration 0.133 M) was added, and the cells were

shaken gently at 4°C for 30 min. Cells were collected by centrifugation, and the pellets

were resuspended in 1 ml buffer CJ2 [100 mM Na_2HPO_4, 600 mM NaCl, 0.05% tween[TM],

and 30 mM imidazole (pH 8.0)] after washing twice with same CJ1 buffer. Cells were

then lysed by sonication, followed by addition of Micrococcal nuclease (100 U ml^{-1},

NEB cat# M0247S) and RNase A ($1\square mg\square ml^{-1}$) for fragmentation of genomic DNA

and removal of RNA. 950 µl of supernatant was collected after centrifugation.

For DNA-VxrB complex purification, HisPur^{TM} Ni-NTA magnetic beads (cat#88831-

Thermo Scientific) were equilibrated with six bed volumes of CJ2 buffer [100 mM

Na_2HPO_4, 600 mM NaCl, 0.05% tween^{TM}, and 30 mM imidazole (pH 8.0)]. 500 µg of

crude lysate (protein concentration determined by Bradford assay) was mixed with the

Ni-NTA resin and incubated at 4°C for 4 hours on rotation mixer. The bead slurry was

recovered via magnetic stand for 5 min standing and washed twice with 400 µl buffer

CJ3 [100 mM Na_2HPO_4, 600 mM NaCl, 0.05% tween^{TM}, and 50 mM imidazole (pH 8.0)].

The VxrB-DNA complexes were then eluted from the beads by addition of CJ4 elution

buffer [100 mM Na_2HPO_4, 600 mM NaCl, 0.05% tween^{TM}, and 500 mM imidazole (pH

8.0)]. Enriched target DNAs were purified by using PCR Clean-Up Kits (Thermo Fisher

Scientific) after reversal of crosslinking at 65°C overnight. The desired DNA fragment

size (200~300 bps) was then purified by gel extraction. For illumina HiSeq, each DNA

library was constructed with specific barcode sequences, according to the

manufacturer's instruction of TruSeq ChIP Library Preparation Kit (IP-202-1012,

Illumina). The constructed libraries were sequenced using 100-bp paired-end read

protocol on an Illumina HiSeq 2500 platform (Illumina) at the Biotechnology Resource

Center (BRC) at Cornell University.

ChIP-Seq, binding peak detection, and motif analysis. Sequencing data was

processed by CLC Genomics Workbench (CLC Bio). First, raw sequencing reads were

quality trimmed using a Trim Sequence Tool with a quality limit of 0.05. Any reads with

more than two ambiguous bases were discarded at this step. Then, quality trimmed

reads were mapped onto the *V. cholerae* reference genome (NCBI accession

NC_002505.1 and NC_002506.1) with mismatch cost of 2, insertion/deletion cost of 3,

length fraction of 0.9, and similarity fraction of 0.9. Non-specific matches were mapped

randomly. Mapping files (.bam) were submitted to the Model-based Analysis of ChIP-

Seq 2 (MACS2) software with a mapping file of a mock sample as a control. Peaks were

detected with shift size of 50 and q-value cutoff of 10^{-10}. For motif analysis, peaks with

q-value lower than 10^{-40} were selected and 100 bp DNA sequences centered at peak

summits were submitted to MEME Suite software v4.11.4.

RNA-Seq analysis

Sequencing data (obtained previously) were also processed on a CLC Genomics

Workbench (CLC Bio). Raw sequencing reads were quality trimmed with the same

method with ChIP-Seq reads. Quality trimmed reads were mapped strand specifically

on the reference genome sequence by Transcriptome Analysis Tool with the following

parameters: maximum number of mismatches of 2, length fraction, and similarity

fraction of both 0.9. Number of mapped reads on each gene was normalized by

DESeq2 package in Bioconductor v3.4. Genes with p-value (padj, DESeq2) lower than

0.001 were scored as differentially expressed genes (DEGs).

Purification of VxrB protein. Wild-type VxrB protein was purified from *E. coli* BL21

(DE3/pLysS) cells containing pET15b based recombinant plasmid. The coding

sequence for WT VxrB was cloned into pET15b (Novagen) into the *Nde*I and *Bam*HI

restriction sites. For purification of VxrB from this construct, an overnight culture from a

single colony was used to inoculate 1 liter of LB medium. Cells were grown with

vigorous shaking at 37°C to an optical density at 600 nm (OD_{600}) of 0.5 and were

induced with 1 mM (final concentration) isopropyl-β-D-thiogalactopyranoside (IPTG) for

6 h at 30°C. Harvested cells were re-suspended in binding buffer [20 mM Tris-HCl (pH

7.9), 0.5 M NaCl, 1 mM TCEP (tris(2-carboxyethyl)-phosphine), and 5 mM imidazole]

and cell extracts were prepared by sonication and centrifugation at 16,000 rpm for 45

min. Cell extracts were loaded onto a cobalt-charged NTA column and then washed

with 6 volumes of binding buffer followed by 6 volumes of washing buffer [20 mM Tris-

HCl (pH 8.0), 0.5 M NaCl, and 50 mM imidazole]. VxrB protein was eluted with 10

volumes of elution buffer [20 mM Tris-HCl (8.0) and 0.5 M NaCl] containing linear

imidazole gradients from 100 to 500 mM. Fractions containing VxrB protein were pooled

and dialyzed against buffer J1 [20 mM Tris-HCl (pH 8.0), 250 mM NaCl, 5% (vol/vol)

glycerol, and 4 mM EDTA] to remove imidazole and cobalt and then transferred to two

extra dialysis buffer J2 [20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol, and 0.2

mM DTT] and buffer J3 [20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 30% glycerol, and 1

mM DTT]. The purified VxrB protein was concentrated by centrifugal filter devices

(Millipore, 3,000 MW CO) before injection onto High load TM (16/60) pg Superdex G75

column in FPLC system (Pharmacia). The column was equilibrated with buffer SG [20

mM Tris-HCl (pH 8.0), 150 mM NaCl, and 1 mM DTT]. Eluted fractions were monitored

through UV detector. Concentration of purified VxrB was estimated in triplicate by

Bradford assay (Bio-Rad) using BSA (bovine serum albumin, Sigma-Aldrich) as the

calibration standard at A595. Measurement of UV absorbance at 280 nm, combined

with calculated molar extinction coefficient (ε_{280} = 46470 M^{-1} cm^{-1}, http://expasy.org/cgi-

bin/protparam), gave nearly identical values. The purity of protein was confirmed

through Coomassie blue staining of loaded protein samples in the SDS-PAGE gel. The

protein was stored in final storage buffer JHS [20 mM Tris-HCl (pH 8.0), 150 mM NaCl,

1 mM DTT, 30% Glycerol, 1 mM MgCl$_2$, and 1 mM ATP] at -80°C.

DNase I foot printing assay. The probe DNAs were prepared by using the "crush and

soaking method" (Sambrook *et al.*, 1989) after PCR with the primer pairs shown in

Table S2. The 5′ end of the probes was labeled with 6-FAM (Fluorescein). Binding

reactions were performed as described below for EMSA, except using 250 ng probe

DNA in a 40 µl reaction mixture. DNA probes were incubated with increasing

concentrations of VxrB (0.17, 0.34, 0.68, and 1.38 µM). After 20 min incubation at

room temperature, 40 µl of 5 mM CaCl$_2$ and 10 mM MgCl$_2$ was added, followed by

adding 200 U of RQI RNase-free DNase I (Promega) for 1 min. The cleavage

reaction was stopped by adding 90 µl stop solution (200 mM NaCl, 30 mM EDTA,

1% sodium dodecyl sulfate, 125 µg ml^{-1} of glycogen), followed by DNA extraction

and precipitation. Samples were analyzed by capillary electrophoresis on an ABI

3730x1 DNA analyzer (Life Technologies). The DNA probe only with no added VxrB

was analyzed in parallel. The core protected regions were indicated with red lines and

its corresponding DNA sequences were shown at the bottom of peak profiles.

Preparation of total RNA and S1 nuclease mapping analysis. Wild type and mutant

strains were cultured to mid-logarithmic phase (at an OD$_{600}$ of 0.4 to 0. 5) in LB

medium. Following application of a stress (PenG or H_2O_2), total RNA was extracted

using the "hot phenol method", albeit without lysozyme pretreatment. RNA was

quantified and quality-controlled by absorbance spectroscopy and confirmed by

resolving RNA samples on 1.3% formaldehyde-agarose gels. Gene-specific DNA

oligonucleotide probes for *vxrA, vxrB, pbp1A, zapB, flaA, murJ, oxyR1, sodB, sodA,*

katG, *katB*, *prxA*, *oxyR2*, *ahpC*, *dps*, *fur*, and *ryhB* transcripts were generated by PCR

amplification using *V. cholerae* wild-type genomic DNA as template. The appropriate

primer pairs are listed in **Table S2**. Each specific DNA probe was radiolabeled with (γ-

^{32}P) ATP and T4 polynucleotide kinase and 30,000-40,000 cpm of labeled probe was

used in each reaction. For S1 assays, 100 μg of total RNA was pelleted and lyophilized.

The total RNA pellet was then carefully resuspended in 20 μl hybridization buffer [40

mM PIPES (pH 6.4), 400 mM NaCl, 1 mM EDTA, 80% (v/v) formamide]. Individual

samples were incubated at 95°C for 25 min and cooled to 42°C. Following overnight

incubation, 300 μl of S1 nuclease mix containing 100 units of S1 nuclease in S1

nuclease buffer [280 mM NaCl, 30 mM NaOAc (pH 4.4), 4.5 mM ZnOAc] was added

and incubated at 37°C for 45 min. The reaction was terminated by addition of 75 μl of

S1 nuclease termination solution (2.5 M NH_4OAc, 0.05 M EDTA). The DNA-RNA hybrid

was precipitated by adding 400 μl of isopropanol (100 % v/v) and the pellet was washed

with 70% (v/v) ethanol, vacuum dried, and re-suspended in 13 μl alkaline loading dye.

The protected DNA fragments were then resolved by 6% (w/v) polyacrylamide gels

containing 7 M urea. Gels were dried using a Hoefer Scientific Drygel Slab Gel Dryer

01385 and then imaged on a phosphor imaging screen (Typhoon FLA 7000; GE); bands

were quantified using Multi Gauge V3.0 (Fuji). Quantified average fold intensity

(compared to non-treated sample) from three independent experiments were presented

under each signal with standard deviation values.

Killing Assay. For most time-dependent killing experiments, overnight cultures of wild

type and each mutant strains were grown at 37°C in LB medium in the presence or

absence of additional supplement such as IPTG,0.2% glucose, or 10% sucrose. At each

time-point, samples were collected, serially diluted (sterile PBS) and spot-plated for

CFU/mL determination (plates were incubated at 30 °C for 20 – 24 h). Where indicated,

PenG (100 µg/ml), 250 µM Bip, 250 µM FeCl$_2$, 100 µM, or 1mM FeSO$_4$ were added. For

the growth phase dependent experiment, wild type, *ΔvxrAB*, *Δfur*, *ΔsodB* and

ΔoxyR1ΔkatGΔkatB mutants were cultured in the presence of 1µg/ml catalase for

overnight seed (catalase is required for *ΔoxyR1* growth) in 100 mL LB in flasks. At

designated time points, 5ml cell cultures were transferred to new tubes and cultured

with or without PenG for 3 hours; CFU/mL was determined before and after addition of

the antibiotic. Where applicable, cells were stained with 5 µM of the membrane dye

FM4-64 (Invitrogen, cat# LST3166) for 5 min. Collected images were processed in

ImageJ by subtracting background and adjusting brightness/contrast uniformly across

all fluorescent images.

Streptonigrin sensitivity assay

Cells were diluted 100 fold from overnight cultures and grown to a density of ~2 x 10^8

CFU/mL. At each time point (before addition of 100 µg/ml PenG and 3 h after), 200 µL

of culture were spread on 25 ml LB agar plates using glass beads. At the 3 h time point,

1 mg/ml (final concentration) β-lactamase (Sigma, catalog # P4524) was added prior to

plating. Following plating, a sterile filter disk containing 5 µL of 5 mg/ml streptonigrin

solution (Sigma, catalog # S1014, dissolved in DMSO) was placed in the center of the

plate. Plates were then incubated 24 h at 37 °C. Measurements are averages of six

biological replicates.

ICP-MS analysis for intracellular metal concentration measurement. A single

colony of the indicated *V. cholerae* strains from LB plates was inoculated into 5 mL LB

with streptomycin (200 µg/mL) and grown overnight. The next morning, 0.1% of

overnight cultured seeds were inoculated in 200 ml LB supplemented with antibiotics.

These cells were grown shaking at 37°C for 2~3 hours until they reached an OD_{600} of

~0.5. Samples were collected before and at indicated time points after PenG treatment.

At the indicated time points, cells were pelleted by centrifugation (10 min, 20000 rcf)

and washed three times with phosphate buffered saline (PBS) buffer containing 0.1 M

EDTA followed by two chelex (BioRad, cat # 1422822)-treated PBS buffer washes to

remove all external metal ions. Cells were then resuspended in 400 µl of chelex-treated

PBS buffer with 1% Triton X-100 containing 75 mM NaN3 to induce lysis and incubated

at 37°C for 90 min. 10 µl of each supernatant was used to measure total cell protein

concentration by Bradford assay. 600 µl of 5% HNO3 with 0.1% (v/v) Triton X-100 were

then added to the rest of the supernatant, which was boiled at 95°C for 30 min. After

centrifugation (10 min, 20000 rcf), the supernatant was then diluted 1000fold with 1%

HNO3. Metal levels were measured by ICP-MS (Perkin-Elmer ELAN DRC II using

ammonia as the reaction gas and gallium as an internal standard) and normalized

against total cell protein concentration. Data represent with average values from

biological triplicates. Significant differences between wild type and *Δvxr*AB after PenG

exposure were determined by paired R t-test as indicated: **P=0.001; ***P<0.0003.

Catalase and SOD activity assay. Catalase activity staining was performed. Briefly,

cell extracts (30 µg per well, obtained by 20 cycles of sonication on ice at 20 % power, 3

s pulse, followed by 10 s cooling) were separated on a 5% native polyacrylamide gel

and electrophoresed at 4°C with 80 volts. The gel was washed in distilled water and

then treated with 1 mM H_2O_2 for 20 min. After treatment, the gel was rinsed and

transferred to 1% (w/v) ferric chloride and 1% (w/v) potassium ferricyanide solution. The

reaction was stopped by washing twice in distilled water. Superoxide dismutase activity

staining was performed as described in [96]. The cell extracts (30 µg/well) were

electrophoresed on an 8% native polyacrylamide gel under the same conditions as

described above for catalase activity, after which the gel was incubated in a solution

containing 2.5 mM nitroblue tetrazolium (NBT) for 25 min and then in 50 mM phosphate

buffer (pH 7.8) containing 28 µM riboflavin and 28 mM tetramethyl ethylene diamine for

20 min in the dark. The gel was placed in distilled water and exposed on a light box for

15 min until the dark blue background color appeared. To quantify SOD and catalase

activity, stained gels were scanned and analyzed by Odyssey Imaging Systems (LI-

COR Biosciences) and Image Studio Lite software. Quantified average values from

three independent experiments are presented as a fold activity against wild type non-

treated sample for SODs and Catalases.

Measurement of total cellular H_2O_2 level. Total cellular hydrogen peroxide (H_2O_2)

level was measured using the Amplex^{TM} Red Hydrogen Peroxide/Peroxidase Assay Kit

(Invitrogen™ cat#A22188) and normalized to total cellular proteins. Wild type and

indicated mutant stains were cultured in LB medium (in the absence or presence of

additional 0.2% glucose when it needed) until OD_{600} ~0.5. PenG (100 µg/ml) was then

added and 1ml cell culture was withdrawn and pelleted at each indicated time point.

Pellets were resuspended in 100 µl cold 1xPBS buffer and lysed by sonication, followed

by centrifugation at 4°C (10 min, 20000 rcf). 3 µl of each supernatant was used to

measure total cell protein concentration by Bradford assay. 20µl extracts from cleaned

supernatant were reacted with kit components following manufacturer's instruction. For

H_2O_2 quantification, the reaction product resorufin was read by fluorometer (QUBIT 3,

Invitrogen) at Ex/Em of: ~571/585 nm. Obtained values were normalized to total protein

concentration.

Detection of oxidized proteins. Oxidized proteins containing carbonyl groups were

detected using the OxyBlot Protein Oxidation Detection Kit (Sigma-Aldrich: cat#S7150).

Briefly, wild type and *ΔvxrAB* mutant were grown to exponential growth phase and

harvested at different times after PenG treatment. 0.5 mM H_2O_2 treated cells were used

as a positive protein oxidation control with oxidized standard marker ($_{OX}$M). The protein

sample (10µg/well) was incubated with dinitrophenylhydrazine (DNPH) (derivatization

reaction) for 15 min, followed by neutralization with a solution containing glycerol and β-

mercaptoethanol. Both DNP-derivatized or nonderivatized (for negative control) protein

samples were electrophoresed on a 10% SDS-PAGE and then transferred to a

nitrocellulose membrane. After blocking, the membrane was incubated for 1 h with a

rabbit antibody against DNP (cat# D9656), followed by incubation with secondary

antibody: goat anti-rabbit IgG (Li-COR, IRDye® 800CW cat# 926-32211) for 1 h at room

temperature as recommended by the manufacturer. The signals were scanned on an

Odyssey CLx imaging device (LI-COR Biosciences) and visualized using image studio

lite software.

Microscopy. All images were taken on a Leica MDi8 microscope (Leica Microsystems,

GmbH, Wetzlar, Germany) with a PECON Temp-Controller 2000-1 (Erbach, Germany),

heated stage at 37 °C for growth experiments, or room temperature. Time-lapse

microscopy was performed by imaging frames five minutes apart and data were

analyzed by ImageJ program. The msfGFP expressing cells were imaged at

$485_{Ex}/530_{Em}$ for two second exposure time. Collected images were processed for

quantification in ImageJ by subtracting background and adjusting brightness/contrast

uniformly across all fluorescent images. Representative images were taken at least

three independent experiments.

***In vivo* mouse experiments with *V. cholerae*.**

All mouse-related protocols, sample, and data collection were reviewed and approved

by the Cornell University Institutional Animal Care and Use Committee (IACUC),

protocol number 2016-0088. Female 7-week-old BALB/c mice were acquired from the

Jackson Lab. All mice were housed in groups of up to five animals (upon receipt) or

individually (upon completion of Sm pre-treatment) under standard barrier conditions in

individually ventilated cages with autoclaved food and autoclaved water *ad libitum* and

monitored in accordance with the rules of the Weill Hall Barrier Animal Facility. For

experiments, animals between 8-9 weeks of age were given Streptomycin (Sm: Sigma,

cat# 1623003) in their drinking water (5 mg/ml) for 48 hours and then orally inoculated

with ca 10^{10} Sm-resistant *V. cholerae* El Tor strain N16961. 24 h p.i., mice in AMX

group received oral gavage of Amoxicillin 50 mg/kg (AMX: Sigma, cat# 1031503), mice

in DFOA group received oral gavage of Deferoxamine 100 μM 200ul (DFOA: Sigma,

cat# 1166003), mice in AMX+DFOA received both AMX and DFOA, mice in the Vc

group received water only. Mice were euthanized 24 hrs post treatment. Fecal samples

were collected at 6 time points (e.g. 0, 2, 4, 6, 8, and 10 hrs) and processed to monitor

the level of *V. cholerae* excretion. For the challenge of mice, bacteria were grown

overnight and harvested by centrifugation (x 2717g for 20 min), and washed in

phosphate-buffered saline (PBS: Thomas Scientific, cat# C999G06) at the same speed

and immediately administered to individual mice in PBS (ca 10^{10} *V. cholerae*). To

enumerate *V. cholerae* cells in mouse feces, fresh fecal pellets were collected from an

individual mouse at the indicated time points p.i. and weighed before resuspending into

1 mL PBS, homogenized by vortexing and pipetting. The sample homogenates were

centrifuged at low speed (x 480g for 30 sec) and the resulting supernatant was plated in

serial dilutions onto agar plates containing Sm (200 μg/mL) for CFU counting. The

number of bacterial CFU was normalized per gram of feces (CFU/mg).

1.   Spellberg, B. The future of antibiotics. *Crit Care* **18**, 228 (2014).
2.   Fair, R.J. & Tor, Y. Antibiotics and bacterial resistance in the 21st century.
Perspect Medicin Chem **6**, 25-64 (2014).

3.   Lewis, K. Persister cells: molecular mechanisms related to antibiotic tolerance. *Handb Exp Pharmacol*, 121-133 (2012).

4.   Lewis, K. Persister cells. *Annu Rev Microbiol* **64**, 357-372 (2010).

5.   Lewis, K. Persister cells, dormancy and infectious disease. *Nat Rev Microbiol* **5**, 48-56 (2007).

6.   El-Halfawy, O.M. & Valvano, M.A. Antimicrobial heteroresistance: an emerging field in need of clarity. *Clin Microbiol Rev* **28**, 191-207 (2015).

7.   Wilmaerts, D., Windels, E.M., Verstraeten, N. & Michiels, J. General Mechanisms Leading to Persister Formation and Awakening. *Trends Genet* **35**, 401-411 (2019).

8.   Balaban, N.Q.*, et al.* Definitions and guidelines for research on antibiotic persistence. *Nat Rev Microbiol* (2019).

9.   Kitano, K., Tuomanen, E. & Tomasz, A. Transglycosylase and endopeptidase participate in the degradation of murein during autolysis of Escherichia coli. *J Bacteriol* **167**, 759-765 (1986).

10. Flores-Kim, J., Dobihal, G.S., Fenton, A., Rudner, D.Z. & Bernhardt, T.G. A switch in surface polymer biogenesis triggers growth-phase-dependent and antibiotic-induced bacteriolysis. *Elife* **8**(2019).

11. Dörr, T., Davis, B.M. & Waldor, M.K. Endopeptidase-mediated beta lactam tolerance. *PLoS Pathog* **11**, e1004850 (2015).

12. Cho, H., Uehara, T. & Bernhardt, T.G. Beta-lactam antibiotics induce a lethal malfunctioning of the bacterial cell wall synthesis machinery. *Cell* **159**, 1300-1311 (2014).

13. Monahan, L.G.*, et al.* Rapid conversion of Pseudomonas aeruginosa to a spherical cell morphotype facilitates tolerance to carbapenems and penicillins but increases susceptibility to antimicrobial peptides. *Antimicrob Agents Chemother* **58**, 1956-1962 (2014).

14. Cross, T.*, et al.* Spheroplast-mediated carbapenem tolerance in Gram-negative pathogens *Antimicrob Agents Chemother* (2019).

15. Errington, J., Mickiewicz, K., Kawai, Y. & Wu, L.J. L-form bacteria, chronic diseases and the origins of life. *Philos Trans R Soc Lond B Biol Sci* **371**(2016).

16. Errington, J. Cell wall-deficient, L-form bacteria in the 21st century: a personal perspective. *Biochem Soc Trans* **45**, 287-295 (2017).

17. Dörr, T.*, et al.* A cell wall damage response mediated by a sensor kinase/response regulator pair enables beta-lactam tolerance. *Proc Natl Acad Sci U S A* **113**, 404-409 (2016).

18. Dörr, T., Lewis, K. & Vulic, M. SOS response induces persistence to fluoroquinolones in Escherichia coli. *PLoS Genet* **5**, e1000760 (2009).

19. Raivio, T.L. Everything old is new again: an update on current research on the Cpx envelope stress response. *Biochim Biophys Acta* **1843**, 1529-1541 (2014).

20. Helmann, J.D. Bacillus subtilis extracytoplasmic function (ECF) sigma factors and defense of the cell envelope. *Curr Opin Microbiol* **30**, 122-132 (2016).

21. Radeck, J., Fritz, G. & Mascher, T. The cell envelope stress response of Bacillus subtilis: from static signaling devices to dynamic regulatory network. *Curr Genet* **63**, 79-90 (2017).

1032   22.   Wall, E., Majdalani, N. & Gottesman, S. The Complex Rcs Regulatory Cascade.
1033   *Annu Rev Microbiol* **72**, 111-139 (2018).
1034   23.   Guest, R.L. & Raivio, T.L. Role of the Gram-Negative Envelope Stress Response
1035   in the Presence of Antimicrobial Agents. *Trends Microbiol* **24**, 377-390 (2016).
1036   24.   Bhagirath, A.Y.*, et al.* Two Component Regulatory Systems and Antibiotic
1037   Resistance in Gram-Negative Pathogens. *Int J Mol Sci* **20**(2019).
1038   25.   Bury-Mone, S.*, et al.* Global analysis of extracytoplasmic stress signaling in
1039   Escherichia coli. *PLoS Genet* **5**, e1000651 (2009).
1040   26.   Cheng, A.T., Ottemann, K.M. & Yildiz, F.H. Vibrio cholerae Response Regulator
1041   VxrB Controls Colonization and Regulates the Type VI Secretion System. *PLoS*
1042   *Pathog* **11**, e1004933 (2015).
1043   27.   Teschler, J.K., Cheng, A.T. & Yildiz, F.H. The Two-Component Signal
1044   Transduction System VxrAB Positively Regulates Vibrio cholerae Biofilm
1045   Formation. *J Bacteriol* **199**(2017).
1046   28.   Dwyer, D.J., Kohanski, M.A., Hayete, B. & Collins, J.J. Gyrase inhibitors induce
1047   an oxidative damage cellular death pathway in Escherichia coli. *Mol Syst Biol* **3**,
1048   91 (2007).
1049   29.   Kohanski, M.A., Dwyer, D.J., Hayete, B., Lawrence, C.A. & Collins, J.J. A
1050   common mechanism of cellular death induced by bactericidal antibiotics. *Cell*
1051   **130**, 797-810 (2007).
1052   30.   Kohanski, M.A., Dwyer, D.J., Wierzbowski, J., Cottarel, G. & Collins, J.J.
1053   Mistranslation of membrane proteins and two-component system activation
1054   trigger antibiotic-mediated cell death. *Cell* **135**, 679-690 (2008).
1055   31.   Dwyer, D.J., Kohanski, M.A. & Collins, J.J. Role of reactive oxygen species in
1056   antibiotic action and resistance. *Curr Opin Microbiol* **12**, 482-489 (2009).
1057   32.   Imlay, J.A. Diagnosing oxidative stress in bacteria: not as easy as you might
1058   think. *Curr Opin Microbiol* **24**, 124-131 (2015).
1059   33.   Liu, Y. & Imlay, J.A. Cell death from antibiotics without the involvement of
1060   reactive oxygen species. *Science* **339**, 1210-1213 (2013).
1061   34.   Keren, I., Wu, Y., Inocencio, J., Mulcahy, L.R. & Lewis, K. Killing by bactericidal
1062   antibiotics does not depend on reactive oxygen species. *Science* **339**, 1213-1216
1063   (2013).
1064   35.   Hong, Y., Zeng, J., Wang, X., Drlica, K. & Zhao, X. Post-stress bacterial cell
1065   death mediated by reactive oxygen species. *Proc Natl Acad Sci U S A* **116**,
1066   10064-10071 (2019).
1067   36.   Dwyer, D.J., Collins, J.J. & Walker, G.C. Unraveling the physiological
1068   complexities of antibiotic lethality. *Annu Rev Pharmacol Toxicol* **55**, 313-332
1069   (2015).
1070   37.   Dwyer, D.J.*, et al.* Antibiotics induce redox-related physiological alterations as
1071   part of their lethality. *Proc Natl Acad Sci U S A* **111**, E2100-2109 (2014).
1072   38.   Jacoby, G.A. AmpC beta-lactamases. *Clin Microbiol Rev* **22**, 161-182, Table of
1073   Contents (2009).
1074   39.   Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes.
1075   *Nucleic Acids Res* **28**, 27-30 (2000).
1076   40.   Bailey, T.L.*, et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic*
1077   *Acids Res* **37**, W202-208 (2009).

41. Dörr, T*., et al.* Differential requirement for PBP1a and PBP1b in in vivo and in
vitro fitness of Vibrio cholerae. *Infect Immun* **82**, 2115-2124 (2014).
42. Dörr, T*., et al.* A novel peptidoglycan binding protein crucial for PBP1A-mediated
cell wall biogenesis in Vibrio cholerae. *PLoS Genet* **10**, e1004433 (2014).
43. Ruiz, N. Lipid Flippases for Bacterial Peptidoglycan Biosynthesis. *Lipid Insights*
8, 21-31 (2015).
44. Teo, A.C. & Roper, D.I. Core Steps of Membrane-Bound Peptidoglycan
Biosynthesis: Recent Advances, Insight and Opportunities. *Antibiotics (Basel)* **4**,
495-520 (2015).
45. Davies, B.W., Bogard, R.W. & Mekalanos, J.J. Mapping the regulon of Vibrio
cholerae ferric uptake regulator expands its known network of gene regulation.
Proc Natl Acad Sci U S A **108**, 12467-12472 (2011).
46. White, J.R. & Yeowell, H.N. Iron enhances the bactericidal action of streptonigrin.
Biochem Biophys Res Commun **106**, 407-411 (1982).
47. Yeowell, H.N. & White, J.R. Iron requirement in the bactericidal mechanism of
streptonigrin. *Antimicrob Agents Chemother* **22**, 961-968 (1982).
48. Yeom, J., Imlay, J.A. & Park, W. Iron homeostasis affects antibiotic-mediated cell
death in Pseudomonas species. *J Biol Chem* **285**, 22689-22695 (2010).
49. Leger, L*., et al.* beta-Lactam Exposure Triggers Reactive Oxygen Species
Formation in Enterococcus faecalis via the Respiratory Chain Component DMK.
Cell Rep **29**, 2184-2191 e2183 (2019).
50. Seo, S.W*., et al.* Deciphering Fur transcriptional regulatory network highlights its
complex role beyond iron metabolism in Escherichia coli. *Nat Commun* **5**, 4910
(2014).
51. Masse, E. & Gottesman, S. A small RNA regulates the expression of genes
involved in iron metabolism in Escherichia coli. *Proc Natl Acad Sci U S A* **99**,
4620-4625 (2002).
52. Kehres, D.G., Janakiraman, A., Slauch, J.M. & Maguire, M.E. Regulation of
Salmonella enterica serovar Typhimurium mntH transcription by H(2)O(2),
Fe(2+), and Mn(2+). *J Bacteriol* **184**, 3151-3158 (2002).
53. Martin, J.E., Waters, L.S., Storz, G. & Imlay, J.A. The Escherichia coli small
protein MntS and exporter MntP optimize the intracellular concentration of
manganese. *PLoS Genet* **11**, e1004977 (2015).
54. Ikeda, J.S., Janakiraman, A., Kehres, D.G., Maguire, M.E. & Slauch, J.M.
Transcriptional regulation of sitABCD of Salmonella enterica serovar
Typhimurium by MntR and Fur. *J Bacteriol* **187**, 912-922 (2005).
55. Fisher, C.R., Wyckoff, E.E., Peng, E.D. & Payne, S.M. Identification and
Characterization of a Putative Manganese Export Protein in Vibrio cholerae. *J*
Bacteriol **198**, 2810-2817 (2016).
56. Wyckoff, E.E. & Payne, S.M. The Vibrio cholerae VctPDGC system transports
catechol siderophores and a siderophore-free iron ligand. *Mol Microbiol* **81**,
1446-1458 (2011).
57. Zeng, L*., et al.* Deferoxamine therapy for intracerebral hemorrhage: A systematic
review. *PLoS One* **13**, e0193615 (2018).

58.   Fisher, S.A.*, et al.* Desferrioxamine mesylate for managing transfusional iron
1123   overload in people with transfusion-dependent thalassaemia. *Cochrane
1124   Database Syst Rev*, CD004450 (2013).
1125   59.   Goodwin, J.F. & Whitten, C.F. Chelation of Ferrous Sulphate Solutions by
1126   Desferrioxamine B. *Nature* **205**, 281-283 (1965).
1127   60.   Miyajima, H.*, et al.* Use of desferrioxamine in the treatment of
1128   aceruloplasminemia. *Ann Neurol* **41**, 404-407 (1997).
1129   61.   Richardson, D., Ponka, P. & Baker, E. The effect of the iron(III) chelator,
1130   desferrioxamine, on iron and transferrin uptake by the human malignant
1131   melanoma cell. *Cancer Res* **54**, 685-689 (1994).
1132   62.   Imlay, J.A. The molecular mechanisms and physiological consequences of
1133   oxidative stress: lessons from a model bacterium. *Nat Rev Microbiol* **11**, 443-454
1134   (2013).
1135   63.   Imlay, J.A. The mismetallation of enzymes during oxidative stress. *J Biol Chem*
1136   **289**, 28121-28128 (2014).
1137   64.   Xia, X.*, et al.* OxyR-activated expression of Dps is important for Vibrio cholerae
1138   oxidative stress resistance and pathogenesis. *PLoS One* **12**, e0171201 (2017).
1139   65.   Wang, H.*, et al.* Catalases promote resistance of oxidative stress in Vibrio
1140   cholerae. *PLoS One* **7**, e53383 (2012).
1141   66.   Wang, H.*, et al.* OxyR2 Modulates OxyR1 Activity and Vibrio cholerae Oxidative
1142   Stress Response. *Infect Immun* **85**(2017).
1143   67.   Storz, G., Tartaglia, L.A. & Ames, B.N. The OxyR regulon. *Antonie Van
1144   Leeuwenhoek* **58**, 157-161 (1990).
1145   68.   Johnson, L.A. & Hug, L.A. Distribution of reactive oxygen species defense
1146   mechanisms across domain bacteria. *Free Radic Biol Med* (2019).
1147   69.   Muras, V., Dogaru-Kinn, P., Minato, Y., Hase, C.C. & Steuber, J. The Na+-
1148   Translocating NADH:Quinone Oxidoreductase Enhances Oxidative Stress in the
1149   Cytoplasm of Vibrio cholerae. *J Bacteriol* **198**, 2307-2317 (2016).
1150   70.   Steuber, J.*, et al.* The structure of Na(+)-translocating of NADH:ubiquinone
1151   oxidoreductase of Vibrio cholerae: implications on coupling between electron
1152   transfer and Na(+) transport. *Biol Chem* **396**, 1015-1030 (2015).
1153   71.   Tuomanen, E., Cozens, R., Tosch, W., Zak, O. & Tomasz, A. The rate of killing of
1154   Escherichia coli by beta-lactam antibiotics is strictly proportional to the rate of
1155   bacterial growth. *J Gen Microbiol* **132**, 1297-1304 (1986).
1156   72.   Weaver, A.I.*, et al.* Genetic Determinants of Penicillin Tolerance in Vibrio
1157   cholerae. *Antimicrob Agents Chemother* **62**(2018).
1158   73.   Kawai, Y.*, et al.* Cell growth of wall-free L-form bacteria is limited by oxidative
1159   damage. *Curr Biol* **25**, 1613-1618 (2015).
1160   74.   Kawai, Y.*, et al.* Crucial role for central carbon metabolism in the bacterial L-form
1161   switch and killing by beta-lactam antibiotics. *Nat Microbiol* (2019).
1162   75.   Schnell, S. & Steinman, H.M. Function and stationary-phase induction of
1163   periplasmic copper-zinc superoxide dismutase and catalase/peroxidase in
1164   Caulobacter crescentus. *J Bacteriol* **177**, 5924-5929 (1995).
1165   76.   Longo, V.D., Gralla, E.B. & Valentine, J.S. Superoxide dismutase activity is
1166   essential for stationary phase survival in Saccharomyces cerevisiae.

Mitochondrial production of toxic oxygen species in vivo. *J Biol Chem* **271**, 12275-12280 (1996).

77. Gort, A.S., Ferber, D.M. & Imlay, J.A. The regulation and role of the periplasmic copper, zinc superoxide dismutase of Escherichia coli. *Mol Microbiol* **32**, 179-191 (1999).

78. Martins, D.*, et al.* Superoxide dismutase activity confers (p)ppGpp-mediated antibiotic tolerance to stationary-phase Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **115**, 9797-9802 (2018).

79. Haugan, M.S., Charbon, G., Frimodt-Moller, N. & Lobner-Olesen, A. Chromosome replication as a measure of bacterial growth rate during Escherichia coli infection in the mouse peritonitis model. *Sci Rep* **8**, 14961 (2018).

80. Keren, I., Kaldalu, N., Spoering, A., Wang, Y. & Lewis, K. Persister cells and tolerance to antimicrobials. *FEMS Microbiol Lett* **230**, 13-18 (2004).

81. Sharma, B., Brown, A.V., Matluck, N.E., Hu, L.T. & Lewis, K. Borrelia burgdorferi, the Causative Agent of Lyme Disease, Forms Drug-Tolerant Persister Cells. *Antimicrob Agents Chemother* **59**, 4616-4624 (2015).

82. Brynildsen, M.P., Winkler, J.A., Spina, C.S., MacDonald, I.C. & Collins, J.J. Potentiating antibacterial activity by predictably enhancing endogenous microbial ROS production. *Nat Biotechnol* **31**, 160-165 (2013).

83. Gutierrez, A.*, et al.* beta-Lactam antibiotics promote bacterial mutagenesis via an RpoS-mediated reduction in replication fidelity. *Nat Commun* **4**, 1610 (2013).

84. Sheldon, J.R. & Skaar, E.P. Metals as phagocyte antimicrobial effectors. *Curr Opin Immunol* **60**, 1-9 (2019).

85. Yang, J.H.*, et al.* A White-Box Machine Learning Approach for Revealing Antibiotic Mechanisms of Action. *Cell* **177**, 1649-1661 e1649 (2019).

86. Radlinski, L. & Conlon, B.P. Antibiotic efficacy in the complex infection environment. *Curr Opin Microbiol* **42**, 19-24 (2018).

87. Yang, J.H., Bening, S.C. & Collins, J.J. Antibiotic efficacy-context matters. *Curr Opin Microbiol* **39**, 73-80 (2017).

88. Stokes, J.M., Lopatkin, A.J., Lobritz, M.A. & Collins, J.J. Bacterial Metabolism and Antibiotic Efficacy. *Cell Metab* (2019).

89. Papenfort, K., Forstner, K.U., Cong, J.P., Sharma, C.M. & Bassler, B.L. Differential RNA-seq of Vibrio cholerae identifies the VqmR small RNA as a regulator of biofilm formation. *Proc Natl Acad Sci U S A* **112**, E766-775 (2015).

90. Krin, E.*, et al.* Expansion of the SOS regulon of Vibrio cholerae through extensive transcriptome analysis and experimental validation. *BMC Genomics* **19**, 373 (2018).

91. Donnenberg, M.S. & Kaper, J.B. Construction of an eae deletion mutant of enteropathogenic Escherichia coli by using a positive-selection suicide vector. *Infect Immun* **59**, 4310-4317 (1991).

92. Dalia, A.B. Natural Cotransformation and Multiplex Genome Editing by Natural Transformation (MuGENT) of Vibrio cholerae. *Methods Mol Biol* **1839**, 53-64 (2018).

93. Hubbard, T.P.*, et al.* A live vaccine rapidly protects against cholera in an infant rabbit model. *Sci Transl Med* **10**(2018).

1213   94.   Gibson, D.G.*, et al.* Enzymatic assembly of DNA molecules up to several
1214   hundred kilobases. *Nat Methods* **6**, 343-345 (2009).
1215   95.   Dörr, T., Cava, F., Lam, H., Davis, B.M. & Waldor, M.K. Substrate specificity of
1216   an elongation-specific peptidoglycan endopeptidase and its implications for cell
1217   wall architecture and growth of Vibrio cholerae. *Mol Microbiol* **89**, 949-962
1218   (2013).
1219   96.   Beauchamp, C. & Fridovich, I. Superoxide dismutase: improved assays and an
1220   assay applicable to acrylamide gels. *Anal Biochem* **44**, 276-287 (1971).
1221

A

Meropenem Killing Dynamics

Log$_{10}$ (CFU/mL)

- *V. cholerae* N16961
- *P. aeruginosa* PA14
- *E. cloacae* WCM0001
- *K. aerogenes* WCM0001
- *K. pneumoniae* WCM0001
- *E. coli* WCM0001
- *E. coli* MG1655

Clinical isolates from patients

Meropenem (10 μg/mL)/Time (hour)

B

Meropenem

	No antibiotics	1 hour	3 hour
	Phase FM4-64 Merge	Phase FM4-64 Merge	Phase FM4-64 Merge

V. cholerae N16961

P. aeruginosa PA14

E. cloacae WCM0001

K. aerogenes WCM 0001

K. pneumoniae WCM0001

E. coli WCM0001

E. coli MG1655

β−lactam tolerance

◄lysis

Figure 2

A

Chromosome I

VxrB, 100 ug/ml PenG 3H

(LPS) VC0215   (PBP1B) VC0602   (PBP5,6/RlpA) VC0947/8   (Flagella assembly) VC2188-93   (PBP1A) VC2635

VxrB+His, 100 ug/ml PenG 3H

Chromosome II

VxrB, 100 ug/ml PenG 3H

(ShyA) VCA0079   (FhuD) VCA0227   (VxrAB) VCA0565/6   (DcuC) VCA0665

VxrB+His, 100 ug/ml PenG 3H

B

Log$_2$(Fold-change; over exp./control) vs Log$_2$ (Mean Expression)

- DEGs (n= 818)
- non-DEG (n=2870)

VC0157, vxrB, pbp1b, vxrA, pbp1a, hutA

C

RNA-Seq (818 genes)   ChIP-Seq (424 peaks)

p-value < 0.001 q-value<10^{-10}

678 genes   140 genes (127 peaks)   297 peaks

Up (102 genes)
Down (38 genes)

D

0.5 kb   VC2634   Penicillin-binding protein 1A VC2635   VC2636

VxrB, 100PG 3H

RNA-Seq normalized mapped reads (x10^3)
WT control
VxrB O/E

Transcripts (S1 nuclease mapping)
Wild Type   ΔvxrAB
0 20 180   0 20 180 min (100 µg/ml)
pbp1A
rRNAs

Directly activated gene

0.25 kb   VC2685   ZapB VC2686   VC2687

VxrB, 100PG 3H

RNA-Seq normalized mapped reads (x10^2)
WT control
VxrB O/E

Transcripts (S1 nuclease mapping)
Wild Type   ΔvxrAB
0 20 180   0 20 180 min (100 µg/ml)
zapB
rRNAs

Directly repressed gene

0.3kb   VC2187   Flagellin VC2188   VC2189

VxrB, 100PG 3H

RNA-Seq normalized mapped reads (x10^2)
WT control
VxrB O/E

Transcripts (S1 nuclease mapping)
Wild Type   ΔvxrAB
0 20 60 180   0 20 60 180 min (100 µg/ml PenG)
flaA
rRNAs

Poor binding of VxrB or indirectly regulated gene

Figure 3

A

VxrB RNA-Seq (818 genes)
P-value<0.001

Fur regulon (63 genes)
RegPrecise (Regulon DB)

771   **47**   16

VxrB binding (8 peaks)
q-values <10^{-10}

20/47

B

Vibriobactin synthesis

Fur VxrB

VC2209   VC2210   VC2211   VC2212

Fe(III) uptake transport

Fur VxrB

VCA0227   VCA0228   VCA0229   VCA0230

Heme transport / Phosphate transport

Fur VxrB   Fur VxrB

VCA0066   VCA0067   VCA0068   VCA0070-VCA0073

TonB2/ExbB2/ExbD2(Iron transport)

Fur VxrB

VC1543-VC1548

C

** / ***

WT   ΔvxrAB

***

***   ***

Total Cellular Iron (ng mg^{-1} protein)

PenG Exposure (min)

0   20   40   60   90   180

D

WT

- PenG   + PenG   + PenG

SNG   SNG   DMSO

26.5 +/- 1.3 mm   32.3 +/- 1.3 mm   0 +/- 0 mm
(n = 6)   (n = 6)   (n = 3)

A

B

C

A

B

C