Short paired-end reads trump long single-end reads for expression analysis

Adam H. Freedman, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

John M. Gaspar, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

Timothy B. Sackton*, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

* Corresponding author: tsackton@g.harvard.edu

ABSTRACT

Background: Typical experimental design advice for expression analyses using RNA-seq

generally assumes that single-end reads provide robust gene-level expression estimates in a cost-

effective manner, and that the additional benefits obtained from paired-end sequencing are not

worth the additional cost. However, in many cases (e.g., with Illumina NextSeq and NovaSeq

instruments), shorter paired-end reads and longer single-end reads can be generated for the same

cost, and it is not obvious which strategy should be preferred. Using publicly available data, we

test whether short-paired end reads can achieve more robust expression estimates and differential

expression results than single-end reads of approximately the same total number of sequenced

bases.

Results: At both the transcript and gene levels, 2x40 paired-end reads unequivocally provide

expression estimates that are more highly correlated with 2x125 than 1x75 reads; in nearly all

cases, those correlations are also greater than for 1x125, despite the greater total number of

sequenced bases for the latter. Across an array of metrics, differential expression tests based

upon 2x40 consistently outperform those using 1x75.

Conclusion: Researchers seeking a cost-effective approach for gene-level expression analysis

should prefer short paired-end reads over a longer single-end strategy. Short paired-end reads

will also give reasonably robust expression estimates and differential expression results at the

isoform level.

BACKGROUND

For over a decade, RNA-seq has empowered gene expression analysis. This has led to

fundamental advances in our understanding of diverse phenomena, including the evolution of

gene regulation across species (1), alternative splicing (2,3), relative importance of cis vs -trans

regulation (4,5), genetic underpinnings of heritable disease (6,7), and the genetic architecture of

phenotypes in natural populations (8,9). While the cost of sequencing has steadily decreased,

making expression analyses feasible even on modest research budgets, maximizing cost

efficiency remains high priority. Cost effectiveness is particularly important as studies scale up

to hundreds or even thousands of samples. Maximizing performance given a particular study

design has already received much attention, with investigations of optimal strategies for quality

trimming (10), the effects of quality trimming on expression estimates and differential

expression testing (11), optimal sequencing depth (12,13), importance of biological replicates

(12,14,15), and the relative performance of expression and differential expression tools and

pipelines (16–20). Less attention has focused on how to improve performance given a fixed

number of biological replicates and total sequenced bases.

While the typical length of sequenced reads has increased as costs have decreased, when

an annotated genome is available it remains standard procedure to sequence single end reads to

estimate gene expression, typically around 75 base pairs. Yet, this strategy does not leverage

information provided by the full length of the library fragments, particularly the greater

specificity in read mapping (or pseudo-alignment) that would be provided by sequencing from

both ends of a fragment. Given a fixed cost per base for the Illumina NextSeq, HiSeq X Ten, and

NovaSeq instruments, paired-end 75bp reads would be twice as expensive as single-end 75bp

sequencing, assuming an equivalent number of sequenced fragments. However, a sizeable

fraction of the long-range information provided by paired-end sequencing should still be retained

by sequencing shorter paired-end reads. In this study, we evaluate whether that intuition is

correct, by assessing whether expression estimates and differential expression results obtained

with 2x40 paired end sequencing are more consistent with 2x125 paired end sequencing than

either 1x75 or 1x125 single-end sequencing. Our particular focus is on 1x75, as the cost of 1x75

and 2x40 are the same per read when sequencing using the Illumina NextSeq (75 cycle kit), a

common instrument for RNA-seq studies. Given the fixed cost per base for the HiSeq X Ten and

NovaSeq instruments, there are analogous choices to make between single-end vs. paired-end

sequencing. For example, for the 100bp NovaSeq kit, a sequencing strategy trade-off exists

between sequencing 2x50 vs. 1x100 reads. We conduct these analyses with publicly available

Illumina sequence data across 12 different SRA (Additional file 1: Table S1). accessions

encompassing diverse experimental designs and multiple model organisms with high quality

annotated genomes.

RESULTS AND DISCUSSION

Our approach to evaluating the performance of short paired-end reads vs. longer single-end ones

is to compare expression estimates and differential expression results derived from these

strategies to a truth set. Because we cannot know the "truth", we assume that its closest

approximation are results obtained from long, paired-end reads. Thus, we define our gold

standard as the results obtained from 2x125 paired-end reads. We then trim these data down to

two data sets with read lengths of 40 and 75 base pairs, and calculate Spearman's rank

correlations between transcripts-per-million (TPM) estimates based upon 2x40, 1x75 and 1x125

with the 2x125 gold standard. We present these correlations in two different ways. First, we

examine the distributions of those correlation coefficients across alternative sequencing

strategies. Second, we plot the correlations obtained using 2x40 over those obtained with 1x75

and 1x125.

Kallisto transcript-level TPM estimates made with 2x40 are always more highly

correlated with those made with the 2x125 gold standard than those made with data trimmed to

1x75 reads (Fig. 1a,c), and perform better than estimates made with data trimmed to 1x125 for

all but a few samples from one SRA accession (Fig. 1a,c,e). A similar performance advantage for

the 2x40 strategy compared to the 1x75 strategy is observed at the gene level (Fig. 1 b,d,f),

although there is one accession where 1x125 performs better than 2x40.

RSEM-based TPM estimates show an overall pattern generally consistent with that

observed with kallisto (Fig. 2). A handful of samples for which 2x40 performed less well than

both 1x75 and 1x125 had low bowtie2 alignment rates to the reference transcripts (for the 2x125

libraries, overall alignment rates of 37.1-51.0%), suggesting that library quality issues were

responsible, but raising a broader issue concerning whether variation in alignment rates between

strategies could explain the relative performance of 2x40 to 1x75 and 1x125.

Kallisto does not conduct formal sequence alignment, so for each sample-strategy

combination we summed the kallisto transcript-level counts as a proxy for the number of

informative reads used in estimating expression. The summed kallisto counts for 2x40 have a

weak correlation with performance (with respect to TPM) relative to 1x75 the transcript level,

and a moderate correlation at the gene level (Fig 3a,b); the relative differences in counts between

2x40 and 1x75 appear uncorrelated with the relative performance of 2x40 and 1x75 (Fig. 3c,d).

A similar pattern is observed with respect to 1x125 (Additional file 2: Fig. S1). For RSEM-

derived expression estimates, we found no impact of the percent of reads uniquely aligning on

the relative performance of 2x40, but at the gene level found that relative differences in overall

alignment rate had some effect on the robustness of TPM estimates of 2x40 relative to both 1x75

and 1x125 (Additional file 3: Fig. S2,S3). For both kallisto and RSEM, while there are no

obvious taxonomic effects there are clear experiment-specific effects, manifested as differences

in relative 2x40 performance among accessions derived from the same species. Similar to the

above results based upon expression correlations, looking across all accessions, for most samples

root-mean-square error of expression estimates based upon 1x75 and 1x125 was greater than for

2x40 (Fig. 4).

We next evaluated the extent to which differences in performance with respect to

estimating expression levels translated into downstream effects on tests of differential

expression. Within each SRA accession and for each pair of conditions and for each sequencing

strategy, we conducted Wald tests with, sleuth, limma-voom, and DESeq2. Similar to our

evaluation of expression estimate robustness, we defined "true" differential expression signals as

those recovered with 2x125, given a false discovery rate (FDR) of 0.01. We then calculated

seven performance metrics on a per-accession basis for each differential expression method -

sequencing strategy combination.

Sequencing strategy clearly impacts downstream differential expression analyses.

Looking across all three differential expression methods, at both the transcript and gene level, for

the majority of accessions false negative rates are lower for 2x40 compared to 1x75 (Figure 5;

Additional file 4: Fig. S4, S5; Additional file 5: Table S2). The empirical false discovery rate—

that is, for a strategy being evaluated, the proportion of putative differentially expressed features

with a Benjamini-Hochberg adjusted p-value below our chosen FDR threshold of 0.01for which

the adjusted p-value is >0.01 with 2x125—was also lower for 2x40 compared to both 1x75 and

x125, although it was consistently higher than 0.01, the threshold used to classify Wald tests as

significant, including the 2x125 analyses with which we defined true positives and negatives

(Figure 5; Additional file 4: Fig. S4, S5; Additional file 5: Table S2). FDR control was poorer at

the transcript than the gene level, and, as previously reported, DESeq2 did not control FDR as

well as limma-voom and sleuth (18). At both the transcript and gene level, AUC is greater for

2x40 than 1x75 for all but a handful of method-accession combinations (Figure 5, Additional file

4: Fig. S4, S5; Additional file 5: Table S2). Additional metrics indicate a similar overall

performance advantage of 2x40 compared to 1x75 (Additional file 4: Table S2). For all metrics,

the short paired-end strategy outperforms 1x125 for the majority of method-accession

combinations, although the magnitude of performance advantage of 2x40 compared to 1x125 is

smaller than when compared with 1x75 smaller (Figure 5, Additional file 4: Fig. S4,S5;

Additional file 5: Table S2).

CONCLUSIONS

When designing a gene expression study using bulk RNA-seq, given a fixed budget, more robust

expression estimates and differential expression test results can be obtained from short paired-

end reads compared to longer single-end reads. In other words, the extra information provided by

reads pairs during the mapping or pseudoalignment phase outweighs any potential penalty of

short reads. In almost all cases, short paired-end reads trump single-end reads that sequence

approximately the same number of bases. Short paired-end reads provide an additional benefit in

that they can identify a large fraction of differentially expressed isoforms, for the cost of gene-

level analysis with longer single-end reads. From a practical perspective, neither NextSeq nor

NovaSeq sequencing kits distinguish between single and paired-end libraries. In the experience

of support staff at the Bauer sequencing core facility at Harvard, one can get 92 bp from the

NextSeq 75 bp kit, such that 2x40 is a viable sequencing design. Alternatively, the smallest

NovaSeq kit is for 100 cycles, which can be used to sequence paired 50bp reads, allowing one to

leverage that instrument's much smaller cost per sequenced base. Overall, our findings suggest

that for a given number of sequencing cycles, the most efficient design for RNA-seq expression

analysis is sequencing shorter paired reads, rather than longer single-end reads.

METHODS

Datasets

From NCBI's Short Read Archive, we downloaded paired-end RNA-seq reads from 12 different

studies (Table 1), representing model organisms with well-annotated reference genomes: mouse

(*Mus musculus*), human (*Homo sapiens*), yeast (*Saccharomyces cerevisiae*), fly (*Drosophila*

melanogaster), and worm (*Caenorhabditis elegans*). In selecting data sets, we required minimum

pre-processing read lengths to be ≥ 125 bp, and experimental designs to contain ≥ 2 experimental

conditions.

Short-read processing

We removed adapters and trimmed reads with NGmerge (21). For each sample, we trimmed

reads to 125, 75, and 40bp using awk. For downstream analyses, we used 2x125, 2x40,1x125,

and 1x75 trimmed data sets. Single-end data sets were comprised of the first read from the

original paired reads. Reads were trimmed from the distal ends, such that trimming mimicked

how sequencing to different lengths would be carried out on an Illumina sequencer.

Expression analyses

For each sample-sequencing strategy combination, we estimated transcript-level expression with

kallisto (22). We designated abundance estimates derived from 2x125 as our gold standard

against which to benchmark all other strategies. We used Spearman's rank correlations to

quantify the degree of concordance between alternative sequencing strategies and the gold

standard in terms of transcript per million (TPM). We performed similar analyses at the gene

level by aggregating transcript-level TPMs using the R package Tximport (19). For yeast, which

has no alternative splicing, analyses were only conducted at the transcript level. As an additional

estimate of the correspondence between alternative sequencing strategies with 2x125 we also

calculated root-mean-square error (RMSE).

Because a large proportion of RNA-seq applications involve tests of differential

expression (DE), we compared differential expression test results between the gold standard

2x125 and alternative strategies. Specifically, for each experiment we conducted all possible

pairwise Wald tests of transcript-level DE with sleuth (18) version 0.30.0, and used p-value

aggregation (https://pachterlab.github.io/sleuth_walkthroughs/pval_agg/analysis.html) for gene-

level analyses. To compare performance of alternative strategies relative to the gold standard, we

calculated false negative and false positive rates, defining significant tests as those with

Benjamini-Hochberg adjusted p-values ≤ 0.01.

In order to evaluate whether our findings were pipeline-specific, perhaps related to

particular features of tools such as kallisto that rely on pseudo-alignment, we also performed

expression analyses with an alternative pipeline, involving estimation of expression levels with

RSEM (23) version 1.2.29 based upon alignments with bowtie2 version 2.3.2, (24). When SRA

metadata enabled us to determine that libraries were stranded, we included the relevant RSEM

command line arguments for restricting bowtie2 alignments to the correct read pair

configuration. Using these expression estimates, we conducted differential expression tests with

two different methods: limma-voom (25) version 3.38.3 and DESeq2 (26) version 1.22.2. For

limma-voom, we only conducted tests where, for each gene or transcript,there were ≥ 2 samples

with counts per million (CPM) ≥ 1. For limma-voom, we employed TMM normalization, and the

voomWithQualityWeights option. Sleuth and limma-voom have previously been demonstrated to

be top performers that correctly control FDR (22), and while DESeq2 has been shown to not

control FDR as well as these two methods (22), we included it because it is a commonly used

differential expression method. For all differential expression analyses, we calculated

performance metrics for Wald tests where, at the specified false discovery rate (FDR) threshold

of 0.01, there were ≥ 50 significantly differentially expressed features in all sequencing

strategies. Metrics were calculated relative to 2x125, i.e. significant and non-significant results

obtained with 2x125 were treated as true positives and negatives, respectively. Those metrics

were the empirical false discovery rate, false positive rate, false negative rate, sensitivity,

specificity, precision, and area under the curve (AUC) based upon receiver-operator statistics.

These metrics were calculated with python scripts available at our associated github repository

(see below). As the multiple-test adjusted p-values are the standard metric with which to

determine whether particular transcripts or genes are differentially expressed, for AUC

calculations we used these values as the response variable relative to the "true" classification of

differences derived from tests based upon the 2x125 data. All reference genomes and gtf-

formatted annotations against which to estimate expression were downloaded from Ensembl, and

transcripts for kallisto-based analyses were extracted using the *extract-reference-transcripts*

utility in RSEM.

DECLARATIONS

Availability of data and materials

The datasets analyzed during the current study and which support the conclusions of this article

are publicly available Illumina short read data available on NCBI's Short Read Archive

(https://www.ncbi.nlm.nih.gov/sra). Specific accessions are listed in Additional File 1: Table S1.

Command lines for executing NGmerge, kallisto, RSEM, sleuth, and limma-voom analyses are

available on GitHub at https://github.com/harvardinformatics/rnaseq_readlength_assessment.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by the Harvard University Faculty of Arts and Sciences.

Authors' contributions

TBS and AHF designed the experiments, AHF and JMG analyzed the data, and AHF wrote the paper with TBS.

Acknowledgments

The computations in this paper were run on the Odyssey cluster supported by the FAS Division of Science, Research Computing Group at Harvard University. We thank Claire Reardon, director of the Bauer Core at Harvard University for helpful discussions during the study design phase of our study, and for feedback on preliminary results. We thank Emma White at the Bauer Core for details concerning available sequencing read lengths and kit sizes for Illumina instruments. We also thank members from the Harvard FAS Informatics Group for feedback on preliminary results produced by this study. This work was conducted on the traditional territory of the Wampanoag and Massachusett peoples.

REFERENCES

1. Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, et al. The evolution of gene expression levels in mammalian organs. Nature. 2011 Oct 19;478(7369):343–8.

2. Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, et al. The evolutionary landscape of alternative splicing in vertebrate species. Science. 2012 Dec 21;338(6114):1587–93.

3. Frumkin I, Yofe I, Bar-Ziv R, Gurvich Y, Lu Y-Y, Voichek Y, et al. Evolution of intron splicing towards optimized gene expression is based on various Cis- and Trans-molecular mechanisms. PLOS Biol. 2019 Aug 23;17(8):e3000423.

4. Crowley JJ, Zhabotynsky V, Sun W, Huang S, Pakatci IK, Kim Y, et al. Analyses of allele-specific gene expression in highly divergent mouse crosses identifies pervasive allelic imbalance. Nat Genet. 2015 Apr;47(4):353–60.

5. Verta J-P, Jones FC. Predominance of cis-regulatory changes in parallel expression divergence of sticklebacks. de Meaux J, Tautz D, editors. eLife. 2019 May 15;8:e43785.

6.   Castel SE, Cervera A, Mohammadi P, Aguet F, Reverter F, Wolman A, et al. Modified
penetrance of coding variants by cis-regulatory variation contributes to disease risk. Nat
Genet. 2018;50(9):1327–34.

7.   Gandal MJ, Zhang P, Hadjimichael E, Walker RL, Chen C, Liu S, et al. Transcriptome-wide
isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. Science. 2018 Dec
14;362(6420):eaat8127.

8.   Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV. Winter
storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard.
Science. 2017 Aug 4;357(6350):495–8.

9.   Mallarino R, Henegar C, Mirasierra M, Manceau M, Schradin C, Vallejo M, et al.
Developmental mechanisms of stripe patterns in rodents. Nature. 2016 24;539(7630):518–
23.

10. MacManes MD. On the optimal trimming of high-throughput mRNA sequence data. Front
Genet [Internet]. 2014 Jan 31 [cited 2019 Sep 16];5. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3908319/

11. Williams CR, Baccarella A, Parrish JZ, Kim CC. Trimming of sequence reads alters RNA-Seq
gene expression estimates. BMC Bioinformatics. 2016 Feb 25;17:103.

12. Tarazona S, García-Alcalde F, Dopazo J, Ferrer A, Conesa A. Differential expression in RNA-
seq: a matter of depth. Genome Res. 2011 Dec;21(12):2213–23.

13. Liu Y, Zhou J, White KP. RNA-seq differential expression studies: more sequence or more
replication? Bioinforma Oxf Engl. 2014 Feb 1;30(3):301–4.

14. Todd EV, Black MA, Gemmell NJ. The power and promise of RNA-seq in ecology and
evolution. Mol Ecol. 2016 Mar;25(6):1224–41.

15. Robles JA, Qureshi SE, Stephen SJ, Wilson SR, Burden CJ, Taylor JM. Efficient experimental
design and analysis strategies for the detection of differential expression using RNA-
Sequencing. BMC Genomics. 2012 Sep 17;13(1):484.

16. Fonseca NA, Marioni J, Brazma A. RNA-Seq Gene Profiling - A Systematic Empirical
Comparison. PLOS ONE. 2014 Sep 30;9(9):e107026.

17. Kanitz A, Gypas F, Gruber AJ, Gruber AR, Martin G, Zavolan M. Comparative assessment of
methods for the computational inference of transcript isoform abundance from RNA-seq
data. Genome Biol. 2015 Jul 23;16:150.

18. Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of RNA-seq
incorporating quantification uncertainty. Nat Methods. 2017 Jul;14(7):687–90.

19. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000Research. 2015 Dec 30;4:1521.

20. Soneson C, Delorenzi M. A comparison of methods for differential expression analysis of
RNA-seq data. BMC Bioinformatics. 2013 Mar 9;14(1):91.

21. Gaspar JM. NGmerge: merging paired-end reads via novel empirically-derived models of
sequencing errors. BMC Bioinformatics. 2018 Dec 20;19(1):536.

22. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol. 2016 May;34(5):525–7.

23. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics. 2011;12:323.

24. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012
Apr;9(4):357–9.

25. Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model analysis
tools for RNA-seq read counts. Genome Biol. 2014;15:R29.

26. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

Fig. 1. Spearman's rank correlations for kallisto-derived transcripts per million (TPM) between the gold standard paired-end 2x125 strategy and alternative strategies. Violin plots of **(a)** transcript and **(b)** gene-level inference. Comparison of correlations with 2x125 between 2x40 and 1x75 for **(c)** transcript and **(d)** genes, and between 2x40 and 1x125 for **(e)** transcript and **(f)** genes. For **c**-f, points above the red dotted line are samples where estimates of expression from 2x40 is more highly correlated with the gold standard than the contrasted single-end strategy.

Fig. 2. Spearman's rank correlations for RSEM-derived transcripts per million (TPM) between the gold standard paired-end 2x125 strategy and alternative strategies. Violin plots of **(a)** transcript and **(b)** gene-level inference. Comparison of correlations with 2x125 between 2x40 and 1x75 for **(c)** transcript and **(d)** genes, and between 2x40 and 1x125 for **(e)** transcript and **(f)** genes. For **c**-f, points above the red dotted line are samples where estimates of expression from 2x40 is more highly correlated with the gold standard than the contrasted single-end strategy.

Fig. 3. Effects of kallisto pseudo-alignment efficiency on differences in TPM estimate Spearman's rank correlations between 2x40 vs. 1x75. Efficiency of 2x40 is presented as the total kallisto count for 2x40 for **(a)** transcripts and **(b)** genes, and contrasted with the efficiency of 1x75 in the form of count ratios for **(c)** transcripts and **(d)** genes.

Fig. 4. Histograms of differences in root-mean-square error (RMSE) between 2x40 and either 1x75 or 1x125, where RMSE is calculated using 2x125 as the gold standard. RMSE differences for kallisto for **(a)** transcripts and **(b)** genes, and RSEM for **(c)** transcripts and **(d)** genes. The vertical black dashed line indicates the threshold above which RMSE for either 1x75 or 1x125 is greater than that for 2x40.

Fig. 5. Variation across SRA accessions in **(a,b)** false negative rate, **(c,d)** empirical false discovery rate, and **(e,f)** AUC between 2x40, 1x75 and 1x125 sequencing strategies based upon differential expression testing with sleuth. Differences are plotted as means for Wald tests within accessions, for **(a,c,e)** transcripts and **(b,d,f)** genes.