Short paired-end reads trump long single-end reads for expression analysis

Adam H. Freedman, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

John M. Gaspar, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

Timothy B. Sackton*, Faculty of Arts and Sciences Informatics Group, Harvard University,

Cambridge, Massachusetts 02138

* Corresponding author: tsackton@g.harvard.edu

ABSTRACT

Background: Typical experimental design advice for expression analyses using RNA-seq

generally assumes that single-end reads provide robust gene-level expression estimates in a cost-

effective manner, and that the additional benefits obtained from paired-end sequencing are not

worth the additional cost. However, in many cases (e.g., with Illumina NextSeq and NovaSeq

instruments), shorter paired-end reads and longer single-end reads can be generated for the same

cost, and it is not obvious which strategy should be preferred. Using publicly available data, we

test whether short-paired end reads can achieve more robust expression estimates and differential

expression results than single-end reads of approximately the same total number of sequenced

bases.

Results: At both the transcript and gene levels, 2x40 paired-end reads unequivocally provide

expression estimates that are more highly correlated with 2x125 reads; in nearly all cases, those

correlations are greater than for 1x125, despite the greater total number of sequenced bases for

the latter. In addition, differential expression false negative rates are consistently lower for 2x40

compared to 1x125.

Conclusion: Researchers seeking a cost-effective approach for gene-level expression analysis

should prefer short paired-end reads over a longer single-end strategy. False negative rates for

gene-level differential expression tests with 2x40 relative to 2x125 are in fact quite modest. In

addition, short paired-end reads will also give reasonably robust expression estimates and

differential expression results at the isoform level.

BACKGROUND

For over a decade, RNA-seq has empowered gene expression analysis. This has led to

fundamental advances in our understanding of diverse phenomena, including the evolution of

gene regulation across species (1), alternative splicing (2,3), relative importance of cis vs -trans

regulation (4,5), genetic underpinnings of heritable disease (6,7), and the genetic architecture of

phenotypes in natural populations (8,9). While the cost of sequencing has steadily decreased,

making expression analyses feasible even on modest research budgets, maximizing cost

efficiency remains high priority. Cost effectiveness is particularly important as studies scale up

to hundreds or even thousands of samples. Maximizing performance given a particular study

design has already received much attention, with investigations of optimal strategies for quality

trimming (10), the effects of quality trimming on expression estimates and differential

expression testing (11), optimal sequencing depth (12,13), importance of biological replicates

(12,14,15), and the relative performance of expression and differential expression tools and

pipelines (16–20). Less attention has focused on how to improve performance given a fixed

number of biological replicates and total sequenced bases.

While the typical length of sequenced reads has increased as costs have decreased, when

an annotated genome is available it remains standard procedure to sequence single end reads to

estimate gene expression, typically around 75 base pairs. Yet, this strategy does not leverage

information provided by the full length of the library fragments, particularly the greater

specificity in read mapping (or pseudo-alignment) that would be provided by sequencing from

both ends of a fragment. Given a fixed cost per base for a given Illumina instrument, paired-end

75bp reads would be twice as expensive as single-end 75bp sequencing, assuming an equivalent

number of sequenced fragments. Intuitively, however, a sizeable fraction of the long-range

information provided by paired-end sequencing should still be retained by sequencing shorter

paired-end reads. In this study, we evaluate whether that intuition is correct, by assessing

whether expression estimates and differential expression results obtained with 2x40 paired end

sequencing are more consistent with 2x125 paired end sequencing than either 1x75 or 1x125

single-end sequencing. Our particular focus is on 1x75, as the cost of 1x75 and 2x40 are

approximately the same per read when sequencing using the Illumina NextSeq, a common

instrument for RNA-seq studies. As with NextSeq, NovaSeq sequencing kits do not distinguish

between single and paired-end libraries, and are configured a particular number of bases. Thus,

for the 100bp NovaSeq kit, a similar sequencing strategy trade-off exists between sequencing

2x50 vs. 1x100 reads. We conduct these analyses with publicly available Illumina sequence data

across 12 different SRA (Additional file 1: Table S1). accessions encompassing diverse

experimental designs and multiple model organisms with high quality annotated genomes.

RESULTS AND DISCUSSION

Our approach to evaluating the performance of short paired-end reads vs. longer single-end ones

is to compare expression estimates and differential expression results derived from these

strategies to a truth set. Because we cannot know the "truth", we assume that its closest

approximation are results obtained from long, paired-end reads. Thus, we define our gold

standard as the results obtained from 2x125 paired-end reads. We then trim these data down to

two data sets with read lengths of 40 and 75 base pairs, and calculate spearman's rank

correlations between transcripts-per-million (TPM) estimates based upon 2x40, 1x75 and 1x125

with the 2x125 gold standard. We present these correlations in two different ways. First, we

examine the distributions of those correlation coefficients across alternative sequencing

strategies. Second, we examine relative differences in these correlations in the form of "ratio

plots" that have, on the x-axis, the ratio of the 1x125 expression correlation with 2x125 to that of

2x40, and on the y-axis, the ratio of the expression correlation between 1x75 and 2x125 to that of

2x40. In this manner, we can visualized in bivariate space, the frequency of cases where 2x40

outperforms both 1x75 and 1x125, where 2x40 only outperforms 1x75 (or 1x125), or where 2x40

is inferior to both single-end strategies.

Kallisto transcript-level TPM estimates made with 2x40 are always more highly

correlated with those made with the 2x125 gold standard than those made with data trimmed to

1x75 reads (Fig. 1a,c), and perform better than estimates made with data trimmed to 1x125 for

all but a few samples from one SRA accession (Fig. 1c). A similar performance advantage for

the 2x40 strategy compared to the 1x75 strategy is observed at the gene level (Fig. 1 b,d).

RSEM-based TPM estimates show a generally consistent overall pattern (Fig. 2), although at the

transcript level four samples have higher performance (better correlations with estimated from

untrimmed data) for one or both single-end strategies than the 2x40 strategy. These samples are

widely dispersed in the rank correlation ratio plots (Fig. 2c); two of the four samples have an

aberrant pattern where TPM estimates from 1x75 reads more highly correlated with estimated

from 2x125 reads than with estimates from 1x125 reads, despite having 67% more nucleotides

sequenced per sample. Upon closer inspection, we discovered that samples comprising this

unusual pattern had low bowtie2 alignment rates to the reference transcripts (for the 2x125

libraries, overall alignment rates of 37.1-51.0%), suggesting library quality issues were

responsible.

More generally, a number of different library-specific factors could potentially associate

with variation in the relative performance of a 2x40 strategy. One important faction is simply the

overall number of informative reads in an experiment. To estimate this, we summed the kallisto

transcript-level counts for each sample. We find that as the number of informative reads

increases, the relative performance of a 2x40 strategy (compared to 1x125) improves (Fig 3). We

did not perform the same analysis for the 1x75 strategy, as using 2x40 reads always

outperformed 1x75 reads at the transcript level with kallisto. The correlation between total

number of sequenced reads and relative performance is slightly weaker ($\rho = 0.46$, $p = 4.4 \times 10^{-10}$),

indicating that read mappability in addition to read number, is important for robustness of

expression estimates for short PE reads. While there are no obvious taxonomic effects, there are

clear library-specific effects, manifested as differences in relative 2x40 performance among

accessions derived from the same species.

We next evaluated the extent to which differences in performance with respect to

estimating expression levels translated into downstream effects on tests of differential

expression. Within each SRA accession and for each pair of conditions and for each sequencing

strategy, we conducted Wald tests with both sleuth and limma-voom. Similar to our evaluation of

expression estimate robustness, we defined "true" differential expression signals as those

recovered with 2x125, given a false discovery rate (FDR) of 0.01.

Sequencing strategy clearly impacts downstream differential expression analyses. Using

either the kallisto-sleuth or RSEM-limma pipeline, at both the transcript and gene level, false

negative rate relative to the 2x125 gold standard was elevated in 1x75 relative to 2x40 (Fig. 4),

while false positive rates remained relatively consistent (Additional file 1: Table S1). With few

exceptions, false negative rate was very similar between 2x40 and 1x125, or slightly lower in the

former.

CONCLUSIONS

When designing a gene expression study using bulk RNA-seq, given a fixed budget, more robust

expression estimates and differential expression test results can be obtained from short paired-

end reads compared to longer single-end reads. In other words, the extra information provided by

reads pairs during the mapping or pseudoalignment phase outweighs any potential penalty of

short reads. In almost all cases, short paired-end reads trump single-end reads that sequence

approximately the same number of bases. Short paired-end reads provide an additional benefit in

that they can identify a large fraction of differentially expressed isoforms, for the cost of gene-

level analysis with longer single-end reads. From a practical perspective, neither NextSeq nor

NovaSeq sequencing kits distinguish between single and paired-end libraries. In the experience

of support staff at the Bauer sequencing core facility at Harvard, one can get 80 bp from the

NextSeq 75 bp kit, such that 2x40 is a viable sequencing design. Alternatively, the smallest

NovaSeq kit is for 100 cycles, which can be used to sequence paired 50bp reads, allowing one to

leverage that instrument's much smaller cost per sequenced base. Overall, our findings suggest

that for a given number of sequencing cycles, the most efficient design for RNA-seq expression

analysis is allocating 50% of the cycles to paired end reads, instead of 100% of the cycles to a

longer, single-end read.

METHODS

Datasets

From NCBI's Short Read Archive, we downloaded paired-end RNA-seq reads from 12 different

studies (Table 1), representing model organisms with well-annotated reference genomes: mouse

(*Mus musculus*), human (*Homo sapiens*), yeast (*Saccharomyces cerevisiae*), fly (*Drosophila*

melanogaster), and worm (*Caenorhabditis elegans*). In selecting data sets, we required minimum

pre-processing read lengths to be ≥ 125 bp, and experimental designs to contain ≥ 2 experimental

conditions.

Short-read processing

We removed adapters and trimmed reads with NGmerge (16–20). For each sample, we trimmed

reads to 125, 75, and 40bp using awk. For downstream analyses, we used 2x125, 2x40,1x125,

and 1x75 trimmed data sets. Single-end data sets were comprised of R1 from the original paired

reads.

Expression analyses

For each sample-sequencing strategy combination, we estimated transcript-level expression with

kallisto (22). We designated abundance estimates derived from 2x125 as our gold standard

against which to benchmark all other strategies. We used spearman's rank correlations to

quantify the degree of concordance between alternative sequencing strategies and the gold

standard in terms of transcript per million (TPM). We performed similar analyses at the gene

level by aggregating transcript-level TPMs using the R package Tximport (19).

Because the majority of RNA-seq applications involve tests of differential expression

(DE), we compared differential expression test results between the gold standard 2x125 and

alternative strategies. Specifically, for each experiment we conducted all possible pairwise Wald

tests of transcript-level DE with sleuth (18) version 0.30.0, and used p-value aggregation

(https://pachterlab.github.io/sleuth_walkthroughs/pval_agg/analysis.html) for gene-level

analyses. To compare performance of alternative strategies relative to the gold standard, we

calculated false negative and false positive rates, defining significant tests as those with

Benjamini-Hochberg adjusted p-values ≤ an FDR of 0.01.

In order to evaluate whether our findings were pipeline-specific, perhaps related to

particular features of tools such as kallisto that rely on pseudo-alignment, for 5 of the 12 SRA

accessions, we performed expression analyses with an alternative pipeline, involving estimation

of expression levels with RSEM (23) version 1.2.29 based upon alignments with bowtie2 version

2.3.2, (24) and differential expression testing with limma-voom (25) version 3.38.3. For limma-

voom, we only conducted tests where, for each gene or transcript,there were ≥ 2 samples with

counts per million (CPM) ≥ 1. In addition, we employed TMM normalization, and the

voomWithQualityWeights option. Sleuth and limma-voom have previously been demonstrated to

be top performers that correctly control FDR (22). When SRA metadata enabled us to determine

that libraries were stranded, we included the relevant RSEM command line arguments for

restricting bowtie2 alignments to the correct read pair configuration. For both sleuth and limma-

voom based analyses, we only calculated performance metrics for Wald tests where, at the

specified false discovery rate (FDR) of 0.01, there were ≥ 50 significantly differentially

expressed features. All reference genomes and gtf-formatted annotations against which to

estimate expression were downloaded from Ensembl, and transcripts for kallisto-based analyses

were extracted using the *extract-reference-transcripts* utility in RSEM.

CODE AVAILABILITY

Command lines for executing NGmerge, kallisto, RSEM, sleuth, and limma-voom analyses are

available on GitHub at https://github.com/harvardinformatics/rnaseq_readlength_assessment.

ACKNOWLEDGMENTS

The computations in this paper were run on the Odyssey cluster supported by the FAS Division

of Science, Research Computing Group at Harvard University. We thank Claire Reardon,

director of the Bauer Core at Harvard University for helpful discussions during the study design

phase of our study, and for feedback on preliminary results. We also thank Emma White at the

Bauer Core for details concerning available library preparation options on Illumina instruments.

We also thank members from the Harvard FAS Informatics Group for feedback on preliminary

results produced by this study. This work was conducted on the traditional territory of the

Wampanoag and Massachusett peoples.

AUTHORS' CONTRIBUTIONS

TBS and AHF designed the experiments, AHF and JMG analyzed the data, and AHF wrote the

paper with TBS.

REFERENCES

1. Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, et al. The evolution

    of gene expression levels in mammalian organs. Nature. 2011 Oct 19;478(7369):343–8.

2. Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, et al. The

    evolutionary landscape of alternative splicing in vertebrate species. Science. 2012 Dec

    21;338(6114):1587–93.

3. Frumkin I, Yofe I, Bar-Ziv R, Gurvich Y, Lu Y-Y, Voichek Y, et al. Evolution of intron

    splicing towards optimized gene expression is based on various Cis- and Trans-molecular

    mechanisms. PLOS Biol. 2019 Aug 23;17(8):e3000423.

4.   Crowley JJ, Zhabotynsky V, Sun W, Huang S, Pakatci IK, Kim Y, et al. Analyses of allele-

specific gene expression in highly divergent mouse crosses identifies pervasive allelic

imbalance. Nat Genet. 2015 Apr;47(4):353–60.

5.   Verta J-P, Jones FC. Predominance of cis-regulatory changes in parallel expression

divergence of sticklebacks. de Meaux J, Tautz D, editors. eLife. 2019 May 15;8:e43785.

6.   Castel SE, Cervera A, Mohammadi P, Aguet F, Reverter F, Wolman A, et al. Modified

penetrance of coding variants by cis-regulatory variation contributes to disease risk. Nat

Genet. 2018;50(9):1327–34.

7.   Gandal MJ, Zhang P, Hadjimichael E, Walker RL, Chen C, Liu S, et al. Transcriptome-wide

isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. Science. 2018 Dec

14;362(6420):eaat8127.

8.   Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV. Winter

storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard.

Science. 2017 Aug 4;357(6350):495–8.

9.   Mallarino R, Henegar C, Mirasierra M, Manceau M, Schradin C, Vallejo M, et al.

Developmental mechanisms of stripe patterns in rodents. Nature. 2016 24;539(7630):518–

23.

10. MacManes MD. On the optimal trimming of high-throughput mRNA sequence data. Front

Genet [Internet]. 2014 Jan 31 [cited 2019 Sep 16];5. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3908319/

11. Williams CR, Baccarella A, Parrish JZ, Kim CC. Trimming of sequence reads alters RNA-

Seq gene expression estimates. BMC Bioinformatics. 2016 Feb 25;17:103.

12. Tarazona S, García-Alcalde F, Dopazo J, Ferrer A, Conesa A. Differential expression in

RNA-seq: a matter of depth. Genome Res. 2011 Dec;21(12):2213–23.

13. Liu Y, Zhou J, White KP. RNA-seq differential expression studies: more sequence or more

replication? Bioinforma Oxf Engl. 2014 Feb 1;30(3):301–4.

14. Todd EV, Black MA, Gemmell NJ. The power and promise of RNA-seq in ecology and

evolution. Mol Ecol. 2016 Mar;25(6):1224–41.

15. Robles JA, Qureshi SE, Stephen SJ, Wilson SR, Burden CJ, Taylor JM. Efficient

experimental design and analysis strategies for the detection of differential expression using

RNA-Sequencing. BMC Genomics. 2012 Sep 17;13(1):484.

16. Fonseca NA, Marioni J, Brazma A. RNA-Seq Gene Profiling - A Systematic Empirical

Comparison. PLOS ONE. 2014 Sep 30;9(9):e107026.

17. Kanitz A, Gypas F, Gruber AJ, Gruber AR, Martin G, Zavolan M. Comparative assessment

of methods for the computational inference of transcript isoform abundance from RNA-seq

data. Genome Biol. 2015 Jul 23;16:150.

18. Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of RNA-seq

incorporating quantification uncertainty. Nat Methods. 2017 Jul;14(7):687–90.

19. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. F1000Research. 2015 Dec 30;4:1521.

20. Soneson C, Delorenzi M. A comparison of methods for differential expression analysis of

RNA-seq data. BMC Bioinformatics. 2013 Mar 9;14(1):91.

21. Gaspar JM. NGmerge: merging paired-end reads via novel empirically-derived models of

sequencing errors. BMC Bioinformatics. 2018 Dec 20;19(1):536.

22. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

quantification. Nat Biotechnol. 2016 May;34(5):525–7.

23. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. BMC Bioinformatics. 2011;12:323.

24. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012

Apr;9(4):357–9.

25. Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model analysis

tools for RNA-seq read counts. Genome Biol. 2014;15:R29.

Fig. 1. Spearman rank correlations for kallisto-derived transcripts per million (TPM) between the gold standard paired-end 2x125 strategy and alternative strategies for **(a)** transcript and **(b)** gene-level inference. Comparison of correlations with 2x125 in the form of ratios for **(c)** transcript and **(d)** gene levels. Points above the red dotted line and to the right of the black dotted line indicate samples (colored by SRA accession) where 1x75 and 1x125, respectively, are more highly correlated with the gold standard that 2x40.

Fig. 2. Spearman rank correlations for RSEM-derived transcripts per million (TPM) between the gold standard paired-end 2x125 strategy and alternative strategies for **(a)** transcript and **(b)** gene-level inference. Comparison of correlations with 2x125 in the form of ratios for **(c)** transcript and **(d)** gene levels. Points above the red dotted line and to the right of the black dotted line indicate samples (colored by SRA accession) where 1x75 and 1x125, respectively, are more highly correlated with the gold standard that 2x40.

Fig. 3. Ratio of the Spearman rank correlation between 2x40 transcript-level TPM and 2x125 TPM to that between 1x125 and 2x125, as a function of the sum of kallisto counts across all transcripts for a given library.

Fig. 4. Violin plots of false negative rate at FDR = 0.01 for differential expression Wald tests, along with median trend "reaction norms" per data set across sequencing strategy for sleuth transcript **(a)** and **(b)** gene, and limma voom **(c)** transcript and **(d)** gene-level inference.

Fig. S1. Violin plots of false positive rate at FDR = 0.01 (red horizonal line) for differential expression Wald tests, along with median trend "reaction norms" per data set across sequencing strategy for sleuth transcript **(a)** and **(b)** gene, and limma voom **(c)** transcript and **(d)** gene-level inference.

Table S1. SRA accessions used in this study.

Accession	Organism	Instrument	Paired-end read length	Conditions	Biological replicates per condition	Kallisto	RSEM
SRP133853	*H. sapiens*	HiSeq 2500	125	6	3	x	x
SRP115815	*H. sapiens*	HiSeq 2500	125-126	9	3	x	
SRP105271	*M. musculus*	HiSeq 2000	125	5	3-4	x	
SRP143508	*M. musculus*	HiSeq X Ten	150	2	3	x	
SRP096374	*M. musculus*	HiSeq 4000	150	2	3	x	x
ERP017328	*D. melanogaster*	HiSeq 2500	126	2	3	x	
SRP128516	*D. melanogaster*	HiSeq 4000	151	2	3	x	x
SRP089981	*C. elegans*	NextSeq 500	151	4	5	x	
SRP092256	*C. elegans*	HiSeq 2500	126	7	5	x	
SRP129557	*C. elegans*	HiSeq 3000	126	4	3	x	x
SRP133093	*S. cerevisiae*	HiSeq 2000	151	4	3	x	x
SRP142501	*S. cerevisiae*	HiSeq X Ten	150	2	3	x	