Overcoming coarse coding in visual cortex via multiplexing:
neural correlations differ dramatically when stimulus bundles are presented

Na Young Jun (1-3)

Douglas A. Ruff (4-5)

Lily E. Kramer (4-5)

Brittany Bowes (4-5)

Surya T Tokdar (6)

Marlene R Cohen (4-5)

Jennifer M Groh (1-3, 7-9)

1. Department of Neurobiology, Duke University
2. Center for Cognitive Neuroscience, Duke University
3. Duke Institute for Brain Sciences
4. Department of Neuroscience, University of Pittsburgh
5. Center for the Neural Basis of Cognition, University of Pittsburgh
6. Department of Statistical Science, Duke University
7. Department of Psychology and Neuroscience, Duke University
8. Department of Biomedical Engineering, Duke University
9. Department of Computer Science, Duke University

Acknowledgments:

Thanks to: Valeria C. Caruso, Yunran Chen, Jeffrey T. Mohl, Meredith N. Schmehl, Shawn M. Willett, for helpful comments on the analysis and manuscript. This work was supported by: National Institutes of Health grant nos. R00EY020844 (MRC), R01EY022930 (MRC); Core Grant P30 EY008098s (MRC); R01DC013906 (JMG, STT); and R01DC016363 (JMG, STT); and by support to MRC from the McKnight, Whitehall, Sloan and Simons Foundations.

Abstract

Sensory receptive fields are large enough that they can contain more than one perceptible stimulus. How, then, can the brain encode information about *each* of the stimuli that may be present at once? We recently showed that when more than one stimulus is present, single neurons can fluctuate between coding one vs. the other(s) across some time period, suggesting a form of neural multiplexing of different stimuli (Caruso et al., 2018). Here we investigate how such fluctuations are coordinated across the neural population by analyzing the spike count ("noise") correlations observed between pairs of visual cortex (V1, V4) neurons when stimuli are presented jointly. We found a novel pattern of V1 noise correlations: distinct distributions of positive and negative values, depending on whether the two neurons in the pair have similar or different preferences for the component stimuli. Distinct distributions of spike count correlations based on tuning preferences were also seen in V4 for adjacent

stimuli, but were not observed in either structure for single stimuli or when the two gratings were superimposed and formed a single plaid. Importantly, the bimodal correlation patterns were most pronounced among pairs of neurons showing the strongest evidence for multiplexing. These findings suggest multiple stimuli evoke different response dynamics than those evoked by single stimuli, lending support to the multiplexing hypothesis and suggesting a means by which information about multiple stimuli can be preserved despite the apparent coarseness of sensory coding.

Significance Statement

How the brain separates information about multiple objects despite overlap in the neurons responsive to each item is not well understood. Here we show that pairs of neurons in primate V1 and V4 show unique patterns of correlated firing when there are multiple objects in the visual scene. These patterns were most pronounced in neurons that showed fluctuating activity consistent with switching between encoding each object across time (i.e. time division multiplexing). Both strongly positive and strongly negative correlations were observed, depending on whether the neurons in the pair exhibited similar or different stimulus preferences. These patterns were absent when only one object was presented, suggesting that they may play a key role in preserving information about multiple items.

Key Words
Spike count correlations; variability; V1; multiplexing; attention; population coding

Introduction

Receptive fields are too large to account for our perceptual sensitivity. For example, in primary visual cortex, receptive fields are considerably larger than the spatial limits of visual acuity (Dow et al., 1981; Keliris et al., 2019). The letters on the page you are reading now are < 0.25 degree apart, but foveal V1 receptive fields are approximately 0.5-2 degrees in diameter (Alonso and Chen, 2009). In the auditory system, mammalian neurons may be responsive to nearly any location in space (e.g. Woods et al., 2001; Groh et al., 2003; McAlpine and Grothe, 2003; Werner-Reiss and Groh, 2008; Grothe et al., 2010; Higgins et al., 2010), and even frequency tuning is broad: between 40-80% of multiunit clusters in the primate inferior colliculus will respond to a pure tone of frequency ~500 Hz at the fairly quiet sound level of 50 dB SPL (Bulkin and Groh, 2011). Such breadth of tuning means that individual neurons commonly experience more than one stimulus to which they could in principle respond, making it unclear how information about multiple objects is preserved. Much previous work concerning the coarseness of neural tuning has occurred in the motor domain, where the concept of a population code in which many neurons "vote" for a particular course of action has provided substantial explanatory power (e.g. (Georgopoulos et al., 1986; Lee et al., 1988)). However, such coarse population codes seem most

appropriate in a motor context because only one movement involving a particular set of muscles will be made at a time, and the same strategy seems unlikely to apply in sensory contexts where information preservation is a goal.

Logic suggests that information about each distinct stimulus must be <u>segregated</u> within the neural code in some fashion, either into exclusive neural subpopulations, different epochs of time, or some combination of both. We have recently presented the hypothesis that the nervous system might employ a form of neural turn-taking, (time division multiplexing) in which individual neurons fluctuate between responding to each of the items in or near their receptive fields across various epochs of time (Caruso et al., 2018; Mohl et al., 2020). Such a coding scheme could preserve information about each stimulus across time and/or across neural subpopulations.

One open question raised by this theory is whether and how activity patterns fluctuate together in a neural ensemble. For example, pairs of neurons that exhibit fluctuating activity in response to multiple stimuli might show either positive or negative correlations with each other. The pattern of such correlations across the population, and in particular how the observed correlations depend on the sensitivity of the two neurons in the pair to the individual component stimuli in the stimulus "bundle", can reveal insights into the degree to which the overall neural representation on any given stimulus presentation may be biased towards one stimulus vs. another.

A second open question concerns how the pattern of fluctuations and their correlation across the population might depend on whether multiple "stimuli" fuse as a single object or are perceived as distinct from each other (Milner, 1974; Gray and Singer, 1989; Von Der Malsburg, 1994; Singer and Gray, 1995; Gray, 1999) (but see Palanca and DeAngelis, 2005). Stimuli that segregate into distinct objects may be more likely to be associated with fluctuations in neural activity and their attendant correlations across neurons, whereas stimuli that fuse into a single distinct object may cause activity patterns that are akin to those observed when only one stimulus is present. Such a pattern would specifically implicate activity fluctuations in playing a role in the perceptual process of object segregation.

To address these questions, we turned to the primary visual cortex (V1). V1 allows for a strong test of these hypotheses since V1 neurons have comparatively small receptive fields and are therefore less subject to the multiple-stimulus-overlap problem than more broadly tuned areas such as the inferior colliculus or inferotemporal (IT) cortex that were assessed in our previous report (Caruso et al., 2018). Furthermore, the responses of V1 neurons are typically much less modulated by cognitive or other modulatory processes than neurons in higher visual areas (e.g. (Luck et al., 1997; Cook and Maunsell, 2002; Buffalo et al., 2010))

We evaluated activity in V1 while monkeys viewed either single stimuli (gratings) or two different types of combined stimuli (superimposed vs. adjacent gratings). When the two gratings were superimposed, they

presumably appeared as a single object, or plaid (Adelson and Movshon, 1982; Rodman and Albright, 1989; Heeger et al., 1996; Busse et al., 2009; Lima et al., 2010), whereas when they were adjacent they appeared as two distinct objects. We then evaluated the degree and sign of the spike count correlations (commonly referred to as "noise" correlations (Cohen and Kohn, 2011)) observed between pairs of simultaneously recorded units in response to presentations of particular stimulus conditions. We report here that the pattern of correlations varied dramatically depending on whether the stimuli were superimposed or presented side-by-side (one object vs. two). In the two-object case, the distribution of spike count correlations was markedly different from previous reports involving single stimuli (c.f. Table 1, Cohen and Kohn, 2011) , and encompassed a range spanning many negative correlations in addition to positive ones. Whether the correlations tended to be positive vs. negative depended on whether the two neurons in the pair preferred the same stimulus (median correlation +0.25) or preferred different stimuli (median correlation -0.05).   The distribution of spike count correlation values was even more widely spread among pairs of neurons that showed demonstrably fluctuating activity across stimulus presentations (using the statistical analyses we previously developed; same preference: +0.49 and different preference: -0.14).   In contrast, in the single object case, positive correlations predominated (median value +0.22). Distinct tuning-preference-related distributions of spike count correlations for adjacent stimuli but not for individual or fused stimuli were also seen in a smaller additional dataset in V4.

Overall, this pattern of results is consistent with the possibility that when two visual objects are presented in close proximity, a subpopulation of visual cortical neurons fluctuates in a coordinated fashion and biases the population code in favor of one object over the other on individual stimulus presentations.

Results

General experimental design and theoretical rationale

Recordings: The activity of neurons in visual cortex was recorded in three experimental designs in six monkeys (N=2 per experiment per brain area) using chronically implanted multielectrode arrays (Fig 1a, Table 1 in Methods). In the "superimposed" data set, the activity of neurons in V1 and V4 was recorded while monkeys passively fixated (for details, see (Ruff et al., 2016). In the "adjacent" data sets, the activity of V1 and V4 neurons were recorded while monkeys either passively fixated (V4 recordings) or fixated while performing an orientation change discrimination task involving either one of these stimuli or a third stimulus presented in the ipsilateral hemifield (V1) (for details, see (Ruff and Cohen, 2016).   Microsaccades were identified using a velocity detection algorithm (Engbert and Kliegl, 2003), and all of the stimulus presentations during which microsaccades were detected were excluded from the analysis (Ruff and Cohen, 2016). In the "superimposed" data set, the gratings were large, spanning the receptive fields of the recorded neurons, and were presented either individually or in combinations of two orthogonal gratings at a consistent location on every trial (Fig 1c). When the two gratings

were presented, they superimposed and formed a "plaid" stimulus. In the "adjacent" data sets, the stimuli were smaller Gabor patches (V1, V4, Figure 1d-e) or natural images (V4, Figure 1e, stimuli from (Long et al., 2018)) and were presented either individually or adjacent to one another. Together they spanned the receptive fields of the V1 or V4 neurons being recorded in a fashion similar to the "superimposed" experiment. For data collected during performance of the attention task (V1), we focused our analyses on trials in which the monkeys attended to the third stimulus and judged its orientation, i.e. attention was consistently directed away from either of the two adjacent Gabor patches that elicited responses in the neurons under study (Figure 1d). Trials in which the monkey was required to attend to one or the other of the adjacent Gabor patches were excluded from the analyses, as were incorrectly performed trials.

Rationale: The possible results and their interpretation are illustrated schematically in Figures 2 and 3. For simplicity, we show only the "adjacent" stimulus case (left column), with the two unattended adjacent Gabor patches "A" and "B", and we focus on hypothetical neurons that appear to fluctuate between responding to "A" and responding to "B" on different stimulus presentations. A statistical method for formally identifying such patterns is presented in Figure 5 and associated text (Caruso et al., 2018; Mohl et al., 2020).

In Figure 2, the column labeled "Neuron 1" depicts how one such neuron might respond to the "A" and "B" Gabor patches presented individually or in combination. For example, this neuron might respond rather modestly to "A", with spike counts ranging from 3-5 spikes (blue) across a series of three stimulus presentations. When "B" is presented, perhaps it responds more vigorously, with 7-9 spikes (red). When "A" and "B" are presented in combination, it is theoretically possible that Neuron 1's spike counts are drawn from either its "A-like" or "B-like" spike count distributions, and which distribution is somehow selected varies across presentations. These spikes are illustrated in red and blue to indicate the distribution they are presumably drawn from. This is referred to as a "mixture" pattern (Caruso et al., 2018; Mohl et al., 2020). It is important to note that the mixture pattern depicted in Figure 2 is an exaggeration of what is likely to be found in the neural data, where actual response patterns can rarely be this easily identified as likely drawn from one distribution vs. another. It is also important to note that neurons whose responses are demonstrably consistent with fluctuating between A-like and B-like response patterns may represent a small fraction of the overall population of neurons – but will tend to coincide with those labelled as showing strong normalization, as previously described in this data set (Ruff et al., 2016). These points will be revisited in greater detail below.

Figure 1. Experimental design. A. Multiunit activity was recorded in V1 and V4 using chronically implanted 10x10 electrode arrays in six monkeys. B. In both "superimposed" and "adjacent" datasets, the stimuli were positioned to overlap with ("adjacent" dataset) or completely span ("superimposed" dataset) the centers of the receptive fields of the recorded neurons. C. In the "superimposed" dataset, gratings were presented either

individually or in combination at a consistent location and were large enough to cover the V1 and V4 receptive fields (stimulus size range: 2.5-7o). The combined gratings appeared as a plaid (rightmost panel). Monkeys maintained fixation throughout stimulus presentation and performed no other task. D. In the V1 "adjacent" dataset, Gabor patches were smaller (typically <1o) and were presented individually or side-by-side roughly covering the region of the V1 receptive fields. Monkeys maintained fixation while performing an orientation change detection task. The data analyzed in this study involved trials in which the monkeys were attending a third Gabor patch located in the ipsilateral hemifield to perform the orientation change detection. E. In the V4 "adjacent" dataset, the stimuli consisted of either Gabor patches or natural image stimuli, and monkeys performed a fixation task. Incorrectly performed trials and stimulus presentations during which we detected microsaccades are excluded from all analyses.

Figure 2. Schematic depiction of possible response patterns and interpretation. Three hypothetical neurons and their possible spiking activity on individual stimulus presentations are shown, for three presentations each of stimulus "A" alone, stimulus "B" alone, and stimuli "A" and "B" together. If neurons "pick" from their "A"-like and "B"-like response distributions when "A" and "B" are presented together, then several possible patterns of correlation might emerge between neurons, depending on whether the individual neurons in the pair "picks" the same stimulus ("Encode same") or different stimuli ("Encode different") and whether they share similar stimulus preferences (Neurons 1 and 2) vs. different stimulus preferences (Neurons 1 and 3). Neurons may also be uncorrelated with one another (not shown).

With these caveats in mind, let us now consider a second neuron, Neuron 2. Suppose this neuron, like Neuron 1, responds more vigorously to "B" (red spikes) than to "A" (blue spikes), and also shows activity that appears

drawn from either its A-like or B-like response distributions on combined AB presentations (bottom groupings of red and blue spike trains). We can then ask whether, on presentations when Neuron 1 "picks" A, does Neuron 2 also tend to "pick" A? And when Neuron 1 "picks" B, does Neuron 2 also tend to "pick" B?[1] If so, then there would tend to be a positive spike count correlation between Neuron 1 and Neuron 2 on combined stimulus presentations ("encode same" branch). Note that this correlation could be larger than customary values of spike count correlations typically observed on single stimulus presentations, depending on how reliably the two neurons "choose" the same stimulus and how different their firing rates are on "A" vs. "B" stimulus presentations.

In contrast, if Neuron 2 and Neuron 1 tend to "pick" differently, with Neuron 2 exhibiting spike counts that appear to be drawn from its "B"-like distribution on presentations that Neuron 1 exhibits spike counts that appear drawn from its "A"-like distribution and vice versa ("encode different" branch), then the spike count correlations between this pair on combined stimulus presentations will be negative, and could be strongly so depending again on the reliability of the underlying anticorrelation together with the difference in the response distributions on the "A" and "B" presentations alone.

Now imagine a third neuron, Neuron 3, which shows an opposite stimulus selectivity as Neurons 1 and 2: this neuron responds more vigorously to "A" (blue spikes) than to "B" (red spikes). If this neuron also fluctuates between "A"-like activity and "B"-like activity on the combined "AB" presentations, it too can be either positively or negatively correlated with Neuron 1. If it "encodes same" (left branch), its spike count correlations will be *negative* in sign (when paired with Neuron 1). If it "encodes different", it will now be *positively* correlated in spike count with Neuron 1: The larger number of spikes associated with stimulus A in Neuron 3 will tend to occur on presentations in featuring a large number of spikes associated with stimulus B in Neuron 1, and with the smaller numbers of B-like spikes for Neuron 3 co-occurring with smaller numbers of A-like spikes occurring for Neuron 1.

In short, positive vs. negative spike count correlations in response to combined stimuli will have different interpretations depending on whether the two neurons in the pair both respond more vigorously to the same component stimulus or to different component stimuli.

---

[1] We set aside as beyond our present scope what it means to "pick" one or the other distribution and what the neural mechanism that might accomplish such an operation might look like; see Caruso et al. (2018) for discussion of this point.

Figure 3. Population-level predictions under different scenarios. Schematic histograms of the possible patterns of spike count correlations across all the pairs of recorded neurons as a function of whether they share the same stimulus preferences or have different preferences, and whether they tend to "encode" the same stimulus or different stimuli on each presentation (panel f is panel a + panel c; g = a + d; h = b + e).

The possible patterns of spike count correlations across a population of pairs of neurons are illustrated in Figure 3. If the population of pairs of neurons that prefer the same individual stimulus reliably follow the "encode-same" pattern, then the distribution of spike count correlations for this subpopulation should be positively skewed (Fig 3A). If the pairs of same-preference neurons include a mixture of "encode-same" and "encode-different" response patterns, then the overall distribution will range from negative to positive (Fig 3B).

Finally, we note that it is not possible for all same-preference neuron pairs to involve "encode-different" response patterns for the two-stimulus case. Imagine another neuron, Neuron 4, which is like Neurons 1 and 2 in its response pattern to the individual stimuli "A" and "B". If Neurons 1 and 2 "encode different" and Neurons 2 and 4 also "encode-different" in relation to each other, then Neurons 1 and 4 will "encode-same" with respect to each other. Neurons 1 and 2 will be negatively correlated and neurons 2 and 4 will be negatively correlated on "AB" presentations, but Neurons 1 and 4 will tend to be positively correlated.

In contrast, if "different preference" pairs tend to "encode-same", then the overall distribution of spike count correlations in this subgroup will be skewed towards negative values (Figure 3C), whereas if they tend to "encode-different", the distribution will skew positive (Figure 3D). A mixture of "encode-same" and "encode-

different" will be a broader distribution ranging from negative to positive, depicted here schematically as two distinct categories for simplicity (Figure 3E).

Next we consider how the "same preference" and "different preference" subpopulations might look when combined. If both subgroups preferentially "encode same", then there will be two distinct distributions, negatively-skewed in the "different preference" subgroup and positively skewed in the "same preference" subgroup (Figure 3F). "Encode same" among "same preference" and "encode different" among "different preference" would yield an overall positively skewed distribution (Figure 3G), and a mixture of different strategies in both subpopulations would yield a mix of positive and negative values for both subgroups (Figure 3H).

It should be noted that it is likely that all spike count correlations between pairs of neurons ride on an overall wave of at least slight positivity, due to shared sensitivity to non-stimulus-related factors like overall arousal level or satiety-related signals that might accompany task performance. Thus, the negative- and positive-modes of a broad distribution may not be symmetric around 0 but slightly shifted towards the positive side.

With two objects, distinct distributions of positive and negative spike count correlations occur in V1

We now turn to the actual results, with these predictions in mind. We examined the Pearson's correlation between the spike count responses observed during presentations of superimposed or adjacent stimuli for pairs of single or multiunits in each data set. This spike count correlation (r_{sc}), also commonly called a noise correlation, was computed for a given pair of units and for a given single stimulus or combination of simultaneously presented stimuli. Each pair of units contributed multiple r_{sc} values to the population distribution, one value for each relevant stimulus condition. See methods for additional details.

Figure 4 A-F illustrates results involving three V1 example units from the adjacent-stimuli dataset, selected for showing spike count distributions on dual stimulus presentations (black lines) that reflect a mixture of the distributions evident on the corresponding single stimulus presentations (red and blue lines; the formal test for "mixtures" is described in next section). The dual-stimulus distributions of spike counts are over-dispersed compared to what would be expected if the spikes on dual stimulus presentations were generated from a similar Poisson process as the single stimulus presentations, and a tendency for bimodality with modes near the modes for each of the individual stimulus presentations is evident.

The units in panels a and b share a similar stimulus preference to each other: stimulus "A", elicits higher spike counts (red line) than stimulus "B" (blue line) for both. The unit in panel c has the opposite preference, with higher spike counts for "B" than for "A". Panels D-F show the activity of each of these units on individual stimulus presentations plotted against the others. The pattern of spike count correlation on individual stimulus

presentations varies depending on the stimulus tuning preferences, with the one pairing between units that shared the same preference being positive (0.56, panel d) and the two pairings involving different tuning preferences yielding negative spike count correlations (-0.45, -.34, panels e and f). This pattern is borne out when the full set of pairings involving the unit in (a) and other units recorded at the same time that also showed "mixture" response patterns is considered (panel g): all the pairings that involve "same" tuning preferences yield positive correlations, and all of these correlations are individually significant (green bars, $p<0.05$). In contrast, all the pairings that involve "different" tuning preferences yield negative correlations (brown bars); as expected, these are slightly more weakly negative than the "same" pairings are positive, but 1 of 5 reaches individual significance (darker brown, $p<0.05$).

While it is visually evident that the spiking responses of these three V1 example units on combined AB stimulus presentations appear drawn from a mixture of the A-like and B-like response distributions, evaluating this systematically across the population requires a formal statistical assessment.   We developed such an assessment in our previous work concerning fluctuating activity in the context of encoding of multiple simultaneous stimuli (Caruso et al., 2018; Mohl et al., 2020; Glynn et al., 2021).

Figure 4. Example units and correlation patterns. A-C. Distribution of spike counts on single stimuli (red, blue) and dual adjacent stimulus presentations (black) for three units recorded simultaneously, selected for showing a bimodal/overdispersed spike count pattern on the dual stimulus presentations ("mixtures", see main text). The units in panels A and B shared a similar stimulus preference (the red curve is shifted to the right of the blue curve for both), whereas the unit shown in C had a different stimulus preference compared to A and B (red curve is to the left of the blue curve). Dashed lines indicate the baseline activity for these units. D-F. Spike counts on individual stimulus presentations for the two units that shared a similar tuning preference exhibited positively

correlated spike count variation on individual stimulus presentations for the dual stimulus condition (green bars and Rsc value), whereas both Unit 1 and Unit 2 exhibited a negative correlation with the differently-tuned Unit 3 in c (brown bars and Rsc values). G. Distribution of Rsc values for the simultaneously-recorded units that, like Unit 1, were also classified as "mixtures"), and color coded according to whether the stimulus preference was the same (green) or different (brown). All of the "same" pairs exhibited positive correlations, and 5 of 5 were individually significant (p<0.05). All of the "different" pairs exhibited negative correlations, and 1 of 5 was individually significant (p<0.05).

In particular, we can model the firing rate behavior of neurons when two simultaneous grating stimuli A and B are presented in relation to the firing rates that occur when stimuli A and B are presented individually. We assume that each single-stimulus condition induces Poisson distributed spike counts and we exclude cases where this assumption is violated. We use a Bayesian model comparison framework to consider four hypotheses concerning the combined AB stimulus presentations (Figure 5A): 1) The responses to A and B together appear drawn from the same distribution as either A or B and consistently so on every stimulus presentation, as if the unit responded to only one of the two stimuli ('*single*'). 2) Responses to A and B together appear drawn from a distribution '*outside*' the range spanned by the A and B response distributions; this is the predicted pattern if neurons were to linearly sum the A and B responses or if one stimulus strongly suppressed the response to the other. 3) The responses to A and B together are drawn from a single distribution with a mean at an '*intermediate*' value between the A-like and B-like response rates. This is the response pattern that would be expected under theories such as divisive normalization in which the responses of an individual neuron to a more favored stimulus are reduced when other stimuli are also present, but can also represent fluctuating activity on a fast, sub-stimulus-duration timescale, as shown for some neurons in the IT face patch system and inferior colliculus (Caruso et al., 2018). 4) The responses to A and B together appear to be drawn from a '*mixture*' of the A-like and B-like response distributions. Mixtures are the category of interest for this analysis, as they indicate the presence of activity fluctuations on the stimulus presentation timescale.

We first consider the overall pattern of results for the adjacent vs. superimposed stimulus datasets in V1. Figure 5B shows the results for conditions that produced a winning model that was at least twice as likely as its nearest competitor (the full results are provided in Supplementary Figure 1). The most obvious difference between these two datasets is the dominance of "singles" in the adjacent stimulus dataset vs. "outsides" in the superimposed dataset. This suggests that the most common pattern of response in the adjacent gratings case was for V1 neurons to respond to only one of the two gratings, whereas in the superimposed gratings dataset responses were generally greater in response to the plaid than to either grating presented alone. This difference almost certainly stems from differences in the size of the stimuli being presented in these two datasets -- typically only one of the two adjacent gratings was located within the classical receptive field of a given V1 unit whereas this was not the case for the superimposed dataset. This "single" vs. "outside" difference, then, is a side note to our main focus on any possibly fluctuating activity patterns.

We found that "mixtures" were not evident in V1 when only one "object" was present, i.e. the superimposed gratings dataset (Figure 5B, rightward bars). The majority of the tested conditions were best explained by the 'outside' or 'single' hypotheses, while only about 2% were best fit by the 'mixture' hypothesis (see also Supplementary Figure 1). These results indicate that fluctuating activities are not evident in V1 when two stimuli are superimposed on each other and likely perceived as one object.

Figure 5. Bayesian model comparison regarding spike count distributions and results. *A. We evaluated the distribution of spike counts on combined stimulus presentations in relation to the distributions observed on when individual stimuli were presented alone. Four possible models were considered as described in the equations and text. Only one case each of the "single" (B-like) and "outside" ($\lambda^{AB} > max(\lambda^A, \lambda^B)$ is shown. B. Best spike count models for the adjacent and superimposed stimulus datasets, meeting a minimum winning probability of at least 0.67, i.e. the winning model is at least twice as likely as the best alternative. While "singles" dominated in the adjacent stimulus dataset and "singles" and "outsides" dominated in the superimposed stimulus datasets,*

we focus on the presence of a "mixtures" as an important minority subpopulation present nearly exclusively in the "adjacent" stimulus dataset.

However, a substantially greater (though still overall modest) number of mixtures did appear when two "objects" were present: the two adjacent stimulus dataset yielded 14-33% "mixtures" (14% overall; 33% when limiting to just cases in which the winning model was at least twice as likely as the next best alternative; Figure 5B left panel and Supplementary Figure 1). Although the proportion of conditions showing mixtures of activity patterns remained small (as expected given that V1 has comparatively small receptive fields which tends to minimize the evoked-activity overlap problem between multiple objects), it was a sufficiently large subgroup to permit analyses of spike count correlations between pairs of neurons both classified as mixtures.

Focusing now on the formally identified "mixtures" subgroup in the adjacent stimulus dataset (Figure 6A), we can see that the pattern observed for the example cells in Figure 4 holds at the population level: neural pairs in which both units responded better to the same individual stimuli tended to have positive correlations with each other (green bars, median 0.486), whereas those that had different stimulus preferences tended to exhibit negative correlations (brown bars, median -0,14). The spread of values is broad, with many of the pairs of "different-preference" neurons in particular exhibiting positive values (a point we will return to in Figure 8).

Because of the lack of an adequate population of "mixture"-classified pairs in the V1 superimposed gratings dataset, we next compared the populations as a whole (Figure 6B-C). The patterns are quite different between these two datasets. In the adjacent-stimulus dataset, the overall broad distribution and distinction between same-preference and different-preference subgroups holds even when not selecting for "mixture" fluctuating patterns (green bars vs. brown bars, median rsc 0.252, -0.052). However, this is much less true of the superimposed-gratings dataset: here, there is very little difference between the "same-preference" and "different preference" pairs of neurons (median "same-preference" rsc = 0.143, median "different-preference" rsc = 0.133), nor is there much difference between the spike count correlations observed on dual-gratings presentations vs. individual grating presentations for this dataset (orange dashed line, Figure 6C). In contrast, there is a distinct difference between spike count correlations observed on the dual-stimuli vs. individual-stimulus presentations in the adjacent-stimulus dataset (orange dashed line, Figure 6AB).

We next considered whether this overall pattern was robust to the classification categories emerging from the Bayesian model comparison, especially given that "singles" reflect the largest category in the adjacent stimulus dataset. Figure 7 illustrates the median spike count correlation by model classification category, for "same-preference" and "different-preference" pairs of neurons. We excluded the "outside" category from this analysis as there were too few units that were classified as such. We found that all 9 combinations of classifications yielded positive median spike count correlations among "same preference" pairs and negative correlations among "different preference pairs". Thus, the overall pattern of results described above does not rest critically

on the particular details of the model comparison we implemented here, and is present even among units that could not be formally shown to be fluctuating fully between "A"-like and "B"-like response distributions.

Returning to the predictions laid out in Figure 3, the implication of same-preference units being on average positively correlated and different-preference units being on average negatively correlated from a coding perspective is that V1's representation (among "mixture" units) may be slightly biased towards one or the other stimulus on each individual stimulus presentation, most closely resembling the schematic depiction in Figure 3F. However, it should be noted that the actual data involves a broad distribution with positive spike count correlations among the different-preference pairs and negative spike count correlations among the same-preference pairs, which is therefore also consistent with the schematic depiction in Figure 3H, and suggests that while the overall pattern of activity among "mixture" units is biased towards one stimulus over the other on individual stimulus presentations, there are ample cases of units that do not follow this pattern, and these exceptions may be sufficient to preserve information about the other stimulus.

Figure 6. Patterns of spike count correlations among pairs of V1 neurons in different subgroups and conditions. A. Neural pairs in which both units met the "mixture" classification showed distinct positive and negative patterns of correlation in response to adjacent stimuli. Positive correlations were more likely to occur among pairs of neurons that responded more strongly to the same individual stimuli (green bars, median rsc = 0.486), and negative correlations were more likely to occur among pairs of neurons that responded more strongly to different individual stimuli (brown bars, median rsc = -0.14, p<0.0001, see Methods). This bimodal distribution did not occur when only a single stimulus was presented (dashed orange line). B. This pattern of results held even when all the unit pairs were considered in aggregate ("same preference" pairs, median rsc = 0.252; "different preference" pairs, median rsc = -0.052, p<0.0001). C. However, among pairs recorded during presentation of superimposed gratings, this pattern was not apparent: unit pairs tended to show positive correlations in both cases ("same preference" median rsc =0.143, "different preference" median rsc = 0..133), and there was little evident difference compared to when a single grating was presented (orange line).

Figure 7. Median spike count correlations as a function of same-different preference (panel a vs. panel b) and as a function of the response profile classification resulting from the Bayesian model comparison for the adjacent stimulus dataset. The "mixture"-"mixture" combinations produced the strongest positive (same preference pairs) and strongest negative (different preference pairs) median spike count correlations, but all other combinations also involved positive median correlations for same preference pairs and negative median correlations for different preference pairs.

To visualize this in another way, we repeated the calculation of Pearson's correlations between pairs of unit-conditions classified as mixtures using not the spike counts on each stimulus presentation but an assignment score concerning how "A"-like vs "B"-like the spike count was on an individual stimulus presentation (ranging from 0 to 1; see methods). Plotted this way, a positive correlation indicates that the two units in the pair tended to exhibit response patterns consistent with the same object at the same time, whereas a negative correlation indicates that the two units tended to exhibit responses consistent with different objects at the same time. The overall pattern in the data is positively skewed (Figure 8), but with a long tail on the negative side, consistent with the population of units giving an edge to one stimulus over the other on each individual presentation, but not to the complete exclusion of the other stimulus.

We note that this correlation pattern cannot be accounted for by any obvious confounds. As mentioned previously, all stimulus presentations with microsaccades were excluded from the analyses, limiting the degree to which shared dependence on eye movements could affect the correlation patterns. Furthermore, any variability in fixation position across stimulus presentations might affect the assessment of spike count correlations within a particular pair of neurons, but would not be expected to produce (a) a bimodal distribution

of spike count correlations at the population level, that (b) only occurs when two distinct objects are presented. For example, if variability in fixation caused positive correlations between pairs of neurons whose receptive fields were aligned (likely a very small subset of our data), this effect should be present on both single stimulus presentations when the stimulus is in those receptive fields and on double stimulus presentations. Yet, as can be seen in Fig 6B, the positive extent of the correlations on double stimulus presentations among "same preference" pairs is higher than is observed on single stimulus presentations (green bars extend to higher values than the orange curve), and vice versa for the "different preference" pairs.

Figure 8. Activity fluctuations in "mixture" pairs of unit-conditions are consistent with a bias towards both units in the pair tending to signal the same stimulus at the same time. This analysis involved the Pearson's correlation coefficients computed on assignments scores (r_{as}), which take into account whether the response on combined "AB" stimulus presentations is more "A-like" vs. "B-like". For two units that share a similar preference (e.g both respond better to A or both respond better to B), this correlation will have the same sign as the spike count correlation (panel a, green points, positively sloped best fit line). For two units that prefer different stimuli, this correlation will be opposite in sign to the spike count correlation (panel a, brown points, negatively sloped best fit line). The overall positive skew in the assignment score correlations for both the "same" and "different" preferring unit-condition pairs (panel b) therefore indicates a bias for the same stimulus at the same time. The negative tail indicates the other stimulus is nevertheless also represented in a (smaller) subpopulation of neurons.

With two objects, distinct distributions of spike count correlations occur in V4

We next assessed V4, which showed both similarities and differences in comparison to V1. Like V1, activity patterns differed considerably in the superimposed vs. adjacent stimuli cases. However, the details of these differences differed: while "mixtures" were present in both the superimposed and adjacent stimulus conditions in V4, "intermediates" were more prevalent in the adjacent stimulus case than in the superimposed stimulus case. Given that "intermediates" could also reflect fluctuations (like "mixtures" but on a faster-than-stimulus-presentation time scale), we considered both mixtures and intermediates as subcategories of particular interest for the V4 dataset (Figure 9).

Figure 9. Results of the response pattern classification analysis for V4 units. Shown here are classifications for all units regardless of confidence level; see Supplementary Figure 2 for a breakdown by confidence level. "Mixtures" were seen in both datasets, but "Intermediates" were seen primarily in the adjacent-stimulus dataset. These two categories can in principle both contain fluctuating activity, and are grouped here as "between" (i.e. the average response for dual stimuli for these two categories is between the average responses to single stimuli). As with V1, the relative proportions of "singles" vs "outsides" also differed across these datasets. The

combined incidence of these "not between" categories was higher for the superimposed dataset than for the adjacent dataset.

The patterns of spike count correlations across mixture-mixture pairs in V4 varied considerably based on whether the stimuli were adjacent vs. superimposed and, for adjacent stimuli, whether the two units in the pair exhibited the same or different stimulus preferences (Figure 10). For adjacent mixture-mixture pairs (Panel A), the same-preferring units again tended to show positive spike count correlations, whereas for different-preferring pairs, the distribution appeared centered around zero. It is unclear whether these pairs are truly uncorrelated or if they might appear uncorrelated due to a negative, stimulus-related, correlation being cancelled out by a comparable, globally shared positive correlation that could stem from other factors (e.g. shared reward sensitivity). When intermediate-intermediate patterns are included, the overall pattern of a difference between the same-preferring and different-preferring distributions is preserved (panel B), although now the different-preference pairs are slightly positive. This pattern was still present when no selection for response pattern was applied (panel C), and is perhaps best appreciated by comparing the medians of the distributions (Figure 10f): there is a distinct difference between the median spike count correlation for same-preference and different-preference pairs for the adjacent dataset.

However, again like V1, when the two stimuli were presented in a superimposed fashion, this difference was no longer evident. This was the case across the whole dataset (Figure 10e-f) as well as for mixture-mixture pairs (Figure 10d,f), suggesting that the when fluctuations do occur for superimposed/bound stimuli, they likely reflect a somewhat different underlying mechanism or purpose than when distinct stimuli are presented.

Figure 10. Like V1, pairs of V4 units show different patterns of spike count correlations when there are two adjacent stimuli vs. when there is one superimposed stimulus, depending on the tuning preferences of the pair. A. Mixture-mixture pairs for adjacent stimuli (gabors or images), color coded by whether the two units in the pair shared the same or had different tuning preferences. The "same preference" and "different preference" median correlations differed (p<0.002, see Methods). B. Similar but including intermediate-intermediate pairs, since they too may be fluctuating (median difference p<0.0001). C. All unit pairs tested with adjacent stimuli, regardless of classification in the modeling analysis (median difference p<0.0001). Orange line shows the results for single stimulus presentations. D. Similar to A but for superimposed stimuli (median difference not significant). E. Similar to C but for superimposed stimuli (median difference not significant). F. Comparison of median spike count correlations in the adjacent vs. superimposed datasets, color coded by tuning preference.

Discussion

The central observation in this paper is that the patterns of spike count correlations observed in visual cortex differs based on the stimulus/stimuli that are presented and whether the two neurons in a pair share the same or different tuning preferences for those stimuli. Superimposed gratings yielded correlation patterns very similar to previous reports in the literature (Cohen and Kohn, 2011; Ruff et al., 2016; Ruff and Cohen, 2016), as well as being very similar to presentations involving single gratings, and the correlations did not greatly depend on whether the two units in the pair preferred the same individual stimulus or different ones. At the population level, there was little evidence that V1 units exhibited "mixture"-type response patterns on superimposed gratings trials, i.e. fluctuating across presentations between the responses observed in conjunction with each grating alone. In V4, such "mixture"-type response patterns were seen for superimposed gratings, but as in V1, there was little evidence of a bimodal spike count correlation pattern for these stimuli.

In contrast, when two stimuli were presented adjacent to one another other, the spike count correlations differed from the superimposed-stimulus dataset. In these datasets, two distinct distributions emerged based on whether the two units in the pair preferred the same individual stimulus (associated with generally positive spike count correlations) vs different individual stimuli (associated with generally negative spike count correlations in V1 or simply less positive spike count correlations in V4). This pattern was observed in the population as a whole, but was especially pronounced in the subset of units that exhibited "mixture"-type response patterns indicating fluctuating across stimulus presentations between the response distributions associated with each of the individual stimuli.

We interpret these observations under the conceptual framework of the challenge that visual cortex faces when representing a visual scene that contains either individual stimuli, combinations of stimuli that bind to form a single object, or combinations of stimuli that remain perceptually distinct from each other. The pattern of positive and negative (or less positive) correlations exhibited between pairs of such units is consistent with a population code biased towards one of the two stimuli on any given stimulus presentation, but that preserves information about the other stimulus as well.

There has been a rich literature concerning correlated activity between visually-responsive neurons in recent decades. One school of thought considers correlations in the context of the variability of neural firing. Under this "noise correlation" view, positive correlations have historically been seen as detrimental for encoding information at the population level (Shadlen and Newsome, 1994; Zohary et al., 1994). Such views have also seen notable refinement and qualification since these early studies (Romo et al., 2003; Averbeck and Lee, 2004; Averbeck et al., 2006; Moreno-Bote et al., 2014; Kanitscheider et al., 2015; Kohn et al., 2016). Arguably closer to the current work is a different school of thought, the temporal correlation hypothesis (Milner, 1974; Gray and Singer, 1989; Von Der Malsburg, 1994; Singer and Gray, 1995; Gray, 1999). This theory focused on the need to connect the brain's representation of different attributes of a given object together, and proposed that such binding might be mediated through precise synchrony of spikes among neurons responding to the same object. This view, then, sees correlated activity as both useful and specifically relevant to object vision. Studies exploring this hypothesis have, however, primarily focused on within-trial temporal synchrony of spikes on the order of milliseconds, whereas the noise correlation literature has focused on spike counts in the domain of hundreds of milliseconds and analysis at the level of the ensemble of trials or stimulus presentations. By evaluating spike-count variation at the level of stimulus presentations and comparing the results as a function of the number of stimuli/objects, the present work forges a bridge between these two areas of the literature.

It should be noted that the adjacent-stimulus and superimposed-stimulus datasets also differed along other dimensions besides whether the gratings were likely perceived as one object or two. In V1, the use of an

attentional task differed across these two datasets. Indeed, our results might appear consistent with the hypothesis that on the timescale of stimulus presentations (or likely fixations in natural viewing), populations of neurons shift focus from one stimulus to another when the two stimuli are perceived as separate objects (adjacent condition) but not when they are perceived as the same object (superimposed condition). Under this possibility, perhaps the animals were also switching attention between the two objects in the V1 receptive field zone (despite the attentional target being in the ipsilateral hemifield) which could contribute to the observed activity patterns (Ecker et al., 2016; Engel et al., 2016; Denfield et al., 2018). However, mildly arguing against this interpretation is that all of the analyzed trials were correctly performed, indicating that the monkey was maintaining attention in the ipsilateral hemifield on those trials. Furthermore, V4 was tested with similar stimuli but no attention task, and still showed a distinctly different correlation pattern in the adjacent vs. superimposed stimuli datasets. This suggests that performance of an attention task per se is not the determining factor in producing this pattern.

Neural responses to attended and unattended stimuli can often be modeled as a weighted combination of the responses to those stimuli when presented alone (Boynton, 2009; Lee and Maunsell, 2009; Reynolds and Heeger, 2009; Ni et al., 2012; Ni and Maunsell, 2017, 2019). Furthermore, the magnitude of the attention effects has been found to be correlated with the values of those weights, with the strongest effects of attention occurring in neurons that appeared to exhibit averaging responses (Lee and Maunsell, 2009; Ni et al., 2012; Ni and Maunsell, 2017, 2019). Such averaging responses (see also Xiao et al., 2014; Xiao and Huang, 2015) would be most consistent with the mixture and intermediate classification we used here, and may actually indicate fluctuations occurring on either the stimulus presentation or sub-stimulus presentation time scales. In future studies, it will be interesting to determine the relationship between the fluctuating responses we show here, response normalization, and attention.

Recent work suggests another more perceptual tie between studies of attention and the findings we report here. The likelihood of detecting a brief near-threshold visual stimulus varies with the phase of the brain wave oscillations at the time the stimulus is presented (Busch et al., 2009; Busch and VanRullen, 2010; Vanrullen et al., 2011; Fiebelkorn et al., 2013; Fiebelkorn et al., 2018; Helfrich et al., 2018; Fiebelkorn and Kastner, 2019) (see also Engel et al., 2016). This might reflect a perceptual consequence of a brain mechanism in which neurons are slightly biased towards representing some stimuli in the visual scene over others in a naturally-occurring oscillatory fashion. In our previous study (Caruso et al., 2018), we found that the LFP signal prior to stimulus onset was predictive of whether neurons would "pick" A vs. B on a given trial. Future work will be needed to ascertain whether a similar phenomenon occurs in V1 or V4.

Finally, it is worth noting here that considering how the brain preserves information about *two* visual stimuli presented is still a far cry from understanding how the myriad details present in a natural scene are encoded.

When the number of objects gets too great, it is unlikely that neurons can fluctuate between all of them, and this is likely to have consequences for perception, perhaps accounting for well-known limits on the number of objects we can perceive, attend to and remember (e.g. Miller, 1956; Whitney and Levi, 2011; Henry and Kohn, 2020). Future studies incorporating many stimuli and investigating how this changes the pattern of fluctuating activity and correlations between units are needed to shed light on how our brains operate outside the rarefied environment of the laboratory.

Methods

Electrophysiological recordings and visual stimuli

The full experimental procedures are described in Ruff et al.(2016) and Ruff & Cohen (2016) and summarized below. All animal procedures were approved by the Institutional Animal Care and Use Committees of the University of Pittsburgh and Carnegie Mellon University.   Each of the data sets consisted of multielectrode recordings from two adult male rhesus monkeys for each brain area (Table 1).   Recordings were made using chronically implanted a 10 x 10 microelectrode arrays (Blackrock Microsystems) in V1 and V4 (Figure 1A). The electrode shafts were 1mm long, and the minimum distance between the nearest electrodes was 400 μm. In some sessions, recordings were also made using other electrodes in areas MT and 7a, but these data are not included in the current analyses.

The visual stimuli and behavioral experiment for the superimposed stimulus dataset are fully described in Ruff et al.(2016). Monkeys were rewarded for passively viewing individual or superimposed orthogonal drifting gratings, positioned to span the receptive fields of the entire population of neurons under study (size range: 2.5-7°). Stimulus presentations with microsaccades were excluded from further analysis. In the full dataset, multiple contrast levels were presented, most of which were not included for analysis in the present study. Here, we included trials in which one grating had a contrast of 0 (i.e. was not visible) and the other had a contrast of 0.5 ("A"-alone and "B"-alone cases) or both gratings had a contrast of 0.5 ("AB"). In most sessions, each stimulus lasted for 200 ms; a few sessions with 1000 ms stimuli were also included but only the first 200 ms were analyzed.

The visual stimuli and behavioral experiment for the V1 adjacent stimulus dataset are fully described in (Ruff and Cohen, 2016). The animals performed an attentional task in which they were cued in blocks of trials to attend to small drifting Gabor patches (~1°) in various locations and respond when the orientation of the attended location changed. In the current study, we analyzed trials in which attention was directed to a Gabor patch located in the hemisphere ipsilateral to the recorded V1 neurons (i.e. well away from those neurons' receptive fields). On these trials, two unattended Gabor patches were presented in close proximity to each other within the area covered by the receptive fields of the recorded V1 neurons. These patches had the same orientation but drifted in opposite directions. The patches flashed on for 200 ms and off for 200-400 ms within the V1 receptive fields, and we

analyzed responses to all stimuli before the orientation change, excluding the first stimulus in every trial. Again, only correctly performed trials with no microsaccades during stimulus presentations were included for analysis.

The adjacent stimulus dataset for V4 involved two types of stimuli, small drifting Gabor patches as above or natural images of animals or common objects, from (Long et al., 2018). The monkeys performed a fixation task.

Analysis of spike count distributions

The full description of the statistical evaluation of spike count distributions on combined stimulus presentations can be found in (Caruso et al., 2018; Mohl et al., 2020). Briefly, we deployed a Bayesian procedure for modeling the distribution of spike counts in response to combined stimuli. Assuming that the spike counts corresponding to condition A and condition B are both Poisson-distributed with the rate parameters λ^A and λ^B respectively (and excluding exceptions, see below), the four hypotheses for the spike count distributions for condition AB consist of:

1. "**single**": A single Poisson distribution $Poi(\lambda^{AB})$, where λ^{AB} exactly equals either λ^A or λ^B.
2. "**outside**": A single Poisson distribution $Poi(\lambda^{AB})$, where λ^{AB} lies outside the range between λ^A and λ^B.
3. "**intermediate**": A single Poisson distribution $Poi(\lambda^{AB})$, where λ^{AB} lies inside the range between λ^A and λ^B.
4. "**mixture**": A mixture of the two single-stimulus distribution with an unknown mixing rate α: $\alpha\, Poi(\lambda^A) + (1-\alpha)\, Poi(\lambda^B)$

For each "triplet" or combination of A, B, and AB conditions, the hypothesis with the highest posterior probability was selected, based on the intrinsic Bayes factor(Berger and Pericchi, 1996) and using the equal prior (¼) among the four hypotheses, Jeffrey's prior(Berger and others, 2006) for the Poisson rate parameters, and a uniform prior in [0, 1] for the mixing weight α.

Only the triplets satisfying two additional criteria are used: (1) the single-stimulus distributions follow Poisson distributions, and (2) the single-stimulus rate parameters λ^A and λ^B are substantially separated. The first criterion was tested using Monte Carlo p-value calculation for a Chi-square goodness-of-fit test, and the second criterion was tested by whether the intrinsic Bayes factor of the model $\lambda^A \neq \lambda^B$ is more than three times higher than that of the model $\lambda^A = \lambda^B$. These exclusion criteria were applied to all the analyses in the paper, even those that did not build specifically on this model classification, to ensure that comparisons between subpopulations of the data were not affected by differences in data selection criteria.

Stimuli	Brain area	Task	Monkeys	Units	Sessions	Units included	Sessions included

Adjacent	V1	Attention	ST, BR	1604	16	935	16
	V4	Fixation	BA, HO	991	17	274	17
Superimposed	V1	Fixation	ST, BR	2304	25	770	23
	V4	Fixation	JD, SY	1744	21	817	21

Table 1. Summary of included data. Analyses were conducted on "triplets", consisting of a combination of A, B, and AB conditions. If the spikes evoked by the A and B stimuli failed to follow Poisson distributions with substantially separated means, the triplet was excluded from analysis. This table shows the numbers of distinct units and sessions that provided triplets for the analysis for each brain area and type of stimulus condition.

Correlation analysis

We calculated spike count correlations between pairs of units recorded at the same time in the same experiment. The Pearson correlation coefficient was calculated on the spike counts for each presentation of each relevant stimulus combination. Stimulus presentations in which one or both units in the pair exhibited an "outlier" response, i.e. more than 3 standard deviations from the mean, were excluded from the analysis. The spike count correlations for particular unit pairs for different stimulus combinations were included in the population analyses as separate observations and were not averaged together. For example, in the V1 adjacent stimulus dataset, pairs were typically tested with two separate adjacent stimulus combinations, differing in the direction of motion, potentially yielding two values of the spike count correlation (assuming both conditions passed the Poisson and response-separation exclusion criteria noted above). Similarly, V4 neurons tested with different combinations of drifting gabors and/or images contributed values of spike count correlations for each stimulus set to the population.

Same or different preference

Preference of a unit for a particular stimulus was determined by higher spike counts. Unit pairs that both exhibited more spikes in response to stimulus A than to B, or both exhibited more spikes in response to stimulus B than to A, were defined as "same preference". Unit pairs in which one responded with more spikes to A and the other with more spikes to B were defined as "different preference".

Comparison of distributions of spike count correlations

The medians of the "same preference" vs. "different preference" distributions of spike count correlations were statistically compared using Monte Carlo methods in which the same/different preference assignments were randomly shuffled and the medians recalculated 10,000 times. When the true difference between the medians was greater than any of the shuffled versions, the p value can be said to be less than 1/10,000 or 0.0001.

Author Contributions:

Experimental design: DAR, MRC

Data collection: DAR, LEK, BB

Data analysis: NYJ, STT, DAR, JMG

Manuscript preparation: NYJ, JMG

Manuscript editing: NYJ, DAR, STT, MAR, JMG

Authors LEK and BB contributed equally.

References

Adelson EH, Movshon JA (1982) Phenomenal coherence of moving gratings. Nature, Lond 300:523-525.

Alonso J-M, Chen Y (2009) Receptive field. Scholarpedia 4:5393.

Averbeck BB, Lee D (2004) Coding and transmission of information by neural ensembles. Trends Neurosci 27:225-230.

Averbeck BB, Latham PE, Pouget A (2006) Neural correlations, population coding and computation. Nat Rev Neurosci 7:358-366.

Boynton GM (2009) A framework for describing the effects of attention on visual responses. Vision Res 49:1129-1143.

Buffalo EA, Fries P, Landman R, Liang H, Desimone R (2010) A backward progression of attentional effects in the ventral stream. Proc Natl Acad Sci U S A 107:361-365.

Bulkin DA, Groh JM (2011) Systematic mapping of the monkey inferior colliculus reveals enhanced low frequency sound representation. J Neurophysiol 105:1785-1797.

Busch NA, VanRullen R (2010) Spontaneous EEG oscillations reveal periodic sampling of visual attention. Proc Natl Acad Sci U S A 107:16048-16053.

Busch NA, Dubois J, VanRullen R (2009) The phase of ongoing EEG oscillations predicts visual perception. J Neurosci 29:7869-7876.

Busse L, Wade AR, Carandini M (2009) Representation of concurrent stimuli by population activity in visual cortex. Neuron 64:931-942.

Caruso VC, Mohl JT, Glynn C, Lee J, Willett SM, Zaman A, Ebihara AF, Estrada R, Freiwald WA, Tokdar ST, Groh JM (2018) Single neurons may encode simultaneous stimuli by switching between activity patterns. Nature communications 9:2715.

Cohen MR, Kohn A (2011) Measuring and interpreting neuronal correlations. Nat Neurosci 14:811-819.

Cook EP, Maunsell JH (2002) Attentional modulation of behavioral performance and neuronal responses in middle temporal and ventral intraparietal areas of macaque monkey. J Neurosci 22:1994-2004.

Denfield GH, Ecker AS, Shinn TJ, Bethge M, Tolias AS (2018) Attentional fluctuations induce shared variability in macaque primary visual cortex. Nature communications 9:2654.

Dow BM, Snyder AZ, Vautin RG, Bauer R (1981) Magnification factor and receptive field size in foveal striate cortex of the monkey. Exp Brain Res 44:213-228.

Ecker AS, Denfield GH, Bethge M, Tolias AS (2016) On the Structure of Neuronal Population Activity under Fluctuations in Attentional State. J Neurosci 36:1775-1789.

Engel TA, Steinmetz NA, Gieselmann MA, Thiele A, Moore T, Boahen K (2016) Selective modulation of cortical state during spatial attention. Science 354:1140-1144.

Fiebelkorn IC, Kastner S (2019) A Rhythmic Theory of Attention. Trends Cogn Sci 23:87-101.

Fiebelkorn IC, Saalmann YB, Kastner S (2013) Rhythmic sampling within and between objects despite sustained attention at a cued location. Curr Biol 23:2553-2558.

Fiebelkorn IC, Pinsk MA, Kastner S (2018) A Dynamic Interplay within the Frontoparietal Network Underlies Rhythmic Spatial Attention. Neuron 99:842-853 e848.

Georgopoulos A, Schartz A, Kettner R (1986) Neuronal population coding of movement direction. Science 233:1416-1419.

Glynn C, Tokdar S, Zaman A, Caruso VC, Mohl JT, Willett SM, Groh JM (2021) Analyzing second order stochasticity of neural spiking under stimuli-bundle exposure. Annals of Applied Statistics, arXiv:191104387 15:41-63.

Gray CM (1999) The temporal correlation hypothesis of visual feature integration: still alive and well. Neuron 24:31-47, 111-125.

Gray CM, Singer W (1989) Stimulus-specific neuronal oscillations in orientation columns of cat visual cortex. Proc Natl Acad Sci U S A 86:1698-1702.

Groh JM, Kelly KA, Underhill AM (2003) A monotonic code for sound azimuth in primate inferior colliculus. Journal of Cognitive Neuroscience 15:1217-1231.

Grothe B, Pecka M, McAlpine D (2010) Mechanisms of sound localization in mammals. Physiol Rev 90:983-1012.

Heeger DJ, Simoncelli EP, Movshon JA (1996) Computational models of cortical visual processing. Proceedings of the National Academy of Sciences of the United States of America 93:623-627.

Helfrich RF, Fiebelkorn IC, Szczepanski SM, Lin JJ, Parvizi J, Knight RT, Kastner S (2018) Neural Mechanisms of Sustained Attention Are Rhythmic. Neuron 99:854-865 e855.

Henry CA, Kohn A (2020) Spatial contextual effects in primary visual cortex limit feature representation under crowding. Nature communications 11:1687.

Higgins NC, Storace DA, Escabi MA, Read HL (2010) Specialization of binaural responses in ventral auditory cortices. J Neurosci 30:14522-14532.

Kanitscheider I, Coen-Cagli R, Pouget A (2015) Origin of information-limiting noise correlations. Proc Natl Acad Sci U S A 112:E6973-6982.

Keliris GA, Li Q, Papanikolaou A, Logothetis NK, Smirnakis SM (2019) Estimating average single-neuron visual receptive field sizes by fMRI. Proceedings of the National Academy of Sciences 116:6425-6434.

Kohn A, Coen-Cagli R, Kanitscheider I, Pouget A (2016) Correlations and Neuronal Population Information. Annu Rev Neurosci 39:237-256.

Lee C, Rohrer WH, Sparks DL (1988) Population coding of saccadic eye movements by neurons in the superior colliculus. Nature 332:357-360.

Lee J, Maunsell JH (2009) A normalization model of attentional modulation of single unit responses. PLoS One 4:e4651.

Lima B, Singer W, Chen NH, Neuenschwander S (2010) Synchronization dynamics in response to plaid stimuli in monkey V1. Cereb Cortex 20:1556-1573.

Long B, Yu CP, Konkle T (2018) Mid-level visual features underlie the high-level categorical organization of the ventral stream. Proc Natl Acad Sci U S A 115:E9015-E9024.

Luck SJ, Chelazzi L, Hillyard SA, Desimone R (1997) Neural mechanisms of spatial selective attention in areas V1, V2, and V4 of macaque visual cortex. J Neurophysiol 77:24-42.

McAlpine D, Grothe B (2003) Sound localization and delay lines--do mammals fit the model? Trends Neurosci 26:347-350.

Miller GA (1956) The magical number seven, plus or minus two: some limits on our capacity for processing information. Psychological Review 63:81-97.

Milner PM (1974) A model for visual shape recognition. Psychol Rev 81:521-535.

Mohl JT, Caruso VC, Tokdar S, Groh JM (2020) Sensitivity and specificity of a Bayesian single trial analysis for time varying neural signals. Neurons, Theory, Data Analysis, and Behavior.

Moreno-Bote R, Beck J, Kanitscheider I, Pitkow X, Latham P, Pouget A (2014) Information-limiting correlations. Nat Neurosci 17:1410-1417.

Ni AM, Maunsell JHR (2017) Spatially tuned normalization explains attention modulation variance within neurons. J Neurophysiol 118:1903-1913.

Ni AM, Maunsell JHR (2019) Neuronal Effects of Spatial and Feature Attention Differ Due to Normalization. J Neurosci 39:5493-5505.

Ni AM, Ray S, Maunsell JH (2012) Tuned normalization explains the size of attention modulations. Neuron 73:803-813.

Palanca BJ, DeAngelis GC (2005) Does neuronal synchrony underlie visual feature grouping? Neuron 46:333-346.

Reynolds JH, Heeger DJ (2009) The normalization model of attention. Neuron 61:168-185.

Rodman HR, Albright TD (1989) Single-unit analysis of pattern-motion selective properties in the middle temporal visual area (MT). Exp Brain Res 75:53-64.

Romo R, Hernandez A, Zainos A, Salinas E (2003) Correlated neuronal discharges that increase coding efficiency during perceptual discrimination. Neuron 38:649-657.

Ruff DA, Cohen MR (2016) Attention Increases Spike Count Correlations between Visual Cortical Areas. J Neurosci 36:7523-7534.

Ruff DA, Alberts JJ, Cohen MR (2016) Relating normalization to neuronal populations across cortical areas. J Neurophysiol 116:1375-1386.

Shadlen M, Newsome W (1994) Noise, neural codes and cortical organization. **Current Opinion in Neurobiology** 4:569-579.

Singer W, Gray CM (1995) Visual feature integration and the temporal correlation hypothesis. Annu Rev Neurosci 18:555-586.

Vanrullen R, Busch NA, Drewes J, Dubois J (2011) Ongoing EEG Phase as a Trial-by-Trial Predictor of Perceptual and Attentional Variability. Front Psychol 2:60.

Von Der Malsburg C (1994) The correlation theory of brain function. In: Models of neural networks, pp 95-119: Springer.

Werner-Reiss U, Groh JM (2008) A rate code for sound azimuth in monkey auditory cortex: implications for human neuroimaging studies. J Neurosci 28:3747-3758.

Whitney D, Levi DM (2011) Visual crowding: a fundamental limit on conscious perception and object recognition. Trends Cogn Sci 15:160-168.

Woods TM, Su TK, Recanzone GH (2001) Spatial tuning as a function of stimulus intensity of single neurons in awake macaque monkey auditory cortex. Soc Neurosci Abstr 27:512.517.

Xiao J, Huang X (2015) Distributed and Dynamic Neural Encoding of Multiple Motion Directions of Transparently Moving Stimuli in Cortical Area MT. J Neurosci 35:16180-16198.

Xiao J, Niu YQ, Wiesner S, Huang X (2014) Normalization of neuronal responses in cortical area MT across signal strengths and motion directions. J Neurophysiol 112:1291-1306.

Zohary E, Shadlen MN, Newsome WT (1994) Correlated neuronal discharge rate and its implications for psychophysical performance. Nature 370:140-143.