Reversible Reorganization Of Basolateral Amygdala Dendrites in Male Rats Accompanies

Stress Resilience

Sheldon D. Michaelson[1], Ana Pamela Miranda Tapia[1], Amanda McKinty[1], Heika Silveira

Villarroel[1], James P. Mackay[1], Janice H. Urban[2], William F. Colmers[1,3]*

[1] Department of Pharmacology, and the Neuroscience and Mental Health Institute, University of

Alberta, Edmonton, Alberta, T6G 2H7, Canada

[2] Center for the Neurobiology of Stress Resilience and Psychiatic Disorders; Discipline of

Physiology and Biophysics, Chicago Medical School, Rosalind Franklin University of Medicine

and Science, North Chicago, IL, 60064 USA

[3] Lead contact.

Corresponding Author: Dr. William F. Colmers, Department of Pharmacology and Institute for

Neuroscience and Mental Health, University of Alberta, 9-36 MSB, Edmonton, AB. T6G 2H7,

Canada. wcolmers@ualberta.ca

Number of Figures: 13

Number of Tables :   2

Number of words in Abstract: 147

ABSTRACT
23
Basolateral amygdala (BLA) responses to stress are regulated by endogenous neuropeptide

Y (NPY) and corticotrophin-releasing factor (CRF), associated respectively with resilience and

vulnerability to stress. We studied mechanisms underlying effects of repeated NPY and CRF

treatment on structure and function of BLA principal neurons (PN) in organotypic slice cultures

(OTC) of rat BLA. NPY attenuated excitatory input and induced dendritic hypotrophy via Y_5-

receptors, while CRF increased excitatory input and induced hypertrophy of BLA PNs. Structural

responses to NPY and CRF required calcineurin and CaMKII, respectively. Finally, repeated *in*

vivo treatment of BLA with NPY or a Y_5 receptor agonist increased social interaction and

recapitulated structural changes in BLA neurons seen in OTCs. These results implicate the Y_5

receptor in the long-term, anxiolytic-like effects of NPY in the BLA. Moreover, BLA OTCs offer

a robust model to study mechanisms associated with resilience and vulnerability to stress in BLA.

Keywords: Organotypic slice cultures; Basolateral amygdala; Stress resilience; Neuropeptide Y;

CRF, dendritic plasticity, Social interaction

INTRODUCTION

Acute stress elicits a coordinated, adaptive response to real or perceived threats, which is soon terminated via neural and endocrine systems that buffer the stress and restore homeostasis. While the stress response is essential to survival, persistent or extreme stress can disrupt this balance, predisposing the individual to maladaptive, affective- and anxiety-related disorders (McEwan, 2003). Stress-related signals are processed in part through the complex network of inhibitory and excitatory neurons in the amygdala, particularly the basolateral amygdala (BLA), which receives input related to stress, emotion and memory (Phelps and Ledoux, 2005). BLA projections, in turn, control and modulate anxiety-, reward- and emotionally-driven behaviors (McDonald, 1985, McDonald et al., 1989). Extreme or prolonged stress can overwhelm this homeostatic control, resulting in an imbalance between the glutamatergic excitatory and GABAergic inhibitory control of BLA activity, and enhancing not only the anxiogenic output of BLA projection neurons (Roozendaal et al., 2009), but causing hypertrophy of their dendritic arbors, accompanied by increased numbers of excitatory synaptic spines (Mitra et al., 2005, Padival et al., 2013, Adamec et al., 2012, Hill et al., 2011, Vyas et al., 2002, 2006). Such changes can persist for weeks (Vyas et al., 2004).

In addition to GABA and glutamate, the neuropeptides corticotropin-releasing factor (CRF) and Neuropeptide Y (NPY) provide regulation of the BLA, with CRF being excitatory and NPY inhibitory to BLA activity. While CRF is associated with the generation of stress responses (Rainnie et al., 2004; Sajdyk et al., 2004) NPY is an important endogenous anxiolytic compound in rodents (Sajdyk et al., 2002, 2004) and in humans where NPY is also linked to resilience (Yehuda et al., 2006; Sajdyk et al., 2008; Zhou et al., 2008). These behavioral effects of NPY are linked to some of its actions within the amygdala, as administration of NPY decreases the activity of BLA PNs

through activation of NPY Y_1 receptors (Y_1R), which acutely hyperpolarizes and inhibits BLA PNs

through suppression of somatodendritic I_h (Giesbrecht et al., 2010). Moreover, NPY acutely

decreases BLA activity by enhancing $GABA_A$-mediated inhibitory postsynaptic currents (IPSCs),

and reducing NMDA-mediated excitatory postsynaptic currents (EPSCs) (Molosh et al., 2013) in

BLA PNs.

Together a balance between CRF and NPY signaling within the BLA is important for

various behavioral responses (Sajdyk et al., 2004; Heilig 1994; 2004; Gutman et al., 2008). When

subjected to a stressor, CRF is released to generate an appropriate behavioral stress response (e.g.,

freezing, fleeing), these neural and behavioral responses are in turn buffered by NPY release which

temporally occurs after the stressor to curtail the actions of CRF and limit the duration of the stress

response (Heilig et al., 1994). A predominance of either NPY or CRF tone can tip this balance and

induce anxiolysis and anxiogenesis, respectively (Sajdyk et al., 2004, Heilig et al., 1994). This

balance is maintained in part through differential modulation of the tonically-active H-current (I_h):

CRF acutely depolarizes and excites BLA PNs by enhancing the (I_h), whereas NPY suppresses I_h

in the same neurons (Giesbrecht et al, 2010). Subjecting animals to either repeated stress (Padival

et al 2013) or daily injections of urocortin (UCN- a CRF-R1 and CRF-R2 agonist) induces anxiety

that long outlasts the stimulus; conversely, generation of a persistent stress resilience is generated

by a similar treatment with NPY (Sajdyk et al,, 2008, Rainnie et al., 2004). In all, these cellular

responses likely underlie the ability of NPY to mitigate CRF- and stress-induced, BLA-mediated

behaviors (Sajdyk et al., 2006; Rainnie et al., 2004).

As repeated injections of CRF produce stress vulnerability (increased sensitivity to stressful

situations; Rainnie et al., 2004), repeated application of NPY induces a persistent stress resilience

that lasts for at least two months (Sajdyk et al., 2008). Resilience is defined as the ability to rebound

from trauma or severe stress and is associated with positive mental health outcomes (Yehuda et al.,

2006). Using these models provides a unique opportunity to identify and exploit molecular and

cellular mechanisms associated with neuroplasticity in the BLA associated with stress-related

behavioral outcomes. We recently showed that a reduction in I_h in BLA PNs contributes to, and

mimics, this long-term resilience (Silveira Villarroel et al., 2018). Beyond this however, the cellular

mechanisms mediating long-term NPY and CRF effects on BLA neurons, which drive adaptive and

maladaptive coping strategies to stress exposure are poorly understood, in part because acute brain

slice preparations, while useful, are short-lived. To get around this issue, organotypic slice cultures

(OTCs) derived from hippocampus, spinal cord and cerebellum have been shown to reliably retain

gross tissue organization, properties of synaptic transmission, neuronal morphology and

neurochemical phenotypes (De Simoni et al., 2003, Avossa et al., 2003, Lu et al., 2009, Gähwiler,

1981) permitting comparison and validation relative to acute slice preparations (De Simoni et al.,

2003, Gähwiler, 1981, Stoppini et al., 1991). Moreover, OTCs permit the rigorous study of chronic

treatments and long-term effects of acute manipulations in a well-defined system (Humpel, 2015).

Here we report OTC preparations containing the BLA, a novel *in vitro* model in which to study

stress-related circuitry and more clearly understand the interactions of NPY and CRF and the

mechanisms underlying their persistent regulation of BLA PN output.

We found that repeated incubations of BLA-OTCs with NPY, designed to mimic *in vivo*

NPY injection protocols that induce stress resilience (Sajdyk et al., 2008, Silveira Villarroel et al.,

2018), reduced synaptic excitation and increased inhibition of PNs and induced a persistent, marked

hypotrophy of their dendritic trees. Structural effects of NPY were mediated entirely by NPY_5

receptors (Y_5R), and were calcineurin-dependent. Similar incubations with CRF resulted in effects

opposite to those of NPY that were mediated by CaMKII; treatment with either peptide reversed or

prevented the effects of the other. Finally, repeated *in vivo* injections of NPY or of Y_5R-selective

agonists into the BLA not only induced an enduring state of behavioral stress resilience, but also

replicated the dendritic remodeling seen in BLA OTCs, an action mediated entirely by the Y_5R.

These studies indicate a novel structural correlate of persistent stress resilience, and highlight the

use of the BLA OTC model in identifying mechanisms of adaptive and maladaptive coping

strategies to stress and anxiety-like behavior.

RESULTS

Repeated NPY and CRF treatments cause dendritic hypotrophy and hypertrophy,

respectively, in OTC PNs

Repeated restraint stress induces lasting dendritic hypertrophy in BLA neurons (Padival et

al., 2013). Whereas *in vivo* administration of CRF into the BLA elicits long-term stress

vulnerability, NPY administration persistently decreases the animals' vulnerability to stress (Sajdyk

et al., 2008, Rainnie et al., 2004, Silveira Villarroel et al., 2018). The persistence of NPY's effects

may parallel the persistence of the stress-induced dendritic hypertrophy in BLA neurons (Padival

et al., 2013), but mechanisms underlying the effects of either endogenous neuropeptide remain

unclear. To test our hypothesis that repeated application of either NPY or CRF to BLA OTCs will

alter electrophysiological properties of PNs and result in their respective hypotrophy or

hypertrophy, we monitored PN properties with electrophysiology and post-recording

morphological analysis. Initial validation of the OTC preparation of BLA (Material and Methods)

included an extensive comparision of PN neuronal properties with neurons from acute *ex vivo* brain

slices of P14 and P70 animals **(Table 1)**; the PNs from OTCs were more compact but electrically

more similar to those from acute slices from the adult animals.

Repeated application of NPY (1 – 100 nM) reduced PN whole-cell capacitance relative to

vehicle controls (**Fig. 1a**). Consistent with this, NPY ($\geq$ 1 nM) reduced and simplified the dendritic

arbor (**Fig. 1b-j**), decreasing the total dendritic length (**Fig. 1b**) and complexity of dendritic

branching (**Fig. 1c,d**). NPY's effects on dendritic morphology were already significant at 1 nM (p

<0.0001 vs. control). Although dendritic spine density was not altered by 100 nM NPY, estimates

of total spine numbers showed a trend toward reduction compared to controls (**Fig. 2**), consistent

with the significant reduction in EPSC frequencies observed (**Fig. 3b,c,e**). Neurons treated with the

inactive, desamido NPY analogue (daNPY – 100 nM) (Wahlestedt et al., 1986), did not differ in

any aspect from untreated controls (**Fig. 1a-f**).

Figure 1. NPY treatment of BLA OTC causes pyramidal neuron hypotrophy.   a. Scatter plot of
mean capacitance of OTC pyramidal neurons treated with vehicle ("Control" n=42), NPY at 100
pM (n=31), 1 nM (n=43), 10 nM (n=47), 100 nM (n=49) and the inactive analogue, desamido-NPY
at 100 nM ("da" n=30) ($H_{(5)}$ = 40.72, *p* =$1.07e^{-7}$). ***Inset*** Representative capacitative transients for
control- or 100 nM NPY-treated BLA OTC neurons. **b**. Scatter plot of mean total dendritic length
for OTC neurons treated as in **a**. Vehicle (n=26), NPY at 100 pM (n=26), 1 nM (n=25), 10 nM (n=21),
100 nM (n=24) and da-NPY 100 nM (n=20) ($H_{(5)}$ = 19.04, *p* = 0.0019). **c.** Scatter plot of mean number
of branch points in neurons in **b** ($H_{(5)}$ = 17.02, *p* = 0.0045). **d**. Sholl analysis of neurons in **b** and **c**
treated as indicated (<u>Treatment:</u> $F_{(2, 72)}$ = 9.96, *p* = 0.00015; <u>Distance:</u> $F_{(40, 2880)}$ = 185.4, *p* ~ 0;
<u>Interaction:</u> $F_{(80, 2880)}$ = 2.09, *p* =$8.57e^{-8}$). **e-j.** Representative reconstructions of neurons treated as
indicated above. **k.** Comparative timelines for experiments with BLA OTCs (upper timeline) and *in*
vivo (lower timeline) models. N-values for **c** and **d** are as in **b**. For **a-c**, Kruskal-Wallis H tests with
Dunn's post hoc tests. For **d**, 2-way RM ANOVA with Tukey's post-hoc test. For **a-c**, circles
represent neurons, black bars represent population means, while in **d**, symbols represent
population means and for all, error bars indicate s.e.m. All tests were two-sided. **P<0.05*,
**P<0.01, ****P<0.001*, *****P<0.0001* for post hoc multiple comparisons. Scale bars in e-j = 100
µm.

Figure 2. Treatment with NPY, CRF or the Y5R agonist, cPP, affects the estimated total number of spines per cell, but not spine density of principal neurons in BLA OTCs. a – c.. Representative images of spines on pyramidal neurons treated with vehicle control, 100 nM NPY, 100 nM cPP and 30 nM CRF, as indicated. **d**. Scatter plot of mean spine density of pyramidal neurons treated as in **a**. vehicle control (n=15), 100 nM NPY (n=15), 100 nM cPP (n=15) and 30 nM CRF (n=15); 1-way ANOVA with Dunnett's multiple comparisons test: $F_{(3, 56)}$ = 1.437, p = 0.24. **e**. Scatter plot of mean estimated total number of spines for neurons in **d** (One-way ANOVA with Welch's correction and Dunnett's T3 multiple comparisons test $W_{(3, 29.15)}$ = 10.61, p = 7.09e^{-5}). For **d,e,** circles represent neuron means, black bars represent population means and for all, error bars indicate s.e.m. All tests were two-sided. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 3. **Changes in principal neuron synaptic properties with repeated NPY or CRF treatment in BLA OTCs. a – b** Representative electrophysiological recordings of spontaneous synaptic currents in BLA OTC principal neurons treated repeatedly with vehicle (**a**) and (**b**) 100 nM NPY, **c.** Mean sIPSC amplitudes in neurons in vehicle control (n=25) or 100 nM NPY (n=25) (Mann-Whitney U test; U = 171.0, *p* = 0.0054). **d**. Scatter plots of mean sEPSC interevent interval from recordings of cells in **c.** Vehicle control (n=25) or 100 nM NPY (n=25) (Mann-Whitney U test; U = 155.5, *p* = 0.0019) **e – f** Representative electrophysiological recordings of neurons treated repeatedly with vehicle or with 30 nM CRF **g**. Mean sEPSC interevent interval in vehicle control (n=25) or 100 nM 30 nM CRF (n=25). (Mann-Whitney U test; U = 194, *p* = 0.0208). For **c,d, g**, circles represent neuron means, black bars represent population means and for all, error bars indicate s.e.m. All statistical tests were two-sided. **P<0.01.

Repeated treatment of OTCs with CRF induced a diametrically opposite set of changes in

BLA PN physiology and structure. Thus, application of 30 nM CRF increased sEPSC frequency

compared to vehicle-treated PNs (**Fig. 3 f-h**). Furthermore, compared with controls, CRF increased

whole-cell capacitance (**Fig. 4a**) and caused hypertrophy of the dendritic arbor (**Fig. 4b**), including

increased branching (**Fig. 4c**) and dendritic complexity (**Fig. 4d-g**). The effects of CRF on dendritic

morphology occurred in a more clearly concentration-dependent manner than did those of NPY.

Although CRF-treatment did not alter spine density, estimates of total spine numbers were

significantly increased compared to controls (**Fig. 2**).

Figure 4. CRF treatment of BLA OTC causes pyramidal neuron hypertrophy. a. Scatter plot of
mean capacitance of OTC pyramidal neurons treated with vehicle control (n=39) and CRF at 300
pM (n=40), 3 nM (n=36) and 30 nM (n=38) ($H_{(3)}$ = 22.67, p = 4.72e^{-5}). *Inset*. Representative
capacitance transients from control- or 30 nM CRF-treated BLA OTC neurons. **b.** Scatter plot of
mean total dendritic length for OTC neurons treated as in **a**. Vehicle control (n=33) and CRF at 300
pM (n=31), 3 nM (n=26) and 30 nM (n=28) ($H_{(3)}$ = 22.71, p =4.63e^{-5}). **c**. Scatter plot of mean number
of branch points in neurons in **b** ($H_{(3)}$ = 19.90, p = 0.0002). **d**. Sholl analysis of neurons in **b** and **c**
treated as indicated (<u>Treatment:</u> $F_{(2, 84)}$ = 11.43, p = 4.05e^{-5}; <u>Distance:</u> $F_{(40, 3360)}$ = 137.4, p ~ 0;
<u>Interaction:</u> $F_{(80, 3360)}$ = 2.58, p = 1.45e^{-12}). **e-g**. Representative reconstructions of neurons treated

as indicated above. For **a-c**, Kruskal-Wallis H tests with Dunn's post hoc tests. For **d**, 2-way RM ANOVA with Tukey's post-hoc test. For **a-c**, circles represent neurons, black bars represent population means, while in **d**, symbols represent population means and for all, error bars indicate s.e.m. All tests were two-sided. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$ for post hoc multiple comparisons. Scale bars in e-g = 100 µm

Bi-directional PN dendritic remodeling after successive treatments with NPY and CRF in BLA OTCs

BLA output, and thus the physiological and behavioral responses to stress are governed in part by the opposing actions of NPY and CRF neuropeptide systems. Disruption of this homeostatic mechanism may in part underlie some persistent anxiety-related disorders (Schmeltzer et al., 2016). We therefore next determined if NPY and CRF can also act as counter-regulatory signals in BLA OTCs.

BLA OTCs were incubated for one week with either CRF or NPY, then some were incubated the following week respectively with either NPY or CRF, and maintained in control media for one additional week (**Fig. 5a**) before electrophysiological and morphological analyses were performed on both single- and double-treated populations. Specifically, when PNs were reciprocally treated with the two peptides in either order, there were no differences either in cell capacitance (**Fig. 5a**) or dendritic size or complexity (**Fig. 5b-g**) from vehicle controls. As in the previous experiments, application of NPY or CRF alone either decreased or increased the frequency of sEPSCs observed in the OTCs, whereas sEPSC frequency in the CRF,NPY- or NPY,CRF-treated groups was not significantly different from vehicle controls (**Fig. 5h-k**). Assessment of BLA PN morphology in the CRF,NPY and NPY,CRF groups was consistent with the neurons undergoing a bi-directional structural plasticity.

Figure 5. Sequential treatment with NPY or CRF reverses or prevents the effects of each neuropeptide alone on BLA OTC pyramidal neuron morphology. a. Scatter plot of mean capacitance of OTC pyramidal neurons treated with media control (n=42), 30 nM CRF followed by 100 nM NPY (n=31) and 100 nM NPY followed by 30 nM CRF (n=32), with the mean capacitance for 30 nM CRF and 100 nM NPY alone (data from figs 1 and 2) indicated by dashed lines ($H_{(2)}$ = 0.73, p = 0.69). **_Inset_** Representative capacitance transients from control, CRF, NPY- and NPY, CRF-treated BLA OTC neurons. **b**. Scatter plot of mean total dendritic length for OTC neurons treated as in **a**. with control (n=26), CRF and NPY (n=23) and NPY and CRF (n=23) with the mean total length for 30 nM CRF and 100 nM NPY alone from Figs 1 and 2 as dashed lines ($H_{(2)}$ = 0.80, p =

0.67). **c.** Scatter plot of mean number of branch points in neurons in **b** with the mean values for
30 nM CRF and 100 nM NPY alone from Figs 1 and 2 as dashed lines ($H_{(2)}$ = 0.36, p = 0.84). **d.** Sholl
analysis of neurons in **b** and **c** with the Sholl analysis for 30 nM CRF and 100 nM NPY from Figs 1
and 2 as dashed lines (<u>Treatment</u>: $F_{(2, 69)}$ = 0.21, p = 0.81; <u>Distance</u>: $F_{(40, 2760)}$ = 157.3, $p \sim$ 0;
<u>Interaction</u>: $F_{(80, 2760)}$ = 0.70, p =0.98). **e-g.** Representative traces from neurons treated as
indicated.. **h.** Mean sEPSC interevent interval in neurons treated with 30 nM CRF followed by 100
nM NPY (n=25), 100 nM NPY followed by 30 nM CRF (n=25), 30 nM CRF alone (n=25) and 100 nM
NPY alone (n=25) ($H_{(3)}$ = 28.18, p = 3.33e^{-6}). **i-k** representative recordings from neurons treated as
indicated. Statistical tests: For **a-c**, Kruskal-Wallis H tests with Dunn's post hoc tests. For **d**, 2-way
RM ANOVA with Tukey's post-hoc tests. For **h**, Kruskal-Wallis H test with Dunn's post hoc test. For
a-c, circles represent individual neurons, black bars represent population means, while in **d**,
symbols represent population means and for all, error bars indicate s.e.m. All statistical tests were
two-sided. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. Scale bars in e-j = 100 μm.

The Y_5R but not the Y_1R mediates dendritic hypotrophy in BLA OTCs

While NPY plays a key role in behavioral stress resilience, the receptor subtype(s) mediating

the long-term effect on behavior is unknown. The BLA expresses Y_1, Y_2 and Y_5 NPY receptors

(Kopp et al., 2002; Wolak et al., 2003; Stanic et al., 2011) and while the acute anxiolytic effects of

NPY in the BLA are mediated predominantly via the Y_1R (Sajdyk et al., 2002a), evidence also

suggests a role for the Y_5R (Sajdyk et al., 2002a). Unexpectedly, selective activation of Y_2

receptors in the BLA acutely increases PN excitability (Mackay et al., 2019) and facilitates the

anomalous expression of anxiety-like behaviors (Sajdyk et al., 2002b). We thus examined the roles

of different NPY receptors in a pharmacological experiment using BLA OTCs.

In a repeated incubation experiment as with NPY above, the Y_1R-selective agonist, F^7,P^{34}-

NPY (1–100 nM) had no measurable long-lasting effects on the properties of BLA OTC PNs

compared to controls at any concentration tested. Thus, whole-cell capacitance (**Fig. 6a**), sEPSC

frequency (**Fig. 7 a-c**), or any of the morphological parameters affected by NPY itself in this system

were unchanged (**Fig. 6b-d**). The Y_2R-selective agonist, [ahx$^{5-24}$]NPY^{36} (1 – 100 nM; Beck-

Sickinger et al, 1992), had only minor effects on the same parameters at lower concentrations, but

at higher concentrations (100 nM) we observed a significant increase, relative to vehicle, both in

whole-cell capacitance and in dendritic arborization (total dendritic length and number of branches;

Fig. 8a-g) which was similar to that seen with CRF. These results are consistent with previously

reported anxiogenic actions of Y_2 receptor agonists (Sajdyk et al, 2002a,b), and indicate the Y_2

receptor is unlikely to mediate NPY-induced hypotrophy.

Figure 6. Treatment with the selective Y_1 receptor agonist F^7P^{34}-NPY does not affect BLA OTC
pyramidal neuron morphology. a. Scatter plot of mean capacitance of OTC pyramidal neurons
treated with vehicle control (n=30) and F^7P^{34}-NPY at 1 nM (n=28), 10 nM (n=34) and 100 nM (n=34)
($H_{(3)}$ = 0.58, p = 0.90). ***Inset*** Representative capacitance transients from Control- and F^7P^{34} NPY-
treated BLA OTC neurons. **b**. Scatter plot of mean total dendritic length for neurons treated as in
a. vehicle control (n=27) and F^7P^{34}-NPY at 1 nM (n=19), 10 nM (n=18) and 100 nM (n=22) ($H_{(3)}$ =
1.04, p = 0.79). **c**. Scatter plot of mean number of branch points for neurons in **b** ($H_{(3)}$ = 4.25, p =
0.24). **d**. Sholl analysis for neurons in **b** and **c** treated as indicated (Treatment: $F_{(1, 48)}$ = 1.82, p =
0.18; Distance: $F_{(40, 1920)}$ = 109.3, p ~ 0; Interaction: $F_{(40, 1920)}$ = 0.33, p =0.99). For **a-c**, Kruskal-Wallis

H tests with Dunn's post hoc tests. For **d**, 2-way RM ANOVA with Tukey's post-hoc test. For **a-c**,
circles represent neurons, black bars represent population means, while in **d**, symbols represent
population means and for all, error bars indicate s.e.m. All tests were two-sided.

284
285

Figure 7. Treatment with the Y_1R agonist F^7P^{34}-NPY does not affect the sEPSC frequency, while the Y_5 agonist cPP reduces sEPSC frequency onto BLA OTC pyramidal neurons. **a-b,d-e.** Representative electrophysiological recordings of spontaneous synaptic currents in BLA OTC pyramidal neurons treated with (**a,d**) vehicle, (**b**) F^7P^{34}- NPYand (**e**) cPP. **c.** Scatter plot of mean interevent sEPSC intervals onto pyramidal neurons treated with vehicle (n=25) and 100 nM F^7P^{34}-NPY (n=25) (Mann-Whitney U test; U = 299.0, p = 0.80). **f.** Scatter plot of mean inter-event interval of sIPSC (t-test with Welch's correction: t($_{36.20}$) = 2.92, p = 0.0060) and sEPSCs (Mann-Whitney U test; U = 111.0, p <0.0001) from vehicle control (n=25) and cPP-treated neurons (n=25). **g.** Scatter

 plot of mean amplitudes of sIPSC (Mann-Whitney U test; U = 259.0, *p* =0.30) and sEPSCs (Mann-
Whitney U test; U = 269.0, *p* =0.40) from vehicle control (n=25) and cPP (100 nM)-treated (n=25)
neurons. For **c,f-g**, circles represent neuron means, black bars represent population means and
for all, error bars indicate s.e.m. All statistical tests were two-sided. ns = P>0.05,*P<0.05,
**P<0.01, ***P<0.001, ****P<0.0001.

Figure 8. Treatment of BLA OTCs with the Y₂R agonist ahx causes pyramidal neuron hypertrophy. **a**. Scatter plot of mean capacitance of OTC pyramidal neurons treated with vehicle (n=26), ahx at 1 nM (n=23), 10 nM (n=21), and 100 nM (n=25) ($H_{(3)}$ = 11.31, *p* = 0.010). ***Inset*** Representative capacitative transients for control- or 10nM ahx-treated BLA OTC neurons. **b**. Scatter plot of mean total dendritic length for OTC neurons treated as in **a** ($H_{(3)}$ = 13.25, *p* = 0.0041). Vehicle (n=23), ahx at 1 nM (n=19), 10 nM (n=23), and 100 nM (n=23). **c**. Scatter plot of mean number of branch points in neurons in **b** ($H_{(3)}$ = 17.94, *p* = 0.0005). **d**. Sholl analysis of neurons in **b** and **c** treated as indicated (Treatment: $F_{(2, 66)}$ = 4.06, *p* = 0.022; Distance: $F_{(40, 2640)}$ = 119.5, *p* ~ 0;

Interaction: $F_{(80, 2640)}$ = 2.27, p = 1.92e^{-9}; Purple = control vs 10 nM ahx; Black = control vs 100 nM
ahx). **e-g.** Representative reconstructions of neurons treated as indicated above. N-values for **c**
and **d** are as in **b**. For **a-c**, Kruskal-Wallis H tests with Dunn's post hoc tests. For **d**, 2-way RM ANOVA
with Tukey's post-hoc test. For **a-c**, circles represent neurons, black bars represent population
means, while in **d**, symbols represent population means and for all, error bars indicate s.e.m. All
tests were two-sided. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$ for post hoc multiple
comparisons. Scale bars in e-g = 100 μm

Finally, we tested the Y$_5$R-selective agonist, [cPP$^{1-7}$,NPY$^{19-23}$,Aib^{32},Gln^{34}] hPP (cPP)

(Cabrele et al., 2001) (1–100 nM). At 100 nM, cPP caused significant reductions in PN sEPSC

frequency and increased sIPSC frequency compared to controls, with no changes in sIPSC or sEPSC

amplitude (**Fig. 7d-g**). Concentrations of cPP ≥ 10 nM also caused robust decreases in PN whole-

cell capacitance (**Fig. 9a**), total dendritic length, and numbers of branch points (**Fig. 9b-g**). While

spine density was not altered by 100 nM cPP, estimates of total spine numbers per neuron were

significantly reduced when compared with vehicle controls (**Fig. 2**). Furthermore, co-incubation of

the Y$_5$R antagonist, CGP71683A (CGP, 30 nM) with a maximally effective concentration (30 nM)

of NPY not only prevented NPY effects on whole-cell capacitance and dendritic extent, but actually

caused dendritic hypertrophy (**Fig. 9a-g**). This effect is consistent with the Y$_5$ antagonist unmasking

the Y$_2$-mediated actions of NPY reported above.

Figure 9. Treatment of BLA OTC with the Y_5R agonist cPP results in pyramidal neuron hypotrophy, which is blocked by the Y_5R antagonist CGP71683A (Y_5A). a. Scatter plot of mean capacitance of OTC pyramidal neurons treated with vehicle control (n=30), cPP at 1 nM (n=32), 10 nM (n=47) and 100 nM and 10 nM NPY with 30 nM CGP (n=45) ($H_{(4)} = 33.61$, $p = 8.97e^{-7}$). ***Inset*** Representative capacitance transients from Control-, cPP- and NPY + Y_5Antagonist-treated BLA OTC neurons. **b**. Scatter plot of mean total dendritic length of neurons treated as in **a**. vehicle control (n=26), cPP at 1 nM (n=23), 10 nM (n=31) and 100 nM (n=32) and 10 nM NPY with 30 nM CGP (n=34) ($H_{(4)} = 49.88$, $p = 3.83e^{-10}$). **c.** Mean number of branch points in neurons in **b** ($H_{(4)} = 36.99$, $p = 1.81e^{-7}$). **d.** Sholl analysis for neurons in **b** and **c** treated as indicated (Treatment: $F_{(2, 90)} = 23.98$, $p = 4.49e^{-9}$; Distance: $F_{(40, 3600)} = 175.1$, $p \sim 0$; Interaction: $F_{(80, 3600)} = 3.91$, $p = 2.68e^{-28}$). Black = control vs. 100 nM cPP; Green = control vs. NPY + Y_5A; Blue = 100 nM cPP vs. NPY + Y_5A. **e-g**. Representative traces from neurons treated as indicated. For **a-c**, Kruskal-Wallis H tests with Dunn's post hoc tests. For **d**, 2-way RM ANOVA with Tukey's post-hoc test. For **a-c**, circles represent neurons, black bars represent population means, while in **d**, symbols represent

In vivo treatment of BLA with NPY or cPP results both in long-term behavioral stress resilience and dendritic hypotrophy

Based on the *in vitro* results, we hypothesized that NPY will cause similar effects on PN

structure in BLA *in vivo* that would also correlate with persistent increases in social interaction, a

measure of anxiety (File and Seth, 2003, Silveira et al, 2018), which permits repeated longitudinal

measures in individual animals, unlike many other paradigms such as the elevated plus maze. Five

sequential daily injections either of NPY (10 pmol/100 nL), the Y_1-agonist F^7P^{34}NPY (10 pmol/100

nL), the Y_5-agonist, cPP (10 pmol/100 nL), or NPY co-applied with the Y_5-antagonist CGP (each

10 pmol/100 nL) were given into bilateral BLA of 8 week-old male rats (Sajdyk et al., 2008). All

NPY agonist treatments in this experiment acutely increased social interaction (SI) on injection

days one and five relative both to pre-treatment baseline and to vehicle controls (**Fig. 10a**).

However, only animals treated with NPY or the Y_5-agonist retained this increase in SI at 2 and 4

weeks (**Fig. 10a**). In recordings from PNs in acute *ex vivo* BLA slices from these same animals at

4 weeks post-treatment, prior treatment with either NPY or the Y_5R agonist decreased whole-cell

capacitance (**Fig. 10b)** and significantly reduced dendritic extent and complexity relative to PNs

from the vehicle-treated animals (**Fig. 10c-j**). Co-administration of the Y_5R antagonist with NPY

prevented the long-term effects of NPY both on behavior and PN morphology (**Fig. 10a-e, j**); no

anxiogenic behavior was observed at any timepoint, nor were increases in PN capacitance or

arborization as seen in the OTCs. Therefore, the Y_5R is both necessary and sufficient to mediate

both the NPY-induced long term structural plasticity and decreased behavioral stress responses *in*

vivo.

Figure 10. Repeated *in vivo* injection of NPY or the Y_5 receptor agonist, but not the Y_1 receptor
agonist results in increased social interaction times and dendritic hypotrophy. **a**. Scatter plot
depicting social interaction time shown as percent of individual animals' baselines for treatment
with vehicle control (100 nL) at day 1 (n=50), day 5 (n=49), week 2 (n=43) and week 4 (n=19); NPY
(10 pmol/100 nL) at day 1 (n=42), day 5 (n=40), week 2 (n=29) and week 4 (n=11); the Y_1 agonist

F^{7}P^{34}-NPY (10 pmol/100 nL) at day 1 (n=15), day 5 (n=14), week 2 (n=14) and week 4 (n=10); the
Y$_5$ agonist cPP (10 pmol/100 nL) at day 1 (n=7), day 5 (n=7), week 2 (n=7) and week 4 (n=7); and
NPY (10 pmol/100 nL) immediately preceded by the Y$_5$ receptor antagonist CGP71683A (Y$_5$A, 10
pmol/100 nL) at day 1 (n=5), day 5 (n=5), week 2 (n=5) and week 4 (n=5) (Linear Mixed-model;
Treatment: $F_{(4, 102.30)}$ = 14.11, p = 3.33e^{-9}; Time: $F_{(3, 93.88)}$ = 1.66, p = 0.18; Treatment * Time: $F_{(12,}$
$_{91.12)}$ = 1.87, p = 0.049). **b**. Scatter plot of mean capacitance of BLA pyramidal neurons in acute
slices treated with vehicle control (n=64), NPY (n=53), F^{7}P^{34}-NPY (n=57), cPP (n=57) and NPY with
CGP (n=51) (Kruskal-Wallis H test with Dunn's post hoc test; $H_{(4)}$ = 47.74, p = 1.07e^{-9}). **c**. Scatter
plot of mean total dendritic length for cells treated as in **b** (ANOVA with Tukey's post hoc; $F_{(4, 205)}$
= 17.70, p = 1.70e^{-12}). vehicle (n=52), NPY (n=38), F^{7}P^{34}-NPY (n=32), cPP (n=50) and NPY with CGP
(n=39). **d**. Scatter plot of mean number of branch points for neurons in **c** (Kruskal-Wallis H test
with Dunn's post hoc test; $H_{(4)}$ = 56.53, p = 1.55e^{-11}). **e**. Sholl analysis for neurons in **c** and **d** as
indicated (2-way RM ANOVA with Tukey's post hoc test; <u>Treatment:</u> $F_{(4, 181)}$ = 13.09, p = 2.18e^{-9};
<u>Distance:</u> $F_{(49, 8869)}$ = 815.7, p = ~ 0; <u>Interaction:</u> $F_{(196, 8869)}$ = 4.81, p = 1.23e^{-91}). Blue = vehicle vs.
NPY; Green = vehicle vs. cPP. **f-j**. Representative traces from neurons treated as indicated above.
For **a**, symbols represent animals and black bars represent population means. For **b-d**, circles
represent neurons and black bars represent population means. For **e**, symbols represent
population means and for all, error bars indicate s.e.m. All tests were two-sided. *P<0.05,
**P<0.01, ***P<0.001, ****P<0.0001 for post hoc multiple comparisons. Scale bars in f-j = 100
µm.

Dendritic remodeling by NPY or CRF requires calcineurin or CaMKII, respectively

Because the long term stress -vulnerability and -resilience effects of CRF and NPY are

mediated via CaMKII- and calcineurin, respectively in the BLA *in vivo* (Rainnie et al., 2004, Sajdyk

et al., 2008,), we hypothesized that the effects of NPY or CRF on BLA PN dendritic structure

require the actions of calcineurin or CaMKII activity, respectively. We first repeatedly incubated

BLA OTCs with NPY (10 nM) together with cyclosporin A (CsA, 2 µM), a protein phosphatase

(PP) inhibitor which blocks calcineurin (Liu et al, 1991). Addition of CsA inhibited NPY-mediated

dendritic hypotrophy and decreases in capacitance (**Fig. 11a-d**), resulting instead in PN dendritic

hypertrophy, possibly by unmasking a Y$_2$R activation (**Fig. 11b-d**). While these results are

consistent with a role for calcineurin, CsA is a general PP inhibitor and is not selective for

calcineurin (PP2b). Okadaic acid (OA) selectively inhibits both PP1 and PP2a but not calcineurin

 407 (Cohen et al., 1990). Treatment of the BLA OTCs with the Y_5-agonist cPP (100 nM) in the presence

408 of OA (10 nM) did not alter the effects of the Y_5R agonist on either capacitance or dendritic

409 morphology, consistent with a specific role for calcineurin not only in NPY-mediated anxiolysis

410 (Sajdyk et al., 2008) but also in NPY-mediated PN dendritic hypotrophy (**Fig. 11a-d**).

Figure 11. Inhibitors of Calcineurin (CsA) and of CaMKII (AIP) respectively block NPY- and CRF-mediated dendritic remodeling in BLA OTCs. a. Scatter plot of mean capacitance of BLA OTC pyramidal neurons treated with vehicle control (n=28) compared to CsA (2 nM) alone (n=35), CsA (2 nM) and NPY (10 nM) (n=31), OA (10 nM) alone (n=39) and OA (10 nM) and cPP (10 nM) (n=38) ($H_{(4)}$ = 22.80, p = 0.0001). Dashed lines represent mean for neurons treated with NPY (10 nM) (blue) and cPP (100 nM) (green). **b**. Scatter plot of mean total dendritic length of pyramidal neurons treated as in **a**, vehicle (n=26), CsA (n=26), CsA and NPY (n=30), OA (n=26) and OA and cPP (n=24) ($H_{(4)}$ = 27.88, p = $1.32e^{-5}$). **c.** Scatter plot of mean number of branch points for neurons in **b** and **c,** with dashed lines representing the mean for neurons treated with NPY (blue) or cPP (green) ($H_{(4)}$ = 28.93, p = $8.10e^{-6}$). **d**. Sholl analysis for neurons in **b** treated as indicated, data for neurons treated with NPY (blue) or cPP (green). Black = control vs. OA + cPP; Gray = control vs. CsA + NPY (<u>Treatment:</u> $F_{(2, 78)}$ = 11.86, p = $3.18e^{-5}$; <u>Distance:</u> $F_{(40, 3120)}$ = 150.0, p ~ 0.; <u>Interaction:</u> $F_{(80, 3120)}$ = 2.39, p = $1.15e^{-10}$). **e**. Scatter plot of mean capacitance values for BLA OTC pyramidal cells treated with vehicle (n=28), AIP (40 nM) (n=27) and AIP (40 nM) and CRF (30 nM) (n=29), with dashed line representing neurons treated with CRF ($H_{(2)}$ = 0.77, p = 0.68). **f**. Scatter plot of mean total dendritic length for pyramidal cells treated as in **e** with neurons treated with CRF (dashed line) ($H_{(2)}$ = 1.23, p = 0.54). Vehicle (n=26), AIP (n=21) and AIP and CRF (n=21). **g**. Scatter plot of mean number of branch points in neurons in **f** with the mean for neurons treated with CRF (dashed line) ($H_{(2)}$ = 0.81, p = 0.67). **h**. Sholl analysis for neurons in **f** and **g** treated as indicated compared with neurons treated with CRF (dashed line) (<u>Treatment:</u> $F_{(2, 65)}$ = 0.50, p = 0.61; <u>Distance:</u> $F_{(40, 2600)}$ = 165.3, p ~ 0; <u>Interaction:</u> $F_{(80, 2600)}$ = 0.48, p = 0.99). For **a-c,e-g**, Kruskal-Wallis H tests with Dunn's post hoc tests. For **d,h**, 2-way RM ANOVA with Tukey's post-hoc tests. For **a-c,e-g**, circles represent neurons, black bars represent population means, while in **d,h**, symbols represent population means and for all, error bars indicate s.e.m. All tests were two-sided. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 for post hoc multiple comparisons.

We then tested the role of CaMKII in mediating the hypertrophic actions of CRF in BLA OTC PNs by using a cell-permeant CaMKII inhibitor, the myristoylated version of autocamtide-2-related inhibitory peptide (AIP –Ishida et al., 1995). Inhibiting CaMKII activity with AIP (40 nM) prevented the CRF-mediated (30 nM) increase in cell capacitance and dendritic hypertrophy (**Fig. 11e-h**). This is consistent with a role for CaMKII in CRF-mediated dendritic hypertrophy and the resultant stress vulnerability generated by repeated stress and CRF treatment (Rainnie et al., 2004).

444
445 **DISCUSSION**
446

Unraveling mechanisms mediating adaptive and pathophysiological responses to stress is

key to developing treatments for stress-related disorders. The countervailing actions of CRF and

NPY on BLA PNs represent a suite of mechanisms contributing to adaptive stress responses in the

amygdala. Although both neuropeptides induce long-term neural changes linked to the initiation

or mitigation of stress responses, the mechanisms underlying the effects of these neuropeptides are

unclear. Here, we identify three novel and important insights into the function of NPY and CRF in

amygdala physiology. First, we found that NPY-induced stress resilience ultimately involves a

novel, persistent form of homeostatic plasticity, specifically, the reversible remodeling of dendritic

structure in BLA pyramidal output neurons mediated by the Y_5R, which counterbalances the

persistent hypertrophy seen with stress or repeated CRF exposure. Second, different NPY receptors

play nuanced and complex roles in the short-and long-term regulation of BLA excitability, structure

and behavior. Finally, the BLA OTC preparation is a robust *in vitro* model, that mimics and predicts

long-term *in vivo* responses of BLA neurons.

Chronic stress or UCN treatment causes dendritic hypertrophy and increases excitatory

inputs to BLA PNs (Padival et al., 2013, Rainnie et al., 2004); these increases in dendritic length

and spine number are associated with enhanced excitatory drive onto BLA PNs which increases

anxiety (Mitra et al., 2005, Padival et al., 2013, Adamec et al., 2012, Hill et al., 2011 Vyas et al.,

2002; 2006). Repeated intra-BLA administration of NPY *in vivo* decreases the overall excitability

of BLA PNs (Silveira Villarroel et al., 2018). Here we demonstrate similar effects of NPY on PNs

in BLA OTC, as well as its ability to reverse or constrain CRF's actions. Following repeated NPY

or CRF treatment, PNs in BLA OTCs also demonstrated respective decreases and increases in

sEPSC frequencies, consistent with their respective behavioral effects. NPY and CRF thus appear

to act via multiple, parallel mechanisms to mediate their opposing forms of plasticity, which when

combined, drive BLA PN structure toward a resting state similar to that of naïve BLA. Consistent

with a key role in stress homeostasis, NPY is thus poised to restore balance to the organism once a

threat has dissipated.

The predominant role of the Y_5R in driving the long-term effects on BLA physiology and

behavior was unanticipated. The Y_1R, which to date has received considerable attention in stress-

mitigation, contributes only to the acute actions of NPY on SI behavior and fear extinction

(Giesbrecht et al., 2010, Gutman et al., 2008). Because all NPY receptors couple to $G_{i/o}$ proteins

(Michel et al., 1998), the different roles played by these two receptors might result from independent

trafficking into separate signaling compartments within BLA neurons (Marley et al., 2013).

Moreover, while the Y_1R desensitizes in somatic cell lines (Berglund et al., 2003) and intact systems

(Holliday et al., 2005), evidence indicates that the Y_5R does not desensitize (Böhme et al., 2008).

This potentially significant difference suggests that the Y_5R is more likely to be available for lasting

mitigation of stress responses.

Figure 12. Proposed model for the mechanism of action of NPY and CRF leading to dendritic hypotrophy or hypertrophy respectively in BLA pyramidal cells. a. NPY treatment results in net activation of the Y_5R and PP2B (Calcineurin), resulting in a reduction in dendritic arborization (illustrated right). **b.** CRF treatment results in net activation of CaMKII and hypertrophy of the dendritic arbor. (Illustrated right).

The long-term effects of NPY or CRF require the respective activation of the Ca^{2+}-dependent enzymes calcineurin and CaMKII, indicating a critical role for alterations in intracellular Ca^{2+} levels in BLA PNs as demonstrated both *in vivo* (Sajdyk et al., 2008, Rainnie et al., 2004) and

now here in BLA OTCs. The acute and long-term enhancement of excitability caused by CRF

would result in robust elevations in intracellular Ca^{2+} ($[Ca^{2+}]_i$), consistent with CaMKII activation

requirements (Rainnie et al., 2004). However, how NPY achieves the more modest $[Ca^{2+}]_i$ elevation

needed by calcineurin is less obvious. In this context, the differential contributions of the multiple

NPY receptor subtypes within the BLA may provide some insight. Acute activation of BLA Y_2Rs

paradoxically results in anxiogenic behavioral responses (Sajdyk et al., 2004a; 2004b), and resulted

in CRF-like changes in BLA OTC neurons here. In both rat and mouse BLA, the Y_2R agonist

reduces tonic $GABA_BR$ activation in BLA PNs by a presynaptically-mediated reduction in dendritic

GABA release. This causes an increase in dendritic excitability and a disinhibition of dendritic Ca^{2+}

channels, which in turn permits a moderately elevated Ca^{2+} influx in about half of BLA PNs

(Mackay et al., 2019). These effects suggest that NPY release in BLA activates all 3 NPY receptors:

Y_1Rs, which acutely hyperpolarize BLA PNs, Y_2Rs, which reduce dendritic $GABA_BR$ activation

and Y_5Rs, whose long-term effects rely on the preferential activation of calcineurin. In this scenario,

the actions of the 3 separate NPY receptors would synergize to faciliate the Y_5R effect by modestly

elevating Ca^{2+} influx into BLA PNs (**Fig. 12a)** while preventing the larger increases in $[Ca^{2+}]_i$ that

would preferentially activate the CaMKII pathway favored by the CRF system (**Fig. 12b**). This

model could in part also explain the greater potency of the pan-agonist NPY to affect dendritic

structure. Nonetheless, further work is needed to more completely understand the pathways and

mechanisms involved in this long-term structural plasticity.

Regulation of cytosolic protein phosphorylation plays a key role in the structural plasticity

seen here, consistent with evidence that calcineurin and CaMKII act as downstream targets of

anxiolytic and anxiogenic molecules in the BLA (Rainnie et al., 2004, Mineur et al., 2014, Lin et

al., 2003). Both enzymes are inherently associated with synaptic plasticity in the BLA, so CRF- and

NPY-mediated alterations in dendritic structure might also engage mechanisms that underlie LTP

(long term potentiation) and LTD (long term depression), respectively (Pape and Pare, 2010). As

LTP and LTD each require different levels of $[Ca^{2+}]_i$, studies on intra-dendritic Ca^{2+} dynamics could

address the contributions made by acute or chronic changes in Ca^{2+} influx, either via NMDA

receptors (Rainnie et al., 2004) or voltage-dependent Ca^{2+} channels, to dendritic remodeling (Pape

and Pare, 2010).

Figure 13. Correlation between whole-cell capacitance and dendritic length. a. Whole-cell
capacitance plotted against total dendritic length for 259 BLA OTC neurons from 100 different
OTCs taken from all experimental groups.

Total dendritic length and whole-cell capacitance were tightly correlated in BLA OTC PNs

over a variety of treatments (**Fig. 13**), were seen in BLA both *in vitro* and *in vivo*, and corresponded

with long-term behavioral changes. Despite the potent acute behavioral effects of Y_1R agonists,

they altered neither dendritic structure nor long-term behavior. It appears reasonable to speculate

that structural changes mediated by repeated peptide treatments result in the long-term changes in

behavioral response to stressors. In any case, this structural plasticity could provide a novel *in vitro*

bioassay predicting efficacy of drug candidates.

OTCs have proved useful in the study of longer-term nervous system changes such as

synaptic plasticity (Debanne et al., 1999, Selcher et al., 2012), and those associated with pathologies

such as chronic pain (Lu et al., 2006), or even prion disorders (Campeau et al., 2013). Since such

changes can involve alterations not only in synaptic microanatomy but also more robust changes in

cellular architecture, the preservation of anatomical relations within the OTC is a distinct advantage.

Mature PNs in BLA OTCs share more electrophysiological attributes with those in age-matched

(10W) *ex vivo* BLA than those in *ex vivo* P14 rats, despite their increased compactness compared

with neurons from either *ex vivo* preparation. Capacitance values were smaller, I_h activation

kinetics (tau$_{fast}$ and tau$_{slow}$) were faster, and sEPSC and sIPSC amplitudes were larger in BLA OTCs

compared with the acute *ex vivo* neurons **(Table 1)**. However, the properties of BLA OTC PNs as

indicated by action potential properties, firing rates and AHPs suggests that growth within OTCs

constrains the physical extent of otherwise physiologically mature OTC neurons.

	Acute P14	Acute 10W	OTC EP 10W	ANOVA
RMP (mV)	-76.8 ± 0.5 *** n=56	-81.3 ± 0.7 n=44	-77.9 ± 1.0** n=54	$F_{(2, 151)} = 9.05$, p= 0.0002
Rheobase (pA)	126.3 ± 5.2 **** n=56	309.2 ± 13.8 n=44	248.2 ± 24.8*, #### n=54	$F_{(2, 151)} = 29.84$, p = 0.00019
I_R (MΩ)	176.8 ± 6.4 **** n=56	71.5 ± 5.2 n=22	103.4 ± 6.3*, #### n=49	$F_{(2, 124)} = 64.16$, p = 7.43e-20
I_h max (pA)	-175.2 ± 7.3*** n=52	-362.5 ± 25.9 n=48	-290.4 ± 20.1*, ### n=49	$F_{(2, 146)} = 25.09$, P = 4.31e-10
I_h density (pA/pF)	-0.96 ± 0.03 * n=52	-1.36 ± 0.10 n=22	-1.64 ± 0.13 #### n=49	$F_{(2, 120)} = 14.38$, p=2.52e-6
C_m (pF)	180.2 ± 3.7 **** N=56	230.6 ± 7.3 N=44	150.0 ± 5.8 ****, ### N=58	$F_{(2, 155)} = 49.98$ P=1.75e-18,
sIPSC AMP (pA)	25.5 ± 1.1 n=25	24.13 ± 1.8 n=25	49.8 ± 4.9 ****, #### n=25	$F_{(2, 72)} = 21.59$, p=4.51e-8

sIPSC IEI (ms)	2249 ± 417 **** n=25	213.9 ± 22.1 n=25	155.6 ± 19.3 #### n=25	F(2, 72) = 24.41 p= 8.07e-9
sEPSC AMP (pA)	25.9 ± 0.9 n=25	26.87 ± 2.3 n=25	44.8 ± 2.1 ****, #### n=25	F(2, 72) = 32.15 p = 1.05e-10
sEPSC IEI (ms)	1241 ± 347.2 ** n=25	171.1 ± 26.4 n=25	221.5 ± 32.2 ## n=25	F(2, 72) = 8.95 p=0.0003
Total Dendritic Length (µm)	5986 ± 337 **** n=17	8644 ± 284 n=36	3625 ± 187.4****, ####, n=39	F(2, 89) = 113.6 p = 3.16e-25
# of branches	37.5 ± 2.3 * n=17	46.3 ± 1.9 n=36	20.5 ± 1.5 ****, #### n=39	F(2, 89) = 60.17 p = 2.95e-17

Table 1. Comparison of electrophysiological and morphological properties of acute P14, acute 10W and EP 10W OTC pyramidal neurons in the BLA.

Data presented as mean ± s.e.m. One-way ANOVA with Tukey's post-hoc test for multiple comparisons were used for all analyses; *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 vs Acute 10W, #P<0.05, ##P<0.01, ###P<0.001, ####P<0.0001 vs Acute P14 for multiple comparisons.

In summary, we report a novel, readily and completely reversible alteration in PN dendritic structure that accompanies the long-term increase in behavioral stress resilience caused by NPY in the BLA, the mirror opposite of equivalent CRF actions. These robust changes are readily observed in BLA OTCs, and are reflected nearly perfectly *in vivo*. The acute and longer-term actions both of NPY and CRF involve regulation of the same elements (I_h, dendrites) in the PN, changes that correlate with alterations in excitability. The coordinated, countervailing regulation of these PN properties by NPY and CRF suggests other signals could access this machinery to regulate stress responses mediated by the BLA.

MATERIALS AND METHODS

Animals

All animal procedures were approved by the University of Alberta Animal Care and Use Committee: Health Sciences, in accordance with the guidelines of the Canadian Council on Animal

Care. Litters of postnatal day (P) 14 and 5-week old male Sprague-Dawley rats were used. P14

pups were given complete access to the dams for maternal care; then were removed from the dam

just prior either to preparation of OTCs or for a subset of acute slice studies. Five week old rats

were grouped housed (2-3 animals per cage) with 12:12 hours light:dark schedule and *ad libitum*

access to food and water.

Development and validation of organotypic slice cultures (OTCs) of BLA

Postnatal development of the rat BLA continues until P28 (Ehrlich et al., 2012; 2013).

Generation of OTCs from late adolescent or early adult-stage animals would be ideal, but the

viability of OTCs declines steeply with postnatal age of origin (Humpel, 2015, Kim et al,, 2013).

Our and others' previous work used larger (250 - 275g) Wistar or older (P42) Sprague-Dawley rats

in studies of NPY and UCN effects on stress-resilience/vulnerability (Molosh et al., 2013), and

acute and persistent NPY actions (Giesbrecht et al., 2010, Silveira Villarroel et al., 2018).

OTC preparation

Organotypic cultures were prepared using the interface method described by Stoppini et al.

(1991) with minor modifications. Briefly, P14 rats were decapitated and their brains rapidly

removed under sterile conditions and submerged in ice-cold slicing solution [Hank's balanced salt

solution (HBSS) + D-glucose (0.6% final) + kynurenic acid (30 µM final)]. The brain was imbedded

in agarose and trimmed agarose-brain blocks were secured with cyanoacrylate glue to a custom

slicing chamber, immersed in cold (<4ºC) slicing solution. Coronal slices (350 µm) were cut serially

from rostral to caudal using a vibratome and trimmed to size. Four slices per hemisphere containing

the BLA (as determined visually with a dissecting microscope), were allowed to rest in fresh slicing

solution at 4°C for 30-45 minutes, and mounted on individual semiporous membrane inserts and

placed in 24-well plates with 300 µL of culture media (50% Minimal Essential Medium (MEM),

25% heat-inactivated horse serum, 25% HBSS, supplemented with: 1 mM Glutamax, 1% D-

glucose, 0.5 mM L-ascorbic acid, and 25 U/mL penicillin/streptomycin). After 48 hours in culture,

an anti-mitotic solution of 1:1:1 cytosine-β-D-arabino-funanoside, uridine, and 5-Fluro-2'-

deoxyuridine was added to the media (0.5 µM final concentration) for 24 hours to prevent glial

proliferation. Slices were maintained at 37°C in 5%/95% CO_2/air. Media was changed three times

per week for the duration of the experiment, except as indicated.

BLA OTCs prepared from rats at P21 and P28 were poorly viable, but those from P14 (2

week) rats remained robustly viable to at least 8 weeks in culture, the equivalent postnatal (EP) age

of P70 (**Fig. 3a**), permitting comparison of the electrophysiological and morphological properties

of PNs from the OTCs (see below). Such OTCs were used for this entire study

OTC drug incubations

To mimic EP 6 weeks (4 weeks in culture), slices were treated with fresh media containing

indicated concentrations of reagents, changed daily for five consecutive days; the last change

remained until the normal Day 7 media change. When cultures were serially treated as above first

with NPY then with CRF (or vice-versa), incubations proceeded with one peptide as above,

followed with an identical one-week daily protocol using the opposing peptide starting with the

Day 7 media change. A control subset of cultures was treated with the inactive desamido NPY

analogue (daNPY – 100 nM). After incubations, media changes were again performed three times

weekly (**Fig. 3a**).

Acute slice preparation

P14 or P70 rats were decapitated and their brains rapidly submerged in cold (4°C) artificial

cerebrospinal fluid (aCSF) slicing solution, containing (in mM): 118 NaCl, 3 KCl, 1.3 $MgSO_4$, 1.4

NaH_2PO_4, 5.0 $MgCl_2$, 10 D-glucose, 26 $NaHCO_3$, 2.5 $CaCl_2$ and 1.0 kynurenic acid and bubbled

with carbogen (95% O2, 5% CO_2) (Silveira Villarroel et al., 2018). Coronal sections (300 μm)

containing the BLA were prepared using a vibrating slicer (HR2; Sigmann Elektronik, Hüffenhardt,

Germany). Brain slices were then transferred to a room temperature (22°C), carbogenated aCSF

containing the following (in mM): 124 NaCl, 3 KCl, 1.3 $MgSO_4$, 1.4 NaH_2PO_4, 10 D-glucose, 26

$NaHCO_3$, and 2.5 $CaCl_2$. Osmolality was adjusted to 300 mOsm/kg.

Comparison with PNs from acute slice preparations

PNs from BLA OTCs are morphologically complex and electrophysiologically more similar to

PNs in acute slices from P70 than from P14 rats, though OTC neurons are considerably smaller
than those from either acutely ex vivo (**Table 1**). Moreover, repeated NPY (100 nM) treatment of
OTCs *in vitro* (**Fig. 3a**) increased the mean amplitude of spontaneous inhibitory postsynaptic
currents (sIPSCs), while increasing the mean inter-event interval (IEI) of spontaneous excitatory
postsynaptic currents (sEPSCs). In all, repeated NPY treatment resulted in similar a net increase
in synaptic inhibition and reduction of synaptic excitation onto BLA PNs in OTCs as *in vivo* (**Fig.**
3b-e).**Stereotaxic surgery and intracranial injections.**

Stereotaxic surgery was carried out as described in detail elsewhere (Silveira Villarroel et

al., 2018). Briefly, 5-week old animals were acclimatized to the animal facility and handled daily

for 1 week prior to experimental manipulation. Rats were anesthetized with ketamine/xylazine

(90:10 mg/kg) or with isoflurane using a SomnoSuite low flow vaporizer (Kent Scientific,

Torrington, CT, USA) and placed in a stereotaxic apparatus (Kopf, Tujunga, CA, USA).

Anaesthesia was monitored carefully to maintain surgical plane throughout, and animals were kept

warm with a temperature-controlled blanket (Harvard Apparatus, Holliston MA, USA). Bilateral

guide cannulas (26 gauge; Plastics One, Roanoke, VA, USA) were implanted to just above (2 mm)

the BLA [anteroposterior (AP): -2.3; mediolateral (ML): ±5.0; dorsoventral (DV): -6.4; incisor bar:

-3.2 mm] (Paxinos and Watson, 1986). The cannulas were secured to the skull with four stainless

steel screws (2.8 mm; Plastics One, Roanoke, VA, USA) and self-curing acrylic resin (Lang Dental

Manufacturing Company, Inc., IN, USA). After completion of surgery, all animals received

meloxicam (Boerhringer Ingelheim, Burlington, Ontario, Canada; 1 mg/kg, s.c.), 1 ml of saline

0.9% s.c., and were placed in a warm environment until they had fully recovered from the

anesthetic. Rats remained singly housed until they fully recovered from surgery, then were housed

in pairs. Pairs were separated and animals housed singly one day prior to behavioral testing (below).

Intracranial injections

As reported elsewhere (Silveira Villarroel et al., 2018), all compounds were delivered

bilaterally via 33 gauge injection cannulas (Plastics One) that extended 2 mm beyond the guide

cannulas, using a dual-channel infusion pump (PHD ULTRA, Harvard Apparatus, Holliston, MA,

USA). Vehicle (sterile saline) alone or containing drug(s) as described was delivered at 100 nL/30s

per side; cannulae were left in place for 1 minute afterward to prevent backflow. All injections were

performed once daily between 8:00 AM and 10:00 AM for successive 5 days. Animals rested in

their home cage for 30 min following injections, then tested (if in their protocol) for social

interaction (SI).

Behavioral testing

The SI test was performed according to Sajdyk et al., (2008), with minor modifications. Briefly,

an experimental animal was placed into the SI box (96 cm long x 96 cm wide x 30 cm high with

an open top) simultaneous with a partner rat of the same sex, age and weight, housed under

identical conditions but which had not encountered the experimental animal previously. Seventy-

two hours after implantation surgery, rats were given a sham injection, then allowed to spend 10

minutes alone in the arena to acclimate. Twenty-four hours later they received another mock

injection, and SI was tested to establish baseline anxiety-like levels prior to any experimental

manipulations. Two days later, rats received a bilateral intra-BLA injection (vehicle or drug) and

were tested for SI 30 min later (day 1). Injections continued daily at the same time and the SI test

was repeated 30 min after the fifth injection (day 5), then at 2 weeks and 4 weeks from the first

injection. Behavior was video recorded and later analysed by individuals blinded to treatment.

Whole-cell patch clamp electrophysiology

Acutely prepared slices or EP 9-10 week OTC slices (with culture insert membrane still

attached), were transferred to a fixed recording chamber beneath a movable upright microscope

(Axioskop FS2; Carl Zeiss, Germany)(Giesbrecht et al., 2010, Silveira Villarroel et al., 2018).

Slices were perfused at 2-3 mL/min for at least 10 minutes before recording, with warmed (32-

34°C), carbogenated aSCF (300 mOsm/L acute slices and 320 mOsm/L for OTCs - adjusted with

NaCl). Patch pipettes were pulled from borosilicate glass (TW150F; World Precision Instruments,

Sarasota, FL, USA) with a two-stage puller (PP-83; Narishige, East Meadow, NY, USA) and had a

resistance of 4-6 MΩ with an internal solution (mM): 126 K-gluconate, 4 KCl, 10 HEPES, 5

MgATP, 0.3 NaGTP, 1 EGTA, 0.3 $CaCl_2$ and 0.2% neurobiotin (pH 7.27, 275 mOsm/L for acute

slices and 300 mOsm/L for OTCs).

The BLA was identified under a 5x objective and PNs identified visually with infrared-

differential interference contrast (60x) optics. Neurons were selected as previously reported

(Giesbrecht et al., 2010, Silveira Villarroel et al., 2018) based on morphological and

electrophysiological criteria. PNs chosen for study were in random locations throughout the BLA;

1-3 neurons per slice were studied. Either a Multiclamp 700B or AxoClamp2A amplifier was used

together with a DigiData 1322 or Digidata 1440 interface and pCLAMP v10.4 software (all

Molecular Devices, Sunnyvale CA, USA). Gigohm seal, whole-cell recordings were low-pass

filtered at 2 kHz and digitized at 10 kHz. Access resistance was measured throughout the

experiment, and only those cells with changes less than 20% were kept for further analysis. The

membrane potential reported was corrected offline for the calculated 15 mV liquid junction

potential (Silveira Villarroel et al., 2018, Chee et al., 2010). Recordings (> 20 min) were terminated

by gradually withdrawing the pipette until the membrane re-sealed.

Capacitance was calculated offline with pClamp, from the area under the curve of the current

amplitude from a 200 ms, 10 mV hyperpolarizing voltage step (Taylor, 2012). No capacitance

compensation was applied during these measurements. Spontaneous synaptic activity was assessed

in voltage clamp at a (corrected) holding potential of -55 mV and sampled at 20 kHz.

Cell processing and labeling

Neurons from *ex vivo* and OTC slices were all processed, stained, imaged, reconstructed

and subjected to the same morphological analyses. Following patch clamp recordings, OTCs and

acute slices were fixed with 10 % formalin (Thermo Fisher Scientific, Ottawa, Ontario, Canada) for

24-72 h at 4°C, and then transferred to 0.2% sodium azide in PBS for subsequent storage at 4 °C.

Slices were processed within 4 weeks following patch clamp recordings. Free-floating slices were

washed for 3 x 10 min periods in PBS before blocking and permeabilization for 2 h with 0.3 %

Triton X-100 in PBS and 4% normal goat serum (Sigma-Aldrich, Oakville, Ontario, Canada). Slices

were incubated with streptavidin conjugated with Alexa Fluor 555 or 546 (1:1000; Life

Technologies, Grand Island, New York, USA) in 0.3 % Triton X-100 in PBS with 4% normal goat

serum for 3 h at room temperature. Slices were washed 4 x10 min each in PBS, mounted on

Superfrost Plus slides and coverslips were applied with Prolong Gold mounting media (Thermo

Fisher). Slides were allowed to air dry in the dark at room temperature before being imaged via

confocal microscopy.

Imaging and neuronal reconstruction

For morphological analysis of filled neurons, z-stack images were obtained with a 20x

objective, 1024 x 1024 resolution, 100 Hz, excitation wavelength 543 nm, for a series of 0.8 μm

steps with a laser scanning confocal microscope (Leica TCS SP5; Leica Microsystems, Canada).

To determine the position of each neuron within the slice, multiple single-plane images were taken

at 10x magnification, 512 x 512 resolution and 100 Hz, and stitched together using Leica LAS AF

software. Only those neurons whose dendritic arbor was clearly and completely filled and cell body

was clearly within the boundaries of the BLA were used for analysis. Neuronal reconstruction was

performed using the simple neurite tracer function in FIJI (NIH, Bethesda, MD, USA). Following

tracing, Sholl analysis (Sholl, 1953) of the entire 3-dimensional dendritic arbor was performed with

another FIJI software module using concentric circles with increasing radii of 10 μm to determine

dendritic intersections vs. distance from the soma. Total dendritic length and dendritic branching

was calculated using FIJI and Excel software. Quantification of dendritic spine density was

achieved by manually counting dendritic spines in z-stack images taken at 100x magnification, 1024

x 1024 resolution and 100 Hz for a series of 0.5 um steps. Spines were counted on three separate,

randomly selected, 100 μm dendrite segments for each neuron analysed in a treatment group. All

protrusions that were connected to the dendrite segment were considered as spines, but were not

categorized by spine morphologies. Total spine number was estimated from the average spine

density and total dendritic length for an individual neuron. Experimenters were blinded to culture

treatment at the time of reconstructions.

Reagents and Drugs

NPY (human, rat) was purchased from Polypeptide Group (San Diego, CA, USA) while

CRF was obtained from Phoenix Pharmaceuticals Inc. (Burlingame, CA, USA). The Y_1-agonist

$(F^7,P^{34}$-NPY), the Y_5-agonist ($[cPP^{1-7},NPY^{19-32},Ala^{31},Aib^{32},Gln^{34}]hPP$) and the Y_2 agonist ($[ahx^{5-}$

$^{24}]$ NPY) were generous gifts from Dr. A.G. Beck-Sickinger (Leipzig, Germany), while the Y_5-

antagonist, CGP71683, was purchased from Tocris Bioscience (Bristol, UK). Cyclosporine A and

okadaic acid were gifts from Drs. Shairaz Baksh and Charles Holmes (University of Alberta,

Canada). The cell permeable CaMKII inhibitor, myristoylated-autocamtide-2- related inhibitory

peptide, was obtained from Enzo Life Sciences, Inc (Farmingdale, NY, USA). Culture reagents

were all obtained from Gibco, except for Cytosine–β-D-arabino-furanoside, Uridine and 5'Fluro-

2'deoxyuridine, which were from Sigma-Aldrich Canada. Pipette solution reagents were all from

Sigma-Aldrich, except for Na-GTP (Roche Diagnostics).

Data analysis and statistics

Electrophysiological recordings were viewed offline using pCLAMP. Traces from

electrophysiological recordings, statistical analysis and graphs were prepared using versions 5 – 8

of Prism software (GraphPad, San Diego, CA, USA) or SPSS (version 20 - IBM). Neuronal

reconstruction and Sholl analysis was performed using modules of the FIJI software suite (National

Institutes of Health). Unless otherwise stated, all data are represented as mean ± s.e.m. D'Agostino-

Pearson omnibus normality test was applied to determine data distributions. One-way analysis of

variance (ANOVA) with Tukey's post-hoc test or Kruskal-Wallis H tests with Dunn's post hoc test

were used as indicated, to compare cell capacitance, total dendritic lengths, branch points and spine

density/total spine estimates between treatments. For Sholl analysis, two-way repeated measures

ANOVA were used with Tukey's post-hoc test for multiple comparisons. A Linear Mixed-Model

was used for analysis of SI studies. Treatment and time (with interaction term) were used as fixed

effects, while we used intercepts for subjects as random effects. For data in **Table 1**, one-way

ANOVA with Tukey's post-hoc test for multiple comparisons was used. For sEPSC and sIPSC

analyses, two-sided Student's t-test or Mann-Whitney U tests were performed when applicable.

OTC cultures for any given experiment were obtained from at least 3 animals (typically more), with

data acquired from 1-3 neurons per OTC. With the extent of time in culture, and minimal number

of neurons recorded from per slice, we opted to treat each neuron as an independent observation for

statistical analysis. The number of animals, cultures and neurons per experimental group are

indicated in **Table 2**. No statistical test was used to predetermine sample sizes, but our sample sizes

are similar to those previously reported in the field (Adamec et al., 2012, Silveira Villarroel et al.,

2018, DeSimoni et al., 2003). All analyses were two-sided. Significance was set at $P<0.05$ (*

P<0.05, **P<0.01, ***P<0.001, ****P<0.0001).

Acknowledgments: We are grateful for the generous contributions of NPY receptor-selective

agonists by Prof. Annette Beck-Sickinger, University of Leipzig. SDM received funding from

Alberta Innovates Health Solutions (AIHS) (Doctoral studentship). JPM received Canadian

Institute of Health Research (CIHR) (Doctoral and Masters studentships), and a Doctoral

studentship from AIHS. WFC was a Medical Scientist of the Alberta Heritage Foundation for

Medical Research for most of this period.

This work was supported by NIH grants MH081152 and MH090297 (to JHU & WFC), and

support by the University of Alberta Hospitals Foundation to WFC.

Author Contributions: . SDM, APMT, JHU and WFC designed experiments. SDM, APMT,

AM, HSV and JM performed experiments. SDM, APMT and AM analyzed data. SDM, JHU and

WFC wrote the manuscript. SDM and WFC conceived the study. All authors edited the

manuscript.

Competing Interests: The authors declare no competing financial interests.

Data availability: The data that support the findings of this study are available from the

corresponding author upon reasonable request.

	Electrophysiological Figures					Morphology Figures			
Figure	**Treatment**	**Neurons**	**Cultures**	**Animals**	**Figure**	**Treatment**	**Neurons**	**Cultures**	**Animals**
1a	Control	42	20	18	1b-d	Control	26	19	16
1a	NPY ($1e^{-10}$ M)	31	15	3	1b-d	NPY ($1e^{-10}$ M)	26	14	3
1a	NPY ($1e^{-9}$ M)	43	25	23	1b-d	NPY ($1e^{-9}$ M)	25	16	15
1a	NPY ($1e^{-8}$ M)	47	27	20	1b-d	NPY ($1e^{-8}$ M)	21	16	14
1a	NPY ($1e^{-7}$ M)	49	26	21	1b-d	NPY ($1e^{-7}$ M)	24	16	13
1a	da-NPY ($1e^{-7}$ M)	30	14	13	1b-d	da-NPY ($1e^{-7}$ M)	20	12	11
3d,e	Control	25	11	5	2d,e	Control	15	11	6
3d,e	NPY ($1e^{-7}$ M)	25	13	7	2d,e	NPY ($1e^{-7}$ M)	15	10	6
3h	Control	25	14	8	2d,e	CRF ($3e^{-8}$ M)	15	10	5
3h	CRF ($3e^{-8}$ M)	25	16	9	2d,e	cPP ($1e^{-7}$ M)	15	8	5
4a	Control	39	19	14	4b-d	Control	33	19	14
4a	CRF ($3e^{-10}$ M)	40	21	11	4b-d	CRF ($3e^{-10}$ M)	31	19	11
4a	CRF ($3e^{-9}$ M)	36	18	11	4b-d	CRF ($3e^{-9}$ M)	26	15	11
4a	CRF ($3e^{-8}$ M)	38	20	11	4b-d	CRF ($3e^{-8}$ M)	28	18	10
5a	Control	42	24	21	5b-d	Control	26	21	16
5a	CRF-NPY	31	15	6	5b-d	CRF-NPY	23	13	6
5a	NPY-CRF	32	15	7	5b-d	NPY-CRF	23	12	6
5h	CRF-NPY	25	12	5	6b-d	Control	27	15	13
5h	NPY-CRF	25	10	4	6b-d	F^7P^{34}-NPY ($1e^{-9}$ M)	19	14	12
6a	Control	30	22	19	6b-d	F^7P^{34}-NPY ($1e^{-8}$ M)	18	15	10
6a	F^7P^{34}-NPY ($1e^{-9}$ M)	28	19	15	6b-d	F^7P^{34}-NPY ($1e^{-7}$ M)	22	14	13
6a	F^7P^{34}-NPY ($1e^{-8}$ M)	34	20	14	8b-d	Control	23	11	9
6a	F^7P^{34}-NPY ($1e^{-7}$ M)	34	22	18	8b-d	ahx ($1e^{-9}$ M)	19	11	10
7c	Control	25	15	9	8b-d	ahx ($1e^{-8}$ M)	23	15	13
7c	F^7P^{34}-NPY ($1e^{-7}$ M)	25	18	13	8b-d	ahx ($1e^{-7}$ M)	23	13	10
7f,g	Control	25	15	9	9b-d	Control	26	17	13
7f,g	cPP ($1e^{-7}$ M)	25	14	8	9b-d	cPP ($1e^{-9}$ M)	23	17	11
8a	Control	26	12	9	9b-d	cPP ($1e^{-8}$ M)	31	17	12
8a	ahx ($1e^{-9}$ M)	23	10	8	9b-d	cPP ($1e^{-7}$ M)	32	17	13
8a	ahx ($1e^{-8}$ M)	21	12	8	9b-d	NPY ($1e^{-8}$ M) + CGP ($3e^{-8}$ M)	34	17	5
8a	ahx ($1e^{-7}$ M)	25	14	10	11b-d	Control	26	17	4
9a	Control	30	20	13	11b-d	CsA ($2e^{-9}$ M)	26	14	4
9a	cPP ($1e^{-9}$ M)	32	18	12	11b-d	CsA ($2e^{-9}$ M) + NPY ($1e^{-8}$ M)	30	17	4
9a	cPP ($1e^{-8}$ M)	47	26	16	11b-d	OA ($1e^{-8}$ M)	26	12	4
9a	cPP ($1e^{-7}$ M)	34	23	15	11b-d	OA ($1e^{-8}$ M) + cPP ($1e^{-7}$ M)	24	13	4
9a	NPY ($1e^{-8}$ M) + CGP ($3e^{-8}$ M)	45	25	6	11f-g	Control	26	13	10
11a	Control	28	17	4	11f-g	AIP ($4e^{-8}$ M)	21	10	4
11a	CsA ($2e^{-9}$ M)	35	16	4	11f-g	AIP ($4e^{-8}$ M) + CRF ($3e^{-8}$ M)	21	11	4
11a	CsA ($2e^{-9}$ M) + NPY ($1e^{-8}$ M)	31	18	4					
11a	OA ($1e^{-8}$ M)	39	15	4					
11a	OA ($1e^{-8}$ M) + cPP ($1e^{-7}$ M)	38	15	4					
11e	Control	28	15	11					
11e	AIP ($4e^{-8}$ M)	27	13	4					
11e	AIP ($4e^{-8}$ M) + CRF ($3e^{-8}$ M)	29	14	4					

779 **Table 2. Numbers of neurons, OTCs and animals used for capacitance and morphological**
780 **analysis, indexed by Figure.**

References

782
783
784 Adamec, R., Hebert, M., Blundell, J. and Mervis, R.F. (2012). Dendritic morphology of amygdala
785 and hippocampal neurons in more and less predator stress responsive rats and more and less
786 spontaneously anxious handled controls. Behav. Brain Res. 226, 133-146. doi:
787 10.1016/j.bbr.2011.09.009
788
Avossa, D., Rosato-Siri, M.D., Mazzarol, F. and Ballerini, L. (2003). Spinal circuits formation: a
study of developmentally regulated markers in organotypic cultures of embryonic mouse spinal
cord. Neuroscience 122, 391-405.

Beck-Sickinger AG, Grouzmann E, Hoffmann E, Gaida W, van Meir EG, Waeber B, Jung G.
(1992). A novel cyclic analog of neuropeptide Y specific for the Y2 receptor. Eur. J. Biochem.
206:957-964.

Berglund, M.M., Schober, D.A., Statnick, M.A., McDonald, P.H. and Gehlert, D.R. (2003). The
use of bioluminescence resonance energy transfer 2 to study neuropeptide Y receptor agonist-
induced beta-arrestin 2 interaction. J. Pharmacol. Exp. Ther. 306, 147-156.
798
799 Böhme I, Stichel J, Walther C, Mörl K, Beck-Sickinger AG.(2008). Agonist induced receptor
800 internalization of neuropeptide Y receptor subtypes depends on third intracellular loop and C-
801 terminus. Cell Signal. 20:1740-1749 doi: 10.1016/j.cellsig.2008.05.017.
802
803 Cabrele, C. and Beck-Sickinger, A.G. (2000). Molecular characterization of the ligand-receptor
804 interaction of the neuropeptide Y family. J Pept Sci 6, 97-122.
805
806 Cabrele, C., Wieland, H.A., Langer, M., Stidsen, C.E. and Beck-Sickinger, A.G. (2001). Y-
807 receptor affinity modulation by the design of pancreatic polypeptide/neuropeptide Y chimera led
808 to Y(5)-receptor ligands with picomolar affinity. Peptides 22, 365-378.
809
810 Campeau, J.L., Wu, G., Bell, J.R., Rasmussen, J. and Sim, V.L.(2013). Early increase and late
811 decrease of purkinje cell dendritic spine density in prion-infected organotypic mouse cerebellar
812 cultures. PLoS One 8, e81776. doi: 10.1371/journal.pone.0081776.
813
814 Chee, M.J., Myers, M.G., Jr., Price, C.J. and Colmers, W.F. (2010). Neuropeptide Y suppresses
815 anorexigenic output from the ventromedial nucleus of the hypothalamus. J. Neurosci. 30, 3380-
816 3390. doi: 10.1523/JNEUROSCI.4031-09.2010.
817
818 Cohen P., Holmes, C.F., Tsukitani, Y. (1990) Okadaic acid: a new probe for the study of cellular
819 regulation. Trends Biochem Sci. 15:98-102.
820
821 De Simoni, A., Griesinger, C.B. and Edwards, F.A. (2003) Development of rat CA1 neurones in
822 acute versus organotypic slices: role of experience in synaptic morphology and activity. J.
823 Physiol. 550, 135-147.
824

825 Debanne, D., Gahwiler, B.H. and Thompson, S.M. (1999). Heterogeneity of synaptic plasticity at
826 unitary CA3-CA1 and CA3-CA3 connections in rat hippocampal slice cultures. J. Neurosci. 19,
827 10664-10671.
828
829 Ehrlich, D.E., Ryan, S.J. and Rainnie, D.G. (2012). Postnatal development of electrophysiological
830 properties of principal neurons in the rat basolateral amygdala. J. Physiol. 590, 4819-4838. oi:
831 10.1113/jphysiol.2012.237453
832
833 Ehrlich, D.E., Ryan, S.J., Hazra, R., Guo, J.D. and Rainnie, D.G. (2013). Postnatal maturation of
834 GABAergic transmission in the rat basolateral amygdala. J. Neurophysiol. 110, 926-941. doi:
835 10.1152/jn.01105.2012
836
837 File S.E. and Seth, P. (2003). A review of 25 years of the social interaction test. Eur J Pharmacol.
838 463:35-536.
839
840 Gahwiler, B.H. (1981). Morphological differentiation of nerve cells in thin organotypic cultures
841 derived from rat hippocampus and cerebellum. Proc. R. Soc. Lond. B. Biol. Sci. 211, 287-290.
842
843 Giesbrecht, C.J., Mackay, J.P., Silveira, H.B., Urban, J.H. and Colmers, W.F. (2010)
844 Countervailing modulation of Ih by neuropeptide Y and corticotrophin-releasing factor in
845 basolateral amygdala as a possible mechanism for their effects on stress-related behaviors. J.
846 Neurosci. 30, 16970-16982. doi: 10.1523/JNEUROSCI.2306-10.2010
847
848 Gutman, A.R., Yang, Y., Ressler, K.J. and Davis, M. (2008). The role of neuropeptide Y in the
849 expression and extinction of fear-potentiated startle. J. Neurosci. 28, 12682-12690. doi:
850 10.1523/JNEUROSCI.2305-08.2008.
851
852 Heilig, M., Koob, G.F., Ekman, R. and Britton, K.T. (1994). Corticotropin-releasing factor and
853 neuropeptide Y: role in emotional integration. Trends Neurosci. 17, 80-85.
854
855 Hill, M.N., Hillard, C.J. and McEwen, B.S. (2011). Alterations in corticolimbic dendritic
856 morphology and emotional behavior in cannabinoid CB1 receptor-deficient mice parallel the
857 effects of chronic stress. Cereb. Cortex 21, 2056-2064. doi: 10.1093/cercor/bhq280
858
859 Holliday, N.D., Lam, C.W., Tough, I.R. and Cox, H.M. (2005). Role of the C terminus in
860 neuropeptide Y Y1 receptor desensitization and internalization. Mol. Pharmacol. 67, 655-664.
861
862 Humpel, C. (2015). Organotypic brain slice cultures: A review. Neuroscience 305, 86-98. doi:
863 10.1002/cpim.59
864
865 Ishida, A., Kameshita, I., Okuno, S., Kitani, T. and Fujisawa, H. (1995). A novel highly specific
866 and potent inhibitor of calmodulin-dependent protein kinase II. Biochem. Biophys. Res. Commun.
867 212, 806-812.
868

869   Kim, H., Kim, E., Park, M., Lee, E. and Namkoong, K. (2013) Organotypic hippocampal slice
870   culture from the adult mouse brain: a versatile tool for translational neuropsychopharmacology.
871   Prog. Neuropsychopharmacol. Biol. Psychiatry 41, 36-43. doi: 10.1016/j.pnpbp.2012.11.004
872

873   Lin, CH, Yeh SH, Leu TH, Chang WC, Wang ST and Gean PW. (2003). Identification of
874   calcineurin as a key signal in the extinction of fear memory. J. Neurosci. 23, 1574-1579.
875

876   Liu, J., Farmer JD Jr, Lane WS, Friedman J, Weissman I and Schreiber SL. (1991). Calcineurin is
877   a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. Cell 66, 807-815.
878

879   Lu, V.B., Colmers, W.F. and Smith, P.A.(2009). Long-term effects of brain-derived neurotrophic
880   factor on the frequency of inhibitory synaptic events in the rat superficial dorsal horn.
881   Neuroscience 161, 1135-1143. doi: 10.1016/j.neuroscience.2009.04.030.
882

883   Lu, V.B., Moran TD, Balasubramanyan S, Alier KA, Dryden WF, Colmers WF and Smith PA.
884   (2006). Substantia Gelatinosa neurons in defined-medium organotypic slice culture are similar to
885   those in acute slices from young adult rats. Pain 121, 261-275.
886

887   Mackay, J.P., Bompolaki M, DeJoseph MR, Michaelson SD, Urban JH and Colmers WF. (2019).
888   NPY2 Receptors Reduce Tonic Action Potential-Independent GABAB Currents in the Basolateral
889   Amygdala. J. Neurosci. 2019; 10.1523/Jneurosci.2226-18.2019 .
890

891   Marley, A., Choy, R.W. and von Zastrow, M. (2013). GPR88 reveals a discrete function of
892   primary cilia as selective insulators of GPCR cross-talk. PLoS One 8, e70857. doi:
893   10.1371/journal.pone.0070857.
894

895   McDonald, A.J. (1985). Immunohistochemical identification of gamma-aminobutyric acid-
896   containing neurons in the rat basolateral amygdala. *Neurosci. Lett.* **53**, 203-207.
897

898   McDonald, A.J., Beitz AJ, Larson AA, Kuriyama R, Sellitto C and Madl JE. (1989). Co-
899   localization of glutamate and tubulin in putative excitatory neurons of the hippocampus and
900   amygdala: an immunohistochemical study using monoclonal antibodies. *Neuroscience* **30**, 405-
901   421.
902

903   McEwen, B.S. (2003). Mood disorders and allostatic load. *Biol. Psychiatry* **54**, 200-207.
904

905   Michel, M.C., Beck-Sickinger A, Cox H, Doods HN, Herzog H, Larhammar D, Quirion R,
906   Schwartz T and Westfall T. (1989). XVI. International Union of Pharmacology recommendations
907   for the nomenclature of neuropeptide Y, peptide YY, and pancreatic polypeptide receptors.
908   Pharmacol. Rev. 50, 143-150.
909

910   Mineur, Y.S., Taylor, S.R. and Picciotto, M.R. (2014). Calcineurin downregulation in the
911   amygdala is sufficient to induce anxiety-like and depression-like behaviors in C57BL/6J male
912   mice. Biol. Psychiatry 75, 991-998. doi: 10.1016/j.biopsych.2014.03.009.
913

914   Mitra, R., Jadhav, S., McEwen, B.S., Vyas, A. and Chattarji, S. (2005) Stress duration modulates
915   the spatiotemporal patterns of spine formation in the basolateral amygdala. *Proc. Natl. Acad. Sci.*
916   *U. S. A.* **102**, 9371-9376.
917
918   Molosh, A.I., Sajdyk TJ, Truitt WA, Zhu W, Oxford GS and Shekhar A. (2013). NPY Y1
919   receptors differentially modulate GABAA and NMDA receptors via divergent signal-transduction
920   pathways to reduce excitability of amygdala neurons. Neuropsychopharmacology 38, 1352-1364.
921   doi: 10.1038/npp.2013.33.
922
923   Padival, M.A., Blume, S.R. and Rosenkranz, J.A. (2013) Repeated restraint stress exerts different
924   impact on structure of neurons in the lateral and basal nuclei of the amygdala. Neuroscience 246,
925   230-242. doi: 10.1016/j.neuroscience.2013.04.061
926
927   Pape, H.C. and Pare, D. (2010). Plastic synaptic networks of the amygdala for the acquisition,
928   expression, and extinction of conditioned fear. Physiol. Rev. 90, 419-463. doi:
929   10.1152/physrev.00037.2009
930
931   Paxinos, G. and Watson, C. (1986). The rat brain in stereotaxic coordinates (Academic Press,
932   Sydney;
933
934   Phelps, E.A. and LeDoux, J.E. (2005). Contributions of the amygdala to emotion processing: from
935   animal models to human behavior. Neuron 48, 175-187.
936
937   Rainnie, D.G., Bergeron R., Sajdyk T., Patil M., Gehlert D. and Shekhar A. (2004).
938   Corticotrophin releasing factor-induced synaptic plasticity in the amygdala translates stress into
939   emotional disorders. J. Neurosci. 24, 3471-3479.
940
941   Roozendaal, B., McEwen, B.S. and Chattarji, S. (2009). Stress, memory and the amygdala. *Nat.*
942   *Rev. Neurosci.* **10**, 423-433. doi: 10.1038/nrn2651
943
944   Sajdyk, T.J., Johnson PL, Leitermann RJ, Fitz SD, Dietrich A, Morin M, Gehlert DR, Urban JH,
945   and Shekhar A. (2008). Neuropeptide Y in the amygdala induces long-term resilience to stress-
946   induced reductions in social responses but not hypothalamic-adrenal-pituitary axis activity or
947   hyperthermia. J. Neurosci. 28, 893-903 (2008). doi: 10.1523/JNEUROSCI.0659-07.2008.
948
949   Sajdyk, T.J., Fitz, S.D. and Shekhar, A. (2006). The role of neuropeptide Y in the amygdala on
950   corticotropin-releasing factor receptor-mediated behavioral stress responses in the rat. Stress 9,
951   21-28.
952
953   Sajdyk, T.J., Schober, D.A. and Gehlert, D.R. (2002a). Neuropeptide Y receptor subtypes in the
954   basolateral nucleus of the amygdala modulate anxiogenic responses in rats. Neuropharmacology
955   43, 1165-1172.
956
957   Sajdyk, T.J., Schober, D.A., Smiley, D.L. and Gehlert, D.R. (2002b). Neuropeptide Y-Y2
958   receptors mediate anxiety in the amygdala. Pharmacol. Biochem. Behav. 71, 419-423.
959

Sajdyk, T.J., Shekhar, A. and Gehlert, D.R. (2004). Interactions between NPY and CRF in the
amygdala to regulate emotionality. Neuropeptides 38, 225-234.

Schmeltzer SN, Herman JP, **Sah R**. (2016) Neuropeptide Y (NPY) and posttraumatic stress
disorder (PTSD): A translational update. Exp Neurol. 284:196-210. Doi:
10.1016/j.expneurol.2016.06.020

Selcher, J.C., Xu, W., Hanson, J.E., Malenka, R.C. and Madison, D.V. (2012). Glutamate receptor
subunit GluA1 is necessary for long-term potentiation and synapse unsilencing, but not long-term
depression in mouse hippocampus. Brain Res. 1435, 8-14. doi: 10.1016/j.brainres.2011.11.029.

Sholl, D.A. (1953). Dendritic organization in the neurons of the visual and motor cortices of the
cat. J. Anat. 87, 387-406.

Silveira Villarroel, H., Bompolaki M, Mackay JP, Miranda Tapia AP, Michaelson SD,
Leitermann RJ, Marr RA, Urban JH and Colmers WF. (2018). NPY Induces Stress Resilience via
Downregulation of Ih in Principal Neurons of Rat Basolateral Amygdala. J. Neurosci. 38, 4505-
4520. doi: 10.1523/JNEUROSCI.3528-17.2018

Soll, R.M., Dinger, M.C., Lundell, I., Larhammer, D. and Beck-Sickinger, A.G. (2001). Novel
analogues of neuropeptide Y with a preference for the Y1-receptor. Eur. J. Biochem. 268, 2828-
2837.

Stanić D, Mulder J, Watanabe M, Hökfelt T. (2011). Characterization of NPY Y2 receptor protein
expression in the mouse brain. II. Coexistence with NPY, the Y1 receptor, and other
neurotransmitter-related molecules. J Comp Neurol. 519:1219-1257. doi: 10.1002/cne.22608.

Stoppini, L., Buchs, P.A. and Muller, D. (1991). A simple method for organotypic cultures of
nervous tissue. J. Neurosci. Methods 37, 173-182.

Taylor, A.L. (2012). What we talk about when we talk about capacitance measured with the
voltage-clamp step method. J. Comput. Neurosci. 32, 167-175 (2012). doi: 10.1007/s10827-011-
0346-8.

Vyas, A., Jadhav, S. and Chattarji, S. (2006). Prolonged behavioral stress enhances synaptic
connectivity in the basolateral amygdala. Neuroscience 143, 387-393.

Vyas, A., Mitra, R., Shankaranarayana Rao, B.S. and Chattarji, S. (2002). Chronic stress induces
contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. J. Neurosci.
22, 6810-6818.

Vyas, A., Pillai, A.G. and Chattarji, S. (2004). Recovery after chronic stress fails to reverse
amygdaloid neuronal hypertrophy and enhanced anxiety-like behavior. Neuroscience 128, 667-
673.
1002

1003   Wahlestedt C, Yanaihara N and Håkanson R. (1986). Evidence for different pre-and post-
1004   junctional receptors for neuropeptide Y and related peptides. Regul Pept. 13:307-318.
1005
1006   Yehuda R, Brand S, Yang RK. (2006). Plasma neuropeptide Y concentrations in combat exposed
1007   veterans: relationship to trauma exposure, recovery from PTSD, and coping. Biol Psychiatry.
1008   59:660-663.
1009
1010   Zhou Z, Zhu G, Hariri AR, Enoch MA, Scott D, Sinha R, Virkkunen M, Mash DC, Lipsky RH,
1011   Hu XZ, Hodgkinson CA, Xu K, Buzas B, Yuan Q, Shen PH, Ferrell RE, Manuck SB, Brown SM,
1012   Hauger RL, Stohler CS, Zubieta JK, Goldman D. (2008) Genetic variation in human NPY
1013   expression affects stress response and emotion. Nature 452:997-1001