Evidence for reduced CTX-M carriage in cattle-associated *Escherichia coli* at

low temperatures and on publicly accessible farmland: implications for

surveillance and potential for farm-to-human transmission

Hannah Schubert[1], Jacqueline Findlay[2], Katy Morley[1], Emma F. Puddy[1,2],

Robert Arbon[3], Virginia C. Gould[1,2], Oliver Mounsey[2], Madeleine Evans[1], Gwen

M. Rees[1], David C. Barrett[1], Katy M. Turner[1], Tristan A. Cogan[1], Matthew B.

Avison[2]*, Kristen K. Reyher[1]*

1) Bristol Veterinary School, University of Bristol

2) School of Cellular and Molecular Medicine, University of Bristol

3) Jean Golding Institute, University of Bristol

* Corresponding authors: kristen.reyher@bristol.ac.uk; bimba@bristol.ac.uk

Running title: CTX-M in cattle-associated *E. coli*

Abstract

Little is known about the drivers of critically important antibacterial resistance in species with zoonotic potential present on farms (e.g. CTX-M β-lactamase-positive *Escherichia coli*). There is also debate about the influence of farms on the circulation of resistance in local human populations. This was a two-year surveillance study on 53 dairy farms. *E. coli* positive for bla_{CTX-M} were detected in 224/4145 (5.4%) of all samples from faecally-contaminated sites. *E. coli* positive for bla_{CTX-M} were more prevalent (98/631; 15.5%) in calf samples and less prevalent (12/630; 1.9%) in samples collected from pastureland, including publicly accessible sites. Multilevel, multivariable logistic regression showed antibiotic dry cow therapeutic choice to be associated with risk of bla_{CTX-M} positivity, including use of cefquinome or framycetin; 74% of bla_{CTX-M}-positive *E. coli* were framycetin-resistant. Low temperature was associated with low risk of bla_{CTX-M} positivity. This was additional to the effect of temperature on total *E. coli* density, a finding with profound implications for surveillance. There was no evidence that study farms had a significant impact on circulating bla_{CTX-M} plasmids in the local human population: across 296 fully sequenced *E. coli* isolates, two cattle isolates shared bla_{CTX-M} plasmids with eight urinary isolates collected in parallel.

Introduction

Antimicrobial resistance (AMR) - and particularly antibacterial resistance (ABR) - is a significant global challenge. Many countries are implementing plans to reduce the use of antibacterial drugs (ABs) in food-producing animals. For example, the most recent UK five-year National Action Plan includes a target to reduce AB use (ABU) in the treatment of food-producing animals by 25% (1). In Europe, AB sales for food-producing animals fell by 20% from 2011 to 2016 (2). In the UK dairy industry, overall ABU dropped from 24 mg/kg in 2015 to 17 mg/kg in 2017(3, 4). In 2018, additional industry-led policies were enforced in the UK that aim to almost eliminate the use of highest priority critically important antimicrobials (HP-CIAs) such as third- and fourth-generation cephalosporins (3GCs and 4GCs) and fluoroquinolones on dairy farms. One reason for reducing ABU in farming is the belief that such measures will reduce the prevalence of ABR bacteria carried by farm animals. However, there is a need for better data on drivers of ABR in farming. More granularity of understanding is required concerning the risks of using individual ABs and other management practices. This is especially important in terms of drivers of HP-CIA resistance. One key focus is on 3GC resistance in *Escherichia coli*, a species commonly found in animal faeces and considered one of the most significant potential zoonotic threats to humans (5).

3GC-resistance is increasingly common in *E. coli* causing infections in humans (6) and is also found in farmed and domestic animals around the world (7). The production of CTX-M (an extended-spectrum β-lactamase) is the most common mechanism of 3GC-resistance in *E. coli* in humans in the UK; for example, in a recent study of urinary *E. coli* from humans in South West England, 82.2% of 3GC-resistant isolates carried *bla*CTX-M (8).

The objective of this study was to describe the prevalence of *E. coli* carrying bla_{CTX-M}

found in faecally contaminated environments of dairy cattle in a geographically

restricted population of UK dairy farms in South West England. Furthermore, this

study investigated ABU and management practice risk factors for the presence of

such *E. coli.* Finally, this study used extensive molecular epidemiology based on

whole genome sequencing (WGS) to help explain risk factors identified and to

investigate evidence for transmission of bla_{CTX-M}-encoding plasmids between farm-

and human-associated *E. coli* collected in parallel in a relatively small (50 x 50 km)

region.

Materials and Methods

Farm recruitment and ethical approval

A convenience sample of 53 dairy farms was recruited through personal contacts,

local veterinary practices and milk processors. Details of the study population are

presented in Supplementary. Of these, 43 farms were in a 50 x 50 km area defined

based on the locations of 146 general practices that referred routine urine samples

from human patients to the microbiology reference lab at Severn Pathology,

Southmead Hospital (8). A further 10 study farms were clustered in a separate

region in South West England. All farmers gave fully informed consent to participate

in the study. Ethical approval was obtained from the University of Bristol's Faculty of

Health Sciences Research Ethics Committee (ref 41562).

Farm sampling and sample processing

Farms were visited monthly between January 2017 and December 2018. Samples

were collected using sterile overshoes (over-boot socks) traversing farm areas.

Where access was restricted (e.g. for pens containing single or pairs of calves),

samples were collected directly from the ground using gloved hands. Details of the

six types of samples collected are in Supplementary; these represent faecally-

contaminated environments representative of milking cows (Adult), cows between

periods of lactation (Dry Cow), heifers after weaning (Heifer), heifers before weaning

(Calf). Samples of these types collected from pastureland were designated as such

for separate analysis (Pasture) which also included samples collected from publicly

accessible land on the farm (Footpath). Samples were refrigerated from collection to

processing. They were transferred into individual labelled sterile stomacher bags and

suspended in 10 mL/g of phosphate buffered saline (PBS Dulbecco A; Oxoid,

Basingstoke, UK). Samples were then mixed for one min in a stomacher (Stomacher

400, Seward, Worthing, UK). Samples were mixed 50:50 with 100% sterile glycerol

and aliquots stored at -80°C.

Microbiology and WGS analysis

Twenty microlitres of sample (diluted 1:10) were spread onto tryptone bile X-

glucuronide agar (TBX; Scientific Laboratory Supplies); 20 µL of undiluted sample

were spread onto TBX agar containing 16 mg/L cephalexin. Plates were incubated at

37°C, and the number of blue colonies (*E. coli*) counted. Samples yielding no *E. coli*

colonies on antibiotic-free agar were excluded from further analysis. Up to five *E. coli*

isolates from each cephalexin (16 mg/L) TBX agar plate were transferred onto

cefotaxime (CTX, 2 mg/L) TBX agar. Concentrations were chosen as those which

define clinically relevant resistance in humans according to EUCAST (9). Confirmed

CTX-resistant isolates were subjected to multiplex PCR to detect bla_{CTX-M} (testing for

bla_{CTX-M} groups 1, 2, 8, 9 and 25; [10]). WGS was performed by MicrobesNG

(https://microbesng.uk/) on a HiSeq 2500 instrument (Illumina, San Diego, CA, USA)

using 2x250 bp paired end reads. Reads were trimmed using Trimmomatic (11),

assembled into contigs using SPAdes (12) 3.13.0

(http://cab.spbu.ru/software/spades/) and contigs were annotated using Prokka (13).

Resistance genes, plasmid replicon types, sequence types and fim types were

assigned using the ResFinder (14), PlasmidFinder (15), MLST (16) 2.0 and

FimTyper on the Center for Genomic Epidemiology

(http://www.genomicepidemiology.org/) platform.

Risk factor data analysis

The risk factors examined fall into four categories: farm management, ABU, sample

characteristics and meteorological. Four management practice questionnaires were

developed (details provided in Supplementary). ABU was extracted from prescribing

and sales data between Jan 2016 to Dec 2018 obtained from veterinary practices

servicing the study farms. For two farms, sales data were not available and on-farm

records were used. The full method for obtaining, cleaning and processing ABU data

can be found in Supplementary. Local meteorological data were extracted from

publicly available UK Met Office data

(https://www.metoffice.gov.uk/pub/data/weather/uk/climate/stationdata/yeoviltondata.

txt).

Sample processing and data analysis workflows are illustrated in **Figure S1**. All data

analysis was performed using R (https://www.r-project.org/). Two modelling

approaches were used: 1) variable selection via univariable screening and stepwise

model selection with a multilevel, multivariable logistic regression model and 2) a

regularised Bayesian model. Both were used to analyse risk factors associated with

bla_{CTX-M} *E. coli* positivity. A sensitivity analyses was performed to test for

measurement bias, to account for the fact that bla_{CTX-M} is more likely to be found in a

sample if there is a higher density of bacterial colony forming units. Further details of

variable selection and development of the models and model checking are provided

in Supplementary.

Results

Farm- and sample-level risk factors for bla_{CTX-M} *E. coli* positivity

4581 samples were collected: 4145 were positive for growth of *E. coli* on non-

selective agar. Of these, 384/4145 (9.3%) samples representing 47/53 (88.7%) of

farms were positive for growth of *E. coli* that were resistant to cefotaxime. Overall,

5.4% (224/4145) of samples representing 42/53 (79.2%) of farms contained

cefotaxime-resistant *E. coli* confirmed to carry bla_{CTX-M} using PCR. Positivity for

bla_{CTX-M} *E. coli* was three times higher in Calf samples (98/631 [15.5%] of samples)

than overall (**Table 1**).

A separate risk factor analysis using only Calf data was performed, given the high

positivity rate for bla_{CTX-M} *E. coli* in these samples. One farm-level fixed effect and

three sample-level fixed effects were retained in the final model (**Table S1**, **Table 2**).

The use of cefquinome or framycetin dry cow therapies were both associated with

increased risk of bla_{CTX-M} *E. coli* positivity, as was higher average monthly

temperature. Plotting sample-level positivity for *E. coli* carrying bla_{CTX-M} versus

average monthly temperature revealed that the relationship between positivity and

temperature was primarily driven by low bla_{CTX-M} *E. coli* positivity rates in months

where the average temperature was below 10°C (**Figure 1A**).

Risk factor analysis was next performed for the full dataset. One farm-level fixed

effect and three sample-level fixed effects were retained in the final model (**Table**

S2, **Table 3**). Interestingly, this model revealed that bla_{CTX-M} *E. coli* was less likely to

be found in samples obtained from pastureland, which includes publicly accessible

farmland (Footpaths) compared with other sample types. Analysis of the full dataset

confirmed what was seen with the Calf dataset: that higher average monthly

temperature was associated with an increased risk of bla_{CTX-M} *E. coli* positivity.

Again, visualisation of the data confirmed that this was primarily driven by a

reduction in bla_{CTX-M} *E. coli* positivity rate in months with an average temperature

below 10°C (**Figure 1B**).

A Bayesian logistic regression model was also constructed in which the effect of total

farm ABU and specifically total 3GC and 4GC use were tested as predictors for

bla_{CTX-M} *E. coli* positivity in the total dataset, with 102 potential confounders. The

impacts of temperature (Odds Ratio 1.71 [1.42, 2.08]) and of samples being

collected from pastureland (Odds Ratio 0.51 [0.22, 1.02]) on bla_{CTX-M} *E. coli* positivity

were also retained in this alternative model (**Table S3**).

Defining sample-level positivity for bla_{CTX-M} *E. coli* depends on finding bla_{CTX-M} using

PCR in *E. coli* colonies that have grown on agar containing cefotaxime. If bla_{CTX-M} *E.*

coli in a sample exist at such a low density that they are not detected using selective

agar, the sample will be falsely identified as negative for bla_{CTX-M} *E. coli*. This impact

of bacterial density on assay sensitivity/specificity is an important consideration in

the context of the finding that bla_{CTX-M} positivity is low at low temperatures. To

account for this, the logistic link function was adjusted (see Supplementary). This

only modestly altered the effect sizes or the p values for the risk factors (**Figure S2**)

confirming that the effect of low temperature on bla_{CTX-M} *E. coli* positivity was

additional to its effect on *E. coli* prevalence. All values in **Tables 2**, **3** and **S3** come

from models with this adjusted logistic link function applied.

Molecular analysis and evidence of limited zoonotic transmission of *bla*CTX-M-

encoding plasmids

Based on WGS for 115 *bla*CTX-M *E. coli* isolates from farm samples, 77.4%, 84.3%

and 81.7% also carried genes encoding resistance to framycetin, streptomycin or

tetracycline, respectively. In contrast, 14.7% had trimethoprim resistance genes,

11.3% carried phenicol resistance genes and only 8.7% had plasmid-mediated

fluoroquinolone resistance genes. Thirty-seven out of 107 *bla*CTX-M-positive *E. coli*

isolates from farm samples were found to carry *bla*CTX-M variants also seen amongst

189 *bla*CTX-M-positive urinary *E. coli* cultured from people living in the same 50 x 50

km region during the same time period (8). By filtering sequenced isolates by their

*bla*CTX-M variants and plasmid replicon types, plasmids that were almost identical in

farm and human isolates were identified. One plasmid type, found in a single

sequence type (ST)345 farm isolate, harboured *bla*CTX-M-1 carried on an IncI1-ST3

plasmid. BLAST analysis of the *bla*CTX-M-1 contig showed that it was most similar to

part of an unpublished ~106 kb plasmid - pTC_N40607 (GenBank Accession No.

CP007651) - found in *E. coli* obtained from meat/cattle isolates in the USA. Mapping

of the ST345 farm isolate sequencing reads against pTC_N40607 showed it

exhibited 100% coverage and 97.5% sequence identity. Six human urinary *E. coli*

isolates representing STs 23, 127, 131, 141 and 2015 harboured *bla*CTX-M-1 on an

IncI1-ST3 plasmid that exhibited 99.4-100% coverage and 96.4-98.7% identity when

sequence reads were mapped to pTC_N40607.

Another plasmid type - again obtained from a single farm isolate, in this case of ST58

- exhibited 100% coverage and 98.5% identity by read mapping to a published IncK

plasmid pCT (GenBank Accession No. NC_014477). pCT is ~94 kb and is known to

harbour $bla_{CTX-M-14}$; pCT-like plasmids have been reported in both human and

veterinary *E. coli* isolates across three continents (17, 18). Amongst human urinary *E.*

coli isolates found in this study, two also carried pCT-like plasmids. Both isolates were

the pandemic clone ST131, and their pCT-like plasmids exhibited 96.4 or 97.2%

identity and 100% coverage to pCT.

Discussion

Prevalence of bla_{CTX-M} *E. coli* and impact of temperature - implications for

surveillance studies

This study is unique in its scale: extensive management practice and ABU data

along with multiple samples from multiple farms were collected over multiple time

points across a two-year period. Overall, 224/4145 (5.4%) of samples were positive

for *E. coli* carrying bla_{CTX-M} with considerable farm- and sample-level variation

(**Tables 1-3**). Of studies using similar methodology (phenotypic selection followed by

PCR analysis), Snow et al. (2012) (19) found 17/48 (35%) of randomly selected UK

dairy farms were positive for bla_{CTX-M} *E. coli*, whereas Mollenkomf et al. (2012) (20)

found 5/25 (20%) of farms in Ohio to be positive. This study found 42/53 (79%) of

farms to be positive, although many samples were collected each month over two

years, hence the chances of finding a positive sample on each farm may have been

greater than in these earlier point-prevalence studies. To further elucidate this, farm-

level positivity for bla_{CTX-M} *E. coli* was plotted on a month-by-month basis (**Figure 2**)

and revealed the highest prevalence for a single monthly survey to be 22.5%.

The prevalence of bla_{CTX-M} *E. coli* varied across sample types in this study: although

prevalence was low overall (5.4%), prevalence was higher (15.5%) in Calf samples.

In contrast, Horton et al. 2016 (21) reported a very high prevalence (>90%) of bla_{CTX-}

$_M$ *E. coli* in samples taken from faecal pats, even from adult milking cows. Brunton et

al. (2014) (22) reported a prevalence of 50% in calves. The finding of relatively low

prevalence in this study could be due to the large number of samples collected,

particularly over winter, given low temperature was associated with low bla_{CTX-M} *E.*

coli positivity (**Fig. 1A; 1B**). Indeed, the observation that average monthly

temperature had a significant effect on bla_{CTX-M} *E. coli* positivity highlights problems

with studies where a single time-point or sampling season is used. This has

potentially profound implications for surveillance studies performed to identify risk

factors, to identify general trends in ABR levels, to benchmark farms or to test the

effects of policy interventions. **Figure 2** shows the stark impact of this in real terms:

bla_{CTX-M} *E. coli* positivity at farm level was zero and 1.9%, respectively in February

and March, the coldest months of the year (based on average temperature). Whilst

average annual temperature found at locations across an entire continent has

previously been shown to impact average ABR levels at those locations (23), the

finding that periods of low temperatures were associated with lower prevalence of a

dominant cause of HP-CIA resistance at a given location during the course of a year

is particularly important; this observation also leads to concern about the impact of

climate change - and especially increasing temperatures - on attempts to reduce

ABR. Whilst temperature was associated with the total number of *E. coli* found in

each sample, this was accounted for using a novel measurement error method

incorporated into the model; as such, the effect of temperature on bla_{CTX-M} *E. coli*,

whilst in part driven by the effect on total *E. coli* number, also had an independent

association suggestive of a temperature-dependent fitness burden of carrying bla_{CTX-M}.

In terms of sample-level effects, there were clear differences in the risk of

encountering bla_{CTX-M} *E. coli* at different sites on a farm (e.g. 15.5% in Calf samples,

4.1% in Adult samples). Watson et al. (2012) (24) also found that CTX-M prevalence

was much higher in calves, but Horton et al. (2016) (21) did not report such a

difference; however, bla_{CTX-M} status in this latter study was presumptive based on the

phenotypic identification of cefotaxime resistance, which would make these results

less comparable with those presented here. In other farmed species, Agerso et al.

(2011) (25) found a prevalence of bla_{CTX-M} *E. coli* carriage of approximately 7% in

Danish slaughter pigs and Randall et al. (2010) (26) found a bla_{CTX-M} *E. coli*

prevalence of 3.6% in UK broiler chickens and turkeys. Various studies have

identified much higher prevalence in chicken meat, but this could be due to cross-

contamination at slaughter and in the food chain (27, 28).

Studies examining the prevalence of bla_{CTX-M} *E. coli* in human populations have

shown mixed results. Luvsansharav et al. (2012) (29) found a prevalence of 65.7%

amongst commensal isolates in Thailand. In the UK, a study across four regions

reported commensal faecal carriage of bla_{CTX-M} *E. coli* to be approximately 7% (30).

A recent analysis of human urinary samples from the same region as the farms

surveyed in this study gave a sample-level prevalence of bla_{CTX-M} *E. coli* of

approximately 5% (8). It should be noted that all farm samples in the present study

were from faecally contaminated sites, not individual animals, and so it is possible

that the number of animals carrying bla_{CTX-M} *E. coli* was much lower than the

reported sample-level prevalence. Direct comparison with human and other farm

animal carriage studies should therefore be made with caution.

AB contamination of colostrum as a possible driver of bla_{CTX-M} *E. coli* positivity

in dairy calves - evidence of direct and co-selection

It has been shown experimentally that waste (AB-contaminated) milk feeding to

calves increases faecal excretion of ABR bacteria (31). This practice is reducing on

UK dairy farms and, in the analysis presented here, waste milk feeding was not

associated with an increased risk of finding bla_{CTX-M} *E. coli*. In contrast, the choice of

dry cow therapy (an antibacterial preparation inserted into a cow's udder between

lactations to help treat or prevent mastitis) was associated with bla_{CTX-M} *E. coli*

positivity in Calf samples. Colostrum management is a hugely important part of early

life for most farmed mammals and is universally encouraged in dairy farming. In this

study, cefquinome (a 4GC) dry cow therapy was most significantly associated with

bla_{CTX-M} *E. coli* in Calf samples (**Table 2**), and it has previously been shown that

colostrum from cows given cefquinome dry cow therapy is heavily contaminated with

cefquinome (32). There was also a clear positive association between the usage of

framycetin as part of a dry cow therapy combination and the risk of finding bla_{CTX-M}

E. coli in Calf samples (**Table 2**). Whilst no work has been published on the

contamination of colostrum with framycetin, its use as a mastitis therapy for milking

cows results in identifiable residues in milk (33), so it is highly likely to also

contaminate colostrum. It is possible, therefore, that feeding of colostrum, which can

be contaminated with AB used for dry cow therapy, is a driver of bla_{CTX-M} *E. coli* in

calves. An alternative (or indeed an additional) explanation for this observed

association is that *E. coli* (a species known to be found in the udders of dairy cows

(34) that carry bla_{CTX-M} are selected within the udder during AB dry cow therapy and

contaminate colostrum alongside the AB used.

WGS showed that bla_{CTX-M} was co-located with framycetin resistance genes in

77.4% of 115 fully sequenced *E. coli* isolates across study farms. Accordingly, it is

hypothesised that framycetin dry cow therapy drives bla_{CTX-M} *E. coli* positivity in

calves because of co-selection of bacteria carrying bla_{CTX-M} and a framycetin

resistance gene, just as cefquinome use drives bla_{CTX-M} *E. coli* positivity by direct

selection of bla_{CTX-M}. Whilst regional differences in the ecology of circulating

resistance may mean that the specific observation made here is not universally

applicable, this is still an important "real-world" example of the frequently identified

laboratory phenomenon of co-selection and illustrates why policy changes to reduce

the usage of HP-CIAs such as cefquinome may not remove all selective pressure for

presence of HP-CIA resistance on farms. Snow et al. (2012) (19) also identified

overall use of 3/4GCs as a risk factor for bla_{CTX-M} *E. coli* presence on dairy farms.

Other studies have not made a link between the usage of framycetin and prevalence

of bla_{CTX-M} *E. coli*. However, it is not always clear whether other studies have

separated out different dry cow therapies since they have tended to focus on

systemic AB use.

Low risk of finding bla_{CTX-M} *E. coli* on publicly accessible farm sites and little

evidence of sharing with locally resident people

This study identified a lower chance of detecting bla_{CTX-M} *E. coli* in samples collected

on pastureland than elsewhere on the farm. Because pastureland may be more

affected by the elements, this finding may be partly linked with the finding that lower

average monthly temperature was associated with decreased risk of detecting

bla_{CTX-M} *E. coli*. Importantly, 395/630 (62.7%) of samples from pastureland were from

publicly accessible sites so the finding of a low positivity for bla_{CTX-M} *E. coli* on

pastureland led to the hypothesis that dairy farms are unlikely to be sources of

transmission of bla_{CTX-M} *E. coli* into the local human population. Analysis of fully

sequenced genomes of 296 bla_{CTX-M} *E. coli* from study farms and from urine samples

provided by people living in the same geographical range as the farms over the

same time period did not suggest any evidence of direct sharing of *E. coli* between

farms and the local human population (i.e. none were the same ST, carrying the

same plasmid with the same bla_{CTX-M} variant). However, two farm *E. coli* isolates

carried a bla_{CTX-M} plasmid almost identical to a plasmid circulating in the local human

population. One plasmid (pCT) is already known to be spread widely amongst *E. coli*

from humans and animals on three continents (Cottell 2011). Whilst unpublished, the

other plasmid (pTC_N40607) has been identified in *E. coli* in the USA so is likely to

be similarly widespread. Accordingly, whilst there is some evidence of shared

circulating plasmids, as reported in a number of recent studies (35-37), the overall

level of overlap between farm and human bla_{CTX-M} *E. coli* identified in this study was

very small and not suggestive of any novel or recent transmission events.

Acknowledgements

We wish to thank all the farmers and veterinary surgeons who participated in this

study. In addition, we would like to thank the milk processing companies who

supported our recruitment process and the veterinary students who helped with data

processing and sample collection. Genome sequencing was provided by

MicrobesNG (http://www.microbesng.uk), which is supported by the BBSRC (grant

number BB/L024209/1).

Funding

This work was funded by grant NE/N01961X/1 to M.B.A., K.K.R., K.M.T., T.A.C. and

D.C.B. from the Antimicrobial Resistance Cross Council Initiative supported by the

seven United Kingdom research councils. R.A. was supported by the Jean Golding

Institute for data-intensive research at the University of Bristol. G.M.R. was funded

by the Langford Trust.

Transparency declarations

D.C.B. was president of the British Cattle Veterinary Association 2018-19.

Otherwise, the authors declare no competing interests. Farming and veterinary

businesses who contributed data and permitted access for sample collection were

not involved in the design of this study or in data analysis and were not involved in

drafting the manuscript for publication.

Author Contributions

Conceived the Study: D.C.B., K.K.R., M.B.A.

Collection of Data: H.S., J.F., K.M., E.F.P., O.M., V.C.G., supervised by T.A.C.,

K.K.R., G.M.R., M.B.A.

Cleaning and Analysis of Data: H.S., J.F., R.A., V.C.G., E.F.P, M.E., supervised by

K.M.T., K.K.R., M.B.A.

Initial Drafting of Manuscript: H.S., K.K.R., M.B.A.

Corrected and Approved Manuscript: All Authors

Tables

Table 1. Prevalence of *E. coli* carrying *bla*CTX-M at farm and sample levels

Sample type			Farm level	Sample level
Overall	**Total sample size**		53	4145
	Total (%) positive for CTX-M-carrying *E. coli*		42 (79.2%)	224 (5.4%)
Adult	**Total sample size**		52	1835
	Total (%) positive for CTX-M-carrying *E. coli*		25 (48.1%)	76 (4.1%)
Dry Cow	**Total sample size**		46	282
	Total (%) positive for CTX-M-carrying *E. coli*		7 (15.2%)	8 (2.8%)
Calf	**Total sample size**		51	631
	Total (%) positive for CTX-M-carrying *E. coli*		33 (64.7%)	98 (15.5%)
Heifer	**Total sample size**		41	1235
	Total (%) positive for CTX-M-carrying *E. coli*		18 (44%)	40 (3.2%)
Pastureland	**Total sample size**		47	630
	Total (%) positive for CTX-M-carrying *E. coli*		8 (17%)	12 (1.9%)
Pastureland that is publicly accessible (Footpath)	**Total sample size**		41	395
	Total (%) positive for CTX-M-carrying *E. coli*		8 (20.0%)	11 (2.8%)

375 **Table 2**. Fixed effects from the multilevel, multivariable logistic regression model

376 performed on Calf samples

Risk factor	Odds ratio [95% confidence interval]	p
Use of cefquinome dry cow therapy in the last six months	4.12 [2.11, 8.25]	0.00003
Daily water trough cleaning	0.44 [0.29, 0.69]	0.0002
Average monthly temperature	1.57 [1.20, 2.06]	0.0008
Use of framycetin dry cow therapy in the last six months	1.91 [1.01, 3.61]	0.04

380 **Table 3**. Fixed effects from the multilevel, multivariable logistic regression model

381 performed on the full dataset

Risk factor	Odds ratio [95% confidence interval]	p
Sample taken from the environment of pre-weaned heifers	4.51 [3.25, 6.27]	0
Average monthly temperature	1.60 [1.34, 1.89]	0.00000001
Sample taken from pastureland	0.32 [0.17,0.61]	0.0000001
Feeding of maize silage	3.28 [1.50, 7.18]	0.002

Figures

Figure 1A. Average monthly temperature vs. presence (1) or absence (0) of CTX-M

in samples from pre-weaned calves. A multilevel, multivariable logistic regression

model revealed a positive association with increased temperature (p=0.0008).

Figure 1B. Average monthly temperature vs. presence (1) or absence (0) of CTX-M

in all samples of faecally contaminated dairy farm environments. A multilevel,

multivariable logistic regression model revealed a positive association with increased

temperature (p=0.00000001).

Figure 2. Percentage of farms positive for CTX-M by month and by average monthly

temperature representing a year during the middle period of this study. Samples

from calves have been excluded.

References

References

1.   https://www.gov.uk/government/collections/antimicrobial-resistance-amr-

information-and resources. 2019.

2.   ESVAC. European Surveillance of Veterinary Antimicrobial Consumption.

3.   Veterinary Medicines Directorate & Animal and Plant Health Agency. UK

Veterinary Antibiotic Resistance and Sales Surveillance. 2017.

4.   Veterinary Medicines Directorate & Animal and Plant Health Agency. UK

Veterinary Antibiotic Resistance and Sales Surveillance. 2015.

5.   Lazarus B, Paterson DL, Mollinger JL, Rogers BA. Do Human Extraintestinal

Escherichia coli Infections Resistant to Expanded-Spectrum Cephalosporins

Originate From Food-Producing Animals? A Systematic Review. Clinical Infectious

Diseases. 2014;60(3):439-52.

6.   Bevan ER, Jones AM, Hawkey PM. Global epidemiology of CTX-M β-

lactamases: temporal and geographical shifts in genotype. Journal of Antimicrobial

Chemotherapy. 2017;72(8):2145-55.

7.   Madec J-Y, Haenni M. Antimicrobial resistance plasmid reservoir in food and

food-producing animals. Plasmid. 2018;99:72-81.

8.   Findlay J GV, North P, Bowker KE, Williams OM, MacGowan, AP, Avison MB.

Characterisation of cefotaxime-resistant urinary Escherichia coli from primary care in

South-West England 2017-2018. J Antimicrob Chemother. 2019.

9.

http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_ta

bles/v_9.0_Breakpoint _Tables.pdf. 2019.

10.   Woodford N, Fagan EJ, Ellington MJ. Multiplex PCR for rapid detection of

genes encoding CTX-M extended-spectrum β-lactamases. Journal of Antimicrobial

Chemotherapy. 2005;57(1):154-5.

11.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30(15):2114-20.

12.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.

SPAdes: a new genome assembly algorithm and its applications to single-cell

sequencing. J Comput Biol. 2012;19(5):455-77.

13.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.

2014;30(14):2068-9.

14.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O,

et al. Identification of acquired antimicrobial resistance genes. Journal of

Antimicrobial Chemotherapy. 2012;67(11):2640-4.

15.   Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L,

et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid

multilocus sequence typing. Antimicrob Agents Chemother. 2014;58(7):3895-903.

16.   Wirth T, Falush D, Lan R, Colles F, Mensa P, Wieler LH, et al. Sex and

virulence in Escherichia coli: an evolutionary perspective. Mol Microbiol.

2006;60(5):1136-51.

17.   Cottell JL, Webber MA, Coldham NG, Taylor DL, Cerdeño-Tárraga AM,

Hauser H, et al. Complete sequence and molecular epidemiology of IncK epidemic

plasmid encoding blaCTX-M-14. Emerg Infect Dis. 2011;17(4):645-52.

18.   Dhanji H, Khan P, Cottell J, Piddock L, Zhang J, Livermore D, et al.

Dissemination of pCT-Like IncK Plasmids Harboring CTX-M-14 Extended-Spectrum

-Lactamase among Clinical Escherichia coli Isolates in the United Kingdom2012.

3376-7 p.

19.   Snow LC, Warner RG, Cheney T, Wearing H, Stokes M, Harris K, et al. Risk

factors associated with extended spectrum beta-lactamase Escherichia coli (CTX-M)

on dairy farms in North West England and North Wales. Preventive Veterinary

Medicine. 2012;106(3):225-34.

20.   Mollenkopf DF, Weeman MF, Daniels JB, Abley MJ, Mathews JL, Gebreyes

WA, et al. Variable within- and between-herd diversity of CTX-M cephalosporinase-

bearing Escherichia coli isolates from dairy cattle. Appl Environ Microbiol.

2012;78(13):4552-60.

21.   Horton RA, Duncan D, Randall LP, Chappell S, Brunton LA, Warner R, et al.

Longitudinal study of CTX-M ESBL-producing E. coli strains on a UK dairy farm.

Research in Veterinary Science. 2016;109:107-13.

22.   Brunton LA, Reeves HE, Snow LC, Jones JR. A longitudinal field trial

assesing the impact of feeding waste milk containing antibiotic residues on the

prevalence of ESBL-producing Escherichia coli in calves. Preventive Veterinary

Medicine. 2014;117(2):403-12.

23.   MacFadden DR, McGough SF, Fisman D, Santillana M, Brownstein JS.

Antibiotic Resistance Increases with Local Temperature. Nat Clim Chang.

2018;8(6):510-4.

24.   Watson E, Jeckel S, Snow L, Stubbs R, Teale C, Wearing H, et al.

Epidemiology of extended spectrum beta-lactamase E. coli (CTX-M-15) on a

commercial dairy farm. Veterinary Microbiology. 2012;154(3):339-46.

25.   Agersø Y, Aarestrup FM, Pedersen K, Seyfarth AM, Struve T, Hasman H.

Prevalence of extended-spectrum cephalosporinase (ESC)-producing Escherichia

coli in Danish slaughter pigs and retail meat identified by selective enrichment and

association with cephalosporin usage. Journal of Antimicrobial Chemotherapy.

2011;67(3):582-8.

26.   Randall LP, Clouting C, Horton RA, Coldham NG, Wu G, Clifton-Hadley FA, et

al. Prevalence of Escherichia coli carrying extended-spectrum β-lactamases (CTX-M

and TEM-52) from broiler chickens and turkeys in Great Britain between 2006 and

2009. Journal of Antimicrobial Chemotherapy. 2010;66(1):86-95.

27.   Overdevest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, et al.

Extended-spectrum β-lactamase genes of Escherichia coli in chicken meat and

humans, The Netherlands. Emerg Infect Dis. 2011;17(7):1216-22.

28.   Kola A, Kohler C, Pfeifer Y, Schwab F, Kühn K, Schulz K, et al. High

prevalence of extended-spectrum-β-lactamase-producing Enterobacteriaceae in

organic and conventional retail chicken meat, Germany. Journal of Antimicrobial

Chemotherapy. 2012;67(11):2631-4.

29.   Luvsansharav U-O, Hirai I, Nakata A, Imura K, Yamauchi K, Niki M, et al.

Prevalence of and risk factors associated with faecal carriage of CTX-M β-

lactamase-producing Enterobacteriaceae in rural Thai communities. Journal of

Antimicrobial Chemotherapy. 2012;67(7):1769-74.

30.   McNulty CAM, Lecky DM, Xu-McCrae L, Nakiboneka-Ssenabulya D, Chung

K-T, Nichols T, et al. CTX-M ESBL-producing Enterobacteriaceae: estimated

prevalence in adults in England in 2014. J Antimicrob Chemother. 2018;73(5):1368-

88.

31.   Awosile BB, Smith BA. Risk assessment modelling of fecal shedding caused

by extended-spectrum cephalosporin-resistant Escherichia coli transmitted through

waste milk fed to dairy pre-weaned calves. Journal of Dairy Science.

2017;100(12):9667-73.

32.   Bachmann J, Helmschrodt C, Richter A, Heuwieser W, Bertulat S. Residue

concentration of cefquinome after intramammary dry cow therapy and short dry

periods. Journal of Dairy Science. 2018;101(8):7540-50.

33.   Luo PJ, Zhang JB, Wang HL, Chen X, Wu N, Zhao YF, et al. Rapid and

Sensitive Chemiluminescent Enzyme Immunoassay for the Determination of

Neomycin Residues in Milk. Biomedical and Environmental Sciences.

2016;29(5):374-8.

34.   Bradley AJ, Green MJ. Adaptation of Escherichia coli to the bovine mammary

gland. J Clin Microbiol. 2001;39(5):1845-9.

35.   Pietsch M, Irrgang A, Roschanski N, Brenner Michael G, Hamprecht A, Rieber

H, et al. Whole genome analyses of CMY-2-producing Escherichia coli isolates from

humans, animals and food in Germany. BMC Genomics. 2018;19(1):601-.

36.   Valcek A, Roer L, Overballe-Petersen S, Hansen F, Bortolaia V,

Leekitcharoenphon P, et al. IncI1 ST3 and IncI1 ST7 plasmids from CTX-M-1-

producing Escherichia coli obtained from patients with bloodstream infections are

closely related to plasmids from E. coli of animal origin. Journal of Antimicrobial

Chemotherapy. 2019;74(8):2171-5.

37.   Ludden C, Raven KE, Jamrozy D, Gouliouris T, Blane B, Coll F, et al. One

Health Genomic Surveillance of Escherichia coli Demonstrates Distinct Lineages and

Mobile Genetic Elements in Isolates from Humans versus Livestock. MBio.

2019;10(1):e02693-18.