Reduced antibacterial drug resistance and *bla*CTX-M β-lactamase gene carriage

in cattle-associated *Escherichia coli* at low temperatures, at sites dominated

by older animals and on pastureland: implications for surveillance.

Hannah Schubert[1], Katy Morley[1], Emma F. Puddy[1,2], Robert Arbon[3], Jacqueline

Findlay[2], Oliver Mounsey[2], Virginia C. Gould[1,2], Lucy Vass[1,4], Madeleine Evans[1],

Gwen M. Rees[1], David C. Barrett[1], Katy M. Turner[1], Tristan A. Cogan[1], Matthew

B. Avison[2]*, Kristen K. Reyher[1]

1) Bristol Veterinary School, University of Bristol

2) School of Cellular and Molecular Medicine, University of Bristol

3) Jean Golding Institute, University of Bristol

4) Bristol Population Health Science Institute, University of Bristol

* Corresponding author: bimba@bristol.ac.uk

Running title: Resistance in cattle-associated *E. coli*

Abstract

Little is known about the drivers of critically important antibacterial resistance in species with zoonotic potential present on farms (e.g. CTX-M β-lactamase-positive *Escherichia coli*). Here, we collected samples, monthly over a two-year period, on 53 dairy farms in the South West of England, and data for 610 variables concerning antimicrobial usage, management practices and meteorological factors. We detected *E. coli* resistant to amoxicillin, ciprofloxacin streptomycin and tetracycline, respectively, in 2754/4145 (66%), 263/4145 (6%), 1475/4145 (36%) and 2874/4145 (69%) of all samples from faecally contaminated sites. *E. coli* positive for bla_{CTX-M} were detected in 224/4145 (5.4%) of samples. Multilevel, multivariable logistic regression showed antibiotic dry cow therapeutic choice (including use of cefquinome or framycetin) to be associated with increased odds of bla_{CTX-M} positivity. Low temperature was associated with reduced odds of bla_{CTX-M} *E. coli* positivity in samples and to reduced odds of finding *E. coli* resistant to each of the four test antibacterials. This was additional to the effect of temperature on total *E. coli* density. Furthermore, samples collected close to calves had increased odds of having *E. coli* resistance to each antibacterial or positive for bla_{CTX-M}. Samples collected on pastureland had reduced odds of having *E. coli* resistant to amoxicillin or tetracycline, and being positive for bla_{CTX-M}.

Importance

Antibacterial resistance poses a significant threat to human and animal health and global food security. Surveillance for resistance on farms is important for many reasons, including to track the impacts of interventions aimed at reducing the prevalence of resistance. In this epidemiological survey of dairy farm antibacterial

resistance, we show that local temperature, as it changes over the course of a year,

is associated with the prevalence of antibacterial resistant *E. coli*. Also, that

prevalence of resistant *E. coli* is higher in indoor environments and in environments

inhabited by young animals. These findings have profound implications for routine

surveillance and for surveys carried out for research. They provide important

evidence that sampling at a single time-point and/or single location on a farm is

unlikely to be adequate to accurately determine the status of the farm with regard to

the presence or number of resistant *E. coli*.

Introduction

Antimicrobial resistance - and particularly antibacterial resistance (ABR) - is a significant global challenge. Many countries are implementing plans to reduce the use of antibacterial drugs (ABs) in food-producing animals. For example, the most recent UK five-year National Action Plan includes a target to reduce AB use (ABU) in the treatment of food-producing animals by 25% (1). In Europe, AB sales for food-producing animals fell by 20% from 2011 to 2016 (2). In the UK dairy industry, overall ABU dropped from 24 mg/kg in 2015 to 17 mg/kg in 2018 (3, 4). In 2018, additional industry-led policies were enforced in the UK that aimed to almost eliminate the use of highest-priority critically important antimicrobials (HP-CIAs) such as third- and fourth-generation cephalosporins (3GCs and 4GCs) as well as fluoroquinolones on dairy farms. One reason for reducing ABU in farming is to reduce the prevalence of ABR bacteria carried by farm animals. However, there is a need for better data on drivers of ABR in farming. More granularity of understanding is required concerning the risks of using individual ABs and other management practices. This is especially important in terms of drivers of HP-CIA resistance. A focus among HP-CIAs is on 3GC and fluoroquinolone resistance in *Escherichia coli*, a species commonly found in animal faeces and considered one of the most significant potential zoonotic threats to humans (5).

3GC resistance is increasingly prevalent in *E. coli* causing infections in humans (6) and is also found in farmed and domestic animals around the world (7). The production of CTX-M (an extended-spectrum β-lactamase) is the most common mechanism of 3GC resistance in *E. coli* in humans in the UK; for example, in a recent study of urinary *E. coli* from humans in South West England, 82.2% of 3GC-resistant isolates carried *bla*$_{CTX-M}$ (8).

The objective of this study was to describe the prevalence of 3GC-resistant *E. coli*

carrying bla_{CTX-M} as well as *E. coli* resistant to amoxicillin, tetracycline, streptomycin

and the fluoroquinolone ciprofloxacin found in faecally contaminated environments of

dairy cattle in a geographically restricted population of dairy farms in South West

England. Furthermore, this study investigated environmental, ABU and management

practice risk factors for the presence of such *E. coli.*

Results

Prevalence and PCR characterisation of 3GC-resistant *E. coli* from dairy farms

4581 samples from faecally contaminated sites were collected from 53 dairy farms.

Samples were collected on each farm monthly between January 2017 and

December 2018. 4145 samples were positive for growth of *E. coli* on non-selective

agar. Of these, 384/4145 (9.3%) samples representing 47/53 (88.7%) of farms were

positive for growth of *E. coli* on agar containing the 3GC cefotaxime. From these,

1226 3GC-resistant isolates were taken forward for PCR testing for possible

cephalosporinase genes of interest (GOIs): bla_{CTX-M} (groups 1, 2, 8, 9 and 25),

bla_{CMY}, bla_{DHA} and bla_{SHV}. Over half (648/1226; 52.7%) of all isolates tested were

found to harbour bla_{CTX-M} genes. Of these, 547/648 (84.4%) were of group 1, 99/648

(15.3%) were of group 9, and, in one case, both gene groups were identified. Twelve

isolates harboured a bla_{CMY} gene – one alongside bla_{CTX-M} group 1 – and one isolate

was bla_{DHA-1}-positive. No isolates were positive for bla_{SHV} and the remaining

566/1226 (46.2%) isolates were PCR-negative for all GOIs. These isolates were

hypothesised to hyper-produce the chromosomally encoded AmpC β-lactamase;

some of these isolates have been characterised in detail in a separate study (9).

Farm- and sample-level risk factors for bla_{CTX-M} *E. coli* positivity

Based on PCR, carriage of bla_{CTX-M} was the most common mechanism of 3GC-resistance in *E. coli* from dairy farms in this study. Identifying management practice- and AMU-associated risk factors for bla_{CTX-M} *E. coli* positivity was therefore considered to be an important objective. Overall, 5.4% (224/4145) of samples representing 42/53 (79.2%) of farms contained 3GC-resistant *E. coli* confirmed to carry bla_{CTX-M} using PCR. Positivity for bla_{CTX-M} *E. coli* was three times higher in Calf samples (samples collected from the environments of calves) (98/631 [15.5%] of samples) than overall (**Table 1**).

Given the high positivity rate for bla_{CTX-M} *E. coli* in Calf samples, a separate risk factor analysis using only Calf data was performed. One farm-level fixed effect and three sample-level fixed effects were retained in the final multilevel, multivariable logistic regression model (**Table S1**, **Table 2**). The use of cefquinome or framycetin dry cow therapies were both associated with increased odds of bla_{CTX-M} *E. coli* positivity, as was higher average monthly temperature. Plotting sample-level positivity for *E. coli* carrying bla_{CTX-M} versus average monthly temperature revealed that the relationship between positivity and temperature was primarily driven by low bla_{CTX-M} *E. coli* positivity rates in months where the average temperature was below 10°C (**Figure 1A**).

Risk factor analysis was also performed for the full dataset. One farm-level fixed effect and three sample-level fixed effects were retained in the final model (**Table S2**, **Table 3**). Interestingly, this model revealed that bla_{CTX-M} *E. coli* was less likely to be found in samples obtained from pastureland, which included publicly accessible farmland (Footpaths) compared with other sample types. Analysis of the full dataset confirmed what was seen with the Calf dataset: higher average monthly temperature

was associated with an increased odds of bla_{CTX-M} *E. coli* positivity. Again,

visualisation of the data confirmed that this was primarily driven by a reduction in

bla_{CTX-M} *E. coli* positivity rate in months with an average temperature below 10°C

(**Figure 1B**).

A Bayesian logistic regression model was also constructed in which the effect of total

farm ABU and specifically total 3GC and 4GC use were tested as predictors for

bla_{CTX-M} *E. coli* positivity in the total dataset, with 102 potential confounders included.

The impact of temperature (odds ratio 1.71 [1.42, 2.05]) on bla_{CTX-M} *E. coli* positivity

was also retained in this alternative model (**Table S3**).

Defining sample-level positivity for bla_{CTX-M} *E. coli* is dependent upon finding bla_{CTX-M}

using PCR in *E. coli* colonies that have grown on agar containing cefotaxime. If

bla_{CTX-M} *E. coli* in a sample exist at such a low density that they are not detected

using selective agar, the sample will be falsely identified as negative for bla_{CTX-M} *E.*

coli. This impact of bacterial density on assay sensitivity is an important

consideration in the context of the finding that bla_{CTX-M} positivity is low at low

temperatures. To account for this, the logistic link function was adjusted (see

Supplementary). This only modestly altered the effect sizes and the p-values for the

risk factors (**Figure S2**), confirming that the effect of low temperature on bla_{CTX-M} *E.*

coli positivity was additional to its effect on *E. coli* prevalence. All values in **Tables 2**

and **3** come from models with this adjusted logistic link function applied.

Prevalence and risk factor analysis for *E. coli* resistant to other antibacterial

classes

All 4145 samples positive for growth of *E. coli* on non-selective agar were also tested

for detectable numbers of *E. coli* resistant to four non-cephalosporins: amoxicillin,

tetracycline, streptomycin and ciprofloxacin, the last being representative of the HP-

CIA class, the fluoroquinolones. Resistance to amoxicillin and tetracycline were the

most prevalent types of resistance found, with ciprofloxacin resistance being the

least commonly detected (**Table 4**).

Using a Bayesian logistic regression method, factors associated with the risk of a

sample being positive for *E. coli* resistant to each of the test ABs were identified from

the total dataset. As seen for bla_{CTX-M} *E. coli*, where and when the samples were

collected were more significantly associated with the odds of finding resistant *E. coli*

in a sample than farm-level management practices or ABU, with all four models

showing a positive association between average monthly temperature and the odds

of finding resistant *E. coli* in a sample (**Table 5**). Also consistent across all models

was the significance of sampling different areas of the farm. Again, as with bla_{CTX-M}

E. coli, environments of calves were more likely (as a proportion of all *E. coli* than

elsewhere on the farm) to harbour all four types of resistant bacteria. Samples

collected from pastureland were, like bla_{CTX-M} *E. coli*, negatively associated with the

presence of amoxicillin and tetracycline resistance (**Table 5**).

Full results for all variables tested can be found in **Table S4**. Re-running the models

with sceptical priors did not affect the results; only very small differences in the

model coefficients were observed (**Table S5**). Details of all model checking can be

found in Supplementary.

Discussion

Prevalence of bla_{CTX-M} positive *E. coli*

This study is unique in its scale: extensive management practice and ABU data along with multiple samples from multiple farms were collected monthly across a two-year period. Overall, 224/4145 (5.4%) of samples were positive for *E. coli* carrying bla_{CTX-M}. This compares well with previously calculated bla_{CTX-M} *E. coli* carriage of approximately 7% in Danish slaughter pigs (10) and 3.6% in UK broiler chickens and turkeys (11). Various studies have identified much higher prevalence in chicken meat, but this could be due to cross-contamination at slaughter and in the food chain (12, 13).

Studies examining the prevalence of bla_{CTX-M} *E. coli* in human populations have shown mixed results. A prevalence of 65.7% was found amongst commensal isolates in Thailand (14). In the UK, a study across four regions reported commensal faecal carriage of bla_{CTX-M} *E. coli* to be approximately 7% (15). A recent analysis of human urinary samples from the same region as the farms surveyed in this study gave a sample-level prevalence of bla_{CTX-M} *E. coli* of approximately 5% (8). It should be noted that all farm samples in the present study were from faecally contaminated sites, not individual animals, and so it is possible that the number of animals carrying bla_{CTX-M} *E. coli* was much lower than the reported sample-level prevalence. Direct comparison with human and other farm animal carriage studies should therefore be made with caution.

Impact of temperature on the odds of finding resistant and bla_{CTX-M}-positive *E. coli*

This study found 42/53 (79%) of farms to be positive, based on phenotypic analysis

and PCR, for bla_{CTX-M} *E. coli*. This was higher than seen in other using similar

methodology e.g. 17/48 (35%) of randomly selected UK dairy farms (16) and 5/25

(20%) of farms in Ohio (17) were positive. In the present study, samples were

collected each month over two years, hence the chances of finding a positive sample

on each farm may have been greater than in these earlier point-prevalence studies

(16,17). When farm-level positivity for bla_{CTX-M} *E. coli* was plotted on a month-by-

month basis (**Figure 2**) the highest prevalence for a single monthly survey was

22.5%, which fits more closely with these other studies.

Sample-level prevalence of bla_{CTX-M} *E. coli* was low overall (5.4%). This contrasts

with >90% (18) or 50% (19) of bla_{CTX-M} *E. coli* in samples taken from bovine faecal

pats. This difference could be due to the large number of samples collected here,

particularly over winter, given low temperature was associated with low bla_{CTX-M} *E.*

coli positivity (**Fig. 1A; 1B**). Indeed, the observation that average monthly

temperature had a significant effect on bla_{CTX-M} *E. coli* positivity (**Table 3**), as well as

positivity for resistance to amoxicillin, ciprofloxacin, streptomycin and tetracycline

(**Table 5**) highlights problems with studies where a single time-point or sampling

season is used. **Figure 2** shows the stark impact of this in real terms: bla_{CTX-M} *E. coli*

positivity and positivity for ciprofloxacin-resistant *E. coli* at farm level was zero in

February, the coldest month of the year (based on average temperature; **Fig 2A, B**).

Whilst average annual temperature found at locations across an entire continent has

previously been shown to impact average ABR levels at those locations (20), the

finding that periods of low temperatures were associated with lower prevalence of a

dominant cause of ABR - and particularly HP-CIA resistance at a given location

during the course of a year - is particularly important. This observation also leads to

concern about the impact of climate change - and especially increasing temperatures

- on attempts to reduce ABR. Whilst temperature was associated with the total

number of *E. coli* found in each sample, this was accounted for using a

measurement error method incorporated into the model; as such, the effect of

temperature on ABR or bla_{CTX-M}-positive *E. coli*, whilst in part driven by the effect on

total *E. coli* number, also had an independent association suggestive of a

temperature-dependent fitness burden of resistance.

High levels of ABR and bla_{CTX-M} positive *E. coli* in farm locations dominated by

young animals and low levels on pastureland

There were clear differences in the risk of encountering bla_{CTX-M} *E. coli* at different

sites on a farm (e.g. 15.5% in Calf samples, 4.1% in Adult samples). The Calf

environment was also much more likely to have *E. coli* resistant to amoxicillin,

tetracycline, streptomycin, and ciprofloxacin, so this seems to be a universal effect.

Other studies have also generally found high levels of resistance in samples

collected from or in the environment of younger calves (21-24). There may also be

an association with temperature here since calves are generally kept in warmer

environments than are adult cows but may also be due to some physiological

change with age.

This study also identified a decreased odds of detecting bla_{CTX-M} *E. coli* in samples

collected on pastureland compared with those collected elsewhere on the farm. This

relationship also held for *E. coli* resistant to amoxicillin and tetracycline. Because

pastureland may be more affected by the elements, this finding may be partly linked

with the association between temperature and ABR.

AB contamination of colostrum as a possible driver of bla_{CTX-M} E. coli positivity in dairy calves - evidence of direct and co-selection

It has been shown experimentally that feeding waste (AB-contaminated) milk to calves increases faecal excretion of ABR bacteria (25). This practice is reducing on UK dairy farms and, in the analysis presented here, waste milk feeding was not associated with an increased risk of finding ABR or bla_{CTX-M} positive E. coli. In contrast, the choice of dry cow therapy (an AB preparation inserted into a cow's udder between lactations to help treat or prevent mastitis) was associated with bla_{CTX-M} E. coli positivity in Calf samples (**Table 2**). It has previously been shown that colostrum from cows given cefquinome dry cow therapy is heavily contaminated with cefquinome (26), and colostrum management is a hugely important part of early life for most farmed mammals and is universally encouraged in dairy farming. In this study, cefquinome (a 4GC) dry cow therapy was most significantly associated with bla_{CTX-M} E. coli in Calf samples (**Table 2**). This can be explained by direct selection because production of CTX-M confers 4GC resistance in E. coli (27). There was also a clear positive association between the usage of framycetin as part of a dry cow therapy combination and the odds of finding bla_{CTX-M} E. coli in Calf samples (**Table 2**). Whilst no work has been published on the contamination of colostrum with framycetin, its use as a mastitis therapy for milking cows results in identifiable residues in milk (28), so it is highly likely to also contaminate colostrum. It is possible, therefore, that feeding of colostrum - which can be contaminated with AB used for dry cow therapy - is a driver of bla_{CTX-M} E. coli in calves. An alternative (or indeed an additional) explanation for this observed association is that E. coli (a species known to be found in the udders of dairy cows; 29) that carry bla_{CTX-M} are selected within the udder during AB dry cow therapy and contaminate colostrum

alongside the AB used. Others (16) also identified overall use of 3/4GCs as a risk

factor for bla_{CTX-M} *E. coli* presence on dairy farms but have not made a link between

the usage of framycetin and prevalence of bla_{CTX-M} *E. coli*. However, it is not always

clear whether other studies have separated out different dry cow therapies since

they have tended to focus on systemic AB use. Clearly, an aminoglycoside like

framycetin cannot directly select for bla_{CTX-M} *E. coli*, but aminoglycoside resistance

genes are common on plasmids (30), as is bla_{CTX-M} (27) and so this may be an

example of co-selection, where selection of resistance to one antibacterial class

increases resistance to another.

Conclusions

Overall, we provide important evidence that sampling at single time-points and/or

limited locations on a farm are unlikely to be adequate to accurately determine the

status of the farm concerning the prevalence of ABR *E. coli*. This makes

comparisons between surveillance studies on farms, designed for research or for

regulatory purposes, extremely difficult, and we urge for a standardised, multi-

sample framework accounting for the differential risk factors identified here to be

used in the design of future studies.

Materials and Methods

Farm recruitment and ethical approval

A convenience sample of 53 dairy farms was recruited through personal contacts, local veterinary practices, and milk processors. Details of the study population are presented in Supplementary. Of these, 43 farms were in a 50 x 50 km area defined based on the locations of 146 general practices that referred routine urine samples from human patients to the microbiology reference lab at Severn Pathology, Southmead Hospital (8). A further 10 study farms were clustered in a separate region in South West England. All farmers gave fully informed consent to participate in the study. Ethical approval was obtained from the University of Bristol's Faculty of Health Sciences Research Ethics Committee (ref 41562).

Farm sampling and sample processing

Farms were visited monthly between January 2017 and December 2018. Samples were collected using sterile overshoes (over-boot socks) traversing farm areas. Where access was restricted (e.g. for pens containing single or pairs of calves), samples were collected directly from the ground using gloved hands. Details of the six types of samples collected are in Supplementary; these represent faecally contaminated environments representative of milking cows (Adult), cows between periods of lactation (Dry Cow), heifers after weaning (Heifer), heifers before weaning (Calf). Samples of these types collected from pastureland were designated as such for separate analysis (Pasture) which also included samples collected from publicly accessible land on the farm (Footpath). Samples were refrigerated from collection to processing, were transferred into individual labelled sterile stomacher bags, and

suspended in 10 ml.g^{-1} of phosphate buffered saline (PBS Dulbecco A; Oxoid,

Basingstoke, UK). Samples were then mixed for one min in a stomacher (Stomacher

400, Seward, Worthing, UK). Samples were mixed 50:50 with sterile glycerol and

aliquots stored at -80°C.

Microbiology and PCR analysis

Twenty microlitres of sample (diluted 1:10) were spread onto tryptone bile X-

glucuronide agar (TBX; Scientific Laboratory Supplies); 20 µL of undiluted sample

were spread onto TBX agar containing 16 mg.L^{-1} tetracycline, 8 mg.L^{-1} amoxicillin,

0.5 mg.L^{-1} ciprofloxacin, 16 mg.L^{-1} streptomycin or 16 mg.L^{-1} cephalexin. Plates were

incubated at 37°C, and the number of blue colonies (*E. coli*) counted. Samples

yielding no *E. coli* colonies on antibiotic-free agar were excluded from further

analysis. Up to five *E. coli* isolates from each cephalexin TBX agar plate were

transferred onto cefotaxime (CTX, 2 mg.L^{-1}) TBX agar. All AB concentrations were

chosen as those which define clinically relevant resistance in humans according to

EUCAST (31). Two multiplex PCRs were performed to screen for β-lactamase genes

in CTX-R *E. coli*. The first was to detect *bla*$_{CTX-M}$ groups as previously described (32)

and the second was to detect the following additional β-lactamase genes: *bla*$_{CMY}$,

bla$_{DHA}$, *bla*$_{SHV}$, *bla*$_{TEM}$, *bla*$_{OXA-1}$ (8).

Risk factor analyses

The risk factors examined fell into four categories: farm management, ABU, sample

characteristics and meteorological. Four management practice questionnaires were

developed (details provided in Supplementary). ABU was extracted from prescribing

and sales data between Jan 2016 to Dec 2018 obtained from veterinary practices

servicing the study farms. For two farms, sales data were not available and on-farm

records were used. The full method for obtaining, cleaning, and processing ABU

data can be found in Supplementary. Local meteorological data were extracted from

publicly available UK Met Office data

(https://www.metoffice.gov.uk/pub/data/weather/uk/climate/stationdata/yeoviltondata.

txt).

Sample processing and data analysis workflows are illustrated in **Figure S1**. All data

analysis was performed using R (https://www.r-project.org/). Two modelling

approaches were used: 1) variable selection via univariable screening and stepwise

model selection with a multilevel, multivariable logistic regression model and 2) a

regularised Bayesian model. Both were used to analyse risk factors associated with

bla_{CTX-M} *E. coli* positivity; the second was also used to analyse risk factors

associated with positivity for *E. coli* resistant to amoxicillin, ciprofloxacin,

streptomycin, and tetracycline. Sensitivity analyses were performed to test for

measurement bias, to account for the fact that resistant *E. coli* were more likely to be

found in a sample if there was a higher density of bacterial colony forming units.

Further details of variable selection and development of the models and model

checking are provided in Supplementary.

Acknowledgements

We wish to thank all the farmers and veterinary surgeons who participated in this

study. In addition, we would like to thank the milk processing companies who

supported our recruitment process and the veterinary students who helped with data

processing and sample collection. Genome sequencing was provided by

MicrobesNG (http://www.microbesng.uk), which is supported by the BBSRC (grant

number BB/L024209/1).

361 **Funding**

362 This work was funded by grant NE/N01961X/1 to M.B.A., K.K.R., K.M.T., T.A.C. and

363 D.C.B. from the Antimicrobial Resistance Cross Council Initiative supported by the

364 seven United Kingdom research councils. R.A. was supported by the Jean Golding

365 Institute for data-intensive research at the University of Bristol. G.M.R. was funded

366 by the Langford Trust.

368 **Transparency declarations**

369 D.C.B. was president of the British Cattle Veterinary Association 2018-19.

370 Otherwise, the authors declare no competing interests. Farming and veterinary

371 businesses who contributed data and permitted access for sample collection were

372 not involved in the design of this study or in data analysis and were not involved in

373 drafting the manuscript for publication.

375 **Author Contributions**

Conceived the Study: D.C.B., K.K.R., M.B.A.

Collection of Data: H.S., J.F., K.M., E.F.P., O.M., V.C.G., supervised by T.A.C.,

K.K.R., G.M.R., M.B.A.

Cleaning and Analysis of Data: H.S., J.F., R.A., V.C.G., E.F.P, L.V., M.E., supervised

by K.M.T., K.K.R., M.B.A.

Initial Drafting of Manuscript: H.S., K.K.R., M.B.A.

Corrected and Approved Manuscript: All Authors

Tables

Table 1. Prevalence of *E. coli* carrying *bla*$_{CTX-M}$ at farm and sample levels

Sample type			Farm level	Sample level
Overall	**Total sample size**		53	4145
	Total (%) positive for CTX-M-carrying *E. coli*		42 (79.2%)	224 (5.4%)
Adult	**Total sample size**		52	1835
	Total (%) positive for CTX-M-carrying *E. coli*		25 (48.1%)	76 (4.1%)
Dry Cow	**Total sample size**		46	282
	Total (%) positive for CTX-M-carrying *E. coli*		7 (15.2%)	8 (2.8%)
Calf	**Total sample size**		51	631
	Total (%) positive for CTX-M-carrying *E. coli*		33 (64.7%)	98 (15.5%)
Heifer	**Total sample size**		41	1235
	Total (%) positive for CTX-M-carrying *E. coli*		18 (44%)	40 (3.2%)
Pastureland	**Total sample size**		47	630
	Total (%) positive for CTX-M-carrying *E. coli*		8 (17%)	12 (1.9%)
Pastureland that is publicly accessible (Footpath)	**Total sample size**		41	395
	Total (%) positive for CTX-M-carrying *E. coli*		8 (20.0%)	11 (2.8%)

387 **Table 2**. Fixed effects from the multilevel, multivariable logistic regression model

388 predicting bla_{CTX-M} *E. coli* positivity performed on Calf samples

Risk factor	Odds ratio [95% confidence interval]	p
Use of cefquinome dry cow therapy in the last six months	4.18 [2.11, 8.25]	0.00003
Daily water trough cleaning	0.44 [0.29, 0.69]	0.0002
Average monthly temperature	1.57 [1.20, 2.06]	0.0008
Use of framycetin dry cow therapy in the last six months	1.91 [1.01, 3.61]	0.04

390   **Table 3**. Fixed effects from the multilevel, multivariable logistic regression model

391   predicting bla_{CTX-M} *E. coli* positivity performed on the full dataset

Risk factor	Odds ratio [95% confidence interval]	p
Sample taken from the environment of pre-weaned heifers	4.52 [3.25, 6.27]	<0.00000001
Average monthly temperature	1.61 [1.36, 1.90]	0.00000001
Sample taken from pastureland	0.32 [0.17,0.61]	0.0004
Feeding of maize silage	3.28 [1.50, 7.18]	0.002

393   **Table 4.** Farm and sample-level prevalence of resistance to non-cephalosporins

Antibacterial drug	Farm level resistance	Sample level resistance
Amoxicillin	53/53 (100%)	2754/4145 (66%)
Ciprofloxacin	49/53 (92%)	263/4145 (6%)
Streptomycin	53/53 (100%)	1475/4145 (36%)
Tetracycline	53/53 (100%)	2874/4145 (69%)

395   **Table 5.** Fixed effects from a Bayesian model predicting resistant *E. coli* positivity

396   performed on the full dataset

Risk factor	Odds ratio [95% credible interval]
Amoxicillin resistance	
Average monthly temperature	1.91 [1.57, 2.35]
Sample taken from the environment of pre-weaned heifers	1.99 [1.29, 2.98]
Sample taken from pastureland	0.27 [0.20, 0.37]
Ciprofloxacin resistance	
Average monthly temperature	2.14 [1.63, 2.87]
Sample taken from the environment of pre-weaned heifers	4.13 [2.79, 6.46]
Streptomycin resistance	
Average monthly temperature	1.53 [1.32, 1.77]
Sample taken from the environment of pre-weaned heifers	1.95 [1.46, 2.51]
Tetracycline resistance	
Average monthly temperature	1.98 [1.55, 2.55]
Sample taken from the environment of pre-weaned heifers	3.39 [2.00, 5.82]
Sample taken from pastureland	0.24 [0.15, 0.35]
Sample taken during the calving season	2.02 [1.10, 3.29]
Average monthly rainfall	1.26 [1.08, 1.47]
Total farm streptomycin use in 2017 measured in mg/PCU*	0.76 [0.60, 0.97]
Total farm ABU in 2017 measured in mg/PCU*	1.47 [1.01, 2.13]

References

1. https://www.gov.uk/government/collections/antimicrobial-resistance-amr-information-and-resources. 2019.

2. ESVAC. European Surveillance of Veterinary Antimicrobial Consumption.

3. Veterinary Medicines Directorate & Animal and Plant Health Agency. 2015. UK Veterinary Antibiotic Resistance and Sales Surveillance.

4. Veterinary Medicines Directorate & Animal and Plant Health Agency. 2018. UK Veterinary Antibiotic Resistance and Sales Surveillance.

5. Lazarus B, Paterson DL, Mollinger JL, Rogers BA. 2014. Do Human Extraintestinal *Escherichia coli* Infections Resistant to Expanded-Spectrum Cephalosporins Originate From Food-Producing Animals? A Systematic Review. Clin Infect Dis 60:439-452.

6. Bevan ER, Jones AM, Hawkey PM. 2017. Global epidemiology of CTX-M β-lactamases: temporal and geographical shifts in genotype. J Antimicrob Chemother 72:2145-2155.

7. Madec J-Y, Haenni M. 2018. Antimicrobial resistance plasmid reservoir in food and food-producing animals. Plasmid 99:72-81.

8. Findlay J, Gould VC, North P, Bowker KE, Williams OM, MacGowan AP, Avison MB. 2020. Characterisation of cefotaxime-resistant urinary *Escherichia coli* from primary care in South-West England 2017-2018. J Antimicrob Chemother 75:65-71.

9. Alzayn M, Findlay J, Schubert H, Mounsey O, Gould VC, Heesom KJ, Turner KM, Barrett DC, Reyher KK, Avison MB. 2020. Characterisation of AmpC Hyper-Producing *Escherichia coli* from Humans and Dairy Farms Collected in Parallel in the Same Geographical Region. bioRxiv doi:10.1101/784694:784694; J Antimicrob Chemother (In Press) doi: 10.1093/jac/dkaa207.

10. Agersø Y, Aarestrup FM, Pedersen K, Seyfarth AM, Struve T, Hasman H. 2011. Prevalence of extended-spectrum cephalosporinase (ESC)-producing *Escherichia coli* in Danish slaughter pigs and retail meat identified by selective enrichment and association with cephalosporin usage. J Antimicrob Chemother 67:582-588.

430   11.   Randall LP, Clouting C, Horton RA, Coldham NG, Wu G, Clifton-Hadley FA,
431   Davies RH, Teale CJ. 2010. Prevalence of *Escherichia coli* carrying extended-
432   spectrum β-lactamases (CTX-M and TEM-52) from broiler chickens and
433   turkeys in Great Britain between 2006 and 2009. J Antimicrob Chemother
434   66:86-95.

435   12.   Overdevest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, Heck
436   M, Savelkoul P, Vandenbroucke-Grauls C, van der Zwaluw K, Huijsdens X,
437   Kluytmans J. 2011. Extended-spectrum β-lactamase genes of *Escherichia coli*
438   in chicken meat and humans, The Netherlands. Emerg Infect Dis 17:1216-
439   1222.

440   13.   Kola A, Kohler C, Pfeifer Y, Schwab F, Kühn K, Schulz K, Balau V, Breitbach
441   K, Bast A, Witte W, Gastmeier P, Steinmetz I. 2012. High prevalence of
442   extended-spectrum-β-lactamase-producing Enterobacteriaceae in organic and
443   conventional retail chicken meat, Germany. J Antimicrob Chemother 67:2631-
444   2634.

445   14.   Luvsansharav U-O, Hirai I, Nakata A, Imura K, Yamauchi K, Niki M,
446   Komalamisra C, Kusolsuk T, Yamamoto Y. 2012. Prevalence of and risk
447   factors associated with faecal carriage of CTX-M β-lactamase-producing
448   Enterobacteriaceae in rural Thai communities. J Antimicrob Chemother
449   67:1769-1774.

450   15.   McNulty CAM, Lecky DM, Xu-McCrae L, Nakiboneka-Ssenabulya D, Chung
451   K-T, Nichols T, Thomas HL, Thomas M, Alvarez-Buylla A, Turner K, Shabir S,
452   Manzoor S, Smith S, Crocker L, Hawkey PM. 2018. CTX-M ESBL-producing
453   Enterobacteriaceae: estimated prevalence in adults in England in 2014. J
454   Antimicrob Chemother 73:1368-1388.

455   16.   Snow LC, Warner RG, Cheney T, Wearing H, Stokes M, Harris K, Teale CJ,
456   Coldham NG. 2012. Risk factors associated with extended spectrum beta-
457   lactamase *Escherichia coli* (CTX-M) on dairy farms in North West England
458   and North Wales. Prev Vet Med 106:225-234.

459   17.   Mollenkopf DF, Weeman MF, Daniels JB, Abley MJ, Mathews JL, Gebreyes
460   WA, Wittum TE. 2012. Variable within- and between-herd diversity of CTX-M
461   cephalosporinase-bearing *Escherichia coli* isolates from dairy cattle. Appl
462   Environ Microbiol 78:4552-4560.

18. Horton RA, Duncan D, Randall LP, Chappell S, Brunton LA, Warner R,
Coldham NG, Teale CJ. 2016. Longitudinal study of CTX-M ESBL-producing
E. coli strains on a UK dairy farm. Res Vet Sci 109:107-113.

19. Brunton LA, Reeves HE, Snow LC, Jones JR. 2014. A longitudinal field trial
assesing the impact of feeding waste milk containing antibiotic residues on
the prevalence of ESBL-producing *Escherichia coli* in calves. Prev Vet Med
117:403-412.

20. MacFadden DR, McGough SF, Fisman D, Santillana M, Brownstein JS. 2018.
Antibiotic Resistance Increases with Local Temperature. Nat Clim Chang
8:510-514.

21. Watson E, Jeckel S, Snow L, Stubbs R, Teale C, Wearing H, Horton R,
Toszeghy M, Tearne O, Ellis-Iversen J, Coldham N. 2012. Epidemiology of
extended spectrum beta-lactamase *E. coli* (CTX-M-15) on a commercial dairy
farm. Vet Microbiol 154:339-346.

22. Duse A, Persson Waller K, Emanuelson U, Ericsson Unnerstad H, Persson Y,
Bengtsson B. 2016. Occurrence and Spread of Quinolone-Resistant
Escherichia coli on Dairy Farms. Appl Environ Microbiol 82:3765-3773.

23. Pereira RV, Siler JD, Ng JC, Davis MA, Grohn YT, Warnick LD. 2014. Effect
of on-farm use of antimicrobial drugs on resistance in fecal *Escherichia coli* of
preweaned dairy calves. J Dairy Sci 97:7644-7654.

24. Afema JA, Davis MA, Sischo WM. 2019. Antimicrobial use policy change in
pre-weaned dairy calves and its impact on antimicrobial resistance in
commensal *Escherichia coli*: a cross sectional and ecological study. BMC
Microbiol 19:217-217.

25. Awosile BB, Smith BA. 2017. Risk assessment modelling of fecal shedding
caused by extended-spectrum cephalosporin-resistant *Escherichia coli*
transmitted through waste milk fed to dairy pre-weaned calves. J Dairy Sci
100:9667-9673.

26. Bachmann J, Helmschrodt C, Richter A, Heuwieser W, Bertulat S. 2018.
Residue concentration of cefquinome after intramammary dry cow therapy
and short dry periods. J Dairy Sci 101:7540-7550.

27. Peirano G, Pitout JDD. 2019. Extended-Spectrum β-Lactamase-Producing
Enterobacteriaceae: Update on Molecular Epidemiology and Treatment
Options. Drugs. 79:1529-1541.

28. Luo PJ, Zhang JB, Wang HL, Chen X, Wu N, Zhao YF, Wang XM, Zhang H,
Zhang JY, Zhu L, Jiang WX. 2016. Rapid and Sensitive Chemiluminescent
Enzyme Immunoassay for the Determination of Neomycin Residues in Milk.
Biomed Environ Sci 29:374-378.

29. Bradley AJ, Green MJ. 2001. Adaptation of *Escherichia coli* to the bovine
mammary gland. J Clin Micro 39:1845-1849.

30. Schultsz C, Geerlings S. 2012. Plasmid-mediated resistance in
Enterobacteriaceae: changing landscape and implications for therapy. Drugs.
72:1-16.

31. European Committee on Antimicrobial Susceptibility Testing. 2020.
http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_ta
bles/v_9.0_Breakpoint _Tables.pdf. Last accessed May 2020.

32. Woodford N, Fagan EJ, Ellington MJ. 2005. Multiplex PCR for rapid detection
of genes encoding CTX-M extended-spectrum β-lactamases. J Antimicrob
Chemother 57:154-155.

Figure Legends

Figure 1. Average monthly temperature vs. presence (1) or absence (0) of *E. coli* positive for bla_{CTX-M} in samples from (**A)** pre-weaned calves and (**B**) all faecally contaminated dairy farm environments. Each sample is represented by a dot. A multilevel, multivariable logistic regression model revealed a positive association with increased temperature in both cases (p=<0.0001).

Figure 2. Percentage of farms positive for (**A**) *E. coli* positive for bla_{CTX-M} in samples (**B**) ciprofloxacin resistant *E. coli*. Data are presented by month (bars) and overlayed by a graph of average monthly temperature (dots) representing a year during the middle period of this study. Samples from calves have been excluded.

Figure 1

A

B

Figure 2

A

B