Molecular characterization of carbapenem-resistant *Klebsiella pneumoniae* isolates with focus on antimicrobial resistance

Running title: Genomic characterization of CRKP

Xiaoling Yu[1], Wen Zhang[2], Zhiping Zhao[1], Chengsong Ye[3], Shuyan Zhou[4], Shaogui Wu[4], Lifen Han[1], Zhaofang Han[5,6*] and Hanhui Ye[1*]

[1] Department of Infectious Diseases, Mengchao Hepatobiliary Hospital of Fujian Medical University, Xihong Road 312, Fuzhou 350025, Fujian, P.R. China.

[2] Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA.

[3] Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, Fujian, P.R. China.

[4] Department of Microbiology, Mengchao Hepatobiliary Hospital of Fujian Medical University, Xihong Road 312, Fuzhou 350025, Fujian, P.R. China.

[5] State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen, 361102, Fujian, PR China.

[6] Xiamen Cingene Science and Technology co., LTD, Xiamen 361021, Fujian, PR China.

Corresponding authors:

Zhaofang Han, E-mail: zhaofang_han@foxmail.com; Hanhui Ye, E-mail: 15960102808@163.com .

Abstract

The enhancing incidence of carbapenem-resistant *Klebsiella pneumoniae* (CRKP)-mediated

infections in Mengchao Hepatobiliary Hospital of Fujian Medical University in 2017 is the

motivation behind this investigation to study gene phenotypes and resistance-associated genes of

emergence regarding the CRKP strains. In current study, seven inpatients are enrolled in the

hospital with complete treatments. The carbapenem-resistant *K. pneumoniae* whole genome is

sequenced using MiSeq short-read and Oxford Nanopore long-read sequencing technology.

Prophages are identified to assess genetic diversity within CRKP genomes. The investigation

encompassed eight CRKP strains that collected from the patients enrolled as well as the

environment, which illustrate that bla_{KPC-2} is responsible for phenotypic resistance in six CRKP

strains that *K. pneumoniae* sequence type (ST11) is informed. The plasmid with IncR, ColRNAI

and pMLST type with IncF[F33:A-:B-] co-exist in all ST11 with KPC-2-producing CRKP

strains. Along with carbapenemases, all *K. pneumoniae* strains harbor two or three extended

spectrum β-lactamase (ESBL)-producing genes. *fosA* gene is detected amongst all the CRKP

strains. The single nucleotide polymorphisms (SNP) markers are indicated and validated among

all CRKP strains, providing valuable clues for distinguishing carbapenem-resistant strains from

conventional *K. pneumoniae*. In conclusion, ST11 is the main CRKP type, and bla_{KPC-2} is the

dominant carbapenemase gene harbored by clinical CRKP isolates from current investigations.

Keywords: *Klebsiella pneumoniae*; Carbapenem-resistant; Prophage; whole-genome sequencing

Introduction

Antibiotic resistance is amongst the extremely severe public health challenges nowadays. Carbapenem-resistant *Enterobacteriaceae* (CRE) is reported as a consequence mainly due to acquisition of carbapenemase genes, and CRE is inferred as an urgent threat to human health by the Centers for Disease Control and Prevention (CDC), USA in 2013[1]. Carbapenems such as imipenem, meropenem, and biapenem represent the first-line treatment of serious infections caused by multi-resistant Enterobacteriaceae including *Klebsiella pneumoniae* (*K. pneumoniae*) and *Escherichia coli* (*E. coli*)[2]. Whereas carbapenems can be hydrolyzed by carbapenemase in carbapenem-resistant *K. pneumoniae* (CRKP)[3], which results in resistance to β-Lactam antibiotics including carbapenem. Carbapenemases can be divided into Ambler class A β-lactamases (e.g. *Klebsiella pneumoniae* carbapenemases (KPC)), class B metallo-β-lactamases (MBLs), verona integrin-encoded metallo-β-lactamase (VIM), New Delhi metallo-β-lactamase (NDM) type, and Class D Enzymes of the OXA-48 type[4]. Among Ambler class A β-lactamases, plasmid-mediated KPC has been identified in all gram-negative members of the ESKAPE pathogens[5], and KPC is the most clinically indispensable enzyme due to its prevalence in Enterobacteriaceae[6]. Moreover, pathogens harboring KPC-2 are resistant to all β-lactams and β-lactamase inhibitors except ceftazidime/avibactam, which extremely limit treatment options as well as lead to high mortality rates[7]. Additionally, NDM has become a serious threat to public health due to the rapid global dissemination of NDM-bearing pathogens and the presence on mobile genetic elements in an extensive series of species[8]. Consequently, it is imperative and urgent to investigate the CRKP characteristics for better controlling pathogens and diagnosing as well as treating patients.

In current investigation, seven CRKP strains are extracted from patients during their

hospitalizations and another one CRKP strain is obtained from the dining car in Mengchao

Hepatobiliary Hospital of Fujian Medical University (**Supplementary Table 1**). The whole

genome of CRKP is sequenced using MiSeq short-reads and Oxford Nanopore long-reads

sequencing technology. We conduct surveillance of the CRKP-mediated infection prevalence in

Mengchao Hepatobiliary Hospital of Fujian Medical University, investigate the molecular

epidemiological characteristics of the strains that obtained, and identify gene phenotypes as well

as resistance-associated genes of the strain emergence. The detected single nucleotide

polymorphisms (SNP) markers would be helpful for recognizing CRKP strain from general *K.*

pneumoniae. Data of this study provide essential insights into effective strategy developments

for controlling CRKP and nosocomial infection reductions.

Materials & Methods

Patient clinical information

In total, seven patients received treatments during their hospitalizations and the data of them

were completely classified and studied. One bacterium was extracted from the dining car in the

hospital and since the carrier was not human, there was no clinical data relating to it. All patients,

except patient 1567P that was diagnosed as abdominal infection, were diagnosed as severe

pneumonia or suffered lung infections (**Supplementary Table 1**). We further give

Supplementary Table 2-8 to in detail provide all patients' treatment records as well as the

phenotype measurement results and data.

All patients received systematic medical examinations such as whole blood cell test, blood

routine test, blood electrolyte test, blood clotting, fungal D-glucan detection, galactomannan

detection, etc. All the records are archived in detail for further investigations.

Bacterial Isolates and Antimicrobial Resistance

Single patient isolates are obtained from specimens that received from inpatients admitted to

Mengchao Hepatobiliary Hospital of Fujian Medical University in 2017. A total of eight CRKP

isolates (**Supplementary Table 1**), which are resistance to all the antibiotics tested, such as

cephalosporins, penicilins, quinolones, aminoglycosides and carbapenems (Imipenem with MICs

$\geq$16 μg/ml) (Table 1), were processed following standard operating procedures: the isolates are

extracted according to the aseptic operating procedures and cultured in the bacterial culture

medium with Columbia Agar + 5% sheep blood. The study has been performed in accordance

with the Institutional Ethical Committee of the Faculty of Medicine, Mengchao Hepatobiliary

Hospital of Fujian Medical University, which approved this study.

K. pneumoniae isolates are confirmed by Matrix-assisted Laser Desorption Ionization-time of

Flight Mass Spectrometry (MALDI-TOF-MS) mass spectrometry (BioMerieux SA, BioMerieux

Inc., France). The resistance of pathogenic bacteria is identified by Automatic Microbial

Identification & Drug Sensitivity Analysis System (VITEK-2 Compact, BioMerieuxInc., France)

with Gram-Negative identification card (VITEK2 AST-GN13, BioMerieuxInc., France). The

results of antimicrobial susceptibility testing are interpreted based upon Clinical and Laboratory

Standards Institute (CLSI) M100-S24[9]. The standard strain under quality control is *K.*

pneumoniae isolates ATCC700603 (American Type Culture Collection, ATCC).

Whole Genome Sequencing (WGS) and Assembly

The isolated seven CRKP bacteria are sequenced on Illumina MiSeq (Illumina, San Diego, CA, USA) platform. MiSeq short-read sequencing library is generated with 1 ng purified DNA. Inserting a phosphate to 5' UTR end and "A" to 3' UTR end produces end-repair, and PCR fragments (300 ~ 600 bp) are collected from bar-coded adapter ligation. The library is purified via AMPure XP (Beckman Coulter) , which is then sequenced on MiSeq platform. In sum, a total of 40.5 million reads (2×300 bp) with a size of 1.36 Gb data are yielded (**Supplementary Table 9**). All short reads are first filtered for the low-quality sequences and then assembled into contigs using SPAdesv3.11.1 software[10].

Subsequently, we select an isolate of 1567D to perform long-read sequencing on Oxford Nanopore MinION (Oxford, UK) platform to easily sequence across repeat regions. The sequencing library is constructed with 1.5 μg purified DNA using the LSK-108 Oxford Nanopore Technologies (ONT) ligation protocol, and the prepared library is sequenced following the standard protocol of Oxford Nanopore MinION. A total of 7.48 Gb ultra-long reads are generated with N50 length of 25,890 bp (**Supplementary Table 10**). The long reads that 'passed' during the Nanopore base calling are used to assemble into complete genomic sequences via Canu software[11]. The long-read sequencing data of the same individual are used to correct base errors of assembled genome using Nanopolish (https://github.com/jts/nanopolish).

Detecting Prophages in the CRKP Genomes

The putative prophages within contigs of the CRKP genomic sequences are identified using the PHAST web server (PHAge Search Tool)[12]. The prophage completeness and categorization

127 (intact, incomplete, or questionable) are presented applying over sequences to check homology,

128 and to detect, annotate, and graphically display prophages.

Carbapenemase-resistance Gene Identifications

To predict the protein-coding genes and functional proteins in the CRKP genomes, all assembled

sequences are annotated by a web-based package RAST (Rapid Annotations using Subsystems

Technology)[13]. The antibiotic resistance and virulence genes, plasmids, phenotyping and

genotyping of CRKP genomes are scanned using the Bacterial Analysis Pipeline[14].

Carbapenemase-resistance genes are further identified from above annotated sequences

according to Simner et al.[15].

The protein-coding genes of long-read assembled genome are predicted using GLIMMER (Gene

Locator and Interpolated Markov ModelER) v3.02[16]. To functionally annotate the predicted

genes and perform the pathway analysis, we align them to NR, COG, Swiss-Prot, GO and KEGG

databases using blastX (E-value: 10^{-5}). The annotated genes serve to improve the completeness

of some important carbapenemase-resistance genes.

Comparisons of strain similarity are performed using the Harvest Tools Suite[17] (version 1.1.2).

For all of the isolates sequenced on a particular platform, parsnp is utilized to compare all the

assembled isolates against each other and known reference strain. Results are visualized using

EvolView.

SNP Identification and Validation

We download *K. pneumoniae* genome from NCBI as the reference to identify SNP markers[18].

All high-quality data (Q value >20, reads length > 50 bp, number of uncertain bases < 5%) of

eight CRKP strains are aligned to the reference genome sequences using BWA v0.7.17[19], and

aligned reads are sorted by coordinates via SAMTOOLS v1.4[20]. The GATK (Genome Analysis

Tool Kit) software v3.8.0[21] is utilized to detect SNPs, which is described as following: (1)

duplicated reads are removed; (2) reads around insertions/deletions are realigned; (3) base

quality is recalibrated using default parameters; (4) all variants are identified using

HaplotypeCaller method in GATK with emitting and calling standard confidence thresholds at

10.0 and 30.0, respectively. To validate the detected SNPs in the seven CRKPs, we select 20 loci

within each sample that are located in exonic regions and sequence them with high read depth.

All chosen markers are designed primers for amplification using Sequenom MassARRAY

iPLEX platform.

Results

Antimicrobial Susceptibilities of the CRKP Strains

The source of isolates is supplied in **Table 1**, which denotes the infectious type and the result of

susceptibility testing during the patients' hospitalization. All eight strains involved in the study

are confirmed to be *K. pneumoniae*, with five strains from sputum, one from bile, one from blood,

and one from the environment (**Supplementary Table 1**). Clinical data demonstrate that seven

of the eight patients are referred due to pulmonary infection, and another one is referred due to

abdominal infection. The susceptibility testing data in **Table 1** reveals that all the *K. pneumoniae*

strains are resistant to almost all antibiotics, such as cephalosporins, penicilins, quinolones and

carbapenems (imipenem with MICs $\geq$16 μg/ml). For aminoglycosides antibiotics, except that

1567D isolate is sensitive to amikacin and tobramycin, all other isolates are resistant to

aminoglycosides antibiotics. The strains including 1566D, 2038D, 2039D and 2040D are

resistant to sulfamethoxazole/trimethoprim with MICs $\geq$320, and the other strains (1567D,

2035D, 2036D, 2037D) are sensitive to sulfamethoxazole/trimethoprim with MICs $\leq$ 20.

Genome assembly and annotation

The short-read sequenced seven CRKP strains are assembled into contigs. As listed in **Table 2**, the assembled genome size of all trains ranged from 5.4 Mb to 5.8 Mb, with mean length of 5.7 Mb and average contigs numbering 199. The N50 length of genomes is from 176.6 kb to 251.6 kb with an average N50 length of 220.4 kb and mean GC content of 57.2%. To obtain a more complete genome, the 1567D strain is resequenced via long-read sequencing technology and assembled into three contigs with size of 5.6 Mb (**Supplementary Figure 1**). A total of 5,841 protein-coding genes are predicted with length between 37 to 1,649 bp (**Supplementary Figure 2**). Totals of 4,657, 5,097, 4,714, 3,179 and 3,099 predicted genes are functionally annotated in NR, COG, Swiss-Prot, GO and KEGG databases, respectively (**Supplementary Figures 3, 4, 5**).

Characteristics of the CRKP Isolates

The isolated seven CRKP bacteria are sequenced through Illumina MiSeq platform and assembled into whole genomes. To understand genetic diversity, mobile genetic elements of 24 prophages are identified in seven CRKP genomes, with sizes ranging from 8.4 kb to 49.3 kb (**Figure 1**). According to the criterion that the length of an intact prophage should be more than 20 kb[22]. Prophages detected in most strains (except for 2036D) are complete with a size of at least 20.2 kb with an average GC percentage of 52.7%. Additionally, three prophages are respectively identified in 3 strains at the same time, revealing the genomic sequence homology among all isolates. The 2036D strain is comprised of just one prophage probably because of the small genome size and distinct sequence characteristics, which is expected to have less neutral targets for prophage integration[22].

Furthermore, multilocus-sequence typing (MLST) analysis reveals that there are two unrelated sequence type (ST) in *K. pneumoniae* strains isolated from different patients. 2036D *K.*

pneumoniae strain correlates with ST2632, and the other six strains are relevant to ST11 (**Table**

3). pMLST analysis reveals that all of the six ST11 *K. pneumoniae* strains are associated with

IncF[F33:A-:b-] and the ST2632 *K. pneumoniae* strain is relevant to IncHI1 and IncF. Majority

of *K. pneumoniae* strains are ST11, with IncF [F33:A-:b-] type.

Plasmid analysis[23] shows different circular plasmids carried by the individual strains. All strains

harbored IncR and ColRNAI plasmids with no virulence genes but contain several resistance-

associated genes that cause resistance to carbapenems, which is demonstrated in **Table 3**. The

IncR plasmid is identified as multidrug-resistant plasmids and has variable copy numbers of

certain resistance genes among *K. pneumoniae* isolates.

Detection of Antibiotic Resistance Genes of CRKP Isolates

The resistance-associated genes of seven CRKP bacteria (**Table 3**) are sequenced on Illumina

MiSeq platform among the patient and environmental isolates. As illustrated in **Table 3**, some

antimicrobial resistance genes are mediated by plasmid such as β-lactamase correlative genes

(*bla*$_{\text{CTX-M}}$, *bla*$_{\text{KPC}}$, *bla*$_{\text{LEN}}$, *bla*$_{\text{TEM}}$) and those genes which encoded aminoglycoside [*aac(3)-IId,*

rmtB], chloramphenicol (*catA1, catA2*), trimethoprim (*dfrA1,dfrA17*), and fluoroquinolone

[*QnrS1*]. The other antimicrobial resistance genes are encoded by chromosome including *bla*$_{\text{SHV}}$

(narrow-spectrum β-lactamasein *K. pneumoniae*), *oqxA* (1,176 bp), *oqxB* (3,153 bp) (efflux

pumps), and *fosA* (420 bp, fosfomycin resistance) genes.

Except 2036D, all the other *K. pneumoniae* strains harbor the associated carbapenemases-

producing resistance gene, *bla*$_{\text{KPC-2}}$. Extended-spectrum β-lactamases (ESBLs) resistance genes

such as *bla*$_{\text{CTX-M}}$, *bla*$_{\text{TEM}}$, *bla*$_{\text{LEN}}$ and *bla*$_{\text{SHV}}$ are also informed. *bla*$_{\text{TEM}}$ is one of the genes that

produce ESBL. *bla*$_{\text{CTX-M}}$ with different types (*bla*$_{\text{CTX-M-14}}$, *bla*$_{\text{CTX-M-3,}}$*bla*$_{\text{CTX-M-55}}$ and *bla*$_{\text{CTX-M-65}}$)

is found among all the *K. pneumoniae* strains. *bla*$_{\text{CTX-M-3}}$ is observed in 2036D strains. *bla*$_{\text{CTX-M-55}}$

ɪ

is observed in 1567D strain and $bla_{CTX-M-14}$ is observed in 1566D and 2040D strains. $bla_{CTX-M-65}$

is detected in the other four (2035D, 2037D, 2038D, 2039D) *K. pneumoniae* strains. bla_{LEN12}

gene is exclusively found in 1566D strain, and there is no bla_{SHV} gene in it. Nevertheless, bla_{SHV-}

$_{93}$ and bla_{SHV-11} genes are detected in 2036D strain and the other five *K. pneumoniae* strains,

respectively. Except for the 2036D strain, bla_{TEM-1B} gene is observed in all the other six *K.*

pneumoniae strains. *Aac(3)-IId* and *rmtB* encoding fluoroquinolone resistance are observed

among all strains. *fosA* resulting in fosfomycin resistance[24] is also informed among all strains.

Characterizing CRKP SNPs

The SNP markers are identified for all strains that sequenced using the short-read MiSeq data.

The data demonstrate that 33,716 markers are detected in the 2036D strain, which is more

significant than the other strains with an average of 8,289 SNPs. The cSNPs located in exonic

regions are in slightly higher amounts among all detected SNPs of a minimal ratio of 85.5%

(**Supplementary Table 11**). In addition, the pairwise comparison analysis reveals that 2036D

isolate is disparate with the other strains based on clusters of sequence similarities using

subprogram of Trinity[25] (**Figure 2a**). Furthermore, the 2036D strain share few SNP loci with the

others, which coincides with strain clusters (**Figure 2b**).

For validations, all strains have a high detection rate in that approximately 153 out of 200 SNPs

(76.4%) that have amplifications, which demonstrate the analysis accuracy (**Supplementary**

Table 11). After filtering SNP loci that are not located in exome regions, containing no-alleles

locus, and comprise all-wild SNP loci in each isolate, we eventually obtain 92 SNPs among 200

validated loci. A total of 40 out of 92 SNPs are all-variation loci in all isolates, which could be

utilized for recognizing CRKP strain from ordinary *K. pneumoniae* (**Supplementary Table 12**).

In addition, 24 SNPs of strain's unique loci, including strains of 2036D (18 loci), 2035D (3 loci),

1566D (2 loci) and 2037D (1 loci), would be helpful resources for specific strain identification of

clinical analysis.

Phylogeny analysis results

The phylogenic tree shows that 1567D strain is most distantly related to the other strains, and

2036D is more closely related to the reference strain comparing with all the other strains

(**Supplementary Figure 6**).

GWAS analysis

To further identify significant SNPs and genes, we perform genome-wide association study

(GWAS) analysis. The patients' body temperature and counts of leukocyte are selected as

phenotypic character. The short-sequencing reads of six strains (**Figure 3**) are aligned to the

1567D genome using BWA v0.7.17 software. We call SNPs using Platypus v0.8.1[26], and then

filter the SNPs through plink v1.9 according to the following conditions: (i) missing loci, (ii)

minor allele frequency (MAF) < 0.05 and (iii) significant deviation from the Hardy-Weinberg

equilibrium (HWE) ($P < 0.01$). A total of 698 SNP markers are remained and utilized for GWAS

analysis. As a result, 9 loci are identified ($P < 0.05$). Two loci (ygbI and murB) are related with

temperature and the other seven loci (IsrD, SufD, yrkF, fabI, sppA, entF and ttuB) are relevant to

leukocyte (**Figure 3**).

Discussion

Data of current study confirm that all CRKP strains hold two types of plasmids with no virulence

gene whereas harbor an abundance of associated resistance genes such as ESBLs and

carbapenemases. One genotype of carbapenemases with bla_{KPC-2} and two ST types with ST11

and ST2632 are identified in the study, and the ST11 with KPC-2-positive is a prevalent strain

accounting in all the six strains. The plasmid with IncR, ColRNAI and pMLST type with

IncF[F33:A-:B-] co-exist in all ST11 with KPC-2-producing CRKP strains. The initial detection

of a KPC-2-producing *K. pneumoniae* isolate from a hospital in China is reported in 2007[27].

Since then, bla_{KPC-2}-bearing *K. pneumoniae* isolates have become more prevalent and reported in

China as well as other countries and areas [28]. Recently, one patient is found to have susceptible *K.*

pneumoniae bacteraemia in US[15]. While that case is relatively specific since the patient might be

affected during the visit and hospitalization in India, which would add more complex

environmental factors to confound the results. CRKP of ST11 associated with bla_{KPC-2} is

disseminated widely across China[18,29], which is concordant with the results of our study. These

findings all suggest that the CRKP-mediated infections in our hospital result from ST11 with

KPC-2-positive *K. pneumoniae* isolates. Continuous monitoring will be necessary to prevent

further dissemination of carbapenemase-resistance genes.

Besides carbapenemases, a variety of ESBLs such as bla_{CTX-M}, bla_{SHV}, bla_{LEN}, bla_{TEM} are present

in CRKP strains of this study. *K. pneumoniae* is one of the most indispensable infectious agents

in the ICU[30]. There are "classic" and hypervirulent strains of *K. pneumoniae*[31,32,33]. The "classic"

non-virulent strain of *K. pneumoniae* (C-KP) can produce ESBLs related to nosocomial

infectious outbreaks especially in the ICU of a hospital. C-KP more easily acquires antimicrobial

resistance such as ESBLs. bla_{CTX-M} with different type is found among all the CRKP strains.

Chromosome-mediated bla_{SHV} and plasmid-mediated bla_{TEM} are also positive for ESBLs

production and are observed in six *K. pneumoniae* strains. Co-occurrence of bla_{CTX-M}, bla_{KPC-2},

bla_{SHV-11} and bla_{TEM-1B} are observed among five *K. pneumoniae* strains. All *K. pneumoniae*

strains harbor two or three ESBLs-producing genes (bla_{CTX-M}, bla_{SHV} and bla_{TEM}), which indicate

all isolates contained multiple ESBLs resistance genes. Previous reports noted consistent results

that co-occurrence of bla_{TEM}, bla_{SHV} and bla_{CTX-M} (any two or all three) was observed among

Klebsiella isolates[34].

fosA is frequently identified in the *E. coli* and *K. pneumoniae* genomes [35,36]. The *fosA5* gene is

first found in *E. coli* in 2014[37]. In 2017, it was reported that all of 73 carbapenem-resistant *K.*

pneumoniae isolates were positive for *fosA5* in one Chinese area: Zhejiang Province[38].

Antimicrobial susceptibility testing about fosfomycin is not conducted in this study though *fosA*

is also found among all the CRKP strains, which might indicate that fosfomycin-modifying

enzymes account for a majority of the fosfomycin resistance, and that fosfomycin is resistant to

CRKP strains. It is reported that *fosA* gene is transferred from *E. coli* to *K. pneumoniae* through

whole plasmid transmission or mobile genetic element transmission, which raise doubts whether

fosfomycin can be used as a supplementary drug for urinary tract infection caused by

carbapenem-resistant *E. coli* in the hospital, as *fosA* exists in all CRKP strains from our study.

Continuous monitoring will be necessary to prevent further dissemination of fosfomycin-

resistant bacteria together with prudent use of fosfomycin in clinical settings.

OqxA and *oqxB* genes are relevant to efflux pumps, which means that antibiotics such as

cephalosporins, carbapenems and fluoroquinolones are almost completely expelled from *K.*

pneumoniae through its cell membrane[39]. Although there are no carbapenemases that observed in

2036D strain, *oqxA* and *oqxB* genes are identified in it. To our knowledge, these two genes are

mainly reported to be responsible for the resistance to fluoroquinolones. They do have been

previously reported to be associated with the nitrofurantoin resistance.

The genome sequences of the seven strains include massive contigs which are highly fragmented.

Upon further investigation, we sequence the 1567D strain using long-read sequencing platform,

which could help us assemble the genome with considerable improvement in completeness and

contiguity. The carbapenem-resistant genes including *fosA*, *oqxA* and *oqxB* and 40 all-variation

SNP loci are also identified in the above genome demonstrating the high-quality assembly. The

assembly and annotation information will be beneficial in understanding the whole genomic

characterization of CRKP strain for future study.

Conclusions

In conclusion, ST11 is the main CRKP type, and *bla*$_{KPC-2}$ is the dominant carbapenemase gene

harbored by clinical CRKP isolates from present study. The plasmid with IncR, ColRNAI and

pMLST type with IncF[F33:A-:B-] exist in all ST11 with KPC-2-producing CRKP strains.

Besides carbapenemases, all *K. pneumoniae* strains harbor two or three ESBLs-producing genes

(*bla*$_{CTX-M}$, *bla*$_{SHV}$ and *bla*$_{TEM}$), which indicate all isolates contain multiple ESBLs resistance

genes. *fosA* genes are also found among all the CRKP strains, which may infer that fosfomycin-

modifying enzymes account for a majority of the fosfomycin resistance and that CRKP strains

are resistant to fosfomycin. The differential expressions of *oqxA* and *oqxB* in CRKP strain might

possibly result in carbapenem-resistant, but this presumption needs more solid experimental

evidences. The 40 all-variation SNP loci in all isolates could be employed and referred for

distinguishing CRKP strain from ordinary *K. pneumoniae*.

Declarations

Ethics approval

The study has been performed in accordance with the Institutional Ethical Committee of the

Faculty of Medicine, Mengchao Hepatobiliary Hospital of Fujian Medical University.

Consent to participate

Not applicable.

Consent to publish

All authors read and approved the final manuscript.

Availability of data and materials

The genome shotgun sequencing data and long reads of Oxford Nanopore data are deposited at

NCBI/GenBank as BioProject of PRJNA506754.

Competing interests

The authors declare that they have no competing interests.

Funding

This study is sponsored by Key Clinical Specialty Discipline Construction Program of Fuzhou,

P.R. China (Grant No. 201510301), Clinical Medicine Center Construction Program of Fuzhou,

Fujian, P.R.C. (Grant No. 2018080306), Health Research Innovation Team Cultivation Project of

Fuzhou, P.R.C. (Grant No. 2019-S-wt4) and Key Clinical Specialty Discipline Construction

Program of Fujian, P.R. China.

Authors' Contributions

X.Y., Z.Z., L.H. and H.Y. conceived of the method. H.Y. supervised the study. S.Z., S.W. and

C.Y. implemented the bacteria culture. X.Y., W.Z. and Z.H. performed the bioinformatics

analysis. X.Y. and Z.H. optimized and performed the sequencing. X.Y., W.Z. and Z.H. drafted

the article with inputs and feedbacks from all the other authors. All authors read and approved

the final manuscript.

Acknowledgements

References

1. Queenan AM, Bush K. 2007. Carbapenemases: the versatile beta-lactamases. Clinical microbiology
reviews 20:440-458.

2. Lee C, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. 2016. Global Dissemination of
Carbapenemase-Producing Klebsiella pneumoniae: Epidemiology, Genetic Context, Treatment
Options, and Detection Methods. Frontiers in Microbiology 7:895.

3. Xu L, Sun X, Ma X. 2017. Systematic review and meta-analysis of mortality of patients infected with
carbapenem-resistant Klebsiella pneumoniae. Annals of Clinical Microbiology and Antimicrobials
16:18.

4. Nordmann P, Naas T, Poirel L. 2011. Global Spread of Carbapenemase-producing Enterobacteriaceae.
Emerging Infectious Diseases 17:1791-1798.

5. Pendleton JN, Gorman SP, Gilmore B. 2013. Clinical relevance of the ESKAPE pathogens. Expert
Review of Anti-infective Therapy 11:297-308.

6. Nordmann P, Cuzon G, Naas T. 2009. The real threat of Klebsiella pneumoniae carbapenemase-
producing bacteria. Lancet Infectious Diseases 9:228-236.

7. Tzouvelekis LS, Markogiannakis A, Psichogiou M, Tassios PT, Daikos GL. 2012. Carbapenemases in
Klebsiella pneumoniae and Other Enterobacteriaceae: an Evolving Crisis of Global Dimensions.
Clinical Microbiology Reviews 25:682-707.

8. Göttig S, Hamprecht A, Christ S, Kempf V, A Wichelhaus T. 2013. Detection of NDM-7 in Germany,
a new variant of the New Delhi metallo- -lactamase with increased carbapenemase activity, vol 68.

9. Clinical and Laboratory Standards Institute C. 2014. Performance standards for antimicrobial
susceptibility testing; twenty-fourth informational supplement, vol 34.

10. Bankevich A, Nurk S, Antipov D, Gurevich A, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjibelski AD. 2012. SPAdes: A New Genome Assembly Algorithm and Its Applications to
Single-Cell Sequencing. Journal of Computational Biology 19:455-477.

11. Koren S, Walenz B, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Research
27:722-736.

12. Zhou Y, Liang Y, Lynch KH, Dennis J, Wishart DS. 2011. PHAST: A Fast Phage Search Tool.
Nucleic Acids Research 39:347-352.

13. Brettin T, Davis JJ, Disz T, Edwards R, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch
GD. 2015. RASTtk: A modular and extensible implementation of the RAST algorithm for building
custom annotation pipelines and annotating batches of genomes. Scientific Reports 5:8365-8365.

14. Thomsen MCF, Ahrenfeldt J, Cisneros J, Jurtz VI, Larsen MV, Hasman H, Aarestrup FM, Lund O.
2016. A Bacterial Analysis Platform: An Integrated System for Analysing Bacterial Whole Genome
Sequencing Data for Clinical Diagnostics and Surveillance. PLOS ONE 11.

15. Simner PJ, Antar AAR, Hao S, Gurtowski J, Tamma PD, Rock C, Opene BNA, Tekle T, Carroll KC,
Schatz MC. 2018. Antibiotic pressure on the acquisition and loss of antibiotic resistance genes in
Klebsiella pneumoniae. Journal of Antimicrobial Chemotherapy 73:1796-1803.

16. Kelley DR, Liu B, Delcher AL, Pop M, Salzberg SL. 2012. Gene prediction with Glimmer for
metagenomic sequences augmented by classification and clustering. Nucleic Acids Research 40.

17. Treangen TJ, Ondov BD, Koren S et al. 2014. The Harvest suite for rapid core-genome alignment and
visualization of thousands of intraspecific microbial genomes. Genome Biol 15: 524.

18. Bi D, Jiang X, Sheng ZK, Ngmenterebo D, Tai C, Wang M, Deng Z, Rajakumar K, Ou HY. Mapping
the resistance-associated mobilome of a carbapenem-resistant Klebsiella pneumoniae strain reveals
insights into factors shaping these regions and facilitates generation of a 'resistance-disarmed' model
organism. J Antimicrob Chemother. 2015 Oct;70(10):2770-4.

19. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics 25:1754-1760.

20. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth GT, Abecasis GR, Durbin R.
2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078-2079.

21. Mckenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky AM, Garimella K, Altshuler
D, Gabriel SB, Daly MJ. 2010. The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. Genome Research 20:1297-1303.

22. Vale F, Nunes A, Oleastro M, Gomes J, A. Sampaio D, Rocha R, Vítor J, Engstrand L, Pascoe B,
Berthenet E, K. Sheppard S, Hitchings M, Megraud F, Vadivelu J, Lehours P. 2017. Genomic
structure and insertion sites of Helicobacter pylori prophages from various geographical origins.
Scientific Reports 7:42471.

23. Li X, Xie Y, Liu M, Tai C, Sun J, Deng Z, Ou HY. oriTfinder: a web-based tool for the identification
of origin of transfers in DNA sequences of bacterial mobile genetic elements. Nucleic Acids Res.
2018 Jul 2;46(W1):W229-W234.

24. Ito R, Mustapha MM, Tomich AD, Callaghan JD, Mcelheny CL, Mettus RT, Shanks RMQ,
Sluiscremer N, Doi Y. 2017. Widespread Fosfomycin Resistance in Gram-Negative Bacteria
Attributable to the Chromosomal fosA Gene. Mbio 8.

25. Grabherr M, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q. 2011. Full-length transcriptome assembly from RNA-Seq data without a
reference genome. Nature Biotechnology 29:644-652.

26. Rimmer, A. J., Phan, H., Mathieson, I., et al. 2014. Integrating mapping-, assembly- and haplotype-
based approaches for calling variants in clinical se- quencing applications. Nature Genetics, 46, 912-
918.

27. Wei Z, Du X, Yu Y, Shen P, Chen Y, Li L. 2007. Plasmid-mediated KPC-2 in a Klebsiella
pneumoniae isolate from China. Antimicrobial Agents and Chemotherapy 51:763-765.

28. Zhang R, Liu L, Zhou H, Chan EW, Li J, Fang Y, Li Y, Liao K, Chen S. 2017. Nationwide
Surveillance of Clinical Carbapenem-resistant Enterobacteriaceae (CRE) Strains in China.
EBioMedicine 19:98-106.

29. Liu J, Yu J, Chen F, Yu J, Simner P, Tamma P, Liu Y, Shen L. 2018. Emergence and establishment of
KPC-2-producing ST11 Klebsiella pneumoniae in a general hospital in Shanghai, China. European
Journal of Clinical Microbiology & Infectious Diseases 37:293-299.

30. Meropol SB, Haupt AA, Debanne SM. 2018. Incidence and Outcomes of Infections Caused by
Multidrug-Resistant Enterobacteriaceae in Children, 2007–2015. Journal of the Pediatric Infectious
Diseases Society 7:36-45.

31. Patel PK, Russo TA, Karchmer AW. 2014. Hypervirulent Klebsiella pneumoniae. Open Forum
Infectious Diseases 1.

32. Xie Y, Tian L, Li G, Qu H, Sun J, Liang W, Li X, Wang X, Deng Z, Liu J, Ou HY. Emergence of the
third-generation cephalosporin-resistant hypervirulent Klebsiella pneumoniae due to the acquisition of
a self-transferable blaDHA-1-carrying plasmid by an ST23 strain. Virulence. 2018 Dec 31;9(1):838-
844.
451
33. Wang X, Xie Y, Li G, Liu J, Li X, Tian L, Sun J, Ou HY, Qu H. Whole-Genome-Sequencing
characterization of bloodstream infection-causing hypervirulent Klebsiella pneumoniae of capsular
serotype K2 and ST374. Virulence. 2018 Jan 1;9(1):510-521.

34. Hossain Mondal A, Siddiqui MT, Sultan I, Mohd. Rizwanul Haq Q. 2018. Prevalence and diversity of
bla TEM, bla SHV and bla CTX-M variants among multidrug resistant Klebsiella spp. from an urban
riverine environment in India. International Journal of Environmental Health Research 6:1-13.

35. Ito R, Mustapha M, D. Tomich A, D. Callaghan J, McElheny C, T. Mettus R, Shanks R, Sluis-Cremer
N, Doi Y. 2017. Widespread Fosfomycin Resistance in Gram-Negative Bacteria Attributable to the
Chromosomal fosA Gene. mBio 8:e00749-17.

36. Li G, Zhang Y, Bi D, Shen P, Ai F, Liu H, Tian Y, Ma Y, Wang B, Rajakumar K, Ou HY, Jiang X.
First report of a clinical, multidrug-resistant Enterobacteriaceae isolate coharboring fosfomycin
resistance gene fosA3 and carbapenemase gene blaKPC-2 on the same transposon,
Tn1721.Antimicrob Agents Chemother. 2015 Jan;59(1):338-43.

37. Ma Y, Xu X, Guo Q, Wang P, Wang W, Wang M. 2015. Characterization of fosA5, a new plasmid-
mediated fosfomycin resistance gene in Escherichia coli. Letters in Applied Microbiology 60:259-264.

38.   Huang L, Yan Hu Y, Zhang R. 2017. Prevalence of fosfomycin resistance and plasmid-mediated
fosfomycin-modifying enzymes among carbapenem-resistant Enterobacteriaceae in Zhejiang, China.
Journal of Medical Microbiology 66:1332-1334.
39.   Zheng J-x, Lin Z-w, Sun X, Lin W-h, Chen Z, Wu Y, Qi G-b, Deng Q-w, Qu D, Yu Z-j. 2018.
Overexpression of OqxAB and MacAB efflux pumps contributes to eravacycline resistance and
heteroresistance in clinical isolates of Klebsiella pneumoniae. Emerg Microbes Infect 7:139.

Figures

494   **Figure 1. Intact prophages identified in seven CRKP strains.**

499 **Figure 2. Assessing the genetic relatedness of the CRKP by WGS.** (a) Clustering of all isolates based on

500 sequence similarity. (b) Communal SNP markers detected by pairwise comparison analysis.

Isolates	1566D	2035D	2036D	2037D	2038D	2039D	2040D
1566D		6922	2275	6895	7952	7848	8029
2035D	6922		2304	7249	7103	6989	7152
2036D	2275	2304		2287	2333	2342	2344
2037D	6895	7249	2287		7050	6964	7132
2038D	7952	7103	2333	7050		8417	8549
2039D	7848	6989	2342	6964	8417		8467
2040D	8029	7152	2344	7132	8549	8467	

Z

Figure 3. GWAS results of the analysis.

Tables

Table 1. Antibiotic susceptibility profiles of *K. pneumoniae*. The results of antimicrobial susceptibility testing - antibiotics MIC (mg/L) and breakpoint interpretation or epidemiological cut-off value. S: susceptible; I: intermediate; R: resistant.

Isolates	1566D	1567D	2035D	2036D	2037D	2038D	2039D	2040D
source	sputum	bile	sputum	sputum	blood	sputum	sputum	environment
Infection	Pulmonary	Abdominal	Pulmonary	Pulmonary	Pulmonary	Pulmonary	Pulmonary	N/A
ampicillin	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)
ampicillin/ sulbactam	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)	≥32(R)
piperacillin/ tazobactam	≥128(R)	≥128(R)	≥128(R)	≥128(R)	≥128(R)	≥128(R)	≥128(R)	≥128(R)
cefazolin	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)
cefotetan	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)
ceftazidime	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)
ceftriaxone	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)
cefepime	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)
aztreonam	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)
imipenem	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)
amikacin	≥64(R)	≤4（S)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)	≥64(R)

gentamicin	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)
tobramycin	≥16(R)	≤2（S）	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)	≥16(R)
ciprofloxacin	≥4(R）	≥4(R）	≥4(R）	≥4(R）	≥4(R）	≥4(R）	≥4(R）	≥4(R）
levofloxacin	≥8(R)	≥8(R)	≥8(R)	≥8(R)	≥8(R)	≥8(R)	≥8(R)	≥8(R)
macrodantin	256(R）	256(R)	≥512(R）	≥512(R）	≥512(R）	≥512(R）	≥512(R）	≥512(R）
sulfamethoxazole/ trimethoprim	≥320(R)	≤20（S）	≤20（S）	≤20（S）	≤20（S）	≥320(R)	≥320(R)	≥320(R)

Table 2. Assembly statistics of seven CRKP strains via short-read sequencing.

Assembly	1566D	2035D	2036D	2037D	2038D	2039D	2040D
Contig number	195	163	215	242	184	196	195
Total length (bp)	5,758,754	5,633,502	5,432,179	5,670,795	5,833,697	5,831,354	5,835,044
Largest contig (bp)	380,781	381,132	929,110	380,795	381,132	381,056	381,056
GC (%)	57.33	57.39	57.18	57.35	57.21	57.21	57.2
N50	176,606	196,479	251,620	196,479	183,862	190,289	183,862
L50	12	11	7	11	11	11	11
Total number of Ns	20	30	20	30	130	130	30

Table 3. Resistance genes among the patient and environmental isolates.

Isolates	1566D	2035D	2036D	2037D	2038D	2039D	2040D
MLST	ST11	ST11	ST2632	ST11	ST11	ST11	ST11
pMLST	IncF[F33:A-:B-]	IncF[F33:A-:B-]	IncHI1, IncF	IncF[F33:A-:B-]	IncF[F33:A-:B-]	IncF[F33:A-:B-]	IncF[F33:A-:B-]
Plasmids	IncR, ColRNAI	IncR, ColRNAI	IncR, ColRNAI	IncR, ColRNAI	IncR, ColRNAI	IncR, ColRNAI	IncR, ColRNAI
Penicillins: Ampicillin/ Narrow-Spectrum	*blaCTX-M-14* *blaKPC-2*	*blaCTX-M-65* *blaKPC-2*	*blaCTX-M-3*	*blaCTX-M-65* *blaKPC-2*	*blaCTX-M-65* *blaKPC-2*	*blaCTX-M-65* *blaKPC-2*	*blaCTX-M-14* *blaKPC-2*
Cephalosporins: cefazolin and cefotetan	*blaLEN12* *blaTEM-1B*	*blaSHV-11* *blaTEM-1B*	*blaSHV-93*	*blaSHV-11* *blaTEM-1B*	*blaSHV-11* *blaTEM-1B*	*blaSHV-11* *blaTEM-1B*	*blaSHV-11* *blaTEM-1B*
β-lactam inhibitors/ Carbapenems	*blaKPC-2*	*blaKPC-2*	-	*blaKPC-2*	*blaKPC-2*	*blaKPC-2*	*blaKPC-2*
Extended-Spectrum Cephalosporins/ Monobactam	*blaCTX-M-14*	*blaCTX-M-65*	*blaCTX-M-3*	*blaCTX-M-65*	*blaCTX-M-65*	*blaCTX-M-65*	*blaCTX-M-65*
Aminoglycosides	*rmtB*	*rmtB*	*aac(3)-IId*	*rmtB*	*aac(3)-IId* *rmtB*	*aac(3)-IId* *rmtB*	*aac(3)-IId* *rmtB*
Fluoroquinolones	*QnrS1*	-	*oqxA* *oqxB*	-	*QnrS1*	*QnrS1*	*QnrS1*
Phosphonic Acid	*fosA*	*fosA*	*fosA*	*fosA*	*fosA*	*fosA*	*fosA*
Phenicol	-	*catA2*	*catA1*	*catA2*	*catA2*	*catA2*	*catA2*
Folate-pathway Inhibitors	*dfrA1*	-	*dfrA17*	-	*dfrA1*	*dfrA1*	*dfrA1*