1

Common gene expression signatures in Parkinson's disease are driven by changes in cell composition

Gonzalo S. Nido[1,2], Fiona Dick[1,2], Lilah Toker[1,2], Kjell Petersen[1,3], Guido Alves[4,5], Ole-Bjørn Tysnes[1,2], Inge Jonassen[1,3], Kristoffer Haugarvoll[1,2], Charalampos Tzoulis[1,2,*]

[1] Neuro-SysMed, Department of Neurology, Haukeland University Hospital, 5021 Bergen, Norway

[2] Department of Clinical Medicine, University of Bergen, Pb 7804, 5020 Bergen, Norway

[3] Computational Biology Unit, Department of Informatics, University of Bergen, Pb 7803, 5020 Bergen, Norway

[4] The Norwegian Centre for Movement Disorders and Department of Neurology, Stavanger University Hospital, Pb 8100, 4068 Stavanger, Norway

[5] Department of Mathematics and Natural Sciences, University of Stavanger, 4062 Stavanger, Norway.

* Corresponding author

E-mail: charalampos.tzoulis@nevro.uib.no (CT)

Abstract

Background

The etiology of Parkinson's disease (PD) is largely unknown. Genome-wide transcriptomic studies in bulk brain tissue have identified several molecular signatures associated with the disease. While these studies have the potential to shed light into the pathogenesis of PD, they are also limited by two major confounders: RNA post mortem degradation and heterogeneous cell type composition of bulk tissue samples. We performed RNA sequencing following ribosomal RNA depletion in the prefrontal cortex of 49 individuals from two independent case-control cohorts. Using cell-type specific markers, we estimated the cell-type composition for each sample and included this in our analysis models to compensate for the variation in cell-type proportions.

Results

Ribosomal RNA depletion results in substantially more even transcript coverage, compared to poly(A) capture, in post mortem tissue. Moreover, we show that cell-type composition is a major confounder of differential gene expression analysis in the PD brain. Correcting for cell-type proportions attenuates numerous transcriptomic signatures that have been previously associated with PD, including vesicle trafficking, synaptic transmission, immune and mitochondrial function. Conversely, pathways related to endoplasmic reticulum, lipid oxidation and unfolded protein response are strengthened and surface as the top differential gene expression signatures in the PD prefrontal cortex.

42 **Conclusions**

43 Differential gene expression signatures in PD bulk brain tissue are significantly confounded by

44 underlying differences in cell-type composition. Modeling cell-type heterogeneity is crucial in

45 order to unveil transcriptomic signatures that represent regulatory changes in the PD brain and

46 are, therefore, more likely to be associated with underlying disease mechanisms.

Introduction

Introduction

Parkinson's disease (PD) is the second most prevalent neurodegenerative disorder, affecting ~1.8

% of the population above 65 years [1]. PD is a complex disorder, caused by a combination of

genetic and environmental factors, but the molecular mechanisms underlying its etiology remain

largely unaccounted for. Genome-wide transcriptomic studies can identify expression signatures

associated with PD. While not able to establish causality, these studies hold the potential to

highlight important biological mechanisms, some of which may be exploited as targets for

therapeutic modulation.

A recent systematic review identified 33 original genome-wide transcriptomic studies in the

PD brain, of which 6 were performed on laser microdissected neurons from the *substantia nigra*

pars compacta (SNc) and the remaining in bulk tissue from various brain regions [2]. These

studies show surprisingly low replicability at the level of individual genes, however, and only

partial concordance for pathways. The most consistent alterations have been found in pathways

related to energy metabolism/mitochondrial function and protein degradation, followed by

synaptic transmission, vesicle trafficking, lysosome/autophagy and neuroinflammation [2].

While these processes commonly show differential expression signatures in PD, it remains

unknown whether this is because they truly reflect the biology of PD or due to systematic bias

and confounding factors. Two major sources of bias for transcriptomic studies in the human

brain are the *post mortem* degradation of RNA and the highly heterogeneous cell type

composition of bulk tissue samples.

RNA degradation of variable extent occurs in *post mortem* tissue. To further complicate the

picture, it has been shown that different cell types exhibit different degrees of susceptibility to

RNA degradation [3], potentially confounding differences in cellular composition with

differences in RNA quality. Access to high-quality brain tissue is generally limited, and thus an

optimal choice of experimental platforms becomes paramount to maximize sensitivity. While

RNA microarrays are being gradually superseded by RNA-seq technology, only 3 out of the 33

studies identified by an up-to-date review [2] used RNA-seq, and all of them employed poly(A)

capture, a widely used protocol to restrict the analysis to mature mRNA [4–6]. However, this

library preparation method only picks up RNA fragments with a poly-A tail, introducing

substantial bias in low quality RNA samples [7–10]. A well-established approach to mitigate this

limitation, is whole RNA-seq following ribosomal RNA (rRNA) depletion [11]. To our

knowledge there are no genome-wide transcriptomic studies on PD brain employing rRNA

depletion methods to date.

Systematic differences in sample cell composition represent another important confounding

factor. These typically originate from two sources: biological differences (e.g. secondary to

neurodegeneration) and technical variation in sample dissection and preparation. Brain areas

affected by neurodegeneration are characterized by neuronal loss and gliosis, resulting in a

systematically increased glia-to-neurons ratio in patients. This confounder is strongest in areas

with severe changes, such as the SNc, but is also present to a variable degree in less affected

areas, such as the neocortex. In addition, technical sources of variation due to sampling may

affect any brain region and cause an uneven distribution of gray and white matter, resulting in a

variable fraction of oligodendrocytes. Thus, transcriptional signatures associated with PD in bulk

brain tissue may reflect changes in cellular composition rather than disease-specific

transcriptional modulation. This observation has already been put forward using

neurodegenerative mouse models and re-analysis of human brain transcriptomic data [12].

Heterogeneous cell composition is, hence, a major confounder that needs to be considered and

appropriately addressed in transcriptomic studies in bulk brain samples.

We report the first genome-wide transcriptomic study in the PD brain employing RNA-seq

following rRNA depletion. We show that this approach is able to substantially mitigate the bias

of *post mortem* degradation, resulting in substantially better transcript coverage compared to

poly(A) capture. Moreover, by estimating the relative cell type proportion in our samples, we

confirm that cellular composition is a major source of variation in bulk tissue data, confounding

the differential gene expression profile even in the less affected prefrontal cortex. By

incorporating the estimated cell type proportions into our analysis models, we were able to

unveil transcriptomic signatures which are more likely to be associated with the underlying

disease mechanisms.

Results

Ribosomal RNA depletion is superior to poly(A) selection in *post mortem* brain

We carried out RNA-seq following rRNA depletion in fresh-frozen prefrontal cortex (Brodmann

area 9) from a total of 49 individuals from two independent cohorts: the Norwegian ParkWest

study (PW, n = 29) [13] and the Netherlands Brain Bank (NBB, n = 20). A detailed description

of our cohorts is provided in the methods. Comparison of our data to a published poly(A) capture

dataset of similar characteristics [4] revealed important differences of mapping coverage.

Mapping efficiency was slightly higher in the poly(A) dataset (median = 0.976, range = 0.971-

0.980) compared to the rRNA-depleted datasets (PW: median = 0.952, range = 0.940-0.962;

NBB: median = 0.959, range = 0.947-0.965). The counts per million (CPMs) of rRNA regions,

as defined by Ensembl release 75, was very low in all samples (PW: median = 3,099, range =

1,047-7,071; NBB: median = 1,583, range = 1,129-5,024) and, as expected, significantly lower

in the rRNA-depleted cohorts than in the poly(A) capture (poly(A) cohort: median = 40,058,

range = 10,701-95,183) (S1 Fig).

In both datasets, the RNA integrity number (RIN) was positively correlated with mapping

efficiency to mRNA regions, but not to intergenic and/or intronic regions (S2 Fig). Despite

having higher mean RIN values, the poly(A) capture cohort showed a marked unevenness of

transcript body coverage compared to the rRNA depletion samples (Fig 1A). The median

coefficient of variation in coverage was significantly lower in the rRNA-depleted samples and

the 5'- and 3'- ends of the transcripts showed substantially better coverage compared to the

poly(A) sample (Fig 1B). Moreover, in the rRNA-depleted samples both the 3'- and 5'-end

coverage loss showed a significant inverse correlation with the RIN values. In contrast, RIN

showed no correlation with the 5'-bias and a positive correlation with the 3'-bias in the poly(A)

capture samples (Fig 1C). Thus, rRNA depletion results in substantially better and more even

coverage of the transcriptome in *post mortem* brain tissue, providing a better alternative to

poly(A) capture and minimizing the prospect of transcript quantification biases downstream.

Cell composition is a major confounder of gene expression in bulk brain samples

The observed gene expression profiles in bulk brain tissue can be dramatically influenced by

differences in cellular composition. Such differences can be a result of variation in gray/white

matter ratios introduced during tissue extraction, inter-subject variability or represent disease

related alterations [14–16]. To study the contribution of various technical and biological sources

of variation in our dataset we first estimated marker gene profiles (MGPs) for the major classes

of cortical cell types (astrocytes, microglia, oligodendrocytes, and neurons) in our samples by

summarizing the expression of the cell type-specific marker genes as previously described [15].

Next, we examined the Pearson's correlation between potential sources of biological variation in

our data, including technical and demographic factors (RIN, *post-mortem* interval (PMI), sex,

age, and disease status) and MGPs. As expected, MGPs for neuronal cell types were significantly

anticorrelated with the other main cortical cell types in both cohorts ($p < 0.05$, Fig 2A). In

agreement with previous studies [3,17], cellular estimates were also confounded with RNA

quality. In both cohorts RIN was significantly correlated with neuronal (positive correlation) and

astrocyte (negative correlation) MGPs. Significant negative correlation of RIN with microglia

MGPs was observed in the NBB cohort (Fig 2A). Most concerning was the detection of a

significant association between the oligodendrocyte MGP and the disease status in the PW

cohort (Fig 2A). The main axis of variation in gene expression (which explained 44% and 45%

of the total variance in PW and NBB, respectively) was significantly correlated with RNA

quality and cellular composition in both cohorts (Fig 2B), singling out RNA quality and cellular

composition as the main drivers of transcriptional change in bulk brain tissue.

We next looked for differences in cellular proportions between PD and controls adjusting for

the known experimental covariates (see methods). In the NBB cohort, PD subjects exhibited a

significant increase in the microglia MGP ($p = 0.015$, Wilcoxon test), while a significant increase

in the oligodendrocyte MGP ($p = 5.5 \times 10^{-3}$, Wilcoxon test) was observed in PD subjects from the

PW cohort. In both cohorts, these changes were accompanied by a non-significant decrease in

neuronal MGPs (Fig 2C).

Since the oligodendrocyte and microglia MGPs were significantly associated with the

disease status in at least one of the cohorts, we re-estimated group differences in astrocyte and

neuronal MGPs, adjusting for the significant MGPs, in addition to the original covariates. This

analysis showed no significant differences between the groups (S3 Fig). Therefore, only MGPs

of oligodendrocytes and microglia were included in the statistical model of differential

expression analysis.

Differential gene expression

Differential gene expression analysis of a total of ~31,000 pre-filtered genes was carried out

using experimental covariates (sex, age, PMI, RIN, and sequencing batch) with or without

oligodendrocyte and microglia MGPs. In the PW cohort, 595 genes were defined as differentially

expressed (Benjamini-Hochberg FDR < 0.05) without adjusting for cell type composition.

Inclusion of oligodendrocyte and microglia MGPs in the model decreased the number of

differentially expressed genes to a total of 220. In total, 74 genes remained significant both with

and without adjustment for cell type composition. No transcripts with FDR < 0.05 were

identified in the NBB cohort, irrespective of adjustment for cell type composition. A list with the

nominally significant genes overlapping between the two cohorts is provided in S1 File.

Comprehensive results of differential expression analysis are available in S2 File.

Functional enrichment

Functional enrichment analysis of the differential gene expression results without MGP

adjustment indicated 476 significantly enriched (Benjamini-Hochberg FDR < 0.05) pathways in

PW (107 up-regulated and 369 down-regulated) and 992 in NBB (421 up-regulated and 571

down-regulated). MGP adjustment reduced the number of significant pathways to 89 in PW (35

up-regulated and 54 down-regulated) and 248 in NBB (115 up-regulated and 133 down-

regulated). Of these, 34 pathways replicated across the two cohorts. Concordant pathways

comprised protein folding, ER-related processes and lipid oxidation (Fig 3). The complete results

are provided in S3 File.

As expected, scoring each pathway according to the change in p-value when accounting for

cellularity, revealed a marked downplay of the relevant cell-type-specific functions (Table 1). In

the PW cohort, which was characterized by a skewed oligodendrocytes/neurons proportion, the

function with the largest attenuation (i.e. increase in p-value) was seen for up-regulation of

myelination and other oligodendrocyte related functions and for down-regulation of neuronal

pathways. For NBB, correcting for cell-composition resulted in attenuation of immunity and

neuronal pathways, consistent with the unbalanced microglial/neuronal proportions seen in that

cohort (Table 1). Strikingly, pathways linked to mitochondrial respiration, including respiratory

complex I, were among the down-regulated processes that lost statistical significance when

controlling for cellularity. The attenuation of the mitochondrial signal was observed in both

cohorts. Conversely, up-regulation of protein folding-related pathways gained significance in

both cohorts (Table 2 and Fig 3). Complete results are provided in S4 File.

Table 1. Loss of significance in enriched pathways.

PW			
Up-regulated		**Down-regulated**	
Pathway	*Delta*	*Pathway*	*Delta*
myelination	-8.85	regulation of synaptic vesicle exocytosis	-9.63
ensheathment of neurons	-8.67	intrinsic component of synaptic membrane	-9.60
axon ensheathment	-8.67	regulation of synaptic vesicle cycle	-9.58
detection of chemical stimulus involved in sens…	-7.95	positive regulation of synaptic transmission	-9.48
oligodendrocyte differentiation	-6.94	Schaffer collateral - CA1 synapse	-9.46
oligodendrocyte development	-6.58	regulation of synaptic plasticity	-9.44
apical junction complex	-5.57	regulation of neurotransmitter secretion	-9.41
glial cell development	-5.54	presynaptic membrane	-9.27
glial cell differentiation	-5.52	regulation of synaptic vesicle transport	-9.19
tight junction	-4.16	protein transport within lipid bilayer	-9.11

NBB			
Up-regulated		**Down-regulated**	
Pathway	*Delta*	*Pathway*	*Delta*
activation of innate immune response	-8.54	ribonucleoside monophosphate metabolic process	-9.09
regulation of leukocyte proliferation	-8.11	purine nucleoside triphosphate metabolic process	-9.03
regulation of lymphocyte proliferation	-8.01	mitochondrial membrane part	-8.99
regulation of mononuclear cell proliferation	-7.79	ATP metabolic process	-8.92
innate immune response-activating signal transd…	-7.43	regulation of synaptic vesicle exocytosis	-8.87
regulation of adaptive immune response	-7.16	inner mitochondrial membrane protein complex	-8.82
response to interferon-gamma	-7.15	purine ribonucleoside triphosphate metabolic proc…	-8.77
adaptive immune response based on somatic rec…	-7.07	respiratory chain	-8.70
blood microparticle	-6.75	regulation of synaptic vesicle transport	-8.68
regulation of T cell proliferation	-6.73	cellular respiration	-8.43

197 Tables representing the top 10 pathways with the lowest delta for up- and down-regulated
198 pathways for both cohorts. The delta value represents the change in the enrichment $-\log_{10}(p$-
199 value) between the results with and without MGP adjustment (negative values of delta imply a
200 loss of significance when accounting for cellularity). Complete results are provided in S4 File.

201 **Table 2. Gain of significance in enriched pathways.**

PW			
Up-regulated		**Down-regulated**	
Pathway	*Delta*	*Pathway*	*Delta*
protein folding	5.77	DNA packaging complex	3.76
'de novo' protein folding	5.73	basement membrane	3.47
unfolded protein binding	5.54	positive regulation of epithelial cell proliferation	2.52
chaperone-mediated protein folding	5.32	negative regulation of gliogenesis	2.49
'de novo' posttranslational protein folding	4.68	fatty acid beta-oxidation	2.30
heat shock protein binding	4.34	nucleosome	2.18
response to unfolded protein	4.10	glomerulus development	2.16
response to topologically incorrect protein	3.53	aorta development	2.06
oxidoreductase activity, acting on paired…	2.74	endothelium development	1.95

NBB			
Up-regulated		**Down-regulated**	
Pathway	*Delta*	*Pathway*	*Delta*
positive regulation of cardiac muscle tissue dev…	1.99	tertiary granule	5.22
regulation of smooth muscle cell differentiation	1.98	ficolin-1-rich granule membrane	5.00

negative regulation of protein serine/threonine kin…	1.98	regulation of myeloid leukocyte mediated immunity	4.55
hormone-mediated signaling pathway	1.95	regulation of leukocyte degranulation	4.34
lung alveolus development	1.71	specific granule	4.22
positive regulation of striated muscle tissue dev…	1.69	ficolin-1-rich granule	4.15
positive regulation of muscle organ development	1.69	tertiary granule membrane	3.57
positive regulation of muscle tissue development	1.62	regulation of mast cell activation	3.52
negative regulation of MAP kinase activity	1.48	vacuolar lumen	3.05
regulation of cardiac muscle cell differentiation	1.48	regulation of mast cell degranulation	3.05

203 Tables representing the top 10 pathways with the highest delta for up- and down-regulated
204 pathways for both cohorts. The delta value represents the change in the enrichment $-\log_{10}$(p-
205 value) between the results with and without MGP adjustment (positive values imply an increase
206 in p-value when accounting for cellularity). Complete results are provided in S4 File

Discussion

We present the first genome-wide transcriptomic study in the PD brain employing whole RNA-

seq after rRNA depletion. Our findings show that PD-associated differential gene expression

signatures in bulk brain tissue are influenced to a great extent by the underlying differences in

cell-type composition of the samples. Modeling cell-type heterogeneity allowed us to highlight

transcriptional signatures that are likely to represent aberrant gene expression within the cells of

the PD brain, rather than changes in cell composition.

Our results suggest that rRNA depletion is superior to the more commonly used poly(A)

capture approach and allows for a more accurate mapping and quantification of the transcriptome

in *post mortem* brain tissue. The rRNA depletion method provides substantially higher evenness

of coverage and effectively mitigates the 3'- and 5'-end coverage bias associated with poly(A)

capture [7–10]. Ultimately, the unevenness of coverage will influence transcript quantification,

affecting the sensitivity of the differential expression estimates.

Our study supports the notion that cell composition can be a major confounder in bulk brain

tissue transcriptomics. The observed expression profiles in both cohorts were driven

predominantly by a combination of technical factors associated with RNA quality, and

differences in cellular composition between PD and controls. This difference was primarily due

to oligodendrocytes in PW and microglia in NBB. Since oligodendrocyte proliferation is not a

pathological feature of PD, it is plausible that the difference in oligodendrocyte MGPs in PW

was due to technical variation in gray/white matter content introduced during tissue sampling.

Microglial infiltration does occur in affected areas of the PD brain [18]. It is noteworthy,

however, that increased microglial MGP was only observed in one of the cohorts (NBB),

highlighting the biological heterogeneity of PD. Accounting for relative cell proportions reduced

the number of differentially expressed genes and attenuated the calculated enrichment of cell-

type-specific pathways between PD and controls. In the PW cohort, this alleviated a substantial

false positive signature of oligodendrocyte genes presumably caused by skewed grey/white

matter sampling bias. Similar sampling bias could be responsible for oligodendroglia-specific

functions appointed to PD brain in previous transcriptomic studies [6].

Intriguingly, accounting for cellular proportions downplayed several of the transcriptomic

signatures that have been previously associated with PD. For instance, the signal from vesicle

trafficking- and synaptic transmission-related processes [19–24] was significantly attenuated in

both cohorts, suggesting that the signal was primarily driven by changes in neuronal proportions

between PD and controls, rather than modulation of these pathways within neurons. Moreover,

we observed an attenuation in the down-regulation of mitochondrial pathways, including the

respiratory chain and oxidative phosphorylation, which are among the most consistent

transcriptomic signatures in PD [2,4,19,21,25–28]. The loss of transcriptional signal in these

pathways is intriguing, because there is compelling evidence that decreased complex I protein

levels occur in PD neurons [29]. Our results suggest that the previously reported transcriptional

down-regulation of the respiratory chain is at least partly driven by altered cellular composition

(due to decreased number of neurons which highly express these genes) and may therefore not be

the sole mechanism by which neuronal complex I deficiency occurs in PD. Indeed, it has been

suggested that complex I deficiency in PD may be mediated by proteolytic degradation by the

LON-ClpP protease system, rather than transcriptional regulation [30].

Similar changes in the cell-composition of the brain are seen in all neurodegenerative

diseases, including PD, Alzheimer disease, ALS and Huntington disease. Interestingly, common

and overlapping transcriptional signatures have been reported across these neurodegenerative

diseases, including mitochondrial, neuronal-specific, and immunity-related pathways [31,32].

Our findings suggest that these common transcriptional signatures of neurodegeneration may be

largely confounded by a common pattern of altered cellularity, involving neuronal loss and glial

proliferation, rather reflect biological processes of causal nature.

Correcting for cell-type composition in our samples highlighted processes related to the

endoplasmic reticulum, unfolded protein response and lipid/fatty acid oxidation as the top

differential gene expression signatures in the PD prefrontal cortex. Unfolded protein response is

indeed one of the most consistently reported transcriptomic signatures in PD [2,4,21,25,33,34].

Moreover, endoplasmic reticulum stress and aberrant proteostasis have been associated with the

accumulation of misfolded proteins, including α-synuclein, in both *in vitro* studies and animal

models of PD [35]. While less is known regarding the role of lipid metabolism in PD, evidence

of aberrant fatty acid oxidation has been found by metabolomic studies in serum [36] and urine

[37] of patients. Our results corroborate these findings and indicate that aberrant fatty acid

metabolism occurs in the PD prefrontal cortex.

Based on our findings, we advocate that modeling cell-type heterogeneity is crucial in order

to unveil transcriptomic signatures reflecting regulatory changes in the PD brain. It is, however,

important noting that modeling of cellular estimates cannot completely mitigate the cell-

composition bias in bulk tissue. Moreover, cell-type correction complicates identification of

transcriptional events co-occurring with changes in cellular composition. Single-cell or cell-

sorting based methods will be key to overcoming this limitation and deciphering transcriptomic

signatures directly associated with underlying disease mechanisms in PD.

Materials and methods

Subject cohorts

All experiments were conducted in fresh-frozen prefrontal cortex (Brodmann area 9) from a total

of 49 individuals from two independent cohorts. The first cohort (n = 29) comprised individuals

with idiopathic PD (n = 18) from the Park-West study (PW), a prospective population-based

cohort which has been described in detail [38] and neurologically healthy controls (Ctrl, n = 11)

from our brain bank for aging and neurodegeneration. Whole-exome sequencing had been

performed on all patients [39] and known/predicted pathogenic mutations in genes implicated in

Mendelian PD and other monogenic neurological disorders had been excluded. None of the study

participants had clinical signs of mitochondrial disease or use of medication known to influence

mitochondrial function (S5 File). Controls had no known neurological disease and were matched

for age and gender. The second cohort comprised samples from 21 individuals from the

Netherlands Brain Bank (NBB) including idiopathic PD (n = 10) and demographically matched

neurologically healthy controls (n = 11). Individuals with PD fulfilled the National Institute of

Neurological Disorders and Stroke [40] and the UK Parkinson's disease Society Brain Bank [41]

diagnostic criteria for the disease at their final visit.

To investigate the effect of rRNA depletion compared to the prevailing poly(A) capture

method, we re-analyzed an RNA-seq dataset from a previous publication which employed a

poly(A) tail selection kit on post-mortem tissue of the same brain area and same disease (poly(A)

cohort, n = 29 PD samples, n = 44 neurologically healthy controls, all males; GEO: GSE68719)

[4].

Informed consent was available from all individuals. Ethical permission for these studies

was obtained from our regional ethics committee (REK 2017/2082, 2010/1700, 131.04).

Tissue collection and neuropathology

Brains were collected at autopsy and split sagittaly along the *corpus callosum*. One hemisphere

was fixed whole in formaldehyde and the other coronally sectioned and snap-frozen in liquid

nitrogen. All samples were collected using a standard technique and fixation time of ~ 2 weeks.

There was no significant difference in PMI (independent t-test, PW cohort p = 0.16; NBB cohort

p = 0.92), age (independent t-test, PW cohort p = 0.18; NBB cohort p = 0.074) or gender

(independent t-test, PW cohort p = 0.94; NBB cohort p = 0.53) between PD subjects and

controls. Subject demographics and tissue availability are provided in S5 File. Routine

neuropathological examination including immunohistochemistry for α-synuclein, tau and beta

amyloid was performed on all brains. All cases showed neuropathological changes consistent

with PD including degeneration of the dopaminergic neurons of the SNc in the presence of Lewy

pathology. Controls had no pathological evidence of neurodegeneration.

RNA sequencing

Total RNA was extracted from prefrontal cortex tissue homogenate for all samples using RNeasy plus mini kit (Qiagen) with on-column DNase treatment according to manufacturer's protocol. Final elution was made in 65 µl of dH2O. The concentration and integrity of the total RNA was estimated by Ribogreen assay (Thermo Fisher Scientific), and Fragment Analyzer (Advanced Analytical), respectively and 500 ng of total RNA was used for downstream RNA-seq applications. First, rRNA was removed using Ribo-Zero™ Gold (Epidemiology) kit (Illumina, San Diego, CA) using manufacturer's recommended protocol. Immediately after the rRNA removal the RNA was fragmented and primed for the first strand synthesis using the NEBNext First Strand synthesis module (New England BioLabs Inc., Ipswich, MA). Directional second strand synthesis was performed using NEBNExt Ultra Directional second strand synthesis kit. Following this the samples were taken into standard library preparation protocol using NEBNext® DNA Library Prep Master Mix Set for Illumina® with slight modifications. Briefly, end-repair was done followed by poly(A) addition and custom adapter ligation. Post-ligated materials were individually barcoded with unique in-house Genomic Services Lab (GSL) primers and amplified through 12 cycles of PCR. Library quantity was assessed by Picogreen Assay (Thermo Fisher Scientific), and the library quality was estimated by utilizing a DNA High Sense chip on a Caliper Gx (Perkin Elmer). Accurate quantification of the final libraries for sequencing applications was determined using the qPCR-based KAPA Biosystems Library Quantification kit (Kapa Biosystems, Inc.). Each library was diluted to a final concentration of 12.5 nM and pooled equimolar prior to clustering. 125 bp Paired-End (PE) sequencing was performed on an Illumina HiSeq2500 sequencer (Illumina, Inc.). RNA quality, as measured by the RNA integrity number (RIN), varied across samples (mean = 5.3, range = 3.0-7.2 for PW;

mean = 6.8, range = 3.2-9.1 for NBB, S5 File), although the difference between conditions did

not reach statistical significance in any of the cohorts (t-test P = 0.72 and 0.90 for PW and NBB

cohorts, respectively).

Data quality control

FASTQ files were trimmed using Trimmomatic version 0.36 [42] to remove potential Illumina

adapters and low quality bases with the following parameters:

ILLUMINACLIP:truseq.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15.

FASTQ files were assessed using fastQC version 0.11.5 [43] prior and following trimming. For

an in-depth quality assessment, we mapped the trimmed reads using HISAT2 version 2.1.0 [44]

against the hg19 human reference genome (using --rna-strandness RF option) preserving lane-

specific information. To discard potential lane-specific sequencing batch effects we inspected the

output of the CollectRnaSeqMetrics tool of Picard Tools version 2.6 [45]. Mapping efficiency

and proportion of reads mapping to rRNA, intronic, intergenic and coding regions were obtained

from the output of the CollectRnaSeqMetrics (S1 and S2 Figs).

For the poly(A) capture dataset [4], raw FASTQ files were obtained from the Gene

Expression Omnibus (GEO:GSE68719) and analyzed exactly as described for our cohorts (with

the exception of --rna-strandness in HISAT2, which was turned off to take into account that the

cDNA library of this cohort was unstranded).

RNA expression quantification and filtering

We used Salmon version 0.9.1 [46] to quantify the abundance at the transcript level with the

fragment-level GC bias correction option (--gcBias) and the appropriate option for the library

type (-l ISR) against the Ensembl release 75 transcriptome. Transcript-level quantification was

collapsed onto gene-level quantification using the tximport R package version 1.8.0 [47]

according to the gene definitions provided by the same Ensembl release. We filtered out genes in

non-canonical chromosomes and scaffolds, and transcripts encoded by the mitochondrial

genome. To further reduce the potential for artifacts we filtered out transcripts with unusually

high expression by removing transcripts that gathered more than 1% of the reads on more than

half of the samples, which resulted in the removal of 3 and 4 transcripts from the PW and NBB

cohorts, respectively. Additionally, low-expressed (i.e. genes whose expression was below the

median expression in at least 20% of the samples) were filtered out from downstream analyses.

Samples were then marked as outliers if their median correlation in gene expression (log counts

per million) with the other samples was below Q_1-1.5*IQR or above Q_3+1.5*IQR (*Tukey's*

fences; Q_1: first quartile, Q_3: third quartile, IQR: inter-quartile range). As a result, 3 samples

were marked as outliers in the PW cohort and 3 in the NBB cohort, and were not included in

downstream analyses (resulting sample sizes: $N_{PW} = 26$, $N_{NBB} = 18$, S4 Fig).

Estimation of cell composition

For each sample we calculated correlates for the main cortical cell type types (neurons,

oligodendroglia, microglia and astrocytes) by summarizing the expression of their respective

marker genes as the first principal component of their expression (marker gene profiles, MGPs

(Mancarci et al., 2017)). Cortical cell type markers were selected using the homologene R

package version 1.5.68 [48]. To unravel potential complex interactions between MGPs and other

experimental covariates, including disease status, we calculated the pairwise correlation between

all the variables and also their association with the main axes of variation of gene expression. To

assist us in choosing an optimal set of MGPs to include as covariates, we quantified the group

differences in the cellular proportions between PD and controls using linear models adjusting for

the known experimental covariates (i.e. RIN, PMI, sex, age, and sequencing batch). Significant

association with disease status was found for oligodendrocyte MGP in the PW cohort and for

microglia in the NBB cohort. Thus, these were included in the downstream analyses.

Differential gene expression and functional enrichment analyses

We performed differential gene expression analyses using the DESeq2 R package version 1.22.2

[49] with default parameters. Experimental covariates (sex, age, RIN, PMI, and sequencing

batch) as well as oligodendrocyte and microglia MGPs were incorporated into the statistical

model. Multiple hypothesis testing was performed with the default automatic filtering of DESeq2

followed by FDR calculation by the Benjamini-Hochberg procedure. Analyses were carried out

independently for the two cohorts. Genes were scored according to their significance by

transforming the p-values to account for direction of change. For each gene, the up-regulated

score was calculated as $S_{\text{up}} = 0.5 + \text{sign}(\text{LFC}) \cdot p/2$, and the down-regulated score as $S_{\text{down}} =$

$1 - S_{up}$, where LFC corresponds to the log fold change and p to the nominal p-value of the gene.

Genes were then tested for enrichment using alternatively S_{up} and S_{down} scores employing the

gene score resampling method implemented in the ermineR package version 1.0.1 [50], an R

wrapper package for ermineJ [51] with the complete Gene Ontology (GO) database annotation

[52] to obtain lists of up- and down-regulated pathways for each cohort.

In order to characterize the main biological processes affected by the cell type correction, we

scored pathways based on the loss of significance caused by the addition of cellular estimates to

the gene expression model. We quantified the difference in the level of significance in the up-

and down-regulated enrichment results for each significant pathway as $\Delta = \log(p_0) - \log(p_{CT})$,

where p_{CT} and p_0 are the enrichment p-values for the model with cell types (CT) and without (0),

respectively. Only pathways that were significant in either one of the models were analyzed in

this manner ($p_0 < 0.05$ or $p_{CT} < 0.05$).

The source code for the analyses is available in the GitLab repository

(https://git.app.uib.no/neuromics/cell-composition-rna-pd) under the GPL public license v3.0.

References

1. de Rijk MC, Launer LJ, Berger K, Breteler MM, Dartigues JF, Baldereschi M, et al.
Prevalence of Parkinson's disease in Europe: A collaborative study of population-based cohorts.
Neurologic Diseases in the Elderly Research Group. Neurology. 2000;54:S21-3.

2. Borrageiro G, Haylett W, Seedat S, Kuivaniemi H, Bardien S. A review of genome-wide
transcriptomics studies in Parkinson's disease. Eur J Neurosci. 2018;47:1–16.

3. Jaffe AE, Tao R, Norris AL, Kealhofer M, Nellore A, Shin JH, et al. qSVA framework for
RNA quality correction in differential expression analysis. Proc Natl Acad Sci U S A.
2017;114:7130–5.

4. Dumitriu A, Golji J, Labadorf AT, Gao B, Beach TG, Myers RH, et al. Integrative analyses of
proteomics and RNA transcriptomics implicate mitochondrial processes, protein folding
pathways and GWAS loci in Parkinson disease. BMC Med Genomics. 2016;9:5.

5. Henderson-Smith A, Corneveaux JJ, De Both M, Cuyugan L, Liang WS, Huentelman M, et al.
Next-generation profiling to identify the molecular etiology of Parkinson dementia. Neurol
Genet. 2016;2:e75.

6. Riley BE, Gardai SJ, Emig-Agius D, Bessarabova M, Ivliev AE, Schüle B, et al. Systems-
based analyses of brain regions functionally impacted in Parkinson's disease reveals underlying
causal mechanisms. PloS One. 2014;9:e102909.

7. Adiconis X, Borges-Rivera D, Satija R, DeLuca DS, Busby MA, Berlin AM, et al.
Comparative analysis of RNA sequencing methods for degraded or low-input samples. Nat
Methods. 2013;10:623–9.

8. Gallego Romero I, Pai AA, Tung J, Gilad Y. RNA-seq: impact of RNA degradation on
transcript quantification. BMC Biol. 2014;12:42.

9. Schuierer S, Carbone W, Knehr J, Petitjean V, Fernandez A, Sultan M, et al. A comprehensive
assessment of RNA-seq protocols for degraded and low-quantity samples. BMC Genomics.
2017;18:442.

10. Zhao W, He X, Hoadley KA, Parker JS, Hayes D, Perou CM. Comparison of RNA-Seq by
poly (A) capture, ribosomal RNA depletion, and DNA microarray for expression profiling. BMC
Genomics. 2014;15:419.

11. Huang R, Jaritz M, Guenzl P, Vlatkovic I, Sommer A, Tamir IM, et al. An RNA-Seq
Strategy to Detect the Complete Coding and Non-Coding Transcriptome Including Full-Length
Imprinted Macro ncRNAs. Gaetano C, editor. PLoS ONE. 2011;6:e27288.

12. Srinivasan K, Friedman BA, Larson JL, Lauffer BE, Goldstein LD, Appling LL, et al.
Untangling the brain's neuroinflammatory and neurodegenerative transcriptional responses. Nat
Commun. 2016;7:11295.

13. Alves G, Muller B, Herlofson K, HogenEsch I, Telstad W, Aarsland D, et al. Incidence of
Parkinson's disease in Norway: the Norwegian ParkWest study. J Neurol Neurosurg Psychiatry.
2009;80:851–7.

14. Capurro A, Bodea L-G, Schaefer P, Luthi-Carter R, Perreau VM. Computational
deconvolution of genome wide expression data from Parkinson's and Huntington's disease brain
tissues using population-specific expression analysis. Front Neurosci. 2014;8:441.

15. Mancarci BO, Toker L, Tripathy SJ, Li B, Rocco B, Sibille E, et al. Cross-Laboratory
Analysis of Brain Cell Type Transcriptomes with Applications to Interpretation of Bulk Tissue
Data. eNeuro. 2017;ENEURO.0212-17.2017.

16. Toker L, Mancarci BO, Tripathy S, Pavlidis P. Transcriptomic Evidence for Alterations in
Astrocytes and Parvalbumin Interneurons in Subjects With Bipolar Disorder and Schizophrenia.
Biol Psychiatry. 2018;84:787–96.

17. Bauer M. RNA in forensic science. Forensic Sci Int Genet. 2007;1:69–74.

18. Dickson DW. Parkinson's Disease and Parkinsonism: Neuropathology. Cold Spring Harb
Perspect Med. 2012;2:a009258–a009258.

19. Hauser MA, Li Y-J, Xu H, Noureddine MA, Shao YS, Gullans SR, et al. Expression
profiling of substantia nigra in Parkinson disease, progressive supranuclear palsy, and
frontotemporal dementia with parkinsonism. Arch Neurol. 2005;62:917–21.

20. Miller RM, Kiser GL, Kaysser-Kranich TM, Lockner RJ, Palaniappan C, Federoff HJ.
Robust dysregulation of gene expression in substantia nigra and striatum in Parkinson's disease.
Neurobiol Dis. 2006;21:305–13.

21. Bossers K, Meerhoff G, Balesar R, van Dongen JW, Kruse CG, Swaab DF, et al. Analysis of
gene expression in Parkinson's disease: possible involvement of neurotrophic support and axon
guidance in dopaminergic cell death. Brain Pathol Zurich Switz. 2009;19:91–107.

22. Botta-Orfila T, Tolosa E, Gelpi E, Sànchez-Pla A, Martí M-J, Valldeoriola F, et al.
Microarray expression analysis in idiopathic and LRRK2-associated Parkinson's disease.
Neurobiol Dis. 2012;45:462–8.

23. Edwards YJK, Beecham GW, Scott WK, Khuri S, Bademci G, Tekin D, et al. Identifying
consensus disease pathways in Parkinson's disease using an integrative systems biology
approach. PloS One. 2011;6:e16917.

24. Chandrasekaran S, Bonchev D. A network view on Parkinson's disease. Comput Struct
Biotechnol J. 2013;7:e201304004.

25. Zhang Y, James M, Middleton FA, Davis RL. Transcriptional analysis of multiple brain
regions in Parkinson's disease supports the involvement of specific protein processing, energy

metabolism, and signaling pathways, and suggests novel disease mechanisms. Am J Med Genet
Part B Neuropsychiatr Genet Off Publ Int Soc Psychiatr Genet. 2005;137B:5–16.

26. Papapetropoulos S, Ffrench-Mullen J, McCorquodale D, Qin Y, Pablo J, Mash DC.
Multiregional gene expression profiling identifies MRPS6 as a possible candidate gene for
Parkinson's disease. Gene Expr. 2006;13:205–15.

27. Duke DC, Moran LB, Kalaitzakis ME, Deprez M, Dexter DT, Pearce RKB, et al.
Transcriptome analysis reveals link between proteasomal and mitochondrial pathways in
Parkinson's disease. Neurogenetics. 2006;7:139–48.

28. Zheng B, Liao Z, Locascio JJ, Lesniak KA, Roderick SS, Watt ML, et al. PGC-1α, a
potential therapeutic target for early intervention in Parkinson's disease. Sci Transl Med.
2010;2:52ra73.

29. Flønes IH, Fernandez-Vizarra E, Lykouri M, Brakedal B, Skeie GO, Miletic H, et al.
Neuronal complex I deficiency occurs throughout the Parkinson's disease brain, but is not
associated with neurodegeneration or mitochondrial DNA damage. Acta Neuropathol (Berl).
2018;135:409–25.

30. Pryde KR, Taanman JW, Schapira AH. A LON-ClpP Proteolytic Axis Degrades Complex I
to Extinguish ROS Production in Depolarized Mitochondria. Cell Rep. 2016;17:2522–31.

31. Arneson D, Zhang Y, Yang X, Narayanan M. Shared mechanisms among neurodegenerative
diseases: from genetic factors to gene networks. J Genet. 2018;97:795–806.

32. Cooper-Knock J, Kirby J, Ferraiuolo L, Heath PR, Rattray M, Shaw PJ. Gene expression
profiling in human neurodegenerative disease. Nat Rev Neurol. 2012;8:518–30.

33. Grünblatt E, Mandel S, Jacob-Hirsch J, Zeligson S, Amariglo N, Rechavi G, et al. Gene
expression profiling of parkinsonian substantia nigra pars compacta; alterations in ubiquitin-
proteasome, heat shock protein, iron and oxidative stress regulated proteins, cell
adhesion/cellular matrix and vesicle trafficking genes. J Neural Transm Vienna Austria 1996.
2004;111:1543–73.

34. Moran LB, Duke DC, Deprez M, Dexter DT, Pearce RKB, Graeber MB. Whole genome
expression profiling of the medial and lateral substantia nigra in Parkinson's disease.
Neurogenetics. 2006;7:1–11.

35. Colla E. Linking the Endoplasmic Reticulum to Parkinson's Disease and Alpha-
Synucleinopathy. Front Neurosci. 2019;13:560.

36. Burté F, Houghton D, Lowes H, Pyle A, Nesbitt S, Yarnall A, et al. metabolic profiling of
Parkinson's disease and mild cognitive impairment. Mov Disord Off J Mov Disord Soc.
2017;32:927–32.

37. Luan H, Liu L-F, Meng N, Tang Z, Chua K-K, Chen L-L, et al. LC-MS-based urinary
metabolite signatures in idiopathic Parkinson's disease. J Proteome Res. 2015;14:467–78.

38. Alves G, Muller B, Herlofson K, HogenEsch I, Telstad W, Aarsland D, et al. Incidence of
Parkinson's disease in Norway: the Norwegian ParkWest study. J Neurol Neurosurg Psychiatry.
2009;80:851–7.

39. Gaare JJ, Nido GS, Sztromwasser P, Knappskog PM, Dahl O, Lund-Johansen M, et al. Rare
genetic variation in mitochondrial pathways influences the risk for Parkinson's disease:
Mitochondrial Pathways In PD. Mov Disord. 2018;33:1591–600.

40. Gelb DJ, Oliver E, Gilman S. Diagnostic criteria for Parkinson disease. Arch Neurol.
1999;56:33–9.

41. Ward CD, Gibb WR. Research diagnostic criteria for Parkinson's disease. Adv Neurol.
1990;53:245–9.

42. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30:2114–20.

43. Andrews S. FastQC: a quality control tool for high throughput sequence data. Babraham
Bioinformatics;

44. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. Nat Methods. 2015;12:357–60.

45. Broad Institute. Picard tools [Internet]. Available from: http://broadinstitute.github.io/picard

46. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. Nat Methods. 2017;14:417–9.

47. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000Research. 2016;4:1521.

48. Mancarci O. Homologene (R package) [Internet]. Available from:
https://github.com/oganm/homologene

49. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15:550.

50. Mancarci O. ErmineR (R package) [Internet]. Available from:
https://github.com/PavlidisLab/ermineR

51. Gillis J, Mistry M, Pavlidis P. Gene function analysis in complex data sets using ErmineJ.
Nat Protoc. 2010;5:1148–59.

52. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology:
tool for the unification of biology. Nat Genet. 2000;25:25–9.

Supporting information

S1 Figure. Read mapping efficiency. (A) Proportion of reads uniquely aligned to the genome.

(B) Counts per million (CPMs) for rRNA sequences (PW: ParkWest, NBB: Netherlands Brain

Bank, PA: poly(A) capture cohort). Asterisks indicate significance at (*) $p > 0.05$, (**) $p \leq 0.01$,

(***) $p \leq 0.001$, (****) $p \leq 0.0001$, Wilcoxon test.

S2 Figure. Read mapping statistics. (A) Percentage of bases mapping to different genomic

regions for the two cohorts analyzed using rRNA depletion (PW: ParkWest, NBB: Netherlands

Brain Bank) and the poly(A) selection cohort (PA). (B) Scatterplots of samples show RIN values

versus percent of bases (unaligned, intergenic, intronic and mRNA). Linear regression trend

represented by a black line. Only significant linear regression p-values ($p < 0.05$) are shown in

the panels (PW: ParkWest, NBB: Netherlands Brain Bank, PA: poly(A) capture kit). Asterisks

indicate significance at (*) $p > 0.05$, (**) $p \leq 0.01$, (***) $p \leq 0.001$, (****) $p \leq 0.0001$, Wilcoxon

test.

S3 Figure. Cellular estimates grouped by status. Astrocyte and neuronal MGPs adjusted for all the

experimental variables (sex, age, PMI, RIN, and sequencing batch) and oligodendrocyte and

microglia MGPs, but not for disease status. PW: ParkWest, NBB: Netherlands Brain Bank. P-values

calculated with Wilcoxon test.

S4 Figure. Sample clustering. Samples were marked as outliers if the median correlation with the

other samples was below Q1-1.5*IQR or above Q3+1.5*IQR, where IQR stands for inter-quartile

range. For each of the rRNA depletion cohorts, the heatmaps represent the pairwise correlation

coefficients between the samples based on the log CPM. Marker gene profile (MGP) estimates for

the main cortical cell types are represented on the top of each heatmap, together with RIN, disease

status, and the outlier status for each sample (PW: ParkWest, NBB: Netherlands Brain Bank).

S1 File. Significant genes overlapping PW and NBB.

S2 File. Complete results of the differential expression analyses.

S3 File. ErmineR pathway enrichment analyses before and after correcting for cell

composition.

S4 File. Up- and down-regulated pathways ranked by the change in p-value resulting from

correcting for cell composition.

S5 File. Cohort demographic and experimental information.

S6 File. Read count matrix.

Ethics declarations

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and its

supplementary files. The source code for the analyses is available in the GitLab repository

(https://git.app.uib.no/neuromics/cell-composition-rna-pd) under the GPL public license v3.0.

Competing interests

The authors declare that they have no competing interests.

Funding

This work is supported by grants from The Research Council of Norway (288164, ES633272)

and Bergen Research Foundation (BFS2017REK05).

Author contributions

GSN participated in the study conception, preprocessed the RNA-seq data, designed and

performed the RNA-seq computational analyses. FD participated in the RNA-seq computational

analyses. KH contributed to data interpretation and critical revision of the manuscript. IJ and KP

participated in the conception and design of the study, data interpretation and critical revision of

the manuscript. OBT and GA contributed part of the tissue material, supplementary data, and

critical revision of the manuscript. CT conceived, designed and directed the study, contributed to

data analyses and interpretation and acquired funding for the study. GSN and CT wrote the

manuscript with the active input and participation of LT, KH, IJ, and KP All authors have read

and approved the manuscript.

Figures

Fig 1. Transcript coverage profiles with rRNA depletion compared to poly(A) capture. (A)

Heatmaps of transcript coverage in our two cohorts (PW, NBB) and a poly(A) capture sample

(PA). The y-axis shows samples sorted by RIN (top: lowest RIN; bottom highest RIN). The x-

axis represents the transcript body percentiles (5' to 3'). The shading for a given row represents

the sample-normalized coverage averaged across all transcripts. (B) Boxplots for different

coverage quality metrics: median 5'-bias, median 3'-bias and median coefficient of variation

(CV) for each cohort. The bias metric is calculated by Picard tools on the 1000 most highly

expressed transcripts and corresponds to the mean coverage of the 3'(or 5')-most 100 bases

divided by the mean coverage of the whole transcript (asterisks indicate significance at (*) $p >$

0.05, (**) $p \leq 0.01$, (***) $p \leq 0.001$, (****) $p \leq 0.0001$, Wilcoxon test). The same metrics are

expanded in (C), with sample scatterplots showing RIN values against the coverage quality

metrics. Linear regression trends are indicated with black lines. P-values for the F-statistic of the

linear model are also shown in the panels. Panels are organized in columns (cohorts) and quality

metrics (rows). CV = coefficient of variation; PW = ParkWest cohort; NBB = Netherlands Brain

Bank cohort; PA = poly(A) cohort

Fig 2. Analysis of sample covariates. (A) Pearson correlation coefficients for each pair of variables are shown in correlograms. Sizes of the circles in the upper triangular of the correlograms are proportional to the Pearson correlation coefficient, with color indicating positive (blue) or negative (red) coefficients. The precise values for the Pearson coefficients are indicated in the lower triangular. Non-significant pairwise correlations ($p \geq 0.05$) are represented with a cross. (B) Heatmaps showing the association between the sample variables with the first 5 principal components of the gene expression. Only significant p-values ($p < 0.05$) are represented (linear regression F-test). (C) Cell type estimates based on MGPs for the main cortical cell types controlling for all the experimental variables except disease status (i.e. sex, age, PMI, RIN, and sequencing batch). P-values calculated with Wilcoxon tests. PW = ParkWest cohort; NBB = Netherlands Brain Bank cohort

Fig 3. Functional enrichment. The treemap shows the concordant enriched pathways between

PW and NBB accounting for experimental covariates and MGPs (same direction of gene

expression change and Benjamini-Hochberg FDR < 0.05). Pathways are grouped with a white

border if their gene overlap is above 0.5 (Szymkiewicz–Simpson coefficient). Darker shades of

red/blue represent lower enrichment p-values for up-/down-regulated pathways. Sizes of the

rectangles are proportional to pathway sizes