Title:

# Oscillatory multi-timescale mechanisms underlying multisensory sequence prediction

Running title: Multisensory sequence prediction

Peng Wang[a,*], Alexander Maÿe[a], Jonathan Daume[a,b], Gui Xue[c] and Andreas Karl Engel[a]

[a] Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Martinistr. 52, 20246 Hamburg, Germany

[b] Department of Neurosurgery, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[c] State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research, Beijing Normal University, 100875, Beijing, China

[*] Correspondence address:

Peng Wang, Phone: +49-40-741054680, Fax: +49-40-741057126, E-mail: p.wang@uke.de

Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Martinistr. 52, 20246 Hamburg, Germany.

# Abstract

Neural oscillations have been proposed to be involved in predictive processing by frequency-specific modulation of power, phase, or both. While this is supported by substantial evidence on unimodal processing, only a few studies have addressed the role of oscillatory activity in multisensory predictions. In the present study, we recorded cortical activity using magnetoencephalography during a multisensory serial prediction task in which participants had to acquire stimulus sequences and to monitor whether subsequent probe items complied with the sequence. The sequences comprised combinations of visual and auditory stimuli, but in a given trial, only one modality was task-relevant, and the other had to be ignored. Task-related changes of power and coupling of neural oscillations were analyzed using a data-driven clustering approach. We observed prediction-related theta-band coupling in a network involving the cingulate, premotor, prefrontal, and superior temporal regions. The behavioral performance of participants was associated with the phase delay of the coupling between different regions in this network. Two additional clusters were detected in which beta power was stronger if the previous or the following stimuli were the same as the current one in the sequence. These clusters interacted with the theta-band prediction network through cross-frequency coupling. Our results corroborate the notion that the oscillation phase is important in coordinating brain regions. Furthermore, they offer novel insights into the multi-timescale neural dynamics underlying multisensory predictions, suggesting that oscillations in multiple frequency ranges, as well as coupling within and across frequencies, may be critical for multisensory sequence processing.

# Introduction

While natural events and objects are often jointly detected by more than one sensory system, the perceptual and behavioral consequences of which are often not just the sum of those evoked by each sensory component alone. However, our knowledge is still limited regarding the mechanisms underlying multisensory interactions in the human brain, reviewed in [1-4]. Fewer studies have so far addressed the issue of dynamic neural interactions in multisensory processing [5-9]. Recent EEG and MEG studies have shown that the integration of visual and auditory or visual and tactile information can be associated with changes in phase coherence [5, 8-10]. Furthermore, studies in humans have provided evidence for changes of crossmodal coupling during audiovisual speech perception [6, 11] and during evaluation of the duration of audiovisual stimuli. Evidence is available for the modulation of dynamic crossmodal coupling by factors such as attention, memory, emotion, or expectation [1, 2, 12]. In the present study, we have addressed the modulation of neural dynamics in the context of predictions on upcoming multisensory stimuli.

A role of neural oscillatory activity in predictive processing has been postulated early on and is supported by work on unimodal processing, for review, see e.g. [13-17]. It has been suggested that predictions regarding stimulus identity or stimulus timing may involve oscillatory mechanisms in different frequency ranges. While the latter may be associated with low frequency (delta, theta) oscillations, the former might predominantly involve oscillations in higher frequency (beta, gamma) bands [13, 15]. The relation between oscillatory activity and predictions has been addressed by only a few studies in the context of multisensory processing, for review, see [15, 18]. While some of these studies have addressed temporal predictions [19, 20], the fast dynamics underlying prediction of multisensory stimulus sequences has not yet been studied. Furthermore, large-scale dynamic coupling of oscillatory activity across brain areas has not yet been investigated in the context of multisensory predictions.

Here, we used magnetoencephalography (MEG) to investigate the neural mechanisms underlying the prediction of multisensory stimulus sequences. Participants were presented with 5-item sequences of audiovisual stimulus pairs, which were repeated (see Fig. 1 for more details). At the end of each trial, they had to judge whether the last item matched or violated the sequence. Sequence items were always audiovisual, but only one sensory modality was task-relevant in any given trial. As a control, we used a memory task in which participants were requested to judge whether the last item was identical to the item before. We used this experimental design to investigate how intrinsically generated oscillatory signals are involved in predictions of stimulus sequences, and to what extent the coupling between different brain areas may relate to predictive processing.

**Fig. 1. Experimental paradigm.** (A) Trial structure: visual gratings (upper row) and auditory tones (middle row) were presented simultaneously in a sequential manner (timing in bottom row). The first five items constituted the sample sequence, which was repeated in the trial, with the exception of a potential change in the last item. Note that the last repetition could comprise less than five items: each trial comprised a total of 8 to 20 items. The number of items was randomized across trials, while hazard rates remained constant. When the fixation cross changed its color to red, subjects had to respond by button press. Data were segmented according to the onset of each stimulus and grouped into three categories: sampling segments (1st to 4th items), prediction segments (6th to penultimate item), and response segments (final item). Each stimulus lasted 150 ms, followed by a 550 ms interval. The fixation cross turned red, as a response cue, 1200 ms after the last stimulus. (B) Correct answers for the example trial shown in (A). In the prediction task, subjects had to judge whether the last item matched with the sample sequence in the attended modality; in the memory task, subjects had to indicate whether the last item was identical to the item before (N-back) in the attended modality. In each block, the subjects performed one of the two tasks and attended to either the visual or the auditory modality. The order of tasks and attended modalities was counterbalanced across subjects.

# Results

**Behavioral data**

Participants performed the task with an average hit ratio of 90.5%. Both the hit ratio and response times were submitted to a 2 by 2 ANOVA, with two factors: task (prediction or memory) and attended modality (visual or auditory). For the hit ratio, neither the interaction between the two factors nor the main effect of factor modality was significant. However, the main effect of task was significant ($F$ (1, 28) = 6.80, $p$ = 0.01): the hit ratio was higher in the memory task ($\eta^2_p$ = 0.20) (Fig. 2). There was a significant interaction between the task and modality with respect to the response time ($F$ (1, 28) = 13.31, $p$ = 0.001, $\eta^2_p$ = 0.32). Post-hoc analysis showed that, in the prediction task, responses were slower ($p$ = 0.005, Cohen's $d$ = 2.21) when participants attended to the auditory modality. In the memory task, the difference was even larger ($p$ < 0.001, Cohen's $d$ = 2.27) in the same direction (Fig. 2). The main effect of factor task was also significant ($F$ (1, 28) = 4.52, $p$ = 0.04): the responses were faster in the prediction task ($\eta^2_p$ = 0.14).

**Fig. 2. Behavioral performance**. Comparison of means and their standard errors for both percentage of hits (left) and response time (right) across tasks and attended modalities. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$

**Audiovisual stimuli lead to power changes in multiple frequency ranges**

Since in all conditions we always presented similar audiovisual stimuli (gratings accompanied by tones), we pooled the data across conditions with stratified trial numbers and examined the power change caused by the first stimuli of the sequences, as compared to baseline (100 ms before trial

onset) with the clustering approach. The results are illustrated in Fig. 3. The stimuli generally caused a power increase in most of the brain areas, which was most pronounced in bilateral occipital areas, fusiform gyri, transverse temporal gyri, superior temporal gyri, superior temporal sulci, intraparietal sulci, precunei, and cingulate cortices. The stimulus-related activation occurred in a broad range of frequencies, including the theta, beta, and gamma range. The activations were strongest immediately after stimulus onset and showed a second peak after stimulus offset.

**Fig. 3. Stimulus-related power changes**. (Left) Spatial distribution of the cluster, see method for details. Color intensity is scaled to the integral of *t* values across time bins and frequency bins for each voxel. (Right) Time-frequency distribution of the cluster; color intensity is scaled to the integral of *t* values across all voxels for each time and frequency bin. Time 0 indicates stimulus onset. Inset plots show integrated *t* values over time (top, x-axis: time in ms, y-axis: *t* value) or frequency (right, x-axis: *t* value, y-axis: frequency in Hz).

**Prediction task induces less beta power than memory task**

We observed main effects regarding power differences between prediction and memory tasks in both sampling (permutation test, $p = 0.006$, Fig. 4A) and prediction segments (permutation test, $p = 0.02$, Fig. 4B), in a 2 x 2 ANOVA design with the two factors task and attended modality. Since there was no interaction between these two factors, we will focus on the prediction-related dynamics and present a more detailed analysis of the attention-related effects in a separate paper. The main cluster with task-related power differences involved bilateral superior temporal gyri, fusiform gyri, precentral gyri, insula, cingulate, and frontal cortices. Further analysis revealed that power in this cluster was always stronger in the memory task. In the sampling segments, significant

differences were observed in the alpha and beta bands. In the prediction segments, the alpha component was absent. This alpha component was mainly observed in occipital areas.

**Fig. 4. Power differences across tasks.** The clusters were obtained with 2 x 2 ANOVA design on oscillation power, during both sampling (A) and prediction (B) segments. A significant main effect of factor task was observed. For each cluster, both spatial (left) and time-frequency (right) distributions are illustrated. Color intensity indicates accumulated *F* values. Side plots in the box are integrated *F* values plotted against time (top, x-axis: time in ms, y-axis: *F* value) or frequency (right, x-axis: *F* value, y-axis: frequency in Hz).

**Prediction is associated with stronger theta coupling and weaker gamma coupling**

As the next step, we analyzed functional connectivity, as measured by the imaginary part of coherency [21], which minimizes the influence of volume conduction. We employed the same ANOVA design as for the power analysis in the preceding section. We observed significant coupling differences between prediction and memory task in both sampling (permutation test, $p = 0.018$, Fig. 5A) and prediction segments (permutation test, $p = 0.026$, Fig. 5B). The clusters were similar for both segment types and extended to superior temporal gyri, pre- and postcentral cortices, cingulate cortices, prefrontal areas, insulae, and precunei in both hemispheres. Coupling differences between the two tasks were observed in the theta, alpha, and gamma frequency ranges. Interestingly, further analysis revealed that the theta and alpha coupling was stronger in the prediction task (Fig. 5C and 5D), whereas gamma coupling was stronger for the memory task (Fig. 5E and 5F). The spatial distribution of low-frequency components was similar to that of the high-frequency

components. This suggests that, in these brain areas, theta and alpha coupling supports the prediction task, whereas gamma coupling may be involved in the memory task.

**Fig. 5. Coupling differences between prediction and memory tasks**. (A, B) Main effects of the factor task in a 2 x 2 ANOVA design. (C-F) Results of paired t-test for the same contrast. The left column shows data from sampling segments and the right column shows data from prediction segments. In each panel, the spatial distribution of the cluster is displayed on the left. Color intensity is scaled to the integral of *F* or *t* values across time bins, frequency bins, and connected locations for each voxel. On the right, the time-frequency distribution of the cluster is shown, with color intensity scaled to integral of *F* or *t* values across voxel pairs for each time and frequency bin. Side plots in the box are integrated *F* or *t* values against time (top, x-axis: time in ms, y-axis: *F* or *t* value) or frequency (right, x-axis: *F* or *t* value, y-axis: frequency in Hz). Color bars are shared between panels A, B, C and D, and E and F.

**Cross-regional theta phase delay modulates behavior**

It has been suggested that brain areas communicate via coherent oscillations when the interaction occurs at an optimal phase relation [22, 23]. Evidence supporting this hypothesis has been obtained from neurophysiological recordings in animal studies, specifically at the level of single neurons or small neuronal populations. In the present study, we tested whether this notion is also supported by the analysis of large-scale data recorded with MEG from the human brain. In the observed theta clusters (Fig. 5C and 5D), we determined the optimal phase relation for each component (as defined by a certain location pair, as well as time and frequency bin) as the average phase lag between the two locations at this time and frequency bin, in all segments corresponding to the data used to generate them (correct trials stratified for each condition). We then classified the phase delays relative to the optimal phase relation into 8 bins (from $-\pi$ to $\pi$), where performance data (response time in the same dataset, and percentage of hits with all data included) were pooled and averaged. Thus, we could obtain the modulation of performance by phase delays for each cluster (Fig. 6A-6D). The results suggest that in both sampling and prediction segments, the percentage of hits was highest at zero delay to the optimal phase relation, and then decreased with increasing delay, forming a cosine-like function for the prediction task (Fig. 6A and 6C, red). Similarly, the response time of the prediction task was shortest at zero delay and then increased with phase delay for the prediction segments, resembling an inverted cosine function (Fig. 6D, red). The response time modulation in the prediction task during sampling segments followed the same trends but was less pronounced (Fig. 6B, red). The modulation of percentage of hit and response time in the memory task exhibited a similar but much weaker pattern in the prediction segments (Fig. 6C and 6D, green), but not in the sampling segments.

There were more data segments in the bin with zero delay to the optimal phase relation, and fewer in those with larger delays (e.g., close to $-\pi$ and $\pi$). To rule out the possibility that the difference of phase modulation was due to the amount of data, we sorted the segments into 8 bins, according to

absolute delay to the optimal phase relation, such that within each bin, there were equal number of segments. Subsequently, we submitted the performance per bin to a non-parametric correlation (Fig. 6E-6H), which confirmed that the effects were not caused by variations in the amount of data.

We further conducted a paired t-test on the data associated with the top 25% and bottom 25% delays for each voxel. These results were correlated with the integral $F$ values of each voxel from the clustering approach, which yielded a significant positive correlation (Fig. 6I-6L). This analysis suggested that the more likely a brain location was part of the theta cluster, the stronger the behavioral modulation by the theta phase.

**Fig. 6. Dependence of behavior on theta coupling phase**. Top row: dependence of percentage of hits (A, C) and response time (B, D) on phase delay to the optimal phase relation. Middle row: dependence of percentage of hits (E, G) and response time (F, H) on absolute delay to the optimal phase relation, with an equal number of segments in each bin. Error bar indicates standard error. Spearman's correlation coefficient ($\rho$) is indicated. In panel A – H the mean was removed for all percentages of hits and response times. Bottom row: Correlation between statistical power for percentage of hits (I, K) and response time (J, L) difference, and statistical power of clusters among the voxels. Pearson's correlation coefficient ($r$) was indicated on the plots. The left two columns show data from the sampling segments while the right two columns show data from the prediction segments. *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$

**Beta power reflects stimulus memory and prediction**

To successfully perform the prediction task, participants needed to remember the past stimulus and make assumptions about the next stimulus. To rule out the influence of the physical properties of the preceding or forthcoming stimulus itself, we analyzed a contrast defined by the stimulus context. Each segment was classified by its relation to the preceding segment, either the same or different, in the attended modality. This yielded a 2 x 2 ANOVA with factors modality (auditory vs. visual) and context (same vs. different). We applied our clustering approach with this design and identified one cluster for the main effect of context (relation to previous item, permutation test, $p < 0.001$, Fig. 7A). The cluster mainly involved lateral occipital areas, intraparietal areas, and precunei. It peaked around 400 ms in time and 13.5 Hz in frequency. Further analysis suggested that the power was stronger for the 'same' condition than the 'different' condition. We defined a modulation index (MI) as (same – different) / (same + different). The MIs were always larger than zero (all four comparisons p $< 0.001$, Cohen's $d = 1.41$ for the prediction task in sampling segments, 1.32 for the memory task in sampling segments, 1.95 for the prediction task in prediction segments, 1.53 for the memory task in prediction segments, Fig. 7B), and were stronger during prediction segments than in sampling segments ($F (1, 28) = 4.95$, $p = 0.034$, $\eta^2_p = 0.15$, Fig. 7B). These results suggest that the information on preceding stimuli could be temporarily stored in this network.

A similar analysis revealed another cluster for the main effects of context (relation to the next item; permutation test, $p < 0.001$; Fig. 7D). It mainly comprised pre- and postcentral areas, supplementary motor areas, and lateral prefrontal regions. It peaked around 600 ms after stimulus onset (or 100 ms before the next stimulus) at a frequency of 19.0 Hz. Further analysis suggested that the power was stronger for the 'same' condition than the 'different' condition. MIs in this cluster were not significantly different from zero in sampling segments but were significantly larger in prediction segments ($p < 0.001$, Cohen's $d = 2.17$ for prediction task, $p = 0.007$, Cohen's $d =$

0.76, for memory task, Fig. 7E). It was stronger for the prediction task than memory task ($F$ (1, 28) = 27.01, $p < 0.001$, $\eta^2_p = 0.49$, Fig. 7E). This suggests that, before the onset of the next stimulus, activity in this network might encode the forthcoming sequence items.

**Fig. 7. Power differences as a function of context**. (A - C) Effects of same vs. different preceding item. (D - F) Effects of same vs. different subsequent item. Left column (A, D): spatial (left) and time-frequency (right) distributions of the identified cluster are illustrated. Color intensity indicates accumulated $F$ values. Side plots in the box are integrated $F$ values over time (top, x-axis: time in ms, y-axis: $F$ value) or frequency (right, x-axis: $F$ value, y-axis: frequency in Hz). Color bars are the same for panel A and D. Middle column (B, E): Modulation indices for same vs. different context conditions. Right column (C, F): Phase amplitude coupling between significant locations shared between the coupling cluster and the context clusters, expressed as difference to the shift predictor. *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$

## Multiple clusters are linked by phase amplitude coupling

A key question is how these different clusters relate to each other, i.e. how the respective neural populations interact to accomplish prediction. One possible mechanism is that the context from preceding items (prior knowledge) is accumulated to form a sequence representation, which is then used to predict the next item (consequence). Thus, the two beta power clusters may reflect the

sources and outcomes of prediction, while the theta coupling cluster might serve as the link, mediating the predictive process. To test this idea, we examined cross-frequency coupling for the areas shared by these clusters, specifically investigating whether the theta phase of the coupling cluster modulated the beta power of the context cluster. We observed that in all cases cross-frequency coupling was stronger than a shift predictor computed as a control (Fig. 7C). Notably, the cross-frequency coupling was stronger for the prediction task than the memory task only in the prediction segments (Fig. 7F).

**Sequence continuation vs. violation induces differences in both power and coupling**

Finally, we examined the differences in power and coupling between sequence continuation and violation for the response segments using paired $t$-tests. The observed power differences are shown in Figure S2. The first cluster (permutation test, $p < 0.001$, Fig. S2A) exhibited stronger power for sequence continuation. The major frequency components of this cluster were in the theta-alpha range and peaked around 900 ms. The activated areas in this cluster included the precunei, cingulate cortices, pre-and postcentral areas, frontal areas, superior temporal gyri, middle temporal sulci, transverse temporal gyri, intraparietal sulci and insulae in both hemispheres.

The second cluster (permutation test, $p < 0.001$, Fig. S2B) involved the anterior cingulate cortices in both hemispheres, and superior temporal cortex, insular, pre-and postcentral areas, inferior precentral sulcus in the right hemisphere, where the theta power was stronger for sequence violation after stimulus onset; this effect peaked around 300 ms.

The third cluster (permutation test, $p < 0.001$, Fig. S2C) included middle cingulate cortex, superior temporal cortex, insular, pre-and postcentral areas in the right hemisphere, where the beta power was stronger for sequence violation just before the response cue; this difference was maximal around 1100 ms after stimulus onset.

We also examined the coupling difference, as measured by the imaginary part of coherency, between sequence continuation and violation for the responding segments with a paired $t$-test. The results are shown in Fig. S3. The first cluster (permutation test, $p = 0.001$, Fig. S3A) exhibited stronger coupling for stimulus continuation in the theta range, peaking around 1000 ms. Activated areas in this cluster included middle cingulate cortices, pre-and postcentral areas and transverse temporal gyri in both hemispheres, and superior temporal gyri, insulae, and superior frontal gyri mainly in the right hemisphere.

The second cluster (permutation test, $p = 0.005$, Fig. S3B) included middle cingulate cortices, pre- and postcentral areas, inferior frontal areas and insulae in both hemispheres and superior temporal cortex mainly in the right hemisphere, where the theta coupling was stronger for sequence violation after stimulus onset; this effect peaked around 400 ms.

# Discussion

In the present study, we have recorded MEG during a serial prediction task in which participants had to acquire stimulus sequences and to monitor whether subsequent probe items complied with the sequence. Presentation of the audiovisual stimulus items revealed activation in a widely distributed set of areas involving early and higher-order visual and auditory areas as well as intraparietal sulci, precunei, and cingulate cortices. Task-related changes of power and coupling of neural oscillations were observed in several frequency bands. We identified prediction-related theta-band coupling in a network involving cingulate, premotor, prefrontal, and superior temporal regions. Behavioral performance of the participants correlated with phase delays between remote regions in this network. In this network, beta power was stronger when either the previous or upcoming stimuli matched the current item in the sequence. Furthermore, the amplitude of these beta signals was modulated by the phase of theta activity, as revealed by analysis of cross-frequency coupling. Overall, our findings suggest that oscillations in multiple frequency ranges, as well as coupling within and across frequency bands, play a crucial role in sequence processing. Given that no interactions between the prediction-related activity and attention-related activities were detected in our data, the attention effects observed with our paradigm will be presented in a separate paper.

Prediction is a key feature of brain function. Based on previous experience, the brain continuously generates predictions about upcoming events, which are then adapted through learning in order to reduce prediction errors [16, 24]. A role of neural oscillations in predictive processing has been postulated early on and is supported by work on unimodal processing, for review see e.g. [13-17]. It has been suggested that predictions regarding stimulus identity or stimulus timing may involve oscillatory mechanisms in different frequency ranges. While the latter may be associated with delta and theta band oscillations, the former might be mediated by beta and gamma band interactions [13, 15].

So far, only relatively few studies have addressed the relation between oscillatory activity and predictions in the context of multisensory processing, for review, see [15, 18]. One of the earliest studies was carried out by Widmann and coworkers [25], who used a matching paradigm in which visual score-like symbols predicted corresponding sound patterns. Compared to incongruent sounds, expected sounds evoked higher gamma-band activity. Van Ede et al [26] presented subjects with a repeated series of concurrent tactile and auditory stimuli in which rare deviant stimuli occurred in one of the two modalities. Interestingly, prediction-related modulation of beta-band oscillations was observed even for unattended tactile stimuli. This modulation was enhanced during states of attentive expectation. Arnal et al [27] investigated a violation of multisensory predictions in speech perception and demonstrated that correct predictions were associated with delta oscillations, whereas prediction errors were predominantly accompanied by beta- and gamma-band activity. Another recent study on crossmodal effects in speech perception [28] has observed modulation of theta-oscillations during auditory speech processing by concomitant visual input from observation of gestures. Low-frequency oscillations have also been found to be associated with temporal prediction in a study where a visual color change predicted the occurrence of an auditory stimulus [20]. A more recent study has observed a phase reset of delta oscillations in a task involving visual-to-tactile temporal predictions [29]. Altogether, only a few studies are currently available that have addressed changes in oscillatory activity during multisensory prediction.

To our knowledge, this study presents the first investigation of neural oscillations and their coupling in networks underlying prediction of multisensory stimulus sequences. In the analysis of task-related power changes, we observed significant differences between the prediction task and the memory control condition. The cluster reflecting these differences was mainly found in the beta range, comprising bilateral superior temporal gyri, sensorimotor cortices, insulae, cingulate and frontal cortices. In addition, there was an alpha component during sampling of the sequence but not in the prediction phase of the trials. This alpha component, which was mainly observed in visual areas, was more pronounced for the memory control task. This suggests that functional inhibition,

as indicated by the alpha oscillations [30], was stronger during the sampling phase in the memory task. In this control condition, the information present in the sampling segments was task-irrelevant, whereas it was vital to construct the sequence in the prediction condition. Thus, it was not surprising that functional inhibition occurred for visual items in the memory task, while inhibition of visual processing was released during the sampling phase in the prediction task. Beta power was stronger during the memory task than during the prediction task, in line with the proposal that beta-band oscillations reflect the maintenance of the task-relevant information in a cognitive state[13]. In the prediction task, the participants needed to constantly update the sequence information after the presentation of an item and predict the upcoming sensory inputs in the interval before the occurrence of the subsequent item; this cognitive updating might relate to the observed reduction of beta power.

In the same network, we also observed power differences in the responding segments, where subjects had to indicate whether the last item continued or violated the acquired sequence. These data suggest a beta power decrease upon violation of the sequence, with a subsequent beta rebound. This dynamic of beta power shows similarities to changes associated with prediction errors, as proposed in recent predictive coding frameworks [15, 16].

In addition to the task-related power differences, we also observed functional connectivity clusters defined by coupling differences between the prediction and the memory task. All clusters were similar in their spatial distribution, and they also shared a large proportion of the involved brain regions with the cluster obtained by comparing power difference. This suggests that these clusters, although formally derived in separate analysis steps, reflect prediction-related dynamics in the same cortical network. This prediction-related network comprised mainly superior temporal gyri, sensorimotor cortices, insulae, cingulate and frontal cortices. in both hemispheres. An interesting observation was that motor and pre-motor areas were quite prominent in the prediction network. This is in line with previous fMRI studies that have demonstrated a critical involvement of motor and premotor regions in prediction tasks[15, 31, 32]. In this network, coupling in the theta-alpha

range was stronger for the prediction task compared to the memory task during both the sampling and the prediction phase of the trials, suggesting that coupling at low frequencies was essential for the prediction process. Interestingly, in the same network, coupling also occurred in the gamma frequency range, which was stronger for the memory condition. A possibility is that, in the memory task, the comparison of low-level visual or auditory features promoted gamma coupling, which has been related to feedforward transfer of sensory information[13, 33]. In contrast, in the prediction task, the information on the 5-item sequence needed to be constantly retrieved and updated, which could require more feedback communication that may primarily be conveyed by coupling of low-frequency oscillations [33, 34]. Interestingly, changes in theta-band coupling were also related to the processing of sequence continuation vs. violation in the response segments of the prediction trials.

Oscillatory coupling has been proposed to mediate communication between remote areas by providing dynamic windows of phase relations optimally geared to permit signal transmission[14, 23, 34-36]. This notion implies that behavioral performance should also be associated with changes in the phase relations in the network and more efficient at particular phases of coupling, but impaired at other phase relations between the recorded sites [22, 34]. We examined how the behavioral performance was tuned by the theta phase in the prediction-related network identified in our cluster analysis. In the prediction condition, both accuracy and response times exhibited a sinusoidal tuning by the theta phase in the prediction phase of the trials. This effect was much weaker in the memory trials. In the prediction task, the participants needed to acquire the information on the sequence and, thus phase relations may have been more strongly employed to coordinate signals across the involved brain areas. In contrast, the memory task possibly did not demand the same amount of coordination across regions of the prediction network. Similar effects have been demonstrated in cellular recordings in animal studies [23, 37]. A recent study showed that when phase relation between V1 and V4 neurons was optimal, monkeys tended to perform faster, during a visual attention task [22]. Our study is, to our best knowledge, the first

demonstration of this sinusoidal tuning of behavior by the phase of oscillatory coupling at the level of whole-brain recordings. Overall, our results strongly support the notion that the phase of neural coupling is of key relevance for coordination of the information flow between separate brain regions [35, 36].

In a serial prediction task, preceding stimuli should act as priors for processing of the current item in the sequence which, in turn, should influence the processing of subsequent stimuli. We therefore investigated whether power changes occurring during the prediction task also reflected the stimulus context, i.e., whether the preceding stimulus would have a contextual effect on the current one, and whether processing of the current item already predicted processing of a subsequent item. Both were observed to be the case and to be reflected in changes of beta power, which was larger if the same item appeared before or after the current item, suggesting that the power difference related to differences in the expectation and storage of information in the network. The storage effect was more pronounced during the prediction than during the sampling phase, possibly reflecting that participants needed to actively access the remembered information of previous items to update the prediction of the forthcoming events. The expectation effect could only be observed in the prediction phase – only when they knew the sequence. Interestingly, this effect mainly involved the motor, supplementary motor, and premotor regions. This strongly supports the idea that prediction is an active process, which relies on areas involved in action control even in the absence of an overt motor response [14, 31]. The results of our phase-amplitude coupling analysis for the theta and beta signals in the prediction network suggest that beta power, through coupling to the local theta phase, might dynamically modulate the information transfer through the long-range theta phase-phase coupling. Similar phase-amplitude coupling between theta and beta signals has been observed in previous studies, e.g., in visual working memory paradigms [38].

Taken together, the power and coupling changes observed for the prediction-related network suggest the following possible scenario underlying the processing of sequence information in our task setting: In the sampling phase, the encoding of the presented items may be differentially

modulated by changes in beta power already in sensory regions, depending on the task (prediction vs. memory) and on the contextual information. In the prediction phase of the trials, incoming new information is compared with the stored information on the sequence, which induces further modulation of local beta power. Local phase-amplitude coupling between beta and theta signals may then shape the transfer of the task-relevant information by long-range theta band coupling in the prediction network. After the transfer of prediction-related information, through theta phase coupling, to target motor-related areas, the information might be recoded again through theta phase to beta power coupling, which then might have an impact on response selection. By demonstrating the involvement of several frequency bands in the multisensory prediction task studied here, our results support recent concepts on multi-timescale information processing which emphasize the need to deploy different time windows for segregation and integration of information [4, 39].

In conclusion, we have investigated the neural mechanisms of multisensory sequence prediction. Our data provide direct evidence that oscillations and phase coupling across remote regions play an important role in prediction, and might serve as a general mechanism for cross-regional communication and collaboration. In particular, our results support the notion that the phase of oscillatory signals can sinusoidally modulate behavioral performance. Furthermore, our data underscore the relevance of cross-frequency coupling for establishing multi-timescale processing dynamics. These findings substantially corroborate and extend previous studies on the functional role of coupled oscillations in large-scale communication in the brain.

# Materials and Methods

## Participants

29 healthy volunteers (17 females, age 26.3 ± 4.2 years) participated in this study. They gave written informed consent prior to commencing the experiment. All participants received 10 €/hour remuneration, and an additional 30 € bonus if they completed all sessions. All volunteers had a normal or corrected-to-normal vision and were right-handed and free of psychiatric or neurological disorders. The experiment was carried out in accordance with the guidelines and regulations, approved by the ethics committee of the Medical Association Hamburg.

## Experimental procedure

A green fixation cross (0.25˚ visual angle) was shown at the center of the screen, and participants were asked to maintain fixation during the stimulation. A set of 5-item-long binary visual-auditory sequences (32 possibilities) was presented to the participants (Fig. 1). Each item appeared for 150 ms, followed by a 550 ms break before the next item. An item was a vertical or horizontal Gabor patch (10˚ visual angle, 0.5 cycles per degree), accompanied by a simultaneously presented sine wave tone which was either high (2000 Hz) or low (1800 Hz) in pitch. The volume of the tone was adjusted to 30 dB above the individual hearing threshold. In each trial, only one of the sequences would be presented and repeated; all sequences were presented with equal probability for all participants. The length of the trials was randomized between 8 and 20 items, with an approximately fixed hazard rate. In the prediction task, participants had to monitor all items and to indicate, by button press, whether the last item of the trial violated the sequence. For different blocks of trials, participants had to attend to either the auditory or the visual stimuli while ignoring the respective other modality. Although only the stimulus changes in the attended modality defined the task-relevant sequence, the combination of visual and auditory features was kept constant except for the last item (Fig. 1). In the control memory task, the physical stimuli were exactly the

same as in the prediction task, but participants had to indicate whether the last item was the same as the item second (1-back) or third (2-back) from last in either the visual or the auditory modality. In the 1-back condition, the participants performed more accurately in the memory task and there was no difference in response time, as compared to the prediction task. In the 2-back condition, the accuracy was the same while the participants were slower in the memory task, as compared to the prediction task. Since there was no substantial difference in the MEG data for these two sets of data, we merged them in the formal analysis. Overall, the study had a two by two design: task (prediction or P, vs. memory, or M) x relevant modality (visual or V, vs. auditory or A). There were 32 test trials in a block, and each participant accomplished 16 blocks for one recording session. Each block was initiated by the participants, who usually took a 0.5 ~ 2 minutes break between blocks. A mandatory break of about 20 minutes was taken in the middle of each session. The order of blocks in a session was counterbalanced with respect to task and modality across participants. For each participant, two sessions were recorded on separate days, no more than 6 days apart. Before the first MEG recording session, a practice session comprising 4 blocks was conducted outside of the MEG chamber to familiarize participants with the task. The combination of visual and auditory features was different between the practice session and the MEG recordings. Only the data from the MEG sessions were used for subsequent analysis.

The main experiment was conducted in a dimly lit, sound attenuated MEG chamber. The participants were seated in a recording chair with armrest. Visual and auditory stimuli were generated from the same computer (Dell Precision T5500) equipped with Psychtoolbox-3 [40] and Matlab (R2012b, MathWorks, Natick, MA), on a Windows 7 Professional 64-bit operating system. The visual stimuli were delivered from a projector outside of the chamber via a mirror system to a thin film screen mounted 60 cm in front of the participants' eyes, with a 100 Hz refresh rate. The auditory stimuli were delivered with MEG-compatible in-ear headphones (SRM-2525, STAX Limited, Fujimi, Japan). The button presses were recorded and merged to MEG data with an fORP system (HHSC-2X2, Current Designs Inc., Philadelphia, USA).

**MEG recording and preprocessing**

MEG signals were recorded with a 275-channel whole-head system (CTF MEG International Services LP, Coquitlam, Canada) in a magnetically shielded chamber, together with electrical eye, and cardiac activity via Ag/AgCl-electrodes in order to control for endogenous artifacts. Three magnetic sensors fixed at nasion, left and right preauricular points were used to track the head position, which was used for both online monitoring [41] and offline artifact rejection. At the beginning of each block, participants were asked to reposition their head to the original position if the deviation was more than 10 mm. The recorded signals were digitized at a sampling rate of 1200 Hz. After removing malfunctioning channels, we obtained data from 274 sensors for each participant.

Raw data were read into Matlab with functions provided by FieldTrip [42]. Extreme noisy events (e.g., squid jumps) were detected, and data within 1-second epochs around the event were excluded from further analysis. Data were then bandpass filtered between 0.5 ~ 140 Hz with a Butterworth filter at the order of 4, in both forward and reverse direction for zero phase distortion. Both semi-automatic z-score based artifact detection (threshold $z > 7$) and visual inspection were applied to label noisy segments for exclusion. The average head position of each session of each participant was computed and used for later source reconstruction. Epochs where the head position deviated from this position more than 5 mm were also excluded. Instead of a notch filter, we removed 50 Hz line noise with sinusoidal signal estimation [43]. The remaining data were submitted to ICA [44] for artifact detection. To decide on the number of ICA components, a PCA was performed. We took the minimum number n, such that the first n PCA components preserved at least 99% of the whole information. Eye movements, cardiac artifacts, remaining 50 Hz line noise, and muscle activities were identified with their distinct spatial extension, spectral signature, waveform properties, and distribution across trials [45] and then removed from the data.

**Event extraction and MEG data segmentation**

Time points of stimulus onsets and responses were extracted from the MEG data, and consistency with the events recorded by the stimulus software was verified. Behavioral performance such as hit ratio and response time was calculated based on these events.

The MEG data were segmented based on these events in several categories: (i) Onset segments, ranging from -700 to +700 ms relative to the beginning of each trial. This category was used to test neural responses to the physical stimulus properties compared to the pre-trial baseline. (ii) Sampling segments, ranging from 0 to 700 ms with reference to the onset of the $1^{st}$ to $4^{th}$ items of each trial. These segments were used to analyze neural data during the acquisition of the sequence. (iii) Prediction segments, ranging from 0 to 700 ms relative to the onset of the $6^{th}$ to the second last item of each trial. This category was used to investigate how the participants predicted the sequences based on the acquired information. We did not include data segments around the $5^{th}$ item in the analyses because there might be confounding components for both learning and predicting: after the onset of the $5^{th}$ item, sampling information on this item and predicting the next one could not be disentangled. (iv) Response segments, ranging from 0 to 1500 ms relative to the onset of the last item. These segments were used to analyze response-related aspects. A segment was rejected if it overlapped with any previously labeled data exclusion period. Data were down-sampled to 400Hz after segmenting the cleaning data.

**MRI data acquisition and pre-processing**

T1-weighted structural MRI scans were obtained for 27 of the 29 participants. MRI recording was performed with a Magnetization Prepared Rapid Gradient Echo (MP RAGE) sequence on a Siemens 3 T scanner (Magnetom Trio, Siemens, Erlangen, Germany). The acquired images were converted from DICOM to NIfTI (Neuroimaging Informatics Technology Initiative) format and then processed using the FreeSurfer software package [46] for cortical segmentation, brain surface

inflation and head shape reconstruction. Subsequently, the nasion and the left and right preauricular points were manually labeled on the reconstructed head shape and co-registered with coordinates of these three landmarks obtained from the MEG system with the aid of the MNE software package [47]. This procedure put the MEG sensors and the structural brain into the same coordinate system.

**Source reconstruction**

We constructed our source space based on the FreeSurfer averaged template brain [46]. To generate the source space, we started from an icosahedron and then recursively subdivided its edges, the middle point of which would be repositioned to its circumscribed sphere as new vertices. The defined sphere, with 162 vertices on it, was morphed onto the outer surface of the white matter (i.e., the inner surface of the gray matter) of each hemisphere, which yielded 324 locations homogeneously covering the entire cortex. By excluding locations that were unlikely to elicit significant brain responses (e.g., the corpus callosum), we reduced the number of source locations to 290 (Fig. S1). This procedure was accomplished with the MNE software package [47]. We chose a relatively low number of source locations for reasons of computational efficiency and in order to be compatible with the physical spatial resolution of the MEG sensors. These source locations were then mapped to each individual anatomical MRI. For those participants without an anatomical MRI, we used the template brain ICBM152 from the Montreal Neurological Institute [48] instead. As a next step, we computed lead fields with these source locations and individual sensor positions for each session. Lead field calculation was realized with a single-shell head model [49]. We did not apply compensation for ICA component reduction with the assumption that the lead field distribution was not altered by removing components related to eye movements or other muscle activity [50]. Making use of linearly constrained minimum variance (LCMV) methods [51], we projected the original sensor-level waveforms to the source locations defined above. For each source location, we computed three orthogonal filters (one for each spatial dimension) that passed activity from the location of interest with unit gain while maximally suppressing activity from all

other locations. Next, we linearly combined the three filters into a single filter in the direction of maximal variance [5]. In order to avoid spurious effects resulting from unequal filters in between-condition comparisons, we used data from all conditions (with the same number of trials for each condition) to compute a common filter. To derive source estimates, we multiplied the cleaned sensor data with the real-valued filter. Then we merged data from different sessions of the same participant. High source correlations can reduce source amplitudes estimated with beamforming due to source cancellation [5, 51] which might influence the magnitude of cortico-cortical coupling estimates. However, within the range of physiological source correlations [52], the identification of cortico-cortical coupling using the beamforming method is possible [53]. Moreover, as source cancellation might reduce the sensitivity to detect coupling, it should not lead to false positive results.

**Statistics**

In general, we performed ANOVA for statistics on multiple factors, and paired t-test for pairwise comparison within one factor, in the analyses of behavioral and MEG data. Besides providing significance measures (e.g., $p$ values), we also show measures of effect sizes ($\eta^2_p$ for ANOVA or Cohen's $d$ for pairwise comparison).

**ERF computation**

For each task (prediction, memory) and modality (visual, auditory) combination (i.e., four conditions), we averaged the waveforms in source space, with trial number stratified for the conditions, to obtain event-related fields (ERF) during each type of segment (onset, sampling, prediction, response).

**MEG time-frequency decomposition**

The time-frequency decomposition was performed in cortical source space. To avoid computing the same effect twice, we removed the ERF from each data segment prior to the spectral analysis of induced activity. We also computed an averaged ERF of all conditions in the onset segments, from which we evaluated the evoked activity by physical stimuli. All spectral estimates were obtained with the multi-taper method [43] and computed across 19 logarithmically scaled frequencies from 5.66 to 128 Hz with 0.25-octave steps. In the induced data, we took time points in steps of 100 ms (the time range of each segment was defined as described in "Event extraction and MEG data segmentation"). In the evoked data, we took time points in steps of 25 ms. We characterized responses relative to the pre-stimulus baseline using the bin centered at -100 ms. The temporal and spectral smoothing was performed as follows. For frequencies larger or equal to 16 Hz, we used temporal windows of 250 ms and adjusted the number of Slepian tapers to approximate a spectral smoothing of 3/4 octave; for frequencies lower than 16 Hz, we adjusted the time window to yield a frequency smoothing of 3/4 octaves with a single taper [5]. In case the window extended outside of the segment range, zero padding was applied. The employed time-frequency transformation ensured a homogenous sampling and smoothing in time and frequency, as required for subsequent network identification within this multi-dimensional space.

**Power estimation and phase coupling analysis**

Power of oscillatory signals was estimated from the spectrum $X(f, t)$ as follows:

$$Power_x(f, t) = \frac{1}{n} \cdot \sum_i X_i(f, t) \cdot X_i^*(f, t) \tag{1}$$

Here $i$ is the trial index, $t$ is the time bin index, and $f$ is the frequency bin index, $*$ represents the complex conjugate.

To quantify frequency-dependent synchronization in source space, we estimated the imaginary part of coherency (*ImC*) – a measure with minimum influence of volume conduction [21]. The *ImC* between pairs of signals *X (f, t)* and *Y (f, t)* was computed according to the following equation:

$$ImC_{xy}(f,t) = Im\left(\frac{1}{n} \cdot \sum_i \frac{X_i(f,t) \cdot Y_i^*(f,t)}{\sqrt{(X_i(f,t) \cdot X_i^*(f,t)) \cdot (Y_i(f,t) \cdot Y_i^*(f,t))}}\right) \qquad (2)$$

Here *i* is the trial index, *t* is the time bin index, and *f* is the frequency bin index, * represents the complex conjugate, *Im* indicates imaginary part of complex numbers.

**Identification of networks by clustering**

The general approach of our network identification closely resembles the method described in [5, 8] and is summarized below. An interaction between two cortical areas can be represented as a point in two-dimensional space and can be extended to four-dimensional space by adding the dimensions time and frequency. Identifying networks of significant interaction is then equivalent to identifying continuous clusters within this 4-dimensional space. In our approach, we first computed the ImC for all pairs of locations in all time and frequency bins and conditions with equation (2). Then t-statistics (for pair-wise comparison) or ANOVA (for multiple-factor analysis) was performed across participants. The resulting multi-dimensional matrix was then thresholded with a *t* or *F* value (e.g., equivalent to $p = 0.01$), setting every connection to either 1 (connected) or 0 (not connected).

Further, we performed a neighborhood filtering using threshold (e.g., 0.3) to remove spurious connections: if the existing neighboring connections of a certain connection were fewer than (e.g., 30%) of all its possible neighboring connections, this connection would be removed. As one connection was defined by four numbers, representing its frequency, time, and the two locations of a pair, two connections were defined as neighbors if they were identical in three of the four numbers, and the other value would only differ by one step, i.e., 100 ms in time, 0.25 octaves in frequency, or immediate neighborhood in source space. Two source locations were defined as immediate neighbors if they were directly connected in the regular polyhedron during source space

construction. Eventually, the remaining neighbor connections formed the candidate clusters for statistics. We defined their sizes as the integral of the $t$ or $F$ values across the volume, time and frequency. To evaluate their significance, random permutation statistics was performed to account for multiple comparisons across the interaction space. To this end, the cluster-identification approach was repeated 1000 times with shuffled condition labels to create an empirical distribution of cluster sizes under the null-hypothesis of no difference between conditions [54]. The null-distribution was constructed from the largest clusters (two-tailed) of each resample, and only clusters with sizes ranked top 5% in these null distributions were considered as significant (i.e., $p = 0.05$) in the comparison. The final clusters corresponded to cortical regions with different synchronization status among comparisons, and were continuous across time, frequency, and pairwise space. Applying this method, we compensated for the relatively arbitrary thresholding procedure during the generation of the 4-D connectivity array: the clusters with fewer statistically stronger connections and clusters with more statistically weaker connections were both considered.

The identification of power difference clusters was similar to the above except that there was only one dimension in space rather than two for coupling.

**Identification of clusters for various contrasts**

Using the methods described above, we applied the cluster identification approach for the analysis of several contrasts. We identified the cluster of regions showing stimulus-related power changes by contrasting the evoked power from each time bin between 0 and 700 ms after the onset of the first item of each trial, with the power averaged between time bins between 125 and 75 ms before stimulus onset, with a pairwise t-test. Here, we use a threshold of $t$ values corresponding to $p = 0.01$; the threshold for the neighborhood spatial filter was 0.3, i.e., at least 30% of neighboring difference had to be significant.

To investigate differences of induced oscillation power between conditions, we employed a 2 x 2 design ANOVA, with factors task (prediction vs. memory) and attended modality (visual vs.

auditory). Here we use a threshold of $F$ values corresponding to $p = 0.05$ for the main effects and interactions. The threshold for the neighborhood spatial filter was 0.3. This means that a test with a significant difference would be still regarded as significant after filtering, only when at least 30% of the neighboring tests were significant. Neighborhood was defined as data differ only one bin in time or frequency, or only one location was different but still was immediately connected of the location in the inspected tests. In this case, orphan-like significant points would be removed. A post-hoc test was applied for each component of the cluster to test the direction of the difference. If directions of effects were not consistent within a cluster, the cluster was split to separate the positive and negative effects. The same approach was used to investigate differences of coupling between conditions.

In order to investigate the effects of stimulus context, we classified the segments according to its previous or next stimulus, which could be the same or different. With this factor and the attended modality, we construct another 2 x 2 ANOVA design. This was submitted to the cluster identification with the threshold of $F$ values corresponding to $p = 0.01$ for the main effects and interactions; the threshold for the neighborhood spatial filter was set to 0.3, i.e., at least 30% of neighboring differences had to be significant. Again, a post-hoc test was applied for each component of the network to test the direction of the difference. If appropriate, the cluster was split to separate the positive and negative effects.

Furthermore, we tested both the power and coupling difference between sequence continuation and violation using paired t-tests and the results were submitted to the clustering approach with the threshold of $t$ values corresponding to $p = 0.01$; in this case, the threshold of neighborhood spatial filter was also set to 0.3.

**Illustration of identified clusters**

To visualize networks identified by the clustering approach, we projected them separately onto two subspaces. We integrated the $t$ or $F$ values among connections over all spatial locations for each

time and frequency bin to illustrate, when and at which frequencies a cluster was active irrespective of the spatial distribution. Complementary to this, we also computed the integral of $t$ values in the connection space over time, frequency, and target locations (only for coupling clusters) for each location in the corresponding cluster. The result was then displayed on the template brain surface to reveal the spatial extent of the cluster independent of its intrinsic coupling structure and time-frequency characteristics. For illustration purposes, the data were interpolated from 290 to ~300,000 vertices on the inflated brain surface. The labeling of brain areas involved in our clusters was based on their relative position to the gyri and sulci of the template brain [55].

**Performance modulation by theta phase within the prediction network**

We tested the modulation of behavioral performance by the phase of the coupling in the theta band clusters related to prediction (Fig. 5C and 5D). We defined the optimal phase relation of the coupling by averaging the segments used to quantify the cluster (all corrected and stratified for conditions) for each component. Each segment was classified into one of the eight bins based on its delay to the optimal phase relation. We averaged the response time from the same data for each component and then averaged these across the whole cluster. Since here the delay was a relative measure (difference to optimal), we averaged the tuning of response time by phase across participants.

We performed a similar analysis as the above to test for effects on accuracy, with two main differences as follows. First, we had to include all the data for computation of the percentage of hits, not only the data that were used to generate the network (corrected and stratified). Second, we used half of the data with correct responses, to generate the optimal phase relation and classify half of the full data, which would not include the trials to generate the optimal phase relation, and this was done as well for the respective other half of the data.

To rule out an influence of the number of segments, which was significantly larger for zero phase delay than a delay of $\pi$, we re-classified the data by absolute values of phase delay to the

optimal phase relation, where the spacing was not exactly evenly distributed, e.g., the data range was smallest for the bin closest to zero so that it could include the same number of segments as the bin farthest from zero, which was much wider. Then a non-parametric correlation was computed to test whether the data followed the same trends. Spearman's rho was calculated.

We also took the top and bottom 25% of data based on the absolute values of difference to optimal phase relation, and then compared the difference of performance with a paired t-test. We pooled the $t$ values for each voxel. Then we also took the $F$ values of each voxel when generating the cluster. The two groups of values were correlated, and Pearson's $r$ was measured.

**Cluster analysis based on stimulus context**

As mentioned in section "Identification of clusters for various contrasts", we detected two clusters related to stimulus context, i.e., the occurrence of the same versus different stimulus as the previous or next item respectively. We calculated the modulation index (MI) by dividing the difference of 'same' and 'different' conditions and then dividing by their sum. MIs were also computed for the corresponding data for sampling segments and memory task, and compared using a 2 x 2 ANOVA.

**Relation between clusters and cross-frequency coupling**

To investigate interactions among clusters, especially between the theta coupling cluster and the clusters defined by the stimulus context, we computed their cross-frequency coupling. We first chose the locations that were most significant in both clusters, i.e., at least 1 standard deviation higher than the mean for the integrated $F$ values. Then we computed the theta (4 - 7 Hz) phase of these voxels in the theta networks and the beta (12 - 32 Hz) power of these voxels in the beta clusters. The cross-frequency coupling among these time courses was computed with the modulation index [8, 56]. As a control, we computed a shift predictor by shuffling the trials, such that the phase from trial A would be coupled with trial B, with all other intact. In this case, the stimulus-locked information would be preserved but the instantaneous correspondence of phase and

amplitude was removed. Subsequently, the cross-frequency coupling of real data was normalized with those obtained from shift predictors. Then they were compared across conditions with ANOVA.

# Acknowledgements

We thank Guido Nolte for methodological discussions and Till Schneider for valuable comments on the experimental design. We thank Roger Zimmermann, Christiane Reissmann, and Karin Reimann for help with MEG data collection and participant recruitment.

# References

1.  Senkowski D, Schneider TR, Foxe JJ, Engel AK. Crossmodal binding through neural coherence: implications for multisensory processing. Trends Neurosci. 2008;31(8):401-9.

2.  Talsma D, Senkowski D, Soto-Faraco S, Woldorff MG. The multifaceted interplay between attention and multisensory integration. Trends Cogn. Sci. 2010;14(9):400-10.

3.  Bauer AKR, Debener S, Nobre AC. Synchronisation of neural oscillations and cross-modal influences. Trends Cogn. Sci. 2020;24(6):481-95.

4.  Senkowski D, Engel AK. Multi-timescale neural dynamics for multisensory integration. Nat. Rev. Neurosci. 2024.

5.  Hipp JF, Engel AK, Siegel M. Oscillatory synchronization in large-scale cortical networks predicts perception. Neuron 2011;69(2):387-96.

6.  Keil J, Muller N, Ihssen N, Weisz N. On the variability of the McGurk effect: audiovisual integration depends on prestimulus brain states. Cereb Cortex 2012;22(1):221-31.

7.  Galindo-Leon EE, Stitt I, Pieper F, Stieglitz T, Engler G, Engel AK. Context-specific modulation of intrinsic coupling modes shapes multisensory processing. Sci. Adv. 2019;5(4): eaar7633.

8.  Wang P, Göschl F, Friese U, König P, Engel AK. Long-range functional coupling predicts performance: oscillatory EEG networks in multisensory processing. Neuroimage 2019;196:114-25.

9.  Cooke J, Poch C, Gillmeister H, Costantini M, Romei V. Oscillatory properties of functional connections between sensory areas mediate cross-modal illusory perception. J. Neurosci. 2019;39(29):5711-8.

10. Doesburg SM, Emberson LL, Rahi A, Cameron D, Ward LM. Asynchrony from synchrony: long-range gamma-band neural synchrony accompanies perception of audiovisual speech asynchrony. Exp. Brain Res. 2008;185(1):11-20.

11.  Lange J, Christian N, Schnitzler A. Audio-visual congruency alters power and coherence of oscillatory activity within and between cortical areas. Neuroimage 2013;79:111-20.

12.  Senkowski D, Hofle M, Engel AK. Crossmodal shaping of pain: a multisensory approach to nociception. Trends Cogn. Sci. 2014;18(6):319-27.

13.  Engel AK, Fries P. Beta-band oscillations – signalling the status quo? Curr Opin Neurobiol. 2010;20(2):156-65.

14.  Engel AK, Fries P, Singer W. Dynamic predictions: oscillations and synchrony in top-down processing. Nat. Rev. Neurosci. 2001;2(10):704-16.

15.  Arnal LH, Giraud AL. Cortical oscillations and sensory predictions. Trends Cogn. Sci. 2012;16(7):390-8.

16.  Bastos AM, Usrey WM, Adams RA, Mangun GR, Fries P, Friston KJ. Canonical microcircuits for predictive coding. Neuron 2012;76(4):695-711.

17.  Hyafil A, Giraud AL, Fontolan L, Gutkin B. Neural cross-frequency coupling: connecting architectures, mechanisms, and functions. Trends Neurosci. 2015;38(11):725-40.

18.  Jessen S, Kotz SA. On the role of crossmodal prediction in audiovisual emotion perception. Front. Hum. Neurosci. 2013;7:369.

19.  Arnal LH, Doelling KB, Poeppel D. Delta-beta coupled oscillations underlie temporal prediction accuracy. Cereb. Cortex 2015;25(9):3077-85.

20.  van Wassenhove V, Grzeczkowski L. Visual-induced expectations modulate auditory cortical responses. Front Neurosci. 2015;9:11.

21.  Nolte G, Bai O, Wheaton L, Mari Z, Vorbach S, Hallett M. Identifying true brain interaction from EEG data using the imaginary part of coherency. Clin. Neurophysiol. 2004;115(10):2292-307.

22.  Rohenkohl G, Bosman CA, Fries P. Gamma synchronization between V1 and V4 improves behavioral performance. Neuron 2018;100(4):1-11.

23.  Womelsdorf T, Schoffelen JM, Oostenveld R, Singer W, Desimone R, Engel AK, et al.

Modulation of neuronal interactions through neuronal synchronization. Science 2007;316(5831):1609-12.

24. Friston K. The free-energy principle: a unified brain theory? Nat. Rev. Neurosci. 2010;11(2):127-38.

25. Widmann A, Gruber T, Kujala T, Tervaniemi M, Schroger E. Binding symbols and sounds: evidence from event-related oscillatory gamma-band activity. Cereb. Cortex 2007;17(11):2696-702.

26. van Ede F, Jensen O, Maris E. Tactile expectation modulates pre-stimulus beta-band oscillations in human sensorimotor cortex. Neuroimage 2010;51(2):867-76.

27. Arnal LH, Wyart V, Giraud AL. Transitions in neural oscillations reflect prediction errors generated in audiovisual speech. Nat. Neurosci. 2011;14(6):797-801.

28. Biau E, Torralba M, Fuentemilla L, de Diego Balaguer R, Soto-Faraco S. Speaker's hand gestures modulate speech perception through phase resetting of ongoing neural oscillations. Cortex 2015;68:76-85.

29. Daume J, Wang P, Maye A, Zhang D, Engel AK. Non-rhythmic temporal prediction involves phase resets of low-frequency delta oscillations. Neuroimage 2021;224:117376.

30. Jensen O, Mazaheri A. Shaping functional architecture by oscillatory alpha activity: gating by inhibition. Front. Hum. Neurosci. 2010;4(186):186.

31. Bubic A, von Cramon DY, Schubotz RI. Prediction, cognition and the brain. Front. Hum. Neurosci. 2010;4:25.

32. Schubotz RI, von Cramon DY. Predicting perceptual events activates corresponding motor schemes in lateral premotor cortex: an fMRI study. Neuroimage 2002;15(4):787-96.

33. Bastos AM, Vezoli J, Bosman CA, Schoffelen JM, Oostenveld R, Dowdall JR, et al. Visual areas exert feedforward and feedback influences through distinct frequency channels. Neuron 2015;85(2):390-401.

34. Fries P. Rhythms for cognition: communication through coherence. Neuron 2015;88(1):220-

35.

35.    Engel AK, Gerloff C, Hilgetag CC, Nolte G. Intrinsic coupling modes: multiscale interactions in ongoing brain activity. Neuron 2013;80(4):867-86.

36.    Fries P. A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. Trends Cogn. Sci. 2005;9(10):474-80.

37.    Ni J, Wunderle T, Lewis CM, Desimone R, Diester I, Fries P. Gamma-rhythmic gain modulation. Neuron 2016;92(1):240-51.

38.    Daume J, Gruber T, Engel AK, Friese U. Phase-amplitude coupling and long-range phase synchronization reveal frontotemporal interactions during visual working memory. J. Neurosci. 2017;37(2):313-22.

39.    Wolff A, Berberian N, Golesorkhi M, Gomez-Pilar J, Zilio F, Northoff G. Intrinsic neural timescales: temporal integration and segregation. Trends Cogn. Sci. 2022;26(2):159-73.

40.    Brainard DH. The psychophysics toolbox. Spatial Vision 1997;10(4):433-6.

41.    Stolk A, Todorovic A, Schoffelen JM, Oostenveld R. Online and offline tools for head movement compensation in MEG. Neuroimage.2013;68(0):39-48.

42.    Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput. Intel. Neurosc. 2011: 156869.

43.    Mitra PP, Pesaran B. Analysis of dynamic brain imaging data. Biophys. J. 1999;76(2):691-708.

44.    Hyvarinen A. Fast and robust fixed-point algorithms for independent component analysis. IEEE Trans. Neural Netw. 1999;10(3):626-34.

45.    Hipp JF, Siegel M. Dissociating neuronal gamma-band activity from cranial and ocular muscle activity in EEG. Front. Hum. Neurosci. 2013;7:338.

46.    Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. Neuroimage 1999;9(2):179-94.

47.    Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MNE

software for processing MEG and EEG data. Neuroimage 2014;86(0):446-60.

48. Mazziotta JC, Toga AW, Evans A, Fox P, Lancaster J. A probabilistic atlas of the human brain: theory and rationale for its development. The International Consortium for Brain Mapping (ICBM). Neuroimage 1995;2(2):89-101.

49. Nolte G. The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalography forward calculation in realistic volume conductors. Phys. Med. Biol. 2003;48(22):3637-52.

50. Hipp JF, Siegel M. Accounting for linear transformations of EEG and MEG data in source analysis. PLoS One 2015;10(4):e0121048.

51. Van Veen BD, van Drongelen W, Yuchtman M, Suzuki A. Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. IEEE Trans. Biomed. Eng. 1997;44(9):867-80.

52. Leopold DA, Murayama Y, Logothetis NK. Very slow activity fluctuations in monkey visual cortex: implications for functional brain imaging. Cereb. Cortex. 2003;13(4):422-33.

53. Kujala J, Gross J, Salmelin R. Localization of correlated network activity at the cortical level with MEG. Neuroimage 2008;39(4):1706-20.

54. Nichols TE, Holmes AP. Nonparametric permutation tests for functional neuroimaging: a primer with examples. Hum. Brain Mapp. 2002;15(1):1-25.

55. Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, et al. Automatically parcellating the human cerebral cortex. Cereb. Cortex 2004;14(1):11-22.

56. Tort AB, Komorowski R, Eichenbaum H, Kopell N. Measuring phase-amplitude coupling between neuronal oscillations of different frequencies. J. Neurophysiol. 2010;104(2):1195-210.

## Supplementary Figures and Captions

**Fig. S1. Definition of the source space.** (A) White matter (left hemisphere) and pial (right hemisphere) surfaces of the template brain. Source locations were evenly distributed on the surface of the white matter, as indicated by the blue dots in the left hemisphere. (B) The inflated view of the template brain. Source locations are indicated by the yellow dots on the inflated surface.

**Fig. S2. Cluster analysis of power during response segments**. (A - C). In each panel, both spatial (left) and time-frequency (right) distribution are illustrated. Color indicates accumulated *t* values. Side plots in the box are integrated *t* values against time (top, x-axis: time in ms, y-axis: *t* value) or frequency (right, x-axis: *t* value, y-axis: frequency in Hz).

**Fig. S3. Cluster analysis of coupling during response segments**. In each panel, both spatial (left) and time-frequency (right) distribution are illustrated. Color indicates accumulated *t* values. Side plots in the box are integrated *t* values against time (top, x-axis: time in ms, y-axis: *t* value) or frequency (right, x-axis: *t* value, y-axis: frequency in Hz).

# Funding Disclosure

A.K.E. acknowledges support for this work from the DFG (SFB936-178316478-A3; TRR169-261402652-B1/B4/Z2) and from the European Union (project cICMs, ERC-2022-AdG-101097402). Views and opinions expressed in this article are those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

# Competing Interests

None.