1 # Consumption of a Western-style diet modulates the response of the
2 # murine gut microbiome to ciprofloxacin
3
4

5 Damien J. Cabral[a,b,+], Jenna I. Wurster[a,c,+], Benjamin J. Korry[a,d], Swathi Penumutchu[a,e], Peter
6 Belenky[a, #]

7
8 Running title: Impact of antibiotics and diet on the gut microbiome
9
10

[a] Department of Molecular Microbiology and Immunology, Brown University, Providence, RI
02906, USA
[b] Email: damien_cabral@brown.edu
[c] Email: jenna_wurster@brown.edu
[d] Email: benjamin_korry@brown.edu
[e] Email: swathi_penumutchu@brown.edu
17
18 [+] Damien J. Cabral and Jenna I. Wurster contributed equally to this work
19
20 [#] Address correspondence to Peter Belenky, peter_belenky@brown.edu

Abstract:

Dietary composition and antibiotic use are known to have major impacts on the structure and function of the gut microbiome, often resulting in dysbiosis. Despite this, little research has been done to explore the role of host diet as a determinant of antibiotic-induced microbiome disruption.

Here, we utilize a multi-omic approach to characterize the impact of Western-style diet consumption on ciprofloxacin-induced changes to gut microbiome community structure and transcriptional activity. We found that mice consuming a Western-style diet experienced a greater expansion of *Firmicutes* following ciprofloxacin treatment than those eating a control diet. At the transcriptional level, we found that ciprofloxacin induced a reduction in the abundance of TCA cycle transcripts on both diets, suggesting that carbon metabolism plays a key role in the response of the gut microbiome to this antibiotic. Despite this shared response, we observed extensive differences in the response of the microbiota to ciprofloxacin on each diet. In particular, at the whole-community level we detected an increase in starch degradation, glycolysis, and pyruvate fermentation following antibiotic treatment in mice on the Western diet, which we did not observe in mice on the control diet. Similarly, we observed diet-specific changes in the transcriptional activity of two important commensal bacteria, *Akkermansia muciniphila* and *Bacteroides thetaiotaomicron*, involving diverse cellular processes such as nutrient acquisition, stress responses, and capsular polysaccharide (CPS) biosynthesis. These findings demonstrate that host diet plays a key role in determining the extent of disruption of microbiome composition and function induced by antibiotic treatment.

Importance:

While both diet and antibiotics are individually known to have profound impacts on gut microbiome composition, little work has been done to examine the effect of these two factors combined. A number of negative health outcomes, including diabetes and obesity, are associated with diets high in simple sugars in fats but low in host-indigestible fiber, and some of these outcomes may be mediated by the gut microbiome. Likewise, treatment with broad-spectrum antibiotics and the resulting dysbiosis is associated with many of the same detrimental side effects. Previous work has shown that nutrient availability, as influenced by host diet, plays an important role in determining the extent of antibiotic-induced disruption to the gut microbiome. Due to the growing incidence of disorders related to antibiotic-induced dysbiosis, it is essential to determine how the prevalence of high fat and sugar "Western"-style diets impacts the response of the microbiome to antibiotics.

Keywords:

Diet, antibiotics, metagenomics, metatranscriptomics, dysbiosis

Background:

The gut microbiome includes the trillions of largely commensal bacteria, archaea, fungi, and viruses and their collective genetic material that inhabit the gastrointestinal tract (1-3). These communities play an important role in numerous biological processes, such as digestion, neurological development, colonization resistance, and immune function (4-18). However, the gut microbiome is exquisitely sensitive to perturbation and the disruption of microbial homeostasis, known as dysbiosis, can result in numerous harmful impacts to the host. In particular, broad-spectrum antibiotic use is known to have numerous detrimental impacts on the gut microbiota. Within hours of treatment, antibiotics induce dramatic reductions in both bacterial loads and diversity within the microbiome (19, 20).

While compositional changes are typically transient and recover following the cessation of therapy, oftentimes the structure and diversity of the microbiota never return to their pre-treatment levels. These changes often result in dysbiosis and have numerous acute and chronic impacts on host health. In particular, dysbiosis may increase the risk of infection with opportunistic fungal and bacterial pathogens by reducing colonization resistance (1, 5-7, 21-25). Most notably, broad-spectrum antibiotic treatment is known to be a major risk factor in *Clostridioides difficile* infection, which is responsible for approximately 29,000 deaths worldwide and 15,000 in the United States alone (21, 23, 26, 27). In addition to short-term complications such as pathogen blooms, persistent dysbiosis is correlated with a number of chronic conditions associated with considerable morbidity and mortality, such as asthma, obesity, and inflammatory bowel disease (5, 8-10, 13, 15, 17, 18, 28).

In addition to eliciting changes in community structure, antibiotic exposure also has a dramatic impact on the functional activity of the gut microbiome by altering the expression of

key metabolic genes and the availability of carbohydrates within the gut (20). Most notably,

amoxicillin treatment has been shown to increase the expression of glycoside hydrolases

responsible for hydrolysis of polysaccharides, while simultaneously decreasing the abundance of

transcripts encoding sugar phosphotransferase systems (PTS) responsible for uptake of simple

sugars (20). Reflecting these changes, amoxicillin also decreases the concentration of glucose

within the ceca of mice (20). Furthermore, dietary intervention experiments have demonstrated

that the response of the microbiota to antibiotics can be impacted by nutrient modulation. For

example, glucose supplementation reduces the absolute abundance of bacteria, particularly

Bacteroides thetaiotaomicron, following amoxicillin treatment in mice (20). Therefore, it is

likely that host diet composition has a major impact on the response of these communities to

perturbation.

It is also known that dietary composition has a profound impact on microbiome diversity

and overall gut health (29-35). Diets high in fat and simple sugars, typically referred to as

"Western" diets, have been associated with a number of negative health states including obesity,

diabetes, allergies, and inflammatory bowel disease (36-46). Such diets have very low levels of

microbiota-accessible carbohydrates (MACs), which are typically found in complex plant

polysaccharides and are indigestible and unabsorbable by the host (40, 44, 47-49). MACs are

typically fermented by the colonic microbiota to produce short-chain fatty acids (SCFAs), which

in turn play important roles in regulating energy homeostasis and inflammation within the host

(40, 45, 50-55). In addition to being associated with increased levels of SCFAs, high-MAC diets

have been shown to increase microbial diversity, a classic benchmark for gut microbiota health.

Conversely, low-MAC Western diets are known to reduce both microbiome diversity and SCFA

production (44, 46, 49, 56). Due to the absence of MACs, such diets also enrich for muciniphilic

microbes that may degrade mucosal lining of the gut, such as *Akkermansia muciniphila* (40, 42,

48). Degradation of the mucosal layer over time is hypothesized to result in compromised gut

barrier function that may ultimately lead to increased inflammation and colitis. Lastly, MAC-

deficient diets have been shown to exacerbate infection with *C. difficile,* suggesting that they

may negatively impact colonization resistance (26, 27).

Individually, antibiotic usage and the consumption of Western-style diets are known to

negatively impact the microbiota, resulting in similar long-term impacts on host health

(including obesity, allergies, and inflammatory bowel disease). Despite this, little work has been

done to explore how diet impacts the response of the microbiota to antibiotics. Previous work has

suggested that dietary composition may play an important role in determining the extent of

antibiotic-induced microbiome disruption (20). In this study, we use a combined metagenomic

and metatranscriptomic approach to characterize the impact of a Western-style diet on the

taxonomic and functional disruption of the microbiome during ciprofloxacin treatment. Using

shotgun metagenomics, we found that ciprofloxacin elicited differential impacts on community

composition in mice at both the phylum- and species-level that were diet-dependent. Using

metatranscriptomics, we observed that consumption of a Western diet itself induced profound

transcriptional changes within the gut microbiomes of mice; furthermore, consumption of this

diet modulated the transcriptional response of these communities to antibiotic treatment. In

particular, dietary composition had a major impact on the abundance of transcripts encoding key

metabolic genes. Lastly, we were able to detect unique species-specific transcriptional changes in

response to both diet and ciprofloxacin treatment in two important commensal bacteria, *A.*

muciniphila and *B. thetaiotaomicron*.

Results:

To determine the impact of dietary composition and antibiotic exposure on the structure and function of the murine gut microbiome, female C57BL/6J mice were randomly assigned to either a high-fat, high-sugar "Western"-style diet, or a low-fat control diet for seven days in multiple cages. At this point, mice from each diet were again randomly split between ciprofloxacin and vehicle control groups and treated for 24 hours in multiple cages (n = 8-to-12 per group). This time point was chosen because previously published work demonstrated that 24 hours of ciprofloxacin treatment was sufficient to induce changes in community structure and transcriptional activity (20). Additionally, this time frame allowed us to profile the acute response of the microbiota to ciprofloxacin exposure, rather than characterizing a post-antibiotic state of equilibrium. Following treatment, the mice were sacrificed to harvest their cecal contents for taxonomic profiling and transcriptional analysis (Figure 1A). Overall, we found that diet and ciprofloxacin treatment had a significant impact on gut microbiome structure (Figure 1B+C, Figure S1).

We first assessed the effects that diet and ciprofloxacin have on the diversity of the gut microbiome using 16S ribosomal RNA (rRNA) sequencing. Mice consuming the Western diet had significantly less diverse gut microbiomes than those fed the control diet (Figure S1A). Interestingly, we also observed that the Western diet exacerbated the reduction in alpha diversity during ciprofloxacin treatment (Figure S1A). Next, we performed principle coordinate analysis using Bray-Curtis dissimilarity paired with PERMANOVA analysis to profile the degree of dissimilarity between our samples and the significance of this distance. Unsurprisingly, our samples formed four distinct clusters, driven by both diet and ciprofloxacin treatment (Figure S1B).

Due to the limited phylogenetic resolution provided by 16S rRNA sequencing, in

addition to its inability to provide functional information about sequenced communities, we

opted to perform shotgun metagenomics and metatranscriptomic analysis on a subset of our

samples, representing mice from multiple cages (n = 4 per treatment group) (20, 57-60). Using

metagenomics, we observed that consumption of the Western diet reduced community diversity

(Figure 1B). Taxonomically, mice fed a Western diet displayed elevated levels of the phyla

Verrucomicrobia and *Bacteroidetes*, and a reduced abundance of *Firmicutes* (Figure 1C). At the

species level, these shifts appear to be largely driven by an expansion of members of the

Bacteroides genus (Figure 1D, Figure S1). Additionally, the Western diet-fed mice displayed an

elevated abundance of several species from the *Proteobacteria* phylum, which has been

previously shown to be suggestive of dysbiosis (61). Two important bacterial species found in

the gut microbiomes of both mice and humans, *B. thetaiotaomicron* and *A. muciniphila*, were

observed at significantly elevated levels in the mice fed a Western diet (Figure 1D, Figure S1).

Notably, both species are known to utilize host-produced mucins; thus, this observation is

consistent with earlier studies that have suggested that the consumption of a low-MAC Western

diet enriches for muciniphilic bacteria (40, 42, 48).

Overall, host diet appears to have a major impact on the structure of the microbiome

during ciprofloxacin treatment. Although ciprofloxacin did not induce a significant reduction in

alpha diversity in the timeframe tested, we found that diet drove differential community

composition following antibiotic exposure. At the phylum level we observed a significant

expansion in the relative abundance of *Firmicutes* following ciprofloxacin treatment on the

Western diet (adjusted p-value = 0.0388) but not on the control diet (adjusted p-value = 0.8718)

(Figure 1C). To determine which species displayed a differential response to ciprofloxacin on the

Western and control diets, we utilized DESeq2 to analyze the interaction between diet and

antibiotic treatment (62). In our analysis, an interaction term describes the direction and

magnitude of the differential impact that host diet has on antibiotic perturbation within the

microbiome. For example, ciprofloxacin reduces the abundance of *Clostridium innocuum* on the

control diet, while it expands following treatment on the Western diet (Figure 1E, Additional File

1); thus, the interaction term in this case is positive, as it indicates an expansion on the Western

diet relative to control following ciprofloxacin exposure.

While most species responded similarly to ciprofloxacin therapy on both diets, there were

several notable exceptions. For example, the expansion of several *Clostridium* species (such as

C. innocuum, Clostridium beijerinckii, and *Clostridium scindens*) following ciprofloxacin tended

to be higher on the Western diet than the control (denoted by a positive interaction value in

Figure 1E, Additional File 1). Conversely, the reduction of several *Bacteroides* species following

antibiotic treatment tended to be exacerbated on the Western diet (negative interaction values,

Figure 1E, Additional File 1). In particular, the reduction of *B. thetaiotaomicron* in response to

ciprofloxacin was enhanced on the Western diet; in contrast, dietary composition had no

significant impact on the response of *A. muciniphila* to ciprofloxacin (Additional File 1).

Ciprofloxacin Elicits Unique Shifts in Gene Expression on Western and Control Diets

We detected clear differences in the susceptibility of the microbiome to treatment on the

two diets. It is possible that the differential metabolic environment between the two diets

changes the activity of the gut microbiome and contributes to altered susceptibility. Though

many studies have examined the impacts of either diet or antibiotic treatment on the gut

microbiome, few have examined their combined effect on the composition or gene expression of

these communities. To address this, we first analyzed our metatranscriptomic dataset using the

HUMAnN2 pipeline, which normalizes the abundance of RNA transcripts against their

corresponding gene abundance in the metagenomic data (63). Thus, this tool normalizes for

differences in community composition between experimental groups and facilitates the

comparison of metabolic pathway expression at the whole-community level. A comparison of

the transcriptional profile of all experimental groups demonstrates that the microbiota of mice

consuming the Western diet display elevated expression of tricarboxylic acid (TCA) cycle and

fatty acid degradation pathways in both vehicle and ciprofloxacin treatments, likely reflective of

the increased fat and sugar content of this diet (Figure 2A, Additional File 2). Additionally, we

found an elevated expression of glycogen degradation genes in the Western diet mice receiving

ciprofloxacin, which was not observed in any other group (Figure 2A, Additional File 2).

Conversely, the microbiota of mice consuming the control diet appeared to have elevated

expression of amino acid biosynthesis pathways (namely isoleucine, aspartate, asparagine,

lysine, and histidine) regardless of antibiotic treatment (Figure 2A, Additional File 2).

Interestingly, we also observed elevated levels of several different pathways of nucleotide

biosynthesis in the control diet samples receiving vehicle while the Western diet mice displayed

elevated levels of adenosine and guanosine nucleotide degradation (Figure 2A, Additional File

2).

A pairwise comparison between the vehicle-treated samples on the Western and control

diets reveals that the microbiota exhibits extensive transcriptional changes in response to dietary

modulation (Additional File 2). Notably, the microbiota on the control diet-fed mice displayed

elevated expression of nucleotide biosynthesis, glycolysis, gluconeogenesis, starch degradation,

and pyruvate fermentation (Figure 2A, Additional File 2). We also observed increased

expression of the *Bifidobacterium* shunt, which is known to play a role in SCFA production and

may provide mechanistic insight into the reduced SCFA levels observed on the Western diet in

other studies (40, 51) (Additional File 2).

When we compared the response of the microbiome to ciprofloxacin on each diet, we

found key differences in the overall transcriptional profiles (Additional File 2). In mice

consuming the Western diet, ciprofloxacin treatment was associated with increased abundance of

transcripts from glycogen and starch degradation, glycolysis, and pyruvate fermentation

(Additional File 2). Notably, the expression of glycogen degradation was elevated in vehicle-

treated samples on the control diet, suggesting that the utilization of this pathway during

ciprofloxacin treatment is diet-dependent (Additional File 2). On the control diet, we observed an

increased abundance of inosine-5-phosphate biosynthesis with ciprofloxacin treatment (Figure

2A, Additional File 2). Interestingly, we observed that TCA cycle expression was reduced in

ciprofloxacin-treated mice compared to the vehicle treatment in both control and Western diet

conditions – the lone commonality between diets (Additional File 2). Previous work has

demonstrated that elevated TCA cycle activity increases sensitivity to bactericidal antibiotics,

including fluoroquinolones, *in vitro* (64-68). Thus, this result suggests that TCA cycle activity

may play a key role in the response of the microbiota to ciprofloxacin treatment *in vivo*, though

more work is required to understand its impact.

Ciprofloxacin has a differential impact on the abundance of iron metabolism and mucin

degradation transcripts on the Western versus control diets

Due to the potential limitations of the use of a single pipeline, we analyzed our

metatranscriptomic dataset with SAMSA2 in parallel with HUMAnN2. While HUMAnN2

normalizes for DNA abundance, SAMSA2 does not have this feature and thus the output is

representative of overall transcript levels rather than relative expression. Despite these

differences, many of the changes observed using HUMAnN2 were detected using SAMSA2 at

the SEED subsystem level. When comparing the vehicle-treated samples on both diets, SAMSA2

detected an increased abundance of transcripts related to respiration in the Western diet,

mirroring the increase in TCA cycle expression found with HUMAnN2 (Figure 2B, Additional

File 3). The microbiota from the Western diet-fed mice also displayed an increased abundance of

transcripts involving fatty acids and iron acquisition, which likely reflect altered nutrient

availability (Figure 2B, Additional File 3). Furthermore, transcripts related to virulence and

disease were elevated in mice consuming the Western diet, further supporting the hypothesis that

consumption of a Western diet may promote dysbiosis and the expansion of enteric pathogens

(Figure 2B, Additional File 3). Interestingly, comparatively few subsystems were changed in

abundance following ciprofloxacin treatment on either diet (Additional File 3). Most notably, we

observed a decrease in transcripts related to dormancy and sporulation in response to

ciprofloxacin on both diets (Additional File 3). A similar finding was observed in a recent study

in which mice were treated with ciprofloxacin while consuming a non-purified diet, suggesting

that these transcripts may play a key role in the response of the microbiota to this antibiotic (20).

An additional strength of the SAMSA2 pipeline is that it enables differential abundance

testing of individual transcripts in addition to pathway- and subsystem-level analysis (69). We

observed that the microbiota of the mice consuming the Western diet displayed increased

abundance of heme b synthase transcripts, which may suggest increased iron acquisition (Figure

2C, Additional File 4). Interestingly, in the untreated mice, we detected large, Western diet-

associated increases in the abundance of two different transcripts encoding sialidases, which play

a key role in the utilization of host-produced mucins (70) (Figure 2C, Additional File 4) . While

other studies have shown that the consumption of a Western diet enriches for muciniphilic taxa,

this observation demonstrates that such a diet also increases transcriptional activity related to

mucin degradation within the microbiome (40, 42). Furthermore, ciprofloxacin increased the

abundance of sialidase transcripts in mice on the control diet, suggesting that this effect may be

exacerbated by antibiotic treatment (Figure 2D, Additional File 4).

In addition to the increased abundance of sialidase transcripts, ciprofloxacin induced a

number of notable transcriptional changes on each diet. Reflecting the overall reduction in

sporulation seen at the subsystem level, we found that the abundance of several sporulation-

related transcripts was reduced on the control diet following ciprofloxacin treatment (Figure 2D,

Additional File 4). We also examined the interaction of diet and antibiotic treatment on

transcript abundance within the microbiome. Notably, we found that the transcript abundance of

several sporulation genes following ciprofloxacin treatment was significantly higher on the

Western diet than the control (Additional File 4). Additionally, transcripts encoding

phosphotransferase system (PTS) transporters of various substrates (such as cellobiose, fructose,

and N-acetylgalactosamine) were also found to be higher on the Western diet following

ciprofloxacin treatment (Additional File 4). Conversely, consumption of the Western diet

significantly reduced the change in transcript abundance of both pectate lyase and a hemin

receptor following ciprofloxacin therapy. Together, these findings demonstrate that dietary

composition significantly impacts the transcriptional response of the microbiome to

ciprofloxacin.

Diet and ciprofloxacin alter gene expression within *B. thetaiotaomicron* and *A. muciniphila*

Next we sought to profile how diet and drug treatment impacted the transcriptional

response of individual species within the microbiota. We used a previously published pipeline to

interrogate the impact of diet and antibiotic treatment on two individual species: *B.*

thetaiotaomicron and *A. muciniphila* (20, 71). We focused on these bacteria because they are

known human gut commensals, were found in relatively high levels in all samples analyzed, and

because they were differentially abundant in a diet-dependent manner.

The Western diet significantly elevated the baseline relative abundance of *A. muciniphila*

(Figure 3A). Interestingly, on this diet *A. muciniphila* displayed increased expression of several

known stress response genes (Figure 3B, Additional File 5). Specifically, we observed elevated

levels of transcripts for catalase HPII (AMUC_RS11055), ATP-dependent chaperone ClpB

(AMUC_RS04500), a universal stress protein (AMUC_RS00415), superoxide dismutase

(AMUC_RS08510), a UvrB/UvrC protein (AMUC_RS00335), and a thioredoxin family protein

(AMUC_RS11695). Together, these changes may indicate that Western diet consumption

induces a general stress response in *A. muciniphila*. Additionally, we observed numerous

changes within respiratory and central carbon metabolism, suggesting broad metabolic changes

in response to the Western diet (Figure 3B, Additional File 5). We specifically detected increased

expression of genes encoding terminal oxidases of the respiratory chain (AMUC_RS09050 -

cytochrome ubiquinol oxidase subunit I and AMUC_RS09045 - cytochrome d ubiquinol oxidase

subunit II), the TCA cycle (AMUC_RS09040 - 2-oxoglutarate dehydrogenase E1 component),

glycolysis (AMUC_RS06320 - phosphopyruvate hydratase, AMUC_RS02385 - pyruvate

kinase), and pyruvate metabolism (AMUC_RS01195 - ubiquinone-dependent pyruvate

dehydrogenase). Of the genes that were significantly reduced on the Western diet, the most

notable were an adenylsuccinate synthase (AMUC_RS11340) and a lyase (AMUC_RS10360),

which both play important roles in purine metabolism (Figure 3B, Additional File 5) (72).

In comparison to diet, ciprofloxacin treatment had a relatively minor impact on *A.*

muciniphila gene expression (Additional File 5). This likely relates to the fact that on both diets,

ciprofloxacin did not impact the relative abundance of *A. muciniphila* (Figure 3A). In total,

ciprofloxacin significantly altered the expression of 14 and 26 genes on the control and Western

diets, respectively (Additional File 5). On the control diet, *A. muciniphila* increased the

expression of the molecular chaperone protein DnaK, which is known to play a role in stress

responses (73-75). Additionally, we observed elevated expression of a MoxR family ATPase

following ciprofloxacin treatment on this diet. Though these proteins are poorly characterized,

other members of this family have been shown to regulate stress responses in other bacteria (76).

On the Western diet, several genes related to tryptophan biosynthesis and metabolism were

elevated following ciprofloxacin treatment (AMUC_RS08210 - tryptophan synthase subunit

beta, AMUC_RS08190 - anthranilate synthase component I family protein, AMUC_RS08215 -

tryptophan synthase subunit alpha); however, their biological significance is unclear at this time

(Additional File 5). Lastly, an examination of the interaction between diet and ciprofloxacin

treatment indicated that only six genes (two of which encoded tRNAs) were significantly altered,

suggesting that diet does not have a major impact on the response of this bacterium to

ciprofloxacin within the microbiome (Additional File 5).

In contrast to *A. muciniphila*, diet had a relatively minor impact on *B. thetaiotaomicron*

gene expression while ciprofloxacin induced extensive changes. It is important to note that *B.*

thetaiotaomicron bloomed in response to the Western diet and was significantly perturbed on the

Western diet but not on the control (Figure 3C). In total, *B. thetaiotaomicron* altered the

expression of 74 genes in response to Western diet consumption (Additional File 6). Of note, this

diet increased the expression of an aminoglycoside efflux pump (BT_0305), the universal stress

protein UspA (BT_0901), and a hemin receptor (BT_0316). However, more than half of the

genes (52.7%) that changed in response to diet are of unknown function and are classified as

"hypothetical proteins;" thus, it is difficult to draw conclusions without improved annotation.

Overall, ciprofloxacin appeared to induce extensive transcriptional changes within *B.*

thetaiotaomicron regardless of diet. On the control diet, we observed an increased abundance of

transcripts encoding a number of proteins involved in capsular polysaccharide (CPS)

biosynthesis and export (Figure 3D, Additional File 6). Within *B. thetaiotaomicron,* CPS

production is encoded by a total of 182 genes distributed among eight loci (typically termed

cps1-8) (77, 78). It is hypothesized that an individual bacterium expresses one of these CPS

configurations at any given time and that these structures play key roles in processes such as

nutrient acquisition and immune evasion (78). Additionally, the two genes with the greatest

increase in expression during ciprofloxacin treatment encoded UDP-glucose 6-dehydrogenase,

which plays a key role in the biosynthesis of glycan precursors that are essential for capsule

production in other bacteria (79-81). Together, these findings may suggest a role for CPS state as

a determinant of ciprofloxacin susceptibility *in vivo*.

On the Western diet, ciprofloxacin elicited profound changes in transcriptional activity,

altering the expression of 442 different genes (Figure 3E, Additional File 6) and this robust

response may be related to the reduction in *B. thetaiotaomicron* in this condition (Figure 3E).

Interestingly, *B. thetaiotaomicron* reduced the expression of a number of genes involved in the

utilization of host-derived carbohydrates (sialic acid-specific 9-O-acetylesterase, endo-beta-N-

acetylglucosaminidase F1, beta-hexosaminidase) and stress responses (universal stress protein

UspA, thioredoxin), mirroring the changes we saw at the whole-community level (Figure 3E,

Additional File 6). Conversely, we observed increased expression of several genes that encode

molecular chaperones (such as GroEL and GroES) or are involved in DNA replication or damage

repair (such as a DNA helicase, DNA gyrase subunit B, DNA mismatch repair protein MutS,

DNA polymerase III subunit alpha, Holliday junction DNA helicase RuvB, DNA-binding

proteins, and DNA primase) (Figure 3E, Additional File 6). Ciprofloxacin, a fluoroquinolone

class antimicrobial, triggers DNA damage via inhibition of DNA gyrase and topoisomerase IV.

Thus, these changes in gene expression may be reflective of the primary mechanism of action of

this antibiotic and are consistent with previously published data (20). Furthermore, our ability to

detect such changes is an important indication that our analysis is detecting ciprofloxacin-

induced transcriptional shifts. Lastly, diet appears to have a significant impact on ciprofloxacin-

induced transcriptional changes within *B. thetaiotaomicron*, modulating the response of 148

genes (Additional File 6). Of note, Western diet consumption in the context of ciprofloxacin

treatment had a negative impact on several genes involved in the acquisition of nutrients, such as

vitamin B_{12} and hemin receptors, and transporters of glucose/galactose, hexuronate, arabinose,

and Na^+ (Additional File 6). Thus, it is likely that the availability of nutrients within the gut plays

a role in the response of these bacteria to antibiotics.

Discussion:

Previous work has demonstrated that host diet, particularly with respect to sugar and fiber content, plays a major role in the extent of antibiotic-induced microbiome disruption (20). In Western societies, many people consume a diet high in added sugars and fat, but low in host-indigestible fiber. It is thought that such a composition promotes the development of metabolic syndrome, heart disease, diabetes, and a number of other chronic conditions (36-46). Furthermore, broad-spectrum antibiotic use and resulting microbiome dysbiosis have been associated with a number of similar co-morbidities along with increased susceptibility to opportunistic infections (1, 5-7, 21-23, 25, 26). Despite this connection, little work has been done examining how host dietary composition impacts the response of the microbiota to antibiotic perturbation. It is known that nutrient availability and metabolic state are a major determinant of antibiotic susceptibility of bacteria *in vitro* (20, 64-67, 82-88). Thus, it is likely that modulating host diet, thus changing the availability of nutrients to the microbiota, would alter the sensitivity of bacteria in these communities to antibiotic therapy.

To address this knowledge gap, we utilized a combined metagenomic and metatranscriptomic approach to profile taxonomic and functional changes in response to both diet and antibiotic treatment. By utilizing these tools in parallel, we are able to link transcriptional changes to observed shifts in community structure on each diet. Using metagenomics, we observed that ciprofloxacin had a differential impact on community composition that was diet-dependent. Specifically, we observed a statistically significant expansion of the *Firmicutes* phylum following ciprofloxacin treatment on the Western, but not control, diet. Using metatranscriptomics, we observed that ciprofloxacin treatment resulted in a decreased abundance of transcripts from the TCA cycle in both diets, suggesting that this

response is diet-independent. Furthermore, this observation is consistent with previous *in vitro*

findings that demonstrate a key role for bacterial respiration as a determinant of susceptibility to

fluoroquinolones (64-66, 68, 82, 85). Conversely, ciprofloxacin had diverging impacts on the

abundance of various iron and mucin utilization transcripts on the Western and control diets.

Most notably, we found that Western diet consumption (alone and in the presence of

ciprofloxacin) influenced the abundance of transcripts encoding genes known to play a role in

virulence, supporting previous literature demonstrating that nutrient availability impacts

virulence of enteric pathogens (20, 89-92). Lastly, we detected species-specific transcriptional

changes in two important commensal bacteria, *B. thetaiotaomicron* and *A. muciniphila*. In

addition to detecting changes in transcript levels that were reflective of stress responses, we also

observed that transcripts involved in diverse cellular processes such as nutrient acquisition,

carbon metabolism, and capsular polysaccharide (CPS) biosynthesis were differentially

expressed as well.

Despite the advantages of a multi-omic approach, there are a number of drawbacks to

these techniques that complicate the interpretation of our results. Most crucially, nearly all

analytical pipelines used to analyze microbiome data are reliant on existing databases that are

largely incomplete. It is hypothesized that approximately half of all genes within the human gut

microbiome have no functional annotation (93). Thus, the ability to accurately profile the

transcriptional activity of these communities is inherently limited by the quality and

completeness of the databases utilized. Additionally, elucidating the biological significance of

taxonomic of functional changes is often difficult in many microbiome analyses. Due to the

complex nature of these communities, it is often difficult to ascertain if the observed

transcriptional changes are the result of the direct action of the antibiotic, or the indirect effect of

changes in host physiology, nutrient availability, or the disruption of ecological networks within

the microbiome. For example, our transcriptional analysis of *B. thetaiotaomicron* showed that

this bacterium differentially expressed receptors for both hemin and vitamin B_{12}, which may

suggest that these nutrients play a role in ciprofloxacin toxicity. Alternatively, it is possible that

these transcriptional changes are reflective of increased availability of these nutrients due to

decreased competition from other members of the microbiota (though these hypotheses are not

mutually exclusive). Additionally, it is possible that dietary composition could play a significant

role in antibiotic absorption or sequestration in the gut, which in turn would impact the extent of

the damage caused to the microbiota.

This study builds on recent work that demonstrates that the availability of metabolites to

the bacteria in the host plays an important role in determining the extent of antibiotic-induced

microbiome disruption (20). Taken together, these results demonstrate the need to consider

dietary composition in the design and interpretation of experiments focused on understanding the

impact of antibiotics on the microbiota. Previous studies have demonstrated that dietary changes

induce rapid shifts in gut microbiome composition (32, 34, 43, 56, 94-97). Therefore, in the

long-term, dietary modulation could represent an attractive strategy to reduce the collateral

damage to commensal bacteria and the resulting complications from dysbiosis caused by clinical

therapy. Despite these promising applications, considerable work is required before these

findings have direct clinical relevance. In particular, the considerable differences in physiology,

microbiome composition, and diet between humans and rodents complicate the direct clinical

relevance of these findings. Furthermore, it is unclear whether short-term dietary modulation has

any long-term consequences on either the host or the microbiome. Thus, additional research is

warranted to fully elucidate how host diet impacts antibiotic-induced microbiome disruption in

humans.

Conclusions:

Using a combined metagenomic and metatranscriptomic approach, we demonstrate that

murine diet composition has a major impact on the response of the murine gut microbiome to

ciprofloxacin therapy. First, we found that the gut microbiome undergoes differential shifts in

community structure in response to antibiotic treatment in a diet-dependent manner. At the

transcriptional level, we found that ciprofloxacin reduced the abundance of TCA cycle

transcripts regardless of diet, suggesting that central carbon metabolism plays a role in the

activity of this antibiotic *in vivo*. Despite this commonality, we observed extensive differences in

the transcriptional response of the microbiome to dietary intervention and/or ciprofloxacin

treatment. Notably, mice consuming a Western diet had a high abundance of transcripts encoding

proteins known to degrade host-derived polysaccharides such as sialic residues of mucin,

suggesting that the consumption of this diet may have detrimental impacts on host physiology.

Lastly, we identified species-specific changes in transcript abundance in two key members of the

gut microbiome, *A. muciniphila* and *B. thetaiotaomicron*. In *A. muciniphila,* consumption of a

Western diet increased the expression of several genes known to play a role in stress responses.

In *B. thetaiotaomicron,* we found that a number of genes involved in CPS biosynthesis were

differentially expressed during ciprofloxacin treatment on the control but not Western diet,

which may suggest a divergent response of this bacterium to ciprofloxacin that is dependent on

nutrient composition. Taken together, these findings demonstrate that host diet is a determinant

of antibiotic-induced microbiome perturbations.

Methods:

Animal Procedures

All animal work was approved by Brown University's Institutional Animal Care and Use Committee (IACUC) under protocol number 1706000283. 4-week-old female C57BL/6 mice were purchased from Jackson Laboratories (Bar Harbor, ME, USA) and given a 2-week habituation period immediately following arrival at Brown University's Animal Care Facility. After habituation, mice were switched from standard chow (Laboratory Rodent Diet 5001, St. Louis, MO, USA) to either a Western diet (D12079B, Research Diets Inc., New Brunswick, NJ, USA) or a macronutrient-defined control diet (D122405B, Research Diets Inc., New Brunswick, NJ, USA) for 1 week. Following dietary intervention, mice were given acidified ciprofloxacin (12.5 mg/kg/day), or a pH-adjusted vehicle, via filter-sterilized drinking water *ad libitum* for 24 hours (n = 8-12 per treatment group). Water consumption was monitored to assure equal consumption across cages. Mice were then sacrificed and dissected in order to collect cecal contents. Cecal contents were immediately transferred to ZymoBIOMICS DNA/RNA Miniprep Kit (Zymo Research, Irvine, CA, USA) Collection Tubes containing DNA/RNA Shield. Tubes were processed via vortex at maximum speed for 5 minutes to homogenize cecal contents, then placed on ice until permanent storage at -80°C.

Nucleic Acid Extraction & Purification

Total nucleic acids (DNA and RNA) were extracted from samples using the ZymoBIOMICS DNA/RNA Miniprep Kit from Zymo Research (R2002, Irvine, CA, USA) using the parallel extraction protocol as per the manufacturer instructions. Total RNA and DNA were eluted in nuclease-free water and quantified using the dsDNA-HS and RNA-HS kits on a

Qubit™ 3.0 fluorometer (Thermo Fisher Scientific, Waltham, MA, USA) before use in library

preparations.

<u>16S rRNA Amplicon Preparation & Sequencing</u>

The 16S rRNA V4 hypervariable region was amplified from total DNA using the

barcoded 518F forward primer and the 816Rb reverse primers from the Earth Microbiome

Project (98). Amplicons were generated using 5X Phusion High-Fidelity DNA Polymerase under

the following cycling conditions: initial denaturation at 98°C for 30 seconds, followed by 25

cycles of 98°C for 10 seconds, 57°C for 30 seconds, and 72°C for 30 seconds, then a final

extension at 72°C for 5 minutes. After amplification, samples were pooled in equimolar amounts

and visualized via gel electrophoresis. Pooled amplicon library was submitted to the Rhode

Island Genomics and Sequencing Center at the University of Rhode Island (Kingston, RI, USA)

for sequencing on the Illumina MiSeq platform. Amplicons were pair-end sequenced (2x250 bp)

using the 500-cycle kit with standard protocols. We obtained an average of 106,135 ± 49,789

reads per sample. Raw reads were deposited in the NCBI Short Read Archive (SRA) under

accession number PRJNA594642.

<u>Analysis of 16S rRNA Sequencing Reads</u>

Raw 16S rRNA reads were subjected to quality filtering, trimming, de-noising, and merging

using the DADA2 package (version 1.8.0) in R (version 3.5.0). Ribosomal sequence variants

were assigned taxonomy using the RDP Classifier algorithm with RDP Training set 16 using the

assignTaxonomy function in DADA2 (99). Alpha diversity (Shannon) and beta diversity (Bray-

Curtis dissimilarity) were calculated using the phyloseq package (version 1.24.2) in R (version

3.5.0).

Metagenomic & Metatranscriptomic Library Preparation

Metagenomic libraries were prepared from DNA (100 ng) using the NEBNext® Ultra II

FS DNA Library Prep Kit (New England BioLabs, Ipswich, MA, USA) > 100ng input protocol

as per the manufacturer's instructions. This yielded a pool of 200 – 1000 bp fragments where the

average library was 250-500 bp. Metatranscriptomic libraries were prepared from total RNA

using the NEBNext® Ultra II Directional RNA Sequencing Prep Kit (New England BioLabs,

Ipswich, MA, USA) in conjunction with the NEBNext® rRNA Depletion Kit for

Human/Mouse/Rat (New England BioLabs, Ipswich, MA, USA) and the MICROBExpress kit

(Invitrogen, Carlsbad, CA, USA). First, up to 1 ug of total RNA was treated with rDNase I and

subsequently depleted of bacterial rRNAs using MICROBExpress as per the manufacturer's

instructions. This depleted RNA was then used to prepare libraries with the NEBNext® Ultra II

Directional RNA Sequencing Prep & rRNA depletion kits as per the manufacturer's instructions.

This yielded libraries that averaged between 200-450 bp. Once library preparation was complete,

both metagenomic and metatranscriptomic libraries were sequenced as paired-end 150 bp reads

on an Illumina HiSeq X Ten. We sequenced an average of 2,278,948,631 (± 2,309,494,556)

bases per metagenomic sample and 14,751,606,319 (± 3,089,205,166) bases per

metatranscriptomic sample. One metagenomic sample from the Western diet + vehicle group had

an abnormally low number of bases sequenced (165,000 bp) and was excluded from all

subsequent analyses. Following the removal of this sample, we obtained an average of

2,430,867,540 (± 2,306,317,898) bases per metagenomic sample. All reads were deposited in

the NCBI Short Read Archive under BioProject number PRJNA563913.

Processing of Raw Metagenomic and Metatranscriptomic Reads

Raw metagenomic reads were trimmed and decontaminated using kneaddata utility

(version 0.6.1) (100). In brief, reads were first trimmed to remove low quality bases and Illumina

TruSeq3 adapter sequences using trimmomatic (version 0.36) using SLIDINGWINDOW value

of 4:20 and ILLUMINACLIP value 2:20:10, respectively (101). Trimmed reads shorter than 75

bases were discarded. Reads passing quality control were subsequently decontaminated by

removing those that mapped to the genome of C57BL/6J mice using bowtie2 (version 2.2) (102).

Additionally, preliminary work by our group detected high levels of reads mapping to two

murine retroviruses found in our animal facility: murine mammary tumor virus (MMTV) and

murine osteosarcoma viruses (MOV) (20). Raw metatranscriptomic reads were trimmed and

decontaminated using the same parameters. However, in addition to removing reads that mapped

to the C57BL/6J, MMTV, and MOV genomes, we also decontaminated sequences that aligned to

the SILVA 128 LSU and SSU Parc ribosomal RNA databases (103).

Taxonomic Classification of Metagenomic Reads

Trimmed and decontaminated metagenomic reads were taxonomically classified against a

database containing all bacterial and archaeal genomes found in NCBI RefSeq using Kraken2

(version 2.0.7-beta) with a default k-mer length of 35 (104). Phylum- and species-level

abundances were subsequently calculated from Kraken2 reports using Bracken (version 2.0.0)

with default settings (105). The phyloseq package (version 1.28.0) in R (version 3.6.0) was used

to calculate alpha diversity using the Shannon Diversity Index (106). Metagenomic data was not

subsampled prior to analysis.

To perform differential abundance testing, species-level taxonomic output was first

filtered to remove taxa that were not observed in >1,000 reads (corresponding to approximately

0.1% of all reads) in at least 20% of all samples using phyloseq in R. Differential abundance

testing was subsequently performed on filtered counts using the DESeq2 package (version

1.24.0) using default parameters (62). All p-values were corrected for multiple hypothesis testing

using the Benjamini-Hochberg method (107).

Annotation of Metatranscriptomic Reads Using SAMSA2

Trimmed and decontaminated metatranscriptomic reads were annotated using a modified

version of the Simple Annotation of Metatranscriptomes by Sequence Analysis 2 (SAMSA2)

pipeline as described previously (20, 69, 108). First, the Paired-End Read Merger (PEAR) utility

was used to merge forward and reverse reads (109). Merged reads were then aligned to databases

containing entries from the RefSeq and SEED Subsystems databases using DIAMOND (version

0.9.12) (110, 111). The resulting alignment counts were subsequently analyzed using DESeq2

(version 1.24.0) using the Benjamini-Hochberg method to perform multiple hypothesis testing

correction (20, 69, 107). Features with an adjusted p-value of less than 0.1 were considered to be

statistically significant.

Metatranscriptomic Analysis using HUMAnN2

To determine the impact of dietary modulation and ciprofloxacin treatment on gene

expression within the gut microbiome, we used the HMP Unified Metabolic Analysis Network 2

(HUMAnN2, version 0.11.1) pipeline (63). First, metagenomic reads were taxonomically

annotated using MetaPhlan2 (version 2.6.0) and functionally annotated against the UniRef90

database to generate gene family and MetaCyc pathway level abundances. To ensure consistent

assignment between paired samples, the taxonomic profile generated from the metagenomic

reads was supplied to the HUMAnN2 algorithm during the analysis of the corresponding

metatranscriptomic reads. Metatranscriptomic reads were subsequently annotated as done for

metagenomic reads. The resulting gene family and pathway level abundance data from the

metatranscriptomic reads was normalized against the metagenomic data from the corresponding

sample and smoothed using the Witten-Bell method (112). Lastly, the resulting RPKM values

were unstratified to obtain whole-community level data, converted into relative abundances, and

analyzed using LEfSe (version 1) hosted on the Galaxy web server (113).

Transcriptional Analysis of *A. muciniphila* and *B. thetaiotaomicron*

A modified version of a previously published pipeline from Deng *et al.* was utilized to

perform transcriptional analysis of individual species within the murine microbiome during

dietary modulation and antibiotic treatment (20, 71). First, Kraken2 (version 2.0.7-beta) was

used to identify the fifty most prevalent bacterial species present within the metatranscriptomic

samples (104). Next, the BBSplit utility within the BBMap package (version 37.96) was used to

extract reads within our metatranscriptomic dataset that mapped to these fifty most abundant

species (114). Reads from *B. thetaiotaomicron* and *A. muciniphila* were subsequently aligned to

their corresponding reference genomes using the BWA-MEM algorithm (version 0.7.15) (115).

Lastly, the featureCounts command within the subread program (version 1.6.2) was used to

analyze the resulting alignment files to generate a count table for differential expression analysis

with DESeq2 (62). All p-values were corrected for multiple hypothesis testing with the

Benjamini-Hochberg method (107). Features with an adjusted p-value of less than 0.1 were

considered to be statistically significant.

List of Abbreviations:

PTS – phosphotransferase systems

MACs – microbiota-accessible carbohydrates

SCFAs – short chain fatty acids

TCA – tricarboxylic acid

HUMAnN2 – Human Microbiome Project (HMP) Unified Metabolic Analysis Network 2

CPS - capsular polysaccharide

GH98 – glycoside hydrolase family 98

SAMSA2 - Simple Annotation of Metatranscriptomes by Sequence Analysis 2

SRA – Sequence Read Archive

Declarations:

Availability of Data and Materials

The datasets generated and analyzed during this study are available from the NCBI Short Read Archive (SRA) under BioProject accession numbers PRJNA563913 (metagenomics and metatranscriptomics) and PRJNA594642 (16S rRNA amplicon sequences). Any additional information is available from the corresponding author upon request.

Competing interests

The authors declare that they have no competing interests

Authors' contributions

DJC planned the study, performed mouse experiments, extracted nucleic acids from cecal samples, conducted analysis of 16S rRNA amplicon, metagenomic, and metatranscriptomic data, and co-wrote the manuscript. JIW assisted with mouse experiments, prepared DNA and RNA into sequencing libraries for metagenomics and metatranscriptomics, assisted in the interpretation of results, and co-wrote the manuscript. BJK assisted with the analysis of metatranscriptomic data. SP assisted in the interpretation of results. PB conceptualized and planned the study, contributed to the writing of the manuscript, and secured funding. All authors have read and approved of the final manuscript.

Acknowledgements

D.J.C, B.J.K, and J.I.W., and S.P. were supported by the Graduate Research Fellowship Program from the National Science Foundation under award number 1644760. P.B. was supported by the

635   National Center for Complementary & Integrative Health of the NIH Award Number

636   R21AT010366, and institutional development awards P20GM121344 and P20GM109035

637   received from the National Institute of General Medical Sciences within the NIH. The funding

638   agencies had no role in the design of the study or the collection, analysis, and interpretation of

639   data.

References:

1. **Rowan-Nash AD**, **Korry BJ**, **Mylonakis E**, **Belenky P**. 2019. Cross-Domain and Viral
Interactions in the Microbiome. Microbiol Mol Biol Rev **83**:51.

2. **Gilbert JA**, **Blaser MJ**, **Caporaso JG**, **Jansson JK**, **Lynch SV**, **Knight R**. 2018.
Current understanding of the human microbiome. Nat Med **24**:392–400.

3. **Ursell LK**, **Metcalf JL**, **Parfrey LW**, **Knight R**. 2012. Defining the human
microbiome. Nutr Rev **70**:S38–44.

4. **Stiemsma LT**, **Michels KB**. 2018. The Role of the Microbiome in the Developmental
Origins of Health and Disease. Pediatrics **141**:e20172437.

5. **Blaser M**. 2011. Antibiotic overuse: Stop the killing of beneficial bacteria. Nature
476:393–394.

6. **Mukherjee PK**, **Chandra J**, **Retuerto M**, **Sikaroodi M**, **Brown RE**, **Jurevic R**, **Salata
RA**, **Lederman MM**, **Gillevet PM**, **Ghannoum MA**. 2014. Oral mycobiome analysis
of HIV-infected patients: identification of Pichia as an antagonist of opportunistic fungi.
PLoS Pathog **10**:e1003996.

7. **Peleg AY**, **Hogan DA**, **Mylonakis E**. 2010. Medically important bacterial-fungal
interactions. Nat Rev Microbiol **8**:340–349.

8. **De Luca F**, **Shoenfeld Y**. 2018. The microbiome in autoimmune diseases. Clin Exp
Immunol **195**:74–85.

9. **Dickerson F**, **Severance E**, **Yolken R**. 2017. The microbiome, immunity, and
schizophrenia and bipolar disorder. Brain Behav Immun **62**:46–52.

10. **Foster JA**, **McVey Neufeld K-A**. 2013. Gut-brain axis: how the microbiome influences
anxiety and depression. Trends Neurosci **36**:305–312.

11. **Foster JA**, **Rinaman L**, **Cryan JF**. 2017. Stress & the gut-brain axis: Regulation by the
microbiome. Neurobiology of Stress **7**:124–136.

12. **Hartstra AV**, **Bouter KEC**, **Bäckhed F**, **Nieuwdorp M**. 2015. Insights into the role of
the microbiome in obesity and type 2 diabetes. Diabetes Care **38**:159–165.

13. **Leong KSW**, **Derraik JGB**, **Hofman PL**, **Cutfield WS**. 2018. Antibiotics, gut
microbiome and obesity. Clin Endocrinol (Oxf) **88**:185–200.

14. **Lynch SV**, **Boushey HA**. 2016. The microbiome and development of allergic disease.
Curr Opin Allergy Clin Immunol **16**:165–171.

15. **Riiser A**. 2015. The human microbiome, asthma, and allergy. Allergy Asthma Clin
Immunol **11**:35.

16. **Rogers MB**, **Firek B**, **Shi M**, **Yeh A**, **Brower-Sinning R**, **Aveson V**, **Kohl BL**, **Fabio
A**, **Carcillo JA**, **Morowitz MJ**. 2016. Disruption of the microbiota across multiple body
sites in critically ill children. Microbiome **4**:66.

17. **Tremlett H**, **Bauer KC**, **Appel-Cresswell S**, **Finlay BB**, **Waubant E**. 2017. The gut
microbiome in human neurological disease: A review. Ann Neurol **81**:369–382.

18. **Vieira SM**, **Pagovich OE**, **Kriegel MA**. 2014. Diet, microbiota and autoimmune
diseases. Lupus **23**:518–526.

19. **Dethlefsen L**, **Relman DA**. 2011. Incomplete recovery and individualized responses of
the human distal gut microbiota to repeated antibiotic perturbation. Proc Natl Acad Sci
USA **108 Suppl 1**:4554–4561.

20. **Cabral DJ**, **Penumutchu S**, **Reinhart EM**, **Zhang C**, **Korry BJ**, **Wurster JI**, **Nilson
R**, **Guang A**, **Sano WH**, **Rowan-Nash AD**, **Li H**, **Belenky P**. 2019. Microbial

Metabolism Modulates Antibiotic Susceptibility within the Murine Gut Microbiome.
Cell Metab **30**:800–823.
21. **Chang JY**, **Antonopoulos DA**, **Kalra A**, **Tonelli A**, **Khalife WT**, **Schmidt TM**,
Young VB. 2008. Decreased diversity of the fecal Microbiome in recurrent Clostridium
difficile-associated diarrhea. J Infect Dis **197**:435–438.
22. **Preidis GA**, **Versalovic J**. 2009. Targeting the human microbiome with antibiotics,
probiotics, and prebiotics: gastroenterology enters the metagenomics era.
Gastroenterology **136**:2015–2031.
23. **Theriot CM**, **Bowman AA**, **Young VB**. 2016. Antibiotic-Induced Alterations of the
Gut Microbiota Alter Secondary Bile Acid Production and Allow for Clostridium
difficile Spore Germination and Outgrowth in the Large Intestine. mSphere **1**:e00045–
15.
24. **Rea MC**, **Dobson A**, **O'Sullivan O**, **Crispie F**, **Fouhy F**, **Cotter PD**, **Shanahan F**,
Kiely B, **Hill C**, **Ross RP**. 2011. Effect of broad- and narrow-spectrum antimicrobials
on Clostridium difficile and microbial diversity in a model of the distal colon. Proc Natl
Acad Sci USA **108**:4639–4644.
25. **Rafii F**, **Sutherland JB**, **Cerniglia CE**. 2008. Effects of treatment with antimicrobial
agents on the human colonic microflora. Ther Clin Risk Manag **4**:1343–1358.
26. **Lessa FC**, **Winston LG**, **McDonald LC**, **Emerging Infections Program C. difficile**
Surveillance Team. 2015. Burden of Clostridium difficile infection in the United States.
N Engl J Med **372**:2369–2370.
27. **Hryckowian AJ**, **Van Treuren W**, **Smits SA**, **Davis NM**, **Gardner JO**, **Bouley DM**,
Sonnenburg JL. 2018. Microbiota-accessible carbohydrates suppress Clostridium
difficile infection in a murine model. Nature Microbiology **3**:662–669.
28. **Durack J**, **Huang YJ**, **Nariya S**, **Christian LS**, **Mark Ansel K**, **Beigelman A**, **Castro**
M, **Dyer A-M**, **Israel E**, **Kraft M**, **Martin RJ**, **Mauger DT**, **Rosenberg SR**, **King TS**,
White SR, **Denlinger LC**, **Holguin F**, **Lazarus SC**, **Lugogo N**, **Peters SP**, **Smith LJ**,
Wechsler ME, **Lynch SV**, **Boushey HA**, **National Heart, Lung and Blood Institute's**
"AsthmaNet." 2018. Bacterial biogeography of adult airways in atopic asthma.
Microbiome **6**:104.
29. **Wu GD**, **Chen J**, **Hoffmann C**, **Bittinger K**, **Chen Y-Y**, **Keilbaugh SA**, **Bewtra M**,
Knights D, **Walters WA**, **Knight R**, **Sinha R**, **Gilroy E**, **Gupta K**, **Baldassano R**,
Nessel L, **Li H**, **Bushman FD**, **Lewis JD**. 2011. Linking long-term dietary patterns with
gut microbial enterotypes. Science **334**:105–108.
30. **Smits SA**, **Leach J**, **Sonnenburg ED**, **Gonzalez CG**, **Lichtman JS**, **Reid G**, **Knight R**,
Manjurano A, **Changalucha J**, **Elias JE**, **Dominguez-Bello MG**, **Sonnenburg JL**.
2017. Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of
Tanzania. Science **357**:802–806.
31. **Smits SA**, **Marcobal A**, **Higginbottom S**, **Sonnenburg JL**, **Kashyap PC**. 2016.
Individualized Responses of Gut Microbiota to Dietary Intervention Modeled in
Humanized Mice. mSystems **1**:105.
32. **Bisanz JE**, **Upadhyay V**, **Turnbaugh JA**, **Ly K**, **Turnbaugh PJ**. 2019. Meta-Analysis
Reveals Reproducible Gut Microbiome Alterations in Response to a High-Fat Diet. Cell
Host Microbe **26**:265–272.
33. **Ley RE**, **Bäckhed F**, **Turnbaugh P**, **Lozupone CA**, **Knight RD**, **Gordon JI**. 2005.
Obesity alters gut microbial ecology. Proc Natl Acad Sci USA **102**:11070–11075.

34.   **Turnbaugh PJ**, **Ridaura VK**, **Faith JJ**, **Rey FE**, **Knight R**, **Gordon JI**. 2009. The
effect of diet on the human gut microbiome: a metagenomic analysis in humanized
gnotobiotic mice. Science Translational Medicine **1**:6ra14–6ra14.

35.   **Xu Z**, **Knight R**. 2015. Dietary effects on human gut microbiome diversity. Br J Nutr
113 Suppl:S1–5.

36.   **Argueta DA**, **DiPatrizio NV**. 2017. Peripheral endocannabinoid signaling controls
hyperphagia in western diet-induced obesity. Physiology & Behavior **171**:32–39.

37.   **Kanoski SE**, **Hsu TM**, **Pennell S**. 2014. Obesity, Western Diet Intake, and Cognitive
Impairment, pp. 57–62. *In* Omega-3 Fatty Acids in Brain and Neurological Health.
Elsevier.

38.   **Sami W**, **Ansari T**, **Butt NS**, **Hamid MRA**. 2017. Effect of diet on type 2 diabetes
mellitus: A review. Int J Health Sci (Qassim) **11**:65–71.

39.   **Qi L**, **Cornelis MC**, **Zhang C**, **van Dam RM**, **Hu FB**. 2009. Genetic predisposition,
Western dietary pattern, and the risk of type 2 diabetes in men. Am J Clin Nutr **89**:1453–
1458.

40.   **Sonnenburg ED**, **Sonnenburg JL**. 2014. Starving our microbial self: the deleterious
consequences of a diet deficient in microbiota-accessible carbohydrates. Cell Metab
20:779–786.

41.   **Sonnenburg ED**, **Sonnenburg JL**. 2019. The ancestral and industrialized gut
microbiota and implications for human health. Nat Rev Microbiol **17**:383–390.

42.   **Sonnenburg JL**, **Xu J**, **Leip DD**, **Chen CH**, **Westover BP**, **Weatherford J**, **Buhler
JD**, **Gordon JI**. 2005. Glycan foraging in vivo by an intestine-adapted bacterial
symbiont. Science **307**:1955–1959.

43.   **Turnbaugh PJ**. 2017. Microbes and Diet-Induced Obesity: Fast, Cheap, and Out of
Control. Cell Host Microbe **21**:278–281.

44.   **Trompette A**, **Gollwitzer ES**, **Yadava K**, **Sichelstiel AK**, **Sprenger N**, **Ngom-Bru C**,
Blanchard C, **Junt T**, **Nicod LP**, **Harris NL**, **Marsland BJ**. 2014. Gut microbiota
metabolism of dietary fiber influences allergic airway disease and hematopoiesis. Nat
Med **20**:159–166.

45.   **Arpaia N**, **Campbell C**, **Fan X**, **Dikiy S**, **van der Veeken J**, **deRoos P**, **Liu H**, **Cross
JR**, **Pfeffer K**, **Coffer PJ**, **Rudensky AY**. 2013. Metabolites produced by commensal
bacteria promote peripheral regulatory T-cell generation. Nature **504**:451–455.

46.   **Cotillard A**, **Kennedy SP**, **Kong LC**, **Prifti E**, **Pons N**, **Le Chatelier E**, **Almeida M**,
Quinquis B, **Levenez F**, **Galleron N**, **Gougis S**, **Rizkalla S**, **Batto J-M**, **Renault P**,
Doré J, **Zucker J-D**, **Clement K**, **Ehrlich SD**, **Consortium AM**. 2013. Dietary
intervention impact on gut microbial gene richness. Nature **500**:585–588.

47.   **Walker AW**, **Ince J**, **Duncan SH**, **Webster LM**, **Holtrop G**, **Ze X**, **Brown D**, **Stares
MD**, **Scott P**, **Bergerat A**, **Louis P**, **McIntosh F**, **Johnstone AM**, **Lobley GE**, **Parkhill
J**, **Flint HJ**. 2011. Dominant and diet-responsive groups of bacteria within the human
colonic microbiota. ISME J **5**:220–230.

48.   **Fischbach MA**, **Sonnenburg JL**. 2011. Eating for two: how metabolism establishes
interspecies interactions in the gut. Cell Host Microbe **10**:336–347.

49.   **Kashyap PC**, **Marcobal A**, **Ursell LK**, **Smits SA**, **Sonnenburg ED**, **Costello EK**,
Higginbottom SK, **Domino SE**, **Holmes SP**, **Relman DA**, **Knight R**, **Gordon JI**,
Sonnenburg JL. 2013. Genetically dictated change in host mucus carbohydrate

landscape exerts a diet-dependent effect on the gut microbiota. Proc Natl Acad Sci USA
110:17059–17064.

50. **Yatsunenko T**, **Rey FE**, **Manary MJ**, **Trehan I**, **Dominguez-Bello MG**, **Contreras**
M, **Magris M**, **Hidalgo G**, **Baldassano RN**, **Anokhin AP**, **Heath AC**, **Warner B**,
Reeder J, **Kuczynski J**, **Caporaso JG**, **Lozupone CA**, **Lauber C**, **Clemente JC**,
Knights D, **Knight R**, **Gordon JI**. 2012. Human gut microbiome viewed across age and
geography. Nature **486**:222–227.

51. **Wong J**, **de Souza R**, **Kendall C**, **Emam A**, **Jenkins D**. 2006. Colonic health:
Fermentation and short chain fatty acids. Journal of Clinical Gastroenterology **40**:235–
243.

52. **Topping DL**, **Clifton PM**. 2001. Short-chain fatty acids and human colonic function:
Roles of resistant starch and nonstarch polysaccharides. Physiol Rev **81**:1031–1064.

53. **Macfarlane S**, **Macfarlane GT**. 2003. Regulation of short-chain fatty acid production.
Proceedings of the Nutrition Society **62**:67–72.

54. **Cani PD**, **Van Hul M**, **Lefort C**, **Depommier C**, **Rastelli M**, **Everard A**. 2019.
Microbial regulation of organismal energy homeostasis. Nat Metab **1**:34–46.

55. **Chambers ES**, **Preston T**, **Frost G**, **Morrison DJ**. 2018. Role of Gut Microbiota-
Generated Short-Chain Fatty Acids in Metabolic and Cardiovascular Health. Curr Nutr
Rep **7**:198–206.

56. **David LA**, **Maurice CF**, **Carmody RN**, **Gootenberg DB**, **Button JE**, **Wolfe BE**, **Ling**
AV, **Devlin AS**, **Varma Y**, **Fischbach MA**, **Biddinger SB**, **Dutton RJ**, **Turnbaugh**
PJ. 2014. Diet rapidly and reproducibly alters the human gut microbiome. Nature
505:559–563.

57. **Poretsky R**, **Rodriguez-R LM**, **Luo C**, **Tsementzi D**, **Konstantinidis KT**. 2014.
Strengths and limitations of 16S rRNA gene amplicon sequencing in revealing temporal
microbial community dynamics. PLoS ONE **9**:e93827.

58. **Ranjan R**, **Rani A**, **Metwally A**, **McGee HS**, **Perkins DL**. 2016. Analysis of the
microbiome: Advantages of whole genome shotgun versus 16S amplicon sequencing.
Biochem Biophys Res Commun **469**:967–977.

59. **Clooney AG**, **Fouhy F**, **Sleator RD**, **Driscoll AO**, **Stanton C**, **Cotter PD**, **Claesson**
MJ. 2016. Comparing Apples and Oranges?: Next Generation Sequencing and Its
Impact on Microbiome Analysis. PLoS ONE **11**.

60. **Tessler M**, **Neumann JS**, **Afshinnekoo E**, **Pineda M**, **Hersch R**, **Velho LFM**, **Segovia**
BT, **Lansac-Toha FA**, **Lemke M**, **DeSalle R**, **Mason CE**, **Brugler MR**. 2017. Large-
scale differences in microbial biodiversity discovery between 16S amplicon and shotgun
sequencing. Sci Rep **7**:–14.

61. **Shin N-R**, **Whon TW**, **Bae J-W**. 2015. Proteobacteria: microbial signature of dysbiosis
in gut microbiota. Trends Biotechnol **33**:496–503.

62. **Love MI**, **Huber W**, **Anders S**. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol **15**:550.

63. **Franzosa EA**, **McIver LJ**, **Rahnavard G**, **Thompson LR**, **Schirmer M**, **Weingart G**,
Lipson KS, **Knight R**, **Caporaso JG**, **Segata N**, **Huttenhower C**. 2018. Species-level
functional profiling of metagenomes and metatranscriptomes. Nat Methods **15**:962–968.

64. **Meylan S**, **Porter CBM**, **Yang JH**, **Belenky P**, **Gutierrez A**, **Lobritz MA**, **Park J**,
Kim SH, **Moskowitz SM**, **Collins JJ**. 2017. Carbon Sources Tune Antibiotic

Susceptibility in Pseudomonas aeruginosa via Tricarboxylic Acid Cycle Control. Cell
Chem Biol **24**:195–206.

65. **Belenky P**, **Ye JD**, **Porter CBM**, **Cohen NR**, **Lobritz MA**, **Ferrante T**, **Jain S**, **Korry BJ**, **Schwarz EG**, **Walker GC**, **Collins JJ**. 2015. Bactericidal Antibiotics Induce Toxic Metabolic Perturbations that Lead to Cellular Damage. Cell Rep **13**:968–980.

66. **Lobritz MA**, **Belenky P**, **Porter CBM**, **Gutierrez A**, **Yang JH**, **Schwarz EG**, **Dwyer DJ**, **Khalil AS**, **Collins JJ**. 2015. Antibiotic efficacy is linked to bacterial cellular respiration. Proc Natl Acad Sci USA **112**:8173–8180.

67. **Dwyer DJ**, **Belenky PA**, **Yang JH**, **MacDonald IC**, **Martell JD**, **Takahashi N**, **Chan CTY**, **Lobritz MA**, **Braff D**, **Schwarz EG**, **Ye JD**, **Pati M**, **Vercruysse M**, **Ralifo PS**, **Allison KR**, **Khalil AS**, **Ting AY**, **Walker GC**, **Collins JJ**. 2014. Antibiotics induce redox-related physiological alterations as part of their lethality. Proc Natl Acad Sci USA **111**:E2100–9.

68. **Dwyer DJ**, **Kohanski MA**, **Hayete B**, **Collins JJ**. 2007. Gyrase inhibitors induce an oxidative damage cellular death pathway in Escherichia coli. Mol Syst Biol **3**:91.

69. **Westreich ST**, **Treiber ML**, **Mills DA**, **Korf I**, **Lemay DG**. 2018. SAMSA2: a standalone metatranscriptome analysis pipeline. BMC Bioinformatics **19**:175.

70. **Corfield AP**, **Wagner SA**, **Clamp JR**, **Kriaris MS**, **Hoskins LC**. 1992. Mucin degradation in the human colon: production of sialidase, sialate O-acetylesterase, N-acetylneuraminate lyase, arylesterase, and glycosulfatase activities by strains of fecal bacteria. Infect Immun **60**:3971–3978.

71. **Deng Z-L**, **Sztajer H**, **Jarek M**, **Bhuju S**, **Wagner-Döbler I**. 2018. Worlds Apart - Transcriptome Profiles of Key Oral Microbes in the Periodontal Pocket Compared to Single Laboratory Culture Reflect Synergistic Interactions. Front Microbiol **9**:124.

72. **Bubić A**, **Mrnjavac N**, **Stuparević I**, **Łyczek M**, **Wielgus-Kutrowska B**, **Bzowska A**, **Luić M**, **Leščić Ašler I**. 2018. In the quest for new targets for pathogen eradication: the adenylosuccinate synthetase from the bacterium Helicobacter pylori. J Enzyme Inhib Med Chem **33**:1405–1414.

73. **Susin MF**, **Baldini RL**, **Gueiros-Filho F**, **Gomes SL**. 2006. GroES/GroEL and DnaK/DnaJ have distinct roles in stress responses and during cell cycle progression in Caulobacter crescentus. J Bacteriol **188**:8044–8053.

74. **Anglès F**, **Castanié-Cornet M-P**, **Slama N**, **Dinclaux M**, **Cirinesi A-M**, **Portais J-C**, **Létisse F**, **Genevaux P**. 2017. Multilevel interaction of the DnaK/DnaJ(HSP70/HSP40) stress-responsive chaperone machine with the central metabolism. Sci Rep **7**:41341.

75. **Ogata Y**, **Mizushima T**, **Kataoka K**, **Kita K**, **Miki T**, **Sekimizu K**. 1996. DnaK heat shock protein of Escherichia coli maintains the negative supercoiling of DNA against thermal stress. J Biol Chem **271**:29407–29414.

76. **Wong KS**, **Houry WA**. 2012. Novel structural and functional insights into the MoxR family of AAA+ ATPases. J Struct Biol **179**:211–221.

77. **Coyne MJ**, **Comstock LE**. 2008. Niche-specific features of the intestinal bacteroidales. J Bacteriol **190**:736–742.

78. **Porter NT**, **Canales P**, **Peterson DA**, **Martens EC**. 2017. A Subset of Polysaccharide Capsules in the Human Symbiont Bacteroides thetaiotaomicron Promote Increased Competitive Fitness in the Mouse Gut. Cell Host Microbe **22**:494–506.

79.   **Petit C**, **Rigg GP**, **Pazzani C**, **Smith A**, **Sieberth V**, **Stevens M**, **Boulnois G**, **Jann K**,
Roberts IS. 1995. Region 2 of the Escherichia coli K5 capsule gene cluster encoding
proteins for the biosynthesis of the K5 polysaccharide. Mol Microbiol **17**:611–620.

80.   **van Selm S**, **Kolkman MAB**, **van der Zeijst BAM**, **Zwaagstra KA**, **Gaastra W**, **van**
Putten JPM. 2002. Organization and characterization of the capsule biosynthesis locus
of Streptococcus pneumoniae serotype 9V. Microbiology **148**:1747–1755.

81.   **Dougherty BA**, **van de Rijn I**. 1993. Molecular characterization of hasB from an
operon required for hyaluronic acid synthesis in group A streptococci. Demonstration of
UDP-glucose dehydrogenase activity. J Biol Chem **268**:7118–7124.

82.   **Cabral DJ**, **Wurster JI**, **Belenky P**. 2018. Antibiotic Persistence as a Metabolic
Adaptation: Stress, Metabolism, the Host, and New Directions. Pharmaceuticals (Basel)
11:14.

83.   **Rowan AD**, **Cabral DJ**, **Belenky P**. 2016. Bactericidal antibiotics induce programmed
metabolic toxicity. Microb Cell **3**:178–180.

84.   **Allison KR**, **Brynildsen MP**, **Collins JJ**. 2011. Metabolite-enabled eradication of
bacterial persisters by aminoglycosides. Nature **473**:216–220.

85.   **Kohanski MA**, **Dwyer DJ**, **Hayete B**, **Lawrence CA**, **Collins JJ**. 2007. A common
mechanism of cellular death induced by bactericidal antibiotics. Cell **130**:797–810.

86.   **Thomas VC**, **Chittezham Thomas V**, **Kinkead LC**, **Janssen A**, **Schaeffer CR**, **Woods**
KM, **Lindgren JK**, **Peaster JM**, **Chaudhari SS**, **Sadykov M**, **Jones J**, **AbdelGhani**
SMM, **Zimmerman MC**, **Bayles KW**, **Somerville GA**, **Fey PD**. 2013. A dysfunctional
tricarboxylic acid cycle enhances fitness of Staphylococcus epidermidis during β-lactam
stress. mBio **4**:e00437–13–e00437–13.

87.   **Adolfsen KJ**, **Brynildsen MP**. 2015. Futile cycling increases sensitivity toward
oxidative stress in Escherichia coli. Metab Eng **29**:26–35.

88.   **Cho H**, **Uehara T**, **Bernhardt TG**. 2014. Beta-lactam antibiotics induce a lethal
malfunctioning of the bacterial cell wall synthesis machinery. Cell **159**:1300–1311.

89.   **Zinöcker MK**, **Lindseth IA**. 2018. The Western Diet-Microbiome-Host Interaction and
Its Role in Metabolic Disease. Nutrients **10**:365.

90.   **Sanchez KK**, **Chen GY**, **Schieber AMP**, **Redford SE**, **Shokhirev MN**, **Leblanc M**,
Lee YM, **Ayres JS**. 2018. Cooperative Metabolic Adaptations in the Host Can Favor
Asymptomatic Infection and Select for Attenuated Virulence in an Enteric Pathogen.
Cell **175**:146–158.e15.

91.   **Chassaing B**, **Van de Wiele T**, **Gewirtz A**. 2017. O-013 Dietary Emulsifiers Directly
Impact the Human Gut Microbiota Increasing Its Pro-inflammatory Potential and Ability
to Induce Intestinal Inflammation. Inflammatory Bowel Diseases **23**.

92.   **Collins J**, **Robinson C**, **Danhof H**, **Knetsch CW**, **van Leeuwen HC**, **Lawley TD**,
Auchtung JM, **Britton RA**. 2018. Dietary trehalose enhances virulence of epidemic
Clostridium difficile. Nature **553**:291–294.

93.   **Human Microbiome Project Consortium**. 2012. A framework for human microbiome
research. Nature **486**:215–221.

94.   **Singh RK**, **Chang H-W**, **Yan D**, **Lee KM**, **Ucmak D**, **Wong K**, **Abrouk M**, **Farahnik**
B, **Nakamura M**, **Zhu TH**, **Bhutani T**, **Liao W**. 2017. Influence of diet on the gut
microbiome and implications for human health. J Transl Med, 4 ed. **15**:73–17.

95.   **Johnson AJ**, **Vangay P**, **Al-Ghalith GA**, **Hillmann BM**, **Ward TL**, **Shields-Cutler**
RR, **Kim AD**, **Shmagel AK**, **Syed AN**, **Personalized Microbiome Class Students**,

Walter J, **Menon R**, **Koecher K**, **Knights D**. 2019. Daily Sampling Reveals
Personalized Diet-Microbiome Associations in Humans. Cell Host Microbe **25**:789–
802.e5.
96. **Turnbaugh PJ**, **Ley RE**, **Mahowald MA**, **Magrini V**, **Mardis ER**, **Gordon JI**. 2006.
An obesity-associated gut microbiome with increased capacity for energy harvest.
Nature **444**:1027–131.
97. **Turnbaugh PJ**, **Bäckhed F**, **Fulton L**, **Gordon JI**. 2008. Diet-induced obesity is linked
to marked but reversible alterations in the mouse distal gut microbiome. Cell Host
Microbe **3**:213–223.
98. **Thompson LR**, **Sanders JG**, **McDonald D**, **Amir A**, **Ladau J**, **Locey KJ**, **Prill RJ**,
Tripathi A, **Gibbons SM**, **Ackermann G**, **Navas-Molina JA**, **Janssen S**, **Kopylova E**,
Vázquez-Baeza Y, **Gonzalez A**, **Morton JT**, **Mirarab S**, **Zech Xu Z**, **Jiang L**,
Haroon MF, **Kanbar J**, **Zhu Q**, **Jin Song S**, **Kosciolek T**, **Bokulich NA**, **Lefler J**,
Brislawn CJ, **Humphrey G**, **Owens SM**, **Hampton-Marcell J**, **Berg-Lyons D**,
McKenzie V, **Fierer N**, **Fuhrman JA**, **Clauset A**, **Stevens RL**, **Shade A**, **Pollard KS**,
Goodwin KD, **Jansson JK**, **Gilbert JA**, **Knight R**, **Earth Microbiome Project**
Consortium. 2017. A communal catalogue reveals Earth's multiscale microbial
diversity. Nature **104**:11436.
99. **Wang Q**, **Garrity GM**, **Tiedje JM**, **Cole JR**. 2007. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ
Microbiol **73**:5261–5267.
100. **McIver LJ**, **Abu-Ali G**, **Franzosa EA**, **Schwager R**, **Morgan XC**, **Waldron L**, **Segata**
N, **Huttenhower C**. 2018. bioBakery: a meta'omic analysis environment. Bioinformatics
34:1235–1237.
101. **Bolger AM**, **Lohse M**, **Usadel B**. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics **30**:2114–2120.
102. **Langmead B**, **Salzberg SL**. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods **9**:357–359.
103. **Pruesse E**, **Quast C**, **Knittel K**, **Fuchs BM**, **Ludwig W**, **Peplies J**, **Glöckner FO**.
2007. SILVA: a comprehensive online resource for quality checked and aligned
ribosomal RNA sequence data compatible with ARB. Nucleic Acids Research **35**:7188–
7196.
104. **Wood DE**, **Salzberg SL**. 2014. Kraken: ultrafast metagenomic sequence classification
using exact alignments. Genome Biol **15**:R46.
105. **Lu J**, **Breitwieser FP**, **Thielen P**, **Salzberg SL**. 2017. Bracken: estimating species
abundance in metagenomics data. PeerJ Computer Science **3**:e104.
106. **McMurdie PJ**, **Holmes S**. 2013. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS ONE **8**:e61217.
107. **Benjamini Y**, **Hochberg Y**. 1995. Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society
Series B (Methodological) **57**:289–300.
108. **Westreich ST**, **Korf I**, **Mills DA**, **Lemay DG**. 2016. SAMSA: a comprehensive
metatranscriptome analysis pipeline. BMC Bioinformatics **17**:399.
109. **Zhang J**, **Kobert K**, **Flouri T**, **Stamatakis A**. 2014. PEAR: a fast and accurate Illumina
Paired-End reAd mergeR. Bioinformatics **30**:614–620.

110.   **Overbeek R**, **Olson R**, **Pusch GD**, **Olsen GJ**, **Davis JJ**, **Disz T**, **Edwards RA**, **Gerdes**
S, **Parrello B**, **Shukla M**, **Vonstein V**, **Wattam AR**, **Xia F**, **Stevens R**. 2014. The
SEED and the Rapid Annotation of microbial genomes using Subsystems Technology
(RAST). Nucleic Acids Research **42**:D206–14.
111.   **Buchfink B**, **Xie C**, **Huson DH**. 2015. Fast and sensitive protein alignment using
DIAMOND. Nat Methods **12**:59–60.
112.   **Witten IH**, **Bell TC**. 1991. The zero-frequency problem: estimating the probabilities of
novel events in adaptive text compression. IEEE Trans Inform Theory **37**:1085–1094.
113.   **Segata N**, **Izard J**, **Waldron L**, **Gevers D**, **Miropolsky L**, **Garrett WS**, **Huttenhower**
C. 2011. Metagenomic biomarker discovery and explanation. Genome Biol **12**:R60.
114.   **Bushnell B**. 2014. BBMap. Genomics of Energy Environment Meeting. Walnut Creek,
CA.
115.   **Li H**, **Durbin R**. 2010. Fast and accurate long-read alignment with Burrows-Wheeler
transform. Bioinformatics **26**:589–595.

Figures:

Figure 1: Impact of diet and ciprofloxacin administration on murine gut microbiome composition

A. Experimental workflow used in this study. Figure was created with Biorender.com.
B. Alpha diversity of experimental groups as measured by the Shannon Diversity Index. Data are represented as mean ± standard error of the mean (SEM). (**p<0.01, Welch ANOVA with Dunnett T3 test for multiple hypothesis testing).
C. Stacked barplot of the five most abundant bacterial phyla in our dataset. Data are represented as mean + SEM for each phylum.
D. Differentially abundant (Benjamini-Hochberg adjusted p-value < 0.1) bacterial species detected in mice consuming the Western diet (WD). Data are represented as log₂ fold change relative to control diet ± standard error. Bar color denotes phylum level taxonomic classification (blue – *Verrucomicrobia*, red – *Firmicutes*, green – *Bacteroidetes*, purple – *Proteobacteria*, orange – *Actinobacteria*).
E. Heatmap of the change in abundance of the top 90 bacterial species in response to ciprofloxacin on control (NC) and Western (WD) diets. The Interaction column represents the interaction term generated by DESeq2, denoting the impact of diet on the change in abundance of each species to ciprofloxacin. Cell color denotes log₂ fold change of a particular species in response to ciprofloxacin. Heatmap rows were sorted by interaction term value from highest to lowest.

For all analyses, n = 4. For full DESeq2 results, see Additional File 1.

Figure 2:
Ciprofloxacin elicits unique shifts in gene expression on Western and control diets at the MetaCyc pathway level

A. Linear discriminant analysis (LDA) of MetaCyc pathways that were differentially associated with each experimental group. Bar size indicates LDA score and color indicates the experimental group that a MetaCyc pathway was significantly associated with. All LDA scores were generated using LEfSe on unstratified pathway outputs from HUMAnN2. For full pathway names and statistics, see Additional File 2.

B. Differentially expressed (Benjamini-Hochberg adjusted p-value < 0.1) level 1 SEED subsystems in the murine cecal metatranscriptome in ciprofloxacin-treated mice consuming the control (NC) diet. Data are represented as $\log_2$ fold change relative to vehicle control ± standard error. Only features with a base mean ≥ 50 were plotted. See Additional File 3 for full results.

(C – E) Volcano plots of the metatranscriptomic profile of the murine cecal microbiome in vehicle-treated mice consuming Western diet (C), ciprofloxacin-treated mice on the control diet (D), and ciprofloxacin-treated mice on the Western diet (E). Data was generated by aligning metatranscriptomic reads to RefSeq using SAMSA2 and analyzing using DESeq2. Points in red represent transcripts for which a statistically significant change in expression was detected (Benjamini-Hochberg adjusted p-value < 0.1). Select genes of interest are labeled. See Additional File 4 for full results.

For all analyses, n = 4.

Figure 3: Diet and ciprofloxacin alter gene expression within *B. thetaiotaomicron* and *A. muciniphila*

A. Normalized counts of *A. muciniphila* in each experimental group. Data are represented as mean ± SEM. Normalized counts were generated with DESeq2 and subsequently used to perform differential abundance testing. (*p<0.05, ****p<0.0001, Wald test with Benjamini and Hochberg correction). See Additional File 1 for full results.

B. Select differentially expressed (Benjamini-Hochberg adjusted p-value < 0.1) genes of interest in *A. muciniphila* within the cecum of vehicle-treated mice consuming the Western (WD) diet. Data are represented as log₂ fold change relative to control diet ± standard error. See Additional File 5 for full results.

C. Normalized counts of *B. thetaiotaomicron* in each experimental group. Data are represented as mean ± SEM. Normalized counts were generated with DESeq2 and subsequently used to perform differential abundance testing. (*p<0.05, ****p<0.0001, Wald test with Benjamini and Hochberg correction). See Additional File 1 for full results.

1072   D. Select differentially expressed (Benjamini-Hochberg adjusted p-value $<$ 0.1) genes of
1073   interest in *B. thetaiotaomicron* within the cecum of ciprofloxacin-treated mice consuming
1074   the control (NC) diet. Data are represented as $\log_2$ fold change relative to control diet $\pm$
1075   standard error. See Additional File 6 for full results.
1076   E. Select differentially expressed (Benjamini-Hochberg adjusted p-value $<$ 0.1) genes of
1077   interest in *B. thetaiotaomicron* within the cecum of vehicle-treated mice consuming the
1078   Western (WD) diet. Data are represented as $\log_2$ fold change relative to control diet $\pm$
1079   standard error. See Additional File 6 for full results.
1080   For all analyses, n $=4$.

Figure S1: Dietary composition and antibiotic treatment impact the diversity of the gut microbiome

A. Alpha diversity of experimental groups as measured by the Shannon Diversity Index. Data are represented as mean ± standard error of the mean (SEM). (*$p<0.05$, **$p<0.01$, ***$p<0.001$, Welch ANOVA with Dunnett T3 test for multiple hypothesis testing)

B. Principle coordinate analysis of experimental groups as measured by Bray-Curtis dissimilarity. (**$p<0.01$, ***$p<0.001$, Permutational ANOVA).

 Additional Files:
 Additional File 1: Full DESeq2 results of differential abundance testing of top 90 species
 detected by shotgun metagenomics
 Table S1 – Differential abundance testing of the impact of Western diet (WD) consumption on
 the abundance of the top 90 bacterial species detected in our dataset. Log_2 fold change values
 were calculated relative to control diet samples.
 Table S2 – Differential abundance testing of the impact of ciprofloxacin treatment on the
 abundance of the top 90 bacterial species in mice consuming the Western diet (WD). Log_2 fold
 change values were calculated relative to vehicle-treated samples on the WD.
 Table S3 - Differential abundance testing of the impact of ciprofloxacin treatment on the
 abundance of the top 90 bacterial species in mice consuming the control diet (NC). Log_2 fold
 change values were calculated relative to vehicle-treated samples on the NC.
 Table S4 – Interaction term analysis generated by DESeq2 for the impact of host diet
 consumption on changes in species abundance following ciprofloxacin therapy. Log_2 fold change
 values were calculated relative to vehicle-treated samples on the NC.

 Additional File 2: Full LEfSe results from the analysis of MetaCyc pathway abundance
 generated by HUMAnN2. "Class" denotes the experimental group a particular pathway was
 associated with.
 Table S5 – LEfSe analysis of all experimental groups.
 Table S6 – Pairwise LEfSe analysis of vehicle-treated samples from mice consuming either the
 Western (WD) or control (NC) diet.
 Table S7– Pairwise LEfSe analysis of ciprofloxacin- and vehicle-treated samples from mice
 consuming the control diet (NC)
 Table S8 – Pairwise LEfSe analysis of ciprofloxacin- and vehicle-treated samples from mice
 consuming the Western diet (WD)

 Additional File 3: Full DESeq2 results of SEED subsystem abundance generated by SAMSA2
 Table S9 – Differential abundance testing of the impact of Western diet (WD) consumption on
 the abundance of SEED subsystems in the murine cecal metatranscriptome. Log_2 fold change
 values were calculated relative to control diet samples.
 Table S10 – Differential abundance testing of the impact of ciprofloxacin treatment on the
 abundance of SEED subsystems in the murine cecal metatranscriptome in animals consuming the
 Western diet (WD). Log_2 fold change values were calculated relative to vehicle-treated samples
 on the WD.
 Table S11 - Differential abundance testing of the impact of ciprofloxacin treatment on the
 abundance of SEED subsystems in the murine cecal metatranscriptome in animals consuming the
 control diet (NC). Log_2 fold change values were calculated relative to vehicle-treated samples on
 the NC.

 Additional File 4: Full DESeq2 results of RefSeq transcript abundance generated by SAMSA2
 Table S12 – Differential abundance testing of the impact of Western diet (WD) consumption on
 the abundance of RefSeq transcripts in the murine cecal metatranscriptome. Log_2 fold change
 values were calculated relative to control diet samples.
 Table S13 – Differential abundance testing of the impact of ciprofloxacin treatment on the
 abundance of RefSeq transcripts in the murine cecal metatranscriptome in animals consuming

the Western diet (WD). Log$_2$ fold change values were calculated relative to vehicle-treated
samples on the WD.
Table S14 - Differential abundance testing of the impact of ciprofloxacin treatment on the
abundance of RefSeq transcripts in the murine cecal metatranscriptome in animals consuming
the control diet (NC). Log$_2$ fold change values were calculated relative to vehicle-treated samples
on the NC.
Table S15 – Interaction term analysis generated by DESeq2 for the impact of host diet
consumption on changes in RefSeq transcripts abundance following ciprofloxacin therapy. Log$_2$
fold change values were calculated relative to vehicle-treated samples on the NC.
1180
Additional File 5: Full DESeq2 results of transcript abundance analysis of *A. muciniphila* during
dietary intervention and ciprofloxacin treatment
Table S16 – Differential abundance testing of the impact of Western diet (WD) consumption on
the abundance of *A. muciniphila* transcripts within the murine cecal metatranscriptome. Log$_2$
fold change values were calculated relative to control diet samples.
Table S17 – Differential abundance testing of the impact of ciprofloxacin treatment on the
abundance of *A. muciniphila* transcripts within the murine cecal metatranscriptome in animals
consuming the Western diet (WD). Log$_2$ fold change values were calculated relative to vehicle-
treated samples on the WD.
Table S18 - Differential abundance testing of the impact of ciprofloxacin treatment on the
abundance of *A. muciniphila* transcripts within the murine cecal metatranscriptome in animals
consuming the control diet (NC). Log$_2$ fold change values were calculated relative to vehicle-
treated samples on the NC.
Table S19 – Interaction term analysis generated by DESeq2 for the impact of host diet
consumption on changes in *A. muciniphila* transcript abundance following ciprofloxacin therapy.
Log$_2$ fold change values were calculated relative to vehicle-treated samples on the NC.
1197
Additional File 6: Full DESeq2 results of transcript abundance analysis of *B. thetaiotaomicron*
during dietary intervention and ciprofloxacin treatment
Table S20 – Differential abundance testing of the impact of Western diet (WD) consumption on
the abundance of *B. thetaiotaomicron* transcripts within the murine cecal metatranscriptome.
Log$_2$ fold change values were calculated relative to control diet samples.
Table S21 – Differential abundance testing of the impact of ciprofloxacin treatment on the
abundance of *B. thetaiotaomicron* transcripts within the murine cecal metatranscriptome in
animals consuming the Western diet (WD). Log$_2$ fold change values were calculated relative to
vehicle-treated samples on the WD.
Table S22 - Differential abundance testing of the impact of ciprofloxacin treatment on the
abundance of *B. thetaiotaomicron* transcripts within the murine cecal metatranscriptome in
animals consuming the control diet (NC). Log$_2$ fold change values were calculated relative to
vehicle-treated samples on the NC.
Table S23 – Interaction term analysis generated by DESeq2 for the impact of host diet
consumption on changes in *B. thetaiotaomicron* transcript abundance following ciprofloxacin
therapy. Log$_2$ fold change values were calculated relative to vehicle-treated samples on the NC.