1   **Title Page**

2   ***Classification***   Biological Sciences

3   *Title*   Large single-locus effects for maturation timing are mediated via condition
4   variation in Atlantic salmon

5   ***Authors***   Paul V Debes[a], Nicolai Piavachenko[a], Annukka Ruokolainen[a], Outi Ovaskainen[a],
6   Jacqueline Emmanuel Moustakas-Verho[a], Noora Parre[a] Tutku Aykanat[a], Jaakko
7   Erkinaro[b], Craig R Primmer[a]

8   ***Author Affiliation***   [a]Organismal & Evolutionary Biology Research Program / Institute of
9   Biotechnology, University of Helsinki, Viikinkaari 9, 00014, Finland; [b]Natural
10   Resources Institute Finland (Luke), Paavo Havaksen tie 3, 90014, Oulu, Finland

11   ***Corresponding Author*** Name: Paul V Debes; Address: Organismal & Evolutionary Biology Research
12   Program / Institute of Biotechnology, 00014, University of Helsinki, Finland;
13   Phone number: +358 2 9415 9537; Email: paul.debes@dal.ca; ORCID:
14   https://orcid.org/0000-0003-4491-9564

15   ***Keywords***   Sexual maturation, life-history evolution, lipid reserves, resource allocation trade-
16   off, large effect locus

Abstract

Sexual maturation is a pivotal life-history trait that balances the probabilities between mortality and reproduction. Environmental vs. genetic contributions to maturation component traits, such as somatic growth and body condition, remain uncertain because of difficulties in determining causality. In Atlantic salmon, maturation timing associates with a large-effect locus around *vgll3*, which also links with growth, condition, and maturation in mammals. We investigate environmental vs. genetic contributions to maturation and its component traits by combining controlled breeding with common-garden experimentation in two temperatures. We test whether *vgll3* associates with first-year maturation of male salmon and, to avoid reverse causality, whether *vgll3* effects express via growth or condition in the males' non-maturing female relatives. Across 41 families, 4% of males matured in the cold vs. 39% in the warm environment. Maturation rate differed 3.3- to 4.6-fold between *vgll3* genotypes, which also explained around 30% of maturation heritability. Female condition differed up to 2% between *vgll3* genotypes, which also explained 6-17% of condition heritability. Non-significant *vgll3* effects on female length were antagonistic to those for condition but of equal proportional size. When accounting for *vgll3* effects, positive genetic correlations between male maturation and female growth increased, whereas those between male maturation and condition decreased, supporting an antagonistic effect of *vgll3* on growth and condition. The results indicate that large *vgll3* effects on maturation are mediated via large condition effects and suggest *vgll3* as a candidate locus for controlling the resource allocation trade-off between somatic growth and body condition.

Significance Statement

Identifying traits that affect sexual maturation timing and quantifying their relative environmental and genetic importance is a major goal in evolutionary research because of their strong links to fitness. However, cause and effect between maturation timing and such traits often remain unknown. We conducted a common-garden quantitative genetic study using Atlantic salmon to test for genetic contributions of body length and body condition expressed in immature females on maturation of their male relatives. We show that the detection of genetic associations between maturation and length or condition depends on both genetic and environmental factors and suggest that the genetic association with condition chiefly underlies a candidate gene with large effect on maturation that is mediated via variation in body condition.

Maturation timing is a central life-history trait that contributes to maximizing individual survival and

reproductive success and, thereby, per *capita* population growth rate (1-3). Maturation timing is assumed

to be genetically and environmentally controlled via factors including growth or size and body condition (1-

5), which may signal current energy status (6, 7). However, disentangling cause and effect is a major

challenge in studies on maturation and associated traits (8-10). For example, as opposed to growth inducing

maturation, ongoing maturation can temporally increase growth, as is the case for the human puberty

growth spurt. Likewise, the maturation process can lower growth and condition by competing with somatic

growth and depleting reserves, respectively, or affect both via appetite (4, 5, 11). Related to this problem,

fundamental evolutionary knowledge on presence and relative importance of genetic vs. environmental

contributions to maturation timing and their link to maturation component traits, such as growth or condition,

remains limited (12-15).

Contrary to the assumption that life-history traits are comprised of several underlying component traits,

each of which are coded by many loci with small effect on phenotypic variation (1), maturation timing in

Atlantic salmon (*Salmo salar*) associates with a locus explaining a large proportion of phenotypic variation

(33-39%; 16, 17). This finding not only bears implications on evolutionary predictions (1, 18) but also offers

outstanding opportunities for understanding the role of age-specific body size and condition and in

quantifying the relative importance of their genetic vs. environmental contributions to maturation timing. The

large-effect locus positions close to a transcriptional co-factor gene (*vgll3*), hypothesized as a strong

candidate gene (16, 17). Known functions of Vgll3, such as inhibition of adipocyte differentiation in cell lines

in favor of somatic growth processes and its negative transcriptional correlation with body mass and fat

content in mice (19), suggest a mechanistic link to maturation via control of resource allocation as energy

reserves vs. somatic growth. Genetic markers around *vgll3* have also been associated with salmon length

(16), human maturation and growth (10), body condition (20), or condition change during maturation (21),

fostering the idea that the link between *vgll3* and maturation, somatic growth, and condition underlies a

common functional phenotype. However, a comprehensive joint assessment of *vgll3* effects on maturation,

growth, and condition, which requires that the latter two traits are estimated free of maturation-induced

changes, is still missing.

In Atlantic salmon, sexual maturation studies have a long history (5, 22, 23) and this species offers features

allowing for a joint assessment of maturation, growth, and condition, thus enabling independent assessment

of their links with *vgll3*. Specifically, readily available pedigreed hatchery populations allow for planned

breeding of highly fecund individuals. Many offspring with tightly connected pedigrees combined with

common garden experimentation followed by quantitative genetic analyses enable i) dissection of genetic

from environmental effects, ii) estimation of genetic correlations between environments or iii) estimation of

genetic, environmental, and phenotypic correlations between different traits. Perhaps the biggest

advantage however, is the observation that Atlantic salmon males, but usually not females, can mature

during their first year (23). This provides opportunities to estimate environmental and genetic contributions

to maturation in males and those to growth and condition in non-maturing females. By comparing the latter

to those of their male relatives (of which some may show maturation-affected traits), and genetically relating

male maturation to female traits, it is possible to infer the presence and relative importance of genetic vs.

environmental components of maturation timing and its maturation-unbiased component traits.

Here, we implemented a quantitative genetic breeding design for 42 pedigreed Atlantic salmon parents with

known homozygous *vgll3* genotypes and created >5,000 offspring males and females from 41 families with

all *vgll3* genotypes (**Fig. S1**). Longitudinal study of these families in a common-garden environment enabled

general estimates of relative genetic vs. environmental contributions for maturation timing of first-year

Atlantic salmon and its component traits in conjunction with an assessment of specific *vgll3* effects on

maturation, growth, and condition. Further, we evaluated whether a known association of the *vgll3* locus

with maturation (co)expresses via growth, condition, or both, by testing whether male maturation probability

correlates genetically with maturation-unbiased female somatic growth or body condition, or with both, and

by quantifying the *vgll3* contribution to the genetic between trait correlation. By replicating all families in two

environments with a life-time 2°C water-temperature difference across a seasonal temperature curve (a

likely global-warming scenario (24); **Fig. S2**), we were also able to assess environmental influences of wide

relevance on our estimates.

Results and Discussion

We used a multivariate generalized animal model to test for associations between *vgll3* and each of the

five investigated traits of male maturation probability (maturation, MAT), sex-specific somatic growth

(length, LEN), and sex-specific body condition (condition, CON). To avoid reverse causality (i.e., inferring

condition- or length-mediated *vgll3* effects on maturation that are truly maturation-mediated *vgll3* effects on

somatic growth or condition) we focused on *vgll3* effects on maturation in males and length and condition

in females (of which none matured). To allow for wider interpretation and relevance of our results, we report

maturation results, consistent with a genetic threshold model, on the probit or observational probability

scale, which are relevant for breeding-value based or phenotypic selection, respectively (25). Likewise, we

report results for length and condition on the proportional scale, which is relevant to growth processes at

the individual level, or on the (phenotypic standard deviation, psd) standardized effects size scale, which is

relevant among individuals at the population level.

When fully accounting for experimental randomization and relatedness, we detected strong *vgll3* effects on

maturation probability and condition, but not on length, in both temperature environments, whereby the

vgll3$^*E$ allele associated with both higher maturation and higher condition (**Fig. 1**). These results provide

the first experimental confirmation that the allele associated with earlier maturation following marine

migration (16, 17, 26) has consistent effects in males maturing in freshwater, and that the same allele also

associates with a higher condition. All focal trait means were much higher in the warm than in the cold

environment (**Fig. 1;** posterior temperature contrasts, 95% credible interval [95% CI]: probit scale, MAT_{Warm}

- MAT_{Cold} = 2.0, 1.5-2.6; psd scale, female LEN_{Warm} - LEN_{Cold}= 1.5, 1.3-1.7; female CON_{Warm} - CON_{Cold}=

0.92, 0.57-1.29). Despite these strong environmental effects on trait means, *vgll3* effect estimates did not

differ between environments for any trait (**Fig. S3**).

Fig. 1. **_Vgll3_ genotypes mean estimates and additive _vgll3_ effect estimates for the sex-specific responses of Atlantic salmon maturation, length, and condition.** The panels show back-transformed _vgll3_ genotype-specific mean estimates with 95% credible intervals for sex-specific maturation probability (**A, B,** no female maturation occurred), body length (**C, D**), and body condition (standardized mass for a geometric mean-sized fish of 9.1 cm, **E, F**) from a common multivariate generalized animal model ($N_{families}$ = 41, N_{male} = 2,534, N_{female} = 2,611). Estimates are shown for either a warm or cold temperature environment. Sex-specific regression lines with 95% credible bands show additive _vgll3_ effects estimates (α) that are either estimated across both temperature environments (maturation, length, female condition) or that are estimated for each environment (male condition). Mean α estimates with 95% credible intervals are also shown. Mean _vgll3_ genotype estimates and _vgll3_ α estimates originate from similar multivariate models differing in whether _vgll3_ effects were specified either as factorial or continuous terms, respectively.

We also formally tested jointly for additive (α) and dominant (δ) effects for all traits, because sex-specific dominance (i.e., _vgll3_-by-sex epistasis) has been reported for sea age at maturity in one of two previous studies (16, but see 26). We did not detect dominance for any trait (**Fig. S4**), which verified a condition for meaningfully associating _vgll3_ effects between male and female traits that would not be given under sex-specific dominance. After removing all dominance effects from the model, we estimated common additive effects for all traits across temperature environments (**Fig. 1 B, D, F**). For female length, we estimated a non-significant but negative additive _vgll3_*E allele effect on the psd scale of α = -0.06 (95% CI = -0.16-0.05). This effect translates to a non-significant 1% lower body length per _vgll3_*E allele (**Fig 1 D**). For female condition, we estimated an additive _vgll3_*E allele effect on the psd scale of α = 0.20 (95% CI = 0.09-0.33). This among-individual additive _vgll3_ effect on condition is much larger than that estimated for length. However, it translates to the same proportional effect as estimated for length, but in the opposite direction. Specifically, a non-significant 1% body length decrease per _vgll3_*E allele is accompanied by a significant 1% body mass increase (**Fig 1 F**). Although the _vgll3_ effect on length is non-significant and of little, if any,

importance at the population level (but may still represent a within-individual trade off, discussed below), the same effect size of 1% body mass (condition) increase per *vgll3**E allele is very likely relevant for maturation. Specifically, male salmon gonad development is dependent on body reserves, such as fat (4, 5), and gonads constitute 5-9% of body mass at spawning (27, 28). Thus, condition variation explained by *vgll3* genotype may finance up to 40% of the required extra mass (ignoring unknown reserve-to-gonad mass conversion factors). These results provide the first indication that the large effect of *vgll3* on maturation is, at least partly, mediated by a large effect of *vgll3* on condition. We strengthen this further by providing additional evidence below where we quantify *vgll3* effects on trait-specific heritabilities and between-trait genetic correlations.

Fig. 2.   Heritability and genetic correlation estimates from a multivariate generalized animal mixed model, for the traits of Atlantic salmon male maturation (MAT), sex-specific length (LEN), and sex-specific condition (CON). The top row shows, for each trait, sex-specific heritabilities (h^2) and the sex-specific or between-sex genetic correlations (R_G), estimated either specific for each temperature environment (**A, B**) or between temperature environments (**C**; genetic between-environment correlations within and, reciprocally, between sexes). The lower row shows phenotypic (R_P) and genetic correlations (R_G) between maturation and length (MAT.LEN) or maturation and condition (MAT.CON) within sexes and between sexes (only for R_G) in the warm (**D**) and cold (**E**) environments. All estimates are depicted with 95% credible intervals. All estimates were obtained by two similar multivariate models differing in whether additive *vgll3* effects were included or excluded as mean effects as indicated in the legends.

The multivariate model also enabled us to estimate environment- and trait-specific heritabilities (h^2) and trait-specific genetic correlations (R_G) between temperature environments (both contribute to genotype-by-environment interactions (29)) and also to estimate how much *vgll3* contributes to either. The heritability of a trait quantifies the proportion of phenotypic variance explained by the sum of all underlying additive genetic effects and predicts the response to selection. The genetic between-environment correlation quantifies the genotype re-ranking between environments and predicts, in conjunction with heritability estimates, the response to selection across environments (30). Fitting each model with and without mean-

effect specified additive *vgll3* effects, enabled us to quantify the relative genetic and phenotypic
contributions of additive genetic *vgll3* effects and thus the trait-specific evolutionary importance of *vgll3*
within and across environments.

Remarkably, male maturation heritability (h^2MAT) on the probit scale was three times larger in the warm
environment than in the 2 °C colder environment (**Fig. 2 A, B;** posterior temperature contrast, 95% CI;
h^2MAT$_{Warm}$ - h^2MAT$_{Cold}$ = 0.32, 0.21-0.41). Translated to the observed scale, maturation heritability was four
times higher in the warm environment (posterior mean, 95% CI; h^2MAT$_{Warm}$ = 0.41, 0.32-0.50; h^2MAT$_{Cold}$ =
0.10, 0.04-0.16; posterior temperature contrast, 95% CI; h^2MAT$_{Warm}$ - h^2MAT$_{Cold}$ = 0.28, 0.17-0.38;). Further,
the between-temperature genetic correlation estimate (R_GMAT) for maturation was < 0.8 (posterior mean,
95% CI; R_GMAT$_{Warm.Cold}$ = 0.70, 0.45-0.93), which is somewhat lower than most previous genotype-by-
temperature environment estimates for maturation of related species (29). Importantly, R_G < 0.8 indicates
breeding value re-ranking between environments with a mere 2°C temperature difference and thus is of
relevance to natural and aquaculture settings (29, 31). This is of particular relevance, in combination with
the pronounced differences in maturation heritability (30), under a 2 °C global-warming scenario (24). For
maturation, additive *vgll3* effects explained 29.3 and 31.2% of the observed-scale heritability and 12.0 and
10.0% of the observed-scale phenotypic variance in the warm and cold environments, respectively. On the
probit scale, additive *vgll3* effects explained 29.9 and 31.9% of heritability and 13.8 and 2.9% of phenotypic
variance, respectively. Additive *vgll3* effects also explained 11.4% of the between-environment genetic
correlation for maturation (**Fig. 2 C**), which supports an interpretation that the abovementioned non-
significant *vgll3*-by-temperature interaction effects imply environmentally stable *vgll3* effects relative to
other additive genetic effects on maturation.

Female length heritability was, in contrast to that for maturation probability, similar between environments
(posterior mean, 95% CI; h^2LEN$_{Warm}$ - h^2LEN$_{Cold}$ = -0.03, -0.19-0.13) and the between-environment genetic
correlation of R_GLEN$_{Warm.Cold}$ = 0.78 (0.58-0.94) indicated only minor genotype re-ranking between
environments (**Fig. 2 A, B, C**). This between-environment genetic correlation contrasts with much lower
estimates in other fish species (29). Accounting for *vgll3* effects, unexpectedly, increased female length
heritability, as opposed to lowering it as observed for maturation heritability, though by only 1.7 and 6.2%
in the warm and cold, respectively (**Fig 2 A, B, C).** When we investigated underlying length variance
components between models, we found that including *vgll3* effects re-allocated variance from the residuals
to the additive genetic component, which is not expected if *vgll3* comprises additive genetic effects on
length.

Female condition heritability was, like that for female length and in contrast to that for maturation probability,
similar between environments (h^2CON$_{Warm}$ - h^2CON$_{Cold}$ = -0.09, -0.30-0.08) and the between-environment
genetic correlation of R_GCON$_{Warm.Cold}$ = 0.84 (95% CI = 0.71-0.96) indicated very little genotype re-ranking
between environments (**Fig. 2 A, B, C**). Additive *vgll3* effects explained 16.2 and 6.5% of the condition
heritability and 5.2 and 2.5% of the phenotypic variance in the warm and cold environments, respectively.
And, *vgll3* effects also explained 4.4% of the between-environment genetic correlation for female condition

(**Fig. 2 C**), which is noticeable but less than what *vgll3* explained for maturation of the same parameter.
Thus, these results support, like for maturation probability, the presence of environmentally stable *vgll3*
effects, but in the presence of a relatively more environmentally stable genetic background for condition
than for maturation.

By quantifying the proportion of the additive *vgll3* effect contribution to trait-specific heritability, we were
unable to support a hypothesized *vgll3* association for male maturation via female length but support an
association for male maturation via female condition. Quantitative contribution estimates of the genetic
marker to the heritability were sufficiently large to consider *vgll3* a large-effect locus for both maturation and
condition. Nonetheless, *vgll3* contributions to the phenotypic maturation-timing variance were smaller than
previous estimates at a later life stage (16, 17, 26), which could have several biological, methodological, or
statistical reasons, discussion of which goes beyond the scope of this manuscript. Furthermore, by having
observed the same direction for *vgll3* effects between male maturation and female condition, we had
obtained the first indication that *vgll3*-induced variation for male maturation and female condition may
underlie a common *vgll3*-related molecular process. We wanted to further foster this idea by quantifying the
vgll3 contribution to the genetic correlation between male maturation and female length (MAT.LEN) and
between male maturation and female condition (MAT.CON). To do so, we had borrowed quantitative
genetic methodology from animal breeding (an extension of methods reviewed in 32), whose
implementation through the covariance structure of the multivariate animal model was possible by our study
design, and estimated the between-sex genetic correlation between different traits.

We did not detect genetic correlations between male maturation and female (or even male) length, although
estimates in both environments were positive and the relationship-controlled phenotypic correlation
estimate (R_P) (33) between male maturation and male length was significant in both environments (**Fig 2**
D, E). All correlation estimates (genetic and phenotypic) between male maturation and female or male
length increased after accounting for *vgll3* effects (**Fig 2 D, E**). Thus, we were able to further exclude length
as a mediator of *vgll3* effects on maturation, which requires inferring a positive genetic covariance among
vgll3 effects between traits, but instead found a statistical behavior that supports a negative genetic
covariance among *vgll3* effects between maturation and length.

Between male maturation and female (and male) condition, we, however, detected a positive genetic
correlation, although only in the warm environment (we discuss this environmental difference below). After
accounting for *vgll3* effects, all genetic correlations between male maturation and female or male condition
decreased, as is expected under scenarios of either the presence of pleiotropic *vgll3* effects on condition
and maturation, or when one trait mediates the genetic-marker associated variation of the other. The
decreasing effect was strongest for the genetic correlation between male maturation and female condition,
which even rendered non-significant when accounting for *vgll3* effects. Ignoring a *vgll3* pleiotropy scenario,
this latter result suggests a positive genetic covariance among *vgll3* effects between male maturation and
female condition, which may even be dominated by *vgll3* effects and, thereby, suggests a *vgll3* effect

mediation on maturation via condition and not *vice versa* (because maturation variation cannot account for
condition variation in immature females).

Although many effects for length were statistically non-significant, we cannot entirely reject a role for length.
Generally, growth or length and maturation initiation are assumed to associate positively (4, 5). By inferring
a negative covariance among *vgll3* effects between maturation and length we were able to rule out that
length mediates *vgll3* effects on maturation. However, the inferred negative genetic covariance among *vgll3*
effects between maturation and length support a scenario for a *vgll3* governed resource allocation trade-
off. Such trade-off was also, although weakly, indicated by the non-significant 1% length decrease per
*vgll3**E allele that was accompanied by a significant 1% condition increase. This result is typical in life-
history research for traits that represent within-individual resource allocation trade-offs in the presence of a
much larger among-individual variation for resource acquisition relative to the within individual resource-
allocation conflict (1). The idea of a *vgll3*-mediated resource allocation trade-off makes also biological
sense. This is not only because resource allocation theory predicts that energy allocated to condition cannot
concurrently be allocated to growth, but also due to the suggested role of Vgll3 in controlling mesenchymal
cell fate into either adipocytes (thus increasing body condition) or bone and cartilage lineages (thus
increasing somatic growth) (19). However, a final conclusion for our tentative results requires additional
research.

A remaining question is why we detected a significant genetic correlation between male maturation and
female condition in the warm, but not the cold, environment. We suspect that the higher maturation rate
and higher maturation heritability in the warm provided sufficient information to detect the genetic correlation
(31). In contrast, many males in the cold environment with high breeding values for condition did not mature
because of the much lower average condition in the cold or due to other lower temperature-related causes,
such as not exceeding growth or size thresholds needed for maturation (4, 5), masking an otherwise
positive genetic correlation.

Extending these thoughts, we detected environmental effects on condition that were consistent with the
notion that both condition and maturation covary positively with temperature. Specifically, relative to the
cold environment and when translated to the proportional scale we detected a 4.6% (95% CI = 2.9-6.4%)
higher female condition in the warm environment, where we had also detected a much higher maturation
probability (**Fig 1 A, B, E, F**). Thus, the environmental effect of a global-warming-relevant 2°C temperature
difference (24) on condition was more than twice the maximum genetic *vgll3* effect (EE vs. LL: 2%) and
may have contributed considerably to the large environmental effect on maturation probability. Similar
positive relationships between body condition and water temperature have previously been observed (34,
35). We suspect that temperature effects on maturation via condition explain, at least partly, unexpectedly
high maturation rates in heated salmon rearing facilities (36) and why effects on size and maturation timing
vary between growth acceleration through increase in feed vs. temperature (37, 38). Important to
consistency of natural and human selection success across temperatures (30), the large average condition
temperature difference observed here was evident in the presence of both high genetic correlations (> 0.8

for both sexes) and similar condition heritabilities between temperatures (**Fig 2 G, H, I**). Furthermore, *vgll3*
effect estimates on both female condition and on male maturation did not differ between temperature
environments, and *vgll3* effects contributed to the between-environment genetic correlations for both traits.
Thus, the environmental stability of *vgll3* effects on condition contributes to the environmental stability of
the remaining additive genetic effects on condition.

These new results suggest - together with the previous results on maturation (16, 17, 26) - the presence of
environmentally stable *vgll3* effects on both condition and maturation. A large share of *vgll3* effects for the
positive genetic correlation between maturation and condition predicts rapid evolutionary co-responses to
selection for either trait, but also predicts that their genetic correlation is sensitive to *vgll3* allele frequencies.
The sex-specific results, together with previous biological knowledge on their causal association (4, 5),
indicate that large *vgll3* effects on maturation are likely mediated via large condition effects and suggest *vgll3*
as a candidate locus for controlling the resource allocation trade-off between somatic growth and body
condition.

Materials and Methods

Fish population, breeding and experimental design, data collection

The experimental cohort was parented by pedigreed hatchery fish maintained by the Natural Resources
Institute Finland (Laukaa, Finland). The hatchery-stock ancestors originated from the River Neva, Russia,
which drains into the Baltic Sea. In November 2017, we crossed 48 parents with known *vgll3* genotype as
12 2x2 factorials of unrelated *vgll3* homozygous individuals; each factorial yielded four reciprocal *vgll3*
offspring genotypes (EE, EL, LE, LL; details on realized design and resulting pedigree in **Fig. S1**). We
reared the experimental cohort in a recirculation system controlled for water temperature, oxygen, dissolved
nitrate components, and natural light cycle, which affect growth or sexual maturation timing (4, 5). We split
each family by randomizing equal number of individuals into two egg-incubator replicates (families
separated; kept in darkness) and, at first feeding and after pooling incubator replicates, into eight similar
tank replicates for each of two water temperatures (totaling four incubators and 16 tanks). Water
temperatures followed a seasonal cycle with a 2°C difference, referred to as "warm" or "cold" (warm, range
= 6.3-17.7 °C; cold, range = 4.1-16.0 °C). Fish were fed *ad libitum* using a commercial salmon diet starting
2018-03-09 (warm) or 2018-04-27 (cold). Once fish size allowed passive integrated transponder tagging
(warm: August 2019; cold: September 2019) to enable re-identification, we anesthetized (using
methanesulfonate), fin clipped, and tagged individuals; fin clips allowed for genotyping individuals to assign
family, determine molecular sex, and confirm *vgll3* genotype. Starting at tagging, we followed trajectories
for fork length (± 1 mm) and wet mass (± 0.01 g) in 3-6-week measurement intervals until final spawning
time (December 2018) when we determined maturity status by checking for extruding milt by gently pressing
the abdomen and confirmed sex and maturity status in 84% of the fish by culling and dissection (N = 4,313).
Once individual identification was possible, we applied a feed-restriction treatment (iterating: *ad libitum*
feeding for two days, no feed for one day, *ad libitum* feeding for two days, no feed for one day) that was
crossed with the temperature treatment for a five-week period (September 2018), but we did not detect any

feed-restriction effect on maturation (**Fig. S5**). Because length and condition data included in models
originated from the earliest common time point, before the feed restriction, we omitted the feed restriction
term from further analyses. Animal experimentation was conducted according to license ESAVI-2778-2018.
Genotypes and molecular sex of both parents and the experimental cohort were determined using a
multiplex-PCR for 177 single nucleotide polymorphisms (SNPs) of a previously described SNP panel (39),
followed by Ion Torrent (988 potential parents) or Illumina sequencing (used parents, experimental offspring
cohort). Using a subset of 141 unlinked, polymorphic SNPs, we reconstructed the parents of potential
parents (the experimental cohort's grandparents) with maximum likelihood (40) and assigned the
experimental individuals to their parents with a likelihood approach (41). Merged information about
reconstructed grandparents and assigned parents of the experimental cohort yielded a three-generation
pedigree (**Fig. S1 B**) on which we based the relationship matrix utilized in animal-model analyses (42).

Data analysis

We fitted a series of uni- and multivariate general and generalized linear animal models to data for
maturation binaries, assessed towards the end of spawning time in December, and length and condition
records, assessed four months before spawning time in late summer. We defined condition as deviation
from the (temperature-specific) slope of logarithmic mass on logarithmic length - a correlate of salmon parr
lipid content (43). Length and condition data were first log-transformed, then mean-centered and variance
scaled to estimate biologically meaningful proportional and phenotypic variance-standardized effects.

To determine the multivariate model structure and allow for comparisons between uni- or bivariate model
estimates with multivariate model estimates (**Fig S7**), we first fit univariate models for the binary response
of male maturation and bivariate models for the continuous responses of sex-specific length or condition.
We estimated means and (co)variances for models with the binary maturation response (including the
multivariate model) using Bayesian Markov Chain Monte Carlo simulations (44) with the R-package
MCMCglmm (45). For the continuous responses, we initially fitted bivariate models under REML (44) using
ASReml (46). We then fit the multivariate model corresponding to the chosen univariate (maturation) or
bivariate models (sex-specific length or condition), and by adding the required between-trait covariances
for the additive genetic effects (2,599 [males] or 5,209 [both sexes] relationship-matrix-predicted breeding
values (Henderson 1973) *alias* animals), common environmental effects (16 tanks), maternal effects (21
dams), and individual environmental effects including measurement error and non-additive genetic effects
(2534 [males] or 5145 [both sexes] residuals). We fitted the models with probit-link function and residual
variance fixed to one for maturation, corresponding to genetic threshold models (47), and with identity link
function for length and condition. We did not detect maternal effects on any trait (**Fig. S6, Table S1, Table
S2**; we removed dam effects as a consequence) and no common environmental effects on maturation (**Fig.
S6;** we still kept the experimental tank effects), but on length and condition, which contributed up to 6 and
15% to the total phenotypic variance, respectively (**Fig. S7**). We started model selection with mean-effect
interactions and removed non-significant effects (**Fig. S3, S5**). Final models followed the general equation
(per trait), with colon indicating term interaction and variance terms in italic: y ~ Intercept + Temperature +

Vgll3 + *Temperature:Tank* + *Temperature:Animal* + *Temperature:Residual*. The Vgll3 model term refers to

either three genotypes (reciprocal heterozygote differences were absent; posterior EL-LE contrast, 95% CI:

MAT_{EL}-MAT_{LE} = 0.00, -0.47-0.49), or to the additive allelic effect (we fitted both). Covariance matrices

across temperatures for initially fit univariate models were diagonal for tank and residual effects and

unstructured for dam and animal effects. The latter allows estimating dam- and additive genotype-by-

environment effects (30). We used (co)variance priors following a χ_1^2 distribution in Bayesian univariate

models (44) or priors that resulted in flat priors for heritabilities and correlations in multivariate models. We

based estimates on 5,000 or 1,000 retained iterations after 50,000 burnin iterations and sampling every

1,250 and 2,500 or 3,000 iterations (uni- and multivariate models, respectively). We confirmed model

convergence by trace plot inspection and ensuring lag-two autocorrelation < 0.1 for mean and (co)variance

estimates. For the multivariate models (five response traits: sex-specific length and condition, male

maturation) we expanded the covariance matrices to temperature-specific block-diagonals for

Trait:Temperature:Tank (two 5x5 blocks), a full *Trait:Temperature:Animal* genetic covariance matrix

(10x10), and temperature- and sex-specific block-diagonals for *Trait:Temperature:Residual* (two 2x2

female and two 3x3 male blocks).

We tested and estimated additive and dominant *vgll3* effects by replacing the factorial Vgll3 term by

appropriate covariates (α: -1, 0, 1; δ = 0, 1, 0). To obtain observed-scale maturation probability parameter

estimates, we integrated over marginal model predictions and used methodology implemented in the R-

package QGglmm (25), applied to each retained MCMC iteration to estimate credible intervals. We

calculated genetic, environmental, and phenotypic correlations following Searle (33). All necessary data

and an R-script mirroring the final uni- and multivariate modeling have been deposited in the DRYAD data

repository (available during review, doi after acceptance).

Acknowledgments

We thank S. Andrew, O. Andersson, Y. Czorlich, A. House, J. Kurko, M. Lindqvist, N. Lorenzen, O. Mehtälä,

K. Mobley, S. Papakostas, V. Pritchard, K. Salminen, M. Sinclair-Waters, S. Tillanen, J.-P. Verta, and K.

Zueva, for help related to gamete stripping, sample processing, tagging, phenotyping, and fish husbandry,

P. de Villemereuil for multivariate modelling advice, and J. Hutchings for improving an earlier manuscript

version. We acknowledge the Academy of Finland, the European Research Council, and the University of

Helsinki for funding.

Author contributions

P.V.D. and C.R.P. designed research; P.V.D., N.P., A.R., O.O., J.M.-V., N.P., T.A., and C.R.P. performed
research; J.E. contributed salmon gametes; P.V.D. analyzed data; P.V.D. and C.R.P. wrote the paper.

1. Roff DA (2002) *Life History Evolution* (Sinauer Associates, Inc, Sunderland, Massachusetts) p 527.
2. Wells JCK, Nesse RM, Sear R, Johnstone RA, & Stearns SC (2017) Evolutionary public health:
introducing the concept. *The Lancet* 390(10093):500-509.
3. Stearns SC (1992) *The Evolution of Life Histories* (Oxford University Press, New York) p 264.
4. Taranger GL*, et al.* (2010) Control of puberty in farmed fish. *Gen. Comp. Endocrinol.* 165(3):483-515.

5. Andersson E, Taranger GL, Wargelius A, & Schulz RW (2018) Puberty in Fish. *Encyclopedia of*
Reproduction (Second Edition), ed Skinner MK (Academic Press, Oxford), pp 426-429.
6. Danielsen ET, Moeller ME, & Rewitz KF (2013) Nutrient signaling and developmental timing of
maturation. *Curr. Top. Dev. Biol.* 105:37-67.
7. Dupont J, Reverchon M, Bertoldo MJ, & Froment P (2014) Nutritional signals and reproduction. *Mol.*
Cell. Endocrinol. 382(1):527-537.
8. Kause A, Ritola O, Paananen T, Mäntysaari E, & Eskelinen U (2003) Selection against early maturity in
large rainbow trout *Oncorhynchus mykiss*: the quantitative genetics of sexual dimorphism and genotype-
by-environment interactions. *Aquaculture* 228(1-4):53-68.
9. Bell JA*, et al.* (2018) Influence of puberty timing on adiposity and cardiometabolic traits: a Mendelian
randomisation study. *PLoS Medicine* 15(8):e1002641.
10.Cousminer DL*, et al.* (2013) Genome-wide association and longitudinal analyses reveal genetic loci
linking pubertal height growth, pubertal timing and childhood adiposity. *Hum. Mol. Genet.* 22(13):2735-
2747.
11.Stearns SC & Koella JC (1986) The evolution of phenotypic plasticity in life-history traits: predictions of
reaction norms for age and size at maturity. *Evolution* 40(5):893-913.
12.Dunlop ES, Heino M, & Dieckmann U (2009) Eco-genetic modeling of contemporary life-history
evolution. *Ecol. Appl.* 19(7):1815-1834.
13.Law R (2007) Fisheries-induced evolution: present status and future directions. *Mar. Ecol. Prog. Ser.*
335:271-277.
14.Enberg K*, et al.* (2012) Fishing-induced evolution of growth: concepts, mechanisms and the empirical
evidence. *Mar. Ecol.* 33(1):1-25.
15.Gjedrem T & Baranski M (2005) *Selective Breeding in Aquaculture: An Introduction* (Springer, London)
p 221.
16.Barson NJ*, et al.* (2015) Sex-dependent dominance at a single locus maintains variation in age at
maturity in salmon. *Nature* 528(7582):405-408.
17.Ayllon F*, et al.* (2015) The *vgll3* locus controls age at maturity in wild and domesticated Atlantic salmon
(*Salmo salar* L.) males. *PLoS Genet.* 11(11):e1005628.
18.Kuparinen A & Hutchings JA (2017) Genetic architecture of age at maturity can generate divergent and
disruptive harvest-induced evolution. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 372(1712).
19.Halperin DS, Pan C, Lusis AJ, & Tontonoz P (2013) Vestigial-like 3 is an inhibitor of adipocyte
differentiation. *J. Lipid Res.* 54(2):473-481.
20.Nakayama K, Ohashi J, Watanabe K, Munkhtulga L, & Iwamoto S (2017) Evidence for very recent
positive selection in Mongolians. *Mol. Biol. Evol.* 34(8):1936-1946.
21.Tu W*, et al.* (2015) Associations between menarche-related genetic variants and pubertal growth in male
and female adolescents. *J. Adolesc. Health* 56(1):66-72.
22.Meerburg DJ (1986) *Salmonid age at maturity* (Department of Fisheries and Oceans, Ottawa) p 118.
23.Marschall EA*, et al.* (1998) A framework for understanding Atlantic salmon (*Salmo salar*) life history.
Can. J. Fish. Aquat. Sci. 55(S1):48-58.
24.IPCC (2014) Future Climate Changes, Risk and Impacts. *Climate Change 2014: Synthesis Report.*
Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental
Panel on Climate Change, eds Core Writing Team, Pachauri RK, & Meyer LA (IPCC, Geneva,
Switzerland), pp 56-74.
25.de Villemereuil P, Schielzeth H, Nakagawa S, & Morrissey M (2016) General methods for evolutionary
quantitative genetic inference from generalized mixed models. *Genetics* 204(3):1281-1294.
26.Ayllon F*, et al.* (2019) The influence of *vgll3* genotypes on sea age at maturity is altered in farmed mowi
strain Atlantic salmon. *BMC Genet.* 20(1):44.
27.Trombley S, Mustafa A, & Schmitz M (2014) Regulation of the seasonal leptin and leptin receptor
expression profile during early sexual maturation and feed restriction in male Atlantic salmon, *Salmo*
salar L., parr. *Gen. Comp. Endocrinol.* 204:60-70.
28.Rowe DK, Thorpe JE, & Shanks AM (1991) Role of fat stores in the maturation of male Atlantic salmon
(*Salmo salar*) parr. *Can. J. Fish. Aquat. Sci.* 48(3):405-413.
29.Sae-Lim P, Gjerde B, Nielsen HM, Mulder H, & Kause A (2016) A review of genotype-by-environment
interaction and micro-environmental sensitivity in aquaculture species. *Rev. Aquacult.* 8(4):369-393.
30.Falconer DS (1952) The problem of environment and selection. *Am. Nat.* 86(830):293-298.

31. Robertson A (1959) The sampling variance of the genetic correlation coefficient. *Biometrics* 15(3):469-
485.
32. Wolak ME, Roff DA, & Fairbairn DJ (2015) Are we underestimating the genetic variances of dimorphic
traits? *Ecol. Evol.* 5(3):590-597.
33. Searle SR (1961) Phenotypic, genetic and environmental correlations. *Biometrics* 17(3):474-480.
34. Dwyer WP & Piper RG (1987) Atlantic salmon growth efficiency as affected by temperature. *Prog. Fish.*
Cult. 49(1):57-59.
35. Tromp JJ*, et al.* (2018) Chronic exposure to increased water temperature reveals few impacts on stress
physiology and growth responses in juvenile Atlantic salmon. *Aquaculture* 495:196-204.
36. Good C & Davidson J (2016) A review of factors influencing maturation of Atlantic salmon, *Salmo salar*,
with focus on water recirculation aquaculture system environments. *J. World Aquacult. Soc.* 47(5):605-
632.
37. Berrigan D & Charnov EL (1994) Reaction norms for age and size at maturity in response to temperature:
a puzzle for life historians. *Oikos* 70(3):474.
38. Shapiro Goldberg D, van Rijn I, Kiflawi M, Belmaker J, & Heino M (2019) Decreases in length at
maturation of Mediterranean fishes associated with higher sea temperatures. *ICES J. Mar. Sci.*
76(4):946-959.
39. Aykanat T, Lindqvist M, Pritchard VL, & Primmer CR (2016) From population genomics to conservation
and management: a workflow for targeted analysis of markers identified using genome-wide approaches
in Atlantic salmon *Salmo salar*. *J. Fish Biol.* 89(6):2658-2679.
40. Jones OR & Wang J (2010) COLONY: a program for parentage and sibship inference from multilocus
genotype data. *Mol. Ecol. Resour.* 10(3):551-555.
41. Anderson EC (2010) Computational algorithms and user-friendly software for parentage-based tagging
of Pacific salmonids. Final report submitted to the PacificSalmonCommission's Chinook Technical
Committee (US Section). p 46.
42. Henderson CR (1973) Sire evaluation and genetic trends. *J. Anim. Sci.* 1973(Symposium):10-41.
43. Sutton SG, Bult TP, & Haedrich RL (2000) Relationships among fat weight, body weight, water weight,
and condition factors in wild Atlantic salmon parr. *Trans. Am. Fish. Soc.* 129(2):527-538.
44. de Villemereuil P, Gimenez O, Doligez B, & Freckleton R (2013) Comparing parent-offspring regression
with frequentist and Bayesian animal models to estimate heritability in wild populations: a simulation
study for Gaussian and binary traits. *Methods Ecol. Evol.* 4(3):260-275.
45. Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. *J. Stat. Soft.* 33(2):1-22.
46. Butler DG, Cullis BR, Gilmour AR, & Gogel BJ (2009) *Mixed models for S language environments*
ASReml-R reference manual (Queensland Department of Primary Industries and Fisheries, NSW
Department of Primary Industries, Brisbane, Australia) p 144.
47. Hadfield JD & O'Hara RB (2015) Increasing the efficiency of MCMC for hierarchical phylogenetic models
of categorical traits using reduced mixed models. *Methods Ecol. Evol.* 6(6):706-714.