1

A processive rotary mechanism couples substrate unfolding and proteolysis in the ClpXP

degradation machinery

Zev A. Ripstein[1,2†], Siavash Vahidi[1,2,3,4†], Walid A. Houry[1,4], John L. Rubinstein[1,2,5] and Lewis E.

Kay[1,2,3,4]

[1]Department of Biochemistry, University of Toronto, Toronto, Ontario, M5S 1A8, Canada.

[2]The Hospital for Sick Children Research Institute, Toronto, Ontario, M5G 0A4, Canada.

[3]Departments of Molecular Genetics, University of Toronto, Toronto, Ontario, M5S 1A8,

Canada.

[4]Department of Chemistry, University of Toronto, Toronto, Ontario, M5S 3H6, Canada.

[5]Department of Medical Biophysics, University of Toronto, Toronto, Ontario, M5G 1L7,

Canada.

† Z.A.R and S.V contributed equally to this work
Correspondence to: kay@pound.med.utoronto.ca; john.rubinstein@utoronto.ca

Abstract

The ClpXP degradation machine consists of a hexameric AAA+ unfoldase (ClpX) and a pair of heptameric serine protease rings (ClpP) that unfold, translocate, and subsequently degrade client proteins. ClpXP is an important target for drug development against infectious diseases. Although structures are available for isolated ClpX and ClpP rings, it remains unknown how symmetry mismatched ClpX and ClpP work in tandem for processive substrate translocation into the ClpP proteolytic chamber. Here we present cryo-EM structures of the substrate-bound ClpXP complex from *Neisseria meningitidis* at 2.3 to 3.3 Å resolution. The structures allow development of a model in which the cyclical hydrolysis of ATP is coupled to concerted motions of ClpX loops that lead to directional substrate translocation and ClpX rotation relative to ClpP. Our data add to the growing body of evidence that AAA+ molecular machines generate translocating forces by a common mechanism.

Introduction

Protein degradation plays a central role in cellular physiology, regulating the timing of cell division, controlling stress responses, and ensuring the timely removal of damaged or aberrantly folded proteins (1, 2). The ClpXP system is central to protein degradation in bacteria and mitochondria and is a key regulator of cellular homeostasis, pathogenesis, and intracellular parasitism (3–5). It also serves as a model for understanding the structure and function of other ATP-dependant proteolytic systems. The ClpXP holoenzyme is composed of a hexameric AAA+ (ATPases Associated with diverse cellular Activities) unfoldase (ClpX) and a tetradecameric serine protease consisting of two heptameric rings (ClpP). Together, the ClpXP complex acts to unfold and degrade protein substrates (Figure 1A) (6). ClpXP function is critical in many bacteria, and consequently small molecules that disrupt either the protease (e.g. β-lactones (7), phenyl esters (8, 9)) or AAA+ unfoldase (e.g. ecumicin (10), lassomycin (11), rufomycin (12), dihydrothiazepines (13)), or mimic the interaction between the two (e.g. acyldepsipeptides [ADEPs] (14–16)) can kill cells, establishing ClpXP as a target for novel therapeutics against a range of different infectious diseases and cancers (17–21).

X-ray crystallographic studies of ClpP alone reveal a barrel-like structure with the protease rings stacked coaxially to form an enclosed degradation chamber that sequesters the fourteen Ser-His-Asp catalytic triads (22, 23), akin to the 20S proteasome (24, 25). Early negative-stain electron microscopy showed that one or two ClpX particles can bind co-axially to ClpP, creating a continuous central channel for substrates from ClpX to the degradation chamber of ClpP (26–28). Biochemical and X-ray crystallographic studies show that the axial entrances to the degradation chamber are gated by flexible N-terminal loops of ClpP that open upon the binding of ClpX or ADEPs, allowing substrates into the degradation chamber (29–31). Binding

of ClpX and ClpP is mediated by loops containing an Ile-Gly-Phe motif (IGF loops) on ClpX

that dock into hydrophobic pockets on the apical interfaces of ClpP subunit pairs (32–34). N-

terminal loops on ClpP also transiently interact with ClpX (33, 35, 36). However, the precise

nature of these interactions and how they allow coordination of the activities of ClpX and ClpP

remains unclear.

ClpX uses energy released by ATP hydrolysis to create a pulling force that unravels

folded protein domains for translocation into the degradation chamber of ClpP (6). This

unfolding activity relies on a series of highly conserved loops in ClpX. These loops include the

'RKH' loops that surround the ClpX entrance pore and recognize substrates for proteolysis (37),

and a pair of "pore loops" in each protomer, termed pore-1 loop (GYVG motif) and pore-2 loop

(RDV motif) (Figure 1 – figure supplement 1A) that line the axial channel of the ClpX ring (38).

Bulky aromatic and aliphatic side chains of the pore loops transmit the pulling force to the

substrate, leading to its unfolding. The mechanism by which some AAA+ ATPases convert

energy from ATP hydrolysis into a mechanical force that unfolds substrates was established

recently (39–44), and structures of ClpP and ClpX in isolation have been determined (45–48).

However, the absence of structures showing ClpXP in the process of unfolding substrate has

prevented an understanding of the conformational changes that are necessary for substrate

engagement, unfolding, and translocation in this system. ClpXP must undergo hundreds of ATP

hydrolysis events to unfold and degrade a single protein, likely via a process involving

hydrolysis of one nucleotide at a time, but the lack of structural information for a ClpXP

holoenzyme has limited understanding of how unfolding and degradation are coupled. For

example, it is not clear how the symmetry mismatch (49–51) between the pseudo-symmetric

hexameric ClpX and the seven-fold symmetric ClpP affects both the binding and interaction of

these two components.

Here we present cryo-EM structures of the ClpXP holoenzyme from the gram-negative

pathogen *Neisseria meningitidis* engaged with a protein substrate. The ClpXP portion of the

structure is at 2.3 to 3.3 Å resolution, allowing construction of a nearly complete atomic model

of the complex. Our findings show how ClpX grips and translocates substrates into the

degradation chamber through a cycle of ATP hydrolysis events involving both concerted motions

of pore loops along the substrate, as well as motions of ClpX on the apical surface of ClpP. The

data lead to a model for interactions between symmetry-mismatched ClpX and ClpP that enable

continuous degradation of substrates as ClpX rotates relative to ClpP.

Results

ATP-dependent binding and degradation of GFP-SsrA by ClpXP

ClpX and ClpP from *N. meningitidis* were expressed separately in *E. coli* before purification with

metal affinity chromatography and size exclusion chromatography (SEC). Degradation assays

using green fluorescent protein bearing an eleven-residue SsrA tag (GFP-SsrA) (43) were

performed to ensure that the heterologously expressed *N. meningitidis* (Nm) ClpX and ClpP

proteins, when mixed, have the well-established activity of the ClpXP holoenzyme. The presence

of ClpP or ClpX alone led to no loss of GFP-SsrA fluorescence relative to background bleaching

in the GFP-alone control assay. ClpX, ClpP, and ATP were all necessary to degrade GFP-SsrA

(Figure 1B), confirming that degradation of GFP-SsrA by ClpXP is ATP-dependent. As shown

below and in previous work (33, 52–54), the presence of MgATP is required for tight binding

between ClpX and ClpP. Therefore, for further structural studies, an E185Q Walker B mutant of

NmClpX (ClpX-WB) was used to slow ATP hydrolysis and prolong the lifetime of the ClpXP complex. SEC was performed to probe the oligomeric state of each component and the formation of the holoenzyme, confirming that ClpP behaves as an oligomeric species, consistent with its well-known tetradecameric architecture (Figure 1C – blue trace). ClpX-WB eluted as a hexamer in the presence of 2 mM MgATP (Figure 1C – green trace). Incubation of ClpX-WB and ClpP in the presence of MgATP resulted in a pair of higher molecular weight SEC peaks that contain both ClpX and ClpP (Figure 1C – red trace and SDS-PAGE gel insert), presumably corresponding to singly and doubly capped ClpXP complexes, showing that the *N. meningitidis* ClpX:ClpP complex can be readily reconstituted *in vitro*.

To isolate a substrate-bound holoenzyme, ClpXP was incubated with GFP-SsrA in the presence of MgATP. SEC fractions corresponding to doubly capped ClpXP bound to GFP-SsrA (Figure 1–figure supplement 1B - denoted with a *) were applied to specimen grids and vitrified for cryo-EM analysis. Visual inspection of the micrographs, as well as 2D classification of particle images, showed doubly capped complexes (Figure 1-figure supplement 2). However, 2D classification showed multiple classes where the two ClpX rings have lower density than the ClpP region, and other classes with one ClpX ring having strong density and the other having weak density. An *ab initio* 3D map showed ClpP along with strong density for one ClpX ring and only weak and fragmented density for the other (Figure 1–figure supplement 2C). Inspection of the map revealed that the ClpX ring is offset and tilted relative to the ClpP symmetry axis (Fig 1D, Video 1). The poor density for the second ClpX hexamer is likely due to a lack of correlation between binding offsets of the two ClpX rings that leads to incoherent averaging of one of the ClpX rings in the map. The map of the intact complex was refined to an overall resolution of 2.8 Å (Figure 1–figure supplement 2C), however the density for the better-resolved ClpX ring

123   remained fragmented. Further local refinement and classification in cryoSPARC (55) improved

124   the density for ClpX and revealed two distinct conformations, at 3.3 Å and 2.9 Å resolution

125   (Figure 1–figure supplement 2), hereafter referred to as Conformations A and B, respectively.

126   Focused refinement of ClpP with D7 symmetry enforced led to a map at 2.3 Å resolution (Figure

127   1–figure supplement 2).

Overall architecture of NmClpXP

Atomic models built into the cryo-EM maps showed that in both Conformations A and B, ClpX

is positioned with an offset and is tilted relative to the seven-fold symmetry axis of ClpP (Figure

1D, Video 1). Nevertheless, a continuous channel for substrate spans the ClpX and ClpP rings.

ClpP is in its active extended conformation (56), and closely resembles a published ADEP-bound

crystal structure with an all atom RMSD of 1.04 Å (56). Even though the full-length constructs

of ClpX included the zinc binding domains (residues 1-62), no density was found for these

domains in either conformation, likely due to their flexibility.

Conformations A and B of ClpX both contained additional density along their axial

channels, adjacent to the pore loops, likely from the GFP-SsrA substrate that co-eluted with

ClpXP from the SEC column (Figure 1E and 1F - orange) (Figure 1 – Figure Supplement 1B).

As the Walker B mutant has low unfolding activity (see below), the density observed likely

corresponds to the unstructured region of the degron tag. The substrate enters ClpX at an angle

of ~15° relative to the ClpP symmetry axis, and is gripped by five of the six ClpX protomers

arranged in a right-handed spiral, akin to other AAA+ ATPases in their substrate-bound

conformations (39–44). The sixth protomer forms a disengaged "seam" that bridges protomers at

the beginning and end of the spiral. In the description that follows we have labeled the protomers

X1 through X6 corresponding to their position in the spiral (Figure 1F), and consider the

configuration in which X1 is the disengaged "seam" protomer of Conformation A.

The main differences between Conformations A and B are within the protomer that

adopts the bridging seam position, and where the seam is located relative to ClpP. In

Conformation A, protomer X1 is disengaged from substrate and in the seam position (Figure 1F

left - dark blue) while in Conformation B protomer X6 is disengaged from substrate and in the

seam position (Figure 1F right - yellow). However, these two seam positions are not equivalent:

in Conformation A the seam is further from the apical surface of ClpP in what we designate as

the "upper seam" (US) position (Figure 1D,F - dark blue), while in Conformation B, the seam is

closer to ClpP in the "lower seam" (LS) position (Figure 1D,F - yellow).

The symmetry-mismatched interaction interface between ClpX and ClpP

ClpX IGF Loops: In both Conformations A and B of ClpXP, there is clear density for most of

the IGF loops that extend from the ClpX ring towards the apical surface of ClpP (Figure 2A). In

Conformation A, clear density is observed for only five of the six IGF loops, with weak density

belonging to the IGF loop of the X6 protomer (white dotted line in Figure 2B left; Figure 2-

figure supplement 1). In Conformation B, the density of the X6 IGF loop is significantly stronger

than in Conformation A, so that all six of the IGF loops are accounted for (residues 264-274,

Figure 2B, right; Figure 2C; Figure 2-figure supplement 1), with the loops binding six of the

seven pockets on ClpP. In both Conformations A and B, the empty ClpP binding pocket is

located between the X5 and X6 protomers of ClpX (Figure 2B). The lack of density for the IGF

loop belonging to protomer X6 in Conformation A (Figure 2-figure supplement 1) is likely due

to conformational flexibility and sub-stoichiometric binding of this IGF loop into the ClpP

binding pocket.

All six IGF loops in ClpX adopt different configurations in Conformations A and B

(Figure 2-figure supplement 1). The ClpX residues immediately following where the IGF loops

contact ClpP (residues 275-280) show weak or no density, likely indicating flexibility. This

property allows the IGF loops to accommodate movement of the ClpX protomers relative to

ClpP (Figure 2D-F), analogous to a set of springs. For instance, in Conformation A the X1

protomer is closer to the ClpP surface than in Conformation B and its IGF loop is thus compact

in this state. In contrast, in Conformation B the loop is fully extended, allowing the X1 protomer

to reach ~7 Å towards the top of the spiral to engage the substrate (Figure 2F; Figure 2-figure

supplement 1). In this way the flexibility of the IGF loops accommodates the spiral arrangement

of ClpX necessary to bind and unfold substrate, while maintaining contact with the planar ring of

ClpP. Additionally, the IGF loops allow the X2 protomer to move laterally away from the ClpP

pore while the X4 and X5 protomers move toward it to form a structure in which the ClpP and

ClpX rings are displaced with respect to each other (Figure 2D-red arrow; Figure 2E).

ClpP N-terminal Gates: Degradation assays (31, 35, 57–59) show that ClpX allows substrate

access to the degradation chamber of ClpP, suggesting that ClpX binding opens the N-terminal

gates of ClpP. Indeed, in both Conformations A and B the gates are in the "up" state, forming

ordered β-hairpins (Figure 2G; Figure 2-figure supplement 2). These β-hairpins closely resemble

the conformation seen in the ADEP-bound structures of NmClpP (56) and are notably different

from the disordered conformations observed in NmClpP when it is not bound to either ClpX or

ADEP (Figure 2–figure supplement 2B). By forming ordered gates, ClpX binding creates a wide

entrance pore with a diameter of ~23 Å for substrates to pass through into the ClpP degradation

chamber (Figure 2G, left; Figure 2–figure supplement 2). While it has long been suspected that

the activating mechanism of ADEPs involves a disorder-to-order transition of these gates (30),

this observation provides direct evidence that ClpX binding has the same effect.

In both Conformations A and B, the rigidification of the ClpP N-terminal gates varies

substantially between protomers. At one extreme, the N-terminal residues of the ClpP protomer

that is not bound to an IGF loop have the weakest density, appearing to possess the most

flexibility at the top of the β-hairpin (Figure 2G center; Figure 2-figure supplement 2E). In

contrast, the N-terminal residues of the ClpP protomer proximal to the X5 ClpX protomer, are

held rigidly through contacts with the ATPase (Figure 2G right). This interaction is mediated by

a pair of α-helices from protomer X5 (residues 254-262 and 291-296) adjacent to the IGF loop,

and by another X5 α-helix that precedes the pore-2 loop (residues 186-192). The three α-helices

stabilize the β-hairpin structure of the ClpP N-terminal residues by interacting with Arg17 at the

top of the hairpin (Figure 2E inset). These interactions are unique for the X5 protomer due to the

lateral displacement of the ClpX ring relative to the axial pore of ClpP (Figure 2A), with the X2

protomer shifted furthest from the pore while the X5 protomer is pulled overtop of the pore

(Figure 2D and F). Even without interacting directly with ClpX, the remaining five ClpP

protomers have nearly rigid N-terminal gates, with reduced density only for residues located in

the turn of the β-hairpin that likely reflects increased flexibility in this region (Figure 2-figure

supplement 2F). Gate rigidification therefore appears to be mediated by an allosteric effect when

IGF loops engage their respective ClpP binding pockets, akin to the effect observed upon ADEP

binding (15, 30). In the ClpXP complex with substrate bound, only a single ClpX protomer (X5)

interacts directly with a single ClpP gate at any given time. The mechanism of substrate

translocation described below involves movement of the ClpX ring on the apical surface of ClpP,

with each ClpX protomer forming contacts with a ClpP gate as translocation proceeds.

Substrate Engagement by ClpX

ClpX fits neatly into the emerging consensus for how AAA+ unfoldases engage substrate (39–43). In both Conformations A and B, five of the six pore-1 loops of ClpX interact tightly with the substrate backbone, with Tyr153 forming the majority of the interface (Figure 3A and B). These protomers generate a right-handed spiral that wraps around an 8 to 10 residue stretch of the substrate along the central channel of ClpX (Figure 3C and D). In Conformation A, the X1 pore-1 loop in the US orientation is disengaged, with the X2 protomer at the top of the spiral. As the spiral continues downwards, protomers X3, X4, and X5 make contacts with the substrate. The spiral terminates with the X6 pore-1 loop that is engaged with the substrate closest to the ClpP apical surface (Figure 3C). In Conformation B, the X6 pore-1 loop is disengaged (LS orientation), while the X1 pore-1 loop engages a new section of substrate (Figure 3D) and the X5 protomer is now located closest to the ClpP surface. When transitioning between substrate-engaged and disengaged states, the X6 pore-1 loop undergoes a large conformational change (Figure 3E). As it disengages the substrate the "lasso" fold (Figure 3E, left) that allows Tyr153 and Val154 of pore 1 to point towards the substrate is replaced by a short α-helix immediately following the GYVG motif of the loop.

Interestingly, the X1 and X5 protomers make additional contacts with the substrate in Conformation B. Whereas all of the pore-2 loops from the other five protomers are radially removed and disengaged from the substrate, the X1 protomer extends its pore-2 loop residues Ile198 and Thr199 to engage the substrate at a position three or four residues down the substrate chain from where its pore-1 loop binds (Figure 3D and F). The RKH loops, located at the top of the ClpX ring (Figure 3–figure supplement 1), are highly conserved motifs among the ClpX family of AAA+ unfoldases and are known to contribute to substrate recognition (60–62). In

Conformation B only the RKH loop of X5 directly engages substrate, with its His230 residue

interacting at a position three or four residues up the substrate chain from where its pore-1 loop

binds (Figure 3D and F). In order to contact substrate at this position the RKH loop of the X5

protomer rearranges to point toward ClpP (Figure 3F), in contrast to the other RKH loops that

point away from the axial channel of ClpX (Figure 3-figure supplement 1). These five RKH

loops are located away from the substrate, and have weaker density in the maps, likely indicating

flexibility.

Nucleotide State and Cycling in ClpX

The cryo-EM maps allowed unambiguous identification of the nucleotide bound to each ClpX

protomer. In Conformation A, there is strong density for ATP and a magnesium cofactor in

protomers X2, X3, X4, and X5, while the X1 and X6 protomers are ADP loaded, with no density

for either the γ-phosphate or magnesium (Figure 4A,4C). As in both the classic and HCLR clades

of AAA+ ATPases (39–43), the orientation between the large and the small AAA+ domains of

ClpX changes considerably depending on the nucleotide state. In Conformation A, both the X1

and X6 protomers adopt conformations that allow the X1 protomer to move away from the pore

in the US position (Figure 4-figure supplement 1) and there is reduced surface area buried

between X1 and its neighboring protomers X2 and X6. In agreement with this finding, the local

resolution in the cryo-EM maps (Figure 4-figure supplement 2) suggests that the X1 protomer in

the US position is significantly more flexible than the other protomers. In Conformation B only

the X6 protomer is ADP-bound. The X1, X2, X3, X4, and X5 protomers are ATP-bound with

clear density for the γ-phosphate and the magnesium cofactor (Fig 4B and D). Interestingly, the

X5 protomer adopts a unique orientation between its large domain and small domain compared

to the other ATP-bound protomers in order to accommodate the translation of the X6 protomer

away from the substrate translocation channel into the LS position (Figure 4B orange box; Figure

4-figure supplement 1).

In Conformation A, the β- and γ-phosphates of the bound ATP are stabilized by

interactions with the sensor-II arginine (Arg369) of the same subunit and the arginine finger

(Arg306) from the adjacent clockwise protomer (Figure 4C). In the ADP-bound sites of the X1

and X6 protomers, both arginines have moved away, allowing for a more open nucleotide

binding pocket (Figure 4C). In Conformation B both the sensor-II arginine and the arginine

finger contact the β and γ-phosphates in only four of the five ATP-bound protomers (X1, X2,

X3, and X4) (Figure 4D). In the X5 protomer (at the bottom of the spiral), the adjacent X6

protomer and its arginine finger have pivoted away from the nucleotide into the LS position,

allowing the sensor-II arginine to move closer to the γ-phosphate of the bound nucleotide. While

there is clear density for the γ-phosphate, the resolution of our experimental map was not

sufficient to differentiate between ATP and a long-lived post hydrolysis ADP/P_i state. Notably,

this key sensor II-priming motif in the nucleotide site adjacent to an ADP bound site has also

been reported for the Lon protease bound to substrate (42), a close relative in the HCLR clade of

AAA+ proteins.

The arrangement of nucleotides in our map suggests that substrate-bound ClpX

hydrolyzes ATP in a sequential manner, contrary to previous reports that ATP hydrolysis by

ClpX is a stochastic process (63). To test these models of ATP hydrolysis we quantified the GFP

degradation rates of ClpX that were doped with varying fractions, F_{WB}, of catalytically-impaired

E185Q Walker B ClpX protomers (Figure 5). The exact composition and population of resulting

complexes can be calculated using combinatorial statistics, as described previously (43, 64, 65).

If ATP hydrolysis and substrate translocation were stochastic, the activity of each protomer

would be independent of others in the ClpX hexamer. Consequently, in the stochastic model, the

expected outcome of introducing Walker B protomers into a WT ClpX hexamer would be a

linear decrease in activity as the ratio of Walker B mutants incorporated randomly into the

complex increases (Figure 5B – green line). However, GFP degradation rates drop sharply as the

Walker B ratio increases (Figure 5B – blue circles), closely following the $y = (1 - F_{WB})^6$ curve

that shows the fraction of ClpX complexes remaining fully WT at each F_{WB} mixing ratio (Figure

5B – red line). This outcome is expected only for a strictly sequential model of substrate

translocation where a single catalytically-impaired protomer blocks the unfolding activity of an

entire ClpX hexamer. It is inconsistent with the previous stochastic model that predicts that

unfolding and translocation are possible with just a single active subunit.

Discussion

In this study we used cryo-EM to build atomic models of a ClpXP complex bound to a SsrA-

tagged GFP substrate. Structures of a variety of different AAA+ rings with substrates have

emerged in the past several years (39–43), providing a description of how these ATP-dependent

molecular machines unfold substrates. In addition, methyl-TROSY NMR studies of a VAT-

substrate complex provided atomic-level insights into structural changes associated with the

substrate as it is pulled into the lumen of the unfoldase (66). Nevertheless, a detailed

understanding of how a pair of symmetry mismatched ClpP/ClpX rings can work in tandem to

unfold, translocate, and subsequently degrade substrate in a processive manner has remained

elusive. Our structural data suggest a translocation mechanism that can be explained in terms of a

pair of conformers, denoted as Conformations A and B here, that represent two steps along the

substrate translocation pathway (Figure 6A). To aid in the following discussion we have color-

coded the ClpX protomers and refer to them with their X1 - X6 labels. As we will discuss, each

of these protomers will cycle between all six protomer positions along the spiral, including,

among others, a state primed to hydrolyze ATP (denoted as ATP*) as well as the LS and US

positions. This cycling is linked to ATP hydrolysis, to the attachment position of ClpX pore-

loops along the substrate, and to the interaction of ClpX IGF loops with ClpP. We begin our

discussion of the unfolding/translocation cycle with the X1 protomer in Conformation A in the

US position, detached from substrate and bridging the lowest and highest positions of the ClpX

spiral. Both X1 and X6 protomers are ADP-bound, while the remaining protomers are in ATP-

bound states. In Conformation B, X1 moves to the highest position of the spiral with respect to

substrate (compare Figure 2F, top and bottom panels) simultaneously exchanging ADP for ATP

in its nucleotide binding pocket (Figure 6B). This transition brings X1 into close proximity with

X2 and allows engagement of X1's pore loops with substrate. The X1 pore-l loop now sits at the

top of the spiral, two substrate residues above the X2 pore loop (Figure 6A and B), thereby

translocating additional residues of substrate into the unfoldase. This transition also leads to

release of substrate by X6, which pivots away from the central pore to adopt the LS position

(second ring, Figure 6A). ATP hydrolysis and phosphate release by X5, which was in the ATP-

hydrolysis primed ATP* state, restores the ClpX spiral back to the Conformation A state, (third

ring of the four in Figure 6A), with X6 now assuming the US position. The complex is thus reset

so that it can take another step along the substrate during the next A to B transition (third ring to

fourth ring in Figure 6A). The cumulative effect of these transitions is a cycling of ClpX

protomers through different positions on the spiral (Figure 6A and B). For example, the protomer

initially in the US position (X1 in our discussion) will, with each step, move along the spiral

such that it transitions from the US position to the top of the spiral (Conformation B) and then

successively moving to lower spiral positions, through the ATP* state and finally the LS

position, as substrate is translocated. Each Conformation A➔Conformation B➔Conformation A

step results in the exchange of ADP for ATP and a single ATP hydrolysis event. Continuous

repetition of these steps leads to a processive "hand-over-hand" translocation of substrate

through the axial pore of ClpX into ClpP (Illustrated in Video 2).

The sequential hydrolysis of ATP at the ATP* position results in unidirectional substrate

translocation and movement of pore loops along the substrate (Figure 6B), with the counter

clockwise hydrolysis cycle translating into a repeated downward pulling force on the substrate.

Both the sequential hydrolysis of ATP as well as the "hand-over-hand" substrate pulling model

appear to be conserved for AAA+ unfoldases (39–43).

The cycle described above, in which each protomer eventually adopts all positions in the

spiral, has implications for how the IGF loops interact with the ClpP binding pockets. If the

spiral position of each ClpX protomer was independent of IGF binding, alignment of the

asymmetric ClpX spiral would lead to averaging of the IGF loops into all the pockets of ClpP

during cryo-EM map refinement. However, our structures clearly show an empty IGF pocket on

ClpP *only* between the ATP* and LS positions of Conformation B (Figure 2B), suggesting that

this pocket must undergo a "cycle" as well (Figure 6C). Thus, following ATP hydrolysis and

phosphate release, the IGF loop of the protomer in the ATP* position (X5 initially) must leave

its engaged pocket and move to bind the adjacent empty site on ClpP (red arrows in Figure 6C),

as the complex transitions back to Conformation A (third ring in Figure 6A). In this state, the

IGF loop appears not to have assumed a structure that allows for tight binding as only weak

density is observed for it in EM maps. In contrast, strong binding is observed for this loop in

Conformation B (Figure 4-figure supplement 1). Consistent with this model, the IGF loop of the

X5 protomer has an "extended/stretched" structure in the ATP* position of Conformation B

(Figure 2F), which likely causes strain and facilitates loop release and subsequent binding to the

corresponding site on the adjacent ClpP protomer (Figure 6C). After seven IGF loop release and

reengagement events, accompanied by seven ATP hydrolysis steps, the 'empty' ClpP binding

site makes a complete cycle, returning to its starting position (Figure 6C). The accompanying

motion of ClpX on the apical surface of ClpP for a given IGF loop transition is complex and

subtle, yet seven of these transitions lead to a net 60° rotation of ClpX with respect to ClpP (note

the positions of the purple ClpX protomer in the first and seventh panels in Figure 6D; Video 3).

The model described above is the simplest one that is consistent with our data. More complicated

models can be envisioned that require additional states for which we have no experimental data.

During preparation of this manuscript two manuscript preprints reported structures of

ClpXP from *L. monocytogenes* (67) and from *E. coli* (68), with ClpX resolutions ranging from

3.9 to 6 Å. While density maps and atomic models are not available for direct comparison with

the NmClpXP, a few important similarities and differences are notable. In all species, the ClpX

ring appears to adopt a similar offset relative to the symmetry axis of ClpP, indicating that the

general architecture of the ClpXP machinery is conserved. Notably, however, the N-terminal

apical loops of *L. monocytogenes* ClpP are not rigidified upon ClpX binding (67), in contrast to

the observation here that the gates rigidify upon binding to create a pore for substrate entry. In

the case of *L. monocytogenes* ClpXP, unusual head-to-head dimers were observed that appeared

to be mediated by the zinc binding domains. Despite the presence of the zinc binding domains in

our *N. meningitidis* construct, no such dimers were observed. While this difference may be due

to differences across species, we note that the glutaraldehyde chemical crosslinker used to

stabilize the *L. monocytogenes* ClpXP complex may have induced the formation of artefactual

dimers.

Notably, substrate engagement is different between the ClpXP from *N. meningitidis*

described here and from *E. coli* (68). In the Conformation A seen with *N. meningitidis*, there is

no engagement of substrate by any of the six pore-2 or six RKH loops, while in Conformation B

only a single RKH loop and a single pore-2 loop contact substrate (Figure 3D and F). This

observation is in contrast to the recently reported *E. coli* ClpXP structures (68) in which

arginines located in the pore-2 loop from three to five different protomers (depending on the

structure), as well as multiple RKH loops, form major contacts with substrate. The differences in

these interactions may reflect the fact that while pore-1 loop and RKH loop motifs are critical for

forming interactions with substrate, and in the case of pore-1 loops in particular, in providing the

force for translocation, additional contacts may differ between species and may be necessary for

recognition of degradation signals and for substrate specificity. The RKH loops are essential for

the recognition of SsrA-tagged substrates by *E. coli* ClpX, while human ClpX, which lacks RKH

loops but contains pore-1 loops with the same sequence as *E. coli* ClpX, fails to recognize

substrates with the SsrA-tag. Notably, however, an engineered human construct bearing the pore-

2 and RKH loops of *E. coli* ClpX is able to unfold SsrA-tagged substrates (61). Electrostatic

interactions appear to play a critical role in substrate recognition by the positively charged RKH

loop. For example, AKH and RAH mutants that reduce the positive charge of the RKH motif

weaken the affinity of *E. coli* ClpX for the negatively charged α-carboxylate of the SsrA tag

while increasing its affinity for the positively charged λO and MuA degradation tags (61).

Reordering of RKH loop residues to KHR and KRH results in significantly impaired substrate

degradation (68) and a more disrupting AAA mutant does not support ClpP degradation of a

SsrA-tagged substrate (68). Furthermore, the single point mutation RKH to RKA results in loss

of activity for both SsrA and λO tagged substrates. The tight contact between H230 and the

substrate in our maps provides a structural basis for understanding these mutagenesis results."

Although in the ClpXP structures from *E. coli* (68) the ClpX ring adopts the same spiral

staircase arrangement observed in many AAA+ ATPases (39–43, 69, 70), strongly supporting

the "hand-over-hand" mechanism of substrate translocation, the authors of this work instead

posit a stochastic mechanism (63, 68) for the HCLR clade of ATPases based on the observation

of only a single bound ADP in their structures. In the stochastic model only a single protomer

pulls the substrate, via a state that has yet to be observed. Such a model, involving stochastic

nucleotide hydrolysis, appears inconsistent with the observations here, with those for the related

Lon protease (42) where two neighbouring protomers are bound with ADP, and with a recent

structural model for a substrate-bound *E. coli* ClpAP complex showing that the unbinding and

rebinding of flexible IGL loops is coupled to substrate translocation as each protomer makes its

way from the bottom to the top of the spiral (71). The proposed mechanism in that case is similar

to the one suggested here for *N. meningitides* ClpXP. The stochastic model is also in conflict

with functional studies presented in this manuscript and for the VAT AAA+ unfoldase (43)

whereby nucleotide hydrolysis was shown to occur in a cycle that could be disrupted through the

substitution of a single hydrolysis-impaired protomer into the ATPase ring. In contrast, the

identification of two conformations in the *N. meningitidis* enzyme and observations of the bound

nucleotides in each ClpX protomer leads to a simple model of cyclic hydrolysis that, in turn,

results in unidirectional substrate translocation. In this model, the substrate is always gripped by

multiple protomers, with only a single protomer disengaged as it traverses from the bottom to the

top of the spiral. While unique features will undoubtedly continue to emerge regarding subtle

functional aspects of different AAA+ unfoldases it is likely that the translocating forces

generated by these molecular machines are based on a common mechanism.

Figure 1. Functional and structural characterization of ClpXP from *N. meningitidis*. (A) Schematic representation of
the ClpXP degradation machinery. The overall positions of the substrate (orange), ClpX unfoldase (green), and the
ClpP protease (blue) are shown; (B) GFP-SsrA degradation by ClpXP is ATP-dependant. The degradation rate of
GFP-SsrA is monitored by measuring loss of fluorescence. The components (ClpX, ClpP, ATP) included for each
measurement are denoted on the plot. All measurements included GFP-SsrA and were performed in triplicate on WT
ClpXP; (C) ClpXP complex formation monitored by size exclusion chromatography (SEC). SEC profiles of isolated
ClpP (blue trace) and ClpX (green trace) are consistent with their expected molecular weights. SEC profile of a 2:1
$ClpX_6{:}ClpP_{14}$ mixture (red trace) incubated for 10 minutes in the presence of MgATP shows the formation of
doubly- and singly-capped ClpXP complexes. The running buffer for all traces contained 2 mM MgATP. SDS-
PAGE gel of SEC fractions shows that they contain both ClpP and ClpX; (D) Overall architecture of the ClpXP
complex as established by cryo-EM. The ClpP double ring component is shown in dark blue, with the exception of a
pair of opposing protomers shown in light blue shade to delineate the IGF loop binding site that is located at the
interface between protomers. Cutaway side (E) and top (F) views of Conformations A and B of ClpX resulting from
focused classification and local refinement. The IGF loops that bind to ClpP, and the presence of substrate (orange)
in the axial channel of ClpX, are highlighted. Each of the ClpX protomers is labeled. In panel (F), the engagement of
the substrate (orange) with five of the six pore loops is indicated (red semi-circle).

439
Figure 2: The interaction interface between ClpX and ClpP. Protomer X1 occupies the US position in Conformation
A and protomer X6 the LS position in Conformation B. (A) Cutaway density map of the overall architecture of the
ClpXP interaction interface. (B) Models for the ClpP apical surface and the ClpX IGF loops (residues 265-275). The
empty IGF binding pocket (dotted oval) resides clockwise to the X5 protomer in both conformations. In
Conformation A an IGF loop was not built into the map for protomer X6 due to weak loop density (yellow dotted
line). (C) Model in map fit for an IGF loop (grey), surrounded by regions of ClpP comprising the IGF binding site
(blue). (D) Interaction of the ClpP N-terminal gates with ClpX. The X2 protomer moves away from the gates, while
the X5 protomer directly contacts the gate of the ClpP protomer to which its IGF loop is bound, shown
schematically in (E). Inset shows the details of this interaction, with model in map fit for the ClpP β-hairpin. (F)
Positions of ClpX protomers relative to ClpP. Images were generated by fitting all ClpP protomers to a common
protomer and displaying the corresponding ClpX protomer. In both conformations ClpX adopts a spiral arrangement
relative to ClpP (dotted red line). In Conformation A, the X2 protomer is located at the top of the spiral, while in
Conformation B the X1 protomer occupies the top position, ~7 Å higher than its position in Conformation A. The
ClpX protomers rotate and translate relative to ClpP; protomers X2 and X3 sit nearly atop their ClpP protomers,
while X4 and especially X5 show large deviations from this position (vertical lines), with X5 sitting overtop the
ClpP axial pore. (G) View down the channel formed by the N-terminal ClpP β-hairpins (left); density and models
for the most flexible and rigid β-hairpins (center & right, respectively).

457
Figure 3. Substrate engagement by the ClpX pore loops. (A and B) Pore-1 loop residues grip the substrate, as
observed in this view looking down the axial channel. The substrate is modelled as poly-Ala. (A) In Conformation
A, the X1 protomer is disengaged from the substrate in the US position (red arrow) and shows noticeable flexibility
in its pore-1 loop. (B) In Conformation B the X1 protomer contacts the substrate but the X6 protomer has
disengaged into the LS position (red arrows). (C and D) Model of substrate and interacting residues, along with
schematic, viewed perpendicular to the axial channel. (C) In Conformation A only the Tyr153 residue of the pore-1
loop makes significant contacts with the substrate; the 5 protomers (X2, X3, X4, X5, X6) form a downward spiral
surrounding the backbone of the substrate chain, extending to the apical surface of ClpP. (D) In Conformation B, in
addition to the five Tyr153 residues from the 5 gripping protomers (X1, X2, X3, X4, X5), more contacts are formed:
Ser197 and Ile198 from the pore-2 loop of the X1 protomer and His230 from the RKH loop of the X5 protomer. (E)
Substrate induced conformational changes of the X6 pore-1 loop upon releasing client. When the X6 protomer is
substrate engaged its pore-1 loop forms a lasso like conformation, with Tyr153 and Val154 both pointing towards
the axial channel (left). When disengaged, a part of the pore-1 loop becomes an α-helix, with Val154 and Tyr153 no
longer pointing outwards (right). (F) Architecture of the additional contacts made by the pore-2 and RKH loops
from the X1 and X5 protomers, respectively.

473
Figure 4. Nucleotide occupancy and interactions with ClpX. (A&B) Model of ClpX in conformations A and B,
looking into the ClpX pore. Nucleotides are shown and color-coded, with ATP red and ADP purple, and bind
between the large and small domains (boxed). Inset shows a schematic of the protomer positions and nucleotide
occupancies. Conformation A has two bound ADP, in the X1 (dark blue, US in Conformation A in this
representation) and X6 (yellow, LS in Conformation B in this representation) protomers, with the X1 protomer
disengaged from the substrate and away from the axial pore (US position). In Conformation B only a single ADP is
bound, protomer X1 is reengaged with substrate and X6 has disengaged. (C) Nucleotide binding sites of
Conformation A, with key interacting residues shown along with the experimental density maps corresponding to
bound nucleotide. In the ADP bound sites, both the arginine finger (R306) and the sensor-II arginine (R369) have
moved away, while in the ATP bound sites they form close contacts with the β and γ-phosphates. (D) Nucleotide
binding in Conformation B. As in Conformation A, R306 and R369 have moved away from the ADP, while in the
X5 protomer only the arginine finger has moved away, while the sensor-II transitions closer to the γ-phosphate.

486
Figure 5. A single defective ClpX protomer derails function. (A) ATPase rates for WT ClpX and the E185Q ClpX
Walker B mutant. Hydrolysis of ATP was measured with an enzyme-coupled assay (72). (B) GFP degradation rates
monitored by the decrease in fluorescence signal as a function of the fraction of added E185Q ClpX protomers to
hexameric ClpX, F_{WB} ([E185Q ClpX]/([E185Q ClpX] + [WT ClpX])). ClpX protomers bearing the slowly-
hydrolyzing E185Q Walker B mutant were mixed with WT protomers at various ratios, F_{WB}, between 5-100% of the
total protomer ClpX concentration. The activity decreases rapidly with increasing E185Q content, and closely
follows the theoretical curve $y = (1 - F_{WB})^6$ that predicts the activity profile for the case where incorporation of one
or more WB ClpX promoters is sufficient to abrogate function, as is predicted from our proposed translocation
model. This curve (red) corresponds to the expected activity if only hexamers with six WT subunits contribute to
GFP degradation while any complex with one or more E185Q mutants incorporated is functionally inactive. This
observation contrasts the expected linear decrease (green line) for a stochastic model. (C) Schematic of the
molecules produced via mixing.

499
Figure 6. Translocation model for ClpXP. (A) Schematic of the ClpX nucleotide dependent conformational cycle.
The transition from Conformation A to Conformation B is mediated through nucleotide exchange in the US position
(1^{st} to 2^{nd} ring), with ATP hydrolysis and phosphate release in the ATP* position restoring the A conformation from
the B state (2^{nd} to 3^{rd} ring) so that the cycle can repeat. In the new Conformation A (3^{rd} ring), protomer X6 (yellow)
adopts the US conformation and the X1 protomer (previous US) moves to the top of the spiral. In this way each
protomer of the complex cycles through all positions as ATP is hydrolyzes in successively new ATP* protomer
states, requiring the hydrolysis of seven ATPs. (B) Hand-over-hand mechanism of pore-loop mediated translocation.
In the first A to B transition depicted in (A) X1 (US protomer) reengages at the top of the spiral and the LS pore 1
loop (X6 in Conformation B) disengages from the bottom of the spiral. As this is repeated the pore loops move "up"
the substrate and pull it into the degradation chamber. (C) IGF loop movements accompanying the hand-over-hand
movements of ClpX. Shown in the small circles are the seven IGF loop binding sites in ClpP with a '+' sign
indicating engagement with an IGF loop of a given ClpX protomer. The binding sites are color coded according to
the color of the protomer that interacts with the corresponding site. In conformation A, the IGF loop of X6 is
unbound or only weakly bound, while in Conformation B it is bound. A new Conformation A is generated when the
IGF loop from the ATP* protomer unbinds. It subsequently reengages with ClpP in a new Conformation B (third
and fourth rings in Figure 5A). In this way the IGF loops "step" around the ClpP ring, with each complete rotation
requiring the hydrolysis of seven ATPs. (D) Schematic of the ClpX-ClpP interactions during seven ATP hydrolysis
steps that lead to a net rotation of ClpX protomers by 60°. Because of the six- to seven-fold symmetry mismatch a
total of forty-two ATPs must be hydrolyzed to return the protomers back to their original starting positions in the
cycle (*i.e.*, ring 1). Each of the seven ring structures corresponds to Conformation A, with the red arrows indicating
the subsequent step of the IGF loop of the ATP* protomer along the cycle. The US protomer in each step is
highlighted by the dashed trace.

522
523 **Figure 1 - Figure Supplement 1**. (A) NmClpX primary sequence domain map. Important structural and functional
524 motifs are denoted. (B) NmClpXP binds a SsrA-tagged GFP. SEC profiles for ClpXP incubated with GFP-SsrA (top
525 panel) and GFP-SsrA alone (bottom panel) measured using absorbance at 280 nm (black trace) and GFP
526 fluorescence (green trace - λ_{ex}:480nm, λ_{em}:508 nm). The region used for SDS PAGE analysis is highlighted by *
527 (Figure 1C).

528
529 **Figure 1 – Figure Supplement 2**. Cryo-EM image processing. (A) Representative cryo-EM micrograph of
530 NmClpXP bound to GFP-SsrA. Example complexes are circled in white. (B) Representative 2D class averages of
531 NmClpXP with GFP-SsrA. (C) Image processing workflow used to obtain final maps.
532

533
534 **Figure 1 – Figure Supplement 3.** Cryo-EM map validation. (A to C) Fourier Shell Correlation (FSC) plots (left)
535 and orientation distribution (right). Resolution values reported are for FSC=0.143.

Conformation A

X2 64-100 X4 64-100

Conformation B

X2 64-100 X4 64-100

X2 316-343 X4 316-343

X2 316-343 X4 316-343

ClpP D7 Chain M

33-58 148-163

536
537 **Figure 1 – Figure Supplement 4.** Examples of regions of the atomic models built into the experimental cryo-EM
538 maps.

Conformation A   Conformation B

539
540 **Figure 1 – Figure Supplement 5**. Experimental density maps and models for the substrate pore-1 loop interaction
541 interface.

Figure 2 - Figure Supplement 1. IGF loop flexibility. (A and B) Large domains of ClpX were aligned and models for the IGF loops are shown. (C and D) Unsharpened density map for Conformations A and B looking at protomer X6.

546
Figure 2 - Figure Supplement 2. The ClpP apical loops extend upwards from the ring surface. (A) Unsharpened
density map of the ClpP N-terminal β-hairpins when bound to ClpX. (B) Cryo-EM map of apo ClpP from *N.*
meningitidis showing that the apical loops no longer extend "up". (C) ClpX-bound and apo forms of ClpP overlaid.
There is no noticeable density for the turn of the ClpP N-terminal β-hairpin in the apo form, and the β-hairpins point
inwards, constricting the axial pore of ClpP. (D) Fourier Shell Correlation (FSC) plot for Apo-NmClpP. (E) Closeup
view of the N-terminal β-hairpins when bound to ClpX. (F) Density map of the β-hairpins in the ClpXP complex,
colored by local resolution.

554
555 **Figure 3 - Figure Supplement 1.** RKH loop positions. (A and B) Unsharpened maps for both conformations A and
556 B of ClpX are shown, with RKH loops highlighted in color (coloring done by a radius of 10 Å around residues 220-
557 240).

558
559
560 **Figure 4 - Figure Supplement 1.** Relative orientation between large and small domains of ClpX. (A and B).
561 Residues 308-343 are shown. Note that protomers X1 and X6 in Conformations A and B, respectively, are displaced
562 from substrate. These show among the largest differences in domain orientations.

563
564 **Figure 4 - Figure Supplement 2.** Local resolution maps of ClpX. (A and B) Density maps colored by local
565 resolution for Conformations A and B. In Conformation A the US protomer is highlighted with a dashed line.

566
567 **Figure 4 – Figure Supplement 3.** ATP binding pocket densities. Experimental density maps and models are shown
568 for the ATP binding pockets of all protomers. Two numerical thresholds are shown to highlight differences in
569 density.

570
Appendix Figure 1. Peptidase rate measurements of NmClpP as a function of pH, monitored using the fluorogenic
substrate PKM-AMC.

Video 1. Overview of the substrate bound ClpXP complex (Conformation B).

Video 2. Hand-over-hand translocation cycle of the ClpXP complex. Interpolation between

states A → B → A → B. Side and top views of the complex are shown (top), along with the

pore-1 loop Tyr153 and substrate (bottom left) as well as IGF loops (bottom right).

Video 3. ClpX precession on the ClpP apical surface. Repeated interpolation between states

reveals a precession of the ClpX ring as its IGF loops are placed successively in different ClpP

binding sites. The position of the empty binding pocket on ClpP undergoes a complete cycle

around the complex for every 60º rotation of the ClpX ring relative to the ClpP ring.

Supplementary File 1. Cryo-EM data acquisition, processing, model statistics, and data

deposition. (A) CryoEM data acquisition and image processing. (B) Map and Model statistics.

(C) Residues excluded in atomic models. (D) Deposited maps and associated coordinate files.

Materials and Methods:

Plasmids and constructs Codon-optimized genes encoding NmClpX (Uniprot entry: Q9JYY3)

bearing an N-terminal His_6-TEV affinity tag and ClpP (Uniprot entry: Q9JZ38) with an N-

terminal His_6-SUMO tag were synthesized by GenScript (Piscataway, NJ, USA) and cloned into

the NdeI and BamHI sites of pET28a+ (Novagen, Madison, WI, USA). Point mutations were

introduced with the Quikchange mutagenesis method (Agilent, Santa Clara, CA, USA).

Expression and purification of NmClpP and NmClpX Transformed Codon+ *E. coli* BL21(DE3)

cells were grown in LB media at 37 °C. Protein over-expression was induced by addition of 0.2

mM IPTG at OD_{600}=1.0 and was allowed to proceed overnight at 18 °C. Cells were lysed in

buffer containing 50 mM Tris, 300 KCl, 10 mM imidazole, 10% glycerol, pH 7.0 and NmClpP

and NmClpX proteins purified by Ni-affinity chromatography [HisTrap HP (GE)] in lysis buffer.

Bound proteins were eluted from the Ni column by increasing the imidazole concentration to 500

mM. The affinity tag was removed by the addition of TEV (for ClpX) or Ulp1 (for ClpP)

protease followed by dialysis against lysis/wash buffer that included 5 mM DTT. Following a

reverse Ni-affinity chromatography step, the flow-through, free from the cleaved tag and other

impurities, was concentrated with an Amicon Ultra-15 50K MWCO (Millipore) concentrator and

subjected to size exclusion chromatography (SEC) with a Superdex 200 Increase 10/300 (GE)

column in SEC buffer (50 mM imidazole, 100 mM KCl, 5 mM DTT, pH 7.0). Fractions

corresponding to NmClpX and NmClpP were pooled and stored at 4 °C in the SEC buffer until

further use. Salts containing magnesium were avoided during the purification of NmClpX to

prevent protein aggregation. Protein concentrations were determined in 8 M GdnCl using

extinction coefficient values (7450 M^{-1} cm^{-1} for ClpP, 8940 M^{-1} cm^{-1} for ClpX) determined with

ExPASy's ProtParam (73).

Expression and purification of GFP-SsrA. Green fluorescent protein (GFP) bearing an 11-

residue SsrA degradation tag at its carboxyl terminus and a non-cleavable N-terminal His×6 tag

was purified by Ni affinity chromatography followed by SEC on a HiLoad 16/60 Superdex 75 pg

column (GE).

Peptidase rate measurements of NmClpP as a function of pH. The peptidase activity of

NmClpP was measured at 37 °C with Acetyl-L-Pro-L-Lys-L-Met bearing a C-terminal

fluorogenic 7-amino-4-methylcoumarin group (PKM-AMC) as substrate. The reaction was

followed with a Synergy Neo2 96-well microplate reader making a measurement every 21

seconds for 60 minutes at λ_{ex}: 355 nm, λ_{em}: 460 nm. Each well contained 1 μM NmClpP

(monomer concentration), 250 μM PKM-AMC, 50 mM citrate, 50 mM phosphate, 50 mM Tris,

100 mM KCl in a total volume of 100 μL adjusted to the appropriate pH. Activities are derived

from initial rates extracted and analyzed using a python script written in-house. Standard errors

are calculated from repeating each reaction in triplicate.

GFP-SsrA degradation assays. Degradation of 1 μM samples of GFP-SsrA was followed by the

loss of GFP fluorescence (λ_{ex}: 480 nm, λ_{em}: 508 nm) with a Synergy Neo2 96-well microplate

reader at 25 °C. The wells included an ATP-regeneration system (74) that contained 1.5 mM

phosphoenolpyruvate, 0.2 mM NADH, 40 μg/mL pyruvate kinase, 40 μg/mL lactate

dehydrogenase, and 2 mM MgATP at pH 8.2. In some assays solutions also contained WT

NmClpX at 0.5 μM (hexamer) and/or WT ClpP at a 0.25 μM (tetradecamer), as indicated in

Figure 1B. All assays were performed in triplicate.

For the mixed protomer ClpX assays, WT ClpX and E185Q ClpX samples were purified

separately as described above (i.e. without the addition of any nucleotide) and mixed at the

indicated ratios. The mixture was left overnight for protomer exchange at room temperature

before being assayed for GFP unfolding and degradation as described. The addition of GdnCl to

promote protomer exchange is not necessary in this case because ClpX is not a stable hexamer in

the absence of nucleotide.

ATP hydrolysis assays: Steady state ATP hydrolysis was measured with an enzyme coupled

assay as described previously (74). The wells included an ATP-regeneration system that

contained 1.5 mM phosphoenolpyruvate, 0.2 mM NADH, 40 μg/mL pyruvate kinase, 40 μg/mL

lactate dehydrogenase, and 2 mM MgATP at pH 8.2. Hydrolysis of ATP was monitored by loss

of absorption at 340 nm with a Synergy Neo2 96-well microplate reader at 25 °C.

Preparation of samples for cryo-EM. A 1 mL mixture containing 10 μM (tetradecamer)

NmClpP and 20 μM NmClpX together with 200 μM GFP-SsrA was incubated with 20 mM

MgATP for 10 minutes at room temperature. This mixture was applied to a Superdex 200

Increase 10/300 (GE) column equilibrated with 50 mM bicine, 100 mM KCl, 2 mM MgATP, pH

adjusted to 8.2 at room temperature (equivalent to pH 8.5 at 4 °C – see Appendix Figure 1), as

the running buffer. Following SEC, a 0.5 mL fraction (denoted with a * in Figure 1 – Figure

Supplement 1) containing doubly capped ClpXP bound to GFP-SsrA was supplemented with 20

μM GFP-SsrA and vitrified immediately without the addition of any cross-linking agent or

detergent.

Sample vitrification. 2.5 μL of the sample mixtures were applied to nanofabricated holey gold

grids (75–77) with a hole size of ~1 μm, that had been glow discharged in air for 15 seconds.

Grids were blotted on both sides using a FEI Vitrobot mark III for 15 seconds at 4 °C and ~100%

relative humidity before freezing in a liquid ethane/propane mixture (78).

Electron Microscopy: NmClpXP was imaged with a Thermo Fisher Scientific Titan Krios G3

microscope operating at 300 kV and equipped with a FEI Falcon III DDD camera. Structures

were calculated from counting mode movies consisting of 30 frames, obtained over a 60 second

exposure with defocuses ranging from 0.9 to 1.7 μm. Movies were at a nominal magnification of

75000× corresponding to a calibrated pixel size of 1.06 Å and with an exposure of 0.8

electrons/pixel/s, giving a total exposure of 43 electrons/Å^2. 2680 movies were collected using

the microscope's *EPU* software. The Apo-NmClpP structure (with no ClpX or GFP present) was

calculated from data obtained using a FEI Tecnai F20 electron microscope operating at 200 kV

and equipped with a Gatan K2 Summit direct detector device camera. Movies consisting of 30

frames over a 15 second exposure were obtained with defocuses ranging from 1.7 to 2.9 μm.

Movies were collected in counting mode at a nominal magnification of 25000× corresponding to

a calibrated pixel size of 1.45 Å and with an exposure of 5 electrons/pixel/s, and a total exposure

of 35 electrons/Å^2. 122 movies were collected using Digital Micrograph software.

EM image analysis: Patch based whole frame alignment and exposure weighting was performed

in *cryoSPARC* v2 (55) with a 10×10 grid and the resulting averages of frames were used for

patch based contrast transfer function (CTF) determination. Templates for particle selection were

generated by 2D classification of manually selected particles. Particle images were extracted in

300×300-pixel boxes for further analysis. *Ab inito* map calculation was performed on a random

subset of 30,000 particle images, generating an initial map showing density for the complex of a

ClpP tetradecamer bound to two ClpX hexamers. A single round of 2D classification was used to

remove images of damaged particles and other contaminants from a dataset of 466,549 particle

images, with selected classes leaving 377,234 particle images for further analysis. Homogeneous

refinement of these particle images using D7 symmetry yielded a map of the complex, with good

density for only the ClpP portion, at 2.3 Å resolution.

To improve the density of the ClpX portion of the map, a round of *Ab initio* classification

was performed using three classes, of which two classes containing 289,144 particle images had

good density for ClpX, with the remaining class containing mostly density for ClpP. Refinement

of these particle images with C1 symmetry resulted in a map of the ClpXP complex at a nominal

resolution of 2.8 Å, but with poorly defined density for much of ClpX. To improve the

interpretability of the map in the ClpX region, local refinement was performed with a mask

around the six ClpX subunits without performing signal subtraction for ClpP. This refinement

greatly improved the map in the ClpX region, while blurring the density at the distal ClpP ring

(indicative of flexibility between ClpX and ClpP). However, density for two of the ClpX

subunits remained fragmented and at lower resolution. To help resolve the heterogeneity of this

region "3D variability analysis" was performed, which utilizes principle component analysis to

separate conformations. Clustering was performed along three eigenvectors. Two clusters were

identified along a single eigenvector corresponding to two conformations of ClpX bound to

substrate. From the trajectory identified, the two endpoints were used to seed a heterogeneous

classification in which the O-EM learning rate was reduced 10-fold to preserve the original

character of the seeds yielding two classes split ~ 40%:60% with 110,696 and 178,448 particle

images for Conformations A and B respectively. Subsequent non-uniform refinement yielded

maps at 3.3 Å and 2.9 Å respectively, which were then used for model building.

For Apo-NmClpP the same preprocessing steps were applied as described above, and

100,132 particle images were extracted in 160×160-pixel boxes for further analysis. Rounds of

2D classification and *ab initio* classification led to a subset of 50,400 particle images that were

used to refined a map to 4.1 Å resolution.

Atomic Model Building and Refinement: To model NmClpP, the crystal structure of NmClpP

(PDBID: 5DKP) (56) was rigidly fit with UCSF Chimera (79) into the 2.3 Å D7 symmetric map,

followed by relaxation with Rosetta using the density map as an additional term in the scoring

function (80), and utilizing D7 non crystallographic symmetry. The best scoring model was then

rigidly fit into the C1 symmetry focussed maps for Conformation A and B. Visual inspection and

real space refinement in *Coot* (81) was then used to better fit the model into the density where it

deviated from the ideal symmetry version, specifically in areas where ClpX contacted ClpP, and

the apical loops (N-terminal β-hairpins).

For NmClpX, Phyre2 (82) was used to perform one-to-one threading onto the previous

crystal structure of ClpX from *E. coli* (PDBID 3HWS chain A) (48). A single chain was then

rigidly docked into the X3 position of the 2.9 Å map, and real space refinement and *Ab initio*

model building of regions that poorly fit the density, as well as for regions missing from the

homology model was performed in *Coot* (81). This model was then relaxed with Rosetta (80),

and rigidly fit into the density for the other five ClpX protomers. Iterative rounds of real space

refinement and *Ab initio* model building in *Coot* (81), relaxation in Rosetta (80), and real space

refinement in Phenix (83), were then used to create the final model. To model Conformation A,

the model for Conformation B was used as a starting point before iterative rounds of real space

refinement and *Ab initio* model building in *Coot*, relaxation in Rosetta, and real space refinement

in Phenix. While the experimental density for the substrate showed some bulky side chains,

attempts to register the SsrA sequence in the density were unsuccessful and the substrate was

modeled as polyalanine. Models were evaluated with Molprobity (84) and EMRinger (85).

Figures and movies were generated in UCSF Chimera (79) and UCSF ChimeraX (86), and colors

chosen with ColorBrewer (87).

Acknowledgements

We thank Dr. Samir Benlekbir for cryo-EM data collection on the Titan Krios. ZAR and SV

were supported by a scholarship and a postdoctoral fellowship from the Canadian Institutes of

Health Research, respectively. LEK and JLR were supported by the Canada Research Chairs

program. This research was funded by Canadian Institutes of Health Research grants FDN-

503573 (LEK), PJT-162186 (JLR), and PJT-148564 (WAH). Titan Krios cryo-EM data were

collected at the Toronto High-Resolution High-Throughput cryo-EM facility supported by the

Canadian Foundation for Innovation and Ontario Research Fund.

References:

1.   A. O. Olivares, T. A. Baker, R. T. Sauer, Mechanistic insights into bacterial AAA+

proteases and protein-remodelling machines. *Nat. Rev. Microbiol.* **14**, 33–44 (2016).

2.   A. L. Goldberg, Protein degradation and protection against misfolded or damaged

proteins. *Nature* **426**, 895–899 (2003).

3.   D. Frees, U. Gerth, H. Ingmer, Clp chaperones and proteases are central in stress survival,

virulence and antibiotic resistance of Staphylococcus aureus. *Int. J. Med. Microbiol.* **304**,

142–149 (2014).

4.   C. M. Pickart, R. E. Cohen, Proteasomes and their kin: proteases in the machine age. *Nat.*

Rev. Mol. Cell Biol. **5**, 177–187 (2004).

5.   V. Bhandari, K. S. Wong, J. L. Zhou, M. F. Mabanglo, R. A. Batey, W. A. Houry, The

Role of ClpP Protease in Bacterial Pathogenesis and Human Diseases. *ACS Chem. Biol.*

13, 1413–1425 (2018).

6.   T. A. Baker, R. T. Sauer, ClpXP, an ATP-powered unfolding and protein-degradation

machine. *Biochim. Biophys. Acta* **1823**, 15–28 (2012).

7.   T. Böttcher, S. A. Sieber, β-Lactones as Privileged Structures for the Active-Site Labeling

of Versatile Bacterial Enzyme Classes. *Angew. Chem. Int. Ed.* **47**, 4600–4603 (2008).

8.   M. W. Hackl, M. Lakemeyer, M. Dahmen, M. Glaser, A. Pahl, K. Lorenz-Baath, T.

Menzel, S. Sievers, T. Böttcher, I. Antes, H. Waldmann, S. A. Sieber, Phenyl Esters Are

Potent Inhibitors of Caseinolytic Protease P and Reveal a Stereogenic Switch for

Deoligomerization. *J. Am. Chem. Soc.* **137**, 8475–8483 (2015).

9.   M. Lakemeyer, E. Bertosin, F. Möller, D. Balogh, R. Strasser, H. Dietz, S. A. Sieber,

Tailored peptide phenyl esters block ClpXP proteolysis by an unusual breakdown into a

heptamer-hexamer assembly. *Angew. Chemie Int. Ed.* **0** (2019).

10. W. Gao, *et al.*, The Cyclic Peptide Ecumicin Targeting ClpC1 Is Active

againstMycobacterium tuberculosis In Vivo. *Antimicrob. Agents Chemother.* **59**, 880

(2015).

11. E. Gavrish, *et al.*, Lassomycin, a Ribosomally Synthesized Cyclic Peptide, Kills

Mycobacterium tuberculosis by Targeting the ATP-Dependent Protease ClpC1P1P2.

Chem. Biol. **21**, 509–518 (2014).

12. M. P. Choules, *et al.*, Rufomycin Targets ClpC1 Proteolysis in Mycobacterium

tuberculosis and M. abscessus. *Antimicrob. Agents Chemother.* **63**, e02204-18 (2019).

13. C. Fetzer, V. S. Korotkov, R. Thänert, K. M. Lee, M. Neuenschwander, J. P. von Kries, E.

Medina, S. A. Sieber, A Chemical Disruptor of the ClpX Chaperone Complex Attenuates

the Virulence of Multidrug-Resistant Staphylococcus aureus. *Angew. Chemie Int. Ed.* **56**,

15746–15750 (2017).

14. J. Kirstein, A. Hoffmann, H. Lilie, R. Schmidt, H. Rübsamen-Waigmann, H. Brötz-

Oesterhelt, A. Mogk, K. Turgay, The antibiotic ADEP reprogrammes ClpP, switching it

from a regulated to an uncontrolled protease. *EMBO Mol. Med.* **1**, 37–49 (2009).

15. M. Gersch, K. Famulla, M. Dahmen, C. Göbl, I. Malik, K. Richter, V. S. Korotkov, P.

Sass, H. Rübsamen-Schaeff, T. Madl, H. Brötz-Oesterhelt, S. A. Sieber, AAA+

chaperones and acyldepsipeptides activate the ClpP protease via conformational control.

Nat. Commun. **6**, 6320 (2015).

16. K. S. Wong, *et al.*, Acyldepsipeptide Analogs Dysregulate Human Mitochondrial ClpP

Protease Activity and Cause Apoptotic Cell Death. *Cell Chem. Biol.* **25**, 1017–1030.e9

(2018).

17. E. Culp, G. D. Wright, Bacterial proteases, untapped antimicrobial drug targets. *J.*

Antibiot. (Tokyo). **70**, 366–377 (2017).

18. I. T. Malik, H. Brotz-Oesterhelt, Conformational control of the bacterial Clp protease by

natural product antibiotics. *Nat. Prod. Rep* **34**, 815–831 (2017).

19. R. M. Raju, A. L. Goldberg, E. J. Rubin, Bacterial proteolytic complexes as therapeutic

targets. *Nat. Rev. Drug Discov.* **11**, 777–789 (2012).

20. C. A. Goard, A. D. Schimmer, Mitochondrial matrix proteases as novel therapeutic targets

in malignancy. *Oncogene* **33**, 2690 (2013).

21. K. S. Wong, W. A. Houry, "Recent Advances in Targeting Human Mitochondrial AAA+

Proteases to Develop Novel Cancer Therapeutics" in *Mitochondria in Health and in*

Sickness, A. Urbani, M. Babu, Eds. (Springer Singapore, 2019), pp. 119–142.

22. J. Wang, J. A. Hartling, J. M. Flanagan, The Structure of ClpP at 2.3 A° Resolution

Suggests a Model for ATP-Dependent Proteolysis. *Cell* **91**, 447–456 (1997).

23. K. Liu, A. Ologbenla, W. A. Houry, Dynamics of the ClpP serine protease: A model for

self-compartmentalized proteases. *Crit. Rev. Biochem. Mol. Biol.* **49**, 400–412 (2014).

24. F. Förster, P. Unverdorben, P. Śledź, W. Baumeister, Unveiling the Long-Held Secrets of

the 26S Proteasome. *Structure* **21**, 1551–1562 (2013).

25. E. Kish-Trier, C. P. Hill, Structural Biology of the Proteasome. *Annu. Rev. Biophys.* **42**,

29–49 (2013).

26. J. Ortega, H. S. Lee, M. R. Maurizi, A. C. Steven, Alternating translocation of protein

substrates from both ends of ClpXP protease. *EMBO J.* **21**, 4938–4949 (2002).

27. J. Ortega, S. K. Singh, T. Ishikawa, M. R. Maurizi, A. C. Steven, Visualization of

Substrate Binding and Translocation by the ATP-Dependent Protease, ClpXP. *Mol. Cell* **6**,

1515–1521 (2000).

28.   M. Kessel, M. R. Maurizi, B. Kim, E. Kocsis, B. L. Trus, S. K. Singh, A. C. Steven,

803   Homology in Structural Organization Between E. coli ClpAP Protease and the Eukaryotic

804   26 S Proteasome. *J. Mol. Biol.* **250**, 587–594 (1995).

805   29.   M. E. Lee, T. A. Baker, R. T. Sauer, Control of Substrate Gating and Translocation into

806   ClpP by Channel Residues and ClpX Binding. *J. Mol. Biol.* **399**, 707–718 (2010).

807   30.   D. H. S. Li, Y. S. Chung, M. Gloyd, E. Joseph, R. Ghirlando, G. D. Wright, Y. Q. Cheng,

808   M. R. Maurizi, A. Guarne, J. Ortega, Acyldepsipeptide Antibiotics Induce the Formation

809   of a Structured Axial Channel in ClpP: A Model for the ClpX/ClpA-Bound State of ClpP.

810   *Chem. Biol.* **17**, 959–969 (2010).

811   31.   G. Effantin, M. R. Maurizi, A. C. Steven, Binding of the ClpA Unfoldase Opens the Axial

812   Gate of ClpP Peptidase. *J. Biol. Chem.* **285**, 14834–14840 (2010).

813   32.   Y.-I. Kim, I. Levchenko, K. Fraczkowska, R. V Woodruff, R. T. Sauer, T. A. Baker,

814   Molecular determinants of complex formation between Clp/Hsp100 ATPases and the

815   ClpP peptidase. *Nat. Struct. Biol.* **8**, 230–233 (2001).

816   33.   S. A. Joshi, G. L. Hersch, T. A. Baker, R. T. Sauer, Communication between ClpX and

817   ClpP during substrate processing and degradation. *Nat Struct Mol Biol* **11**, 404–411

818   (2004).

819   34.   A. J. Amor, K. R. Schmitz, T. A. Baker, R. T. Sauer, Roles of the ClpX IGF loops in ClpP

820   association, dissociation, and protein degradation. *Protein Sci.* **28**, 756–765 (2019).

821   35.   A. Gribun, M. S. Kimber, R. Ching, R. Sprangers, K. M. Fiebig, W. A. Houry, The ClpP

822   Double Ring Tetradecameric Protease Exhibits Plastic Ring-Ring Interactions, and the N

823   Termini of Its Subunits Form Flexible Loops That Are Essential for ClpXP and ClpAP

824   Complex Formation. *J. Bio. Chem.* **280**, 16185–16196 (2005).

825   36.   L. D. Jennings, J. Bohon, M. R. Chance, S. Licht, The ClpP N-Terminus Coordinates

826   Substrate Access with Protease Active Site Reactivity. *Biochemistry* **47**, 11031–11040

827   (2008).

828   37.   A. Martin, T. A. Baker, R. T. Sauer, Diverse Pore Loops of the AAA+ ClpX Machine

829   Mediate Unassisted and Adaptor-Dependent Recognition of ssrA-Tagged Substrates. *Mol.*

830   *Cell* **29**, 441–450 (2008).

831   38.   A. Martin, T. A. Baker, R. T. Sauer, Pore loops of the AAA+ ClpX machine grip

832   substrates to drive translocation and unfolding. *Nat Struct Mol Biol* **15**, 1147–1151 (2008).

833   39.   N. Monroe, H. Han, P. S. Shen, W. I. Sundquist, C. P. Hill, Structural basis of protein

834   translocation by the Vps4-Vta1 AAA ATPase. *Elife* **6** (2017).

835   40.   H. Yu, T. J. Lupoli, A. Kovach, X. Meng, G. Zhao, C. F. Nathan, H. Li, ATP hydrolysis-

836   coupled peptide translocation mechanism of Mycobacterium tuberculosis ClpB. *Proc.*

837   *Natl. Acad. Sci. U. S. A.* **115**, E9560–E9569 (2018).

838   41.   A. H. De la Peña, E. A. Goodall, S. N. Gates, G. C. Lander, A. Martin, Substrate-engaged

839   26S proteasome structures reveal mechanisms for ATP-hydrolysis–driven translocation.

840   *Science (80-.).* **362**, eaav0725 (2018).

841   42.   M. Shin, A. Asmita, C. Puchades, E. Adjei, R. L. Wiseman, A. W. Karzai, G. C. Lander,

842   Distinct Structural Features of the Lon Protease Drive Conserved Hand-over-Hand

843   Substrate Translocation. *bioRxiv*, 617159 (2019).

844   43.   Z. A. Ripstein, R. Huang, R. Augustyniak, L. E. Kay, J. L. Rubinstein, Structure of a

845   AAA+ unfoldase in the process of unfolding substrate. *Elife* **6**, e25754 (2017).

846   44.   S. N. Gates, *et al.*, Ratchet-like polypeptide translocation mechanism of the AAA +

847   disaggregase Hsp104. **1052**, 1–13 (2017).

45. M. Li, O. Kandror, T. Akopian, P. Dharkar, A. Wlodawer, M. R. Maurizi, A. L. Goldberg,

Structure and Functional Properties of the Active Form of the Proteolytic Complex,

ClpP1P2, from Mycobacterium tuberculosis. *J. Biol. Chem.* **291**, 7465–7476 (2016).

46. S. R. Geiger, T. Bottcher, S. A. Sieber, P. Cramer, A Conformational Switch Underlies

ClpP Protease Function. *Angew. Chem. Int. Ed.* **50**, 5749–5752 (2011).

47. J. Ishizawa, *et al.*, Mitochondrial ClpP-Mediated Proteolysis Induces Selective Cancer

Cell Lethality. *Cancer Cell* (2019)

https:/doi.org/https://doi.org/10.1016/j.ccell.2019.03.014.

48. S. E. Glynn, A. Martin, A. R. Nager, T. A. Baker, R. T. Sauer, Structures of Asymmetric

ClpX Hexamers Reveal Nucleotide-Dependent Motions in a AAA+ Protein-Unfolding

Machine. *Cell* **139**, 744–756 (2007).

49. M. C. Bewley, V. Graziano, K. Griffin, J. M. Flanagan, The asymmetry in the mature

amino-terminus of ClpP facilitates a local symmetry match in ClpAP and ClpXP

complexes. *J. Struct. Biol.* **153**, 113–128 (2006).

50. F. Beuron, M. R. Maurizi, D. M. Belnap, E. Kocsis, F. P. Booy, M. Kessel, A. C. Steven,

At Sixes and Sevens: Characterization of the Symmetry Mismatch of the ClpAP

Chaperone-Assisted Protease. *J. Mol. Biol.* **123**, 248–259 (1998).

51. P. Majumder, T. Rudack, F. Beck, R. Danev, G. Pfeifer, I. Nagy, W. Baumeister, Cryo-

EM structures of the archaeal PAN-proteasome reveal an around-the-ring ATPase cycle.

Proc. Natl. Acad. Sci. **116**, 534 LP-539 (2019).

52. G. L. Hersch, R. E. Burton, D. N. Bolon, T. A. Baker, R. T. Sauer, Asymmetric

Interactions of ATP with the AAA+ ClpX6 Unfoldase: Allosteric Control of a Protein

Machine. *Cell* **121**, 1017–1027 (2005).

53.   R. Grimaud, M. Kessel, F. Beuron, A. C. Steven, M. R. Maurizi, Enzymatic and Structural

Similarities between theEscherichia coli ATP-dependent Proteases, ClpXP and ClpAP. *J.*

Biol. Chem. **273**, 12476–12481 (1998).

54.   J. M. Jones, D. J. Welty, H. Nakai, Versatile Action of Escherichia coli ClpXP as Protease

or Molecular Chaperone for Bacteriophage Mu Transposition. *J. Biol. Chem.* **273**, 459–

465 (1998).

55.   A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for

rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–+ (2017).

56.   J. D. Goodreid, J. Janetzko, J. P. Santa Maria, K. S. Wong, E. Leung, B. T. Eger, S.

Bryson, E. F. Pai, S. D. Gray-Owen, S. Walker, W. A. Houry, R. A. Batey, Development

and Characterization of Potent Cyclic Acyldepsipeptide Analogues with Increased

Antimicrobial Activity. *J. Med. Chem.* **59**, 624–646 (2016).

57.   A. Martin, T. A. Baker, R. T. Sauer, Distinct static and dynamic interactions control

ATPase-Peptidase communication in a AAA plus protease. *Mol. Cell* **27**, 41–52 (2007).

58.   J. Ortega, H. S. Lee, M. R. Maurizi, A. C. Steven, ClpA and ClpX ATPases bind

simultaneously to opposite ends of ClpP peptidase to form active hybrid complexes. *J.*

Struct. Biol. **146**, 217–226 (2004).

59.   S. K. Singh, R. Grimaud, J. R. Hoskins, S. Wickner, M. R. Maurizi, Unfolding and

internalization of proteins by the ATP-dependent proteases ClpXP and ClpAP. *Proc. Natl.*

Acad. Sci. **97**, 8898 LP-8903 (2000).

60.   A. Martin, T. A. Baker, R. T. Sauer, Distinct Static and Dynamic Interactions Control

ATPase-Peptidase Communication in a AAA+ Protease. *Mol. Cell* **27**, 41–52 (2007).

61.   C. M. Farrell, T. A. Baker, R. T. Sauer, Altered Specificity of a AAA+ Protease. *Mol. Cell*

25, 161–166 (2007).

62.   S. M. Siddiqui, R. T. Sauer, T. A. Baker, Role of the processing pore of the ClpX AAA+

ATPase in the recognition and engagement of specific protein substrates. *Genes Dev.* **18**,

369–374 (2004).

63.   A. Martin, T. A. Baker, R. T. Sauer, Rebuilt AAA + motors reveal operating principles for

ATP-fuelled machines. *Nature* **437**, 1115–1120 (2005).

64.   S. Vahidi, Z. A. Ripstein, M. Bonomi, T. Yuwen, M. F. Mabanglo, J. B. Juravsky, K.

Rizzolo, A. Velyvis, W. A. Houry, M. Vendruscolo, J. L. Rubinstein, L. E. Kay,

Reversible inhibition of the ClpP protease via an N-terminal conformational switch. *Proc.*

Natl. Acad. Sci. **115**, E6447--E6456 (2018).

65.   R. Huang, F. Perez, L. E. Kay, Probing the cooperativity of Thermoplasma acidophilum

proteasome core particle gating by NMR spectroscopy. *Proc. Natl. Acad. Sci. U.S.A.* **114**,

E9846–E9854 (2017).

66.   R. Augustyniak, L. E. Kay, Cotranslocational processing of the protein substrate

calmodulin by an AAA+ unfoldase occurs via unfolding and refolding intermediates.

Proc. Natl. Acad. Sci. **115**, E4786 LP-E4795 (2018).

67.   C. Gatsogiannis, D. Balogh, F. Merino, S. A. Sieber, S. Raunser, Cryo-EM structure of the

ClpXP protein degradation machinery. *bioRxiv*, 638692 (2019).

68.   X. Fei, T. A. Bell, S. Jenni, B. M. Stinson, T. A. Baker, S. C. Harrison, R. T. Sauer,

Structures of the ATP-fueled ClpXP proteolytic machine bound to protein substrate.

bioRxiv, 704999 (2019).

69.   E. J. Enemark, L. Joshua-Tor, Mechanism of DNA translocation in a replicative

hexameric helicase. *Nature* **442**, 270–275 (2006).

70. R. Huang, Z. a Ripstein, R. Augustyniak, M. Lazniewski, K. Ginalski, L. E. Kay, J. L.

Rubinstein, Unfolding the mechanism of the AAA+ unfoldase VAT by a combined cryo-

EM, solution NMR study. *Proc. Natl. Acad. Sci. U. S. A.* (2016)

https:/doi.org/10.1073/pnas.1603980113.

71. K. E. Lopez, A. N. Rizo, E. Tse, J. Lin, N. W. Scull, A. C. Thwin, A. L. Lucius, J.

Shorter, D. R. Southworth, Conformational Plasticity of the ClpAP AAA+ Protease

Couples Protein Unfolding and Proteolysis. *bioRxiv* (2019) https:/doi.org/10.1101/820209.

72. J. G. Nørby, Coupled assay of Na+,K+-ATPase activity. *Methods Enzymol.* **156**, 116–9

(1988).

73. E. Gasteiger, C. Hoogland, A. Gattiker, S. Duvaud, M. R. Wilkins, R. D. Appel, A.

Bairoch, "Protein Identification and Analysis Tools on the ExPASy Server" in *The*

Proteomics Protocols Handbook, J. M. Walker, Ed. (Humana Press, 2005), pp. 571–607.

74. J. G. B. T.-M. in E. Nørby, "[11] Coupled assay of Na+,K+-ATPase activity" in

Biomembranes Part P: ATP-Driven Pumps and Related Transport: The Na,K-Pump,

(Academic Press, 1988), pp. 116–119.

75. C. R. Marr, S. Benlekbir, J. L. Rubinstein, Fabrication of carbon films with ~500nm holes

for cryo-EM with a direct detector device. *J. Struct. Biol.* **185**, 42–47 (2014).

76. C. J. Russo, L. A. Passmore, Ultrastable gold subtrates for electron cryomicroscopy.

Science (80-.). **346**, 1377–1380 (2014).

77. J. R. Meyerson, P. Rao, J. Kumar, S. Chittori, S. Banerjee, J. Pierson, M. L. Mayer, S.

Subramaniam, Self-assembled monolayers improve protein distribution on holey carbon

cryo-EM supports. *Sci. Rep.* **4**, 7084 (2014).

78. W. F. Tivol, A. Briegel, G. J. Jensen, An improved cryogen for plunge freezing. *Microsc.*

Microanal. **14**, 375–379 (2008).

79.   E. F. Pettersen, T. D. Goddard, C. C. Huang, G. S. Couch, D. M. Greenblatt, E. C. Meng,

T. E. Ferrin, UCSF Chimera—A visualization system for exploratory research and

analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

80.   R. Y. R. Wang, M. Kudryashev, X. Li, E. H. Egelman, M. Basler, Y. Cheng, D. Baker, F.

Dimaio, De novo protein structure determination from near-atomic-resolution cryo-EM

maps. *Nat. Methods* **12**, 335–338 (2015).

81.   P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta*

Crystallogr. Sect. D **60**, 2126–2132 (2004).

82.   L. A. Kelley, S. Mezulis, C. M. Yates, M. N. Wass, M. J. E. Sternberg, The Phyre2 web

portal for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845 (2015).

83.   P. D. Adams, *et al.*, PHENIX: a comprehensive Python-based system for macromolecular

structure solution. *Acta Crystallogr. Sect. D* **66**, 213–221 (2010).

84.   V. B. Chen, W. B. Arendall III, J. J. Headd, D. A. Keedy, R. M. Immormino, G. J. Kapral,

L. W. Murray, J. S. Richardson, D. C. Richardson, MolProbity: all-atom structure

validation for macromolecular crystallography. *Acta Crystallogr. Sect. D* **66**, 12–21

(2010).

85.   B. A. Barad, N. Echols, R. Y.-R. Wang, Y. Cheng, F. DiMaio, P. D. Adams, J. S. Fraser,

EMRinger: side chain–directed model and map validation for 3D cryo-electron

microscopy. *Nat. Methods* **12**, 943 (2015).

86.   T. D. Goddard, C. C. Huang, E. C. Meng, E. F. Pettersen, G. S. Couch, J. H. Morris, T. E.

Ferrin, UCSF ChimeraX: Meeting modern challenges in visualization and analysis.

Protein Sci. **27**, 14–25 (2018).

87. M. Harrower, C. A. Brewer, ColorBrewer.org: An Online Tool for Selecting Colour

Schemes for Maps. *Cartogr. J.* **40**, 27–37 (2003).