1   Transposable elements in individual genotypes of *Drosophila simulans*

6   Sarah Signor[1]

7   [1]Department of Biological Sciences, North Dakota State University

8   sarah.signor@ndsu.edu

Key words: transposable elements, *Drosophila simulans*, inbred lines, copy number, genetic

variation

Abstract

Transposable elements are mobile DNA sequences that are able to copy themselves within a

host's genome. Within insects they often make up a substantial proportion of the genome. While

they are the subject of intense research, often times when copy number is estimated it is

estimated only at the population level, or in a limited number of individuals within a population.

I was interested in whether or not transposable elements accumulate at different rates in different

genetic backgrounds – the variance between individual genotypes in transposable element

accumulation. Using two populations of *Drosophila simulans* from California and Africa I

estimated transposable element copy number in individual genotypes. Some active transposable

elements seem to be a property of the species, while others of the populations. I find that in

addition to population level differences in transposable element load certain genotypes

accumulate transposable elements at a much higher rate than others. Most likely active

transposable elements are fairly rare, and were inherited only by specific genotypes that were

used to create the inbred lines. Whether or not this reflects dynamics in natural populations,

where transposable elements may accumulate in specific genotypes and maintain themselves in

the population rather than being active at low levels population wide, is an open question.

Introduction

Transposable elements are mobile DNA sequences that are able to move around the genome, potentially increasing their copy number in the process. They are abundant and contribute substantially to genetic variation in populations, influencing genome evolution (Biémont *et al.*, 1999; Bennetzen, 2000; Feschotte, 2008). The number and location of transposable element insertions can vary substantially between species, populations, and individuals (Vieira *et al.*, 1999; Vieira & Biémont, 2004; C Vieira, 2008; Kofler & Schlötterer, 2015; Kofler *et al.*, 2015b; Jakšić *et al.*, 2017; Kofler *et al.*, 2018). Introgression of transposable elements between species is also common, and has been reported frequently (Kofler *et al.*, 2015a; Hill & Betancourt, 2018). In addition, they make up a substantial proportion of the genome in many invertebrates and plants (Bennetzen, 2000; Tenaillon *et al.*, 2010; Kofler *et al.*, 2015b). Transposable elements are divided into three categories: long terminal repeat retrotransposons, short interspersed nuclear element retrotransposons, and transposons. The latter moves through DNA excision and repair, while the former two transpose via an RNA intermediate.

The most well studied transposable element population in insects is that of *Drosophila melanogaster*. It has been reported that in *D. melanogaster* total copy number of transposable elements may be similar between individuals, but the location of insertions can vary considerably (Barrón *et al.*, 2014). Overall copy number is thought to be maintained by transposition-selection balance, though new transposable elements may initially have high transposition rates as the host machinery evolves new defenses (Yang & Nuzhdin, 2003; Pasyukova, 2004; Slotkin & Martienssen, 2007; Johnson, 2010; Lee & Langley, 2012; Romero-Soriano & Garcia Guerreiro, 2016). Transposable elements also accumulate in heterochromatic regions, potentially because these regions are thought to be 'transcriptionally silent', and thus the deleterious effects of their

insertion are mitigated (Dimitri, 2003). However, they can also contribute to variation in

quantitative traits, differences in fitness, and changes in gene expression (Mackay 1984; Mackay

1989, Shrimpton et al. 1990, Mackay et al. 1992, Long et al. 2000) . Some transposable elements

have been associated with increases in fitness due to changes in gene regulation (Aminetzach *et*

al. 2005; Mateo *et al.*, 2014)).

The recent explosion of population level genomic data has greatly increased our ability to

estimate transposable element load in populations. Much of this work on transposable element

copy number has occurred in *Drosophila*, where the existence of multiple sequenced inbred

panels lend themselves to estimating copy number and insertion site frequency . However,

differences in the number of transposable element insertions are thought to accumulate in inbred

laboratory lines because active copies of transposable elements are rare and will be inherited

only in some lines (Nuzhdin *et al.*, 1997). This is potentially mirroring what occurs in natural

populations, where transposable element site heterogeneity may potentially be due to frequent

transpositions in rare flies rather than low levels of transposition population-wide (Nuzhdin,

2000). Many studies of transposable element insertions compare populations which have been in

the laboratory for different amounts of time, wild flies and laboratory flies, and they estimate

population-level insertion rate using Pool-seq. While Pool-seq is an effective tool for estimating

population-level frequency, understanding the dynamics of transposable elements in inbred lines

requires the ability to estimate the variance between genotypes in transposable element copy

number, as well as the repeatability of transposable element load between populations. Here we

use previously sequenced lines of *D. simulans* from two populations to directly estimate

transposable element copy number in individual genotypes.

One population was collected in Africa in 2002 and 2006 and I do not know how long

they were held in the lab prior to being sequenced, however six of the stocks were sequenced

from originally collected material (wild). A second Californian population was held in the lab for

two years prior to sequencing while they were being inbred. I will look at variance in

transposable element copy number between inbred genotypes, differences between the wild and

inbred lines, and differences between the populations in the mean and variance of transposable

element copy number.

Methods

Fly lines

The 22 African *D. simulans* isofemale lines were collected by William Ballard in 2002 from

Madagascar and Peter Andolfatto in 2006 from Kenya (Jackson *et al.*, 2017). They were inbred

in the lab for nine generations, during which time six were lost and DNA was collected from the

initial stocks. The raw reads are 90 bp paired end illumina sequencing and they were downloaded

from SRA PRJEB7673 (Jackson *et al.*, 2017). The first read from each pair was used for

mapping. The California lines were collected from the Zuma Organic Orchard in Los Angeles,

CA on two consecutive weekends of February 2012, from a pile of rotting strawberries (Signor *et*

al., 2017; Signor & Nuzhdin, 2018; 2019). Reads were single end 100 bp and this project has

been deposited at the SRA under accession SRP075682.

Mapping and copy number estimation

We mapped the reads using bwa mem version 0.7.15 to the *D. simulans* 2.02 assembly and the

179 consensus transposable element sequences from EMBL, downloaded from Flybase.org (Li

2015). Bam files were sorted and index with samtools v.1.9 and optical duplicates were removed

using picard MarkDuplicates (http://picard.sourceforge.net) (Li *et al.*, 2009; McKenna *et al.*,

2010). Reads with a mapping quality of below 15 were removed (this removes reads which map

equally well to more than one location). Using read coverage to determine copy number has been

compared to other methods and is neither permissive nor conservative (Srivastav & Kelleher,

2017). We therefore estimated transposable element copy number per family by comparing

average coverage of the transposable sequence to average coverage of chromosome arm 2L in R,

and computed population averages and individual copy number (Hill *et al.*, 2015). P-values of

comparisons between means and variances were corrected for multiple testing using Bonferroni

correction.

SNPs and summary statistics

We called SNPs within the transposable elements and the genomes using GATK Haplotypecaller

(McKenna *et al.*, 2010). SNPs were filtered for a minimum depth of four. SNPs were not filtered

for missing calls given that not at all individuals will share insertions. Tajima's *D* was estimated

in windows of 1 kb using VCFtools, and prior to estimation indels and SNPs with more than two

alleles were removed (Danecek *et al.*, 2011). Tajima's D was not intended to estimate selection

on the transposable elements, but rather to understand if some transposable element insertions

were more recent than others, indicated by the frequency and number of SNPs.

Results and Discussion

Population-level variation

Of the 128 elements examined in the population, 85 have different mean numbers of insertions

between the two populations (t-test, Bonferroni corrected p = .05/128). Of those, only 17 are

higher in the African populations, suggesting that overall the CA population has more

transposable element insertion sites. Indeed, overall Californian *D. simulans* have an average of

1,797 insertions per genotype, while African *D. simulans* have 1,496. The five elements with the

largest difference in copy number in Californian *D. simulans* compared to African are the *INE-1*,

Tc1, *transib2*, *1360*, and *Cr1a*. These are present on average in 37 more copies in Californian *D.*

simulans. The elements more common in Africa include *baggins*, *X-element*, *R1A1-element*,

diver2, *Rt1c*, *Rt1b*, *jockey*, *ninja*, *Osvaldo*, *gypsy*, *Bari1*, *gypsy8*, *Tirant*, *mariner*, *BS3*, *G5A*, and

aurora-element. *G5A*, *gypsy8*, *X-element*, *baggins*, *R1A1-element*, *copia*, *G7*, *invader6*, *diver*,

hobo, *invader3*, *Rt1b*, *Rt1c*, *Bari1*, and *Tirant* also have significantly different (and higher)

variance compared to Californian *D. simulans* (F-test, Bonferroni corrected p = 0.05/128). 20

elements have different and larger variance in the Californian *D. simulans* compared to African

(F-test, Bonferroni corrected p = 0.05/128). This includes *Stalker4*, *Tc3*, *frogger*, *F-element*, *412*,

accord, *G4*, *rover*, *17.6*, *HMS-Beagle*, *gypsy6*. The most abundant transposable elements in the

California population were the highly abundant *INE-1*, *Cr1a*, *G6*, *1360*, *transib2*, *Tc1*, *baggins*,

and *roo*. In the African populations this is similar, *INE-1*, *Cr1a*, *G6*, *baggins*, *R1A1-element*,

diver2, *X-element*, and *HB*, followed by *1316* and *roo*.

Stalker4, Stalker, Bari2, Tc3, G7, and *Tart-C* were never present as more than a fraction

of an element in any individual, and in general coverage was limited to small regions within

these elements. This suggests that if these elements are present they are old and degraded. *G3*

and *hopper2* are estimated as being present in ~1 copy per individual in both populations,

however closer inspection reveals that all reads are mapping to the start and end of the elements.

This suggests that the only copy or copies present have internal deletions. For the *G-element* all

but a small fraction of reads map to one 140 bp sequence, but they map abundantly to that

particular sequence, such that the copy number is estimated as close to 6 per individual. Again,

reads which map equally well to more than one location were filtered out, thus this does not

represent non-specific mapping to repetitive elements. These patterns are not seen in the other

transposable elements, where coverage is even across the length of the element (Supplementary

Figures 1,2,4,5 as examples)

The *D. melanogaster pogo* and *Helitron* elements were not present in these populations,

which has been previously noted, suggesting that these transposable elements are not present in

D. simulans (Kofler *et al.*, 2015b). A full-length version of *Quasimodo* (two copies) and *gypsy6*

(one copy) were present in one individual. In other individuals reads map to very specific small

regions of the transposable elements, suggesting old and degraded copies. *Stalker3* is also

present in one genotype as a full-length copy. There is no mapping in the African populations

and very little in Californian populations, suggesting that old or degraded copies are not present

in the rest of the genotypes sampled.

Tajima's *D* of transposable elements is not intended as a measure of selection, given that

the reads are coming from different places in the genome. However, a large negative Tajima's *D*

might indicate recent activity of a single copy of the transposable element, as the copies have

more low frequency SNPs. The most negative Tajima's D in California belonged to *G6*, *TAHRE*,

Juan, *gypsy7*, *flea*, *Bari1* and *frogger* (less than -1.5). In African *D. simulans* this includes

Tabor, *Transpac*, *flea*, *Juan*, *Bari1*, *G6*, and *accord*. Thus in both populations *G6*, *flea*, *Bari1*,

and *Juan* likely have recent activity (Supplementary Figure 1 & 2). This is consistent with other

work on *Juan*, which suggests it is actively transposing in the species (Kofler *et al.*, 2015b). Note

that African *D. simulans* have a largely neutral genomic Tajima's *D* when split between the two

populations or considered together, while Californian *D. simulans* is elevated genome-wide

((Signor *et al.*, 2017), Supplementary Figure 3). An elevated Tajima's *D* could indicate a number

of things, such as older invasions or multiple, diverged active elements in the populations. The

majority of differences in mean and variance in the population are due to transposable elements

with a positive Tajima's *D*, suggesting that these differences are due to different rates of gain and

loss of abundant transposable elements over time or low level spread of multiple diverged

copies.

The p-element

The *p-element* recently invaded *D. simulans* from *D. melanogaster* as described in (Kofler *et al.*,

2015a), however Pool-seq cannot tie *p-element* insertions to specific individuals, only determine

the average number of insertions. What was reported previously was 0.4 insertions in Florida

populations, and 29 in South Africa (Kofler *et al.*, 2015a). What we see in the California

population is an average of 2 insertions, however that is because the majority of individuals do

not have any insertions (137 individuals have less than .3 estimated copies, Figure 1). The

remaining individuals have between .5 and 39 copies. We recovered the single reported SNP in

the *p-element*, a G to A transition at position 2040. It is interesting that it is not invading

genotypes in the population at the same rate, but rather reaching high copy number in some

genotypes and not others (Nuzhdin 2000). However, given the observed 'outliers' (see next

section), one genotype for a given transposable element with high copy number, this may be

reflective of how some transposable elements invade populations. It is also possible that *p-*

elements are just proliferating in strains that were infected prior to collection (Nuzhdin &

Mackay 1997). This was observed previously in lab strains of *D. melanogaster*, though

contamination and introgression may also have played a role (Rahman *et al.* 2015).

Transposable elements in individual genomes

Some transposable elements have considerably higher copy number in particular genotypes

compared to the population average. For example, in one genotype *Dsim\ninja* is present in 29

copies, compared to the population mean of three. *Dsim\ninja* has 10 fixed differences and 27

polymorphisms in this strain from the California population, and the population average is 7.5

fixed differences and 264 polymorphisms (Supplementary Figure 4 & 5). This suggests that

Dsim\ninja was recently active in this genome. This is true of several transposable elements

which have outliers in the population. *Stalker2* has an outlier with 17 fixed SNPs and 8

polymorphisms, compared to a population average of 14 fixed SNPs and 43 polymorphisms. The

gypsy10 element has a single outlier individual with 203 fixed differences compared to *D.*

melanogaster, and 77 polymorphisms. The population mean is 158 fixed differences and 222

polymorphisms. Other transposable elements with large outliers in the California population

include *opus*, *blood*, *GATE*, *diver*, *Tabor*, *INE-1*, *diver2*, *idefix*, *1731*, *412*, and *297*.

Sampling of the African populations was much more limited thus less genotype-specific

variation is sampled, and indeed only two transposable elements had large outliers, both in the

same individual from Madagascar: *copia*, and *diver*. This individual had 11 fixed differences in

copia, and 20 polymorphisms, while the population average is 10 fixed differences and 78

polymorphisms (and 20 copies compared to 4-11 for the rest of the population). For *diver* this

individual had 20 fixed differences and 84 polymorphisms, compared to a population average of

12 fixed differences and 215 polymorphisms (and 30 copies compared to 4-10 for the rest of the

population). The outlier individual is one that was inbred in the lab. In general being inbred in

the lab is not affecting overall transposable element copy number however, as comparing

between lines that were sequenced directly upon collection and those that there inbred, there is

no significant difference between the mean number of transposable elements for any

transposable element family. All of this is to suggest that transposable element load varies

significantly between genotypes within a population, though which transposable elements are

able to 'wake up' and multiply may be shared between genotypes in different populations

(*diver*). That is to say that some recent transposable element activity is shared across the species

(*G6*), populations (*p-element*) and some is specific to genotypes within a population. Those that

'wake up' in individual lines appears to be due to sampling of individuals that are permissive or

contain active transposable elements, rather than an overall increase in transposable element

activity in inbred lines.

Comparison to other studies

Previously *HMS Beagle*, *blood*, *flea*, *opus*, *Stalker*, and *F-element* were described as being

present in 0-1 copies in most populations from a large survey, with some populations having a

significant number of insertions (Biémont *et al.*, 1999). In our populations all of these elements

with the exception of *Stalker* were present in on average more than six insertions. Of these only

flea has more low frequency polymorphisms suggesting that it has spread within the genome

recently (CA, Tajima's D -1.7). (Vieira *et al.*, 1999) detected *17.6* at one insertion site in *D.*

simulans, and here it was present on average in 1 (CA) to 0.5 (AF) copies in these populations,

which is more frequent than recent papers such as (Kofler *et al.*, 2015b) which detected four

copies in the population of ~800 which was sampled. (Vieira *et al.*, 1998) estimated 0-30 copies

of *412* and 20-100 copies of *roo* per genome including heterochromatin. We observed 5-42

copies of *412* in both populations, with a single outlier at 73. For *roo* we observed 20-75 copies

in both populations. Thus our estimates are consistent with previous estimates for these elements

that include heterochromatin. It has also been suggested that *412* is recently active (Biémont *et*

al., 1999).

Tirant has previously been reported as having higher copy number in African *D.*

simulans, potentially due to a recent mobilization of the element (Fablet *et al.*, 2006). We find

that pattern here, including a higher variance in the African populations where copy number

ranges from 2 to 6.68, compared to 2-3.8 in California (Fablet *et al.*, 2006). Viera et al. 1999

sampled diverse *D. simulans* populations and found ~1 *Tirant* element per genome, though this

was limited to euchromatic regions and older *Tirant* insertions may potentially have accumulated

in heterochromatic regions. It has previously been suggested than *Dmau\mariner* activity is

higher in populations that have spread out of Africa, though here the *Dmau\mariner* element has

a higher copy number in Africa than in the Californian *D. simulans*, from 0-5 with an average of

2.33, compared to 0-3 with an average of 1.22 (Picot *et al.* 2008). For *mariner2* we find copy

number to be relatively stable in both populations (9.6 in California, 7.3 in Africa), while in

Kofler et al. 2015 *mariner2* is only reported in six copies in the entire population of ~800,

though again this is limited to euchromatic regions.

The *G6* element has a large difference from previously reported values, with an average

of 66 insertions in Californian *D. simulans* and 69 in African. However only 37 insertions where

reported total for a previously estimated population of ~800 *D. simulans* isofemale lines (Kofler

et al., 2015b). Only 18% of reads in Kofler et al. (2015) map to euchromatin, but this is not a

large enough fraction to explain the difference. In addition, in the populations reported here the

G6 element has primarily low frequency polymorphisms (Tajima's D = -2.4), suggesting that this

is a recent expansion of copy number. The population examined in Kofler et al. (2015) is from

South African in 2013, a later collection date than either population here, though it is possible

that the *G6* activation is geographically limited to more northern populations. Overall our

estimates are higher than the work of Kofler et al. 2015, which only estimates more than one

insertion per line for four transposable elements (*1360, hobo*, *roo* and *Tc-2*). This is conservative

compared to other estimates of euchromatic insertions, for example Vieira et al. 1999 use *in situ*

hybridization to estimate copy number and found more than one insertion on average for

nineteen elements (of 36 measured, *412, blood, burdock, copia, coral, flea, gypsy, HMS beagle,*

mdg3, opus, roo, Tirant, dor, F-element, Helena, I-element, jockey, Bari1, and *hobo*). There is

other evidence this work may be conservative, for example (Kaminker *et al.*, 2002) estimated

146 *roo* insertions in the euchromatin of the *D. melanogaster* reference genome, while (Kofler *et*

al., 2015b) estimates 1966 total for a population of 554 (approximately 3.6 per individual). This

may also be due to inherent variability between populations.

Conclusions

Transposable element load is an attribute of species, populations, and individual genotypes.

Elements which were not active in most of the population wake up in particular genotypes and

spread rapidly, other elements reach high abundance across populations but were not noted in

previous studies as being particularly prominent (*G6*). In inbred laboratory lines active

transposable element copies may be inherited by some lines and not others, and active

transposable elements can accumulate over time (Nuzhdin *et al.*, 1997). This can cause

differences over time in the number of insertions within a line and large differences between

lines in transposable element copy number (Nuzhdin *et al.*, 1997). This may also be reflective of

natural patterns in which transposable elements proliferate in particular genotypes rather than at

low levels in the population as a whole (Nuzhdin, 2000). Now that the Californian *D. simulans*

lines have been held in the laboratory for eight years it would be interesting to see how the

genotype-specific patterns of transposable element insertions have continued to change. Overall,

looking at variance between individuals is an important part of understanding the ways in which

transposable elements maintain themselves in populations.

Acknowledgements: I would like to thank C. and S. Emery for helpful commentary on the

manuscript. I am also thankful to J. Butler and T. Robert for help in the laboratory.

The author declares that she has no conflicting interests.

Figure 1: Estimated copy number for the *p-element* and *G6* in Californian and African *D.*

simulans. Each bar represents an individual from the population. As expected, the *p-element* was

not found in the African population sampled in the early 2000's, but by 2012 when the

Californian *D. simulans* was sampled it had invaded. *G6* has a high copy number in both

populations of *D. simulans*, which wasn't recorded in previous studies on African *D. simulans*.

Figure 2: Estimated copy number for the two non-melanogaster transposable elements included

here, Dsim\ninja and Dmau\mariner. Variance between individual genomes is large for

transposable elements, for example in California Dsim\ninja varies from 1-29, and in Africa

from 2-12. The individual with 29 copies of Dsim\ninja has 10 fixed SNPs within the element

and 27 polymorphisms compared to other population members (for example, 239 polymorphisms

and 5 fixed SNPs), suggesting that it has been recently active within that genome.

Element	CA Average	Variance	AF Average	Variance	Element	CA Average	Variance	AF Average	Variance	Element	CA Average	Variance	AF Average	Variance
297	18.44	7.87	12.11	8.31	GATE	26.98	12.73	28.32	13.07	NOF	1.04	0.21	0.97	0.32
412	27.78	31.66	14.77	5.39	gtwin	4.85	0.73	4.78	0.87	opus	7.64	5.03	6.85	9.54
1360	64.03	37.76	35.07	20.84	gypsy	4.97	1.06	5.92	2.02	Osvaldo	5.44	0.85	6.96	1.38
1731	6.69	6.86	7.10	3.65	gypsy10	8.37	5.13	6.34	3.94	P-element	2.36	44.92	-	-
17.6.	1.10	0.13	0.50	0.00	gypsy11	3.73	0.29	2.42	0.08	pogo	-	-	-	-
3S18	11.89	5.66	12.59	4.38	gypsy12	3.97	0.75	4.51	0.86	Porto1	10.02	0.65	7.49	0.39
accord	6.05	10.35	2.54	1.17	gypsy2	0.95	0.02	0.81	0.01	Q-element	0.95	0.03	0.72	0.01
accord2	4.84	0.81	3.23	0.41	gypsy3	3.45	0.95	3.08	0.73	Quasimodo	0.19	0.03	0.08	0.00
aurora-element	0.70	0.02	0.89	0.02	gypsy4	2.07	0.04	1.98	0.08	R1-2	1.13	0.05	1.21	0.11
baggins	52.65	15.51	62.04	42.68	gypsy5	3.68	1.12	2.26	0.52	R1A1-element	34.07	72.24	49.39	301.81
Bari1	4.92	4.58	18.05	30.07	gypsy6	0.39	0.01	0.31	0.00	R2-element	39.52	65.11	30.05	74.21
Bari2	0.12	0.00	0.11	0.00	gypsy7	0.86	0.01	0.64	0.00	roo	50.89	66.75	33.38	43.96
blood	12.12	11.30	7.98	10.57	gypsy8	4.77	0.69	6.38	1.95	rooA	19.29	1.30	18.61	1.19
BS	2.93	0.27	2.94	0.17	gypsy9	3.14	0.24	2.46	0.23	rover	1.24	0.15	0.58	0.03
BS3	6.41	0.63	6.95	0.51	HB	43.04	20.11	40.52	32.24	Rt1a	3.61	0.26	3.91	0.38
BS4	3.31	0.13	2.93	0.07	Helena	29.20	17.08	22.24	10.19	Rt1b	17.43	6.41	24.71	8.67
Burdock	12.73	4.35	9.27	3.04	Helitron	-	-	-	-	Rt1c	20.60	14.16	27.78	41.37
Circe	6.56	1.79	5.88	2.80	HeT-A	6.52	4.70	1.50	1.59	S-element	1.41	0.06	0.83	0.05
copia	9.22	4.00	6.52	11.85	HMS-Beagle	13.44	2.59	10.49	2.17	S2	0.73	0.01	0.57	0.00
Cr1a	125.69	88.15	97.02	39.36	HMS-Beagle2	0.82	0.01	0.69	0.00	springer	2.10	0.10	2.11	0.09
diver	6.75	8.19	8.14	26.95	hobo	32.87	12.41	31.79	32.23	Stalker	0.28	0.00	0.19	0.00
diver2	35.58	21.62	44.05	32.27	hopper	23.58	16.70	15.88	11.24	Stalker2	2.86	3.49	2.31	1.14
Dm88	8.43	0.45	8.34	0.34	hopper2	0.77	0.06	0.66	0.05	Stalker3	0.08	0.03	-	-
Dmau\mariner	1.22	0.72	2.33	1.54	I-element	37.33	35.49	21.37	13.78	Stalker4	0.09	0.00	0.06	0.00
Doc	25.20	24.77	23.59	37.24	Idefix	3.70	1.91	2.64	0.57	Tabor	9.95	44.94	5.81	3.25
Doc2-element	30.43	6.53	28.56	8.46	INE-1	193.45	296.57	116.27	104.78	TAHRE	3.09	1.10	1.01	1.35
Doc3-element	20.57	1.17	19.21	1.48	invader1	4.62	0.11	4.55	0.18	TART-A	1.05	0.24	0.54	0.09
Doc4-element	3.68	0.25	2.99	0.07	invader2	6.09	1.50	4.29	0.92	TART-B	0.94	0.20	0.36	0.15
Dsim\ninja	3.14	4.99	4.93	5.85	invader3	6.71	1.93	6.47	5.80	TART-C	0.02	0.00	0.01	0.00
F-element	28.41	11.23	19.78	1.48	invader4	3.24	0.08	3.07	0.06	Tc1	54.48	36.25	31.21	10.81
FB	22.44	20.74	13.75	14.15	invader5	0.66	0.01	0.56	0.00	Tc1-2	12.45	2.47	8.46	1.85
flea	13.71	10.09	10.85	13.78	invader6	5.85	1.55	6.20	3.67	Tc3	0.15	0.00	0.06	0.00
frogger	1.05	0.06	0.89	0.01	Ivk	14.98	2.26	12.87	2.72	Tirant	2.77	0.06	3.58	2.17
Fw2	6.50	0.20	5.64	0.14	jockey	8.73	3.10	11.61	4.33	Tom1	26.03	6.41	17.00	7.94
Fw3	3.01	0.07	2.89	0.16	jockey2	5.55	0.15	4.68	0.04	transib1	19.12	8.84	12.41	4.99
G-element	4.22	0.93	4.08	1.26	Juan	21.46	27.13	18.68	13.75	transib2	55.12	44.06	27.48	39.43
G2	25.99	15.12	23.93	16.78	looper1	2.24	0.11	1.56	0.07	transib3	2.81	0.07	2.07	0.07
G3	0.91	0.01	0.76	0.01	mariner2	9.61	0.54	7.34	0.28	transib4	4.14	0.17	2.80	0.16
G4	4.10	0.23	3.52	0.06	Max-element	29.78	17.50	32.43	15.97	Transpac	8.21	4.81	5.35	4.09
G5	15.05	2.32	15.99	2.77	McClintock	13.49	8.00	6.86	2.81	X-element	34.10	8.98	41.49	23.77
G5A	10.93	0.66	11.67	2.70	mdg1	6.43	1.00	4.95	0.85	Xanthias	9.43	3.14	8.90	6.90
G6	66.62	309.64	69.36	532.88	mdg3	7.02	4.73	5.94	3.01	ZAM	2.92	0.12	2.01	0.23
G7	0.18	0.00	0.20	0.01	micropia	8.01	4.05	6.53	2.71					

 Table 1: Summary statistics for copy number of transposable elements from the Californian and

 African *D. simulans*.

Aminetzach YT, Macpherson JM, Petrov DA. 2005. Pesticide resistance via transposition-

mediated adaptive gene truncation in *Drosophila*. *Science* **309**:764–7.

Barrón, M.G., Fiston-Lavier, A.-S., Petrov, D.A. & González, J. 2014. Population genomics of
transposable elements in *Drosophila*. *Annu. Rev. Genet.* **48**: 561–581.

Bennetzen, J.L. 2000. Transposable element contributions to plant gene and genome evolution.
Plant Mol. Biol. **42**: 251–269.

Biémont, C., Vieira, C., Borie, N. & Lepetit, D. 1999. Transposable elements and genome
evolution: the case of *Drosophila simulans*. *Genetica* **107**: 113–120.

C Vieira, C.B. 2008. Selection against transposable elements in *D. simulans* and *D.*
melanogaster. *Genet. Res.* **68**: 9-115

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., *et al.* 2011. The
variant call format and VCFtools. *Bioinformatics* **27**: 2156–2158.

Dimitri, P. 2003. Colonization of heterochromatic genes by transposable elements in *Drosophila*.
Mol. Biol. Evol. **20**: 503–512.

Fablet, M., McDonald, J.F., Biémont, C. & Vieira, C. 2006. Ongoing loss of the *tirant*
transposable element in natural populations of *Drosophila simulans*. *Gene* **375**: 54–62.

Feschotte, C. 2008. Transposable elements and the evolution of regulatory networks. *Nat. Rev.*
Genet. **9**: 397–405.

Hill, T. & Betancourt, A.J. 2018. Extensive exchange of transposable elements in the *Drosophila*
pseudoobscura group. 1–14. Mobile DNA.

Hill, T., Schlötterer, C. & Betancourt, A. 2015. Hybrid dysgenesis in *Drosophila simulans* due to
a rapid global invasion of the P- element. *PLoS* **12**: e1005920.
349

Jackson, B.C., Campos, J.L., Haddrill, P.R., Charlesworth, B. & Zeng, K. 2017. Variation in the
intensity of selection on codon bias over time causes contrasting patterns of base
composition evolution in *Drosophila*. *Genome Biol. Evol.* evw291–22.

Jakšić, A.M., Kofler, R. & Schlötterer, C. 2017. Regulation of transposable elements: interplay
between TE-encoded regulatory sequences and host-specific trans-acting factors in
Drosophila melanogaster. *Mol. Ecol.* **26**: 5149-5159.

Johnson, N.A. 2010. Hybrid incompatibility genes: remnants of a genomic battlefield? *Trends*
Genet. **26**: 317–325.

Kaminker, J.S., Bergman, C.M., Kronmiller, B., Carlson, J., Svirskas, R., Patel, S., *et al.* 2002.
The transposable elements of the *Drosophila melanogaster* euchromatin: a genomics
perspective. *Genome Biology* **3**: research0084.1.
361

362 Kofler, R. & Schlötterer, C. 2015. Low levels of transposable element activity in *Drosophila*
363   *mauritiana*: causes and consequences. *bioRxiv*.

Kofler, R., Hill, T., Nolte, V., Betancourt, A.J. & Schlötterer, C. 2015a. The recent invasion of natural *Drosophila simulans* populations by the *P-element*. *Proc. Nat. Acad. Sci. USA* **112**: 6659–6663.

Kofler, R., Nolte, V. & Schlötterer, C. 2015b. Tempo and mode of transposable element activity in *Drosophila. PLoS Genet.* **11**: e1005406.

Kofler, R., Senti, K.-A., Nolte, V., Tobler, R. & Schlötterer, C. 2018. Molecular dissection of a natural transposable element invasion. *Genome Res.* **28**: 824–835.

Lee, Y.C.G. & Langley, C.H. 2012. Long-term and short-term evolutionary impacts of transposable elements on *Drosophila. Genetics* **192**: 1411–1432.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., *et al.* 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Li, H. 2015. Aligning sequence reads, clone sequences and assemble contigs with BWA-MEM *arXiv* 1–3.

Long, A.D., Lyman, R.F., Morgan, A.H., Langley, C.H., Mackay, T.F. 2000. Both naturally occurring insertions of transposable elements and intermediate frequency polymorphisms at the *achaete-scute* complex are associated with variation in bristle number in *Drosophila melanogaster. Genetics* **154**:1255–69.

Mackay, T.F.C. 1984. Jumping genes meet abdominal bristles: hybrid dysgenesis-induced quantitative variation in *Drosophila melanogaster. Genet. Res.* **44**: 231-237.

Mackay, T.F.C. 1989. Transposable elements and fitness in *Drosophila melanogaster. Genome* **31**: 284-295.

Mackay, T.F., Lyman, R.F., Jackson, M.S., 1992. Effects of P element insertions on quantitative traits in *Drosophila melanogaster. Genetics* **130**: 315-332.

Picot ,S., Wallau, G.L., Loreto, E.L.S., Heredia, F.O., Hua-Van A., Capy P. 2008. The *mariner* transposable element in natural populations of *Drosophila simulans. Heredity* **101**: 53–9.

Rahman R., Chirn, G.-W., Kanodia, A., Sytnikova, Y.A., Brembs, B., Bergman, C.M., *et al.* 2015. Unique transposon landscapes are pervasive across *Drosophila melanogaster. Nucleic Acids Res.* **43**:10655–72.

Mateo, L., Ullastres, A. & González, J. 2014. A Transposable element insertion confers xenobiotic resistance in *Drosophila. PLoS Genet.* **10**: e1004560.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., *et al.* 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**: 1297–1303.

Nuzhdin, S.V. 2000. Sure facts, speculations, and open questions about the evolution of transposable element copy number. In: *Transposable Elements and Genome Evolution* (J. F. McDonald, ed), pp. 129–137. Springer Netherlands, Dordrecht.

Nuzhdin, S.V., Pasyukova, E.G. & Mackay, T.F. 1997. Accumulation of transposable elements in laboratory lines of *Drosophila melanogaster*. *Genetica* **100**: 167–175.

Pasyukova, E.G. 2004. Accumulation of transposable elements in the genome of *Drosophila melanogaster* is associated with a decrease in fitness. *Journal of Heredity* **95**: 284–290.

Romero-Soriano, V. & Garcia Guerreiro, M.P. 2016. Expression of the retrotransposon *Helena* reveals a complex pattern of TE deregulation in *Drosophila* hybrids. *PLoS ONE* **11**: e0147903.

Signor, S. & Nuzhdin, S. 2018. Dynamic changes in gene expression and alternative splicing mediate the response to acute alcohol exposure in *Drosophila melanogaster*. *Heredity* **121:** 342-360.

Signor, S.A. & Nuzhdin, S.V. 2019. Evolution of phenotypic plasticity in response to ethanol between sister species with different ecological histories (*Drosophila melanogaster and D. simulans*). *bioRxiv*.

Signor, S.A., New, F.N. & Nuzhdin, S. 2017. A large panel of *Drosophila simulans* reveals an abundance of common variants. *Genome Biol. Evol.* **10**: 189–206.

Shrimpton, A.E., Mackay, T.F., Brown, A.J. 1990. Transposable element-induced response to artificial selection in *Drosophila melanogaster*: molecular analysis of selection lines. *Genetics* **125**:803–11.

Slotkin, R.K. & Martienssen, R. 2007. Transposable elements and the epigenetic regulation of the genome. *Nat. Rev. Genet.* **8**: 272–285.

Srivastav, S.P. & Kelleher, E.S. 2017. Paternal induction of hybrid dysgenesis in *Drosophila melanogaster* is weakly correlated with both *P-element* and *hobo* element dosage. *G3* **7**: 1487–1497.

Tenaillon, M.I., Hollister, J.D. & Gaut, B.S. 2010. A triptych of the evolution of plant transposable elements. *Trends Plant Sci.* **15**: 471–478.

Vieira, C. & Biémont, C. 2004. Transposable element dynamics in two sibling species: *Drosophila melanogaster* and *Drosophila simulans*. *Genetica* **120**: 115–123.

Vieira, C., Aubry, P., Lepetit, D. & Biémont, C. 1998. A temperature cline in copy number for *412* but not *roo*/B104 retrotransposons in populations of *Drosophila simulans*. *Proc. R. Soc. B* **265**: 1161–1165.

Vieira, C., Lepetit, D., Dumont, S. & Biémont, C. 1999. Wake up of transposable elements following *Drosophila simulans* worldwide colonization. *Mol. Biol. Evol.* **16**: 1251–1255.

437   Yang, H.-P. & Nuzhdin, S.V. 2003. Fitness costs of *Doc* expression are insufficient to stabilize
438   its copy number in *Drosophila melanogaster. Mol. Biol. Evol.* **20**: 800–804.

439