A conserved expression signature predicts growth rate and reveals cell & lineage-specific differences

Zhisheng Jiang*[3], Serena Francesca Generoso*[1], Marta Badia[2], Bernhard Payer[†,1,2], Lucas B. Carey[†,2,3]

[1] Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Dr. Aiguader 88, 08003, Barcelona, Spain.
[2] Universitat Pompeu Fabra (UPF), Barcelona, Spain.
[3] Center for Quantitative Biology and Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China

[†]Correspondence: bernhard.payer@crg.eu and lucas.carey@pku.edu.cn
*These authors contributed equally to the work

Synopsis

By performing RNA-seq on cells FACS sorted by their proliferation rate, this study identifies a gene expression signature capable of predicting proliferation rates in diverse eukaryotic cell types and species. This signature, applied to scRNAseq data from *C.elegans*, reveals lineage-specific differences in proliferation during development. In contrast to the universality of the proliferation signature, mitochondria and metabolism related genes show a high degree of cell-type specificity; mouse pluripotent stem cells (mESCs) and differentiated cells (fibroblasts) exhibit opposite relations between mitochondria state and proliferation. Furthermore, we identified a slow proliferating subpopulation of mESCs with higher expression of pluripotency genes. Finally, we show that fast and slow proliferating subpopulations are differentially sensitive to mitochondria inhibitory drugs in different cell types.

Highlights

1. A FACS-based method to determine the transcriptomes of fast and slow proliferating subpopulations.

2. A universal proliferation-correlated transcriptional signature indicates high protein synthesis and degradation in fast proliferating cells across cell types and species.

3. Applied to scRNA-seq, the expression signature predicts correctly the global slowdown in proliferation during *C. elegans* development, with lineage-specific exceptions.

4. Mitochondria membrane potential predicts proliferation rate in a cell-type specific manner, with ETC complex III inhibitor having distinct effects on the proliferation of fibroblasts vs mESCs.

Abstract:

Isogenic cells cultured together show heterogeneity in their proliferation rate. To determine the differences between fast and slow-proliferating cells, we developed a method to sort cells by proliferation rate, and performed RNA-seq on slow and fast proliferating subpopulations of pluripotent mouse embryonic stem cells (mESCs) and mouse fibroblasts. We found that slowly proliferating mESCs have a more naïve pluripotent character. We identified an evolutionarily conserved proliferation-correlated transcriptomic signature that is common to all eukaryotes: fast cells have higher expression of genes for protein synthesis and protein degradation. This signature accurately predicted growth rate in yeast and cancer cells, and identified lineage-specific proliferation dynamics during development, using *C. elegans* scRNA-seq data. In contrast, sorting by mitochondria membrane potential revealed a highly cell-type specific mitochondria-state related transcriptome. mESCs with hyperpolarized mitochondria are fast proliferating, while the opposite is true for fibroblasts. The mitochondrial electron transport chain inhibitor antimycin affected slow and fast subpopulations differently. While a major transcriptional-signature associated with cell-to-cell heterogeneity in proliferation is conserved, the metabolic and energetic dependency of cell proliferation is cell-type specific.

Introduction

Rates of cell growth and division vary greatly, even among isogenic cells of a single cell-type, cultured in the same optimal environment[1]. Cell-to-cell heterogeneity in proliferation rate has important consequences for population survival in bacterial antibiotic resistance, stress resistance in budding yeast, and chemo-resistance in cancer[2-10]. Time-lapse fluorescence microscopy has shown that cell-to-cell variability in the expression of some genes, such as *p53* and *p21*, is associated with cell-to-cell variability in proliferation and survival[1, 11]. While microscopy can identify dynamic relationships between gene expression and cell fate, it is limited to measurements of one or two genes per cell. Single-cell RNA sequencing measures transcriptome-level heterogeneity but does not directly link this to cell-biological heterogeneity in organelle state, or dynamic heterogeneity in proliferation or drug resistance. Transcriptome-level approaches for understanding within-population cell-to-cell heterogeneity in proliferation and other dynamic processes are lacking. While the presence of intrapopulation variation in proliferation, transcriptome, and organelle-state in both steady-state and in differentiation populations is well established, the relationship among the three remains unclear.

One possibility is that the proliferation-correlated gene expression program is the same, regardless of if one looks at interpopulation variation due to genetic or environmental differences, or intrapopulation heterogeneity due to epigenetic differences and expression noise. However, in the budding yeast *Saccharomyces cerevisiae,* the expression program of intrapopulation heterogeneity in proliferation rate only partially resembles that of cells growing at different rates due to genetic or environmental perturbations[8]. The relation between gene expression and proliferation rate is much less well studied in mammalian cells.

In yeast, in tumors, and in organs, genetic, environmental and developmental changes
cause changes in proliferation rate, and changes in the expression of hundreds or thousands
of genes[12-16]. Unsurprisingly, many of the genes for which changes in expression are
associated with changes in proliferation rate are associated with adverse clinical outcomes in
cancer and with antibiotic and antifungal resistance[17, 18]. Within a population of microbes,
and within a single multicellular organism, the correct balance of proliferation states and
rates is essential. Yet measuring this heterogeneity is difficult, and without such data,
understanding the consequences of this heterogeneity is impossible.

Gene expression is associated with phenotype, but mRNAs themselves do not often
directly cause phenotypes. Instead, they serve as markers for cell-biological differences
between cells. Phenotypes are mostly driven by larger cell-biological differences between
cells, such as differences in metabolic state. Cell-to-cell heterogeneity in mitochondria state
has been linked to differences in transcription rates, growth rates, proliferation and
developmental trajectories[19-21]. Both cancer cells and pluripotent stem cells have atypical
metabolisms and use glycolysis to produce much of their ATP, instead of the
mitochondria-based oxidative phosphorylation, which is the predominant form of
ATP-generation in differentiated cells[22]. It is unknown if this inter-population variation in
proliferation, transcriptome, and mitochondria extends to intra-population variation among
single cells within a single isogenic population.

Pluripotent stem cells exist in various states, such as naïve or primed, based on culture
conditions and embryonic origin[23]. Mouse ESCs reflect the naïve pluripotency state of the
blastocyst epiblast and can be cultured in either serum+LIF or 2i+LIF conditions, the latter
involving inhibitors of FGF/ERK and GSK3 pathways. Culture in 2i+LIF conditions promotes a
ground state more closely mirroring the *in vivo* situation with reduced heterogeneity in
pluripotency gene expression and different cell cycle profile when compared to cells grown
in serum+LIF[24-26]. Nevertheless, even in 2i+LIF conditions, mESCs display a certain amount
of cell-to-cell heterogeneity[27, 28] and it is unclear, how this relates to heterogeneity in
differentiated cell types when it comes to gene expression and its link to proliferation rate.

To understand the relation between intra-population transcriptome heterogeneity and
heterogeneity in proliferation, we developed a FACS-based method to sort cells by
proliferation rate. We applied this method to mouse immortalized fibroblasts and mESCs and
performed RNA-seq on fast, medium and slow proliferating cell sub-populations. We
identified a "proliferation signature", mostly consisting of ribosome-biogenesis (protein
synthesis) and proteasome-related (protein degradation) genes that are highly expressed in
fast proliferating fibroblasts and ESCs. Moreover, the proliferation signature is conserved
across cell-type and species, from yeast to cancer cells, allowing us to predict the relative
proliferation rate from the transcriptome. We used this gene expression signature to predict
proliferation rates in single cells from scRNA-seq data of *C. elegans* development. Unlike
previous models to predict growth rate from gene expression[29], this model has no free
parameters other than the set of genes, and does not suffer from overfitting – it can predict

differences in growth rate in yeast, cancer cells and *C. elegans*, in spite of no data from either
species going into the initial model. When applied to scRNA-seq data from developing *C.*
elegans, this model identified a global slowdown in proliferation rate during development,
with lineage-specific exceptions where some lineages maintain constant proliferation scores,
and others even increase proliferation rate. In contrast to the universality of this main
transcriptional signature, many mitochondria-related genes were upregulated in fast
proliferating fibroblasts, yet down-regulated in fast-proliferating mESCs. Consistent with this,
we found that a high mitochondria membrane potential is indicative of slow proliferating
fibroblasts, while in mESCs this is a property of fast proliferating cells. And the mitochondrial
electron transport chain complex III inhibitor Antimycin treatment cause opposite effects on
the proliferation of fibroblasts and ESCs. Fast, but not slow proliferating fibroblasts are
particularly sensitive to the ATP synthase inhibitor oligomycin. Taken together, these results
show the existence of a core protein-synthesis and protein-degradation expression program
that is conserved across cell types and species, from yeast to mice, and a metabolic and
energy-production program that is highly cell-type specific, with cell-type and
proliferation-rate specific consequences on the effects of mitochondria inhibitors.

Results

A method to sort single mammalian cells by semi-heritable cell-to-cell heterogeneity in
proliferation rate

To understand the causes and consequences of intrapopulation cell-to-cell heterogeneity
in proliferation rate in mammalian cells we developed a method for sorting single
mammalian cells by their proliferation rate **(Figure 1 and Figure S1)**. The cell-permeable dye
carboxyfluorescein succinimidyl ester (CFSE) covalently binds free amines within cells, thus
staining most intracellular proteins at lysine residues. In cell types that divide symmetrically,
such as embryonic stem cells and immortalized fibroblasts[30], the equal dilution of CFSE
into the two daughter cells enables counting of the number of divisions that each cell has
undergone. This method is commonly used to differentiate proliferating from
nonproliferating cells, and to count discrete numbers of cell division, such as in the study of
T- and B-cell proliferation following antigen stimulation[31]. To eliminate confounding effects
due to differences in initial staining we used fluorescence-activated cell sorting (FACS) to
obtain an initially homogeneous cell population of cells with identical CFSE signals **(Figure 1A**
and Figure S1A, B). Thus, the initial CFSE signal is independent of initial cell-to-cell variation
in dye uptake or protein content, as the initial distribution is determined by the FACS gate.
$CFSE_{CFR2}$ conjugates are stable and unable to exit the cell[32]; the dye signal is stable for over
eight weeks in non-dividing lymphocytes[33]. The measured CFSE signal should be relatively
insensitive to cell-to-cell variation in protein degradation. We cultured this sorted starting
cell population for several generations, during which time the CFSE signal decreases with
each cell division **(Figure 1B)**. Consistent with the decrease in CFSE being mostly due to cell
division, the population-level doubling time of each cell type can be calculated based on the
decrease in CFSE signal over time **(Figure 1C, D)**, and these doubling times (19-21 hours for
fibroblasts and 10-12 hours for mESCs) are consistent with those reported by other
methods[34, 35].

To test if the intrapopulation heterogeneity in CFSE that develops after a few doublings
corresponds to intrapopulation heterogeneity in proliferation rates, we stained cells with
CFSE, isolated a homogenous population by FACS, grew ESCs or fibroblasts for 24h or 48h
respectively, and used FACS to isolate the 20% of cells with the highest and lowest CFSE
signal, and measured both viability and the fraction of cells in S phase **(Figure S1C, D)**. We
found that fast proliferating (low CFSE) subpopulations maintain higher proliferation rates for
at least three days **(Figure 1E, F and Figure S1E, F)**, and found no differences in viability
between CFSE subpopulations **(Table S1)**. Thus, intrapopulation differences in CFSE
correspond to semi-heritable differences in proliferation rates.
185
To identify genes whose expression is positively or negatively correlated with
proliferation rate within a single population we grew fibroblasts MEF (mouse embryonic
fibroblast) medium and mouse ESCs in pluripotent ground-state promoting 2i+LIF
medium[36], stained cells with CFSE, performed the initial sort to isolate cells with the same
CFSE signal, and then grew fibroblasts for five days, and ESCs for three days. We then used
FACS to isolate cells with high, medium, and low CFSE signal, and performed RNA-seq on
each sub-population **(Figure 1G)**.
193

Slow-proliferating ESCs are of more naïve pluripotent character than fast-proliferating ESCs
Embryonic stem cells exhibit cell-to-cell heterogeneity in the expression of naïve
pluripotency marker genes such as *Nanog*, *Stella* (*Dppa3*) or *Rex1* (*Zfp42*)[37-39]. Although
this heterogeneity is most apparent in ESCs cultured in serum+LIF, even when cultured in
ground state-pluripotency-promoting 2i+LIF conditions, the sub-population of ESCs with low
NANOG-levels displays a propensity for lineage-priming and differentiation[28, 40]. To
determine if cell-to-cell variation in proliferation rate was caused by a sub-population of
mESCs initiating a differentiation program, we determined the fold-change in expression
between slow and fast proliferating sub-populations for a set of genes that are upregulated
during lineage commitment (see "Differential expression of pluripotency…" in methods). We
found no consistent enrichment of these differentiation genes in fast versus slow
proliferating cells, as they could be found to be expressed in either population **(Figure 2A)**.
However, the slow proliferating subpopulation did have higher expression of genes that are
upregulated in naïve pluripotent cells, and in 2-cell(2C)-like state stem cells **(Figure 2B, C)**,
suggesting that slow proliferating mESCs are in a more naïve pluripotent cell state than their
fast proliferating counterparts.
210

Identification of biological processes correlated with proliferation rate that are conserved
across cell-types and species, and within single populations
To identify functional groups of genes that are differentially expressed between fast and
slow proliferating cells within a single population we performed gene set enrichment analysis
(GSEA)[41, 42] **(Figure 3A, B and Figure S2A, B)** on mRNA-seq data from fast and slow
proliferating subpopulations. We found that in both fibroblasts and ESCs, as well as for
intrapopulation variability in budding yeast FACS-sorted by proliferation rate (data from van
Dijk et al.[8]), genes involved in ribosome-biogenesis and the proteasome are more highly
expressed in fast proliferating cells **(Figure 3C, D and Table S2)**. High expression of ribosomal

genes is a common signature for fast proliferating cells[12, 43], and cancer cells often exhibit high proteasome expression[44-46], but it is not clear if this is related to proliferation in-and-of-itself or due to aneuploidy and other genetic alterations[47]. Our results suggest that coordinated regulation of the ribosome and proteasome is an intrinsic signature of fast proliferating cells that is conserved across cell-types and species.

To test if the coupling between ribosome biogenesis and proteasome expression holds across species and in diverse cell types, we analyzed the bulk RNA-seq data across developmental stages, covering multiple organs in seven species[16]. Ribosome biogenesis and proteasome expression are highly correlated **(Figure 3E)**. The coordinated expression change with developmental stages between ribosome biogenesis genes and proteasome genes across organs and species suggests that the coordination between protein synthesis and degradation is likely to be a conserved feature across a large number of species and cell-types **(Figure S2D)**.

In addition to ribosome-biogenesis and the proteasome, several other gene sets are differentially expressed between fast and slow proliferating cells in both fibroblasts and ESCs **(Figure 3C)**. mTORC1 (mammalian Target Of Rapamycin Complex 1) functions as a nutrient sensor and regulator of protein synthesis, and is regulated by nutrient and cytokine conditions that cause differences in proliferation[48, 49]. We find that, even in the absence of genetic and environmental differences, mTORC1 is more active in fast proliferating cells. Activation of mTORC1 can promote ribosome-biogenesis[48, 50], however, there is still controversy about the regulation of proteasome activity by mTORC1[49, 51-55].

The transcription factor MYC **(Figure S2C)**, and MYC target genes **(Figure 3C and Table S2)** are more highly expressed in fast proliferating cells. MYC is frequently amplified in cancer, regulates the transcription ~15% of all genes[56] and is a master regulator of cell proliferation[57]. Overexpression of MYC promotes ribosome-biogenesis and cell growth rates[58, 59], and active mTORC1 can promote MYC activation[60, 61]. Our data suggest that increased expression of MYC and increased mTORC1 activity are general properties of fast-proliferating cells, and those genetic or environmental perturbations are not necessary to cause differential expression of these pathways.

Defining a proliferation signature to predict the growth rate across species

Expression of typical proliferation markers, such as PCNA and Ki67, did not correlate with intra-population heterogeneity in proliferation **(Figure 4A)**. The high degree of conservation of genes whose expression correlates with intra-proliferation heterogeneity in proliferation, from yeast to mouse, suggests that there should be a set of genes whose expression is predictive of growth rate across all eukaryotes. To build such a set we combined "proliferation correlated genes" – those with a Spearman correlation of rho = 1 in both fibroblasts and ESCs (243 genes) with genes from six ribosome biogenesis and proteasome related gene sets that are significantly enriched in both fibroblasts and ESCs, which result in a final gene set consisting of 370 genes **(Table S4)**, from whose expression we can calculate a proliferation signature score.

To test the ability of this proliferation signature to predict proliferation rates in new data
we used out-of-species cross-validation. While several models have been developed to
predict growth rates from gene expression[29], the performance of these models has been
evaluated using within-experiment cross validation, in which a single sample (e.g., condition
or genotype) was held-out (excluded) and used for testing model performance. Accurate
prediction of growth rates in cells for which actual growth rates cannot be measured, such as
tumors *in-vivo*, or from single-cell RNA-sequencing data, would be more useful. However,
models tested using in-experiment cross-validation (also known as internal validation) are
often over-fit, resulting in poor performance when the model is applied to new data from
new experiments[62].

To overcome this problem, we used the proliferation signature from above, which was
developed using data from mouse cells, to predict growth rate from gene expression in
budding yeast. Our model has correlations of R = 0.82, 0.73 and 0.77 across three different
datasets **(Figure 4B, S3B and S3C)**. In contrast, the model of Wyotck et al.[29], which was
trained on these yeast data, has an out-of experiment predictive power of R < 0.15 (Figure S6
in Wyotck et al.[29]). The Wyotck et al. model is over-fit and cannot predict proliferation
rates in new data. Similarly, our model has better performance (R = 0.65) than a
cancer-specific model[63], which was trained on cancer cell-line data and cannot predict
out-of-experiment **(Figure 4C)** (Figure 4 in Waldman et al.[63]). In contrast to most published
models, our proliferation signature score model performs well on data on which it has not
been trained.

When run on FACS-sorted 2C-like embryonic stem cells (2C::tdTomato+)[64], our
proliferation signature model predicts that 2C-like mESCs proliferate slower **(Figure 4D)**.
2-cell embryos also have uniquely low proliferation scores (**Figure S4H**). These results are
experimentally independent of, and biologically consistent with, our observation that
expression of 2C-like cell state marker genes is higher in slow proliferating mESCs **(Figure 2C).**
This provides further evidence that the proliferation signature we have identified can be
universally applied to predict the proliferative state of many cell types.

Prediction of lineage-specific changes in proliferation rates during C. elegans development by
the proliferation signature
Expression of the most commonly used markers (PCNA and Ki67) for measuring
proliferation rates in bulk populations are cell-cycle regulated; what is really being measured
is the fraction of the population that is in S-phase (PCNA) or is not in G0 (Ki67). Expression of
these markers does not measure proliferation rates in single cells. Single-cell RNA sequencing
is a powerful method for understanding development and differentiation *in vivo*, but it
suffers from high levels of noise at the single-gene level[65]. We reasoned that our
proliferation signature model would be ideal for measuring the proliferation rates of single
cells from scRNA-sequencing data, as the model takes into account the expression of >300
genes, most of which are highly expressed and therefore have low levels of technical noise.
To test the ability of the proliferation signature model on scRNAseq data we used a dataset

of 89,701 cells from *C. elegans* development[66]. We computed the proliferation signature score for each cell and divided the cells into terminal cell types vs preterminal cells. Non-terminally differentiated cells have a higher proliferation signature score (t-test, p = 4.9×10^{-41}) **(Figure 5A)**. A visual comparison of the 89,701 cells in UMAP space, colored by either embryo age[66] or by proliferation signature score **(Figure 5B)**, suggested that, globally, proliferation rates in single cells decreases as development proceeds. However, three clusters of cells did not follow this pattern: germline, intestine and M cells. To quantify the relationship between proliferation rates of single cells and developmental time we binned all cells with same embryo time, and calculated the correlation between proliferation score and developmental time **(Figure 5C)**. The proliferation signature score decreases as the embryo develops (rho = -0.65, p = 9.3 ×10^{-19} and rho = -0.73, p = 8.7×10^{-24} when excluding the three outlier groups **(Figure S4B)**). This conclusion from our single-cell gene expression analysis using the proliferation signature score is therefore quantitatively consistent with lineage-tracing microscopy data from Sulston et al.[67], showing that the rate of cell division within the developing nematode decreases during development **(Figure S4D, S4E)**.

To our surprise, the predicted proliferation rates increased after 650 minutes **(Figure S4C)**. To understand why, we grouped all cells in 650 minutes or older by lineage **(Figure 5D)**. The three outlier groups from UMAP space: germline, M cell and intestine, had the highest proliferation signature among all cell types late in development. Specifically, for these three cell types, the proliferation score did not decrease with the embryo time, but increased or maintained a high level **(Figure 5E)**. This can be explained by lineage-specific characteristics: the germline is the only cell type in *C. elegans* that is continuously proliferating, M cells are a highly proliferative single mesodermal blast cell, and the intestinal cells, although they do not proliferate, continue to increase in both biomass and DNA content through endoreplication. Other cell types with high proliferation scores, such as Z1/Z4, are also known to continue proliferation after 650 minutes[68]. The proliferation signature score decreases with embryo time for most cell types, including body wall cell, hypodermis and ciliated amphid neuron, which are the most prevalent in the single cell RNA-seq dataset **(Figure 5E)**.

Two cell types, hypodermis and seam cells, exhibited very low proliferation signature scores at late time points, while intestinal cells exhibited very high proliferation scores **(Figure 5D, S4F)**. Both these cell types contain multinucleated cells, but these multinucleated cells arise through very different mechanisms: hypodermis and seam cells through cell-fusion, and intestine through endoreplication[68]. Thus, two cell types with seemingly similar properties have highly divergent transcriptomes, and highly divergent mechanisms to reach their similar final state.

Upon development to an L1 larva, a nematode has more than half of the final number of cells present in the adult. In contrast, mammals continue to rapidly increase in cell number even after embryonic development is complete. This difference can be seen in the change in proliferation signature over time, which decreases in *C. elegans*, but increases in human (data from Petropoulos et al.[69]) and mouse (data from Deng et al.[70]) **(Figure 5F, G**

and S4G, H). In conclusion, our proliferation signature genes obtained from mouse fibroblast and ESC data can predict dynamic changes in proliferation rates during the development of various cell types and species, thereby confirming its universal applicability.

Cell-to-cell heterogeneity in mitochondria state predicts variation in proliferation both in ESCs and fibroblasts, but in opposite directions.

While the pattern of within-population proliferation-correlated expression in yeast, mouse fibroblasts and ESCs was broadly similar with regard to genes involved in protein synthesis and degradation, the behavior of metabolic and mitochondria-related genes in fast and slow proliferating subpopulations was highly cell-type specific. Mitochondria membrane and respiratory chain-related gene sets were more highly expressed in fast proliferating fibroblasts, but not in fast proliferating ESCs **(Table 1)**. These results are consistent with differential mitochondrial states in ESCs when compared to differentiated cells like fibroblasts[22], which suggests the existence of different types of metabolism and proliferation-related heterogeneity between pluripotent and differentiated cell-types. We also observed cell-type specific differences in glycolysis, fatty acid metabolism, and other metabolic processes, suggesting fundamental differences in the metabolic pathways required for fast proliferation between pluripotent ESCs and differentiated cells like fibroblasts **(Table 1)**.

The mitochondrial membrane potential is a major predictor of cell-to-cell heterogeneity in proliferation rate in budding yeast[9]. Mitochondria-related genes are more highly expressed in the fast proliferating subpopulation of fibroblasts (**Table 1**). In contrast, these genes are slightly more highly expressed in the slow proliferating subpopulation of ESCs. This suggests that the relation between cell-to-cell heterogeneity in mitochondria state and proliferation may be different in these two cell types. To test the ability of mitochondrial membrane potential to predict proliferation rate in mammalian cells we used the mitochondria membrane potential-specific dye tetramethylrhodamine ethyl ester (TMRE) to stain fibroblasts and ESCs, and performed both RNA-seq and proliferation-rate assays on high and low TMRE sub-populations **(Figure 6A)**.

Unlike the proliferation-based sort **(Figure 1)**, sorting ESCs and fibroblasts by mitochondria-state **(Figure 6 and Figure S5A, B)** resulted in highly divergent expression profiles. ESCs with high TMRE signal had high expression of ribosome-biogenesis, proteasome, MYC-targets and mitochondrial-related genes, while in fibroblasts these gene sets are more highly expressed in the low TMRE sub-population **(Figure 6B, C and Table S5)**. This is consistent with the opposite behavior of mitochondria-related gene sets in proliferation-rate sorted cells from the two cell types **(Table 1)**. We note that the differences between high and low TMRE populations are smaller than the difference between high and low CFSE population **(Figure S5C, D)**, either due to technical limitation of the dye, or because there is less heterogeneity in mitochondria state than there is in proliferation rate.

These expression data make the following prediction: ESCs with high TMRE should have a

shorter doubling time, while fibroblasts with high TMRE should have a longer doubling time. To test this, we sorted fibroblasts and ESCs by TMRE, and found that high TMRE fibroblasts indeed do proliferate more slowly, while high TMRE ESCs proliferate more rapidly **(Figure 6D)**. In addition, we tested the effect of ascorbic acid (vitamin C, an antioxidant) and O_2 levels (ambient 21% atmospheric vs. low 5% physiological levels) on doubling time, but found no significant effects **(Table S6)**.

To investigate additional cell types, we searched for RNA-seq data for cells sorted by mitochondria state, and analyzed RNA-seq data of mouse CD8+ T-lymphocytes that have been sorted by mitochondria membrane potential (TMRM)[21]. CD8+ T cells with high TMRM signal (high ΔΨm) showed higher expression of ribosome-biogenesis and proteasome related genes **(Table S7)**, and proliferate more rapidly[21], thereby behaving in a similar fashion to ESCs.

Thus, across yeast, mouse ESCs, fibroblasts and CD8+ T cells, while mitochondria state and proliferation rate co-vary within a single population, the direction of this correlation is different, with yeast and fibroblasts behaving similarly with each other, and opposite to ESCs and CD8+ T cells.

Perturbation of mitochondria function affects fast and slow proliferating fibroblasts and ESCs in different ways.

In order to investigate the relation between proliferation rate, cell type, and mitochondrial state, we performed perturbation experiments by directly inhibiting mitochondria function. We stained both mouse ESCs and fibroblasts with CFSE and sorted 20% of the viable cells on the peak of CFSE signal to have a homogeneous starting population. After culturing them for 24h or 48h respectively, two bins were sorted: the lowest 20% (fast proliferating cells) and the highest 20% CFSE (slow proliferating cells) **(Figure 7A)**. We then cultured the sorted cells with either medium containing DMSO as mock control, the mitochondrial electron transport chain complex III inhibitor Antimycin, the ATP synthase inhibitor Oligomycin for 16h, washed out the drugs, and measured both viability and proliferation rate.

Both fast fibroblasts and ESCs sorted by CFSE signal maintained a higher fraction of cells in S-phase over two days in growth-media with DMSO, indicating that the proliferation status, fast vs slow, is semi-heritable **(Figure 7B, S6A)**. Interestingly, we found cell-type and proliferation-state specific effects of mitochondria perturbation. Antimycin strongly decreased the fraction of slow-proliferating fibroblasts that were in S-phase but had a weaker effect on fast-proliferating fibroblasts (t-test, p = 0.0089) **(Figure 7C)**. In ESCs, the effect of antimycin appeared somewhat stronger on fast- than on slow-proliferating cells (although it did not differ significantly).

In contrast, fast proliferating fibroblasts were highly sensitive to oligomycin treatment. Specifically, we observed a change in cell morphology upon treatment **(Figure 7D)**. In

comparison with DMSO-treated cells, the cells lost their elongated shape and became more
round and smaller. This led us to hypothesize if that morphology change might be explained
by a mesenchymal to epithelial transition (MET) upon oligomycin treatment. In fact, during
induced pluripotent stem cell (iPSC) reprogramming, MET of fibroblasts is an important early
reprogramming step[71, 72]. In that context, oligomycin treatment has been recently shown
not only to promote a metabolic switch from oxidative phosphorylation to glycolysis, but also
to modulate mesenchymal markers during reprogramming[72, 73]. Therefore we measured
the protein levels of the regulators N-cadherin (mesenchymal marker expressed in
fibroblasts) and E-cadherin (epithelial marker expressed in ESCs) with and without treatment,
by both immunostaining and Western Blot **(Figures 7E and S6A-C)**. We could not detect
E-cadherin in fibroblasts, but we observed reduced expression of N-cadherin in comparison
with DMSO treated-cells in particular in oligomycin-treated fast cycling fibroblasts **(Figure 7E**
and S6A-C). In addition to the change in morphology, oligomycin treatment reduced cell
viability specifically in fast proliferating fibroblasts, but not in slow fibroblasts **(Figure 7F)**. In
conclusion, although we observed both a change in cell viability, morphology and a reduction
in N-cadherin levels, oligomycin treatment did not induce a complete mesenchymal to
epithelial transition in the fast-proliferating fibroblasts as indicated by the lack of E-cadherin
upregulation. Nevertheless, the distinct effects of antimycin treatment on the proliferation of
fibroblasts and ESCs and the subpopulation-specific effect of oligomycin on fast fibroblasts
are in line with a differential dependency on mitochondrial function between the different
subpopulations of fibroblasts and ESCs.
461
462

Discussion

In summary, we have developed a method to sort cells by their proliferation rate and
used these data to identify a pattern of proliferation-correlated gene expression that is
conserved among eukaryotes. We used these data to develop a model that can predict
proliferation rates from gene expression in multiple eukaryotic species and cell types, and for
types of data, such as single-cell RNA sequencing in a developing organism, for which
proliferation rates cannot be measured experimentally.
470
While the CFSE signal is not a measure of the instantaneous proliferation rate, but
instead determined by the average proliferation rate integrated over several days, the fact
that (A) the transcriptomes of the sorted cells are predictive of proliferation rates, and (B)
the cells with low CFSE maintain faster proliferation rates over at least three days, suggests
that there are not likely to be large differences in the instantaneous proliferation rate vs the
average rate, at least for these cell types and experimental timescales.
477
We found that genes involved in protein synthesis (ribosome-biogenesis, translation
initiation), and in protein degradation (the proteasome and proteasome-related protein
degradation) are highly expressed in fast proliferating eukaryotic cells, including mammalian,
nematode and yeast cells. Previous studies have reported that high expression of the
proteasome in fast-growing cells may be necessary in order to degrade misfolded protein,
because the fast protein synthesis in fast-growing cells produce more incorrectly folded

proteins[49, 74, 75]. Even with a constant translation and folding error rate, fast proliferating
cells will produce more protein, and therefore more misfolded protein that needs to be
degraded.
487
488 In all non-cancer mammalian cells we investigated, we also found the mTORC1 signaling
489 pathway enriched in fast proliferating cells and P53-targets enriched in slow proliferating
490 cells. Our results show both upregulations of the mTORC1 signaling pathway and
491 proteasome activity in fast proliferating cells, which is consistent with several previous
492 studies[9, 12-15].
493
Our analysis of fast versus slow proliferating ESCs cultured in 2i+LIF conditions indicated
at several levels that slow proliferating cells were of a more naïve ground state pluripotent
character than fast proliferating cells. First, this was supported by the fact that they displayed
a higher expression of naïve pluripotency marker genes and markers of 2C-like cells **(Figure**
2B, C). Second, we observed enrichment of E2F targets and genes involved in G1 S cell cycle
phase transition **(Table 1)** in our fast cycling ESC population, indicative of a shortened G1
phase and a shorter doubling-time, as described for ESCs cultured in serum+LIF [26]. Finally,
although we could find differentiation genes to be expressed both in fast and slow
proliferating cells **(Figure 2A)**, we saw several differentiation pathways to be enriched
specifically in fast dividing ESCs **(Table 1)**. In summary, even when ESCs are cultured in
ground-state pluripotency promoting 2i+LIF conditions, they display heterogeneity in
proliferation rate, with the slow proliferating being of more naïve pluripotent character when
compared to fast dividing cells.
507
While we observed ESCs behave similarly to other cell types like fibroblasts or yeast
when it comes to gene expression signatures characteristic of fast proliferating cells related
to protein synthesis and turnover **(Figure 3C)**, we found a very different behavior when it
comes to regulation of metabolism. Although the growth rate can be predicted by
mitochondrial membrane potential in *Saccharomyces cerevisiae*[23], where it is negatively
correlated with proliferation rate like in fibroblasts as we show in this study, our results show
mitochondrial membrane potential to be positively correlated with proliferation rate in ESCs
(Figure 6D). This suggests mitochondrial membrane potential has different functions in
pluripotent cells when compared to differentiated cell types or yeast. This is corroborated by
our gene expression analysis of cells with high vs. low mitochondrial membrane potential
(Figure 6B, C), where we found pathways linked with fast proliferating cells to be enriched in
fibroblasts with low mitochondrial membrane potential but on the contrary, enriched in ESCs
with high mitochondrial membrane potential. Surprisingly, primed pluripotent stem cells
have been described to rely more on non-oxidative, glycolysis-based metabolism than naïve
pluripotent stem cells[76-78], which appears in contradiction with our result that our slow
proliferating, mitochondria activity low ESCs being more naïve-like. However, TMRE is not a
direct measure of ATP generation by mitochondria; yeast cells that are respiring and
producing all of their ATP using their mitochondria, and yeast cells unable to respire, both
have high TMRE signals[9]. Differentiated cells in general rely more on oxidative metabolism
than pluripotent cells, therefore our fast proliferating ESCs could potentially reflect a more

differentiation prone state.
529
In vivo, cells exhibit a great degree of variability in proliferation rates, from terminally
differentiated neurons, to slowly proliferating cancer stem cells, to rapidly proliferating
embryonic stem cells. Many cell types, such as hemopoietic stem cells, contain both
proliferating and non-proliferating populations. The proliferation signature model, because
of its applicability across all tested species and cell types, provides a useful tool for
understanding *in vivo* development for systems, in which precise measurements of
proliferation are impossible. Our model has been validated on scRNA-seq data, using
published time-lapse microscopy of cell lineages in *C. elegans* as the ground truth[67].
However, it is technically challenging to do microscopy or to otherwise measure proliferation
of individual cells inside of a developing mouse embryo, or in a tumor in a patient. Models
that can accurately predict difficult to measure properties, such as proliferation rate, from
easy to measure ones, such as gene expression, will therefore aid in our understanding of
complex biological processing during tumor formation, differentiation, and development.
543
544 MATERIALS AND METHODS
545
Cell culture growth conditions
Tail tip fibroblasts (TTFs) were isolated from a female newborn mouse from a *Mus musculus*
x Mus Castaneus cross and immortalized with SV40 large T antigen[79]. The clonal line
68-5-11[80] was established and maintained in DMEM supplemented with 10% serum
(LifeTech), HEPES (30mM, Life Tech), Sodium Pyruvate (1mM, Life Tech), non-essential amino
acids (NEAA) (Life Tech), penicillin-streptomycin (Ibian Tech), 2-mercaptoethanol (0.1mM,
Life Tech).
The mouse embryonic stem cell (ESC) line EL16.7 (40XX, Mus musculus/*M.castaneus* hybrid
background[81] was maintained on gelatin coated tissue culture dishes and passaged every 2
days by seeding around $2x10^6$ cells in 2i+LIF medium. Accutase (Merck Chemicals and Life
Science) was regularly used for cell detachment when passaging cells. 2i+LIF medium
contains a 1:1 mixture of DMEM/F12 supplemented with N2 (LifeTech) and neurobasal
media (LifeTech) supplemented with glutamine (LifeTech), B27 (LifeTech), insulin (Sigma),
penicillin-streptomycin (Ibian Tech), 2-mercaptoethanol (LifeTech), LIF (Orfgenetics),
PD0325901 (Sigma) and CHIR9021 (Sigma). Both TTFs and EL16.7 were cultured at 37C in 5%
CO_2.
562
Proliferation and doubling time analysis
ESCs and fibroblasts were plated on 10 cm plates at $5.3x10^6$ and $7.3x10^5$ concentrations,
respectively. Cells were expanded and counted for 7 days. To monitor distinct generations of
proliferating cells, carboxyfluorescein succinimidyl ester (CFSE, Thermo Fisher Scientific) was
used to stain the cells and the dilution of the dye was detected by flow cytometry every day.
CFSE was dissolved in dimethyl sulfoxide at a concentration of 5 mM as stock solution and
CFSE was added to a 1 ml cell suspension, to a final concentration of 5uM or 10uM. After the
addition of CFSE, cells were incubated at 37°C for 20 min. Then the cells were washed twice
with complete medium and maintained on ice until use in a buffer containing PBS, 2% serum

and 1% pen-strep. Cell viability was determined by DAPI (Biogen Cientifica) staining. Dye
signals were measured on an LSRII flow cytometer.
574
RNA-seq
To collect cells with different growth rates, cells were isolated by sorting at room
temperature according to the CFSE signal (median and high CFSE signal). ESCs and fibroblasts
were sorted into 1.5 ml Eppendorf tubes containing medium and were cultured for 3 days
and 5 days respectively in specific culture conditions as described earlier. For each cell line
three bins were sorted: the lowest 10%, the median 10% and the highest 10% CFSE. Cells
were sorted into prechilled 1.5-ml Eppendorf tubes containing 200 μl medium each. Cells
were then centrifuged at 1000 rpm for 5 min, the media removed and the resulting cell pellet
was used for RNA extraction. All bins were treated identically throughout the process.
Cellular RNA was extracted using the Maxwell RNA Purification Kit and processed for RNA
sequencing. For biological replicates, all experiments were repeated on three or four
different days. Expression was quantified using Kallisto v0.42.3[82] from the raw reads (no
pre-processing) using the gencode.VM18.transcript annotations. We experimented with
multiple methods for batch effect removal using the R package SVA[83] and found that the
results of the GSEA, with regards to which gene sets were differentially expressed between
fast and slow, or high and low TMRE cell populations, did not change. We therefore used the
original data.
PCA on TMRE sorted biological replicates showed that one TMRE sort was an extreme outlier
(Figure S5C); this pair was discarded from all analysis.
594
BrdU Staining
Cell Proliferation was measured by the incorporation of bromodeoxyuridine (BrdU). Every
24h BrdU was added at a final concentration of 10 μM to the cells. Incubation under the
appropriate growth conditions occurred for 30 minutes for ESCs and 45 minutes for
fibroblasts to pulse label the cells. Cells were trypsinized, spun down at 1050 rpm for 5
minutes. After washing them in ice-cold PBS, cells were fixed overnight in ice cold Ethanol
(70%) while maintaining a gentle vortex. The following day the Ethanol fixed cells were
centrifuged and the DNA denatured by adding 2N HCl - 0.5%Triton X-100 for 30min at room
temperature. Then cells were centrifuged and resuspended in 0.1 M $Na_2B_4O_7$, pH 8.5 for 10
minutes at room temperature. After spinning them down the cells were incubated overnight
at 4C with PerCP/Cy5.5 anti-BrdU (1:30 dilution) (BioLegend) in a buffer containing 0.5%
Tween 20 / 1% BSA/PBS and RNase (0.8 mg/ml). The following day cells were washed in PBS
and spun down at 1050rpm for 5min at room temperature. The pellet was resuspended in
PBS with DAPI (1:1000) and analyzed in an BD LSRII flow cytometer.
609
Mitochondria inhibitor treatment assay
For the assessment of chemical inhibitors on membrane potential changes, cells were
incubated with medium containing DMSO (0.1%, mock control), Antimycin A (500 nM),
Oligomycin (1uM) for 16h. Cells with or without treatment were washed with PBS and
trypsinized. After spinning the cells for 5 minutes at 1050 rpm at room temperature, the cell
pellet has been stained with 50nM TMRE for 20 min at 37C. After 2 times washes with PBS,

cells were resuspended in PBS containing DAPI and immediately analyzed by flow cytometer BD LSRII.

Western blot

Cells were lysed in a lysis buffer (20 mM Tris-HCL, pH 8.0, 150 mM NaCl, 1% Triton X-100, supplemented with protease inhibitors cocktail) and centrifuged for 30 minutes at 16000g. The supernatant was boiled in SDS loading buffer. After SDS-PAGE, the samples were transferred to a polyvinylidene difluoride membrane using a transfer apparatus according to the manufacturer's protocols (Bio-Rad). After incubation with 5% nonfat milk in TBST (10 mM Tris, pH 8.0, 150 mM NaCl, 0.5% Tween 20) for 1h, the membrane was washed once with TBST and incubated with antibodies against N-Cadherin (BD Biosciences, 1:1000), E-Cadherin (BD Biosciences, 1:1000), GAPDH (Santa Cruz, 1:5000), at 4 °C for 16 h. Membranes were washed three times for 10 min and incubated with a 1:5000 dilution of Rabbit Anti-Mouse Immunoglobulins/HRP for 1.5 h. Blots were washed with TBST three times and developed with the ECL system (Amersham Biosciences) according to the manufacturer's protocols.

Immunofluorescence staining

Cells were grown in 8-well Lab-Tek chamber slides (Thermo Fisher Scientific) and fixed in 4% paraformaldehyde for 10min at room temperature. Then, washed three times in PBS. Fixed cells were permeabilized in 0,5% Triton X-100 (Sigma-Aldrich) in PBS buffer for 10min at room temperature. And then washed in PBST (PBS with 0.1% Tween (Sigma-Aldrich)) for 2min at RT. Then cells were incubated in a blocking solution containing 10% bovine serum albumin (BSA, Sigma) and 0.01% Triton X-100 for 1h at room temperature. Cells were then left at 4°C overnight in a blocking solution containing the primary antibody: mouse E-Cadherin (BD Biosciences, 1:1000) and mouse N-Cadherin (BD Biosciences, 1:1000). The next day, the cells were washed three times in PBS and then incubated with the secondary antibody for 45min at room temperature. Goat anti-mouse IgG, (1:1000, Life Technologies) conjugated to Alexa Fluor-555, was used as a secondary antibody. Nuclear staining was performed with DAPI (1:1000, Biogen Cientifica). Images were taken with a Leica TCS SP8 confocal microscopy system and were analyzed with Fiji (ImageJ).

Differential expression of pluripotency, 2C-like state and lineage commitment-related genes in mESCs sorted by proliferation rate (CFSE)

Pluripotent state gene markers are chosen from 4 different studies[28, 84-86], only genes that are used as pluripotent state gene marker in at least 2 of these 4 papers are used in this paper. Lineage commitment and 2C-like state gene markers are the same as genes in Figure 5A and key differentiation regulators in Figure 6 of Kolodziejczyk et al.[84]. To see the corresponding pluripotent cell state of fast and slow proliferating mESCs, we calculated the mean expression of naïve pluripotent markers in four fast-proliferating and four slow-proliferating replicates and log2(fast/slow) was calculated to compare genes expression in fast proliferating subpopulation and slow proliferating sub-population. The same method was applied to lineage commitment gene markers and 2C-like state gene markers.

Mitochondrial Membrane Potential Measurements.

The relative mitochondrial transmembrane potential ($\Delta\Psi$m) was measured using the membrane-potential-dependent fluorescent dye TMRE (Tetramethylrhodamine, Ethyl Ester, Perchlorate) (Molecular Probes, Thermo Fisher Scientific)[87]. For TMRE staining fibroblasts and ESCs were grown, washed in PBS, trypsinized and resuspended in PBS with 0.1% BSA and TMRE added at a final concentration of 50nM, from a 10uM stock dissolved in DMSO. Cells were incubated for 20min at 37C, washed with PBS and were analyzed by flow cytometry or sorted.

Cell sorting

Cells at 80% confluence in 10cm plates were trypsinized, centrifuged at 1000rpm for 5min and stained with medium containing 10uM CFSE for 20min. Then cells were washed twice with PBS and stained with DAPI as viability dye. To have a homogeneous starting population, 20% of the viable cells were sorted according to the proliferation rate on the peak of CFSE signal and re-plated. ESCs and fibroblasts have been cultured for 24h or 48h respectively and two bins were sorted: the lowest 20% (fast proliferating cells) and the highest 20% CFSE (slow proliferating cells). Cells were sorted into prechilled 1.5-ml Eppendorf tubes containing 200 µl medium each. Cells were then centrifuged at 1000 rpm for 5 min, the media removed and plated in their culture medium. To monitor their proliferation rate the dilution of the CFSE dye was detected by flow cytometry every day up to 3 days for ESCs and 5 days for fibroblasts. Dye signals were measured on an LSRII flow cytometer.

For the CFSE sort (no TMRE), cells were stained with CFSE and DAPI, and we used FACS to obtain a population of viable cells the same CFSE signal. We then grew cells for 3 or 5 days, and every 24 hours measured the CFSE signal using flow cytometry. Staining did not have a strong effect on cell viability or proliferation **(Figure S7)**.

For the TMRE sort for proliferation rate, cells were stained with CFSE and TO-PRO-3, and we used FACS to obtain a population of G1 cells with the same CFSE signal. We then grow cells for 3 or 5 days, and every 24 hours measured the CFSE signal using flow cytometry.

In order to have a homogeneous starting population, both cell types were stained with Hoechst (10 ug/ml, Life Technologies) to pick cells in G0/G1 phase. Within this population, cells were selected according to the proliferation rate on the peak of CFSE signal prior to staining them with the dye. Then cells were sorted by TMRE into three bins: low, medium and high with a BD Influx cell sorter into prechilled 1.5 ml Eppendorf tubes containing 200 µl medium each. Cells were then centrifuged at 1000 rpm for 5 min, the cell pellet was washed with PBS and used for RNA extraction. All bins were treated identically throughout the process. Cellular RNA was extracted using the Maxwell RNA Purification Kit and processed for RNA sequencing. Cell viability was determined by TO-PRO-3 (Thermo Fisher Scientific) staining.

To test the effect of O_2 levels and ascorbic acid/vitamin C in both cell types, sorted cells from each bin were plated into each of the four different conditions (low O_2 (5%), normal oxygen

growing conditions, and with or without ascorbic acid/vitamin C (25 ug/ml, Sigma-Aldrich))
in duplicate. After one day of recovery from the sorting, the cells were washed in PBS, were
trypsinized, and counted. After seeding the same initial number, the rest of the cells was
analyzed on a BD Fortessa analyzer. Every day a sample from each condition and replicate
was taken for counting, and stained with 50 nM TMRE, up to 3 days for ESCs and 5 days for
fibroblasts, and both TMRE and CFSE were measured by flow cytometry.

Images of CFSE and TMRE stained cells are shown in **Figures S8** and **S9**.

Gene set enrichment analysis (GSEA)
GSEA was performed using the GSEA software and the MSigDB (Molecular Signature
Database v6.2)[88, 89]. We use signal-to-noise (requires at least three replicates) or log2
ratio of classes (for experiments with less than three replicates) to calculate the rank of each
gene. The maximum number of genes in each gene set size was set to 500, the minimum to
15, and GSEA was run with 1000 permutations. We provided all GSEA results in this study
(Table S8).

Enrichment map
Enrichment map of this study (Figure 3C, D and Figure 6B, C) are created using
EnrichmentMap in Cytoscape[90, 91], we refer to Reimand et al's protocol[92] for using
EnrichmentMap.
We imported the output file of GSEA to EnrichmentMap and set FDR threshold as 0.1, other
parameters set as default.

Coordination of expression of ribosome biogenesis and proteasome related genes
We first calculate the mean expression (average of log2(TPM+1)) of ribosome biogenesis
genes (genes in GO preribosome gene set) and proteasome genes (genes in GO proteasome
complex gene set) across organ developmental time course, then we calculate the Pearson
correlation of ribosome biogenesis and proteasome.

Calculation of proliferation signature scores
To obtain proliferation correlated genes, we first calculate, for each gene, the Spearman
correlation with proliferation rate, as measured by the decrease in CFSE signal, in both
fibroblasts and ESCs. We define "proliferation correlated genes" as genes that have a
correlation of 1 in both fibroblasts and ESCs (243 genes). To this set we add genes from six
ribosome biogenesis and proteasome related gene sets that are significantly enriched in
both fibroblasts and ESCs, which result in a final gene set consisting of 370 genes **(Table S3)**
and we called this gene set proliferation signature.
To apply proliferation signature in other species, the R package Biomart[93, 94] was used to
obtain homologous genes of other species in this study and to map across different gene
naming schemes (eg: transfer Ensemble gene id to Entrez gene id).

Prediction of growth rates using proliferation signature

Published expression profiling data for yeasts cultured in the chemostat with controlled
growth rate from Airoldi et al. (dataset1)[95], Slavov et al. (dataset2)[96], Regenberg et al.
(dataset3)[13] and cancer cell lines with corresponding growth rate[63] were downloaded.
For each dataset, we used ssGSEA to calculate the enrichment score of a gene set containing
all proliferation signature genes, and calculated the Pearson correlation of proliferation
signature score with growth rate. ssGSEA[97] is a rank-based method that computes an
overexpression measure for a gene set of interest relative to all other genes in the genome.
We use R package GSVA to apply ssGSEA with default settings[98].
We also used another method to calculate proliferation signature score for 3 yeast datasets.
We use the sum of genes expression for all genes in the proliferation gene set to represent
proliferation signature score **(Figure S3D-F)**, the result is slightly worse than the ssGSEA
method.
759
760 **Proliferation score of 2C-like mESCs and non-2C-like mESCs**
761 RNA-seq data (GSE33923) of 2C-like mESCs are from Macfarlan et al.[64], who FACS
762 separated 2C-like cells (high MuERVpromoter driven expression, 2C::tdTomato$^+$) from
763 non-2C-like mESCs (2C::tdTomato$^-$). We calculated the proliferation signature score for each
764 of the six samples, and used a paired t-test to control for differences between replicates.
765
Brief description of experiments from other papers
In van Dijk et al.[8] *cts1Δ* histone-GFP budding yeast undergo cytokinesis to separate mother
and daughter cells, but these cells remain physically attached to each other by their cell wall.
Thus, starting from an initial population of single cells in G1, variation in proliferation rate
can be measured by variability in histone-GFP signal in physically connected clusters of cells.
Each cluster contains cells descended from the same ancestor cell.
In Dhar et al.[9] wild-type yeast were stained with TMRE, and sorted into four bins with
varying TMRE signal.
In Sukumar et al.[21] pmel-1 T cell receptor (TCR) transgenic mice were injected with
recombinant vaccinia virus encoding hgp100 (gp100-VV). Five days after vaccination, they
isolated CD8+ T cells, stained them with the lipophilic cationic dye tetramethylrhodamine
methyl ester (TMRM) (25 nm for 30 min at 37°C) and FACS-sorted the highest and lowest
7-10% of cells for subsequent RNA-seq.
779
Proliferation scores of preterminal cell lineages vs terminal cell types
Preterminal cell lineage and terminal cell type pseudo-bulk RNAseq data of *C.elegans* were
downloaded from Murray et al.[66], specifically, gene expression profile for terminal cell
types and preterminal cell lineage is in Table S7 and Table S8, annotation file for terminal cell
types and preterminal cell lineage is in Table S2 and Table S4. As there are multiple time
points for one terminal cell type, we only use the sample with maximum time point to
represent corresponding terminal cell type, processed data provided in this study **(Table S9)**.
For each cell we calculate proliferation signature score, and a t-test was used to compare the
mean proliferation signature score of all cells in each of the two groups.
789
C. elegans **scRNA-seq data analysis**

C. elegans scRNA-seq data was provided in R package "VisCello.celegans". After loading the
package, type cello() to load all data into the current environment. We calculated the
proliferation signature score for all single cells, then color them by proliferation signature
score in UMAP space. The calculation of proliferation signature score for single cell data is
different from the calculation for bulk RNA-seq data. We just sum up the expression value of
genes in proliferation signature gene set to get proliferation signature score, but not use
ssGSEA consider ssGSEA is a rank-based method, however most of the genes have 0
expression in this scRNA-seq data set **(Figure S4A)**, which makes it not appropriate to use
ssGSEA.
We binned all single cells by their embryo time. We first calculate the mean proliferation
score for cells with same embryo time, then calculate Spearman correlation of this mean
proliferation score with embryo time, the result is rho = -0.65 (p = 9.3 ×10^{-19}), the correlation
of unbinned data is -0.41 (p < 2.2×10^{-16}). After excluding three special cell types germline, M
cell and intestine, the result is rho = -0.73 (p = 8.6 ×10^{-24}), the correlation of unbinned data is
-0.45 (p < 2.2×10^{-16}).
806
Experimental data for *C. elegans* development
Developmental data of *C. elegans* was extracted from figure 4 of Sulston et al.[67]. This
figure is cell number (live nuclei number) change over embryo time and we use
WebPlotDigitizer[99] to extract data. We use the data to plot log2 cell number change over
embryo time. The difference of log2 cell number for two adjacent time points divided by the
difference of time is the proliferation rate of mean of two time points.
813
814
DATA AND CODE AVAILABILITY
The code for performing analysis is available at
https://github.com/carey-lab/Proliferation_Signature_Public. Raw and processed RNA-seq
data created in this study are available on GEO (Gene Expression Omnibus) with the
accession code GSE139594. Information of all published data used in this study is provided
(Table S10).
821

822 **AUTHOR CONTRIBUTIONS**
823 Z.J. and S.F.G. made the figures. Z.J. and M.B. analyzed the data. S.F.G. and M.B. did the
824 experiments. L.B.C. and B.P. supervised the project. L.B.C., Z.J., S.F.G. and B.P. wrote the
825 manuscript. All authors read and approve of the final manuscript.
826

ACKNOWLEDGMENTS
We thank the CRG/UPF flow-cytometry core and CRG Genomics facilities for help with
experiments, and Yang Zhao for comments on the manuscript.
This work has been funded by the Spanish Ministry of Science, Innovation and Universities
(BFU2014-55275-P and BFU2017-88407-P to B.P. and BFU2015-68351-P to L.B.C.), the AXA
Research Fund and the Agencia de Gestio d'Ajuts Universitaris i de Recerca (AGAUR, 2017
SGR 346 to B.P. and 2014 SGR 0974 & 2017 SGR 1054 to L.B.C.), the National Natural Science
Foundation of China (nos.3191101182 to L.B.C.). We would like to thank the Spanish Ministry

of Economy, Industry and Competitiveness (MEIC) to the EMBL partnership, to the 'Centro de
Excelencia Severo Ochoa', and the Unidad de Excelencia María de Maeztu, funded by the
MINECO (MDM-2014-0370). We also acknowledge the support of the CERCA Programme of
the Generalitat de Catalunya. L.B.C. was supported by funding from Peking University and
from the Peking-Tsinghua Center for Life Sciences, and from the Research Fund for
International Young Scientists (National Natural Science Foundation of China).

841
842
843
844

FIGURE LEGENDS AND TABLE

Figure 1. A CFSE-based method to sort mammalian cells by proliferation rate.
(A) Cells were stained with CFSE and a subpopulation of cells with identical CFSE levels was
collected by FACS. Growth for several generations resulted in a heterogeneous cell
population with a broad CFSE distribution, and cells with high, medium, and low CFSE signal

(slow, medium and fast proliferation, respectively) were sorted by FACS for RNA-sequencing.
(B) The change in the CFSE distribution over time, for fibroblasts and ESCs. **(C, D)** The
population-level doubling time can be calculated by fitting a line to the median of the
log2(CFSE) signal. We discard data from time 0, cells immediately after the sort, because
the CFSE signal decreases in the initial hours, even in the absence of cell division, likely due
to efflux pumps. **(E, F)** BrdU was used to measure the % of cells in S-phase for FACS-sorted
fast and slow proliferating subpopulations. Fibroblasts: 4 replicates, p = 0.0002441. ESCs: 3
replicates for ESCs, p = 0.001953. p-values are for binomial tests across all biological
replicates that the two populations have the same percentage of cells in S-phase. **(G)**
Examples of genes whose expression positively or negatively correlated with proliferation
rate. Each line is one biological replicate, and the error bars are 95% confidence intervals for
each expression value.
863

864
Figure 2. Slow-proliferating ESCs display a more naive pluripotent stemness character than
fast-proliferating ESCs.
(A) Comparison of lineage commitment-related gene expression between fast and slow
proliferating subpopulations. **(B)** Comparison of pluripotency-associated gene expression
between fast and slow proliferating subpopulations. **(C)** Comparison of 2C-like state markers
expression between fast proliferating subpopulation and slow proliferating sub-population.
Dashed lines separate genes expressed preferentially in slow- (left of dashed line) or in
fast-proliferating (right of dashed line) ESCs. P-values are from binomial tests, testing if genes
are more often highly expressed in slow cells than would be expected by chance (53.5% of all
genes are more highly expressed in slow cells).
875

Figure 3. Functional pathways for which cell-to-cell heterogeneity in expression correlates with proliferation rate across cell types and species.

(A) In Gene Set Enrichment Analysis, genes are sorted by their fast/slow expression value (left panel, bottom), and each gene is represented by a single black line (left panel middle). The enrichment score is calculated as follows: for each gene not in the GO preribosome gene set, the value of the green line decreases, and for each gene in the gene set, the value of the green line increases. The ES score will be near zero if the genes in a gene set are randomly distributed across the sorted list of genes, positive if most genes are to the left, and negative if most genes are to the right. **(B)** The heatmap (right panel) shows the expression (z-scored read counts) of preribosome genes in fibroblasts across four biological replicates of the CFSE sorting experiment. **(C)** Gene sets enriched (FDR<0.1) in both fibroblasts and ESCs were mapped as a network of gene sets (nodes) related by mutual overlap (edges), where the color (red or blue) indicates if the gene set is more highly expressed in fast (red) or slow (blue) proliferating cells. Node size is proportional to the total number of genes in each set and edge thickness represents the number of overlapping genes between sets. **(D)** GSEA results

892   (FDR<0.1) of *S. cerevisiae*[8] that sorted by cell-to-cell heterogeneity in proliferation rate. **(E)**
893   Pearson correlations of mean expression (average of log2(TPM+1)) of ribosome biogenesis
894   genes vs proteasome genes across organ developmental time courses in seven species (see
895   also Figure S1).
896

897
Figure 4. A proliferation signature model can predict relative growth rates from gene
expression for species and cell-types on which it was not trained.
(A) Genes and proliferation signature spearman correlation with proliferation rate (sorted by
CFSE). Compare with Ki67 or PCNA, proliferation signature has a better correlation with
proliferation rate. **(B)** Using the proliferation signature to predict growth rate in budding
yeast, we apply ssGSEA to calculate the enrichment score of proliferation signature for each
sample. The Pearson correlation of proliferation signature score with growth rate is 0.82 (p =
8.9×10^{-7}). **(C)** Using the proliferation signature to predict growth rate in cancer cell lines, the
Pearson correlation is 0.65 (p = 1.9×10^{-8}). **(D)** Comparison of proliferation signature score
between 2C-like ESC and non-2C-like ESC (paired t-test, p = 0.04669).
908

Figure 5. Proliferation signature of cell development.

(A) A cartoon showing four terminal cells, and a partial linage showing the final four generations of preterminal cells. Comparison of single-cell proliferation signatures between preterminal cell lineage and terminal cell types (t-test, p = 4.9×10⁻⁴¹). **(B)** UMAP projection of 89,701 cells. Cells in the left panel are colored by estimated embryo times; in the right panel by proliferation signature score. **(C)** To calculate the proliferation signature score (y-axis) at each time point (x-axis) cells are binned by embryo time, and the mean proliferation signature score for all cells in the same bin is calculated. The spearman correlations are -0.65 (p = 9.3 ×10⁻¹⁹) for binned data and -0.42 (p < 2.2e-16) for unbinned data. **(D)** Boxplots (line shows median, boxes interquartile range) of proliferation signature score for all cells with embryo time > 650min. **(E)** Temporal dynamics of proliferation scores of select cell lineages, showing the average proliferation score for all single cells in that lineage, at each time point. **(F-G)** Boxplot of *C. elegans* **(F)** and human **(G)** proliferation signatures as a function of developmental time, from scRNAseq data.

Figure 6. Expression of proliferation-related gene sets in cells sorted by intra-population
heterogeneity in mitochondria membrane potential.
(A) Cells were stained with Hoechst and CFSE and a homogenous population of equally sized
cells in G1 with equal CFSE was obtained by FACS. These cells were stained with TMRE sorted
by TMRE, and then used for RNA-seq, or allowed to proliferate to measure the doubling time
of each TMRE sub-population. **(B, C)** Enrichment maps of fibroblasts and ESCs sorted by
TMRE. **(D)** Doublings times, as estimated by the measured by the decrease in CFSE signal
over time, for high, medium and low TMRE sorted cells. P-values are from ANOVA, testing if
TMRE is predictive of doubling time (see methods).

Figure 7. Growth-rate and cell-type specific effects of mitochondria inhibitors on proliferation rate, cell viability and cell state. (A) Schematic of the experimental setup for measuring the effects of mitochondria inhibitors on slow and fast proliferating cells. **(B)** Fast proliferating Fibroblast and ESCs sorted by CFSE signal maintained a higher fraction of cells in S phase over two days of growth in media+DMSO. **(C)** Effect of antimycin treatment on fast and slow proliferating fibroblasts and ESCs. **(D)** Fast fibroblasts changed morphology after the treatment with oligomycin. Scale bars = 80μm. **(E)** Immunostaining of fibroblasts for N-Cadherin and DAPI after drug treatment. Fast fibroblasts lose N-Cadherin staining specifically after oligomycin treatment. Scale bars = 15μm. **(F)** Effect of oligomycin and antimycin treatment on fibroblast viability. Oligomycin has a specific effect on the viability of fast-proliferating fibroblasts.

 Table 1. Gene sets whose expression exhibits opposite correlations with growth between

 fibroblasts and ESCs. Shown are representative gene sets whose expression is significantly

 correlated with proliferation in either fibroblasts or ESCs, but whose expression changes with

 proliferation in opposing directions.

 SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Fast proliferating subpopulations maintain higher proliferation rates than slow proliferating subpopulations for at least 3 days.

FACS gating strategy for CFSE staining to get cell subpopulations with different proliferation rates.

(A, B) The gating strategy for CFSE staining to get cell subpopulations with different proliferation rates in fibroblasts **(A)** and in ESCs **(B)**. Slow, medium and fast proliferating cell subpopulations were sorted by FACS according to their CFSE signal. Then RNA-seq was performed on each of the three subpopulations. **(C, D)** FACS gating strategy for measuring the heritability of proliferation rates. The gating strategy for measuring the heritability of proliferation rates in fibroblasts **(C)** and in ESCs **(D)**. In all experiments, the laser voltage was increased so that, when sorting high and low CFSE cells, the modal CFSE signal was at least 10^3; the voltage is not the same for the first and second CFSE sorts. **(E)** 3 Replicates of fibroblasts that similar to figure 1E. **(F)** 2 Replicates of ESCs that similar to figure 1F.

Developmental stages (early to late)

Figure S2. Functional pathways for which cell-to-cell heterogeneity in expression correlates with proliferation rate across cell types and species.

(A) GSEA result plot of Go preribosome genes set for ESC. **(B)** The heatmap (right panel) shows the expression (z-scored read counts) of preribosome genes in ESCs across four biological replicates of the CFSE sorting experiment. **(C)** Higher expression of Myc in both fast proliferating ESCs and fibroblasts. log2 fold change of Myc expression between fast and slow proliferating subpopulation in both ESCs and fibroblasts, each cell type 4 replicates. **(D)** Correlated changes in the expression of ribosome biogenesis and proteasome related genes during organ development. Change of average expression of log2(TPM+1) of genes in ribosome biogenesis (Go preribosome) gene set and proteasome complex (Go proteasome complex) gene set with developmental stages across different organs in seven species[16]. Points (circle and triangle) are the mean expression of replicates, error bars represent the maximum and minimum value in the replicates.

Figure S3. Proliferation signature scores predict growth rate, using different methods of calculation, and different species.

(A-C) Using the Normalized Enrichment Score from ssGSEA to predict growth rate in three different data sets. The Pearson correlation of proliferation signature score with growth rate in are R = 0.82 (p = 8.9×10^{-7}), R = 0.73 (p = 1.3×10^{-8}) and R = 0.77 (p = 3.7×10^{-3}). **(D-F)** Similar to figure A-C, but using the sum of expression values for all genes in the proliferation signature gene set to calculate proliferation signature score. The Pearson correlation of proliferation signature score with growth rate are R = 0.83 (p = 7.0×10^{-7}), R = 0.73 (p = 1.6×10^{-8}) and R = 0.55 (p = 0.65×10^{-2}).

Figure S4. Lineage-specific proliferation signature scores during *C. elegans* development.
(A) A density histogram of counts (UMI) across 1000 randomly sampled cells; 95.6% of genes have zero reads. This causes ssGSEA to give unreliable results, so the sum of expression values method is used for calculating the proliferation signature score for single cells. **(B)** Cells are binned by embryo time, and the mean proliferation signature score for all cells not the three outlier cell types (germline, intestine and M cells). The Spearman correlation is rho = -0.73 (p = 8.7×10^{-24}) for binned data, and rho = -0.45 (p < 2.2×10^{-16}) for unbinned data. **(C)** Similar to figure 5C, but only showing cells with an age higher than 650 minutes, rho = 0.5 (p = 1.5×10^{-2}). **(D, E)** The change in cell number, and the rate of change in cell number, during development, as measured by microscopy[67]. **(F)** Change in proliferation signature score for

1014   seam cells, which form multinucleated cells through cell-fusion. **(G)** Single-cell gene
1015   expression data from Petropoulos et al.[69] projected onto the first two principal
1016   components and colored by proliferation signature score or developmental stages. And
1017   boxplot shows the change of proliferation signature score with developmental stages. **(H)**
1018   Similar to figure G, but use scRNA-seq data from Deng et al.[70].
1019
1020

Figure S5. FACS gating strategy for TMRE staining and volcano plots for fibroblasts and ESCs sorted by CFSE or TMRE.

(A, B) The gating strategy for TMRE staining to get cell subpopulations with different mitochondrial states in fibroblasts **(A)** and in ESCs **(B)**. We use Hoechst to get cells in G0/G1, gate by CFSE to get a more uniform cell population, and separate populations with high and low TMRE signal, then do RNA-seq on each of the two subpopulations. **(C, D)** Deseq2 was

1028 used to calculate log2 fold change and adjusted p-values for CFSE sorting **(C)** and TMRE
1029 sorting **(D)**, combining biological replicates. To set the axes to be maximally informative,
1030 genes with $p < 10^{-5}$ had p set to 10^{-5}, and those abs(log2 fold change) > 5 were truncated at
1031 -5 or +5. **(E)** PCA for RNA-seq data of ESCs sorted by TMRE. Low TMRE ESCs of replicate 1 is
1032 an outlier, so we remove replicate 1 for all analysis.
1033
1034

1035
1036 **Figure S6. N-cadherin and E-cadherin levels in oligomycin and antimycin treated cells.**
1037 **(A)** E-cadherin staining of ESCs as positive control for E-cadherin detection, Scale bar = 80μm.
1038 **(B)** Western blot for N-cadherin and E-cadherin (Gapdh = loading control) in DMSO- and
1039 drug-treated fibroblasts and ESCs. **(C)** Immunostaining for E-cadherin does not show
1040 detectable levels in fibroblasts, Scale bars = 15μm.
1041
1042

	Seeded cell number	Harvested cell number (after 24h)	Viability	Doubling time
Unstained	150000	3.4×10^5	97%	≈ 20.3
CFSE	150000	3.52×10^5	98%	≈ 19.5
Hoechst	150000	3.34×10^5	98%	≈ 20.7
CFSE+Hoechst	150000	3.46×10^5	96%	≈ 19.9

1043

	Seeded cell number	Harvested cell number (after 24h)	Viability	Doubling time
Unstained	750000	2.8×10^6	90%	≈ 12.6
CFSE	750000	2.60×10^6	89%	≈ 13
Hoechst	750000	2.72×10^6	88%	≈ 12.9
CFSE + Hoechst	750000	2.65×10^6	90%	≈ 13.1

1044

1045 **Figure S7. The effects of Hoechst and CFSE staining on cell viability and proliferation rates.**
1046 Shown are the estimated doubling times (based on the increased in cell number after 24hrs
1047 growth) and measured viability (trypan blue) for fibroblasts **(A)** and ESCs **(B)**. The microscopy

1048   shows that stained cells maintain the correct morphology.

1049

1050

1051

1052   **Figure S8. The images of CFSE staining for both fast and slow proliferating fibroblasts and**

1053   **ESCs.**

1054   Fibroblasts **(A)** and ESCs **(B)** were stained by CFSE and sorted into two bins: fast proliferating

1055   cells (low CFSE) and slow proliferating cells (high CFSE).

1056

1057

Figure S9. Images of TMRE staining for fibroblasts and ESCs, showing heterogeneity.
Bright-field and TMRE staining images for both Fibroblasts and ESCs. Two pairs of ESC colonies of similar size but showing staining heterogeneity is circled in blue.

References

1.   Min, M. and S.L. Spencer, *Spontaneously slow-cycling subpopulations of human cells originate from activation of stress-response pathways.* PLOS Biology, 2019. **17**(3): p. e3000178.

2.   Wakamoto, Y., et al., *Dynamic persistence of antibiotic-stressed mycobacteria.* Science, 2013. **339**(6115): p. 91-5.

3.   Fridman, O., et al., *Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations.* Nature, 2014. **513**(7518): p. 418-21.

4.   Balaban, N.Q., et al., *A problem of persistence: still more questions than answers?* Nature Reviews Microbiology, 2013. **11**(8): p. 587-591.

5.   Gupta, P.B., et al., *Stochastic state transitions give rise to phenotypic equilibrium in populations of cancer cells.* Cell, 2011. **146**(4): p. 633-44.

6.   Brown, R., et al., *Poised epigenetic states and acquired drug resistance in cancer.* Nat Rev Cancer, 2014. **14**(11): p. 747-53.

7.   Marusyk, A., V. Almendro, and K. Polyak, *Intra-tumour heterogeneity: a looking glass for cancer?* Nat Rev Cancer, 2012. **12**(5): p. 323-34.

8.   van Dijk, D., et al., *Slow-growing cells within isogenic populations have increased RNA*

polymerase error rates and DNA damage. Nat Commun, 2015. **6**: p. 7972.

9.   Dhar, R., et al., *Single cell functional genomics reveals the importance of mitochondria in*
cell-to-cell phenotypic variation. eLife, 2019. **8**: p. e38904.

10.   Yaakov, G., et al., *Coupling phenotypic persistence to DNA damage increases genetic*
diversity in severe stress. Nat Ecol Evol, 2017. **1**(1): p. 16.

11.   Paek, A.L., et al., *Cell-to-Cell Variation in p53 Dynamics Leads to Fractional Killing.* Cell,
2016. **165**(3): p. 631-42.

12.   Brauer, M.J., et al., *Coordination of Growth Rate, Cell Cycle, Stress Response, and*
Metabolic Activity in Yeast. Molecular Biology of the Cell, 2007. **19**(1): p. 352-367.

13.   Regenberg, B., et al., *Growth-rate regulated genes have profound impact on*
interpretation of transcriptome profiling in Saccharomyces cerevisiae. Genome biology,
2006. **7**(11): p. R107-R107.

14.   Im, H.K., et al., *Mixed effects modeling of proliferation rates in cell-based models:*
consequence for pharmacogenomics and cancer. PLoS genetics, 2012. **8**(2): p.
e1002525-e1002525.

15.   Choy, E., et al., *Genetic Analysis of Human Traits In Vitro: Drug Response and Gene*
Expression in Lymphoblastoid Cell Lines. PLOS Genetics, 2008. **4**(11): p. e1000287.

16.   Cardoso-Moreira, M., et al., *Gene expression across mammalian organ development.*
Nature, 2019.

17.   Venet, D., J.E. Dumont, and V. Detours, *Most random gene expression signatures are*
significantly associated with breast cancer outcome. PLoS computational biology, 2011.
7(10): p. e1002240-e1002240.

18.   Levy, S.F., N. Ziv, and M.L. Siegal, *Bet hedging in yeast by heterogeneous, age-correlated*
expression of a stress protectant. PLoS Biol, 2012. **10**(5): p. e1001325.

19.   Johnston, I.G., et al., *Mitochondrial Variability as a Source of Extrinsic Cellular Noise.* PLOS
Computational Biology, 2012. **8**(3): p. e1002416.

20.   das Neves, R.P., et al., *Connecting variability in global transcription rate to mitochondrial*
variability. PLoS biology, 2010. **8**(12): p. e1000560-e1000560.

21.   Sukumar, M., et al., *Mitochondrial Membrane Potential Identifies Cells with Enhanced*
Stemness for Cellular Therapy. Cell Metabolism, 2016. **23**(1): p. 63-76.

22.   Mathieu, J. and H. Ruohola-Baker, *Metabolic remodeling during the loss and acquisition*
of pluripotency. Development, 2017. **144**(4): p. 541.

23.   Nichols, J. and A. Smith, *Naive and Primed Pluripotent States.* Cell Stem Cell, 2009. **4**(6): p.
487-492.

24.   Wray, J., T. Kalkan, and Austin G. Smith, *The ground state of pluripotency.* Biochemical
Society Transactions, 2010. **38**(4): p. 1027.

25.   Kolodziejczyk, Aleksandra A., et al., *Single Cell RNA-Sequencing of Pluripotent States*
Unlocks Modular Transcriptional Variation. Cell Stem Cell, 2015. **17**(4): p. 471-485.

26.   ter Huurne, M., et al., *Distinct Cell-Cycle Control in Two Different States of Mouse*
Pluripotency. Cell Stem Cell, 2017. **21**(4): p. 449-455.e4.

27.   Nair, G., et al., *Heterogeneous lineage marker expression in naive embryonic stem cells is*
mostly due to spontaneous differentiation. Scientific Reports, 2015. **5**: p. 13339.

28.   Abranches, E., et al., *Stochastic NANOG fluctuations allow mouse embryonic stem cells to*
explore pluripotency. Development (Cambridge, England), 2014. **141**(14): p. 2770-2779.

29.   Wytock, T.P. and A.E. Motter, *Predicting growth rate from gene expression.* Proc Natl
Acad Sci U S A, 2019. **116**(2): p. 367-372.

30.   Smith, A., *10 Embryonic Stem Cells.* Cold Spring Harbor Monograph Archive; Volume 40
(2001): Stem Cell Biology, 2001: p. 205-230.

31.   Cho, B.K., et al., *Homeostasis-stimulated proliferation drives naive T cells to differentiate*
directly into memory T cells. J Exp Med, 2000. **192**(4): p. 549-56.

32.   Parish, C.R., *Fluorescent dyes for lymphocyte migration and proliferation studies.*
Immunology & Cell Biology, 1999. **77**(6): p. 499-508.

33.   Weston, S.A. and C.R. Parish, *New fluorescent dyes for lymphocyte migration studies:*
Analysis by flow cytometry and fluorescence microscopy. Journal of Immunological
Methods, 1990. **133**(1): p. 87-97.

34.   Romano, P., et al., *Cell Line Data Base: structure and recent improvements towards*
molecular authentication of human cell lines. Nucleic acids research, 2009. **37**(Database
issue): p. D925-D932.

35.   Tamm, C., S. Pijuan Galitó, and C. Annerén, *A Comparative Study of Protocols for Mouse*
Embryonic Stem Cell Culturing. PLOS ONE, 2013. **8**(12): p. e81156.

36.   Ying, Q.-L., et al., *The ground state of embryonic stem cell self-renewal.* Nature, 2008.
453: p. 519.

37.   Hayashi, K., et al., *Dynamic equilibrium and heterogeneity of mouse pluripotent stem*
cells with distinct functional and epigenetic states. Cell stem cell, 2008. **3**(4): p. 391-401.

38.   Toyooka, Y., et al., *Identification and characterization of subpopulations in*
undifferentiated ES cell culture. Development, 2008. **135**(5): p. 909.

39.   Chambers, I., et al., *Nanog safeguards pluripotency and mediates germline development.*
Nature, 2007. **450**: p. 1230.

40.   Nair, G., et al., *Heterogeneous lineage marker expression in naive embryonic stem cells is*
mostly due to spontaneous differentiation. Scientific reports, 2015. **5**: p. 13339-13339.

41.   Subramanian, A., et al., *Gene set enrichment analysis: A knowledge-based approach for*
interpreting genome-wide expression profiles. Proceedings of the National Academy of
Sciences, 2005. **102**(43): p. 15545.

42.   Liberzon, A., et al., *Molecular signatures database (MSigDB) 3.0.* Bioinformatics (Oxford,
England), 2011. **27**(12): p. 1739-1740.

43.   Athanasiadou, R., et al., *Growth Rate-Dependent Global Amplification of Gene*
Expression. bioRxiv, 2016: p. 044735.

44.   Kumatori, A., et al., *Abnormally high expression of proteasomes in human leukemic cells.*
Proceedings of the National Academy of Sciences of the United States of America, 1990.
87(18): p. 7071-7075.

45.   Chen, L. and K. Madura, *Increased Proteasome Activity, Ubiquitin-Conjugating Enzymes,*
and eEF1A Translation Factor Detected in Breast Cancer Tissue. Cancer Research, 2005.
65(13): p. 5599.

46.   Arlt, A., et al., *Increased proteasome subunit protein expression and proteasome activity*
in colon cancer relate to an enhanced activation of nuclear factor E2-related factor 2
(Nrf2). Oncogene, 2009. **28**: p. 3983.

47.   Cetin, B. and D.W. Cleveland, *How to survive aneuploidy.* Cell, 2010. **143**(1): p. 27-29.

48.   Iadevaia, V., R. Liu, and C.G. Proud, *mTORC1 signaling controls multiple steps in*

1169   *ribosome biogenesis.* Semin Cell Dev Biol, 2014. **36**: p. 113-20.

1170   49.   Zhang, Y., et al., *Coordinated regulation of protein synthesis and degradation by*
1171   *mTORC1.* Nature, 2014. **513**(7518): p. 440-3.

1172   50.   Lempiainen, H. and D. Shore, *Growth control and ribosome biogenesis.* Curr Opin Cell
1173   Biol, 2009. **21**(6): p. 855-63.

1174   51.   Choi, J.-H., et al., *mTORC1 accelerates retinal development via the immunoproteasome.*
1175   Nature Communications, 2018. **9**(1): p. 2502.

1176   52.   Uprety, B., A. Kaja, and S.R. Bhaumik, *TOR Facilitates the Targeting of the 19S*
1177   *Proteasome Subcomplex To Enhance Transcription Complex Assembly at the Promoters*
1178   *of the Ribosomal Protein Genes.* Mol Cell Biol, 2018. **38**(14).

1179   53.   Yun, Y.S., et al., *mTORC1 Coordinates Protein Synthesis and Immunoproteasome*
1180   *Formation via PRAS40 to Prevent Accumulation of Protein Stress.* Mol Cell, 2016. **61**(4): p.
1181   625-639.

1182   54.   Zhang, Y., et al., *Rapamycin extends life and health in C57BL/6 mice.* The journals of
1183   gerontology. Series A, Biological sciences and medical sciences, 2014. **69**(2): p. 119-130.

1184   55.   Zhao, J., G.A. Garcia, and A.L. Goldberg, *Control of proteasomal proteolysis by mTOR.*
1185   Nature, 2016. **529**: p. E1.

1186   56.   Gearhart, J., E.E. Pashos, and M.K. Prasad, *Pluripotency redux--advances in stem-cell*
1187   *research.* N Engl J Med, 2007. **357**(15): p. 1469-72.

1188   57.   Dang, C.V., *MYC, metabolism, cell growth, and tumorigenesis.* Cold Spring Harbor
1189   perspectives in medicine. **3**(8): p. a014217.

1190   58.   *Drosophila myc Regulates Cellular Growth during Development.* Cell, 1999. **98**(6): p. 779
1191   - 790.

1192   59.   van Riggelen, J., A. Yetil, and D.W. Felsher, *MYC as a regulator of ribosome biogenesis*
1193   *and protein synthesis.* Nat Rev Cancer, 2010. **10**(4): p. 301-9.

1194   60.   Csibi, A., et al., *The mTORC1/S6K1 pathway regulates glutamine metabolism through the*
1195   *eIF4B-dependent control of c-Myc translation.* Curr Biol, 2014. **24**(19): p. 2274-80.

1196   61.   Liu, P., et al., *A functional mammalian target of rapamycin complex 1 signaling is*
1197   *indispensable for c-Myc-driven hepatocarcinogenesis.* Hepatology, 2017. **66**(1): p.
1198   167-181.

1199   62.   Riley, R.D., et al., *External validation of clinical prediction models using big datasets from*
1200   *e-health records or IPD meta-analysis: opportunities and challenges.* BMJ, 2016. **353**: p.
1201   i3140.

1202   63.   Waldman, Y.Y., T. Geiger, and E. Ruppin, *A genome-wide systematic analysis reveals*
1203   *different and predictive proliferation expression signatures of cancerous vs.*
1204   *non-cancerous cells.* PLoS Genet, 2013. **9**(9): p. e1003806.

1205   64.   Macfarlan, T.S., et al., *Embryonic stem cell potency fluctuates with endogenous retrovirus*
1206   *activity.* Nature, 2012. **487**(7405): p. 57-63.

1207   65.   Hicks, S.C., et al., *Missing data and technical variability in single-cell RNA-sequencing*
1208   *experiments.* Biostatistics, 2017. **19**(4): p. 562-578.

1209   66.   Packer, J.S., et al., *A lineage-resolved molecular atlas of C. elegans*
1210   *embryogenesis at single-cell resolution.* Science, 2019: p. eaax1971.

1211   67.   Sulston, J.E., et al., *The embryonic cell lineage of the nematode Caenorhabditis elegans.*
1212   Dev Biol, 1983. **100**(1): p. 64-119.

68. Altun, Z.F., Herndon, L.A., Wolkow, C.A., Crocker, C., Lints, R. and Hall, D.H. *WormAtlas.*
(ed.s) 2002-2020.

69. Petropoulos, S., et al., *Single-Cell RNA-Seq Reveals Lineage and X Chromosome*
Dynamics in Human Preimplantation Embryos. Cell, 2016. **167**(1): p. 285.

70. Deng, Q., et al., *Single-cell RNA-seq reveals dynamic, random monoallelic gene*
expression in mammalian cells. Science, 2014. **343**(6167): p. 193-6.

71. Samavarchi-Tehrani, P., et al., *Functional Genomics Reveals a BMP-Driven*
Mesenchymal-to-Epithelial Transition in the Initiation of Somatic Cell Reprogramming.
Cell Stem Cell, 2010. **7**(1): p. 64-77.

72. Li, R., et al., *A Mesenchymal-to-Epithelial Transition Initiates and Is Required for the*
Nuclear Reprogramming of Mouse Fibroblasts. Cell Stem Cell, 2010. **7**(1): p. 51-63.

73. Sun, H., et al., *Metabolic switch and epithelial-mesenchymal transition cooperate to*
regulate pluripotency. EMBO J, 2020. **39**(8): p. e102961.

74. Conn, C.S. and S.B. Qian, *Nutrient signaling in protein homeostasis: an increase in*
quantity at the expense of quality. Sci Signal, 2013. **6**(271): p. ra24.

75. Valvezan, A.J. and B.D. Manning, *Molecular logic of mTORC1 signalling as a metabolic*
rheostat. Nature Metabolism, 2019. **1**(3): p. 321-333.

76. Teslaa, T. and M.A. Teitell, *Pluripotent stem cell energy metabolism: an update.* The
EMBO journal, 2015. **34**(2): p. 138-153.

77. Lu, V. and M.A. Teitell, *Alpha-ketoglutarate: a "magic" metabolite in early germ cell*
development. 2019. **38**(1): p. e100615.

78. Tischler, J., et al., *Metabolic regulation of pluripotency and germ cell fate through*
α-ketoglutarate. 2019. **38**(1): p. e99518.

79. Brown, M., et al., *A recombinant murine retrovirus for simian virus 40 large T cDNA*
transforms mouse fibroblasts to anchorage-independent growth. Journal of virology,
1986. **60**(1): p. 290-293.

80. Minajigi, A., et al., *A comprehensive Xist interactome reveals cohesin repulsion and an*
RNA-directed chromosome conformation. Science, 2015: p. aab2276.

81. Lee, J.T. and N. Lu, *Targeted mutagenesis of Tsix leads to nonrandom X inactivation.* Cell,
1999. **99**(1): p. 47-57.

82. Bray, N.L., et al., *Erratum: Near-optimal probabilistic RNA-seq quantification.* Nat
Biotechnol, 2016. **34**(8): p. 888.

83. Leek, J.T., et al., *The sva package for removing batch effects and other unwanted*
variation in high-throughput experiments. Bioinformatics, 2012. **28**(6): p. 882-3.

84. Kolodziejczyk, A.A., et al., *Single Cell RNA-Sequencing of Pluripotent States Unlocks*
Modular Transcriptional Variation. Cell Stem Cell, 2015. **17**(4): p. 471-85.

85. Boroviak, T., et al., *The ability of inner-cell-mass cells to self-renew as embryonic stem*
cells is acquired following epiblast specification. Nat Cell Biol, 2014. **16**(6): p. 516-28.

86. Young, R.A., *Control of the embryonic stem cell state.* Cell, 2011. **144**(6): p. 940-954.

87. Crowley, L.C., M.E. Christensen, and N.J. Waterhouse, *Measuring Mitochondrial*
Transmembrane Potential by TMRE Staining. Cold Spring Harb Protoc, 2016. **2016**(12).

88. Liberzon, A., et al., *Molecular signatures database (MSigDB) 3.0.* Bioinformatics, 2011.
27(12): p. 1739-40.

89. Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*

interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43): p.
15545-50.

90.   Shannon, P., et al., *Cytoscape: a software environment for integrated models of*
biomolecular interaction networks. Genome Res, 2003. **13**(11): p. 2498-504.

91.   Merico, D., et al., *Enrichment map: a network-based method for gene-set enrichment*
visualization and interpretation. PLoS One, 2010. **5**(11): p. e13984.

92.   Reimand, J., et al., *Pathway enrichment analysis and visualization of omics data using*
g:Profiler, GSEA, Cytoscape and EnrichmentMap. Nat Protoc, 2019. **14**(2): p. 482-517.

93.   Durinck, S., et al., *BioMart and Bioconductor: a powerful link between biological*
databases and microarray data analysis. Bioinformatics, 2005. **21**(16): p. 3439-3440.

94.   Durinck, S., et al., *Mapping identifiers for the integration of genomic datasets with the*
R/Bioconductor package biomaRt. Nat Protoc, 2009. **4**(8): p. 1184-91.

95.   Airoldi, E.M., et al., *Steady-state and dynamic gene expression programs in*
Saccharomyces cerevisiae in response to variation in environmental nitrogen. Mol Biol
Cell, 2016. **27**(8): p. 1383-96.

96.   Slavov, N. and D. Botstein, *Coupling among growth rate response, metabolic cycle, and*
cell division cycle in yeast. Mol Biol Cell, 2011. **22**(12): p. 1997-2009.

97.   Barbie, D.A., et al., *Systematic RNA interference reveals that oncogenic KRAS-driven*
cancers require TBK1. Nature, 2009. **462**(7269): p. 108-12.

98.   Hanzelmann, S., R. Castelo, and J. Guinney, *GSVA: gene set variation analysis for*
microarray and RNA-seq data. BMC Bioinformatics, 2013. **14**: p. 7.

99.   Rohatgi, A., *WebPlotDigitizer.* 2017, Austin, Texas, USA.

1279