1 **Altering PRC2 activity partially suppresses *ddm1* mutant phenotypes in Arabidopsis**
2

3 Martin Rougée*, Leandro Quadrana*, Jérôme Zervudacki*, Vincent Colot, Lionel Navarro
4 and Angélique Deleris †
5

6 * These authors contributed equally to this work
7 † Correspondance : deleris@biologie.ens.fr
8

Institut de biologie de l'Ecole normale supérieure (IBENS), Ecole normale supérieure, CNRS,
INSERM, PSL Research University, 75005 Paris, France

Abstract

In plants and mammals, DNA methylation is a hallmark of transposable element (TE)
sequences that contributes to their epigenetic silencing. In contrast, histone H3 lysine 27
trimethylation (H3K27me3), which is deposited by the Polycomb Repressive Complex 2
(PRC2), is a hallmark of repressed genes. Nevertheless, there is a growing body of evidence for
a functional interplay between these pathways. In particular, many TE sequences acquire
H3K27me3 when they lose DNA methylation and it has been proposed that PRC2 can serve as
a back-up silencing system for hypomethylated TEs. Here, we describe in the flowering plant
Arabidopsis thaliana the gain of H3K27m3 at hundreds of TEs in the mutant *ddm1*, which is
defective in the maintenance of DNA methylation specifically over TE and other repeat
sequences. Importantly, we show that this gain depends solely on CURLY LEAF (CLF), which
is one of two otherwise partially redundant H3K27 methyltransferases active in vegetative
tissues. Finally, our results challenge the notion that PRC2 can be a compensatory silencing
system for hypomethylated TEs, as the complete loss of H3K27me3 in *ddm1 clf* double mutant
plants was not associated with further reactivation of TE expression nor with a burst of
transposition. Instead, and surprisingly, *ddm1 clf* plants exhibited less activated TEs, and a
chromatin recompaction as well as hypermethylation of linker DNA compared to *ddm1*. Thus,
we have described an unexpected genetic interaction between DNA methylation and Polycomb
silencing pathways, where a mutation in PRC2 does not aggravate the molecular phenotypes
linked to TE hypomethylation in *ddm1* but instead partially suppresses them.

Introduction

DNA methylation is an epigenetic mark involved in the stable silencing of transposable elements (TEs) as well as the regulation of gene expression in plants and mammals. When present over TE sequences, it is usually associated with the di- or trimethylation of histone H3 lysine 9 (H3K9me2/H3K9me3) and, in plants, positive feedback loops between the two marks exist, which maintain TE sequences in the heterochromatic state (Johnson *et al*, 2002; Du *et al*, 2015). In *A. thaliana*, DOMAINS REARRANGED METHYLASE 2 (DRM2) establishes DNA methylation in all three sequences contexts (i.e. CG, CHG and CHH) in a pathway referred to as RNA-directed DNA methylation (RdDM) that involves small RNAs (Cao & Jacobsen, 2002; Chan, 2004). Maintenance of DNA methylation over TEs is achieved by the combined and context-specific action of RdDM (CHH methylation), CHROMOMETHYLASES 2 and 3 (CMT2 and CMT3, for CHH and CHG methylation, respectively) (Zemach *et al*, 2013; Stroud *et al*, 2014) and METHYLTRANSFERASE1 (MET1) (CG methylation) (Kankel *et al*, 2003). In addition, the SNF2 family chromatin remodeler DECREASE IN DNA METHYLATION 1 (DDM1) is necessary for DNA methylation of most TE sequences in all cytosine sequence contexts.

In contrast to DNA methylation, histone H3 lysine 27 trimethylation (H3K27me3), which is targeted by the highly conserved Polycomb Group (PcG) proteins, in particular Polycomb Repressive Complex 2 (PRC2), is a hallmark of transcriptional repression of protein-coding and microRNA genes in plants as well as in animals (Marasca *et al*, 2018; Förderer *et al*, 2016; Chica *et al*, 2017; Prete *et al*, 2015) ; it is thought to act by promoting a local compaction of the chromatin that antagonizes the transcription machinery (Prete *et al*, 2015) in order to maintain transcriptional silencing (Holoch & Margueron, 2017). Thus, H3K27me3 and DNA methylation are generally considered as mutually exclusive chromatin marks.

Nonetheless, there is a growing body of evidence of an interplay between the two silencing pathways. In particular, in both plants and mammals, many TE sequences gain H3K27me3 upon their loss of DNA methylation (Mathieu *et al*, 2005; Deleris *et al*, 2012; Saksouk *et al*, 2014; Reddington *et al*, 2013). Moreover, in the filamentous Neurospora, H3K27me3 is redistributed from gene to TE-rich constitutive heterochromatin when the heterochromatic mark H3K9me3 or the protein complexes that bind to it are lost (Basenko *et al*, 2015).

The fact that H3K27me3 can mark TE sequences upon their demethylation led to the idea that PcG could serve as a back-up silencing system for hypomethylated TEs (Deleris *et al*, 2012). Consistent with this notion, subsequent work in mammals showed that H3K27me3 re-

established the repression of thousands of hypomethylated TEs in embryonic stem cells

subjected to rapid and extensive DNA demethylation (Walter *et al*, 2016).

In a previous study, we provided evidence supporting a role of PcG in the transcriptional

silencing of *EVD* (Zervudacki *et al*, 2018), an *A. thaliana* retroelement of the *ATCOPIA93*

family that is tightly controlled by DNA methylation and which transposes in plants mutated

for the chromatin remodeler *DDM1* (Tsukahara *et al*, 2009). We observed that silencing of *EVD*

is dependent on both DNA methylation and H3K27me3, which, at this locus, depends on the

SET-domain protein CURLY LEAF (CLF) (Zervudacki et al., 2018). Whether the dual control

observed at *EVD* is also present at other plant TEs is unknown.

In the present work, we integrated genetics, epigenomics and cell imaging to show that

numerous TEs gain H3K27me3 in response to *ddm1*-induced loss of DNA methylation. We

demonstrate also that this gain is mediated by CLF, with no role for the SET-domain H3K27

methyltransferase SWINGER (SWN), which is also active in vegetative tissues and otherwise

partially redundant with CLF (Chanvivattana *et al*, 2004; Wang *et al*, 2016; Yang *et al*, 2017).

Unexpectedly, the combination of *ddm1* and *clf* mutations was not associated with further

reactivation of TE expression or transposition as compared to *ddm1,* except for *EVD*. Instead

less TEs were expressed and we observed a partial recompaction of heterochromatin coupled

with DNA hypermethylation*,* prominently in linker DNA, in *ddm1 clf* versus *ddm1*. Together,

these results show that altering PRC2 activity partially suppresses ddm1 phenotypes.

Hundreds of hypomethylated TEs gain H3K27me3 in *ddm1*

We previously showed that in *met1* mutants impaired for CG methylation, hundreds of TEs

gain H3K27me3 methylation in Arabidopsis (Deleris *et al*, 2012). While *met1* and *ddm1*

mutants are both globally hypomethylated, the later, in contrast to the former, affects almost

exclusively TE and other repeat sequences, and in the three sequence contexts. Thus, the *ddm1*

mutant appeared to be a more relevant background to directly explore the interplay between

DNA methylation and Polycomb at TE sequences and we conducted a H3K27me3 ChIP-seq

experiment in this mutant. Like in *met1*, hundreds of TEs showed increased accumulation of

H3K27me3 in *ddm1* (**Figure 1A-C**). A stringent subset of 672 TEs showed no H3K27me3

ChIP-seq signal in wild-type plants and significantly gained H3K27me3 over their full length

in *ddm1* (**Figure 1D**). Moreover, the vast majority of those 672 TEs are located in

pericentromeric regions (**Figure 1E**) and were included in the subset of TEs that gain

H3K27me3 in *met1* (Deleris *et al*, 2012) (**Figure supplement 1A**). Four major TE super

families were overrepresented among the 672 TEs (LTR/Gypsy, LTR/Copia, DNA/En-Spm,

Figure 1. A mutation in *DDM1* leads to a gain of H3K27me3 on some heterochromatic TEs.
(**A**) Representative genome browser view of H3K27me3 levels in a 100 kb heterochromatic region of chromosome 1 in wild-type (WT) and two biological replicates (BR) of *ddm1* (blue bar: TE; orange bar: gene). (**B,C,D**) H3K27me3 enrichment in WT, *ddm1*-BR1 and *ddm1*-BR2 over TEs located in euchromatin (**B**), heterochromatin (**C**) and TEs with a significant positive fold change (FC) of H3K27me3 in *ddm1* (Pval<0,1 ; Log 2FC>2) (**D**). The upper graph shows the mean enrichment of H3K27me3 over a given TE subset; the corresponding heatmap below ranks them from top to bottom according to the average enrichment in all genotypes. (**E**) Chromosomal distribution of the 672 TEs that significantly gain H3K27me3 (Pval<0,1 ; Log2FC>2) and the 61 TEs that significantly lose H3K27me3 in *ddm1* (Pval<0,1 ; Log2FC<-2). (**F**) Table showing the occurences of Polycomb Response Elements (PRE) (Xiao et al., 2017) among the 672 TEs with a gain of H3K27me3 in *ddm1*. (**G**) Graph showing the proportion of TEs (TEs with a gain of H3K27me3 in *ddm1* compared to heterochromatic TEs) with a given PRE.

DNA/others) (**Figure supplement 1B**), suggesting the existence of sequence-specific targeting.

Indeed, PRC2 can be targeted to specific genes by the recognition of short sequence motifs

(Xiao et al., 2017). Consistent with this motif-based targeting, we detected a significant

enrichment of the Telobox, CTCC, GA-repeat and AC-rich motifs in the 672 TEs compared to

the rest of the heterochromatic TEs (**Figure 1F-G**). These results suggest the presence of an

instructive mechanism of PRC2 recruitment at TEs with particular motifs used as nucleation

sites. In addition, the detection of continuous blocks of H3K27me3, in particular on

chromosomes 1 and 4 (**Figure supplement 1C**), may suggest that, once nucleated, H3K27me3

domains spread over entire TE sequences and even beyond, into nearby TEs, in a fashion similar

to what was described at genic sequences (Yang *et al*, 2017; Wang *et al*, 2016).

The gain of H3K7me3 over hypomethylated TEs depends on CLF

To test whether CLF is required for the gain of H3K7me3 over TEs we performed a second set

of ChIP-seq experiments in *ddm1 clf*. Because total levels of H3K27me3 are strongly reduced

in both *clf* and *ddm1 clf* mutants (**Figure supplement 2A**) compared to wild-type, we spiked-

in exogenous Drosophila chromatin in the chromatin extracts for normalization (Orlando *et al*,

2014). Levels of H3K27me3 at genes were similar between *ddm1 clf* and *clf* mutant (**Figure**

supplement 2B), consistent with *DDM1* affecting TE and other repeat sequences specifically

(Lippman *et al*, 2004). Conversely, the gain of H3K27me3 observed over TEs in *ddm1* was

completely abolished in *ddm1 clf* (**Figure 2A-C**). Together, these results show that deposition

of H3K27me3 at most TEs in *ddm1* is fully dependent on *CLF*. This is in contrast with the well-

established, partial dependency of H3K27me3 deposition at genes on CLF (Wang *et al*, 2016;

Yang *et al*, 2017), which has been proposed to be due to a specialization of this factor in

mediating amplification and spreading of H3K27me3 marks after their establishment by SWN

(Yang *et al*, 2017). Thus, at least over *ddm1*-dependent hypomethylated TEs, CLF is required

for the nucleation, and perhaps, additionally, spreading of H3K27me3.

TE activation in *ddm1* is partially suppressed in *ddm1 clf*

We previously described the phenotype of *ddm1 clf* plants at the rosette stage (4.5 weeks-old).

By observing the plants at the seedling stage (15 days-old), we could observe additional

phenotypes in two out of three independent *ddm1 clf* lines (chlorosis in one line, growth arrest

in another) that segregated 1:3 and thus evoked the segregation of recessive mutations (**Figure**

3A). We previously showed that there is an enhanced accumulation of *ATCOPIA93* mRNAs in

ddm1 clf double mutant as compared to *ddm1* (Zervudacki et al., 2018), thus we hypothesized

Figure 2. The gain of H3K27me3 in *ddm1* is completely lost in *ddm1 clf*.
(**A**) Genome browser view of H3K27me3 levels on the same region as Figure 1 (**B**) H3K27me3 enrichment in *ddm1* and two *ddm1 clf* replicates over TEs with a H3K27me3 domain in *ddm1* with corresponding heatmap ranking TEs according to the H3K27me3 mean value. (**C**) Metagene representing H3K27me3 enrichment in *ddm1*-BR3, *ddm1 clf*-BR1 and *ddm1 clf*-BR2 of the 672 TEs previously shown to gain H3K27me3 in *ddm1.*

that these segregating phenotypes were caused by the transposition of TEs activated in *ddm1*

clf.

To test whether *ATCOPIA93* activity is increased and whether additional copies of this TE

family accumulate in *ddm1 clf*, we performed Southern blot analysis (**Figure 3B**) using a probe

specific of *ATCOPIA93 EVD/ATR* (*ATR* is an almost identical copy of *EVD*). In the wild-type

as well as the *ddm1* and *clf* single mutants, we could only observe one copy of *EVD* and *ATR*.

By contrast, we observed linear extrachromosomal DNA in all the *ddm1 clf* double mutant lines

tested (progenies of individual F2), consistent with previous results (Zervudacki *et al*, 2018).

In addition, in two out of three double mutant lines, we detected numerous additional bands

corresponding to *EVD/ATR* new insertions (**Figure 3B**), indicating that transposition occurs in

the double mutant, although to various extents which may reflect different dynamics of

mobilization following the initial event (Quadrana *et al*, 2019). Pyrosequencing performed on

the mutants showed that *ATCOPIA93* DNA (extrachromosomal and integrated) was mostly

contributed by *EVD* (**Figure 3C**). Thus, DNA methylation and H3K27me3 act in synergy to

negatively control *EVD* activity and prevent its transposition, in accordance with the dual

control they exert on its transcription (Zervudacki *et al*, 2018). To test whether more TEs are

activated in *ddm1 clf*, we performed TE-sequence capture (Quadrana *et al*, 2016), which

enables sensitive detection of new TE insertions at the genome wide-level, in about 100

seedlings of wild-type, *ddm1* (2nd generation inbred mutants) and *ddm1 clf* (F3 progenies). No

new insertions were detected for any of the 250 TEs tested in this manner, except for *EVD*,

which had accumulated more copies in *ddm1 clf* than in *ddm1* (**Figure 3D**)*,* although this

difference was not statistically significant in average — this was presumably because the *ddm1*

clf line (#1), where only ecDNA accumulates, was included in the experiment along lines #2

and #3 **(Figure supplement 3A**).

To investigate further the relationship between H3K27me3 and TE silencing, we performed

RNA-seq experiments in three and four biological replicates of *ddm1* and *ddm1 clf*,

respectively. In all *ddm1*, about a thousand TEs (929) were upregulated, the large majority of

which (865) did not show a significant gain of H3K27me3 marks in this background. We found

that a similar (*ddm1 clf*-BR3/4) or even lower (*ddm1 clf*-BR1/2) number of TEs were

transcriptionally active in *ddm1 clf* compared to *ddm1* (**Figure 3E**) in keeping with the lack of

enhanced transposition in *ddm1 clf*. Moreover, among the 1957 TEs that lose H3K27me3 in

ddm1 clf, we found only 14 TEs that were upregulated in all *ddm1 clf* lines versus *ddm1* (**Figure**

3F). Thus, with the exception of *EVD* case study, which supports our initial hypothesis of a

dual epigenetic control by DNA methylation and H3K27me3, it appears that *ddm1*-induced

Figure 3. Transposons activation in *ddm1* and *ddm1 clf*.
(**A**) Representative picture of 15 days-old *ddm1 clf* F3 seedlings coming from two independent F2 lines; the white arrows show two different phenotypes segregating 1:3, growth arrest (left) and chlorosis (right) respectively. (**B**) Southern blot analysis on a pool of leaves (8 rosettes for each genotype), using an *ATCOPIA93* probe that recognizes both *EVD* and *ATR* copies; epiRIL454 (Mari-Ordonez et al., 2013) was used as a positive control for activation of *ATCOPIA93* and three independent *ddm1 clf* lines were tested. (**C**) Qualitative analysis of genomic DNA by pyrosequencing. The position interrogated corresponds to the discriminating SNP between *EVD* (C/G) and *ATR* (A/T) and the % indicated represent the % of G (*EVD*, white bar) or T (*ATR*, grey bar) (**D**) Heatmap showing the transposition rates of 40 potentially mobile TEs as determined by TE-capture (Quadrana et al., 2016) using 100 seedlings for each genotype. (**E**) Table showing up-regulated and down-regulated TEs compared to WT as identified by RNA-seq analyses on seedlings. (**F**) Venn diagram showing the overlap between TEs with a loss of H3K27me3 in *ddm1 clf* and TEs up-regulated in the 4 different *ddm1 clf* lines.

activation of TEs is partially suppressed in *ddm1 clf,* with some heterogeneity between the lines

used.

<u>***ddm1 clf* displays chromatin recompaction and increased DNA methylation compared to**</u>

<u>***ddm1***</u>

To understand the mechanisms underlying the antagonism between DNA methylation and

CLF-dependent H3K27me3 deposition, we performed cytogenetic analyses on nuclei from

single and double mutants for *DDM1* and *CLF.* Chromocenters formation and DNA compaction

were unchanged in *clf* but strongly impacted in *ddm1* as previously published (Soppe *et al*,

2002). By contrast, they were only partially affected in the double mutant, indicating that

H3K27me3 loss induces chromatin recompaction specifically in the *ddm1* background (**Figure**

4A). In addition, immunofluorescence experiments showed that this chromatin recompaction

was associated with the recovery of H3K9me2 distribution (**Figure 4B**).

In order to assess whether chromatin recompaction associates with DNA methylation gain, we

compared the DNA methylomes of *ddm1* and *ddm1 clf* mutant plants. Consistent with the

cytogenetic analyses, hundreds of loci exhibited higher DNA methylation in all three sequence

contexts in *ddm1 clf* compared to *ddm1* (**Figure 5A**), with gains being most pronounced at

CHH sites and variable from one line to the other (**Figure 5B**). Furthermore, TEs with increased

CG methylation in *ddm1 clf* versus *ddm1* were slightly but statistically significantly less

expressed compared to those that did not show such an increase in this mutant background

(**Figure supplement 5A**).

To test whether H3K27me3 antagonizes *ddm1*-induced DNA methylation loss in *cis*, we

searched for differential methylated regions (DMRs) using non-overlapping 100bp windows.

There were few hyper DMRs and these tended to be inconsistent across replicates, suggesting

that they were the result of stochastic variation of DNA methylation (Becker *et al*, 2011).

Moreover, TEs showing clear DNA methylation gain in *ddm1 clf* compared to *ddm1* (**Figure**

5A) were not identified by our DMR detection. Visual inspection of these DNA

hypermethylated TEs revealed that increased DNA methylation was in fact confined to short

sequences (~20bp), typically separated by ~140 bp (**Figure supplement 5B**). Indeed, we

identified 1208 TE-specific short (20bp) sequences with higher CG methylation in *ddm1 clf*

compared to *ddm1*. These small-size hyper-DMRs were distributed over 759 TEs, most of

which (N= 571) are pericentromeric, and they were associated with a reduction of H3K27me3

levels (**Figure 5C,** H3K27me3 profiles centered on CG-DMRs). Among these 759 TEs

hypermethylated in *ddm1 clf* compared to *ddm1*, 61 showed a significant gain of H3K27me3 in

Figure 4. The double mutant *ddm1 clf* displays a partial recompaction of DNA and more visible H3K9me2 compared to *ddm1*.
(**A**) Representative picture of DAPI stained nuclei extracted from 50 fixed 10 days-old Arabidopsis seedlings for each genotype (three independent F3 lines from *ddm1 clf*) (**B**) Representative picture of DAPI staining and H3K9me2 immunodetection of nuclei extracted from 50 fixed 10 days-old Arabidopsis seedlings for each genotype.

ddm1: this is a significant 3.8 representation factor enrichment (p<3.644e^{-19}) compared to the

number of TEs in the genome that would be both CG-hypermethylated (in *ddm1 clf*) and

H3K27me3-marked (in *ddm1*) by chance. Although significant, this overlap is small, which

suggests that globally the antagonism between H3K27me3 and DNA methylation is indirect.

This could be explained by the impact of *ddm1 clf* double mutation on the transcriptome;

nevertheless we did not find any consistent expression changes for the components involved in

DNA methylation (**Figure supplement 5C**). Finally, the ~140 bp distance between DMRs at

discrete loci —which is roughly the size of nucleosomal DNA— prompted us to test whether

DNA hypermethylation in *ddm1 clf* versus *ddm1* occurred preferentially between nucleosomes.

Analysis of publicly available MNase data obtained for *ddm1* (Lyons & Zilberman, 2017)

indicated that the periodic DNA hypermethylation in *ddm1 clf* associate with linker DNA

(**Figure 5C**), consistent with *in vitro* and *in vivo* data showing that inter-nucleosomal DNA is

more accessible to DNA methyltransferases than DNA bound to the nucleosome core particle

(Lyons & Zilberman, 2017). These findings suggest therefore that CLF-dependent H3K27me3

marks, although indirectly, limit the accessibility of DNA methyltransferases to linker DNA.

Discussion

In this study, we have shown that hundreds of TEs gain H3K27me3 when hypomethylated by

ddm1. This phenomenon has two important mechanistic implications, the first one being that

DNA methylation can globally exclude H3K27me3. We and others previously showed that loss

of DNA methylation induced by another hypomethylation mutant, *met1*, rather than loss of

H3K9me2, allows for H3K27me3 deposition at TEs (Mathieu *et al*, 2005; Deleris *et al*, 2012).

However, another commonality between *met1* and *ddm1* mutants is the decompaction of their

DNA, which could also contribute to favor the access of PRC2 to chromatin, a possibility that

could be further investigated by identifying and using genetic backgrounds impaired for

chromatin compaction but not DNA methylation. The second mechanistic implication of our

observations is that PRC2 can be recruited to TEs. Here, we have characterized the transposons

marked by H3K27me3 in *ddm1* and found some motifs recently described in the genic targets

of PRC2 and functionally involved in the recruitment of this complex at genes (Xiao *et al*,

2017). Thus, while some TEs could be marked by H3K27me3 due to spreading from adjacent

PcG targets as it is the case for *EVD* (Zervudacki et al., 2018), the presence of these motifs as

well as the localization of many TEs far away from any H3K27me3 domain, suggests a

sequence-based, instructive mode of *cis*-recruitment for PRC2. Alternatively, this recruitment

could also be promoted by the enrichment of H2A.Z over demethylated TEs (Zilberman *et al*,

Figure 5. Periodic hypermethylation is observed in *ddm1 clf* compared to *ddm1*.
(**A**) Representative genome browser view of H3K27me3 and methylation of a TE (blue bar) showing hypermethylation in *ddm1 clf* compared to *ddm1*. (**B**) Meta-TEs showing average DNA methylation levels on all TEs, in the three different contexts, in all the genotypes of interest, with a zoom on *ddm1* and *ddm1 clf* methylated TEs. (**C**) Plot showing average DNA methylation, H3K27me3 and nucleosome occupancy (every peak in *ddm1* Mnase data shows the position of a nucleosome) 1Kb upstream ans downstream of "small" CG DMRs (20bp) in the three different sequence contexts.

2008) since a mechanistic link was found between H2A.Z and H3K27me3 (Carter *et al*, 2018).

These non mutually exclusive scenarios will deserve future investigation.

Subsequently, the compound mutant of DNA methylation and PRC2, *ddm1 clf*, uncovered

unexpected and interesting molecular phenotypes. First, PRC2 targeting of TEs in the *ddm1*

background depends almost completely on CLF. This requirement of CLF contrasts with the

partial dependency on CLF and SWN at H3K27me3 marked genes, where SWN plays a major

role, presumably in nucleating H3K27me3 (Yang *et al*, 2017). A possible explanation for this

result could be that the transcription factors implicated in the nucleation of H3K27me3 (Xiao

et al, 2017) at TEs in *ddm1* recruit a FIE-EMF2-PRC2 complex that does not contain SWN.

Secondly, from a functional point of view and rather unexpectedly, our results did not show

that PRC2 acts as a back-up silencing system in the absence of DNA methylation. Rather, they

are reminiscent of the observations made in Neurospora where the gain of H3K27me3 in

constitutive heterochromatin domains is not compensatory for gene silencing upon loss of

H3K9 methylation (or its reader HP1) but rather leads to growth defects and altered sensitivity

to genotoxic agents (Basenko *et al*, 2015). In our study, in *ddm1 clf*, the loss of H3K27me3 did

not enhance the number of *ddm1*-activated TEs. Instead, molecular phenotypes of partial *ddm1*

suppression were observed: same or lesser number of activated TEs and partial compaction of

DNA compared to *ddm1* associated with hypermethylation of the DNA in the three sequence

contexts, which presumably results in transcriptional TE silencing. Since the hypermethylation

also occurred at TEs that are non-targets of PcG, we could not conclude on a direct antagonism

between H3K27me3 and DNA methylation. Instead, we propose two possible mechanisms for

the global hypermethylation of *ddm1 clf* compared to *ddm1*. Firstly, it is likely that in *ddm1*

mutant, like in *met1* (Zilberman *et al*, 2008), H2AZ is incorporated at TEs. H2AZ is known to

be antagonistic to DNA methylation and given that a mutation of *CLF* leads to loss of H2AZ

(Carter *et al*, 2018), this may favor the reestablishment of DNA methylation. Another non-

exclusive possibility could be that the chromatin decompaction displayed by *ddm1* in the

presence of PRC2 globally restricts DNA remethylation which would occur upon recompaction

in *ddm1 clf* through unknown mecanisms. In addition, small RNAs could also participate in

either of these processes. What favors DNA compaction upon loss of H3K27me3, and,

accordingly what antagonizes chromocenters formation in *ddm1* in the presence of H3K27me3

is also unclear. It could be linked to the replacement of H3.1, the substrate for ATRX5 and 6

and H3K27 monomethylation (a modification associated with chromatin condensation) (Jacob

et al, 2009, 2010) by H3.3 and H3K27me3 in *ddm1*, and the subsequent reincorporation of H3.1

in the absence of CLF. In this respect, the *ddm1 clf* mutant represents a valuable system to test

the causal and functional relationship between DNA methylation and DNA compaction, which

is elusive. As for the possibility of a direct antagonism between H3K27me3 and DNA

methylation, we do not exclude it but the use of PcG mutants to analyze the impact of

H3K27me3 loss at specific loci currently makes it difficult to tear apart cis-effects of this loss

on a given TE from global and indirect effects. In the future, targeted removal of H3K27me3

by epigenome-editing needs to be implemented for this purpose.

Finally, our work in *ddm1 clf* mutant could have important implications for biological situations

where TEs are naturally hypomethylated, DDM1 absent and/or chromatin decompacted. First,

in the pollen vegetative nucleus, where hundreds of TEs are DNA-demethylated due to the

activity of DEMETER glycosylases (Ibarra *et al*, 2012), where DDM1 is not detected (Slotkin

et al, 2009). TE activation in this cell type is contributed by chromatin decondensation and

DNA demethylation-dependent and independent mechanisms (He *et al*, 2019) and has been

proposed to reinforce silencing in the gamete through production of small RNAs that could

transit into the sperm cell (Slotkin *et al*, 2009; Calarco *et al*, 2012; Ibarra *et al*, 2012; Martínez

et al, 2016). Whether PRC2 targets some of the demethylated TE-sequences and whether this

could antagonize chromatin decondensation and modulate TE activation and small RNA

production in this context is an open question. In this scenario, the mechanisms we have

described in this work could play an important role in the reinforcement of silencing in the

gametes and its modulation. Similarly, in the endosperm, the nutritive terminal tissue

surrounding the seed and derived from the central cell, the chromatin is less condensed than in

other types of nuclei (Baroux *et al*, 2007). In addition, hundreds of TEs of the maternal genome

are naturally hypomethylated forming primary imprints and many of these hypomethylated TEs

are targeted by PRC2 forming secondary imprints (Weinhofer *et al*, 2010; Rodrigues &

Zilberman, 2015). These secondary imprints cause, for instance, the silencing of the maternal

PHERES locus while the cognate paternal allele is activated (Makarevich *et al*, 2008; Rodrigues

& Zilberman, 2015). The interplay between DNA methylation and PRC2 that we have

evidenced could thus be particularly relevant in this cell type and modulate the imprinting of

some genes. Finally, an exciting possibility is that PRC2 could target transposons after their

mobilization and integration and this could slow down the establishment of their DNA

methylation-based transgenerational and stable epigenetic silencing.

Material and methods

Plant material and growth condition

14 day-old seedlings grown on MS plates and with an 8h light/16h dark photoperiod were used throughout the study except for Southern Blot where analyses were performed on 5-week-old rosette leaves grown in the same conditions.

Mutant lines

We used the *ddm1-2* allele (Vongs *et al*, 1993) and the *clf-29* allele (Bouveret *et al*, 2006). Double *ddm1 clf* mutants were generated by crossing the above-mentioned mutants (Zervudacki *et al*, 2018). Experiments were performed on F3 progenies using *ddm1*-2 2nd generation plants as controls.

SDS-PAGE and Western blotting

Chromatin-enriched protein fraction was extracted from 10-days-old seedlings as described in (Bourbousse *et al*, 2012), quantified by standard BCA assay and 20 µg were resolved on SDS/PAGE. After electroblotting the proteins on a Polyvinylidene difluoride (PVDF) membrane, H3K27me3 analysis was performed using an antibody against the H3K27me3 (Millipore, 07-449) at a 1:1000 dilution and a secondary antibody against rabbit coupled to HRP (Promega W4011) at a 1:20 000 dilution.

BS-seq libraries

DNA was extracted using a standard CTAB protocol. Bisulfite conversion, BS-seq libraries and sequencing (paired-end 100 nt reads) were performed by BGI Tech Solutions (Hong Kong). Adapter and low-quality sequences were trimmed using Trimming Galore v0.3.3. Mapping was performed on TAIR10 genome annotation using Bismark v0.14.2 (Krueger & Andrews, 2011) and the parameters: --bowtie2, -N 1, -p 3 (alignment); --ignore 5 --ignore_r2 5 -- ignore_3prime_r2 1 (methylation extractor). Only uniquely mapping reads were retained. The methylKit package v0.9.4 (Akalin *et al*, 2012) was used to calculate differential methylation in 100 bp or 20bp non-overlapping windows (DMRs). Significance of calculated differences was determined using Fisher's exact test and Benjamin-Hochberg (BH) adjustment of p-values (FDR<0.05) and methylation difference cutoffs of 40% for CG, 20% for CHG and 20% for CHH. Differentially methylated windows within 100bp or 20bp of each other were merged to

form larger DMRs. 100bp windows with at least six cytosines covered by a minimum of 6 (CG and CHG) and 10 (CHH) reads in all libraries were considered.

RNA extraction and 3'quant-seq (RNA) analyses

Total RNA was extracted using Trizol followed by clean-up on RNeasy Plant Mini kit columns (Macherey-Nagel). 3'end librairies were prepared using the QuantSeq 3' Fwd library prep kit (Lexogen). Libraries were sequenced to acquire 150 bp-reads on a NextSeq Mid output flow cell (Fasteris, Geneva). Expression level was calculated by mapping reads using STAR v2.5.3a (Dobin *et al*, 2013) on the *A. thaliana* reference genome (TAIR10) with the following arguments --outFilterMultimapNmax 50 --outFilterMatchNmin 30 --alignSJoverhangMin 3 -- alignIntronMax 10000. Counts were normalized and annotations (genes and TEs) were declared differentially expressed between samples using DESeq2 (Love *et al*, 2014).

Chromatin immunoprecipitation and ChIP-seq analyses

ChIP experiments were conducted in two to four biological replicates of WT, *ddm1*, *clf* and *ddm1 clf* using an anti-H3K27me3 antibody (Millipore, 07-449). For each biological replicate, two IPs were carried out using 80 μg of Arabidopsis chromatin mixed with 5 μg of Drosophila chromatin, as quantified using BiCinchoninic Acid assay (Thermo Fisher Scientific). DNA eluted and purified from the two technical replicates was pooled before library preparation (Illumina TruSeq ChIP) and sequencing (Illumina sequencing single-reads, 1 × 50 bp) of the resulting input and IP samples performed by Fasteris (Geneva, Switzerland).

Reads were mapped using Bowtie2 v2.3.2 (Langmead & Salzberg, 2012) onto TAIR10 *Arabidopsis thaliana* and Drosophila melanogaster (dm6) genomes. Normalization factor (Rx) for each sample using spiked-in *D. melanogaster* chromatin was calculated (Nassrallah *et al*, 2018) using the following formula Rx = r/Nd_IP, where Nd_IP corresponds to the number of reads mapped on *D. melanogaster* genome in the IP and r corresponds to the percentage of *Drosophila*-derived reads in the input. Genomic regions significantly marked by H3K27me3 were identify using MACS2 (Zhang *et al*, 2008) and genes or TEs overlapping these regions were obtained using bedtools (Quinlan & Hall, 2010). The number of reads over marked genes or TEs were normalized by applying the normalization factor and differentially marked genes between samples were calculated using DESeq2 (Love *et al*, 2014).

TE-sequence capture

TE sequence capture was performed on around 100 seedlings in all cases except *ddm1-clf*#23 NOchl were only 12 seedlings were recovered. Seedlings were grown in plates under control (long-day) conditions and genomic DNA was extracted using the CTAB method (Murray & Thompson, 1980). Libraries were prepared as previously described (Quadrana *et al*, 2019) using 1µg of DNA and TruSeq paired-end kit (Illumina) following manufacturer instructions. Libraries were then amplified through 7 cycles of ligation-mediated PCR using the KAPA HiFi Hot Start Ready Mix and primers AATGATACGGCGACCACCGAGA and CAAGCAGAAGACGGCATACGAG at a final concentration of 2µM. 1µg of multiplexed libraries were then subjected to TE-sequence capture exactly as previously reported (Quadrana *et al*, 2016). Pair-end sequencing was performed using one lane of Illumina NextSeq500 and 75bp reads. About 42 million pairs were sequenced per library and mapped to the TAIR10 reference genome using Bowtie2 v2.3.2 (Langmead & Salzberg, 2012) with the arguments --mp 13 --rdg 8,5 --rfg 8,5 --very-sensitive. An improved version of SPLITREADER (available at https://github.com/LeanQ/SPLITREADER) was used to detect new TE insertions. Briefly, split-reads as well as discordant reads mapping partially on reference and consensus TE sequences (obtained from RepBase update) were identified, soft clipped and remapped to the TAIR10 reference genome using Bowtie2 (Langmead & Salzberg, 2012). Putative insertions supported by at least one split- and/or discordant-reads at each side of the insertion sites were retained. Insertions spanning centromeric repeats or coordinates spanning the corresponding donor TE sequence were excluded. In addition, putative TE insertions detected in more than one library were excluded to retain only sample-specific TE insertions.

Southern blot

DNA from 5 week-old rosette leaves was extracted using a standard CTAB protocol. 1,5 µg of genomic DNA was digested overnight with SSpI restriction enzyme. The digestion was run on a 1% agarose gel, transferred to Hybond N+ membranes, blocked and washed according to manufacturer instructions (GE Healthcare). Membranes were probed with a PCR product (corresponding to a fragment of *EVD* GAG sequence,) radiolabeled with alpha 32P-dCTP using the Megaprime DNA Labeling System. EVD PCR product was generated with the same primers as in (Marí-Ordóñez *et al*, 2013).

Pyrosequencing

ATCOPIA93 DNA from gDNA (same extraction as used in Southern Blot) was analyzed as in (Zervudacki *et al*, 2018).

Cytology

DAPI staining on fixed nuclei was performed as described in Bourbousse et al., 2015 using 10-days-old chopped cotyledons to avoid developmental defects due to accumulation of transposition. For immuno-detection, a primary antibody against H3K9me2 (Abcam 1220) at a 1:200dilution was used, followed by a secondary antibody (Alexa Fluor 488 mouse) at a 1:400 dilution and DAPI staining. Images were acquired with a confocal laser scanning microscope and processed using ImageJ (https://imagej.nih.gov/ij/).

Data availability

Sequencing data has been deposited in the European Nucleotide Archive (ENA) under project PRJEB34363.

Acknowledgments

We thank the Genomics and Informatics facilities at IBENS as well as the Barneche lab (C. Bourbousse and G. Teano) for their help with Drosophila spike-in ChIP experiments and discussions. This work was funded by a Human Frontier Scientific Program Career Development Award (HFSP-CDA-00018/2014) granted to A.D and the Centre National de la Recherche Scientifique (CNRS) Momentum program granted to L.Q. Additional support was from the CNRS.

Conflict of interest: The authors declare that they have no conflict of interest.

Sup Fig.1A Overlap between TEs marked by H3K27me3 in *met1* (Deleris et al., 2012) and TEs with a gain of H3K27me3 in *ddm1*.

Sup Fig.1B Distribution of the different TE families among the TEs that significantly gain H3K27me3 in *ddm1* compared to the family distribution of all TEs.

Sup Fig.1C Genome browser view of chromosomes 1 and 4 showing the two largest H3K27me3 domains in *ddm1* (BR=biological replicate).

Sup Fig.2A Immuno detection of H3K27me3 in chromatin extracts (H4 is used as a loading control), two biological replicates are shown.

H3K27me3 profiles on genes marked at least in one genotypes (N=6571)

Sup Fig.2B Metagene showing H3K27me3 levels on genes marked in at least one genotype

Number of *ATCOPIA93* neo-insertions per 100 plants

Sup Fig.3A Number of *EVD* neo-insertions revealed by TE-capture in several lines per genotype, all derived from a cross between *ddm1* and *clf* mutants.

Sup Fig.5A Graph showing expression changes of TE hypermethylated or not in *ddm1 clf* in different genetic backgrounds.

Sup Fig.5B Representative view of a TE with a periodic DNA hypermethylation every 140bp; the bottom panel is a zoom of the top pane and the double arrow represents a distance of 140bp.

		WT *ddm1clf*-BR4		WT *ddm1clf*-BR3		WT *ddm1clf*-BR1		WT *ddm1clf*-BR2	
		log2FoldChange	padj	log2FoldChange	padj	log2FoldChange	padj	log2FoldChange	padj
AT5G49160	MET1	0,004	1,000	0,65	0,928	-0,20	1,000	-0,69	1,000
AT5G15380	DRM1	NA	NA	NA	NA	NA	NA	NA	NA
AT5G14620	DRM2	-0,588	1,000	-1,25	0,714	-0,01	1,000	0,34	1,000
AT1G69770	CMT3	1,761	0,312	1,80	0,215	-0,02	1,000	-0,75	1,000
AT4G19020	CMT2	0,317	1,000	0,82	0,898	-0,24	1,000	-0,71	1,000
AT1G06760	H1.1	-0,16	1	-0,15	1	0,06	1	0,08	1
AT2G30620	H1.2	1,07	0,8006537	0,22	1	0,21	1	0,83	1

Sup Fig.5C Table showing expression changes of genes involved in DNA methylation in all *ddm1 clf* compared to WT

References

Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A & Mason CE (2012) MethylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol.* **13:** R87 Available at: http://genomebiology.com/2012/13/10/R87

Baillie JK, Barnett MW, Upton KR, Gerhardt DJ, Richmond TA, Sapio F De, Brennan P, Rizzu P, Smith S, Fell M, Talbot RT, Gustincich S, Freeman TC, Mattick JS, Hume DA, Heutink P, Carninci P, Jeddeloh JA & Faulkner GJ (2011) Somatic retrotransposition alters the genetic landscape of the human brain. *Nature* **479:** 534–537 Available at: http://dx.doi.org/10.1038/nature10531

Baroux C, Pecinka A, Fuchs J, Schubert I & Grossniklaus U (2007) The Triploid Endosperm Genome of Arabidopsis Adopts a Peculiar, Parental-Dosage-Dependent Chromatin Organization. *Plant Cell* **19:** 1782 LP – 1794 Available at: http://www.plantcell.org/content/19/6/1782.abstract

Basenko EY, Sasaki T, Ji L, Prybol CJ, Burckhardt RM, Schmitz RJ & Lewis ZA (2015) Genome-wide redistribution of H3K27me3 is linked to genotoxic stress and defective growth. *Proc. Natl. Acad. Sci.* **112:** E6339–E6348

Becker, Claude Koenig D, Stegle O, Borgwardt K & Weigel D (2011) thaliana methylome.

Bourbousse C, Ahmed I, Roudier F, Zabulon G, Blondet E, Balzergue S, Colot V, Bowler C & Barneche F (2012) Histone H2B Monoubiquitination Facilitates the Rapid Modulation of Gene Expression during Arabidopsis Photomorphogenesis. *PLOS Genet.* **8:** e1002825 Available at: https://doi.org/10.1371/journal.pgen.1002825

Bouveret R, Schönrock N, Gruissem W & Hennig L (2006) Regulation of flowering time by Arabidopsis MSI1. *Development* **133:** 1693–1702

Calarco JP, Borges F, Donoghue MTA, Van Ex F, Jullien PE, Lopes T, Gardner R, Berger F, Feijó JA, Becker JD & Martienssen RA (2012) Reprogramming of DNA methylation in pollen guides epigenetic inheritance via small RNA. *Cell* **151:** 194–205

Cao X & Jacobsen SE (2002) Role of the Arabidopsis DRM methyltransferases in de novo DNA methylation and gene silencing. *Curr. Biol.* **12:** 1138–1144

Carter B, Bishop B, Ho K, Huang R, Jia W, Zhang H & Pascuzzi PE (2018) The Chromatin Remodelers PKL and PIE1 Act in an Epigenetic Pathway That Determines H3K27me3 Homeostasis in Arabidopsis. **30:** 1337–1352

Chan SW-L (2004) RNA Silencing Genes Control de Novo DNA Methylation. *Science.* **303:**

1336–1336 Available at: http://www.sciencemag.org/cgi/doi/10.1126/science.1095989

Chanvivattana Y, Bishopp A, Schubert D, Stock C, Moon Y, Sung ZR & Goodrich J (2004)

Interaction of Polycomb-group proteins controlling flowering in Arabidopsis. **:** 5263–

5276

Chica C, Louis A, Roest Crollius H, Colot V & Roudier F (2017) Comparative epigenomics

in the Brassicaceae reveals two evolutionarily conserved modes of PRC2-mediated gene

regulation. *Genome Biol.* **18:** 1–15

Deleris A, Stroud H, Bernatavichute Y, Johnson E, Klein G, Schubert D & Jacobsen SE

(2012) Loss of the DNA Methyltransferase MET1 Induces H3K9 Hypermethylation at

PcG Target Genes and Redistribution of H3K27 Trimethylation to Transposons in

Arabidopsis thaliana. *PLoS Genet.* **8:** e1003062 Available at:

http://dx.plos.org/10.1371/journal.pgen.1003062

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M &

Gingeras TR (2013) STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29:** 15–

21

Du J, Johnson LM, Jacobsen SE & Patel DJ (2015) DNA methylation pathways and their

crosstalk with histone methylation. *Nat. Rev. Mol. Cell Biol.* **16:** 519–532 Available at:

http://www.nature.com/doifinder/10.1038/nrm4043

Förderer A, Zhou Y & Turck F (2016) The age of multiplexity: Recruitment and interactions

of Polycomb complexes in plants. *Curr. Opin. Plant Biol.* **29:** 169–178

Holoch D & Margueron R (2017) Mechanisms Regulating PRC2 Recruitment and Enzymatic

Activity. *Trends Biochem. Sci.* **42:** 531–542 Available at:

http://dx.doi.org/10.1016/j.tibs.2017.04.003

Ibarra CA, Feng X, Schoft VK, Hsieh TF, Uzawa R, Rodrigues JA, Zemach A, Chumak N,

Machlicova A, Nishimura T, Rojas D, Fischer RL, Tamaru H & Zilberman D (2012)

Active DNA demethylation in plant companion cells reinforces transposon methylation

in gametes. *Science.* **337:** 1360–1364

Ingouff M, Rademacher S, Holec S, Xin N, Readshaw A & Foo SH (2010) Report Zygotic

Resetting of the HISTONE 3 Variant Repertoire Participates in Epigenetic

Reprogramming in Arabidopsis. **:** 2137–2143

Jacob Y, Feng S, LeBlanc CA, Bernatavichute Y V, Stroud H, Cokus S, Johnson LM,

Pellegrini M, Jacobsen SE & Michaels SD (2009) ATXR5 and ATXR6 are H3K27

monomethyltransferases required for chromatin structure and gene silencing. *Nat. Struct.*

Mol. Biol. **16:** 763–768 Available at: https://www.ncbi.nlm.nih.gov/pubmed/19503079

505 Jacob Y, Stroud H, Leblanc C, Feng S, Zhuo L, Caro E, Hassel C, Gutierrez C, Michaels SD

506   & Jacobsen SE (2010) Regulation of heterochromatic DNA replication by histone H3

507   lysine 27 methyltransferases. *Nature* **466:** 987–991

508 Johnson L, Cao X & Jacobsen S (2002) Interplay between two epigenetic marks. DNA

509   methylation and histone H3 lysine 9 methylation. *Curr. Biol.* **12:** 1360–1367

510 Kankel MW, Ramsey DE, Stokes TL, Flowers SK, Haag JR, Jeddeloh JA, Riddle NC,

511   Verbsky ML & Richards EJ (2003) Arabidopsis MET1 cytosine methyltransferase

512   mutants. *Genetics* **163:** 1109–1122

513 Krueger F & Andrews SR (2011) Bismark: A flexible aligner and methylation caller for

514   Bisulfite-Seq applications. *Bioinformatics* **27:** 1571–1572

515 Langmead B & Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat. Methods*

516   **9:** 357–9

517 Li Y, Zheng H, Wang Q, Zhou C, Wei L, Liu X, Zhang W, Zhang Y, Du Z, Wang X & Xie

518   W (2018) Genome-wide analyses reveal a role of Polycomb in promoting

519   hypomethylation of DNA methylation valleys. *Genome Biol.* **19:** 1–16

520 Lippman Z, Gendrel A-V, Black M, Vaughn MW, Dedhia N, Richard McCombie W, Lavine

521   K, Mittal V, May B, Kasschau KD, Carrington JC, Doerge RW, Colot V & Martienssen

522   R (2004) Role of transposable elements in heterochromatin and epigenetic control.

523   *Nature* **430:** 471–476 Available at:

524   http://www.nature.com/doifinder/10.1038/nature02651

525 Love MI, Huber W & Anders S (2014) Moderated estimation of fold change and dispersion

526   for RNA-seq data with DESeq2. *Genome Biol.* **15:** 550

527 Lyons DB & Zilberman D (2017) DDM1 and Lsh remodelers allow methylation of DNA

528   wrapped in nucleosomes. **:** 1–20

529 Makarevich G, Villar CBR, Erilova A & Köhler C (2008) Mechanism of PHERES1

530   imprinting in Arabidopsis. **1:** 906–912

531 Marasca F, Bodega B & Orlando V (2018) How Polycomb-Mediated Cell Memory Deals

532   With a Changing Environment. *BioEssays* **40:** 1700137 Available at:

533   https://doi.org/10.1002/bies.201700137

534 Marí-Ordóñez A, Marchais A, Etcheverry M, Martin A, Colot V & Voinnet O (2013)

535   Reconstructing de novo silencing of an active plant retrotransposon. *Nat. Genet.* **45:**

536   1029–1039 Available at: http://www.nature.com/doifinder/10.1038/ng.2703

537 Martínez G, Panda K, Köhler C & Slotkin RK (2016) movement from the surrounding nurse

538   cell. *Nat. Plants* **2:** 1–8 Available at: http://dx.doi.org/10.1038/nplants.2016.30

Mathieu O, Probst A V & Paszkowski J (2005) Distinct regulation of histone H3 methylation

at lysines 27 and 9 by CpG methylation in Arabidopsis. *EMBO J.* **24:** 2783–2791

Available at: http://emboj.embopress.org/cgi/doi/10.1038/sj.emboj.7600743

Murray G & Thompson WF (1980) Rapid isolation of high molecular weight plant DNA.

Nucleic Acids Res. **8:** 4321–4326

Nassrallah A, Rougée M, Bourbousse C, Drevensek S, Fonseca S, Iniesto E, Ait-Mohamed O,

Deton-Cabanillas AF, Zabulon G, Ahmed I, Stroebel D, Masson V, Lombard B,

Eeckhout D, Gevaert K, Loew D, Genovesio A, Breyton C, de Jaeger G, Bowler C, et al

(2018) DET1-mediated degradation of a SAGA-like deubiquitination module controls

H2Bub homeostasis. *Elife* **7:** 1–29

Orlando DA, Chen MW, Bradner JE, Guenther MG, Orlando DA, Chen MW, Brown VE,

Solanki S, Choi YJ & Olson ER (2014) Quantitative ChIP-Seq Normalization Reveals

Global Modulation of the Epigenome Resource Quantitative ChIP-Seq Normalization

Reveals Global Modulation of the Epigenome. *CellReports* **9:** 1163–1170 Available at:

http://dx.doi.org/10.1016/j.celrep.2014.10.018

Prete S, Mikulski P, Schubert D & Gaudin V (2015) One, Two, Three: Polycomb Proteins Hit

All Dimensions of Gene Regulation. **:** 520–542

Quadrana L, Bortolini Silveira A, Mayhew GF, LeBlanc C, Martienssen RA, Jeddeloh JA &

Colot V (2016) The Arabidopsis thaliana mobilome and its impact at the species level.

Elife **5:** 1–25 Available at: http://elifesciences.org/lookup/doi/10.7554/eLife.15716

Quadrana L, Etcheverry M, Gilly A, Caillieux E, Madoui M-A, Guy J, Bortolini Silveira A,

Engelen S, Baillet V, Wincker P, Aury J-M & Colot V (2019) Transposition favors the

generation of large effect mutations that may facilitate rapid adaption. *Nat. Commun.* **10:**

1–10 Available at: http://dx.doi.org/10.1038/s41467-019-11385-5

Quinlan AR & Hall IM (2010) BEDTools: A flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26:** 841–842

Reddington JP, Perricone SM, Nestor CE, Reichmann J, Youngson NA, Suzuki M, Reinhardt

D, Dunican DS, Prendergast JG, Mjoseng H, Ramsahoye BH, Whitelaw E, Greally JM,

Adams IR, Bickmore WA & Meehan RR (2013) Redistribution of H3K27me3 upon

DNA hypomethylation results in de-repression of Polycomb target genes. *Genome Biol.*

14: R25 Available at: http://genomebiology.biomedcentral.com/articles/10.1186/gb-

2013-14-3-r25

Rodrigues JA & Zilberman D (2015) Evolution and function of genomic imprinting in plants.

: 2517–2531

Saksouk N, Barth TK, Ziegler-Birling C, Olova N, Nowak A, Rey E, Mateos-Langerak J,

Urbach S, Reik W, Torres-Padilla ME, Imhof A & Déjardin J (2014) Redundant

Mechanisms to Form Silent Chromatin at Pericentromeric Regions Rely on BEND3 and

DNA Methylation. *Mol. Cell* **56:** 580–594

She W, Grimanelli D & Baroux C (2014) An efficient method for quantitative , single-cell

analysis of chromatin modification and nuclear architecture in whole-mount ovules in

arabidopsis An Efficient Method for Quantitative , Single-cell Analysis of Chromatin

Modification and Nuclear Architect. **:** 1–9

Shengbo He, Martin Vickers, Jingyi Zhang XF (2019) Natural depletion of H1 in sex cells

causes DNA demethylation, heterochromatin decondensation and transposon activation.

*bioRxiv***:** 1–11 Available at:

https://www.biorxiv.org/content/biorxiv/early/2018/10/26/451930.full.pdf

Slotkin RK, Vaughn M, Borges F, Tanurdžić M, Becker JD, Feijó JA & Martienssen RA

(2009) Epigenetic Reprogramming and Small RNA Silencing of Transposable Elements

in Pollen. *Cell* **136:** 461–472

Soppe WJJ, Jasencakova Z, Houben A, Kakutani T, Meister A, Huang MS, Jacobsen SE,

Schubert I & Fransz PF (2002) DNA methylation controls histone H3 lysine 9

methylation and heterochromatin assembly in Arabidopsis. **21:** 6549–6559

Stroud H, Do T, Du J, Zhong X, Feng S, Johnson L, Patel DJ & Jacobsen SE (2014) Non-CG

methylation patterns shape the epigenetic landscape in Arabidopsis. *Nat. Struct. Mol.*

Biol. **21:** 64–72 Available at: http://www.ncbi.nlm.nih.gov/pubmed/24336224

Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A & Kakutani T (2009) Bursts of

retrotransposition reproduced in Arabidopsis. *Nature* **461:** 423–426 Available at:

http://www.nature.com/doifinder/10.1038/nature08351

Vongs A, Kakutani T, Martienssen RA & Richardstt EJ (1993) Arabidopsis thaliana DNA

Methylation Mutants. *Science.* **260:** 1926–1928

Walter M, Teissandier A, Pérez-Palacios R & Bourc'His D (2016) An epigenetic switch

ensures transposon repression upon dynamic loss of DNA methylation in embryonic

stem cells. *Elife* **5:** 1–30

Wang H, Liu C, Cheng J, Liu J, Zhang L, He C, Shen WH, Jin H, Xu L & Zhang Y (2016)

Arabidopsis Flower and Embryo Developmental Genes are Repressed in Seedlings by

Different Combinations of Polycomb Group Proteins in Association with Distinct Sets of

Cis-regulatory Elements. *PLoS Genet.* **12:** 1–25

Weinhofer I, Hehenberger E, Roszak P, Hennig L & K??hler C (2010) H3K27me3 profiling

of the endosperm implies exclusion of polycomb group protein targeting by DNA

methylation. *PLoS Genet.* **6:** 1–14

Xiao J, Jin R, Yu X, Shen M, Wagner JD, Pai A, Song C, Zhuang M, Klasfeld S, He C,

Santos AM, Helliwell C, Pruneda-Paz JL, Kay SA, Lin X, Cui S, Garcia MF, Clarenz O,

Goodrich J, Zhang X, et al (2017) Cis and trans determinants of epigenetic silencing by

Polycomb repressive complex 2 in Arabidopsis. *Nat. Genet.* **49:** 1546–1552 Available at:

http://www.nature.com/doifinder/10.1038/ng.3937

Yang H, Yang H, Berry S, Olsson TSG, Hartley M, Howard M & Dean C (2017) Distinct

phases of Polycomb silencing to hold epigenetic memory of cold in Arabidopsis.

Science. **1121:** 1–9

Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, Harmer SL &

Zilberman D (2013) The arabidopsis nucleosome remodeler DDM1 allows DNA

methyltransferases to access H1-containing heterochromatin. *Cell* **153:** 193–205

Available at: http://dx.doi.org/10.1016/j.cell.2013.02.033

Zervudacki J, Yu A, Amesefe D, Wang J, Drouaud J, Navarro L & Deleris A (2018)

Transcriptional control and exploitation of an immune-responsive family of plant

retrotransposons. *EMBO J.* **37:** e98482 Available at:

http://emboj.embopress.org/lookup/doi/10.15252/embj.201798482

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers

RM, Brown M, Li W & Shirley XS (2008) Model-based analysis of ChIP-Seq (MACS).

Genome Biol. **9:**

Zilberman D, Coleman-Derr D, Ballinger T & Henikoff S (2008) Histone H2A.Z and DNA

methylation are mutually antagonistic chromatin marks. *Nature* **456:** 125–129 Available

at: https://www.ncbi.nlm.nih.gov/pubmed/18815594