Sub-species niche specialization in the oral microbiome is associated with

nasopharyngeal carcinoma risk in an endemic area of southern China

Justine W. Debelius[1] *, Tingting Huang[1, 2] *, Yonglin Cai[3, 4] *, Alexander Ploner[1], Donal

Barrett[1], Xiaoying Zhou[5, 6], Xue Xiao[7], Yancheng Li[3, 4], Jian Liao[8], Yuming Zheng[3, 4],

Guangwu Huang[7], Hans-Olov Adami[1,9], Yi Zeng[10] §, Zhe Zhang[7] §, Weimin Ye[1] §

[1] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,

Sweden

[2] Department of Radiation Oncology, The First Affiliated Hospital of Guangxi Medical

University, Nanning, P. R. China

[3] Department of Cancer Prevention Center, Wuzhou Red Cross Hospital, Wuzhou, P. R.

China;

[4] Wuzhou Health System Key Laboratory for Nasopharyngeal Carcinoma Etiology and

Molecular Mechanism, Wuzhou, P. R. China

[5] Life Science Institute, Guangxi Medical University, Nanning, P. R. China;

[6] Key Laboratory of High-Incidence-Tumor Prevention & Treatment (Guangxi Medical

University), Ministry of Education, Nanning, P. R. China

[7] Department of Otolaryngology-Head & Neck Surgery, First Affiliated Hospital of Guangxi

Medical University, Nanning, P. R. China

[8] Cangwu Institute for Nasopharyngeal Carcinoma Control and Prevention, Wuzhou, P. R.

China

[9]Clinical Effectiveness Research Group, Institute of Health, University of Oslo, Oslo,

Norway

[10] State Key Laboratory for Infectious Diseases Prevention and Control, Institute for Viral

Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing,

P. R. China

Weimin Ye - Department of Medical Epidemiology and Biostatistics, Karolinska Institutet,

Nobels väg 12A, PO Box 281, Stockholm, SE-171 77, Sweden. Tel: +46-8-5248 6184; E-

mail: weimin.ye@ki.se.

Zhe Zhang - Department of Otolaryngology-Head & Neck Surgery, First Affiliated Hospital

of Guangxi Medical University, Nanning, P. R. China (zhangzhe@gxmu.edu.cn)

*First authors Contributed equally;

§Last authors who contributed equally.

Summary

Nasopharyngeal carcinoma (NPC) is a globally rare cancer, with a unique geographic

distribution. In endemic areas including Southern China, the incidence is more than 20

times higher than the rest of the world.[1] Although recent evidence suggests poor oral

hygiene is a risk factor for NPC,[2] it remains unknown whether the disease status is

associated with changes in the oral microbiome. Therefore, we carried out a population-

based case-control study in an endemic area of southern China.[3] We analyzed microbial

communities from 499 untreated incident NPC cases and 495 age and sex frequency-

matched controls. Here, we show the oral microbiome is altered in patients with NPC:

patients have lower microbial diversity and significant changes in the overall structure

of their microbial communities which cannot be attributed to other factors.

Furthermore, the combination of two closely related amplicon sequence variants (ASVs)

from *Granulicatella adiacens* an individual carried were predicted by disease status.

These ASVs sat at the center of a network of closely-related co-excluding organisms,

suggesting that NPC may be associated with subtle changes in the oral microbiome.

Study participants were recruited from the Wuzhou region in Southern China between 2010

and 2014 as part of a large population-based case-control study.[3] Saliva was collected during

interview. After sequencing and denoising to ASVs, samples from 1066 subjects had

sufficiently high-quality sequences and clinical information to be retained for analysis (Figure

S1). Preliminary investigation suggested the microbiota of a small number of former smokers

were highly heterogenous (n=72, 33 cases, 39 controls; Figure S2). We excluded former

smokers from the final analysis, retaining 994 individuals (Table S1; Figure S1).

We aimed to address the relationship between NPC and the oral microbiome, adjusted for

potential confounders. As a result, we looked for factors which might affect the oral

microbiome at a community level. Our primary confounders included oral hygiene and

health,[2,4,5] tobacco use,[6,7] family history of NPC,[8,9] alcohol use,[10,11] and tea consumption.[12,13]

We also considered a history of oropharyngeal inflammation, and the region where an

individual lived[14] as covariates primarily expected to affect the microbiome, as well as salted

fish consumption, which is primarily seen as a risk factor for NPC.[15]

When comparing alpha diversity between cases and controls, we found that NPC cases

showed significantly fewer overall ASVs, reduced phylogenetic diversity, and reduced

Shannon diversity compared to controls (rank sum $p < 0.001$; Figure 1a; Table S2); these

findings did not change after adjustment for covariates which were significantly associated

with alpha diversity (Figure 1b; Tables S3-S5). Hence, this suggests that patients newly

diagnosed with NPC have lower overall microbial diversity than healthy controls. Our results

agree with a small study of the oral microbiome in NPC patients (n=90), which also found

reduced alpha diversity.[16] Unlike other body sites, there is no clear relationship between

salivary microbiome richness and the health of the microbial community.

Figure 1. The oral microbiome differs between patients with nasopharyngeal carcinoma and healthy controls. (**a**) NPC cases (red) have significantly lower microbial richness compared to cases (blue; $p < 1\times10^{-12}$). The horizontal line in the boxlin represents the median, the large box the interquartile region, increasingly smaller boxes are the upper and lower eighths, sixteenths, etc. in the data, reflecting the distribution. This difference is reflected in (**b**) the correlation coefficients from a multivariate regression model. (**c**) Adonis testing with a model adjusted for age, sex, and sequencing run shows that for unweighted UniFrac distance, NPC diagnosis has more than five times the explanatory power of the next most important variable, residential community. For 9999 permutations, FDR-adjusted $p < 0.001$ ***; $p < 0.01$ **; $p < 0.05$*. (**d**) Principal coordinates analysis (PCoA) of unweighted UniFrac shows separation between cases (red) and controls (blue) along PC1 and PC2. Upper and right panels reflect the density distribution along each axis. The axes are labeled with the variation they explain. In unweighted UniFrac, PC1 explains 19.7% and PC2 explains 4.8% of the variation. A volcano plot of (e) the Poisson regression coefficient for disease status vs the log p-value reflects reduced diversity. The horizontal line indicates significant at a Benjamini-Hochberg corrected p-value of less than 0.05.

Similarly, when comparing global community patterns (beta-diversity) via Adonis models minimally adjusted for sex, age and sequencing run, we found significant differences between NPC cases and controls, both based on unweighted UniFrac distance[17] as well as for weighted UniFrac[18] and Bray-Curtis distances (FDR p< 0.001, 9999 permutations; Figures 1c,d and S3a,b). Compared to the potential confounders in the same setting, NPC status was the strongest explanatory factor for unweighted UniFrac distance, more than five times the effect size of the next strongest variable, as well as the second-strongest factor for weighted

UniFrac- and Bray-Curtis distances, just after tobacco use. There was no statistically

significant difference in dispersion between cases and controls in any metric, supporting the

idea that the differences are due to consistent differences between cases and controls ($p >$

0.55, 999 permutations; Figure 1d). Significance persisted in more fully adjusted Adonis

models including potential confounders with robust differences in community patterns.

These findings establish that NPC status and smoking are strongly associated with

differences in the oral microbiome in our population; the association with NPC is especially

strong with regard to presence and absence of organisms (as emphasized by unweighted

UniFrac), but second only to smoking with regard to abundances (as captured by weighted

UniFrac and Bray-Curtis). We found no evidence that these associations are driven by

community heterogeneity; they are, however, robust under adjustment for observed

confounders, and in the case of the unweighted UniFrac distances, unlikely to be the result of

confounding by unobserved factors due to the crushing dominance of the signal for NPC

status. Since we recruited incident, treatment-naive patients,[3,16] it is also implausible that the

observed differences in microbiome composition are treatment-related. Taken together, our

findings provide strong evidence for a clear difference in the oral microbiome between

patients with NPC and healthy controls.

Since the relationship between the microbiome and NPC status was strongest in unweighted

UniFrac distance, which focuses on presence and absence, we evaluated the relationship

between ASV prevalence and disease in a fully adjusted log binomial model. To limit

spurious correlations, we defined presence as a relative abundance greater than 0.02% and

focused on ASVs present in at least 10% of samples (n=245, Figure S4). We identified 53

ASVs which were significantly different between cases and controls (FDR $p < 0.05$; Figure

1e; Table S6). The large majority of these ASVs were more prevalent in controls and came

from a wide variety of taxonomic clades, which may suggest a somewhat stochastic loss of

ASVs in NPC patients, rather than a systematic loss of specific organisms (Table S6). This

finding is in line with our alpha diversity findings, and may indicate overall community

instability. In contrast, two ASVs were more prevalent in NPC cases: a member of genus

Lactobacillus (Lact-eca9) and a *Granulicatella* ASV (Gran-7770).

To evaluate whether NPC status affected abundance-based partitioning of the microbial

community, we applied Phylofactor.[19] Our model looked for phylogenetic clades which

differentiated NPC cases and controls, adjusting for potential confounders (Figure 2, Table

S7). Of the twelve factors examined, nine were associated with disease status. The primary

partition in the data suggested a *Granulicatella* ASV (Gran-7770) was 3.4 (95% CI 2.4, 4.9)

fold more abundant in NPC cases compared to controls. The third factor identified was

second *Granulicatella* ASV (Gran-5a37) as less abundant in cases. Both ASVs were also

associated with smoking status. We identified three large-scale shifts in microbial abundance

associated with NPC status. The remaining factors associated with NPC status were all single

ASVs which differentiated cases and controls, none of which differed in prevalence (Table

S6, S7).

Figure 2. There are significant associations between phylogenetic partitioning of the taxa and NPC status. The phylogenetic tree with the first 12 phylofactor-based clade partitions is shown on the left. The top row is colored by phylum, the associated color is shown below. The isometric log transformation is taken as the ratio of the tips highlighted in pink over those highlighted in gray and passed into the regression model to predict the coefficient shown in the forest plot. Clades which are excluded from that factor appear white in the row. The forest plot to the right shows the estimated increase in the factor associated with case-control status based on fitting the ratio in a linear regression adjusted for age, sex, sequencing run, number of missing or repaired teeth, tobacco use, and residential community. Error bars are 95% confidence intervals for the regression coefficient. Black bars indicate significance at $a < 0.05$, gray indicates a non-significant association.

Based on the significant difference in abundance and prevalence of ASVs from genus *Granulicatella* between cases and controls, we further explored this genus. We identified a total of 14 ASVs in the dataset; three were prevalent enough to be included in our feature-based analyses (Gran-5a37, Gran-7770, and Gran-6959). In 972 (97.8%) individuals, the abundant ASVs were the only Granulicatella present. When blasted against the Human Oral Microbiome Database (HOMD), the ASV sequences mapped to two cultured species with more than 99.5% accuracy to their corresponding assignment: *Granulicatella elegans* (*G. elegans*) which included Gran-6959 and *Granulicatella adiacens* (*G. adiacens*; Gran-7770 and Gran-5a37).[20] Strikingly, we found our two abundant *G. adiacens* ASVs differ by a single nucleotide: Gran-7770 carries a G at nucleotide 119 of our sequence (corresponding approximately to 458 in the full 16s rRNA sequence) while Gran-5a37 carries an A.

Gran-7770 was found to be 26% more prevalent among cases, while Gran-5a37 was among

the 51 ASVs less prevalent in cases (Prevalence Ratio [PR] 0.81 [95% CI 0.74, 0.88]; Table

S6]). Both ASVs were also significantly associated with smoking status: Gran-7770 was

more prevalent in smokers (PR 1.48, [95% CI 1.29, 1.70]) and Gran-5a37 less prevalent (PR

0.74, [95% CI 0.67, 0.81]). There was not a significant relationship between Gran-6959 (*G.*

elegans) and either disease status (PR 0.94 [95% CI 0.88, 1.00]) or tobacco use (PR 0.97

[95% CI 0.90, 1.06]).

We found that 993 out of 994 individuals carried at least one *G. adiacens* with a relative

abundance of at least 0.02%: 330 (33.2%) carried only Gran-5a37, 316 (31.8%) carried Gran-

7770 alone, and 347 (34.9%) carried both. Among individuals who were classified as

carrying only one ASV (Gran-7770 alone or Gran 5a37 alone), the "present" ASV was at

least 50-fold more abundant than the other variant. We used a multinomial logistic regression

to confirm that disease status was significantly associated with variants an individual carried:

compared to the odds of carrying Gran-5a37 alone, cases had significantly higher odds of

carrying both ASVs and, again, significantly higher odds of carrying Gran-7770 alone

(Figure 3a). Although smokers were more likely to have both ASVs or Gran-7770 alone,

there was no significant interaction between smoking and disease status.

Figure 3. The *Granulicatella adiacens* variant predicts community structure. (**a**) NPC cases have
significantly higher odds of carrying both Gran-5a37 and Gran-7770 than Gran-5a37 alone, and again,
significantly higher odds than carrying either Gran-5a37 and Gran-7770 or Gran-7770. (**b**) In unweighted
UniFrac space, we see separation based on the *G. adiacens* variant along PC2.

We also investigated how the presence of a *G. adiacens* variant structured the overall

microbial community. We filtered the full ASV table to remove any *Granulicatella* ASVs

and used the reduced table to re-calculate beta diversity metrics. The *Granulicatella*-free

community recapitulated the patterns seen in the full community well (Mantel $R^2 > 0.91$;

p=0.001, 999 permutations). We found significant differences between individuals who

carried Gran-7770, both, or Gran-5a37 in weighted and unweighted UniFrac distances and

Bray Curtis; all three metrics show clear separation in PCoA space (p=0.001, 999

permutations; Figure 3b; Figure S5). In unweighted UniFrac space (Figure 3b), the separation

was primarily along PC2, likely corresponding to the separation along PC2 seen between

cases and controls (Figure 1d). Furthermore, we found that the *G. adiacens* variant explained

16% of the variation attributed to case-control status in unweighted UniFrac distance and

15% of the variation in weighted UniFrac distance. Our results suggest that the *G. adiacens*

variant carried by an individual is significantly associated with community structure, and may

be a route by which NPC status shapes the oral microbiome.

We used a SparCC-based network analysis to identify other community members

Granulicatella might interact with to exert an effect on the microbiome.[21] We were able to

identify five networks: one pair of co-occurring ASVs, two pairs of co-excluding ASVs, one

three-member network of co-occurring ASVs and a large 29-member network of co-

occurring and co-excluding ASVs (Figures 4a). This main network consisted of two clusters

of a total of 20 organisms which were positively correlated with a *Granulicatella* variant; the

main members of the networks belonged to *Veillonella*, *Streptococcus*, and *Prevotella*.

Blasting against HOMD, we identified two additional pairs of ASVs that co-excluded

between the two nodes but mapped to the same clones: Stre-900d and Stre-0531

(*Streptococcus parasanguinis* clade 411) and *Prevotella melaninogenica* (Prev-b7f2 and

Prev-71e7; Figure 4b; Table S8).[20]

We hypothesize the co-excluding networks of ASVs, centered around *Granulicatella*, may

reflect partial niche specialization. Previous work suggests quorum sensing networks can

form between the core species,[22,23] and that metabolic changes occur in these networks. We

hypothesize these closely correlated organisms occupy the same niches within these

metabolic networks, however, strain-specific variation may either respond to or promote

disease-associated transformation. Without culture-based experimentation, it is difficult to

determine how these organisms may function in concert. One major challenge for in-silico

validation is the limited resolution of existing databases; our results exceed the OTU-based

resolution and span a less frequently characterized hypervariable region.

Figure 4. *Granulicatella adiacens* **variants set at the center of a network of closely related co-occurring organisms. (a)** SparCC-based network analysis for co-occurring and co-excluding ASVs for all subjects showed a large network with two clusters with common core structures. The color and shape of the nodes are genus-specific. The two *G. adiacens* variants are highlighted as stars: Gran-5a37 in purple and Gran-7770 in green. Correlated edges are shown in pink, anti-correlated edges are grey. The sides of each network are labeled with their associated *G. adiacens* **variant. (b)** Phylogenetic tree of the core ASVs from the network (positively correlated with either Gran-7770 or Gran-5a37). Tips are labeled by their association with Gran-7770 (Green) or Gran-5a37 (Purple).

Within the context of NPC in an endemic region, we hypothesize the oral microbiome may act through several potential mechanisms. The oral microbiome has been suggested to contribute to local tumorigenesis through immune regulation or oncogenic metabolites such as acetaldehyde or nitrosamines.[24] An *in silico* study suggested that commercially available

strains of *G. adiacens* and co-abundant organisms encode genes involved in nitrate and nitrite

reduction.[25]

Alternatively, we propose the possibility of an NPC-specific mechanism, in which the

microbiome interacts with the Epstein-Barr Virus (EBV). Infection with EBV is the most

widely accepted etiological factor for NPC, and butyrate, a well-known product of microbial

fermentation, has been linked to EBV reactivation,[26] a necessary step in NPC oncogenesis.[27]

The local microbiota has also been suggested to be involved in the acquisition and

persistence of oncogenic viral infections at other sites, for example, the interaction between

the vaginal microbiome and the human papillomavirus.[28] We therefore hypothesize the oral

microbiome and potentially the nasopharyngeal microbiome, may work in concert to lead to

high risk EBV infection in the nasopharyngeal epithelium, leading to NPC. However,

prospective studies are needed to determine whether the microbiome contributes to EBV

infection, or if differences in the oral microbiota only reflect EBV infection and NPC-related

stress.

In summary, we have demonstrated a difference in the oral microbial community between

NPC patients and healthy controls in an endemic area of southern China, which cannot be

explained by other measured factors. The difference is associated with both a loss of

community richness and differences among specific organisms, including closely related

ASVs from genus *Granulicatella*. In addition, we identified a network of co-occurring and

co-excluding ASVs which included these *Granulicatella* variants. These results strongly

suggest a relationship between the oral microbiome and nasopharyngeal carcinoma status in

untreated patients.

Acknowledgements

The authors wish to thank the study participants, the field work team for the NPCGEE

project, and the Wuzhou Health System Key Laboratory for Nasopharyngeal Carcinoma

Etiology and Molecular Mechanism and the Key Laboratory of High-Incidence-Tumor

Prevention & Treatment (Guangxi Medical University), especially Suhua Zhong, Xiling

Xiao, for the processing of salivary samples. The data was stored in the Department of

Medical Epidemiology and Biostatistics, Karolinska Institutet; the authors wish to thank them

for their assistance.

We acknowledge funding from the Swedish Research Council (2015-02625, 2015-06268,

2017-05814, PI Dr. W. Ye); the National Natural Science Foundation of China (81272983,

PI Dr. Z. Zhang); and the Guangxi Natural Science Foundation (2013GXNSFGA019002, PI

Dr. Z. Zhang). The field work of the NPCGEE study was funded by the National Cancer

Institute of the NIH (Award Number R01CA115873, PI H.-O. Adami). T. Huang is partly

supported by a grant from China Scholarship Council.

Data Availability

Raw sequencing data, feature table, and metadata are available from the corresponding author

upon request.

Author contributions

The study approach was conceived by HA, YZ, GH, ZZ and WY. YC, DB, WY, TH, JWD,

and AP refined the study design for this project. YC, YL, JL and YZ were responsible for

sample collection and management. DB performed the lab work, supervised by TH, XZ, XX,

ZZ, and WY. Bioinformatics and biostatistical analyses were performed by JWD; TH and AP

contributed to statistical modeling and refinement. WY contributed to the supervision and

coordination of the project. JWD and TH wrote the manuscript; AP provided critical edits.

All authors reviewed and approved the final submission.

Methods

Survey metadata and sample collection

Participant recruitment has been previously described.[3] Briefly, incident cases of NPC in

Guangdong Province and Guangxi Autonomous Region between 2010 and 2013 were invited

to participate in the study. Age and sex matched controls were selected from the total

population. The current study was approved by the Institutional Review Board or Ethical

Review Board at all participating centers. All study participants provided written or oral

informed consent.

A questionnaire covering demographics, diet, residential, occupational, medical and family

history was administered in a structured interview. Sample collection occurred at the

interview. Participants were asked not to eat nor chew gum for 30 minutes prior to sample

collection. Saliva samples with volumes (2ml-4ml) were collected into 50ml falcon tubes

with a Tris-EDTA buffer.

Demographic characteristics of the study population were compared using a two-sided t-test

for continuous covariates (age) and a chi-squared test for categorical covariates. Tests were

conducted using scipy 0.19.1[29] in python 3.5.5.

DNA extraction, PCR, and sequencing

Saliva DNA was extracted using a two-step protocol including the sample pre-processing

with lysozyme lysis and bead beating, and the TIANamp blood DNA kit (Beijing, China).

The 16s rRNA amplicon library was amplified with 341F/805R primers

(CCTACGGGNGGCWGCAG, GACTACHVGGGTATCTAATCC).[30,31] Samples were

amplified with 20 cycles of a program with 30 seconds at 98°C for melting, 30 second at

60°C, and 30 seconds at 72°C. Samples were barcoded in a second PCR step.[30] DNA clean-

up was performed using Agentcourt AMPure XP purification kit. DNA volume and purity

were measured on an Agilent 2100 Bioanalyzer system and Real-time polymerase chain

reaction. Sequencing was performed at Beijing Genome Institute (BGI) on an Illumina MiSeq

using a 2x300bp paired end strategy.

Denoising, Annotation and Filtering

Samples were demultiplexed using an in-house script. Adaptors were trimmed and paired end

sequences were joined using VSEARCH (v. 2.7).[32] Paired sequences were loaded into the

November 2018 release of QIIME 2.[33] Sequences were quality filtered (q2-quality-filter)[34]

and denoised using deblur (v. 1.0.4; q2-deblur)[35] with the default parameters on 420 bp

amplicons to generate amplicon sequence variants (ASVs). A phylogenetic tree was built

using fragment insertion into the August 2013 Greengenes 99% identity tree backbone with

q2-fragment-insertion;[36,37] taxonomic assignments were made with a naïve Bayesian

classifier trained against the same reference (q2-feature-classifier).[38] In cases where the

classifier or reference database was unable to describe a taxonomic level (for instance, a

missing genus), the taxonomy was described by inheriting the lowest defined level using a

custom python script. Following sequencing and denoising, 24,763,933 high quality reads

were retained.

Any sample with fewer than 1000 reads after denoising was excluded, leaving 1074 saliva

samples and 9 negative or single organism controls. Additionally, samples missing

information on tobacco use, defined information about tooth brushing frequency, or an

undefined residential region (n=8) were excluded (Figure S1).

Preliminary investigation suggested that the microbial communities for former smokers

(*n*=72) were highly heterogenous (Figure S2). Sensitivity analyses suggest their exclusion

does not alter the major community-level differences. Therefore, they were excluded, leaving

a total of 994 individuals in the analysis.

356
ASV-based analyses were performed on a representative subset: those with at least 0.02%

relative abundance in at least 10% of samples (n=245). A Mantel test[39] was applied to Bray

Curtis distance[40] and showed a correlation of 0.96 between the filtered matrix rarefied to

5000 sequences/sample and the full table distance matrix (p=0.001, 999 permutations); the

mantel corresponding correlation for UniFrac distance[41] was 0.76 (p=0.001, 999

permutations; Figure S3).

The sequences and identifiers for the abundant ASVs are listed in supplemental file 2. ASVs

are identified by the first 4 letters of their lowest taxonomic assignment and the first 4

characters of a MD5 hash of the sequence. The full taxonomic assignment and MD5 hashes

can be found in Table S6.

Diversity Analyses

Diversity analyses were performed using samples rarefied to 6,500 sequences.

Alpha diversity was calculated as observed ASVs, Shannon diversity,[42] and Faith's

phylogenetic diversity[43] using q2-diversity in QIIME 2. Potentially significant alpha diversity

predictors were identified using a rank-sum test in scipy 0.19.1.[29] A p-value of 0.05 was

considered the threshold for borderline significance for inclusion in a subsequent regression

model. Alpha diversity was then evaluated in a multivariate ordinary least squares (OLS)

regression model adjusted for age, sex and sequencing run number. A final model for each

metric was selected by forward selection using models which resulted in decreasing Akaike

information criterion (AIC). We checked for the normality of residuals by plotting. The

relative contribution of each covariate to that metric was estimated by a "leave one out"

approach. Regressions were performed in Statsmodels (v. 0.9.0).[44] For visualization, we

calculated z-normalized alpha diversity using the mean and standard deviation in diversity for

the controls. Alpha diversity was plotted using boxenplots in Seaborn 0.9.0.[45,46]

Beta diversity was measured using the unweighted UniFrac,[17] weighted UniFrac,[18] and Bray-

Curtis[40] metrics on rarefied data (q2-diversity). Beta diversity was compared using Adonis in

the R vegan library (v 2.5-2) adjusted for host age, sex, and sequencing run, with 9999

permutations.[47–49] We used a permdisp test with 999 permutations and the centroid estimate

to test for the presence of differences in within-group variation implemented in scikit-bio

0.5.4 (www.scikit-bio.org).[50] Uncorrected p-values of less than 0.05 were considered to have

significant dispersion, since we were more concerned about false positives than false

negatives. Principal coordinate analyses (PCoA)s were visualized using Emperor[51] (v.

1.0.0b18) and with seaborn[45] v. 0.9.0 in matplotlib v. 2.2.3.

ASV regression model

To look at the relationship between ASV prevalence and disease and smoking status, we used

a log binomial regression which was approximated via a Poisson regression with robust

standard errors,[52] implemented via base function *glm* in R and the robust error mechanism

implemented via packages *lmtest* (v 0.9) and *sandwich* (v. 2.5) in R 3.5.[49,53,54] The model was

adjusted for age, sex, sequencing run, residential community, and the number of missing or

repaired teeth. "Presence" was defined as a relative abundance of 1 / 5000, which

corresponded to the shallowest sequencing depth for the abundant counts. ASVs which were

present in more than 1000 samples were excluded from prevalence analysis. A Benjamini-

Hochberg FDR corrected p-value of 0.05 was considered significant.

Phylofactor

Phylofactor (v. 0.01) was used to look at the relationship between disease status and

phylogenetic partitioning between clades.[19] Phylofactor is a compositionally aware technique

which uses isometric log transforms over an unrooted phylogenetic tree to model differences

in the data. This allows the partitioning of data into polyphyletic clades. The Phylofactor

multivariate model for each partition was modeled with an OLS regression considering

diagnosis, adjusted for residential community, age, sex, number of missing or repaired teeth,

tobacco use, and sequencing run. We looked at the first 12 factors using the default

parameters, which optimized for explaining maximal variance. The cladogram, and

regression coefficient plots were generated in seaborn.[45]

Granulicatella

Total *Granulicatella* was identified by filtering the full ASV table for any ASV assigned to

the genus. Species-level assignments were made by blasting each ASV against the Human

Oral Microbiome Database using the online tool;[20] species-level assignments were taken for

the cultured species with the best match. We treated the abundance of Gran-6959 as the *G.*

elegens abundance and the combined abundance of Gran-5a37 and Gran-7770 as the *G.*

adiacens abundance throughout.

We used a multinomial logistic regression model, implemented in the *nnet* library (v. 0.8) in

R to look at whether the carriage of Gran-5a377 alone, Gran-7770 alone, or both ASVs was

associated with smoking and disease status.[55] The regression was adjusted for age, sex,

sequencing run, number of missing or repaired teeth, residential community, the relative

abundance of *G. adiacens*, and the relative abundance of *G. elegens*. Having Gran-5a37 was

considered the reference group for the multinomial logistic regression.

The effect of *Granulicatella* on alpha and beta diversity was calculated by first, filtering out

all *Granulicatella* ASVs from the table, and then rarifying to 6250 sequences/sample before

diversity calculations. Adonis coefficients were calculated in a model adjusted for *G.*

adiacens abundance, sequencing run, age, sex, residential community, number of missing or

repaired teeth, tobacco use, and disease status. The proportion of disease status explained by

comparing a model excluding the *Granulicatella* variant minus the model including the

variant over the model excluding the variant.

Network Analysis

We used the Sparse Cooccurrence Network Investigation for Compositional data (SCNIC;

https://github.com/shafferm/SCNIC) in QIIME 2 (q2-SCNIC) to perform network analysis on

the abundant ASVs in current and never smokers. The correlation network was built using

SparCC, and the network was built using edges with a correlation co-efficient of at least 0.3,

allowing both co-occurrence and co-exclusion.[21] Network clusters were identified by finding

the most connected node and following all positively correlated nodes in the trimmed SparCC

network. Networks were visualized in Cytoscape (v. 3.7.1) using a perfuse-weighted network

layout.[56] Nodes which were anti-correlated with a single node in the main cluster were

trimmed for the sake of visualization; these are labeled with the correlation coefficient.

The phylogenetic tree of core network members was visualized using ete3 (v. 3.1.1) in

python 3.6.[57]

References

1.   Wei, K.-R. *et al.* Nasopharyngeal carcinoma incidence and mortality in China, 2013.

457   *Chin. J. Cancer* **36**, 90 (2017).

458   2.   Liu, Z. *et al.* Oral Hygiene and Risk of Nasopharyngeal Carcinoma-A Population-

459   Based Case-Control Study in China. *Cancer Epidemiol. Biomarkers Prev.* **25**, 1201–7

460   (2016).

461   3.   Ye, W. *et al.* Development of a population-based cancer case-control study in southern

462   china. *Oncotarget* **8**, 87073–87085 (2017).

463   4.   Kilian, M. *et al.* The oral microbiome – an update for oral healthcare professionals. *Br.*

464   *Dent. J.* **221**, 657–666 (2016).

465   5.   Belstrøm, D. *et al.* Impact of Oral Hygiene Discontinuation on Supragingival and

466   Salivary Microbiomes. *JDR Clin. Transl. Res.* **3**, 57–64 (2018).

467   6.   Long, M., Fu, Z., Li, P. & Nie, Z. Cigarette smoking and the risk of nasopharyngeal

468   carcinoma: a meta-analysis of epidemiological studies. *BMJ Open* **7**, e016582 (2017).

469   7.   Wu, J. *et al.* Cigarette smoking and the oral microbiome in a large study of American

470   adults. *ISME J.* **10**, 2435–46 (2016).

8.   Huang, S.-F. *et al.* Familial aggregation of nasopharyngeal carcinoma in Taiwan. *Oral*

Oncol. **73**, 10–15 (2017).

9.   Blekhman, R. *et al.* Host genetic variation impacts microbiome composition across

human body sites. *Genome Biol.* **16**, 191 (2015).

10.   Chen, L. *et al.* Alcohol Consumption and the Risk of Nasopharyngeal Carcinoma: A

Systematic Review. *Nutr. Cancer* **61**, 1–15 (2009).

11.   Fan, X. *et al.* Drinking alcohol is associated with variation in the human oral

microbiome in a large study of American adults. *Microbiome* **6**, 59 (2018).

12.   Yuan, X. *et al.* Green Tea Liquid Consumption Alters the Human Intestinal and Oral

Microbiome. *Mol. Nutr. Food Res.* **62**, e1800178 (2018).

13.   Hsu, W.-L. *et al.* Lowered risk of nasopharyngeal carcinoma and intake of plant

vitamin, fresh fish, green tea and coffee: a case-control study in Taiwan. *PLoS One* **7**,

e41779 (2012).

14.   He, Y. *et al.* Regional variation limits applications of healthy gut microbiome

reference ranges and disease models. *Nat. Med.* **24**, 1532–1535 (2018).

15.   Barrett, D. *et al.* Past and Recent Salted Fish and Preserved Food Intakes Are Weakly

Associated with Nasopharyngeal Carcinoma Risk in Adults in Southern China. *J. Nutr.*

(2019). doi:10.1093/jn/nxz095

16.   Zhu, X.-X. *et al.* The Potential Effect of Oral Microbiota in the Prediction of Mucositis

During Radiotherapy for Nasopharyngeal Carcinoma. *EBioMedicine* **18**, 23–31 (2017).

17.   Lozupone, C. & Knight, R. UniFrac: a new phylogenetic method for comparing

microbial communities. *Appl Env. Microbiol* **71**, 8228–8235 (2005).

18.   Lozupone, C. A., Hamady, M., Kelley, S. T. & Knight, R. Quantitative and Qualitative

Diversity Measures Lead to Different Insights into Factors That Structure Microbial

Communities. *Appl. Environ. Microbiol.* **73**, 1576–1585 (2007).

19.   Washburne, A. D. *et al.* Phylogenetic factorization of compositional data yields

lineage-level associations in microbiome datasets. *PeerJ* **5**, e2969 (2017).

20.   Escapa, I. F. *et al.* New Insights into Human Nostril Microbiome from the Expanded

Human Oral Microbiome Database (eHOMD): a Resource for the Microbiome of the

Human Aerodigestive Tract. *mSystems* **3**, e00187-18 (2018).

21.   Friedman, J. & Alm, E. J. Inferring correlation networks from genomic survey data.

PLoS Comput. Biol. **8**, e1002687 (2012).

22.   Chalmers, N. I., Palmer, R. J., Cisar, J. O. & Kolenbrander, P. E. Characterization of a

Streptococcus sp.-Veillonella sp. Community Micromanipulated from Dental Plaque.

J. Bacteriol. **190**, 8145–8154 (2008).

23.   Palmer, R. J., Diaz, P. I. & Kolenbrander, P. E. Rapid succession within the

Veillonella population of a developing human oral biofilm in situ. *J. Bacteriol.* **188**,

4117–24 (2006).

24.   Gholizadeh, P. *et al.* Role of oral microbiome on oral cancers, a review. *Biomed.*

Pharmacother. **84**, 552–558 (2016).

25.   Hyde, E. R. *et al.* Metagenomic analysis of nitrate-reducing bacteria in the oral cavity:

implications for nitric oxide homeostasis. *PLoS One* **9**, e88645 (2014).

26.   Luka, J., Kallin, B. & Klein, G. Induction of the Epstein-Barr virus (EBV) cycle in

latently infected cells by n-butyrate. *Virology* **94**, 228–231 (1979).

27.   Hirayama, T. & Ito, Y. A new view of the etiology of nasopharyngeal carcinoma.

Prev. Med. (Baltim). **10**, 614–22 (1981).

28.   Mitra, A. *et al.* The vaginal microbiota, human papillomavirus infection and cervical

intraepithelial neoplasia: what do we know and where are we going next? *Microbiome*

4, 58 (2016).

29.   Jones, E., Oliphant, T., Peterson, P. & others. SciPy: Open source scientific tools for

Python.

30. Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km

salinity gradient of the Baltic Sea. *ISME J.* **5**, 1571–9 (2011).

31. Hugerth, L. W. *et al.* DegePrime, a Program for Degenerate Primer Design for Broad-

Taxonomic-Range PCR in Microbial Ecology Studies. *Appl. Environ. Microbiol.* **80**,

5116–5123 (2014).

32. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile

open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

33. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

34. Bokulich, N. A. *et al.* Quality-filtering vastly improves diversity estimates from

Illumina amplicon sequencing. *Nat. Methods* **10**, 57–9 (2013).

35. Amir, A. *et al.* Deblur Rapidly Resolves Single-Nucleotide Community Sequence

Patterns. *mSystems* **2**, e00191-16 (2017).

36. Janssen, S. *et al.* Phylogenetic Placement of Exact Amplicon Sequences Improves

Associations with Clinical Information. *mSystems* **3**, e00021-18 (2018).

37. McDonald, D. *et al.* An improved Greengenes taxonomy with explicit ranks for

ecological and evolutionary analyses of bacteria and archaea. *ISME J* **6**, 610–8 (2012).

38. Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for

rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ.*

Microbiol. **73**, 5261–7 (2007).

39. Mantel, N. The detection of disease clustering and a generalized regression approach.

Cancer Res. **27**, 209–220 (1967).

40. Sørensen, T. *A method of establishing groups of equal amplitude in plant sociology*

based on similarity of species content and its application to analyses of the vegetation

on Danish commons. (I kommission hos E. Munksgaard, 1948).

41. Lozupone, C. & Knight, R. UniFrac: a new phylogenetic method for comparing

microbial communities. *Appl. Environ. Microbiol.* **71**, 8228–35 (2005).

42. Shannon, C. E. & E., C. A mathematical theory of communication. *ACM SIGMOBILE*

Mob. Comput. Commun. Rev. **5**, 3 (2001).

43. Faith, D. P. & Baker, A. M. Phylogenetic diversity (PD) and biodiversity conservation:

some bioinformatics challenges. *Evol Bioinform Online* **2**, 121–128 (1992).

44. JS Seabold, J. P. Statsmodels: Econometric and Statistical Modeling with Python.

Proc. 9th Python Sci. Conf. (2010).

45. Waskom, M. *et al.* mwaskom/seaborn: v0.9.0 (July 2018). (2018).

doi:10.5281/ZENODO.1313201

46. Hofmann, H., Kafadar, K. & Wickham, H. *Letter-value plots: Boxplots for large data.*

The American Statistican (2011).

47. McArdle, B. H. & Anderson, M. J. FITTING MULTIVARIATE MODELS TO

COMMUNITY DATA: A COMMENT ON DISTANCE-BASED REDUNDANCY

ANALYSIS. *Ecology* **82**, 290–297 (2001).

48. Oksanen, J. *et al.* vegan: Community Ecology Package. (2018).

49. R Core Team. R: A Language and Environment for Statistical Computing. (2018).

50. Anderson, M. J. Distance-Based Tests for Homogeneity of Multivariate Dispersions.

Biometrics **62**, 245–253 (2006).

51. Vázquez-Baeza, Y. *et al.* EMPeror: a tool for visualizing high-throughput microbial

community data. *Gigascience* **2**, 16 (2013).

52. Barros, A. J. & Hirakata, V. N. Alternatives for logistic regression in cross-sectional

studies: an empirical comparison of models that directly estimate the prevalence ratio.

BMC Med. Res. Methodol. **3**, 21 (2003).

53. Zeileis, A. Object-Oriented Computation of Sandwich Estimators. *J. Stat. Softw.* **16**,

1–16 (2006).

54.   Zeileis, A. Econometric Computing with {HC} and {HAC} Covariance Matrix

Estimators. *J. Stat. Softw.* **11**, 1–17 (2004).

55.   Venables, W. N. & Ripley, B. D. *Modern Applied Statistics with S.* (Springer, 2002).

56.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Res.* **13**, 2498–504 (2003).

57.   Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and

Visualization of Phylogenomic Data. *Mol. Biol. Evol.* **33**, 1635–1638 (2016).