1 **Selection-based model of prokaryote pangenomes**

2 Maria Rosa Domingo-Sananes and James O. McInerney
3 School of Life Sciences, University of Nottingham. University Park. Nottingham, NG7 2RD.
4 United Kingdom.
5
For correspondence: Maria.Domingosananes@nottingham.ac.uk;
James.McInerney@nottingham.ac.uk

Abstract

The genomes of different individuals of the same prokaryote species can vary widely in gene
content, displaying different proportions of core genes, which are present in all genomes, and
accessory genes, whose presence varies between genomes. Together, these core and
accessory genes make up a species' pangenome. The reasons behind this extensive diversity
in gene content remain elusive, and there is an ongoing debate about the contribution of
accessory genes to fitness, that is, whether their presence is on average advantageous,
neutral, or deleterious. In order to explore this issue, we developed a mathematical model to
simulate the gene content of prokaryote genomes and pangenomes. Our model focuses on
testing how the fitness effects of genes and their rates of gene gain and loss would affect the
properties of pangenomes. We first show that pangenomes with large numbers of low-
frequency genes can arise due to the gain and loss of neutral and nearly neutral genes in a
population. However, pangenomes with large numbers of highly beneficial, low-frequency
genes can arise as a consequence of genotype-by-environment interactions when multiple
niches are available to a species. Finally, pangenomes can arise, irrespective of the fitness
effect of the gained and lost genes, as long as gene gain and loss rates are high. We argue
that in order to understand the contribution of different mechanisms to pangenome diversity,
it is crucial to have empirical information on population structure, gene-by-environment
interactions, the distributions of fitness effects and rates of gene gain and loss in different
prokaryote groups.

Introduction

Intraspecific variation in gene content in prokaryotes is often so extensive that for a given
species a "typical genome" might not exist[1,2]. Therefore, the concept of a pangenome, defined
as the complete set of genes present in a species, has been put forward as a better description
of the gene content of a prokaryote species[1]. Pangenomes include *core* genes that are present
in all individuals, and *accessory* genes, whose presence varies among individuals. The role
of accessory genes remains puzzling, and a debate has developed centring on whether they
have positive, negative, or neutral effects on the cells that host them[3–8]. Understanding the
contributions of these accessory genes to cell fitness and evolution has important implications,
including insights into the factors that determine the prevalence and maintenance of virulence
genes, antibiotic resistance, and other traits often encoded by accessory genes[1,9].

Pangenomes arise as a consequence of gene acquisition via horizontal gene transfer (HGT),
and gene loss[2]. These processes ultimately result in general patterns observed across
prokaryotes, which include an increase in the number of known accessory genes as more
genomes from the same species are sequenced —along with a slower decrease in the number
of core genes— and a U-shaped *gene frequency distribution* or *spectrum*[2,10,11]. The U shape

of this distribution indicates that there is a large proportion of genes present in a single or very
few genomes, few genes present at intermediate frequencies, and substantial proportion of
core genes. Although across prokaryotic life there is a certain amount of commonality in the
shape of the gene content frequency distributions, different prokaryote species manifest
significant differences in the proportions of core and accessory genes[12–14].
54
Although we know that HGT and gene loss result in pangenomes, what is less clear are the
forces that lead to high diversity in gene content and U-shaped gene frequency distributions.
According to mathematical models, gain and loss of entirely neutral genes along a simulated
phylogeny or population can in principle predict the U-shaped gene frequency distributions of
pangenomes [11,15]. However, the fit to real data improves significantly when at least two classes
of genes, with either slow or fast turnover rates, are considered[11,15,16]. Specifically, including a
class of genes with low turnover rates increases the number of core and very common genes,
thus allowing a better fit to the peak of high frequency genes in the distribution. This implies
that selection is required to retain core genes for long periods of time, exactly as we would
expect if these have important functions. Therefore, these models suggest that high frequency
genes are beneficial on average, while most accessory and very rare genes are likely to be
neutral.
67
Alternative models suggest that accessory genes are on average beneficial, and propose
selection-based mechanisms that can explain the maintenance of diversity in gene
content[12,17–19]. One possibility is that negative-frequency dependent selection (NFDS) acting
on large numbers of accessory genes explains the existence and maintenance of gene
content variation[20,21]. NFDS occurs when the fitness effect of a gene changes from positive to
negative as its frequency in the population increases. That is, the gene is beneficial when rare
but deleterious when common and is therefore maintained at an intermediate frequency in the
population. NFDS may be important for maintaining gene content diversity under some
conditions, such as when rare serotypes become fitter after a vaccination program that only
targets common serotypes[17,18]. A related possibility is the existence of Black Queen effects,
where the combination of pressure to reduce genome size while maintaining common-good
functions results in intermediate frequencies of the genes encoding such functions[22].
Genotype-by-environment (GxE) effects have also been proposed to maintain beneficial
genes at an intermediate frequency in a population[12]. GxE effects imply that particular genes
are beneficial only in some conditions and are therefore only maintained in particular
environments. Migration of cells between different ecological niches, or distinct environments,
could explain variation in gene content and the maintenance of genes at low frequencies due
to differences in their fitness in the different environments[19].
86
Overall, different models propose that the existence of pangenomes can be the consequence
of either neutral processes or selection. We do not currently know which of these models
better explains pangenomes in nature. Most likely, genes with deleterious, neutral and
beneficial effects are all present in the accessory component of pangenomes. The more
difficult question is whether we can determine which fitness class is the most prevalent and,
at the same time, understand what processes and pressures determine the existence and
maintenance of pangenomes and their properties. Here we develop a mathematical model
from the point of view of a single gene that has a particular fitness contribution (that can be
positive or negative) and that can be gained or lost. We analyse the consequences that
variation in fitness effects, and the rates of gene gain and loss, have on the expected

frequencies of a gene in a population where drift is negligible. We then expand the single-
gene model in order to consider many genes, each allowed to have different properties and
using this genome-scale expansion, we simulate pangenomes and their gene frequency
distributions. We show that the distributions of the input parameters directly influence
pangenome properties such as the size of the accessory genome, core genome and the shape
of the gene frequency distribution. We also model the effect that multiple niches, and niche-
dependent gene fitness contributions have on pangenomes. Our results show that the
presence of accessory genes is likely to be overwhelmingly nearly neutral in the absence of
GxE effects, which is likely the case when populations are constrained to a single
environmental niche (*i.e.* there is no migration or exposure to variation in environmental
conditions). However, when multiple environmental niches are considered, accessory genes
can be highly beneficial. Overall, we argue that real pangenomes are an outcome of a
combination of processes and crucially, variability in model parameters, like fitness effects and
rates of gene gain and loss, for different genes and species.

Model description

We propose the model depicted in Figure 1a for the frequency of a gene that can be present
or absent in the genomes of a prokaryote group (for example a population or species). We
assume that cells without the gene divide at rate μ, and that the gene's presence provides a
contribution s to this growth rate, such that cells that possess the gene divide at rate $\mu + s$.
We call s the selection coefficient or *fitness effect* of the gene, and its value can be positive or
negative. For cells that lack the gene, it can be gained at a rate r_g, and lost from cells that
have it with rate r_l (Figure 1a). We assume that the population size is constant and that all
cells in the population have the same death rate ν for cells with and without the gene. We also
assume that genetic drift in the population is negligible and that the only changes in genomes
are the gain and loss of genes. That is, there are no other mutations (that would, for example,
change the value of s). We can then write a differential equation for the frequency of the gene
in the population, x:

126
$$\frac{dx}{dt} = r_g(1-x) - r_l x + (\mu + s)x - \nu x$$

128 where

129
$$\nu = \mu(1-x) + (\mu + s)x$$

131 because the population size is constant $(d(x + (1-x))/dt = 0)$. Therefore, we obtain the
132 following equation, which describes the frequency of the gene independently of the basal
133 growth rate of the population:

$$\frac{dx}{dt} = r_g(1-x) + sx(1-x) - r_l x \qquad \text{Eq. 1}$$

Thus, in this model, the frequency of a gene in a population depends on three parameters: its
fitness effect (s), its rate of gain (r_g) and its rate of loss (r_l). These parameters summarise
important features of the external environment, the recipient cell, and the gene itself:

The *fitness effect* of a gene describes the additive contribution of that gene to fitness, or more
narrowly to the cell's growth rate. That is, it describes the difference in growth provided by the

gene per unit time. Therefore, $s > 0$ if the gene confers a trait that increases fitness, (i.e. if it
is beneficial), $s < 0$ if the gene's presence is deleterious, and $s = 0$ if the gene has a neutral
effect with respect to fitness. The fitness effect is not just a property of the gene itself but
depends on the organism's biology. For example, it depends on whether or not the cell can
make use of that gene, in terms of compatibility with its cell biology and codon usage, for
instance. Importantly, the value of this parameter also depends on the environment. For
example, a gene that allows a cell to metabolise a substrate, or protect it from an antibiotic,
will only be beneficial in the relevant external ecological setting.

The *rate of gain* for a given gene describes the proportion of cells in the population that gain
that gene per unit of time. The value of this parameter is heavily dependent on the route
through which the gene can be acquired. For example, genes acquired by transduction or
conjugation are likely to have high rates of gain, while those acquired through natural
transformation will have lower gain rates. Importantly, this parameter also depends on the
prevalence of a gene in the environment (whether it is environmental DNA, phage, plasmids
or other cells), because the higher the prevalence, the higher the chance a gene can be
acquired by a cell in the focal population. Finally, the rate of gain is also dependent on the
characteristics of the species or cells where the gene is acquired. For example, factors such
as the intrinsic rates of natural transformation and recombination, the prevalence of lysogenic
phage infections and conjugative plasmids, and even the existence and efficiency of defence
mechanisms that can prevent the entry of exogenous DNA, will impact on the gain rate of
genes. The *rate of loss* for a given gene is the proportion of cells that lose that gene per unit
time, and likely depends mostly on the organism's biology, i.e. rates of recombination that lead
to gene loss. However, this rate may also vary with the environment and properties that
enhance or reduce rates or recombination of particular genes, such as the types of sequences
flanking the gene[23,24].

Our model does not explicitly include evolutionary history (*i.e.* phylogeny) and drift, therefore,
the implication is that selection is quite effective. Consequently, our model is at the opposite
end of the spectrum of assumptions when compared to models based exclusively on neutral
evolution[11,15]. We use this model to explore how the values of these parameters and their
distributions determine genome content and pangenome properties.

173
174 ***Figure 1:*** Selection-based model of pangenomes. (a) Model scheme for the frequency of one
175 gene. (b) Model scheme for multiple genes and pangenome simulations.

Results

Effect of model parameters on pangenome gene frequencies

According to the model (Eq 2, Methods) the steady state gene frequency in a prokaryote population is a sigmoid function of the fitness effect s, which means that the frequency of a gene increases monotonically from 0 to 1 as the value of s increases, but changes in a switch-like manner around $s = 0$ (Figure 2a). Overall, this implies that deleterious genes are present at low frequencies, while beneficial genes are found at high frequencies. The rates of gene gain and loss influence the shape of this sigmoid function. As the gain and loss rates decrease, the sharpness or switch-like shape of the function increases (compare purple and blue lines in Figure 2a). The consequence of lower rates of gene gain and loss is a decrease in the frequency of deleterious genes, and an increase in the frequency of beneficial genes. In the limit where the gene gain and loss rates are equal to zero, we obtain a perfect step function (Equation 2), such that all advantageous genes are fixed in the population and all deleterious genes are absent from the population, even if their fitness effects are very small (due to our assumption of strong selection). This result implies that at equilibrium, fitness is maximised when the rates of gene gain and loss are very small compared to fitness effects. However, if the gene gain and loss rates are equal to zero, any beneficial genes not already present in the genome cannot be gained, while any deleterious genes cannot be lost. Additionally, if the rates of gene gain and loss are very small, it takes longer to reach steady state frequencies and the gene frequency dynamics are mainly driven by selection (see time-course simulations in Figure S1). Furthermore, in agreement with evolutionary theory, the smaller the fitness effect, the slower the change in gene frequency (not shown).

The relative balance between the rates of gene gain and loss is also important for determining the equilibrium gene frequency. Figure 2b shows that if the rate of gene loss increases (from blue to purple line) while the rate of gain remains fixed, the frequency of deleterious genes remains low, while the frequency of beneficial and nearly neutral genes decreases. Therefore, according to the model, if rates of loss are higher than rates of gain, deleterious genes are efficiently purged from the population. The drawback to higher rates of loss is that highly beneficial genes can also be lost at high rates and are therefore not fixed in the population at equilibrium. The disparity in the rates of gain and loss also has an effect on gene frequency dynamics. At high loss rates, deleterious genes are quickly lost, but acquisition of advantageous genes is delayed in comparison with the case of equal gain and loss rates (Figure S1). Essentially the opposite situation occurs if the rates of gain are higher than the rates of loss: beneficial genes will be fixed in the population, but neutral and deleterious genes will also be present at high frequencies (Figure 2c). Furthermore, the model shows that highly deleterious genes will be present at low or intermediate frequencies. In terms of dynamics, higher rates of gene gain than loss also implies that advantageous genes quickly fix in the population, but deleterious genes are lost slowly (Figure S1).

220
221 **Figure 2**. Dependence of gene frequency with respect to model parameters. Plots shows the
222 gene frequency with respect to the fitness effect, s, for different rates of gene gain and loss.
223 A. Effect of rates of gene and loss when $r_g = r_l$. B. Effect of the rate of gene loss with fixed
224 rate of gene gain ($r_g = 0.01$). C. Effect of the rate of gene gain with fixed rate of gene loss
225 ($r_l = 0.01$).

226
Overall, according to our model, it is not just the fitness effects of accessory genes that can
determine the frequency of a gene in the population, but the rates of gene gain and loss are
also important, especially if their values are of a magnitude comparable to the fitness effects
of genes. In particular, the frequency of nearly neutral genes in a population is heavily
dependent on the balance between rates: if gain rates are higher than loss rates, neutral genes
are maintained in the population at high frequency, while they will be rare when loss rates are
higher (Figure S2).

234
235 _Effect of parameter distributions on pangenomes_

236
In a population of cells in which multiple genes can be gained and lost, the values of the model
parameters are likely to vary among different genes. Furthermore, even for a single gene, its
fitness effect, rate of gain and rate of loss can vary among different species, cells and
environments. Therefore, each of the gene-specific model parameters is associated with a
parameter distribution. Some point estimates for rates of gene gain and loss have been
obtained[25] and individual examples of fitness effects of accessory genes are known[8,26]
however, there is very little empirical data on these parameter distributions. Consequently, in
our model we rely on hypothetical parameter distributions in order to assess how parameter
variation shapes pangenomes. We sampled gene parameter values from proposed
distributions and simulated gene content for multiple genomes in a population at equilibrium
as described in Methods. Importantly, because we know the parameter values of these
simulated pangenomes, we assessed the contribution to pangenomes of genes with different
fitness effects.

250
Distribution of fitness effects
The distribution of fitness effects (DFE) of genes is central to our expectations of the gene
content variation we see in populations. This distribution is analogous to the distribution of
fitness effects of mutations, which is important in determining the level of nucleotide
polymorphisms that are maintained in populations[8,27]. We currently know little about the DFE
of accessory genes, but as a first approximation we speculate that it may have similar

properties to the DFE for mutations: most genes will have nearly neutral effects and highly
beneficial or highly deleterious genes will be rare.
259
As a first approximation, we assume equal and relatively low rates of gene gain and loss for
all genes. For simplicity, we also assume a finite pool of genes available for acquisition. Figure
3A shows results from a simulation of 100 genomes, sampled from a gene pool of 1000 genes
whose fitness effects are drawn from a normal distribution with a mean of zero and unit
standard deviation (blue density plot in Figure 3Ib). The top panel (Figure 3Ia) shows the
presence/absence of all the genes in the 100 simulated strains as a heatmap, and the scatter
plot (Figure 3Id) shows the frequency of every gene in the pangenome with respect to its
fitness effect. In this case, advantageous genes ($s > 0$) are present at high frequencies, nearly
neutral genes present at intermediate frequencies and deleterious genes ($s < 0$) are very rare
or absent. This simulation thus qualitatively recovers a U-shaped gene frequency distribution
similar to those observed for real pangenomes[2] (Figure 3Ic). In these simulations highly
advantageous genes are essentially fixed and would be considered the 'core' genome, while
nearly neutral genes and some deleterious genes would constitute the 'accessory' genome.
This also results in 'filtering' of a fraction of the deleterious genes, such that the fitness effects
of genes present in the pangenome (orange density plot in Figure 3Ib) are slightly higher than
the original gene pool DFE. These results demonstrate that the pattern of gene content
variation of real pangenomes can be recovered by a model based on selection only. This
contrasts with neutral models, which propose that the U-shaped gene frequency distributions
observed in prokaryote pangenomes can be explained by entirely neutral genes subject only
to drift in an evolving population[11,15]. Therefore, models with opposite assumptions regarding
the effects of drift and selection, are able to qualitatively recover patterns of gene content
variation that are similar to empirically observed patterns.
282
In our model, if we assume most genes are nearly neutral, the gene frequency distribution
starts to deviate from the expected U shape. Compared to the model with larger variance in
fitness effects (Figure 3I), if we significantly reduce the magnitude of fitness effects of the DFE
(~10-fold reduction) the result is that more genes are present at intermediate frequencies and
fewer genes are fixed or present in a single genome (Figure 3II). In the limit where all genes
are truly neutral, the mean frequency of all genes would be expected to be 0.5 (when the rates
of gene gain and loss are equal). This is because each gene has equal chance of being
present in any sampled genome, since we assume the population is at equilibrium and do not
consider the effects of phylogeny and drift.
292
Many other possible DFEs among incoming genes can result in U-shaped gene frequency
distributions arising in a population. Consequently, just changing the DFE results in a myriad
of observed gene frequency distributions. For example, Figure 3III shows the simulated
pangenome of 100 cells, each with 1000 genes whose fitness coefficients are sampled from
an exponential distribution with rate parameter 1 and mean shifted to -1. In this case, most
genes are highly deleterious, and many are not found in the simulated pangenome (Figure
3IIIa). In this case, most accessory genes are very rare, and only a small proportion of genes
that are highly advantageous are present at high frequencies. These results illustrate that
many distributions of fitness effects can potentially replicate the patterns observed in real
pangenomes and would suggest that core genes are highly beneficial and accessory genes
are mostly nearly neutral or slightly deleterious, in agreement with previous modelling

results[16,28]. However, this conclusion also relies on simplifying assumptions that are unlikely
to be met in the real world, such as equal rates of gene gain and loss for all genes.
306

Figure 3. Simulated pangenomes for different distributions of fitness effects. Each panel
represents a simulation with fitness effects sampled from the specified distribution of fitness
effects. **I**. Normal distribution with mean 0 and SD = 1. **II**. Normal distribution with mean 0 and
SD = 0.1. **III**. Exponential distribution with rate 1, mean shifted to -1. For each panel: **a**.
Presence/absence of genes in the simulated pangenome, present genes are indicated in grey.
The orange line indicates the number of strains in which a gene is present. **b**. Density plots
for the DFE of the gene pool (sampled from the specified theoretical distribution, in blue) and
for the fitness effects of genes actually present in the pangenome (orange). **c**. Gene frequency
distribution for the simulated pangenome. **d**. Gene frequencies versus fitness effects for 1000
genes in the simulated pangenome (blue dots) and the theoretical relationship (orange line).
Rates of gene gain and loss are assumed to be equal for all genes ($r_g = r_l = 0.01$)

Variation in gene gain and loss rates

Because the rates of gene gain and loss play an important role in determining the equilibrium
frequency of a gene (as we showed in the previous section), we expect that their distributions
will be important in shaping pangenomes. A reasonable approximation is that for most genes,
the rates of gene gain and loss are very low, while for a small number of genes these rates
can be relatively high. Thus, as an example, we sample these gain and loss rates from an
exponential distribution. Figure 4 shows results from simulated pangenomes with fitness
effects drawn from a normal distribution with mean 0 and standard deviation of 1 (blue density
plot in Figures 4Ic, 4IIc and 4IIIc) and variable rates of gene gain and loss (Figures 4Ib, 4IIb
and 4IIIb). If we fix the rate of gene loss, and draw the rates of gene gain from an exponential
distribution with rate parameter 0.1 (Figure 4Ib), we find that some deleterious genes can be
found at high frequencies (Figure 4Ie), when compared to the assumption of equal rates for
all genes (Figure 3Id). If we fix the rate of gene gain and sample rates of gene loss from a
similar exponential distribution (Figure 4IIb), we find an asymmetric U-shaped distribution in
which many advantageous genes are now found at intermediate or low frequencies (Figure
4IIe). Finally, if we independently sample the three parameters for every gene (Figure 4III),
we recover a symmetric U-shaped gene frequency distribution (Figure 4IIId). The U-shape of
this distribution in this case is less extreme than that of Figure 3Ic, because there are more
genes with intermediate frequencies for all values of fitness coefficient. This is due to both
higher average gene gain and loss rates and the variation in these values. Overall, this

analysis shows that if gene gain and loss rates vary for different genes, the observed gene frequencies are no longer determined by their fitness effects. Therefore, some beneficial genes may be present a low frequencies and deleterious genes at high frequencies.

Figure 4. Simulated pangenomes for different distributions of rates of gene gain and loss. Each panel represents a simulation with rates of gene gain and loss sampled from the specified distribution. Fitness effects were sampled from a normal distribution with mean 0 and SD = 1. **I**. Rates of gain sampled from an exponential distribution with parameter 0.1 and fixed rate of loss ($r_l = 0.01$). **II**. Rates of loss sampled from an exponential distribution with parameter 0.1 and fixed rate of gain ($r_g = 0.01$). **III**. Rates of gain and loss sampled from independent exponential distributions with parameter 0.1. For each panel: **a**. Presence/absence of genes in the simulated pangenome, present genes are indicated in grey. The orange line indicates the number of strains in which a gene is present. **b**. Density plots for the sampled distributions of rates of gene gain and loss **c**. Density plots for the DFE of the gene pool (sampled from the specified theoretical distribution, in blue) and for the fitness effects of genes actually present in the pangenome (orange). **d**. Gene frequency distribution for the simulated pangenome. **e**. Gene frequencies versus fitness effects for 1000 genes in the simulated pangenome (blue dots) and the theoretical relationship (orange line).

Niche-dependent fitness effects

The previous results show that if the focal organism can only live in a single ecological niche, then accessory genes are most likely to be either nearly neutral or deleterious, unless their rates of gene gain and loss are relatively high. However, besides the assumption of strong selection, this relies on the important but hidden assumption that a gene's fitness effect is constant across all individuals of a given species. That is, the previous simulations have assumed that the contribution of a gene to a cell's fitness is the same throughout the range of a species, in all possible environmental conditions and niches. This uniformity of fitness effects across environments is unlikely to be true for many genes[23]. For example, genes that can confer antibiotic resistance are likely to only be beneficial in environments where the antibiotic is present, and either confer no benefit or have a cost in its absence[29]. In other words, as the antibiotic concentration rises, presence of a resistance gene goes from normally deleterious (or neutral) to highly advantageous, to the point of potentially becoming essential under constant antibiotic pressure[30].

To analyse the effect of niche dependence, we simulated pangenomes by sampling genomes from different 'niches', where a gene has different fitness effects, depending on the niche (Figure 5). In this case, to evaluate the *fitness contribution of a gene* to the simulated population sample, we introduce a new measure ϕ, which is the average fitness effect of the gene when present (see Methods). For example, if a gene is beneficial only in a particular niche and only present there, its fitness contribution will be positive. If it is also present in niches where its fitness effect is negative, its fitness contribution to the overall population will be reduced.

We first consider a single niche where gene fitness effects are drawn from a normal distribution as in the previous section. Figure 5Ia displays values of these fitness effects for 1000 genes as a heatmap. As before, this simulation recovers a U-shaped distribution of gene frequencies (Figure 5Id). In this case, the fitness contribution of a gene is identical to its fitness coefficient, because the fitness effect is the same for all genomes (Figure 5Ic). Again, we see that genes present at high frequency are on average highly advantageous (have a high fitness contribution) and rare genes or those present at intermediate frequencies are nearly neutral or deleterious. However, if we allow multiple niches, this situation can change drastically. Figure 5B shows an example where we consider 10 hypothetical niches. We assume that all genes in the gene pool are available for acquisition in all the niches, but genes are on average deleterious in 9 of the 10 niches and on average beneficial in just one. 100 genes are assumed to be on average beneficial in each niche, and we sample 10 genomes from each niche to simulate the pangenome shown in Figure 5IIa. In this case, most genes are only found in an average of 10 genomes (Figure 5IId, and square blocks in Figure 5IIa), those in which the genes are beneficial. Therefore, these relatively rare genes have a high fitness contribution (Figures 5IIe). In a reversal from the previous case, most genes found at high frequency make nearly neutral fitness contributions. Overall, all genes have a positive fitness contribution to the population (Figure 5IIc and 5IIe). By considering even more niches, for example 100 in Figure 5III and a single genome sampled from each, we obtain an even more extreme picture where all the genes that significantly contribute to fitness are very rare and most present in only one or two genomes (Figure 5IIId and 5IIIe). This result is in stark contrast to the simulations with constant fitness effects and low, uniform rates of gene gain and loss, where rare genes had a nearly neutral or deleterious effect. The situation corresponds to that described by McInerney et al.[12], who proposed that where selection is highly effective (in species that have large long-term effective population sizes) and where multiple, diverse niches are occupied by the same species, we should expect accessory genes to accumulate and pangenomes to become quite large.

These simulations are extreme cases and intended as exemplars of the effect that niche-dependent selection can have on our expectations of the correlation between the frequency of a gene in the population and its fitness effect. Real pangenomes are more likely to be closer to the intermediate case presented in in Figure 5IV. In this case we also assume one genome sampled from each of 100 niches, where a single gene is likely highly beneficial, and the rest have fitness effects drawn from a normal distribution with mean zero. Therefore, the overall fitness effects of genes are similar to the single-niche situation presented in Figure 5A, except for the fact that each gene is likely to be highly beneficial in one niche. This simulation again recovers a U-shaped gene frequency distribution. The crucial difference in comparison to the single-niche case, is that now a substantial proportion of rare genes positively contribute to

fitness in particular genomes (Figure 5IVc, 5IVd and 5IVe). This intermediate scenario then shows that accessory genes can be (nearly) neutral or advantageous.

In summary, our analysis shows that accessory genes present in pangenomes can be neutral, deleterious or advantageous, depending on particular properties of the genes, cells and environment. Therefore, the distributions and environment-dependence of gene fitness effects and rates of gene gain and loss are thus crucial for understanding the composition of pangenomes.

Figure 5. Simulated pangenomes for different distributions of rates of gene gain and loss. Each panel represents a simulation with a particular niche and population structure. **I**. 100

genomes from a single niche, 1000 genes with fitness effects drawn from a normal distribution
with zero mean and unit standard deviation). **II**. 10 genomes simulated from each of 10 niches.
1000 mostly-deleterious genes (fitness effects drawn from a normal distribution with -1 mean
and 0.5 standard deviation) can be gained in all niches. In each niche 100 genes (different for
each niche) are mostly beneficial (fitness effect drawn form a normal distribution with mean 2
and 0.5 standard deviation). **III.** 1 genome simulated from each of 100 niches. 1000 mostly-
deleterious genes (fitness effects drawn from a normal distribution with -1 mean and SD = 0.5)
can be gained in all niches. In each niche 1 gene (different for each niche) is mostly beneficial
(fitness effect drawn form a normal distribution with mean 2 and SD = 0.5). **IV**. 1 genome
simulated from each of 100 niches. 1000 genes with fitness effects drawn from a normal
distribution with zero mean and SD=1) can be gained in all niches. In each niche 1 gene
(different for each niche) is mostly beneficial (fitness effect drawn form a normal distribution
with mean = 2 and SD = 0.5). For each panel: **a**. Presence/absence of genes in the simulated
pangenome, present genes are indicated in grey. The orange line indicates the number of
strains in which a gene is present. **b**. Heatmap showing the sampled fitness effect of each
gene in each niche **c**. Density plots for the DFE of the gene pool (sampled from the specified
theoretical distribution, in blue) and for the fitness effects of genes actually present in the
pangenome (orange). **d**. Gene frequency distribution for the simulates pangenome. **e**. Gene
frequencies versus fitness effects for 1000 genes in the simulated pangenome (blue dots) and
the theoretical relationship (orange line). Rates of gene gain and loss are assumed to be equal
for all genes and niches ($r_g = r_l = 0.01$)
458
459 **Discussion**
460
We have presented a model to simulate prokaryote pangenomes that allows us to analyse the
effect of important parameters on gene frequencies and fitness contributions. We have shown
that fitness effects, rates of gene gain and loss and their respective distributions are crucial in
determining pangenome properties. Depending on assumptions and circumstances, the
model shows that accessory genes can be neutral, advantageous or even deleterious.
Therefore, this model contributes to reconcile previous models which suggest that
pangenomes are the result of purely neutral or purely selective processes, and that accessory
genes are mostly neutral or beneficial[11,15–19]. Our aim is to highlight factors that may be
important in shaping pangenomes and that perhaps have received little attention, and
therefore our model is intended as a framework to investigate how basic parameters can affect
gene content, rather than a detailed account of pangenome evolution. By assuming a regime
with strong selection and disregarding drift, our model lies at the opposite end to models that
assume pure neutrality in the evolutionary process. This assumption may be justified for the
large population sizes of many prokaryotes in which selection is expected to be very effective
and drift may be relatively weak, as suggested by high levels of codon adaptation in some
species and the general compactness of prokaryote genomes[12]. Together with the assumption
of equilibrium for gene frequencies, these simplifications allow a flexible framework where we
can incorporate many of the processes that influence pangenome properties.
479
It is known that the distribution of fitness effects of mutations or genes is a major determinant
of the levels of genetic polymorphism we should expect in populations[8,27], and we have shown
how this can impact the gene frequency distribution (Figure 3). Disregarding population
structure and variation in the rates of gene gain and loss, we find that as the proportion of
neutral and deleterious genes in the DFE increases, the proportion of low frequency genes in

the pangenome also increases. However, we show that the DFE is not the only determinant
of gene frequencies, and that rates of gene gain and loss can have a large influence on gene
content and on the types of genes we expect to find in pangenomes[31]. If these rates are low
relative to fitness effects, selection can maintain advantageous genes essentially fixed in the
population, and genes present at low frequencies will be nearly neutral. On the other hand, if
these rates are high, deleterious genes can be common in the population, while advantageous
genes may not reach fixation. Although, in reality, these rates may on average be low with
respect to fitness effects, they are likely to vary for different genes, genomes and species, and
the values of these parameters and their distributions may contribute significantly to gene
content variation in prokaryote genomes. In particular, genes with high gain rates can be
maintained in the population at high frequencies even if they are significantly deleterious. This
is likely the case for some MGEs and lysogenic viruses, what we call 'infectious' genes, which
are most likely deleterious but gained at high rates. For example, deleterious polymorphisms
can become prevalent in a bacterial population due to the effect of high recombination
mediated by conjugative plasmids[32], and inteins with high fitness costs seem to be maintained
in genomes[33].
501
Furthermore, the genetically-determined component of the rates of gene gain and loss could
itself be a target of natural selection, from the perspective of either the receiving cell or an
infectious gene. Besides selection on the rates of natural transformation and DNA
recombination, mechanisms that quickly recognise and destroy foreign DNA, such as
restriction modification and CRISPR systems may not only prevent viral infections, but also
reduce the rate of gene gain in cells that possess these systems, regardless of their fitness
effects. 'Addictive genes' are another example, where a gene (or operon), even if deleterious,
once acquired becomes essential and is difficult to lose. Such may be the case of anti-toxin
genes of toxin/anti-toxin systems and restriction enzymes in restriction modification
systems[34,35]. Another possible example of beneficial change in the loss rate of a gene may be
the expansion of transposon numbers within a genome, and the observed loss of sequences
responsible for transposition for a widely-spread gene encoding colistin resistance[36].
514
We have also highlighted the potential of niche-dependent fitness effects for maintaining
variability in gene content. Although we lack information of how prevalent these effects are,
we know that individuals from the same prokaryote species can survive and thrive in very
different environments[37]. Therefore, it is likely that these fitness by environment interactions
for particular genes are relatively common. Besides previously mentioned antibiotic genes,
certain metabolism genes are good candidates for having niche-specific benefits[38]. Examples
are the association of vitamin B5 synthesis genes with cattle-specific strains of
Campylobacter[39], and the acquisition by HGT of transporter-encoding genes with
environment-dependent fitness effects in yeasts[40]. Moreover, we do not really know how
environments are partitioned, and what may appear as a homogeneous niche may in fact
have distinct compartments with specific selective pressures. For example, pathogens
multiplying in different organisms, parts of the body and even different parts of the same organ
may be subject to different conditions in terms of nutrients, physicochemical properties and
immune responses[41,42]. Even apparently homogeneous environments such as the open ocean
are more heterogeneous than previously thought, having niches which vary significantly in
availability of resources and other environmental conditions[43]. In any case, the existence of
these 'gene by environment' effects could account for the presence of accessory genes that

may be highly advantageous under some, but not all, conditions. In fact, even gene
essentiality can be environment-dependent[44].

Overall, we argue that for an individual gene, its particular circumstances and details matter,
and therefore its contribution to fitness cannot be extrapolated from its population frequency
alone. However, in general, neutral or deleterious genes are unlikely to be part of the core
genome, unless populations are subjected to significant genetic drift, their gain rates are very
large or/and their loss rates very small. Besides these broad conclusions, it is currently difficult
to make strong claims as to whether accessory genes are on average advantageous or not.
Our model shows that very different assumptions can broadly replicate the patterns of gene
presence/absence we observe in nature. Therefore, the issue is not that the data do not fit our
models, but that we lack information to truly distinguish between the different possible
explanations. If we want to understand the average properties of pangenomes, we need more
information about the distributions of fitness effects and rates of gene gain and loss.
Furthermore, it is worth noting that even though we have simulated parameter values as
independent, this will not necessarily be the case for real genes. For example, genes that are
highly beneficial are also likely to be more abundant in the environment and may have higher
gain rates. On the other hand, deleterious genes present in MGEs might also be gained at
high rates. Since we lack information about the individual distributions of these parameters, it
is not surprising that we know little about their joint distributions.

In conclusion, further understanding of the existence and maintenance of pangenomes in
different prokaryote groups, requires not only genome sequences, but extensive metadata,
environmental parameters and phenotypic characteristics. We need to understand the extent
and nature of the structure of bacterial populations, and how much gene flow there is between
subpopulations. Finally, we need longitudinal information about how gene frequencies change
in the population, and the stability of particular polymorphisms and pangenomes in general.
Collecting all this information not only requires extensive environmental sampling but will also
likely rely on experiment inside and outside the lab to determine the values of parameters
under controlled conditions.

Methods

<u>Calculation of gene frequencies</u>
From Eq. 1 in the model description, we obtain the steady state gene frequency, x_{ss} , which
is described by a quadratic equation with one positive and thus meaningful solution:

$$x_{ss} = \frac{(s - r_g - r_l) + \sqrt{(r_g + r_l - s)^2 + 4sr_g}}{2s}$$
    Eq. 2

We use Equation 2 to calculate the expected frequency of a gene in the population. Although
x_{ss} is undefined when $s = 0$, this equation can be used to calculate equilibrium frequencies
for nearly neutral genes, with fitness effects arbitrarily close to zero. The equilibrium
assumption may not be entirely justified in real populations (in particular if gene gain and loss
rates are low and fitness effects are low), but it is a useful assumption that allows us to explore
how parameters affect gene frequencies and their distributions for large numbers of genes.

An analytic solution describing the temporal dynamics of gene frequency also exists for
Equation 1 (see supplementary information).
578
<u>Pangenome simulations</u>
To simulate pangenomes, we assume there is a finite 'gene pool', a set of n genes that can
be gained and lost from cells in the population (Figure 1b). If we make the simplifying
assumption that there is no interaction between genes, the gene frequency model described
above can be used to model multiple independent genes in the population and thus explore
the effects of model parameters on pangenomes. This assumption implies that there are no
epistatic effects between genes, and thus the parameters describing the frequency of one
gene are independent from those describing any other gene (Also see Supplementary
Information for derivation of the same equation from genotypes for two genes). For each
simulation, we assume values for each of the three model parameters (s, r_g, r_l) for each gene.
Depending on the simulation, the values can be identical for all the genes, or sampled from a
specified distribution. Using these values, we calculate the expected equilibrium frequency for
each gene using Equation 2. In order to simulate the presence of each gene in each of N
genomes, a uniformly-distributed random number in the [0,1] interval is generated, and the
presence of the gene is accepted if the random number is smaller than the calculated
equilibrium gene frequency.
595
<u>Simulations with niche-dependent fitness effects</u>
To simulate pangenomes from different niches, we first define a general gene pool with n
genes whose fitness coefficients are drawn from a specified distribution (usually with negative
average values). In each of k niches the fitness effects of the genes are assumed to be the
same as the generalised gene pool, except for a defined subset of genes whose fitness effects
are drawn from a different distribution (usually with positive average values). In the simulations
presented in Figure 5 we consider non-overlapping sets of n/k genes to have niche-
dependent fitness values. Finally, we simulate a specified number of genomes from each
niche as described above.
605
<u>Fitness contribution of a gene</u>
To define the average contribution of a gene to the population fitness, ϕ, we consider a sample
of N genomes. For a particular gene, we define z_i as a variable representing the presence or
absence of a gene in genome i, where its value is equal to 1 if the gene is present, and 0
otherwise. In this case, s_i represents the fitness coefficient of the gene in the niche from which
genome i is sampled. The fitness contribution of a gene ϕ is then defined as:

$$\phi = \frac{\sum_{i=1}^{N} z_i s_i}{\sum_{i=1}^{N} z_i}$$

Acknowledgements
We thank John Brookfield, Caroline Colijn, Jessica Stockdale and Fiona Whelan for comments
on the manuscript and members of the O'Connell and McInerney groups for discussion. JOMc
is supported by a BBSRC Responsive Mode Grant BB/N018044/2.

618
619
620
621

References

1. Vernikos, G., Medini, D., Riley, D. R. & Tettelin, H. Ten years of pan-genome analyses. *Curr. Opin. Microbiol.* **23**, 148–154 (2015).
2. Koonin, E. V. & Wolf, Y. I. Genomics of bacteria and archaea: The emerging dynamic view of the prokaryotic world. *Nucleic Acids Res.* **36**, 6688–6719 (2008).
3. Rocha, E. P. C. Neutral Theory, Microbial Practice: Challenges in Bacterial Population Genetics. *Mol. Biol. Evol.* **35**, 1338–1347 (2018).
4. Shapiro, B. J. The population genetics of pangenomes. *Nat. Microbiol.* **2**, 1574 (2017).
5. McInerney, J. O., McNally, A. & O'Connell, M. J. Reply to 'The population genetics of pangenomes'. *Nat. Microbiol.* **2**, 1575 (2017).
6. Vos, M. & Eyre-Walker, A. Are pangenomes adaptive or not? *Nat. Microbiol.* **2**, 1576 (2017).
7. Gogarten, J. P. & Townsend, J. P. Horizontal gene transfer, genome innovation and evolution. *Nat. Rev. Microbiol.* **3**, 679–687 (2005).
8. Vos, M., Hesselman, M. C., Te Beek, T. A., van Passel, M. W. J. & Eyre-Walker, A. Rates of Lateral Gene Transfer in Prokaryotes: High but Why? *Trends Microbiol.* **23**, 598–605 (2015).
9. Medini, D., Donati, C., Tettelin, H., Masignani, V. & Rappuoli, R. The microbial pan-genome. *Current Opinion in Genetics and Development* **15**, 589–594 (2005).
10. Welch, R. A. *et al.* Extensive mosaic structure revealed by the complete genome sequence of uropathogenic Escherichia coli. *Proc. Natl. Acad. Sci.* **99**, 17020–17024 (2002).
11. Baumdicker, F., Hess, W. R. & Pfaffelhuber, P. The infinitely many genes model for the distributed genome of bacteria. in *Genome Biology and Evolution* **4**, 443–456 (2012).
12. McInerney, J. O., McNally, A. & O'Connell, M. J. Why prokaryotes have pangenomes. *Nat. Microbiol.* **2**, 17040 (2017).
13. Lees, J. A. *et al.* Fast and flexible bacterial genomic epidemiology with PopPUNK. *Genome Res.* **29**, 304–316 (2019).
14. Sela, I., Wolf, Y. I. & Koonin, E. V. Genome plasticity, a key factor of evolution in prokaryotes. *bioRxiv* 357400 (2018). doi:10.1101/357400
15. Haegeman, B. & Weitz, J. S. A neutral theory of genome evolution and the frequency distribution of genes. *BMC Genomics* **13**, (2012).
16. Lobkovsky, A. E., Wolf, Y. I. & Koonin, E. V. Gene frequency distributions reject a neutral model of genome evolution. *Genome Biol. Evol.* **5**, 233–242 (2013).
17. Takeuchi, N., Cordero, O. X., Koonin, E. V. & Kaneko, K. Gene-specific selective sweeps in bacteria and archaea caused by negative frequency-dependent selection. *BMC Biol.* **13**, 1–11 (2015).
18. Corander, J. *et al.* Frequency-dependent selection in vaccine-associated pneumococcal population dynamics. *Nat. Ecol. Evol.* **1**, 1950–1960 (2017).
19. Niehus, R., Mitri, S., Fletcher, A. G. & Foster, K. R. Migration and horizontal gene transfer divide microbial genomes into multiple niches. *Nat. Commun.* **6**, 1–9 (2015).
20. Clarke, B. C. The evolution of genetic diversity. *Proc. R. Soc. London - Biol. Sci.* **205**, 453–474 (1979).
21. Brisson, D. Negative Frequency-Dependent Selection Is Frequently Confounding. *Front. Ecol. Evol.* **6**, 10 (2018).
22. Fullmer, M. S., Soucy, S. M. & Gogarten, J. P. The pan-genome as a shared genomic resource: mutual cheating, cooperation and the black queen hypothesis. *Front. Microbiol.* **6**, 1–5 (2015).
23. Polz, M. F., Alm, E. J. & Hanage, W. P. Horizontal gene transfer and the evolution of bacterial and archaeal population structure. *Trends Genet.* **29**, 170–175 (2013).
24. Vos, M., Hesselman, M. C., te Beek, T. A., van Passel, M. W. J. & Eyre-Walker, A. Rates of Lateral Gene Transfer in Prokaryotes: High but Why? *Trends in Microbiology*

23, 598–605 (2015).

25. Sela, I., Wolf, Y. I. & Koonin, E. V. Estimation of universal and taxon-specific
parameters of prokaryotic genome evolution. *PLoS One* **13**, 1–20 (2018).

26. Vaccaro, B. J. *et al.* Determining Roles of Accessory Genes in Denitrification by
Mutant Fitness Analyses. (2015). doi:10.1128/AEM.02602-15

27. Eyre-Walker, A. & Keightley, P. D. The distribution of fitness effects of new mutations.
Nat. Rev. Genet. **8**, 610–618 (2007).

28. Sela, I., Wolf, Y. I. & Koonin, E. V. Theory of prokaryotic genome evolution. *Proc.*
Natl. Acad. Sci. **113**, 11399–11407 (2016).

29. Hernando-Amado, S., Sanz-García, F., Blanco, P., Jos´, J. & Martínez, J. L. Fitness
costs associated with the acquisition of antibiotic resistance. *Essays Biochem.* 61–98
(2017). doi:10.1042/EBC20160057

30. Durão, P., Balbontín, R. & Gordo, I. Evolutionary Mechanisms Shaping the
Maintenance of Antibiotic Resistance. *Trends Microbiol.* **26**, 677–691 (2018).

31. Shapiro, B. J. How clonal are bacteria over time? *Curr. Opin. Microbiol.* **31**, 116–123
(2016).

32. Maddamsetti, R. & Lenski, R. E. Analysis of bacterial genomes from an evolution
experiment with horizontal gene transfer shows that recombination can sometimes
overwhelm selection. *PLoS Genet.* **14**, 1–30 (2018).

33. Naor, A. *et al.* Impact of a homing intein on recombination frequency and organismal
fitness. *Proc. Natl. Acad. Sci.* **113**, E4654–E4661 (2016).

34. Mruk, I. & Kobayashi, I. To be or not to be: Regulation of restriction-modification
systems and other toxin-antitoxin systems. *Nucleic Acids Res.* **42**, 70–86 (2014).

35. Fullmer, M. S., Ouellette, M., Louyakis, A. S., Papke, R. T. & Gogarten, J. P. The
patchy distribution of restriction–modification system genes and the conservation of
orphan methyltransferases in halobacteria. *Genes (Basel).* **10**, (2019).

36. Wang, R. *et al.* The global distribution and spread of the mobilized colistin resistance
gene mcr-1. *Nat. Commun.* **9**, 1–9 (2018).

37. Tamames, J., Abellán, J. J., Pignatelli, M., Camacho, A. & Moya, A. Environmental
distribution of prokaryotic taxa. *BMC Microbiol.* **10**, (2010).

38. Goyal, A. Metabolic adaptations underlying genome flexibility in prokaryotes. *PLOS*
Genet. **14**, e1007763 (2018).

39. Sheppard, S. K. *et al.* Genome-wide association study identifies vitamin B5
biosynthesis as a host specificity factor in Campylobacter. *Proc. Natl. Acad. Sci.* **110**,
11923–11927 (2013).

40. Milner, D. S. *et al.* Environment-dependent fitness gains can be driven by horizontal
gene transfer of transporter-encoding genes. *Proc. Natl. Acad. Sci. U. S. A.*
201815994 (2019). doi:10.1073/pnas.1815994116

41. Sheppard, S. K., Guttman, D. S. & Fitzgerald, J. R. Population genomics of bacterial
host adaptation. *Nat. Rev. Genet.* **19**, 549–565 (2018).

42. Jorth, P. *et al.* Regional Isolation Drives Bacterial Diversification within Cystic Fibrosis
Lungs. *Cell Host Microbe* **18**, 307–319 (2015).

43. Shapiro, B. J. *et al.* Population Genomics of Early Events in the Ecological
Differentiation of Bacteria. *Science (80-.).* **336**, 48–51 (2012).

44. Martínez-Carranza, E. *et al.* Variability of bacterial essential genes among closely
related bacteria: The case of Escherichia coli. *Frontiers in Microbiology* **9**, 1059
(2018).