# Genetic and epigenetic characteristics associated with

# the rapid radiation of *Aquilegia* species

Zhen-Hui Wang[1,#], Tianyuan Lu[2,3,4,#], Ming-Rui Li[2], Ning Ding[2], Li-Zhen Lan[2], Xiang Gao[5], Ai-Sheng Xiong[6,*],

Jian Zhang[1,7,*] and Lin-Feng Li[2,*]

[1]Faculty of Agronomy, Jilin Agricultural University, Changchun 130118, China;

[2]Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, School of Life

Sciences, Fudan University, Shanghai 200438, China;

[3]McGill University and Genome Quebec Innovation Center, Montreal H3A 0G1, Quebec, Canada;

[4]Lady Davis Institute, SMBD JGH, Montreal H3A 1A3, Quebec, Canada;

[5]Key Laboratory of Molecular Epigenetics of the Ministry of Education, Northeast Normal University,

Changchun 130024, China.

[6]State Key Laboratory of Crop Genetics and Germplasm Enhancement, Ministry of Agriculture and Rural

affairs, Key Laboratory of Biology and Germplasm Enhancement of Horticultural Crops in East China,

College of Horticulture, Nanjing Agricultural University, 1 Weigang, 210095 Nanjing, China

[7]Department of Biology, University of British Columbia, Okanagan, Kelowna, Canada

**\*Correspondence author:**

**Lin-Feng Li (**lilinfeng@fudan.edu.cn**)**

**Jian Zhang** (zhangjian@jlau.edu.cn)

**Ai-Sheng Xiong** (xiongaisheng@njau.edu.cn)

**Abstract**

Elucidating the genetic and epigenetic bases underlying species diversification is crucial to understanding the evolution and persistence of biodiversity. As a well-known horticultural plant grown worldwide, the genus Aquilegia (columbine) is also a model system in adaptive radiation research. In this study, we surveyed the genomes and DNA methylomes of ten representative Aquilegia species from the Asian, European and North American lineages. Our inferences of the phylogenies and population structure revealed clearly high genetic and DNA methylomic divergence across the three lineages. By multi-levelled genome-wide scanning, we identified candidate genes exhibiting lineage-specific genetic or epigenetic variation patterns that are signatures of inter-specific divergence. We demonstrated that these species diversification-associated genetic variations and epigenetic variabilities were partially independent but were both functionally related to various biological processes vital to adaptation, including stress tolerance, cell reproduction and DNA repair. Our study provides an exploratory overview of how the established genetic and epigenetic signatures are associated with the rapid radiation of Aquilegia species.

**Key words:** Adaptive radiation; Aquilegia; Selection; Whole-genome sequencing; DNA methylation

**Introduction**

Adaptive radiation is the rapid diversification of a single ancestral species into a vast array of common descendants that inhabit different ecological niches or use a variety of resources, but differ in phenotypic traits required to exploit diverse environments[1–4]. Disentangling the evolutionary mechanisms underpinning adaptive radiation is fundamental to understanding the evolution and persistence of biodiversity[5,6]. This has been a key focus of many studies which were investigating different animal and plant lineages that diversified through adaptive radiation, including Hawaiian silversword, Caribbean anoles, Darwin's finches, and African cichlids[7–10]. However, it remains under-investigated as to why some lineages could diversify rapidly but their close relatives or other sympatrically distributed lineages did not. In the past decades, accumulating evidence from diverse radiation lineages suggest that both the extrinsic environmental factors (e.g., resource availability) and genetic variations can determine the rate and volume of species diversification[11]. Among the environmental triggers, ecological opportunity is considered as the primary mechanism that causes rapid adaptive radiation through acquisition of key innovations, penetration of new environments and extinction of competitors[2,12]. On the other hand, new species also arise as a result of new genetic variations being preserved which could ultimately influence the phenotypic disparity, where natural selection act on, among closely related species[13]. In the rapid speciation of the African cichlid fishes, extrinsic environmental factors (e.g., ecological specialization) and genetic mechanisms (e.g., adaptive introgression) acted together to provoke the repeated adaptive radiation in geographically isolated lakes[7,11,14,15].

The genus *Aquilegia* L. (columbine) is a well-recognized model system to study the evolutionary mechanisms underlying adaptive radiation[16,17]. This genus includes approximately 70 recently diversified species that are widely distributed in the temperate zones of North America and Eurasia[18]. Phylogenetic and geographic inferences have illustrated two independent adaptive radiations of North American and European lineages from the ancestral Asian species[17,19]. For example, floral diversification of the North American *Aquilegia* species is highly correlated with the pollinator specialization[20–23]. In contrast, ecological adaptation and geographic isolation are considered as the major driving forces promoted rapid radiation of the European species[17,24]. In Asia, changes in pollinator and ecological habitats are both proposed to be the underpinning mechanisms that resulted in the diversification of more than 20 morphologically distinct species[25,26]. These Asian *Aquilegia* species constitute four highly divergent lineages corresponding to their geographic origins and have evolved relatively independently[25,26]. Despite this well-described evolutionary history and crucial role played by environmental factors, how

genetic and epigenetic factors are involved in the rapid speciation in this genus remains poorly
investigated.

In this study, the main objective is to survey the genomes and DNA methylomes of 36 accessions
from ten worldwide *Aquilegia* species from the Asian, European and North American lineages. Among
the Asian species, four phylogenetically distinct species (*A. japonica, A. oxysepala, A. yabeana, and A.*
*viridiflora*) were selected according to their geographic distributions and ecological habitats. *Aquilegia*
*japonica* and *Aquilegia. oxysepala* are sister species inhabiting alpine tundra and low altitude forest
niches in northeastern China, respectively[25,27]. Our previous studies have documented that natural
selection during ecological specialization together with genetic drift under geographic isolation caused
the rapid evolution of reproductive isolation between these two species[25,28]. Here, we further
investigated how diverse evolutionary driving forces shaped genetic and epigenetic architectures of the
two species in the processes of speciation and adaptation. In addition, we also evaluated patterns of
nucleotide variation and cytosine methylation in the *A. yabeana* and *A. viridiflora*. The former species
shares highly similar morphological traits and ecological niches with the *A. oxysepala* but is allopathically
distributed in northern China. In contrast, while the *A. viridiflora* is sympatrically distributed with *A.*
*yabeana* and *A. oxysepala* in northern and northeastern China, it often occupies rocky and sandy
ecological niches. As a supplementary, we also examined nucleotide and cytosine methylation variation
patterns of the North American and European lineages.  Our study will provide a genome-wide view of
how the specific genomic and epigenomic variation patterns are correlated with the diversification of
Aquilegia species.

**Results**

**Population structure and nucleotide variation pattern**

Neighbor-joining (NJ) trees were reconstructed for the 36 *Aquilegia* accessions based on 689,123

homozygous SNPs. The phylogenic analysis suggested that these accessions of the ten species formed

three distinct lineages corresponding to their geographic origins (**Figure 1a** and **Figure S1**). In brief, all 22

accessions of the four East Asian species, *A. japonica*, *A. oxysepala*, *A. yabeana* and *A. viridiflora*,

clustered as a monophyletic lineage, with the first two species and their hybrid forming a clade and the

last two species grouping as a sister clade. In contrast, the West Asian species *A. fragrans* clustered with

the geographically adjoining European species. The principal component analysis (PCA) and population

structure inferences also revealed distinct genetic structure of the three phylogenetic lineages (**Figure**

**1b** and **c**). It should be noted that one *A. alpina* var. alba accession shared the same ancestral genetic

cluster with the North American lineage, while the putative hybrid of the *A. oxysepala* and *A. japonica*

possessed an admixed genetic background (**Figure 1b** and **c**).

To further gain an insight into genome-wide nucleotide variation pattern of the ten *Aquilegia*

species, we calculated nucleotide diversity ($\pi$) and genetic divergence ($F_{ST}$) for each chromosome and for

100-kb sliding windows, respectively. Among the three phylogenetic lineages, the Asian *Aquilegia*

species harbored the highest nucleotide diversity compared to the European and North American

lineages across the seven chromosomes (**Figure S2**). By comparing the nucleotide diversity for each 100-

kb sliding window, we observed a moderate correlation of genome-wide variation pattern among the

three lineages (Spearman R = 0.42-0.56) and a high correlation between the *A. oxysepala* and *A.*

*japonica* (Spearman R = 0.70) (**Figure S3**). In particular, 116 of 241 low genetic diversity genomic regions

(LDGRs, with 5% lowest $\pi$) were shared by at least two of the three lineages (**Figure S4**). Between the *A.*

*oxysepala* and *A. japonica*, while we defined 148 LDGRs and 148 high divergence genomic regions

(HDGRs, with 5% highest $F_{ST}$), only seven candidate genomic regions overlapped (**Table S1**).

**Figure 1**. **Phylogenetic relationship and population structure of the ten worldwide *Aquilegia* species**.

(a) Phylogenetic tree of the 36 accessions constructed by neighbor-joining algorithm based on 689,123

whole-genome SNPs. (b) PCA reveals genetic similarity within each of the three lineages and genetic

disparity between lineages based on 15,988 LD-pruned SNPs. Ellipses of each lineage denote 99%

confidence region estimated from distribution of the first two principal components. (c) Population

admixture of the 36 *Aquilegia* accessions.

**Identification of the genomic regions indicating selection pressure and highly impactful genetic**

**variations**

Candidate genes or genomic regions associated with adaptive divergence were determined from three

perspectives. First, we considered genes localized within the regions that showed low intra-specific

diversity but high inter-specific divergence to be representative of intra-specific genetic differences. We

thus identified 23 genes from the above seven candidate genomic regions that were both HDGRs and

LDGRs shared by *A. oxysepala* and *A. japonica* (**Table S1**). Genes within these genomic regions were

functionally associated with meiotic nuclear division, adenine methyltransferase and basic cellar

activities.

While the first strategy mainly relied on genome-wide scanning for 100-kb non-overlapping sliding

window, we also employed a functional annotation-based approach to identify highly impactful

conservative clade-specific variations (CCVs) from both the within and between lineage comparisons.

Our results revealed that a considerable proportion (17.9-40.5%) of the CCVs were identified in the gene

body regions (**Table S2**). We then examined the potential functional impacts of genes harboring these

identified CCVs. Between the *A. oxysepala* and *A. japonica*, the CCV-carrying genes were enriched in

several vital biological pathways related to cell reproduction, including telomere maintenance, DNA

repair, and DNA helicase activity (**Figure 2** and **Table 1**). For example, two candidate genes

(*Aqcoe6G160300* and *Aqcoe7G062500*) coding for *Xklp2* (*TPX2*) were functionally correlated with spindle

assembly during the mitotic process (27, 28). Among the three phylogenetic lineages, the CCVs-

harboring genes were also functionally involved in the mitotic chromosome condensation, DNA ligase

activity and aminopeptidase activity (**Figure 2** and **Table 1**). For instance, two CCV-containing genes

(*Aqcoe2G276600* and *Aqcoe1G273400*) encoding DNA mismatch repair proteins *MutS*/*MSH* and *MutS2*

(ref. 29) carried one Asian-specific-to-American frameshift variant.

Thirdly, we also derived pair-wise synonymous ($d_S$) and non-synonymous ($d_N$) mutation rate to

identify genes informative of positive or purifying selection pressure. We found that species within the

Asian lineage experienced significantly stronger positive ($d_N/d_S > 1$) and purifying ($d_N/d_S < 1$) selection

pressures compared to the European and North American lineages (Wilcoxon rank sum test, all

Bonferroni-corrected *p* values < $1.5 \times 10^{-16}$) (**Figure S5**). Likewise, the European species showed

 significantly stronger purifying selection (Wilcoxon rank sum test, Bonferroni-corrected $p$ value = 7.8x10$^{-8}$) compared to the North American species.

**Figure 2**. **Functional enrichment of genes harboring highly impactful CCVs and DMGs.** CCV-containing

genes specific to either of the two lineages/species being compared were merged to construct a target

gene set. Ratio denotes proportion of CCV-containing genes or DMGs in the corresponding gene set of

interest. Absence of dot indicates no significant enrichment.

**Table 1**. Information of the high-impact conservative clade-specific variants (CCVs) in the cell

reproduction related genes.

| Gene | Variant-carrying | Reference | Chromosome | Position | Reference allele | Variant | Annotation | Gene function |
|------|------------------|-----------|------------|----------|------------------|---------|------------|---------------|
| Aqcoe1G273400 | Asian | American | Chr1 | 18994915 | GAA | GAAA | frameshift | DNA mismatch repair protein *MutS2* |
| Aqcoe2G151500 | European | American | Chr2 | 15305837 | A | G | splicing | PIF1-like helicase |
| | European | American | | 15307442 | A | C | stop gain | |
| | European | American | | 15309865 | AATATATAT | AATATATATAT | frameshift | |
| | European | Asian | | 15307442 | A | C | stop gain | |
| | European | Asian | | 15309865 | AATATATAT | AATATATATAT | frameshift | |
| | *A. oxysepala* | *A. japonica* | | 15305837 | A | G | splicing | |
| | *A. oxysepala* | *A. japonica* | | 15309267 | AT | A | frameshift | |
| Aqcoe2G177700 | European | American | Chr2 | 21794397 | TATGCACCAAAGGTATCACGATGC | TATGC | frameshift | PIF1-like helicase |
| | European | American | | 21794979 | TT | TTGT | frameshift | |
| | European | Asian | | 21794397 | TATGCACCAAAGGTATCACGATGC | TATGC | frameshift | |
| | *A. oxysepala* | *A. japonica* | | 21795089 | CA | C | frameshift | |
| Aqcoe6G208600 | European | American | Chr6 | 15364081 | A | ATCTCTTCG | frameshift | PIF1-like helicase |
| | European | Asian | | 15364081 | A | ATCTCTTCG | frameshift | |
| | *A. japonica* | *A. oxysepala* | | 15364330 | TAA | TA | frameshift | |
| Aqcoe6G253800 | European | American | Chr6 | 22789898 | C | T | stop gain | DNA helicase |
| | European | American | | 22790012 | G | A | splicing | |
| | European | Asian | | 22789898 | C | T | stop gain | |
| | *A. japonica* | *A. oxysepala* | | 22790012 | G | A | splicing | |
| Aqcoe2G276600 | Asian | American | Chr2 | 33314422 | AGGGGG | AGGGGGG | frameshift | DNA mismatch repair protein *Msh6* |
| Aqcoe6G160300 | *A. japonica* | *A. oxysepala* | Chr6 | 9414625 | G | A | stop gain | *TPX2* |
| Aqcoe7G062500 | *A. oxysepala* | *A. japonica* | Chr7 | 3789055 | G | A | stop gain | cell cycle regulated microtubule associated protein |

**CG methylation patterns and differentially methylated genes**

In parallel with the above genomic analyses, we also investigated CG methylation pattern of the

representative *Aquilegia* species. Despite variability across the 36 *Aquilegia* accessions, the North

American, Asian and European species showed no distinguishable differences (*t* test, all Bonferroni-

corrected *p* values > 0.01) in overall percentage of methylated cytosines (**Figure 3a**). We then performed

PCA to examine the CG-cytosine methylomic diversity of all the *Aquilegia* accessions. The resulting

overall methylation pattern highly resembled the above genomic inferences, with the European and

American species forming two distinct groups and the four Asian species forming three separate clusters

(**Figure S6**). We then assessed the CG methylation patterns for the European and North American

lineages as well as the three Asian species (*A. japonica*, *A. oxysepala* and *A. viridiflora*) separately.

Consistent with the described genomic features, heterogeneous pattern of the CG methylation was also

observed for the seven chromosomes, with the chromosome 4 demonstrating obviously higher overall

CG methylation divergence compared to the other six chromosomes (**Figure 3b**). We further quantified

CG methylation level deposited in the genic regions, putative *cis*-regulatory regions and CG island,

respectively. In genic and regulatory regions, all three lineages shared similar modification patterns with

apparent depletion of CG methylation around the transcription start site (TSS) and transcription end site

(TES) (**Figure 3c**). However, the American lineage exhibited hyper-methylation (more than 10%) around

the center of CG islands and a more drastic decrease throughout the CG island shores compared to the

European and Asian species (**Figure 3d**).

To examine the biological impacts of CG methylation on the species diversification, differentially

methylated regions (DMRs) and differentially methylated genes (DMGs) were identified for both within-

and between-lineage comparisons, respectively (**Tables S3** and **S4**). Within the Asian lineage, 3,622

DMRs in 2,899 DMGs were identified between the *A. japonica* and *A. oxysepala*. Functional enrichment

of these DMGs indicated that the two species may have different activities in photosynthesis-related

pathways, including photosystem I, photosynthesis and chloroplast (**Figure 2**). For example, two

photosynthesis-related genes, *PsaA*/*PsaB* and *CemA*, showed significantly differential methylation

between the two species in the genic regions (**Figure S7a** and **b**). At the inter-lineage level, apparently

more DMGs were identified between the North American and European species (6,087 genes)

compared to those of between the two lineages and Asian species (3,308-5,003 genes) (**Table S3** and **S4**).

DMGs characterized from the inter-lineage comparisons were mainly involved in the plant growth (e.g.,

response to auxin) and defense, response to biotic stimulus and wounding (**Figure 2**).

We then examined whether the candidate genes (CCV-carrying genes and DMGs) superimposed on

the same signature of natural selection. We found while a considerable proportion of the candidate

genes were shared for each of the genetics- and epigenetics-based assessments (**Figure S8**), they

showed a segregated distribution pattern across all comparisons (**Figure S9**). Likewise, the Gene

Ontology (GO) enrichment analyses of the candidate gene identified from the genetic and epigenetic

levels were enriched in functionally complementary pathways (**Figure 2**), suggesting co-existence of

different underlying evolutionary mechanisms.

**Figure 3**. **Patterns of cytosine methylation for the ten worldwide *Aquilegia* species.** (a) Genome-wide

cytosine methylation levels of 36 accessions. (b) MDI illustrates chromosome-level CG methylation

similarity. *Aquilegia viridiflora* was used as the reference. (c) CG methylation profiling in genic region

across the four *Aquilegia* groups. Each row represents one genic region starting at 5-kb upstream of its

TSS and terminating at 5-kb downstream of its TES, sorted by mean methylation level of all analyzed CG

loci. Gene body regions were scaled to have the same length. (d) CG methylation profiling in and around

CG islands.

**Association between epigenetic variability and genetic variations**

Since both genetic variation and differential CG methylation seemed to have crucial and multifaceted influences on the adaptation of the ten *Aquilegia* species, we wondered whether differential epigenetic modifications were dependent on genetic variations. Among the 588,659 CG loci examined, 224,222 (38.09%) carried a CG-loss variation. We then illustrated epigenetic variability for the variation-carrying and non-variant CG loci, respectively. As shown in Figure 4, genetic-epigenetic associations of varying magnitude were observed in both types of CG loci. The variation-carrying CG loci conveyed information that highly resembled their genetic background. The overall methylation pattern was highly conserved within the same species but exhibited obvious divergence across the ten *Aquilegia* species (**Figure 4a**). In contrast, CG methylation divergence at the non-variant CG loci varied with higher variability at both the intra- and inter-specific levels (**Figure 4b**). By examining the correlation of genetic variability and cytosine methylation, we found that CG methylation divergence at variation-carrying CG-site was largely attributable to the CG-loss variations (**Figure 4c**). In particular, 75% of the CG-loss variations occurring at the most highly variable CG-methylated dinucleotides could explain at least 75% of the total epigenetic variability *per se*. Nevertheless, there was still a considerable proportion of epigenetic variability that could not be sufficiently explained by the variant-CG locus (**Figure 4d**).

We also attempted to identify *cis*-driver mutations for each of the 1,229 DMRs between the *A. japonica* and *A. oxysepala*. Our results revealed that only 568 out of the 1,229 (46.2%) DMRs were significantly associated with at least one genetic variation inside or around a 500 base-pair (bp) upstream/downstream genomic region, even under the least stringent *p* value threshold ($5 \times 10^{-5}$), indicating that the epigenetic changes were only partially dependent on *cis*-genetic driving mutations (**Figure 4e**). Moreover, we observed weak yet significant associations between differential CG methylation and selection pressure. In most inter-lineage comparisons, DMGs were significantly more prone to be under positive selection than non-DMGs (**Table 2**), suggesting that epigenetic modifications could probably assist selection pressure in shaping genotypes. In contrast, DMGs were significantly less prone to be under purifying selection (**Table 2**).

**Figure 4**. **Association between CG-loss variations and epigenetic variability**. (a) Top 3,000 most variable CG loci containing CG-loss variations. (b) Top 3,000 most variable non-variant CG loci across 36 accessions show clade-specific methylation patterns. CG methylation in the hybrid tends to be neutralized possibly due to heterozygosity. (c) Linear regression demonstrates that CG-loss variations explain a large proportion of CG methylation variation. (d) Summary of composition of each category with regard to whether each CG locus contains a CG-loss variation. Epigenetic variability was determined by standard deviation in methylation β value across all 36 accessions. CG loci with top 10,000, 10,001-50,000 and 50,001-150,000 largest standard deviation were ordinally labelled as possessing "very high", "high" and "moderate" variability respectively. The rest CG loci were labelled as possessing "low" variability. (e) Association test shows most DMRs were independent of *cis*-acting SNPs. Results under different significance levels are compared in this exploratory analysis.

344 **Table 2.** Significant correlation between differential methylation and natural selection.

| Type of selection | Differential methylation | Jap-Oxy[*] | Jap-Ame | Jap-Eur | Oxy-Ame | Oxy-Eur | Ame-Eur |
|---|---|---|---|---|---|---|---|
| Positive selection | DMG | 7.2% | 7.3% | 11.9% | 6.7% | 8.4% | 8.9% |
| | non-DMG | 4.4% | 5.0% | 5.6% | 4.7% | 5.4% | 5.4% |
| | $p$ value | 0.11 | 7.3e-02 | 3.9e-05 | 6.7e-02 | 1.8e-02 | 2.8e-04 |
| Purifying selection | DMG | 3.1% | 1.8% | 2.4% | 2.3% | 2.0% | 1.9% |
| | non-DMG | 4.3% | 4.3% | 4.7% | 4.9% | 5.1% | 4.0% |
| | $p$ value | 0.53 | 3.2e-02 | 9.1e-02 | 1.3e-02 | 8.4e-03 | 1.0e-02 |

[*]: Each percentage represents the proportion of genes belonging to either DMGs or non-DMGs
compared between the two corresponding clades that are under corresponding or higher strength of
positive selection. For example, 7.2% indicates that 7.2% DMGs compared between *A. japonica* and *A.*
*oxysepala* are under strong selection; 4.4% indicates that 4.4% non-DMGs compared between these two
species are under strong selection. *p* values were obtained from Chi-square tests and were not adjusted
for multiple testing due to dependence arising from overlapping gene sets.
Jap: *A. japonica*; Oxy: *A. oxysepala*; Ame: American; Eur: European

**Discussion**

**Genetically determined mechanisms associated with the rapid diversification of *Aquilegia* species**

Elucidating evolutionary mechanisms underpinning species diversification is crucial to understanding the evolution and persistence of biodiversity[2,5,6]. The genus *Aquilegia* provides an ideal system to address how the diverse evolutionary mechanisms promoted rapid adaptive radiation[16,17]. Although various environmental conditions related to ecological opportunities, such as shifts in pollinator and habitat, have been proposed to facilitate the evolution of reproductive isolation[21,25], genetic basis associated with the rapid diversification of *Aquilegia* species has still remained largely unclear. In this study, we surveyed the genomes of ten worldwide *Aquilegia* species to address whether specific genetic architectures have been involved in the rapid species diversification. Broadly consistent with previously inferred phylogeny[17,19,25,30], the ten *Aquilegia* species from Asia, Europe and North America formed three phylogenetically independent lineages corresponding to their geographic origins. This attribute renders the *Aquilegia* species a suitable system to identify genomic variations associated with the repeated adaptive speciation by extensive comparisons from different facets.

It has been proposed that if a genetic factor is the potential determinant promoting adaptive speciation, one would expect to identify specific genetic architectures in the diversified lineages[8,14,31]. In Darwin's finches, for example, polyphyletic topology was observed as a general pattern in 14 morphologically distinct species, phenotypic diversity of the beak shape was mainly determined by natural selection acting on the *ALX1* gene during the ecological specialization process[8]. A similar phenomenon was observed in the East African cichlid fish where the radiating lineages are more dynamic in terms of gene content and transcriptomic landscape compared to their non-radiating relatives[14,31]. In this study, the genome-wide nucleotide variation pattern highly reflects the evolutionary history that the Asian, European and North American *Aquilegia* species have clearly diverged and evolved allopatrically in respective geographic regions. This also suggests that a considerable proportion of the genetic variations and changes in environment are intertwined during the diversification process. As expected, our genome-wide scanning for selection signatures revealed distinct positive and purifying selection modes in the intra- and inter-lineage comparisons. More importantly, the CCV-carrying genes

identified from the three lineages are associated with cell reproduction (e.g., telomere maintenance and

mitotic chromosome condensation) and other functionally important traits. Similar genomic feature was

also observed in *A. japonica* and *A. oxysepala*. Our previous studies have demonstrated that natural

selection and genetic drift together resulted in the rapid evolution of reproductive isolation[25,30]. Here,

we further demonstrate that candidate genes involved in the adaptive speciation are functionally

enriched in the pathways related to cell reproduction (e.g., telomere maintenance), stress tolerance

(e.g., response to wounding) and basic cellular activities. It should be noted that although a majority of

the enriched pathways are specific to each comparison, enrichment of cell reproduction-related

pathways (e.g., telomere maintenance, DNA repair and DNA helicase activity) and stress tolerance are

shared in the intra- and inter-lineage comparisons. Taken together, these findings indicate that specific

genetic determinants might have conferred high adaptability to the Aquilegia species to cope with

different environmental conditions.

**Evolutionary potential of cytosine methylation in the adaptation of *Aquilegia* species**

The role of epigenetic modification in the long-term evolutionary process has long been debated[32–34]. It

has been proposed that epigenetic variations are frequently under the genetic control which can alter

rapidly as a result of environmental induction and stochastic epimutation[35,36]. Nevertheless, it has also

been recognized that some epigenetic variations can persist over generations and be highly correlated

with phenotypic diversity[32]. As illustrated in Arabidopsis, changes in cytosine methylation can produce

meiotically stable epialleles, which could eventually lead to phenotypic diversity in the absence of

genetic variations[37–39]. Here, we assessed whether the epigenetic modifications were also associated

with the adaptive speciation of the *Aquilegia* species. Consistent with the genomic features detailed

above, high divergence of cytosine methylation was observed across the Asian, European and North

American lineages. Notably, differential cytosine methylation was not only found across the seven

chromosomes but also evident in the gene body of DMGs and CG island region among the three lineages.

Particularly, functional enrichment analyses identified significant associations with adaptation-related

traits, including plant growth, stress tolerance and basic cellular activities. For example, the candidate

DMGs identified between the *A. japonica* and *A. oxysepala*, showed significant enrichment in pathways

related to diverse important phenotypic traits, such as photosynthesis, embryo development and

response to auxin. These features indicated that epigenetic factors might also play a role in response to

diverse environmental conditions.

We noted that some candidate genes and enriched pathways had shared hotspots of both genetic
and epigenetic disparities, especially those related to cell reproduction, plant growth and stress
tolerance. Many studies based on human and mouse have shown that genetic variations can manipulate
*cis*-CG methylation at specific loci to further influence phenotypes, where CG methylation serves as a
mediator[40,41]. By analyzing the associations between genetic and epigenetic variability, we conclude that
while many CG-loss variations can directly lead to depletion of CG methylation, a lot of DMRs are not
manipulated by any *cis*-variations. Since gene body CG methylation in plants generally stabilizes gene
expression and is positively correlated with gene expression[42–45], differential methylation in our study is
indicative of likely differential amount of gene products. Based on these attributes together with the
plausible associations between differential methylation (e.g., DMGs) and positive selection (e.g., $d_N/d_S$),
we propose that epigenetic modification may be a complementary mechanism facilitating phenotypic
diversity of the *Aquilegia* species.
443
**Limitations and future directions**
We realized that this study has some limitations. Fistful, the small sample size in this study may
introduce bias and inflation of false positives, and we postulate that our findings should be interpreted
carefully and considered exploratory. When the association between genetic divergence and
evolutionary events is investigated, it is impossible to deny the roles of other evolutionary forces. We
acknowledge that the lineage-specific allele frequencies are possibly consequences of genetic drift, and
genetic hitchhiking may lead to identification of candidate genes residing in neighboring genomic
regions representing the other driving forces. Therefore, we claim that the candidate genes identified to
be associated with adaptive radiation do not necessarily point towards causal evolutionary mechanisms.
They may also be by-products of the long-term process of adaptive radiation. In addition, we never than
less only focused on analysis of CG methylation as puzzles persist regarding the functional roles of CHG
and CHH methylation. We also expect that future studies with larger sample sizes will be able to
improve the statistical power and investigate *trans*-genetic control.

**Methods and Materials**

**Sample collection, DNA extraction and whole-genome sequencing**

In this study, a total of 36 accessions from ten worldwide *Aquilegia* species were collected (**Table S5**).

Among the Asian species, four phylogenetically distinct species (*A. japonica*, *A. oxysepala, A. yabeana*,

and *A*. *viridiflora*) are selected according to their geographic distributions and ecological habitats.

*Aquilegia japonica* and *A. oxysepala* are sister species inhabiting alpine tundra and low altitude forest

niches in northeastern China, respectively[25,46]. Eighteen accessions were collected to represent these

two Asian species and their putative hybrid. In addition, four accessions were collected from the other

two Asian species, *A. yabeana* and *A. viridiflora*. The former species shares highly similar morphological

traits and ecological niches with *A. oxysepala* but is allopatrically distributed in the northern China. In

contrast, *A. viridiflora* is sympatrically distributed with *A. yabeana* and *A. oxysepala* in the northern and

northeastern China, but often occupies rocky and sandy ecological niches. Furthermore, six and eight

accessions were sampled from the European and North American lineages, respectively. All the 36

accessions were grown in green house under the same conditions (25°C/12 hours, 16°C/12 hours).

Genomic DNA was extracted from fresh mature leaves using TianGen plant genomic DNA kit. Whole

genome resequencing and bisulfite sequencing were performed on the extracted genomic DNA using

the Illumina X-ten platform (Illumina, California, USA). Short-insert (350 bp) DNA libraries of all

accessions were constructed by NovoGene (NovoGene, Tianjin, China). Genome assembly of an admixed

species *A. coerulea* "Goldsmith" was obtained from Phytozome v12.1 (https://phytozome.jgi.doe.gov) as

the reference genome[16].

**Sequence assembly, functional annotation and genetic diversity**

Whole genome sequences of each accession were aligned against the reference genome using default

settings of the BWA-MEM algorithm implemented in Burrows-Wheeler Aligner (BWA)[47]. Raw assemblies

were realigned using IndelRealigner provided in the Genome Analysis Tool Kit by default settings[48].

Single nucleotide polymorphisms (SNPs) and insertions/deletions (INDELs) were reported using

SAMtools[49]. Only the high-quality variants (SNPs and INDELs) (read depth > 3, mapping quality > 20 and

missing allele < 1%) were retained for subsequent population genomics analyses. Genomic annotation of

the identified variants was reported for each of the 36 samples separately. Functional annotation of

each identified variant was performed using SnpEff, based on the reference genome[50].

To infer the phylogenetic relationship between the ten *Aquilegia* species, NJ trees were

reconstructed for each chromosome and the whole genome dataset using MEGA 7(ref. 51). PCA was

carried out to examine the genetic diversity of the 36 *Aquilegia* accessions[52]. Ancestral components

were estimated using ADMIXTURE[53] with different number of populations ranging from one to ten.

Optimal population composition with the least 5-fold cross-validation error was selected to decompose

ancestral admixture. To obtain the genome-wide nucleotide variation pattern, nucleotide diversity ($\pi$)

and genetic differentiation (Weir and Cockerham's $F_{ST}$) were calculated for each 100 kb non-overlapping

sliding window using VCFtools[54,55]. Pair-wise non-synonymous-to-synonymous ($d_N/d_S$) ratios of the ten

species were inferred by yn00 program in the Phylogenetic Analysis by Maximum Likelihood (PAML)

package[56]. Inter-lineage $d_N/d_S$ value for each gene was derived by averaging $d_N/d_S$ values obtained from

all pairwise comparisons of samples belonging to the two lineages under investigation. Candidate genes

with the 5% highest and 5% lowest $d_N/d_S$ values were considered to have undergone strong positive and

purifying selection, respectively.

**Cytosine methylation pattern and epigenetic population structure**

Whole genome bisulfite sequencing data were pre-processed using TrimGalore (https://www.

bioinformatics.babraham.ac.uk/projects/trim_galore/, accessed August 21, 2018). Paired-end reads

were then aligned to the reference genome using Bismark[57] with a moderately stringent minimum-score

function (L,0,-0.3). De-duplicated alignments of the 36 *Aquilegia* accessions were used to report cytosine

methylation level using "Bismark_Methylation_Extractor", on loci with a read depth ≥ 3. Genomic

annotations of the methylated cytosine site were identified based on the reference genome using an in-

house Python script. PCA was conducted for 588,659 loci which were passed the quality control to infer

the CG-methylomic diversity of the ten *Aquilegia* species. Differential cytosine methylation was

determined at the gene and chromosome levels, respectively. At the gene level, we determined DMRs

for each 100 bp non-overlapping sliding window using Cochran-Mantel-Haenszel (CMH) test to account

for imbalanced read depth (Supplementary Notes). Genomic regions that possessed a Benjamin-

Hochberg adjusted *p* value < 0.05 and showed inter-specific or inter-lineage methylation divergence

higher than 25% were defined as significant DMRs. Genes with > 20% of the genic region being DMR(s)

were defined as DMGs. Chromosome level methylation patterns were measured by chromosomal

methylation discrepancy index (MDI)[58]. Methylation patterns of the identified DMGs were visually

confirmed on Integrative Genomics Viewer[59] prior to downstream analyses and biological interpretation.

In addition, we identified CG islands from the *A. coerulea* "Goldsmith" reference genome using EMBOSS

cpgplot with default settings[60]. Only the identified CG-enriched genomic regions with > 200 bp were

defined as CG islands. We then investigated inter-specific and inter-lineage methylation patterns in and

around the CG islands.

**Associations between the genetic variation and cytosine methylation**

We tested for associations between the identified DMGs and genes under positive selection by a Chi-

square test. Linear regression model was adopted to measure the direct causal effect of CG-loss

variation on CG methylation. To further assess whether genetic variations drive the establishment of

DMG, driving mutations of DMRs between the *A. japonica* and *A. oxysepala* were identified using an

Eigenstrat-based method (See Supplementary Notes for more details)[61].

**Identification of conservative clade-specific variant**

CCVs were defined as variants that had a SnpEff-predicted "high" functional impact and that were

conserved across all samples belonging to the same species or lineage, but not present in any sample of

the other species/lineages. Since the biological consequences of heterozygous variants were less

affirmable, only the homozygous point mutations and INDELs were included in the characterization of

CCVs, including frameshift, stop-gain, stop-loss, start-loss and splicing-alteration variations.

**Functional analysis**

The above mentioned genetic and epigenetic analyses helped to identify relevant candidate genes,

which might be associated with the rapid diversification of the *Aquilegia* species from different

perspectives. These candidate genes were employed to conduct functional enrichment analyses using

the R package topGO with default settings[62]. Enriched GO terms that possessed a *p* value <0.05 were

considered statistically significant. Since the statistical tests performed by topGO are not independent,

multiple testing correction does not apply here[62]. Structures of functional domains of targeted genes

were determined based on the InterPro database (https://www.ebi.ac.uk/interpro, accessed January 25,

2019). Distribution patterns of the identified candidate genes and their related functional pathways

were visualized using the R package jvenn[63].

**Data availability**

566  All data generated from the study were submitted to EBI under the accession number PRJEB34182. All

567  scripts for conducting computational analyses are available upon reasonable request to the

568  corresponding author.

**Author Contributions**

571  L.F.L., J.Z and A.S.X conceived this project. Z.H.W, T.L. and M.R.L. developed statistical analysis pipeline.

572  Z.H.W., T.L., M.R.L., N.D., L.Z.L., Y.J.H and X.G. carried out experiments and analyzed the data. T.L.,

573  M.R.L., N.D., Z.H.W., L.Z.L., X.G., and L.F.L. participated in discussion and interpreted the data. Z.H.W, T.L.

574  and L.F.L. wrote the manuscript. All authors read and approved the manuscript.

**Acknowledgments**

577  We are very grateful to Aköz Gökçe for her constructive comments on our manuscript. We appreciate Dr.

578  Peng Jiang, Zhang Zhang, and the USDA for kindly providing the seeds of the *Aquilegia* species.

**Funding:**

581  This work was financially supported by National Natural Science Foundation of China (31670382),

582  Shanghai Pujiang Program (19PJ1401500), Start-up funding at Fudan University (JIH1322105) and the

583  Department of Science and Technology of Jilin Province (20190201299JC). The funders had no role in

584  study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests**

587  The authors have declared that no competing interests exist.

**Supplementary files**

**Figure S1**. Per-chromosome phylogenetic trees reconstructed using neighbor-joining algorithm. Polymorphisms detected on each chromosome were retrieved separately to infer the phylogeny.

**Figure S2**. Distribution of nucleotide diversity (π) at the whole-genome level and the per-chromosome level. Nucleotide diversity was estimated for each lineage pooling corresponding species, as well as for *A. japonica* and *A. oxysepala*.

**Figure S3**. Spearman correlation of the genome-wide nucleotide variation pattern for each 100-kb sliding window between European and Asian (a), North American and Asian (b), European and North American (d), *Aquilegia japonica* and *A. oxysepala* (d). Each dot represents a 100-kb sliding window. Values on the x- and y-axis are the nucleotide diversity (π) for each sliding window.

**Figure S4**. Overlapping of low diversity genomic region (LDGR) between the three lineages. 148 LDGRs with 5% lowest nucleotide diversity were defined as LDNRs in each lineage, totaling 241 unique regions.

**Figure S5**. Pair-wise $d_N/d_S$ ratio for all genes between the species within and between the Asian, European and North American lineages.

**Figure S6**. PCA illustrates three distinct clusters corresponding to the three lineages. Asian species further demonstrated higher inter-specific divergence than the American and the European species. PCA was performed based on 588,659 loci with sufficiently high sequencing quality.

**Figure S7**. Illustration of differential methylation in two photosynthesis genes. CG methylation pattern of two genes, *Aqcoe7G230600* photosystem I *PsaA/PsaB* (a) and *Aqcoe7G231300 CemA* (b) in *A. japonica* and *A. oxysepala* throughout the gene body region. Red bars indicate methylation level (0-100) at CG loci. Genomic coordinates on the chromosome 7 are annotated.

**Figure S8**. Overlapping of the CCVs (a) and DMGs (b) identified in inter-lineage/species comparisons. A considerable proportion of these CCVs (84.1%) and DMGs (51.3%) were shared by two or more inter-lineage/species comparisons.

**Figure S9**. Venn analyses of the candidate genes carrying CCVs and DMGs. Each subpanel indicates the

comparison between the *A. japonica* and *A. oxysepala* (a), *A. japonica* and North American (b), *A.*

*japonica* and European (c), *A. oxysepala* and North American (d), *A. oxysepala* and European (e), North

American and European (f).

**Table S1**. Candidate genomic regions that showed high genetic divergence (top 5% highest $F_{ST}$) between

*Aquilegia japonica* and *A. oxysepala* and low nucleotide diversity (top 5% lowest $\pi$) within each species.

**Table S2**. Summary of the highly impactful clade specific variations (CCVs) at both the species and

lineage levels.

**Table S3**. Statistics for differentially methylated regions (DMRs) among the four *Aquilegia* lineages or

species. Odds ratio estimates the relative methylation level between two lineages or species being

compared in corresponding region. DMRs were sorted by genomic coordinates with hypo-methylated

DMRs in the first lineage/species preceding hyper-methylated DMRs.

**Table S4**. Statistics for differentially methylated genes (DMGs) among the four *Aquilegia* lineages/

species. Only genes harboring a high density of differentially methylated regions (> 2 per kb) were

considered DMGs.

**Table S5**. Information of the 36 *Aquilegia* samples used in this study.

**References**

1. Givnish, T. J. Adaptive radiation versus 'radiation' and 'explosive diversification': Why conceptual

distinctions are fundamental to understanding evolution. *New Phytologist* **207,** 297–303 (2015).

2. Schluter, D. The Ecology of Adaptive Radiation. *Heredity* **86,** 749–750 (2001).

3. Schluter, D. Ecological causes of adaptive radiation. *American Naturalist* **148,** (1996).

4. Givnish TJ. Adaptive radiation and molecular systematics: aims and conceptual issues. Molecular

Evolution and Adaptive Radiation. 1997:1-54.

5. Simpson GG, Olson EC. The major features of evolution. Quarterly Review of Biology.1953;8(1):87-8.

6. Losos, J. B. Adaptive Radiation, Ecological Opportunity, and Evolutionary Determinism. *American*

*Naturalists.* **175,** 623–639 (2010).

7. Irisarri, I. *et al.* Phylogenomics uncovers early hybridization and adaptive loci shaping the radiation of

Lake Tanganyika cichlid fishes. *Nature Communications* **9,** (2018).

8. Lamichhaney, S. *et al.* Evolution of Darwin's finches and their beaks revealed by genome sequencing.

*Nature* **518,** 371–375 (2015).

9. Losos, J. B. & Ricklefs, R. E. Adaptation and diversification on islands. *Nature* **457,** 830–836 (2009).

10. Motley, T. J. Tarweeds & Silverswords, Evolution of the Madiinae (Asteraceae). *Economic Botany* **58,**

123–124 (2004).

11. Wagner, C. E., Harmon, L. J. & Seehausen, O. Ecological opportunity and sexual selection together

predict adaptive radiation. *Nature* **487,** 366–369 (2012).

12. Simpson, E. H. Measurement of diversity. *Nature* **163,** 688 (1949).

13. Berner, D. & Salzburger, W. The genomics of organismal diversification illuminated by adaptive

radiations. *Trends in Genetics* **31,** 491–499 (2015).

14. Brawand, D. *et al.* The genomic substrate for adaptive radiation in African cichlid fish. *Nature* **513,**

375–381 (2015).

15.   Ford, A. G. P. *et al.* Niche divergence facilitated by fine-scale ecological partitioning in a recent

cichlid fish adaptive radiation. *Evolution* **70,** 2718–2735 (2016).

16.  Filiault, D. L. *et al.* The Aquilegia genome provides insight into adaptive radiation and reveals an

extraordinarily polymorphic chromosome with a unique history. *Elife* **7,** (2018).

17.   Fior, S. *et al.* Spatiotemporal reconstruction of the Aquilegia rapid radiation through next-

generation sequencing of rapidly evolving cpDNA regions. *New Phytologist* **198,** 579–592 (2013).

18.  Aquilegia:, M. the cultivated and wild columbines. *Gentes Herbarum* **7,** 1–150 (1946).

19.   Bastida, J. M., Alcántara, J. M., Rey, P. J., Vargas, P. & Herrera, C. M. Extended phylogeny of

Aquilegia: The biogeographical and ecological patterns of two simultaneous but contrasting radiations.

*Plant Systematics and Evolution* **284,** 171–185 (2010).

20.  Sharma, B., Yant, L., Hodges, S. A. & Kramer, E. M. Understanding the development and evolution of

novel floral form in Aquilegia. *Current Opinion in Plant Biology* **17,** 22–27 (2014).

21.  Kramer, E. M. & Hodges, S. A. Aquilegia as a model system for the evolution and ecology of petals.

*Philosophical Transactions of the Royal Society B: Biological Sciences* **365,** 477–490 (2010).

22.   Alcántara, J. M., Bastida, J. M. & Rey, P. J. Linking divergent selection on vegetative traits to

environmental variation and phenotypic diversification in the Iberian columbines (Aquilegia). *Journal of*

*Evolutionary Biology* **23,** 1218–1233 (2010).

23.  Whittall, J. B. & Hodges, S. A. Pollinator shifts drive increasingly long nectar spurs in columbine

flowers. *Nature* **447,** 706–709 (2007).

24.   Garrido, J. L. *et al.* Geographic genetic structure of Iberian columbines (gen. Aquilegia). *Plant*

*Systematics and Evolution* **303,** 1145–1160 (2017).

25.  Li, L. F. *et al.* Phenotypic and genetic evidence for ecological speciation of aquilegia japonica and A.

oxysepala. *New Phytologist.* **204,** 1028–1040 (2014).

26.  Tang, L. L., Yu, Q., Sun, J. F. & Huang, S. Q. Floral traits and isolation of three sympatric Aquilegia

species in the Qinling Mountains, China. *Plant Systematics Evolution* **267,** 121–128 (2007).

27.  Zhang, H. & Dawe, R. K. Mechanisms of plant spindle formation. *Chromosome Research* **19,** 335–344

(2011).

28.  Aguirre-Portolés, C. *et al.* Tpx2 controls spindle integrity, genome stability, and tumor development.

*Cancer Research* **72,** 1518–1528 (2012).

29.  Fukui, K. *et al.* Crystal structure of Muts2 endonuclease domain and the mechanism of homologous

recombination suppression. *Journal of Biological Chemistry* **283,** 33417–33427 (2008).

30.  Li, M. R. *et al.* Rapid Divergence Followed by Adaptation to Contrasting Ecological Niches of Two

Closely Related Columbine Species Aquilegia japonica and A. oxysepala. *Genome Biology Evolution* **11,**

919–930 (2019).

31.  Machado-Schiaffino, G., Henning, F. & Meyer, A. Species-specific differences in adaptive phenotypic

plasticity in an ecologically relevant trophic trait: Hypertrophic lips in midas cichlid fishes. *Evolution* **68,**

2086–2091 (2014).

32.  Verhoeven, K. J. F., VonHoldt, B. M. & Sork, V. L. Epigenetics in ecology and evolution: What we

know and what we need to know. *Molecular Ecology* **25,** 1631–1638 (2016).

33.  Diez, C. M., Roessler, K. & Gaut, B. S. Epigenetics and plant genome evolution. *Current Opinion in*

*Plant Biology* **18,** 1–8 (2014).

34.  Bossdorf, O., Richards, C. L. & Pigliucci, M. Epigenetics for ecologists. *Ecology Letters* **11,** 106–115

(2008).

35.  Richards, C. L. *et al.* Ecological plant epigenetics: Evidence from model and non-model species, and

the way forward. *Ecology Letters* **20,** 1576–1590 (2017).

36.  Richards, C. L., Bossdorf, O. & Verhoeven, K. J. F. Understanding natural epigenetic variation. *New*

*Phytologist* **187,** 562–564 (2010).

37.  Cortijo, S. *et al.* Mapping the epigenetic basis of complex traits. *Science* **343,** 1145–1148 (2014).

38.  Schmitz, R. J. *et al.* Transgenerational epigenetic instability is a source of novel methylation variants.

*Science* **334,** 369–373 (2011).

39.  Becker, C. *et al.* Spontaneous epigenetic variation in the Arabidopsis thaliana methylome. *Nature*

**480,** 245–249 (2011).

40.  Li, L., Wang, C., Lu, T., Lin, S. & Hu, Y.-Q. Indirect effect inference and application to GAW20 data.

*BMC Genetics* **19,** (2018).

41.  Fisher, V. A. *et al.* Do changes in DNA methylation mediate or interact with SNP variation? A

pharmacoepigenetic analysis. *BMC Genetics* **19,** (2018).

42.  Zilberman, D. An evolutionary case for functional gene body methylation in plants and animals.

*Genome Biology* **18,** (2017).

43.  Bewick, A. J. *et al.* On the origin and evolutionary consequences of gene body DNA methylation.

*Proceedings of the National Academy of  Sciences of the United States of America* **113,** 9111–9116

(2016).

44.   Takuno, S. & Gaut, B. S. Gene body methylation is conserved between plant orthologs and is of
evolutionary consequence. *Proceedings of the National Academy of  Sciences of the United States of*
*America* **110,** 1797–1802 (2013).

45.   Bewick, A. J. & Schmitz, R. J. Gene body DNA methylation in plants. *Current Opinion in Plant Biology*
**36,** 103–110 (2017).

46.   Li, L. F., Pang, D., Liao, Q. L. & Xiao, H. X. Genomic and EST microsatellite markers for Aquilegia
flabellata and cross-amplification in A. oxysepala (Ranunculaceae). *American Journal of Botany* **98,**
(2011).

47.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
*Bioinformatics* **25,** 1754–1760 (2009).

48.   McKenna, A. *et al.* The genome analysis toolkit: A MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Research* **20,** 1297–1303 (2010).

49.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–2079
(2009).

50.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff. *Fly (Austin).* **6,** 80–92 (2012).

51.   Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0
for Bigger Datasets. *Molecular Biology and Evolution* **33,** 1870–1874 (2016).

52.   Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal component
analysis of SNP data. *Bioinformatics* **28,** 3326–3328 (2012).

53.   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated
individuals. *Genome Research* **19,** 1655–1664 (2009).

54.   Weir, B. S. & Cockerham, C. C. Estimating F-statistics for the analysis of population structure.
*Evolution* **38,** 1358–1370 (1984).

55.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27,** 2156–2158 (2011).

56.   Yang, Z. PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution* **24,**
1586–1591 (2007).

57.   Krueger, F. & Andrews, S. R. Bismark: A flexible aligner and methylation caller for Bisulfite-Seq
applications. *Bioinformatics* **27,** 1571–1572 (2011).

58.   O'Sullivan, D. E., Johnson, K. C., Skinner, L., Koestler, D. C. & Christensen, B. C. Epigenetic and
genetic burden measures are associated with tumor characteristics in invasive breast carcinoma.
*Epigenetics* **11,** 344–353 (2016).

59.   Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV): High-

performance genomics data visualization and exploration. *Brief Bioinformatics* **14,** 178–192 (2013).

60.   Rice, P., Longden, L. & Bleasby, A. EMBOSS: The European Molecular Biology Open Software Suite.

*Trends in Genetics* **16,** 276–277 (2000).

61.   Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-wide association

studies. *Nature Genetics* **38,** 904–909 (2006).

62.   Alexa, A., Rahnenführer, J. & Lengauer, T. Improved scoring of functional groups from gene

expression data by decorrelating GO graph structure. *Bioinformatics* **22,** 1600–1607 (2006).

63.   Bardou, P., Mariette, J., Escudié, F., Djemiel, C. & Klopp, C. Jvenn: An interactive Venn diagram

viewer. *BMC Bioinformatics* **15,** (2014).

European  Asian

49  7  37

5  17

American

*Size of each list*

148  148  148

0
European  Asian  American

*Number of elements: specific (1) or shared by 2, 3, ... lists*

87  29  125

3  2  1

(a)

*A. japonica*

*A. oxysepala*

20392800                    Aqcoe7G230600                    20394100

(b)

*A. japonica*

*A. oxysepala*

20403000                    Aqcoe7G231300                    20406500

(a)

Jap vs Ame    Jap vs Eur    Ame vs Eur

95    102    52    27    69

49    17

57    37    23

2    17

163    1    Oxy vs Ame

91    29    103

37    38    0

Oxy vs Eur    Jap vs Oxy

*Size of each list*

1416

1172    1091    736    1389    1154    1416

Jap vs Ame    Jap vs Eur    Ame vs Eur    Oxy vs Ame    Jap vs Oxy    Oxy vs Eur

*Number of elements: specific (1) or shared by 2, 3, ... lists*

1152    794    430

6 (17)    3    2    1
5 (134)
4 (178)

(b)

Jap vs Ame    Jap vs Eur    Ame vs Eur

133    37    44    38

0    406

41    79

69    2

26    4

3    14    Jap vs Oxy

15    14    40

83    14    77

72    200

Oxy vs Eur    Oxy vs Ame

*Size of each list*

465.5

578    417    931    295    682    531

Jap vs Ame    Jap vs Eur    Ame vs Eur    Jap vs Oxy    Oxy vs Ame    Oxy vs Eur

*Number of elements: specific (1) or shared by 2, 3, ... lists*

317    507    906

6 (4)    3    2    1
5 (31)
4 (96)

(a) *A. japonica* and *A. oxysepala*

CCV       DMG
1134    20    275

(b) *A. japonica* and North Amrican

CCV       DMG
1128    44    534

(c) *A. japonica* and European

CCV       DMG
1072    19    398

(d) *A. oxysepala* and North American

CCV       DMG
1339    50    632

(e) *A. oxysepala* and European

CCV       DMG
1380    36    495

(f) North American and European

CCV       DMG
895    36    700