Genetic and epigenetic signatures associated with

the rapid radiation of *Aquilegia* species

Tianyuan Lu[1,2,3], Ming-Rui Li[1], Ning Ding[1], Zhen-Hui Wang[4], Li-Zhen Lan[1], Xiang Gao[5], and Lin-Feng Li[1,*]

[1]Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, School of Life

Sciences, Fudan University, Shanghai 200438, China;

[2]McGill University and Genome Quebec Innovation Center, Montreal H3A 0G1, Quebec, Canada;

[3]Lady Davis Institute, SMBD JGH, Montreal H3A 1A3, Quebec, Canada;

[4]Faculty of Agronomy, Jilin Agricultural University, Changchun 130118, China;

[5]Key Laboratory of Molecular Epigenetics of the Ministry of Education, Northeast Normal University,

Changchun 130024, China.

*Correspondence author:

Lin-Feng Li (lilinfeng@fudan.edu.cn**)**

Abstract

Elucidating how the diverse evolutionary mechanisms interact to determine species diversification is

crucial to understanding the evolution and persistence of biodiversity. The genus *Aquilegia* (columbine)

is a model system to identify genetic and epigenetic signatures underpinning the rapid adaptive

radiation. In this study, we surveyed the genomes and DNA methylomes of ten worldwide *Aquilegia*

species to investigate whether specific genomic architectures were associated with rapid species

diversification in the Asian, European and North American lineages. The resulting phylogenies and

population structure inferences revealed clearly high genetic and DNA methylomic divergence among

the three lineages. Genome-wide scanning demonstrated significantly higher positive and purifying

selection pressures among the Asian species compared to the European and North American lineages.

More importantly, candidate genes identified at the genetic and epigenetic levels are functionally

related to diverse adaptation-related traits, such as stress tolerance, photosynthesis and cell

reproduction. While a considerable proportion of the candidate genes share hotspots of intra- and inter-

lineage divergence, genetic and epigenetic factors were found to act on complementary biological

pathways. By assessing the interaction between genetic variations and epigenetic variability, we found

that epigenetic variability is a partially independent factor that intertwines with genetic mechanism to

facilitate the diversification of the *Aquilegia* species. Taken together, these findings suggest that specific

genetic architectures have been involved in vital biological pathways in response to diverse

environmental conditions and epigenetic modifications may play a complementary role in the process of

adaptive speciation. Our study provides an exploratory overview of how the established genetic and

epigenetic signatures are associated with the rapid species diversification of *Aquilegia* species.

Key words: Adaptive radiation; *Aquilegia*; Methylome; Speciation; Species diversification

Author Summary

Disentangling the genetic and epigenetic bases underpinning species diversification is crucial to understanding the evolution and persistence of biodiversity. The columbine (genus *Aquilegia*) is a model system to address biological mechanisms associated with rapid adaptive radiation. We surveyed the genomes and DNA methylomes of ten worldwide columbine species to investigate whether specific genetic and epigenetic architectures were involved in the diversification of Asian, European and North American columbine species. We pinpointed candidate genes featuring rapid adaptive radiation identified at either the genetic or the epigenetic levels. These candidate genes are functionally associated with diverse adaptation-related traits such as cell reproduction, plant growth, and stress tolerance. Such genetic and epigenetic signatures have potential contributed adaptabilities to the columbine species to cope with diverse environmental conditions. In addition, we also showed that epigenetic modifications could act as a complementary factor that intertwined with genetic mechanism to facilitate the diversification of the columbine species. In all, our study provides a genome-wide view of how the genetic and epigenetic factors are associated with the rapid species diversification of the columbine species.

Introduction

Adaptive radiation is the rapid diversification of a single ancestral species into a vast array of common descendants that inhabit different ecological niches or use a variety of resources, but differ in phenotypic traits required to exploit diverse environments [1-4]. Disentangling the evolutionary mechanisms underpinning adaptive radiation is fundamental to understanding the evolution and persistence of biodiversity (5, 6). This has been a key focus of many studies investigating different animal and plant lineages that diversified through adaptive radiation, including Hawaiian silversword, Caribbean anoles, Darwin's finches, and African cichlids (7-10). However, it remains under-investigated as to why some lineages could diversify rapidly but their close relatives or other sympatrically distributed lineages did not. In the past decades, accumulating evidence from diverse radiation lineages suggests that both the extrinsic environmental variables (e.g., resource availability) and intrinsic factors (e.g., genetic variants determine lineage-specific morphological variations) are the potential mechanisms in determining the rate and volume of species diversification [11]. Among the environmental triggers, ecological opportunity is considered as the primary mechanism that causes rapid adaptive radiation through the acquisition of key innovations, invasion of new environments and extinction of competitors [2, 12]. On the other hand, new species can also arise as a result of multiple evolutionary mechanisms (e.g. natural selection) acting on genetic factors (e.g., advantage alleles) that ultimately influence the phenotypic disparity among closely related species [13]. An illustration of rapid adaptive radiation is the African cichlid fishes, on which the extrinsic environmental factors (e.g., ecological specialization) and intrinsic mechanisms (e.g., adaptive introgression) acted together to provoke the repeated adaptive radiation in geographically isolated lakes [10, 11, 14, 15].

The genus *Aquilegia* L. (columbine) is a well-recognized model system to address the evolutionary mechanisms underlying rapid adaptive radiation [16, 17]. This genus includes approximately 70 recently diversified species that are widely distributed in the temperate areas of North America and Eurasia [18]. Phylogenetic and geographic inferences have illustrated two independent adaptive radiations of North American and European lineages from the distinct ancestral Asian species [16, 19]. For example, floral diversification of the North American *Aquilegia* species is highly correlated with the pollinator specialization [20-23]. In contrast, ecological adaptation and geographic isolation are considered as the major driving forces that promoted rapid radiation of the European species [16, 24]. In Asia, changes in pollinator and ecological habitat are both proposed to be the underpinning mechanisms that resulted in the diversification of more than 20 morphologically distinct species [25, 26]. These Asian *Aquilegia* species constitute four highly divergent lineages corresponding to respective geographic origins and

evolved relatively independent under distinct external conditions [25, 26]. This well-described
evolutionary history renders the *Aquilegia* species an ideal system to address how the extrinsic and
intrinsic factors shaped genome-wide genetic and epigenetic variation patterns.

In this study, we survey the genomes and DNA methylomes of 36 accessions from ten worldwide
Aquilegia species to elucidate the genetic and epigenetic bases underlying rapid diversification of the
Asian, European and North American *Aquilegia* species. Among the Asian species, four phylogenetically
distinct species (*A. japonica*, *A. oxysepala, A. yabeana*, and *A. viridiflora*) are selected according to their
geographic distributions and ecological habitats. *Aquilegia japonica* and *A. oxysepala* are sister species
inhabiting alpine tundra and low altitude forest niches in northeastern China, respectively [26, 27]. Our
previous studies have documented that natural selection during ecological specialization together with
genetic drift under geographic isolation caused the rapid evolution of reproductive isolation between
these two species [26, 28]. Here, we further investigate how diverse evolutionary driving forces shaped
genetic and epigenetic architectures of the two species in the processes of speciation and adaptation. In
addition, we also evaluated patterns of nucleotide variation and cytosine methylation in the *A. yabeana*
and *A. viridiflora*. The former species shares highly similar morphological traits and ecological niches
with the *A. oxysepala* but is allopatrically distributed in the northern China. In contrast, while the *A.*
viridiflora is sympatrically distributed with *A. yabeana* and *A. oxysepala* in the northern and
northeastern China, it often occupies rocky and sandy ecological niches. As a supplementary, we also
examined nucleotide and cytosine methylation variation patterns of the North American and European
lineages to investigate how the distinct evolutionary mechanisms shaped genomic and epigenomic
architectures on a genome-wide scale. Our study will provide a genome-wide view of how the genetic
and epigenetic architectures associated with the phenotypic divergence and ecological specialization of
the *Aquilegia* species.

Results

Population structure and nucleotide variation pattern

Neighbor-joining (NJ) trees were reconstructed for the 36 *Aquilegia* accessions based on 689,123
homozygous SNPs. The established phylogenies showed that accessions of the ten species formed three
distinct lineages corresponding to their geographic origins (**Figure 1a** and **Figure S1**). In brief, all the 22
accessions of the four East Asian species, *A. japonica*, *A. oxysepala*, *A. yabeana* and *A. viridiflora*,
clustered as a monophyletic lineage, with the former two species and their hybrid forming a clade and
the latter two species grouping as a sister clade. In contrast, the West Asian species *A. fragrans*

clustered with geographically adjoining European species. Within the North American lineage,

phylogenetic relationships of the four species varied across the seven chromosomes. In particular, one

European accession of the *A. alpina* var. *alba* fell into the North American lineage at the chromosomes 1

and 3. Likewise, the principal component analysis (PCA) and population structure inferences also

revealed distinct genetic structure of the three phylogenetic lineages (**Figure 1b** and **c**). Consistent with

the above phylogenies, one *A. alpina* var. *alba* accession shared the same ancestral genetic cluster with

the North American lineage and the putative hybrid of the *A. oxysepala* and *A. japonica* possessed a

mixed genomic constitution (**Figure 1b** and **c**).

To further gain genome-wide nucleotide variation pattern of the ten *Aquilegia* species, we

calculated nucleotide diversity (π) and genetic divergence (F_{ST}) for each chromosome and 100-kb sliding

window, respectively. Among the three phylogenetic lineages, the Asian *Aquilegia* species harbored

relatively higher nucleotide diversity compared to the European and North American lineages across the

seven chromosomes (**Figure S2**). By comparing the nucleotide diversity for each 100-kb sliding window,

we observed a moderate correlation of genome-wide variation pattern among the three lineages

(Spearman R = 0.40-0.52) and a high correlation between the *A. oxysepala* and *A. japonica* (Spearman R

= 0.83) (**Figure S3**). In particular, 116 of 260 low genetic diversity genomic regions (LDGRs, the top 5%

lowest π) were shared by at least two of the three lineages (**Figure S4**). Between the *A. oxysepala* and *A.*

japonica, while we defined 148 LDGRs and 148 high divergence genomic regions (HDGRs, 5% highest F_{ST}),

only seven candidate genomic regions showed overlap between the HDGRs and LDGRs (**Table S1**).

Identification of the genes indicating selection pressure and highly impactful genetic variations

Candidate genes associated with the adaptive divergence were determined according to the following

three strategies. First, we considered the genes localized within the HDGRs and LDGRs more prone to be

exposed to selection in the adaptive evolution process. Under this hypothesis, we identified 23 genes

from the above seven candidate genomic regions between the *A. oxysepala* and *A. japonica* (**Table S1**).

Genes within these genomic regions were functionally associated with meiotic nuclear division, adenine

methyltransferase and basic cellar activities.

The first strategy mainly relied on genome-wide scanning for 100-kb non-overlapping sliding

window. We next employed a functional annotation-based approach to identify highly impactful

conservative clade-specific variations (CCVs; e.g., frameshift mutations, stop mutation and splicing

alteration) from both the within and between lineage comparisons. Our results revealed that a

considerable proportion (17.9-40.5%) of the CCVs were identified in the gene body regions (**Table S2**).

We then examined the potential impacts of these identified CCVs on gene functions. Between the *A.*

oxysepala and *A. japonica*, the CCV-carrying genes were enriched in several vital biological pathways

related to cell reproduction, including telomere maintenance, DNA repair, and DNA helicase activity

(**Figure 2** and **Table 1**). For example, two candidate genes (*Aqcoe6G160300* and *Aqcoe7G062500*) coding

for *Xklp2* (*TPX2*) were functionally correlated with spindle assembly during the mitotic process [29, 30].

Among the three phylogenetic lineages, the CCVs-harboring genes were also functionally involved in the

mitotic chromosome condensation, DNA ligase activity and aminopeptidase activity (**Figure 2** and **Table**

1). For instance, two CCV-containing genes (*Aqcoe2G276600* and *Aqcoe1G273400*) encoding DNA

mismatch repair proteins *MutS/MSH* and *MutS2* [31] carried one Asian-specific-to-American frameshift

variant.

Thirdly, we also performed genome-wide scanning to identify genes informative of selection

pressure from the within and between lineage comparisons. By estimating the pair-wise synonymous (d_S)

and non-synonymous (d_N) mutation rate, we found that species within the Asian lineage experienced

significantly higher positive ($d_N/d_S > 1$) and purifying ($d_N/d_S < 1$) selection pressures compared to the

European and North American lineages (Wilcoxon rank sum test, all Bonferroni-corrected *p* values <

1.5×10^{-16}) (**Figure S5**). Likewise, the European species showed significant purifying selection (Wilcoxon

rank sum test, Bonferroni-corrected *p* value = 7.8×10^{-8}) compared to the North American species.

Among the three lineages, all the comparisons showed significant purifying and positive selection

pressures (Wilcoxon rank sum test, all Bonferroni-corrected *p* values < 0.0018) except those that

indicated positive selection between the North American and the other two lineages (Wilcoxon rank

sum test, *p* value = 1).

CG methylation patterns and differentially methylated genes

In parallel with the above genomic analyses, we also investigated CG methylation pattern of the

representative *Aquilegia* species. Despite variance across the 36 *Aquilegia* accessions, the North

American, Asian and European species showed no distinguishable differences (t test, all Bonferroni-

corrected *p* values > 0.01) in overall percentage of methylated cytosines (**Figure 3a**). We then performed

PCA to examine CG-cytosine methylomic relatedness of all the *Aquilegia* accessions. The resulting

overall methylation pattern highly resembled the above genomic inferences, with the European and

American species forming two independent groups and the four Asian species forming three separated

clusters (**Figure S6**). We then assessed the CG methylation patterns for the European and North

American lineages as well as the three Asian species (*A. japonica*, *A. oxysepala* and *A. viridiflora*)

separately. Consistent with the described genomic features, heterogeneous pattern of the CG

methylation was also observed among the seven chromosomes, with the chromosome 4 demonstrating

obviously higher overall CG methylation divergence compared to the other six chromosomes (**Figure 3b**).

We further quantified CG methylation deposited in the genic regions, putative *cis*-regulatory regions and

CG island, respectively. In genic and regulatory regions, all three lineages shared similar modification

patterns with apparent depletion of CG methylation around the transcription start site (TSS) and

transcription end site (TES) (**Figure 3c**). However, the American lineage exhibited hyper-methylation

(more than 10%) around the center of CG islands and a more drastic decrease throughout the CG island

shores compared to the European and Asian species (**Figure 3d**).

To examine the biological impacts of CG methylation on the species diversification, differentially

methylated regions (DMRs) and differentially methylated genes (DMGs) were identified from both the

within and between thee three lineages, respectively (**Tables S3** and **S4**). Within the Asian lineage, 3,622

DMRs in 2,899 DMGs were identified between the *A. japonica* and *A. oxysepala*. Functional enrichment

of these DMGs indicated that the two species may have different activities in photosynthesis-related

pathways, including photosystem I, photosynthesis and chloroplast (**Figure 2**). For example, two

photosynthesis-related genes, *PsaA/PsaB* and *CemA*, showed significantly differential methylation

between the two species in the genic regions (**Figure S7a** and **b**). At the inter-lineage level, apparently

more DMGs were identified between the North American and European species (6,087 genes)

compared to those of between the two lineages and Asian species (3,308-5,003 genes) (**Table S3** and **S4**).

DMGs characterized from the inter-lineage comparisons were mainly involved in the plant growth (e.g.,

response to auxin) and defense, response to biotic stimulus and wounding (**Figure 2**).

We then examined whether the candidate genes (CCV-carrying genes and DMGs) superimposed on

the same signature of natural selection. We found while a considerable proportion of the candidate

genes were shared for each of the genetic and epigenetic assessments (**Figure S8**), they showed a

segregated distribution pattern across all comparisons (**Figure S9**). Likewise, the Gene Ontology (GO)

enrichment analyses of the candidate gene identified from the genetic and epigenetic levels were

enriched in functionally complementary pathways (**Figure 2**), suggesting co-existence of different

underlying evolutionary mechanisms.

Association between epigenetic variability and genetic variations

Since both genetic variation and differential CG methylation seemed to have crucial and multifaceted

influences on the adaptation of the ten *Aquilegia* species, we wondered whether differential epigenetic

modifications were dependent on genetic variability. Among the 588,659 CG sites examined, 224,222

(38.09%) carried a CG-loss variation. We then illustrated epigenetic variability for the variation-carrying

and non-variant CG sites, respectively. As shown in Figure 4, genetic-epigenetic associations of varying

magnitude were observed in both types of CG sites. The variation-carrying CG sites conveyed

information that highly resembled their genetic background. The overall methylation pattern was highly

conserved within the same species but exhibited obvious divergence across the ten *Aquilegia* species

(**Figure 4a**). In contrast, CG methylation divergence at the non-variant CG sites varied with higher

variability at both the intra- and inter-specific levels (**Figure 4b**). By examining the correlation of genetic

variability and cytosine methylation, we found that CG methylation divergence at variation-carrying CG-

site was largely attributable to the CG-loss variations (**Figure 4c**). In particular, 75% of the CG-loss

variations occurring at the most highly variable CG-methylated dinucleotides could explain at least 75%

of the total epigenetic variability *per se*. Nevertheless, there was still a considerable proportion of

epigenetic variability that could not be sufficiently explained by the variant-CG site (**Figure 4d**).

We also attempted to identify *cis*-driving mutations for each of the 1,229 DMRs between the *A.*

japonica and *A. oxysepala*. Our results revealed that only 568 out of the 1,229 (46.2%) DMRs were

significantly associated with at least one genetic variation inside or around a 500 base-pair (bp)

upstream/downstream genomic region, even under the least stringent *p* value threshold (5×10^{-5}),

indicating that the epigenetic changes were only partially dependent on *cis*-genetic driving mutations

(**Figure 4e**). Moreover, we observed significant, though weak, associations between differential CG

methylation and selection pressure. In most inter-lineage comparisons, DMGs were significantly more

prone to be under positive selection than non-DMGs (**Table 2**), suggesting that epigenetic modifications

could probably assist selection pressure in shaping genotypes. In contrast, DMGs were significantly less

prone to be under purifying selection (**Table 2**).

Discussion

Genetic-driven mechanisms associated with the rapid diversification of *Aquilegia* species

Elucidating evolutionary mechanisms underpinning species diversification is crucial to understanding the

evolution and persistence of biodiversity [2, 5, 6]. The genus *Aquilegia* provides an ideal system to

address how the diverse evolutionary mechanisms promoted rapid adaptive radiation [16, 17]. Although

various environmental conditions related to ecological opportunities, such as shifts in pollinator and

habitat, have been proposed to facilitate the evolution of reproductive isolation [22, 26], genetic basis

associated with the rapid diversification of *Aquilegia* species has still remained largely unclear. In this

study, we surveyed the genomes of ten worldwide *Aquilegia* species to address whether specific genetic

architectures have been involved in the rapid species diversification. Broadly consistent with previously

inferred phylogenies [16, 19, 26, 28], the ten *Aquilegia* species from Asia, Europe and North America

formed three phylogenetically independent lineages corresponding to their geographic origins. This

attribute offers the ten *Aquilegia* species a suitable system to identify genomic determinants associated

with the repeated adaptive speciation by extensive comparisons from different facets.

It has been proposed that if a genetic factor is the potential determinant promoting adaptive

speciation, one would expect to identify specific genetic architectures in the diversified lineages [9, 14,

32]. In Darwin's finches, for example, polyphyletic topology was observed as a general pattern in 14

morphologically distinct species, phenotypic diversity of the beak shape was mainly determined by

natural selection acting on the *ALX1* gene during the ecological specialization process [9]. A similar

phenomenon was observed in the East African cichlid fish where the radiating lineages are more

dynamic in terms of gene content and transcriptomic landscape compared to their non-radiating

relatives [14, 32]. In this study, while the Asian, European and North American species have clearly

diverged and evolved allopatrically in respective geographic regions, similar genome-wide nucleotide

variation pattern was observed among the three lineages, suggesting that a considerable of the genomic

variants might have evolved naturally during the diversification process. Nevertheless, while both the

North American and European lineages were descendants of the ancestral Asian species, shifts in

pollinator and ecological habitat are supposed to drive the independent adaptive radiations,

respectively [16, 17, 19]. This attribute suggest that specific genetic determinants are potentially related

to the repeated adaptive speciation of the *Aquilegia* species. As expected, our genome-wide scanning of

selection not only revealed higher positive and purifying selection pressures among the Asian species,

but also demonstrated significant differences in selection pressures in all inter-lineage comparisons

excepting these of experienced positive selection in North American and European lineages. More

importantly, the CCV-carrying genes identified from the three lineages are associated with cell

reproduction (e.g., telomere maintenance and mitotic chromosome condensation) and some other

functionally important pathways, indicating distinct selection pressures in specific lineages of these loss

function genes.

Similar genomic feature was also observed in the two ecological species *A. japonica* and *A.*

oxysepala. Our previous studies have demonstrated that natural selection and genetic drift together

resulted in the rapid evolution of reproductive isolation [26, 28]. Here, we further demonstrated that

candidate genes involved in the adaptive speciation are functionally enriched in the pathways related to

cell reproduction (e.g., telomere maintenance), stress tolerance (e.g., response to wounding) and basic

cellular activities. These findings from intra- and inter-lineage comparisons suggest that specific genetic

architectures might have played important roles in the repeated adaptive speciation of the *Aquilegia*

species. It should be noted that although a majority of the enriched pathways are specific to each

comparison, enrichment of cell reproduction-related pathways (e.g., telomere maintenance, DNA repair

and DNA helicase activity) is consistent across the intra- and inter-lineage comparisons. Taken together,

our study indicates that both the common and specific genetic determinants might have conferred high

adaptability to the *Aquilegia* species to cope with diverse environmental conditions.

Evolutionary potential of cytosine methylation in the adaptation of *Aquilegia* species

The role of epigenetic modification in the long-term evolutionary process has long been debated [33-35].

It has been proposed that epigenetic variations are frequently under the genetic control but which can

also alter rapidly as a result of environmental induction and stochastic epimutation [36, 37].

Nevertheless, it has also been recognized that some epigenetic variants can persist over generations and

be highly correlated with phenotypic diversity [35]. As illustrated in Arabidopsis, changes in cytosine

methylation can produce meiotically stable epialleles, which could eventually lead to phenotypic

diversity in the absence of genetic variations [38-40]. Here, we assessed whether the epigenetic

modifications are also associated with the adaptive speciation of the *Aquilegia* species. Consistent with

the genomic features detailed above, high divergence of cytosine methylation was observed across the

Asian, European and North American lineages. Nevertheless, differential cytosine methylation is not

only found across the seven chromosomes but also evident in the gene body of DMGs and CG island

region among the three lineages. Particularly, functional enrichment analyses identified significant

associations with adaptation-related traits, including plant growth, stress tolerance and basic cellular

activities. Between the *A. japonica* and *A. oxysepala*, for example, the identified DMGs showed

significant enrichment in categories related to diverse important phenotypic traits, such as

photosynthesis, embryo development and response to auxin. These epigenomic features indicate that

epigenetic factors may also play a role in response to diverse environmental conditions.

We noted that some candidate genes and enriched pathways shared hotspots of both genetic and

epigenetic disparities, especially those related to cell reproduction, plant growth and stress tolerance.

Many studies based on human and mouse have shown genetic variations can not only directly control

CG methylation patterns [41, 42], but also manipulate long-range CG methylation at specific sites to

further influence phenotypes, where CG methylation serves as a mediator [43, 44]. However, precedent

studies on explaining, in a time-span of evolutionary history, whether CG methylation functions

independent of genetic variations are few to our knowledge. To this end, we asked whether the

epigenetic mechanism works independently without certain genetic control. By analyzing the

associations between genetic and epigenetic variability, we conclude that while many CG-loss variations

can directly lead to depletion of CG methylation, a lot of DMRs are not manipulated by any *cis*-variations.

Since gene body CG methylation in plants generally stabilizes gene expression and is positively

correlated with gene expression [45-48], differential methylation in our study is indicative of the amount

of eventual gene products. Based on these attributes together with the informative associations

between differential methylation (DMGs) and positive selection (d_N/d_S), we propose that epigenetic

modification is a complementary mechanism facilitating phenotypic diversity of the *Aquilegia* species.

Limitations and future directions

Our study has important limitations. First of all, the small sample size in our study may introduce bias

and inflation of false positives, and we postulate that our findings should be interpreted carefully and

considered exploratory. When investigating the association between genetic divergence and

evolutionary events, it is impossible to deny the roles of other evolutionary forces. We acknowledge

that the lineage-specific allele frequencies are possibly consequences of genetic drift and that genetic

hitchhiking may lead to identification of candidate genes residing in neighboring genomic regions

representing the same driving force. Therefore, we claim that the candidate genes identified to be

associated with adaptive radiation do not necessarily point towards driving evolutionary mechanisms.

They may also be by-products of the long-term process of adaptive radiation. Besides, we only focused

on analysis of CG methylation as puzzles persist regarding the functional roles of CHG and CHH

methylation. While our sample size is not sufficient to yield enough statistical power even if such *trans*-

association tests were to be performed, our findings at least suggest that the epigenetic variability also

possesses autonomy to some extent.

Methods and Materials

Sample collection, DNA extraction and sequencing

In this study, a total of 36 accessions from ten worldwide *Aquilegia* species were collected (**Table S5**).

Among these samples, 18 accessions were collected to represent the two Asian species, *A. japonica* and

A. oxysepala, and their putative hybrid. In addition, four accessions were collected from the other two

Asian species *A. yabeana* and *A. viridiflora*. Furthermore, six and eight accessions were sampled from

the European and North American lineages, respectively. All the 36 accessions were grown in green
house under the same conditions (25°C/12 hours, 16°C/12 hours). Genomic DNA was extracted from
fresh mature leaves using TianGen plant genomic DNA kit. Whole genome resequencing and bisulfite
sequencing were performed on the extracted genomic DNA using the Illumina X-ten platform (Illumina,
California, USA). Short-insert (350 bp) DNA libraries of all accessions were constructed by NovoGene
(NovoGene, Tianjin, China). Genome assembly of an admixed species *A. coerulea* "Goldsmith" was
obtained from Phytozome v12.1 (https://phytozome.jgi.doe.gov) as the reference genome [17]. All data
generated from the study were submitted to EBI under the accession number PRJEB34182.

Sequence assembly, population genetic structure and functional annotation
Whole genome sequences of each accession were aligned against the reference genome using Burrows-
Wheeler Aligner (BWA) [49]. Raw assemblies were realigned using IndelRealigner provided in the
Genome Analysis Tool Kit [50].　Single nucleotide polymorphisms (SNPs) and insertions/deletions
(INDELs) were reported using SAMtools [51]. Only the high-quality variants (SNPs and INDELs) (read
depth > 3, mapping quality > 20 and missing allele < 1%) were retained for subsequent population
genomics analyses. Genomic context and predicted functional effects of the identified variants were
reported for each of the 36 samples separately. Intergenic variants occurring within 5 kb up- and down-
stream of TSS or TES were defined as regulatory variations. Functional annotation of each identified
variant was performed using SnpEff [52].
To infer the phylogenetic relationships of the ten *Aquilegia* species, NJ trees were reconstructed for
each chromosome and the whole genome dataset using MEGA 7 [53]. PCA was also carried out to
examine the genetic relatedness of the 36 *Aquilegia* accessions [54]. Ancestral components were
estimated using ADMIXTURE [55] with the assumed population number ranging from one to ten.
Optimal population composition with the least cross-validation error was selected to decompose
ancestral admixture. To obtain the genome-wide nucleotide variation pattern, nucleotide diversity (π)
and genetic differentiation (Weir and Cockerham's F_{ST}) were calculated for each 100 kb non-overlapping
sliding window using VCFtools [56, 57]. Pair-wise non-synonymous-to-synonymous (d_N/d_S) ratios of the
ten species were inferred by yn00 program in the Phylogenetic Analysis by Maximum Likelihood (PAML)
package [58]. Inter-lineage d_N/d_S value for each gene was derived by averaging d_N/d_S values obtained
from all pairwise comparisons of samples belonging to the two lineages under investigation. Candidate
genes with the 5% highest and 5% lowest d_N/d_S values were considered to have undergone strong
positive and purifying selection, respectively.

Cytosine methylation pattern and epigenetic population structure

Whole genome bisulfite sequencing data were pre-processed using TrimGalore (https://www.
bioinformatics.babraham.ac.uk/projects/trim_galore/, accessed August 21, 2018). Trimmed paired-end
reads were aligned to the reference genome using Bismark [59] with a moderately stringent minimum-
score function (L,0,-0.3). De-duplicated alignments of the 36 *Aquilegia* accessions were used to report
cytosine methylation level using bismark_methylation_extractor, on loci with a read depth ≥ 3. Genomic
annotations of the methylated cytosine site were identified based on the reference genome using an in-
house Python script. PCA was conducted to infer the CG-methylomic relatedness of the ten *Aquilegia*
species. Differential cytosine methylation was determined at both the gene and chromosome levels,
respectively. At the gene level, we determined DMRs for each 100 bp non-overlapping sliding window
using Cochran-Mantel-Haenszel (CMH) test to account for imbalanced read depth (Supplementary
Notes). Genomic regions that possess a Benjamini-Hochberg adjusted *p* value < 0.05 and show inter-
specific or inter-lineage methylation divergence higher than 25% are defined as significant DMRs. Genes
with > 20% of the genic region being DMR(s) were defined as DMGs. Chromosome-level methylation
patterns were measured by chromosomal methylation discrepancy index (MDI) [60]. Methylation
patterns of the identified DMGs were visually verified on Integrative Genomics Viewer [61] prior to
downstream analyses and biological interpretation. In addition, we identified CG islands from the *A.*
coerulea "Goldsmith" reference genome using EMBOSS cpgplot with default settings [62]. Only the
identified CG-enriched genomic regions with > 200 bp were defined as CG islands. We then investigated
inter-specific and inter-lineage methylation divergence in and around the CG islands.

Associations between the genetic variation and cytosine methylation

To address how the genetic and epigenetic variations interacted to promote species diversification, we
tested for associations between the identified DMGs and genes under positive selection by a Chi-square
test. Linear regression model was adopted to measure the direct causal effect of CG-loss variation on CG
methylation. To further assess whether genetic variations drive the establishment of DMG, driving
mutations of DMRs between the *A. japonica* and *A. oxysepala* were identified using an Eigenstrat
method (See Supplementary Notes for more details) [63].

Identification of conservative clade-specific variant

CCVs were defined as these that were conserved across all samples belonging to the same

species/lineage but not present in any sample of the other species/lineages. Since the biological

consequences of heterozygous variants are less affirmable, only the homozygous point mutations and

INDELs were considered in the characterization of CCVs, including frameshift, stop-gain, stop-loss, start-

loss and splicing-alteration variations.

Functional enrichment analysis

The aforementioned genetic and epigenetic analyses identified candidate genes that might be

associated with the rapid diversification of the *Aquilegia* species from different perspectives. These

candidate genes were employed to conduct functional enrichment analyses using R package topGO [64].

Enriched GO terms that possessed a *p* value <0.05 and has an occurrence ≥ 5 in the background were

considered statistically significant. Since the statistical tests performed by topGO are not independent,

multiple testing correction does not apply [64]. Structures of functional domains of targeted genes were

determined based on the InterPro database (https://www.ebi.ac.uk/interpro, accessed January 25,

2019). Distribution pattern of the identified candidate genes and their related functional pathways was

visualized using the R package jvenn [65].

Author Contributions

L.F.L. conceived this project. T.L. and M.R.L. developed statistical analysis pipeline. T.L., M.R.L., N.D.,

Z.H.W., L.Z.L., and X.G. carried out experiments and analyzed the data. T.L., M.R.L., N.D., Z.H.W., L.Z.L.,

X.G., and L.F.L. participated in discussion and interpreted the data. T.L. and L.F.L. wrote the manuscript.

All authors read and approved the manuscript.

Acknowledgments

We thank Aköz Gökçe for her constructive comments on our manuscript. This work was financially

supported by National Natural Science Foundation of China (31670382), Shanghai Pujiang Program

(19PJ1401500), Start-up funding at Fudan University (JIH1322105) and the Department of Science and

Technology of Jilin Province (20190201299JC).

Financial Disclosure

The funders had no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript.

Competing Interests

The authors have declared that no competing interests exist.

References

1.   Schluter D. Ecological causes of adaptive radiation. American Naturalist. 1996; 148: S40-S64.

2.   Schluter D. The Ecology of Adaptive Radiation. Heredity. 2000; 86(6): 749-50.

3.   Givnish TJ. Adaptive radiation and molecular systematics: aims and conceptual issues. In: Givnish TJ, Systma KJ, eds. . Molecular Evolution and Adaptive Radiation. 1997: 1-54.

4.   Givnish TJ. Adaptive radiation versus 'radiation' and 'explosive diversification': why conceptual distinctions are fundamental to understanding evolution. New Phytologist. 2015; 207(2): 297-303.

5.   Simpson GG, Olson EC. The major features of evolution. Quarterly Review of Biology. 1953; 8(1): 87-8.

6.   Losos JB. Adaptive radiation, ecological opportunity, and evolutionary determinism. American Society of Naturalists E. O. Wilson award address. American Naturalist. 2010; 175(6): 623-39.

7.   Carlquist, Motley TJ. Tarweeds and Silverswords, Evolution of the Madiinae (*Asteraceae*). Economic Botany. 2003; 58(1): 123-4.

8.   Losos JB, Ricklefs RE. Adaptation and diversification on islands. Nature. 2009;457(7231):830-6.

9.   Lamichhaney S, Jonas B, Markus S, Liman A, Khurram M, Manfred G, Alvaro MB, et al. Evolution of Darwin's finches and their beaks revealed by genome sequencing. Nature. 2015; 518(7539): 371-5.

10.   Irisarri I, Singh P, Koblmüller S, Torresdowdall J, Henning F, Franchini P, et al. Phylogenomics uncovers early hybridization and adaptive loci shaping the radiation of Lake Tanganyika cichlid fishes. Nature Communications. 2018; 9(1): 3159-71.

11.   Wagner CE, Harmon LJ, Seehausen O. Ecological opportunity and sexual selection together predict adaptive radiation. Nature. 2012; 487(7407): 366-U124.

12.   Simpson EH. Measurement of Diversity. Nature. 1949; 163(4148): 688-688.

13.   Berner D, Salzburger W. The genomics of organismal diversification illuminated by adaptive radiations. Trends in Genetics. 2015; 31(9): 491-9.

14.   Brawand D, Wagner CE, Li YI, Malinsky M, Keller I, Fan SH, et al. The genomic substrate for adaptive radiation in African cichlid fish. Nature. 2014; 513(7518): 375-81.

15.   Ford AG, Rüber L, Newton J, Dasmahapatra KK, Balarin JD, Bruun K, et al. Niche divergence
facilitated by fine-scale ecological partitioning in a recent cichlid fish adaptive radiation. Evolution. 2016;
70(12): 2718-35.

16.   Fior S, Mingai L, Bengt O, Roberto V, Hodges SA, Lino O, et al. Spatiotemporal reconstruction of
the *Aquilegia* rapid radiation through next-generation sequencing of rapidly evolving cpDNA regions.
New Phytologist. 2013; 198(2): 579-92.

17.   Filiault DL, Ballerini ES, Mandakova T, Akoz G, Derieg NJ, Schmutz J, et al. The *Aquilegia* genome
provides insight into adaptive radiation and reveals an extraordinarily polymorphic chromosome with a
unique history. elife. 2018; 7.

18.   Munz. Aquilegia: the cultivated and wild columbines. Gentes Herbarum. 1946; 7: 1-150.

19.   Bastida JM, Herrera CM. Extended phylogeny of *Aquilegia*: the biogeographical and ecological
patterns of two simultaneous but contrasting radiations. Plant Systematics and Evolution. 2010; 284(3-
4): 171-85.

20.   Whittall JB, Hodges SA. Pollinator shifts drive increasingly long nectar spurs in columbine
flowers. Nature. 2007; 447(7145): 706-9.

21.   Alcántara JM, Bastida JM, Rey PJ. Linking divergent selection on vegetative traits to
environmental variation and phenotypic diversification in the Iberian columbines (*Aquilegia*). Journal of
Evolutionary Biology. 2010; 23(6): 1218-33.

22.   Kramer EM, Hodges SA. *Aquilegia* as a model system for the evolution and ecology of petals.
Philosophical Transactions Biological Sciences. 2010; 365(1539): 477-90.

23.   Sharma B, Yant L, Hodges SA, Kramer EM. Understanding the development and evolution of
novel floral form in *Aquilegia*. Current Opinion in Plant Biology. 2014; 17(1): 22-7.

24.   Garrido JL, Alcántara JM, Rey PJ, Medrano M, Guitián J, Castellanos MC, et al. Geographic
genetic structure of Iberian columbines (gen. *Aquilegia*). Plant Systematics and Evolution. 2017; 303(9):
1145-60.

25.   Tang LL, Yu Q, Sun JF, Huang SQ. Floral traits and isolation of three sympatric *Aquilegia* species
in the Qinling Mountains, China. Plant Systematics and Evolution. 2007; 267(1/4): 121-8.

26.   Li LF, Wang HY, Pang D, Liu Y, Liu B, Xiao HX. Phenotypic and genetic evidence for ecological
speciation of *Aquilegia japonica* and A. *oxysepala*. New Phytologist. 2014; 204(4): 1028-40.

27.   Li LF, Pang D, Liao QL, Xiao HX. Genomic and EST microsatellite markers for *Aquilegia* flabellata
and cross-amplification in A. *oxysepala* (Ranunculaceae). American Journal of Botany. 2011; 98(8):213-5.

28.   Li MR, Wang HY, Ding N, Lu T, Huang YC, Xiao HX, et al. Rapid divergence followed by adaptation
to contrasting ecological niches of two closely related columbine species *Aquilegia japonica* and A.
oxysepala. Genome Bioligy and Evolution. 2019; 11(3): 919-30.

29.   Zhang H, Dawe RK. Mechanisms of plant spindle formation. Chromosome Research. 2011; 19(3):
335-44.

30.   Aguirre-Portoles C, Bird AW, Hyman A, Canamero M, de Castro IP, Malunnbres M. *Tpx2* controls
spindle integrity, genome stability, and tumor development. Cancer Research. 2012; 72(6): 1518-28.

31.   Fukui K, Nakagawa N, Kitamura Y, Nishida Y, Masui R, Kuramitsu S. Crystal structure of *MutS2*
endonuclease domain and the mechanism of homologous recombination suppression. Journal of
Biological Chemistry. 2008; 283(48): 33417-27.

32.   Machado-Schiaffino G, Henning F, Meyer A. Species-specific differences in adaptive phenotypic
plasticity in an ecologically relevant trophic trait: hypertrophic lips in Midas cichlid fishes. Evolution.
2014; 68(7): 2086-91.

33.   Bossdorf O, Richards CL, Massimo P. Epigenetics for ecologists. Ecology Letters. 2008; 11(2):
106-15.

34.   Diez CM, Roessler K, Gaut BS. Epigenetics and plant genome evolution. Current Opinion in Plant
Biology. 2014; 18(1): 1-8.

35.   Verhoeven KJ, Vonholdt BM, Sork VL. Epigenetics in ecology and evolution: what we know
and what we need to know. Molecular Ecology. 2016; 25(8): 1631-8.

36.   Richards CL, Oliver B, Verhoeven KJF. Understanding natural epigenetic variation. New
Phytologist. 2010; 187(3): 562-4.

37.   Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, Colome-Tatche M, et al. Ecological plant
epigenetics: Evidence from model and non-model species, and the way forward. Ecology Letters. 2017;
20(12): 1576-90.

38.   Becker C, Hagmann J, Jonas M, Daniel K, Oliver S, Karsten B, et al. Spontaneous epigenetic
variation in the *Arabidopsis thaliana* methylome. Nature. 2011; 480(7376): 245.

39.   Schmitz RJ, Schultz MD, Lewsey MG, O'Malley RC, Urich MA, Ondrej L, et al. Transgenerational
epigenetic instability is a source of novel methylation variants. Science. 2011; 334(6054): 369-73.

40.   Cortijo S, Wardenaar R, Colome-Tatche M, Gilly A, Etcheverry M, Labadie K, et al. Mapping the
epigenetic basis of complex traits. Science. 2014; 343(6175): 1145-8.

41.   Smith AK, Kilaru V, Kocak M, Almli LM, Mercer KB, Ressler KJ, et al. Methylation quantitative

trait loci (meQTLs) are consistently detected across ancestry, developmental stage, and tissue type. BMC

Genomics. 2014; 15(1): 145.

42.   Drong AW, George N, Hedman AK, Eshwar M, Elin G, Small KS, et al. The presence of

methylation quantitative trait loci indicates a direct genetic influence on the level of DNA methylation in

adipose tissue. PloS ONE. 2013; 8(2): e55923.

43.   Fisher VA, Wang L, Deng X, Sarnowski C, Cupples LA, Liu CT. Do changes in DNA methylation

mediate or interact with SNP variation? A pharmacoepigenetic analysis. BMC Genetics. 2018; 19(Suppl

1): 70.

44.   Li L, Wang C, Lu T, Lin S, Hu YQ. Indirect effect inference and application to GAW20 data. BMC

Genetics. 2018; 19(Suppl 1): 67.

45.   Bewick AJ, Schmitz RJ. Gene body DNA methylation in plants. Current Opinion in Plant Biology.

2017; 36: 103-10.

46.   Takuno S, Gaut BS. Gene body methylation is conserved between plant orthologs and is of

evolutionary consequence. Proceedings of the National Academy of Sciences of the United States of

America. 2013; 110(5): 1797-802.

47.   Bewick AJ, Ji L, Niederhuth CE, Willing EM, Hofmeister BT, Shi X, et al. On the origin and

evolutionary consequences of gene body DNA methylation. Proceedings of the National Academy of

Sciences of the United States of America. 2016; 113(32): 9111-6.

48.   Zilberman D. An evolutionary case for functional gene body methylation in plants and animals.

Genome Biology. 2017; 18(1): 87.

49.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics, 2010; 25(14): 1754-60.

50.   Mckenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome

Analysis Toolkit: A map reduce framework for analyzing next-generation DNA sequencing data. Genome

Research. 2010; 20(9): 1297-303.

51.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence alignment/map

format and SAMtools. Bioinformatics. 2009;2 5(16): 2078-9.

52.   Cingolani P, Platts A, Le LW, Coon M, Nguyen T, Luan W, et al. A program for annotating and

predicting the effects of single nucleotide polymorphisms, SnpEff. Fly. 2012; 6(2): 80-92.

53.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for

bigger datasets. Molecular Biology and Evolution. 2016; 33(7): 1870.

54.   Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. A high-performance computing
toolset for relatedness and principal component analysis of SNP data. Bioinformatics. 2012; 28(24):
3326-8.

55.   Alexander DH, Novembre JK. Fast model-based estimation of ancestry in unrelated individuals.
Genome Research. 2009; 19(9): 1655-64.

56.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format
and VCFtools. Bioinformatics. 2011; 27(15): 2156-8.

57.   Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population Structure.
Evolution. 1984; 38(6): 1358-70.

58.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Molecular Biology and Evolution.
2007; 24(8): 1586-91.

59.   Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics. 2011; 27(11): 1571-2.

60.   O''Sullivan DE, Johnson KC, Skinner L, Koestler DC, Christensen BC. Epigenetic and genetic
burden measures are associated with tumor characteristics in invasive breast carcinoma. Epigenetics.
2016; 11(5): 344-53.

61.   Thorvaldsdottir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. Brief Bioinformatics. 2013; 14(2): 178-92.

62.   Rice P, Longden I, Bleasby A. EMBOSS: The European molecular biology open software suite.
Trends in Genetics. 2000; 16(6): 276-7.

63.   Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, David R. Principal components
analysis corrects for stratification in genome-wide association studies. Nature Genetics. 2006; 38: 904-9.

64.   Alexa A, Rahnenfuhrer J, Lengauer T. Improved scoring of functional groups from gene
expression data by decorrelating GO graph structure. Bioinformatics. 2006; 22(13): 1600-7.

65.   Bardou P, Mariette J, Escudie F, Djemiel C, Klopp C. jvenn: an interactive Venn diagram viewer.
BMC Bioinformatics. 2014; 15: 293.

Figure legend

Figure 1. **Phylogenetic relationship and population structure of the ten worldwide *Aquilegia* species**.

(a) Phylogenetic tree of the 36 accessions constructed by neighbor joining algorithm based on whole-

genome SNPs. (b) PCA reveals genetic similarity within each of the three lineages and genetic disparity

between lineages based on 15,988 LD-pruned SNPs. Curves for each lineage denote 99% probability

ellipses estimated from distribution of the first two principal components. (c) Population admixture of

the 36 *Aquilegia* accessions. Most accessions exhibit almost pure ancestry composition matching the

presumed evolutionary history of the corresponding species.

Figure 2. **Functional enrichment of genes harboring highly impactful CCVs and DMGs.** CCV-containing

genes specific to either of the two lineages/species being compared were merged to construct a target

gene set. Ratio denotes proportion of CCV-containing genes or DMGs in the corresponding gene set of

interest. Absence of dot indicates no significant enrichment.

Figure 3. **Patterns of cytosine methylation for the ten worldwide *Aquilegia* species.** (a) Genome-wide

cytosine methylation level distributions of 36 accessions. (b) MDI illustrates chromosome-level CG

methylation similarity. *Aquilegia viridiflora* was used as the reference. (c) CG methylation profiling in

genic region across the four *Aquilegia* groups. Each row represents one genic region starting at 5kb

upstream of its TSS and terminating at 5kb downstream of its TES, sorted by mean methylation level of

all analyzed CG sites. Length of gene body was normalized. (d) CG methylation profiling in and around

CG island.

Figure 4. **Association between CG-loss variations and epigenetic variability**. (a) Top 3,000 most variable

CG sites containing CG-loss variations. (b) Top 3,000 most variable non-variant CG sites across 36

accessions show clade-specific methylation patterns. CG methylation in the hybrid tends to be

neutralized possibly due to heterozygosity. (c) Linear regression demonstrates that CG-loss variations

explain a large proportion of CG methylation variation. (d) Summary of composition of each category

with regard to whether each CG site contains a CG-loss variation. Epigenetic variability was determined

by standard deviation in methylation β value across all 36 accessions. CG sites with top 10,000, 10,001-

50,000 and 50,001-150,000 largest standard deviation were ordinally labelled as possessing "very high",

"high" and "moderate" variability respectively. The rest CG sites were labelled as possessing "low"

variability. (e) Association test shows most DMRs are independent of *cis*-acting SNPs. Results under

different significance levels are compared for this exploratory analysis.

Supplementary file

Figure S1. Per-chromosome phylogenetic trees reconstructed using neighbor-joining method.

Polymorphisms detected on each chromosome were retrieved separately to infer the phylogeny.

Figure S2. Distribution of nucleotide diversity (π) at the whole-genome level and the per-chromosome

level. Nucleotide diversity was estimated for each lineage pooling corresponding species, as well as for *A.*

japonica and *A. oxysepala*.

Figure S3. Spearman correlation of the genome-wide nucleotide variation pattern for each 100-kb

sliding window between European and Asian (a), North American and Asian (b), European and North

American (d), *Aquilegia japonica* and *A. oxysepala* (d). Each dot represents a 100-kb sliding window.

Values on the x- and y-axis are the nucleotide diversity (π) for each sliding window.

Figure S4. Overlapping of low diversity genomic region (LDGR) in the three lineages. 148 LDGRs with 5%

lowest nucleotide diversity were defined as LDNRs in each lineage, totaling 241 unique regions.

Figure S5. Pair-wise d_N/d_S ratio for all genes between the species within and between the Asian,

European and North American lineages.

Figure S6. PCA illustrates three distinct clusters corresponding to the three lineages. Asian species

further demonstrated higher inter-specific divergence than the American and the European species. PCA

was performed based on 588,659 loci with sufficiently high sequencing quality.

Figure S7. Illustration of differential methylation in two photosynthesis genes. CG methylation pattern

of two genes, *Aqcoe7G230600* photosystem I *PsaA/PsaB* (a) and *Aqcoe7G231300 CemA* (b) in *A.*

japonica and *A. oxysepala* throughout the gene body region. Red bars indicate methylation level (0-100)

at CG sites. Genomic coordinates on the chromosome 7 are annotated.

Figure S8. Overlapping of the CCVs (a) and DMGs (b) identified in inter-lineage/species comparisons. A

considerable proportion of these CCVs (84.1%) and DMGs (51.3%) were shared by two or more inter-

lineage/species comparisons.

Figure S9. Venn analyses of the candidate genes carrying CCVs and DMGs. Each subpanel indicates the

comparison between the *A. japonica* and *A. oxysepala* (a), *A. japonica* and North American (b), *A.*

japonica and European (c), *A. oxysepala* and North American (d), *A. oxysepala* and European (e), North

American and European (f).

Table 1. Information of the high-impact conservative clade-specific variants (CCVs) in the cell
659 reproduction related genes.

Gene	Variant-carrying	Reference	Chromosome	Position	Reference allele	Variant	Annotation	Gene function
Aqcoe1G273400	Asian	American	Chr1	18994915	GAA	GAAA	frameshift	DNA mismatch repair protein *MutS2*
Aqcoe2G151500	European	American	Chr2	15305837	A	G	splicing	PIF1-like helicase
	European	American		15307442	A	C	stop gain	
	European	American		15309865	AATATATAT	AATATATATAT	frameshift	
	European	Asian		15307442	A	C	stop gain	
	European	Asian		15309865	AATATATAT	AATATATATAT	frameshift	
	A. oxysepala	*A. japonica*		15305837	A	G	splicing	
	A. oxysepala	*A. japonica*		15309267	AT	A	frameshift	
Aqcoe2G177700	European	American	Chr2	21794397	TATGCACCAAAGGTATCACGATGC	TATGC	frameshift	PIF1-like helicase
	European	American		21794979	TT	TTGT	frameshift	
	European	Asian		21794397	TATGCACCAAAGGTATCACGATGC	TATGC	frameshift	
	A. oxysepala	*A. japonica*		21795089	CA	C	frameshift	
Aqcoe6G208600	European	American	Chr6	15364081	A	ATCTCTTCG	frameshift	PIF1-like helicase
	European	Asian		15364081	A	ATCTCTTCG	frameshift	
	A. japonica	*A. oxysepala*		15364330	TAA	TA	frameshift	
Aqcoe6G253800	European	American	Chr6	22789898	C	T	stop gain	DNA helicase
	European	American		22790012	G	A	splicing	
	European	Asian		22789898	C	T	stop gain	
	A. japonica	*A. oxysepala*		22790012	G	A	splicing	
Aqcoe2G276600	Asian	American	Chr2	33314422	AGGGGG	AGGGGGG	frameshift	DNA mismatch repair protein *Msh6*
Aqcoe6G160300	*A. japonica*	*A. oxysepala*	Chr6	9414625	G	A	stop gain	*TPX2*
Aqcoe7G062500	*A. oxysepala*	*A. japonica*	Chr7	3789055	G	A	stop gain	cell cycle regulated microtubule associated protein

660
661
662

663   **Table 2.** Significant correlation between differential methylation and natural selection.

Type of selection	Differential methylation	Jap-Oxy[*]	Jap-Ame	Jap-Eur	Oxy-Ame	Oxy-Eur	Ame-Eur
Positive selection	DMG	7.2%	7.3%	11.9%	6.7%	8.4%	8.9%
	non-DMG	4.4%	5.0%	5.6%	4.7%	5.4%	5.4%
	p value	0.11	7.3e-02	3.9e-05	6.7e-02	1.8e-02	2.8e-04
Purifying selection	DMG	3.1%	1.8%	2.4%	2.3%	2.0%	1.9%
	non-DMG	4.3%	4.3%	4.7%	4.9%	5.1%	4.0%
	p value	0.53	3.2e-02	9.1e-02	1.3e-02	8.4e-03	1.0e-02

[*]: Each percentage represents the proportion of genes belonging to either DMGs or non-DMGs
compared between the two corresponding clades that are under corresponding or higher strength of
positive selection. For example, 7.2% indicates that 7.2% DMGs compared between *A. japonica* and *A.*
oxysepala are under strong selection; 4.4% indicates that 4.4% non-DMGs compared between these two
species are under strong selection. *p* values were obtained from Chi-square tests and were not adjusted
for multiple testing due to dependence arising from overlapping gene sets.
Jap: *A. japonica*; Oxy: *A. oxysepala*; Ame: American; Eur: European

European

Asian

American

Size of each list

European

Asian

American

Number of elements: specific (1) or shared by 2, 3, ... lists

(a)

A. japonica

A. oxysepala

20392800   Aqcoe7G230600   20394100

(b)

A. japonica

A. oxysepala

20403000   Aqcoe7G231300   20406500

(a) *A. japonica* and *A. oxysepala*

CCV   DMG
1134   20   275

(b) *A. japonica* and North Amrican

CCV   DMG
1128   44   534

(c) *A. japonica* and European

CCV   DMG
1072   19   398

(d) *A. oxysepala* and North American

CCV   DMG
1339   50   632

(e) *A. oxysepala* and European

CCV   DMG
1380   36   495

(f) North American and European

CCV   DMG
895   36   700