Genetic and epigenetic mechanisms underpinning the

adaptive radiation of *Aquilegia* species

Tianyuan Lu[1,2,3], Ming-Rui Li[1], Ning Ding[1], Zhen-Hui Wang[4], Li-Zhen Lan[1], Xiang Gao[5], and Lin-Feng Li[1,*]

[1]Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, School of Life

Sciences, Fudan University, Shanghai 200438, China;

[2]McGill University and Genome Quebec Innovation Center, Montreal H3A 0G1, Quebec, Canada;

[3]Lady Davis Institute, SMBD JGH, Montreal H3A 1A3, Quebec, Canada;

[4]Faculty of Agronomy, Jilin Agricultural University, Changchun 130118, China;

[5]Key Laboratory of Molecular Epigenetics of the Ministry of Education, Northeast Normal University,

Changchun 130024, China.

Correspondence author:

Lin-Feng Li (lilinfeng@fudan.edu.cn)

Summary

• Elucidating how the intrinsic factors interact with extrinsic triggers to determine species diversification

is crucial to understanding the evolution and persistence of biodiversity. The genus *Aquilegia* is a model

system to address the evolutionary mechanisms underpinning rapid adaptive radiation.

• We surveyed the genomes and methylomes of ten worldwide *Aquilegia* species to investigate whether

specific genetic and epigenetic architectures are involved in the diversification of Asian, European and

North American columbine species.

• The resulting phylogenies and population structure inferences revealed high divergence among the

Asian, European and North American species. Candidate genes identified at the genetic and epigenetic

levels are functionally correlated with diverse adaptation-related traits such as cell reproduction, flower

development, and stress tolerance. In particular, a considerable proportion of the selection genes and

their associated pathways show overlaps among the intra- and inter-lineage comparisons. Moreover,

while CG-loss variations can lead to depletion of cytosine methylation level, epigenetic modification is a

complementary intrinsic factor that intertwines with genetic mechanism to facilitate the diversification

of *Aquilegia* species.

• Our findings suggest that specific genetic and epigenetic architectures have conferred high

adaptability to the columbine species to cope with diverse external conditions, which eventually led to

the rapid radiation of *Aquilegia* species.

Key words: Adaptive radiation; *Aquilegia*; Methylome; Speciation; Whole genome resequencing

Introduction

Adaptive radiation is the rapid diversification of a single ancestral species into a vast array of common descendants that inhabit different ecological niches or use a variety of resources and differ in phenotypic traits required to exploit diverse environments (Schluter, 1996; Givnish, 1997; Schluter, 2000; Givnish, 2015). Disentangling the evolutionary mechanisms underpinning adaptive radiation is fundamental to understanding the evolution and persistence of biodiversity (Simpson & Olson, 1953; Losos, 2010). Well-known animal and plant lineages that diversified through adaptive radiation include Hawaiian silversword, Caribbean anoles, Darwin's finches, and African cichlids (Carlquist & Motley, 2003; Losos & Ricklefs, 2009; Lamichhaney *et al.*, 2015; Irisarri *et al.*, 2018). However, it still remains under-investigated as to why some lineages can diversify rapidly but their close relatives or other sympatrically distributed lineages do not. In the past decades, accumulating evidence from diverse radiation lineages suggests that extrinsic environmental variables (e.g., resource availability) and intrinsic factors (e.g., lineage-specific genetic variations) can interact to determine the rate and volume of species diversification (Wagner *et al.*, 2012). Among the extrinsic triggers, ecological opportunity is considered as the primary mechanism that causes rapid adaptive radiation through the acquisition of key innovations, invasion of new environments and extinction of competitors (Simpson, 1949; Schluter, 2000). On the other hand, new species can also arise as a result of natural selection acting on intrinsic factors (e.g., advantage alleles) that ultimately generate both the phenotypic disparity and similarity among closely related species (Berner & Salzburger, 2015). An illustration of rapid adaptive radiation is the African cichlid fishes, on which the extrinsic environmental factors (e.g., ecological specialization) and extrinsic traits (e.g., adaptive introgression) have acted together to provoke the repeated adaptive radiation in geographically isolated lakes (Wagner *et al.*, 2012; Brawand *et al.*, 2014; Ford *et al.*, 2016; Irisarri *et al.*, 2018).

The genus *Aquilegia* L. (columbine) is a well-recognized model system to address the evolutionary mechanisms underlying rapid adaptive radiation (Fior *et al.*, 2013; Filiault *et al.*, 2018). It includes approximately 70 recently diversified species that are widely distributed in the temperate areas of North America and Eurasia (Munz, 1946). Phylogenetic and geographic inferences have illustrated two independent adaptive radiations of North American and European lineages from the distinct ancestral Asian species (Bastida & Herrera, 2010; Fior *et al.*, 2013). Floral diversification of the North American species is highly correlated with the pollinator specialization (Whittall & Hodges, 2007; Alcántara *et al.*, 2010; Kramer & Hodges, 2010; Sharma *et al.*, 2014). In contrast, ecological adaptation and geographic isolation are thought as the major driving forces that promoted the rapid radiation of European species

(Fior *et al.*, 2013; Garrido *et al.*, 2017). In Asia, shifts in pollinator and ecological habitat are both
proposed to be the underpinning mechanisms that resulted in the diversification of more than 20
morphologically distinct species (Tang *et al.*, 2007; Li *et al.*, 2014). Unlike the North American and
European lineages, the Asian columbines consist of four highly divergent lineages corresponding to
respective geographic origins and evolved independently under distinct external triggers (Tang *et al.*,
2007; Fior *et al.*, 2013; Li *et al.*, 2014). The independent adaptive radiations of North American and
European lineages from the more ancestral but independently diversified Asian columbine species
provide an ideal system to examine how the extrinsic and intrinsic triggers interact to promote the
diversification of *Aquilegia* species.

In this study, we surveyed the genomes and DNA methylomes of 36 accessions from ten worldwide
columbine species to address the underlying evolutionary mechanisms that caused the diversification of
Asian, European and North American columbine species. Of the Asian lineage, four phylogenetically
distinct species (*A. yabeana*, *A. viridiflora*, *A. oxysepala* and *A. japonica*) were selected according to their
geographic distributions and ecological habitats. *Aquilegia japonica* and *A. oxysepala* are sister species
inhabiting alpine tundra and low altitude forest niches in northeastern China, respectively (Li *et al.*, 2011;
Li *et al.*, 2014). Our previous studies have documented that natural selection during ecological
specialization together with genetic drift under geographic isolation caused the rapid evolution of
reproductive isolation between the two species (Li *et al.*, 2014; Li *et al.*, 2019). Here we further
investigate how the genetic and epigenetic mechanisms conferred adaptability to the two species to
cope with contrasting ecological conditions. In addition, patterns of nucleotide variation and cytosine
methylation were also evaluated for the *A. yabeana* and *A. viridiflora*. The former species shares highly
similar morphological traits and ecological niches with the *A. oxysepala* but is allopatrically distributed in
the northern China. In contrast, while the *A. viridiflora* is sympatrically distributed with *A. yabeana* and *A.*
oxysepala in the northern and northeastern China, it often occupies rocky and sandy ecological niches.
These attributes allow us to experimentally test how the genetic and epigenetic mechanisms
contributed to the phenotypic divergence and ecological specialization of these Asian species. In
addition, we also surveyed the genetic and epigenetic architectures of six North American and European
columbine species. We aimed to examine whether the adaptive radiations of North American and
European lineages are determined by extrinsic factors related to different ecological opportunities or
triggered by intrinsic traits utilizing the lineage-specific genomic and epigenomic architectures. Our
study will provide a genome-wide view of how the diverse evolutionary mechanisms facilitate the rapid
species diversification in the genus *Aquilegia*.

Methods and Materials

Sample collection, DNA extraction and sequencing

In this study, a total of 36 accessions from ten worldwide *Aquilegia* species were collected (**Table S1**).

Among these samples, 18 accessions were collected to represent the two Asian species, *A. japonica* and

A. oxysepala, and their putative hybrid. In addition, four accessions were collected from the other two

Asian species *A. yabeana* and *A. viridiflora*. Furthermore, six and eight accessions were sampled from

the European and North American lineages, respectively. All the 36 accessions were grown in green

house under the same conditions (25°C/12 hours, 16°C/12 hours). Genomic DNA was extracted from

fresh mature leaves using TianGen plant genomic DNA kit. Whole genome resequencing and bisulfite

sequencing were performed on the extracted genomic DNA using the Illumina X-ten platform (Illumina,

California, USA). Short-insert (350bp) DNA libraries of all accessions were constructed by NovoGene

(NovoGene, Tianjin, China). Genome sequences of the American species *A. coerulea* was obtained from

Phytozome v12.1 (Filiault *et al.*, 2018, https://phytozome.jgi.doe.gov). All data generated from the study

were submitted to EBI under the accession number PRJEB34182.

Sequence assembly, population genetic structure and functional annotation

Whole genome sequences of each accession were aligned against the reference genome of *A. coerulea*

using Burrows-Wheeler Aligner (BWA) (Li & Durbin, 2009). Raw assemblies were realigned using

IndelRealigner provided in the Genome Analysis Tool Kit (Mckenna *et al.*, 2010). Single nucleotide

polymorphisms (SNPs) and insertions/deletions (INDELs) were reported using SAMtools (Li *et al.*, 2009).

Only the high-quality variants (SNPs and INDELs) (read depth > 3 and mapping quality > 20) were

retained for subsequent population genomics analyses. Genomic context and predicted functional

effects of the identified variants were reported for each of the 36 samples separately. Intergenic

variants occurring within 5 kb up- and down-stream of transcription start site (TSS) or transcription end

site (TES) were defined as regulatory variations. Functional annotation of each identified variant was

performed using SnpEff (Cingolani *et al.*, 2012).

To infer the phylogenetic relationships of the ten *Aquilegia* species, neighbor-joining (NJ) trees were

reconstructed for each chromosome and the whole genome dataset using MEGA 7 (Kumar *et al.*, 2016).

Principal component analysis (PCA) was also carried out to obtain the clustering pattern of the 36

Aquilegia accessions (Zheng *et al.*, 2012). Ancestral components were estimated using ADMIXTURE

(Alexander & Novembre, 2009) with the assumed population number ranging from one to five. Optimal

population composition with the lowest cross-validation error was selected to account for ancestry. To

obtain the genome-wide nucleotide variation pattern, nucleotide diversity (π) and genetic

differentiation (F_{ST}) were calculated for each 100 kb non-overlapping sliding window using VCFtools

(Danecek *et al.*, 2011). Pair-wise non-synonymous (d_N)/synonymous (d_S) ratios of the ten species were

inferred by yn00 program in the Phylogenetic Analysis by Maximum Likelihood (PAML) package (Yang

2007). Candidate genes that showed the top 5% highest and lowest d_N/d_S values were supposed to

undergo positive and purifying selection, respectively.

Cytosine methylation pattern and epigenetic population structure

Whole genome bisulfite sequencing data were pre-processed using TrimGalore (https://www.

bioinformatics.babraham.ac.uk/projects/trim_galore/, accessed August 21, 2018). Trimmed paired-end

reads were aligned to the reference genome of *A. coerulea* using Bismark (Krueger & Andrews, 2011)

with a moderately stringent minimum-score function (L,0,-0.3). De-duplicated alignments of the 36

Aquilegia accessions were used to report cytosine methylation level using bismark_methylation_

extractor, on loci with a read depth > 3. Genomic annotations of the methylated cytosine site were

identified based on the reference genome using an in-house Python script. Overall CG methylation level

for each accession was calculated by pooling the methylation on all CG site. PCA was conducted to infer

the genome-wide CG methylation clustering pattern of the ten *Aquilegia* species. Differential cytosine

methylation was determined at both the gene and chromosome levels, respectively. At the gene level,

we determined differentially methylated region (DMR) for each 100 bp non-overlapping sliding window

using Cochran-Mantel-Haenszel (CMH) test to account for imbalanced read depth (Supplementary

Information). Only the genomic regions that possess Benjamini-Hochberg adjusted *p* value < 0.05 and

show inter-specific or inter-lineage methylation divergence higher than 25% are defined as significant

DMRs. Genes with > 20% of the genic region being DMR(s) were defined as differentially methylated

gene (DMG). Chromosome-level methylation pattern was measured by chromosomal methylation

discrepancy index (MDI) according to (O'Sullivan *et al.*, 2016). Methylation patterns of the identified

DMGs were visually verified before biological interpretation on Integrative Genomics Viewer

(Thorvaldsdottir *et al.*, 2013). In addition, we identified CG islands from the *A. coerulea* reference

genome using EMBOSS cpgplot with default settings (Rice *et al.*, 2000). Only the identified CG-enriched

genomic regions with > 200 bp were defined as CG islands. We then investigated inter-specific and inter-

lineage methylation divergence in and around the CG islands.

Associations between the genetic variation and cytosine methylation

To address how the genetic and epigenetic variations interact to promote species diversification, we

tested for associations between the identified DMGs and genes under positive selection by Chi-square

test. Linear regression model was adopted to measure the direct causal effect of CG-loss variation on CG

methylation *in situ*. To further assess whether genetic variations drive the establishment of DMG,

driving mutations of DMRs between the *A. japonica* and *A. oxysepala* were identified using the

Eigenstrat method (Supplementary Information) (Price *et al.*, 2006). Given that only a limited number of

DMGs were identified, differentially stringent cutoffs (5×10^{-5}, 5×10^{-8}, and 5×10^{-11}) were adopted to

identify the driver mutations.

Identification of conservative clade-specific variant and functional enrichment analysis

Conservative clade-specific variations (CCVs) were defined as these that were conserved across all

samples belonging to the same species/lineage but not present in any sample of the other

species/lineages. Since the biological consequences of heterozygous variants are less affirmable, only

the homozygous point mutations and INDELs were considered in the characterization of CCVs, including

frameshift, stop-gain, stop-loss, start-loss and splicing-alteration variations. Genes carrying CCVs are

supposed to undergo relaxed selection during the adaptation process. The foregoing genetic and

epigenetic analyses identified candidate genes that might be associated with the rapid diversification of

the *Aquilegia* species. Candidate genes identified at the genetic and epigenetic levels were employed to

conduct enrichment analyses using R package topGO (Alexa *et al.*, 2006). Enriched GO terms that

possessed a *p* value <0.05 and occurred at least five times in the background were considered

statistically significant. Structures of functional domains of targeted genes were determined based on

the InterPro database (https://www.ebi.ac.uk/interpro, accessed January 25, 2019). Distribution pattern

of the identified candidate genes and their related functional pathways was visualized using the R

package Jvenn (Baidou *et al.*, 2014).

Results

Population structure and nucleotide variation pattern

NJ trees were reconstructed for the 36 *Aquilegia* accessions based on 689,123 homozygous SNPs. The

established phylogenies show that accessions of the ten species are broadly clustered as three distinct

lineages corresponding their geographic origins (**Figure 1a** and **Figure S1**). In brief, all the 22 accessions

of the four East Asian species (*A. oxysepala*, *A. japonica*, *A. yabeana* and *A. viridiflora*) cluster as a

monophyletic lineage, with the former two species and their hybrid forming a clade and the latter two

species grouping as a sister clade. In contrast, the West Asian species *A. fragrans* clusters with
geographically adjoining European species. Within the North American lineage, phylogenetic
relationships of the four species vary across the seven chromosomes. In particular, one European
accession of the *A. alpina* var. *alba* falls into the North American lineage at the chromosomes 1 and 3.
Likewise, the PCA and population structure inferences also revealed distinct genetic structure of the
three phylogenetic lineages (**Figure 1b** and **c**). Consistent with the above phylogenies, the *A. alpina* var.
alba accession shares the same ancestral genetic cluster with the North American lineage and the
putative hybrid of the *A. oxysepala* and *A. japonica* possesses a mixed genomic constitution of its
parental species (**Figure 1b** and **c**).

To examine whether specific genomic architectures are involved in the adaptive speciation in the
three lineages, we assessed the nucleotide diversity (π) and genetic divergence (F_{ST}) at both the intra-
and inter-lineage levels. At the inter-lineage level, the Asian columbine species harbored relatively
higher nucleotide diversity compared to the European and North American lineages (**Figure S2**). Notably,
the genetic divergence between the European and North American lineages is obviously higher than
those of the Asian-to-European and Asian-to-North American (**Figure S3**). At the intra-lineage level,
while both the *A. oxysepala* and *A. japonica* possess lower nucleotide diversity, apparently higher overall
genetic divergence (excepting the chromosome 4) is found compared to any of the inter-lineage
comparisons (**Figure S2** and **S3**). We also identified in total 325 high divergence genomic regions (HDGRs,
5% highest F_{ST}) and 241 low diversity genomic regions (LDGRs, 5% lowest π) from the inter-lineage
comparisons. Among these candidate genomic regions, 96 (29.5%) HDGRs and 116 (48.1%) LDGRs are
common between either two or three of these lineages (**Figure S4a** and **b**).

Identification of the selection genes and highly impactful genetic variations
Candidate genes associated with the adaptive divergence at the intra- and inter-lineage levels were
determined according to the following three strategies. First, genes that localized within the HDGR and
LDGR are considered to be exposed to selection in the adaptive evolution process. Between the *A.*
oxysepala and *A. japonica*, we identified seven 100-kb genomic regions that show overlaps between the
HDGRs and LDGRs (**Table S2**). Genes within these genomic regions are functionally associated with the
meiotic nuclear division, adenine methyltransferase and basic cellar activities. At the inter-lineage level,
candidate genes identified from the HDGRs and LDGRs play important roles in telomere maintenance
(e.g., telomeric single stranded DNA binding), plant growth (e.g., gibberellin receptor GID1) and flower
development (e.g., AGAMOUS MADS-box protein) (**Table S2**). In addition, we also find that some

photosynthesis related genes resided in two adjacent HDGRs (chr6:23.9-24.1Mb), including the
cytochrome C protein, NADH-ubiquinone oxidoreductase, photosystem I *ycf3* and photosystem II *PsbT*
(**Table S2**). Of significance, a majority of these candidate genes possess fixation sites at the intra- and
inter-lineage comparisons (**Supplementary file 1**).

The above assessment mainly relied on genome-wide scanning for 100-kb non-overlapping sliding
window, we next employed site-based approach to identify highly impactful CCVs (e.g., frameshift
mutations, stop mutation and splicing alteration) from both the intra- and inter-lineage comparisons.
Our results reveal that a considerable proportion (17.9-40.5%) of the CCVs are identified at the gene
body region (**Table S3**). We then examined the impacts of these identified CCVs on the gene function.
Between the *A. oxysepala* and *A. japonica*, the CCV-carrying genes are enriched in several vital biological
pathways related to cell reproduction, including telomere maintenance, DNA repair, and DNA helicase
activity (**Figure 2** and **Table 1**). For example, two candidate genes (*Aqcoe6G160300* and
Aqcoe7G062500) coding *Xklp2* (*TPX2*) possess species-specific stop-gain variations, which might largely
influence spindle assembly during the mitotic process (Zhang & Dawe, 2011; Aguirre-Portoles *et al.*,
2012). At the inter-lineage level, the CCVs-harboring genes are also functionally involved in the mitotic
chromosome condensation, DNA ligase activity and aminopeptidase activity (**Figure 2** and **Table 1**). Two
CCV-containing genes (*Aqcoe2G276600* and *Aqcoe1G273400*) encoding DNA mismatch repair proteins
MutS/*MSH* and *MutS2* (Fukui *et al.*, 2008) carry one Asian-specific-to-American frameshift variant.

In addition, we also performed genome-wide scanning to identify selection genes from the intra-
and inter-lineage comparisons. Our analyses characterized a total of 4,347 high and 3,566 low d_N/d_S
ratio genes, with 1,773 (40.5%) and 1,584 (44.4%) of the candidate genes sharing between at least two
of the six comparisons (**Figure S5**). Functional analysis of the positive selection genes revealed significant
enrichment in various important pathways, including cell reproduction (e.g., telomere maintenance),
stress tolerance (e.g., defense to fungus and bacterium), and plant development (e.g., response to auxin
and cell growth) (**Figure S6**). Likewise, genes under purifying selection are enriched in diverse
functionally important categories related to cell reproduction (e.g., DNA recombination and mitotic cell
cycle), photosynthesis (e.g., photosystem I reaction center) and basic cellular activities (**Figure S6**).

CG methylation patterns and differentially methylated genes
In parallel with the above genomic analyses, we also investigated CG methylation pattern of the
representative columbine species. Overall, the North American accessions (median CG methylation
level: 91.99%) show significantly higher CG methylation than the European (90.49%, Wilcoxon rank sum

test FDR = 4.1×10^{-3}) and Asian (90.67%, Wilcoxon rank sum test FDR = 2.0×10^{-4}) accessions (**Figure 3a**

and **Table S1**). On the contrary, the difference in CG methylation is not significant between the

European and Asian lineages (FDR = 0.17). To justify pooling samples from the same species/lineage in

downstream analyses, we performed PCA-based clustering to examine CG-cytosine methylation

similarity of all the columbine accessions. The resulting overall methylation pattern highly resembled the

above genomic inferences, with the European and American species falling into two independent groups

while the four Asian species forming three separated clusters (**Figure S7**). Thus, we assessed the CG

methylation pattern for the European and North American lineages as well as the three Asian species (*A.*

japonica, *A. oxysepala* and *A. viridiflora*) separately. Consistent with aforementioned genomic features,

heterogeneous pattern of the CG methylation is also observed among the seven chromosomes, with the

chromosome 4 demonstrating obviously higher overall CG methylation divergence compared to the

other six chromosomes (**Figure 3b**). Then, we further evaluated how CG methylation is deposited at the

genic, putative *cis*-regulatory and CG island regions, respectively. At the genic and regulatory regions, all

three lineages share similar modification pattern with apparent depletion of CG methylation around the

TSS and TES (**Figure 3c**). However, the American lineage exhibits hyper-methylation (more than 10%)

around the center of CG islands and a more dramatic decrease in the CG island shores compared to the

European and Asian species (**Figure 3d**).

To examine the biological impacts of differential CG methylation, DMRs and DMGs were identified

from both the intra- and inter-lineage comparisons, respectively (**Tables S4** and **S5**). At the intra-lineage

level, 3,622 DMRs corresponding to 2,899 DMGs were identified between the *A. japonica* and *A.*

oxysepala. Functional enrichment of these DMGs indicate that the two species may have different

activities in photosynthesis-related pathways, including photosystem I, photosynthesis and chloroplast

pathways (**Figure 2**). For example, two photosynthesis-related genes, *PsaA*/*PsaB* and *CemA*, showed

significantly differential methylation between the two species at the genic regions (**Figure S8a** and **b**). At

the inter-lineage level, apparently more DMGs were identified between the North American and

European species (6,087 genes) compared to those of between the two lineages and Asian species

(3,308-5,003 genes) (**Table S4** and **S5**). DMGs characterized from the inter-lineage comparisons are

mainly involved in the plant growth (e.g., response to auxin) and defense, response to biotic stimulus

and wounding (**Figure 2**). It should be noted that while some of the identified DMGs share by at least

two of the six comparisons (**Figure S9**), candidate genes identified from the genetic and epigenetic levels

show obviously complementary patterns in the functional enrichment analyses (**Figure 2**).

It should be noted that the above genetic and epigenetic assessments identified similar enrichment
pathways from the intra- and inter-lineage comparisons, particularly those related to cell reproduction
and basic cellular activities (**Figure 2**, **Figure S6** and **Table S2**). We then examined how the candidate
genes (CCV-carrying gene, high and low d_N/d_S value gene, and DMG) distribute among the above distinct
strategies. While a considerable proportion of the candidate genes share among the intra- and inter-
lineage comparisons for each of the genetic and epigenetic assessments (**Figure S4** and **S5**), the four
types of candidate gene show a clearly separated distribution pattern in the Venn diagrams (**Figure S10**).
Likewise, the GO enrichment analyses also revealed complementary patterns among the four types of
candidate genes (**Figure 2** and **Figure S6**).
305
Dependence of the epigenetic variability on genetic variations
Both genetic variations and differential CG methylation seemed to have crucial and multifaceted
influences on the adaptation of the ten *Aquilegia* species. We wondered whether differential epigenetic
modifications are dependent on genetic variability. Among the 588,659 CG sites examined, 224,222
(38.09%) carried CG-loss variation(s). We then illustrated epigenetic variability for the mutation-carrying
and non-variant CG sites, respectively. As shown in the Figure 4, different levels of genetic-epigenetic
association are observed in the two types of CG sites. For example, the variation-carrying CG site
conveys information that highly reassembles their genetic background, with the overall methylation
pattern showing highly conservativity at the intra-specific level but exhibiting obvious divergence across
the ten columbine species (**Figure 4a**). In contrast, CG methylation divergence at the non-variant sites
varies with stronger variability at both the intra- and inter-specific levels (**Figure 4b**). By examining the
correlation of genetic variability and cytosine methylation, we find that CG methylation divergence at
variation-carrying CG-site is largely attributable to the CG-loss variations (**Figure 4c**). In particular, 75% of
the CG-loss variations occurring at the most highly variable CG-methylated dinucleotides can explain at
least 75% of the total epigenetic variability *per se*. Nevertheless, there is still a considerable proportion
of epigenetic variability that can not be sufficiently explained by the variant-CG site (**Figure 4d**).
Methylation at non-variant CG site could still reflect population structure, such as the neutralized
methylation levels in the hybrid possibly arising from heterozygosity, though with increased noise and
slightly compromised accuracy (**Figure 4b**). For instance, CG methylation pattern in the hybrid is
neutralized possibly due to heterozygosity. In addition, one *A. japonica* accession (A. japonica9, **Table**
S1) inhabiting low altitude environmental niche shows differential methylation at some CG site
compared to the other con-specific accessions (**Figure 4a** and **b**).

It is arguable that non-variant CG site possesses species-specific methylation pattern that is either

associated with adjacent CG-loss variations or some other non-CG-loss variations. Therefore, we have

attempted to identify driving mutations for each of the 1,229 DMRs between the *A. japonica* and *A.*

oxysepala (**Supplementary Information**). Our results reveal that only 568 out of the 1,229 (46.2%) DMRs

are significantly associated with point mutation inside or around a 500 bp upstream/downstream

window under the least stringent *p* value threshold (5×10^{-5}), indicating that the epigenetic changes are

only partially dependent on or consequences of *cis*-genetic mutations (**Figure 4e**). Notable, we observed

significant associations between differential CG methylation and positive selection at some specific

genes. In most inter-lineage comparisons, DMGs are significantly more prone to be under positive and

purifying selection than non-DMGs (**Table 2**), suggesting that epigenetic modifications could probably

assist selection pressures in shaping genotypes. In contrast, DMGs are significantly less prone to be

under purifying selection (**Table 2**).

Discussion

Genetic-driven mechanisms underpinning the diversification of *Aquilegia* species

Elucidating the extrinsic and intrinsic factors underpinning species diversification is crucial to

understanding the evolution and persistence of biodiversity (Simpson & Olson, 1953; Schluter, 2000;

Losos, 2010). The genus *Aquilegia* provides an ideal system to address how the diverse evolutionary

mechanisms promoted rapid adaptive radiation (Fior *et al.*, 2013; Filiault *et al.*, 2018). Yet, various

external conditions related to ecological opportunities, such as shifts in pollinator and habitat, are

supposed to facilitate the evolution of reproductive isolation (Kramer & Hodges, 2010; Li *et al.*, 2014),

genetic basis underlying the rapid diversification of *Aquilegia* species remained largely unclear. In

particular, it is still unclear whether the adaptive radiations of North American and European lineages

are caused by distinct extrinsic triggers or determined by lineage-specific genetic and epigenetic

architectures. In this study, we surveyed the genomes of ten worldwide columbine species to address

genetic basis underpinning the rapid species diversification. Broadly consistent with previously inferred

phylogenies (Bastida & Herrera, 2010; Fior *et al.*, 2013; Li *et al.*, 2014; Li *et al.*, 2019), the ten columbine

species formed three highly divergent lineages corresponding to their geographic origins, with the Asian

species being separated into three distinct clades but the phylogenetic relationships of European and

North American species varying obviously among the seven chromosomes. These genomic features

allow us to address whether specific genomic architectures contributed to the repeated adaptive

speciation in the genus *Aquilegia*.

It has been proposed that if intrinsic factor is the potential determinant promoted adaptive species,

one would expect to identify specific genetic architectures in the diversified lineages (Brawand *et al.*,

2014; Machado-Schiaffino *et al.*, 2014; Lamichhaney *et al.*, 2015). In Darwin's finches, for example,

polyphyletic topology is observed as a general pattern in the 14 morphologically distinct species.

Phenotypic diversity of the beak shape is mainly determined by natural selection acting on the *ALX1*

gene during the ecological specialization process (Lamichhaney *et al.*, 2015). A similar phenomenon is

observed in the East African cichlid fish wherein the radiating lineages are more dynamic in terms of

gene content and transcriptomic landscape compared to their non-radiating relatives (Brawand *et al.*,

2014; Machado-Schiaffino *et al.*, 2014). In *Aquilegia*, our previous studies have demonstrated that

natural selection and genetic drift together resulted in the rapid evolution of reproductive isolation

between the *A. japonica* and *A. oxysepala* (Li *et al.*, 2014; Li *et al.*, 2019). Here we further revealed that

candidate genes involved in the adaptive speciation are functionally enriched in categories related to

cell reproduction (e.g., telomere maintenance), stress tolerance (e.g., response to wounding) and basic

cellular activities. More importantly, a considerable proportion of the candidate genes (e.g., CCVs and

d_N/d_S) and enrichment pathways share at the intra- and inter-specific comparisons, particularly those of

related to cell reproduction, stress tolerance, and plant growth. Given that these functionally important

pathways are potentially correlated with adaptive traits, we propose that specific genetic architectures

are the intrinsic factor conferred adaptability to the *Aquilegia* species to respond to diverse external

conditions.

It should be noted that while both the North American and European columbines originate from

ancestral Asian species, shifts in pollinator and ecological habitat are supposed to drive the adaptive

radiation of the two lineages, respectively (Bastida & Herrera, 2010; Fior *et al.*, 2013; Filiault *et al.*,

2018). One hypothesis is that the ancestral Asian species carry lineage-specific genetic architectures,

which offered the North American and European species adaptabilities to cope with pollinator and

ecological specializations. With this reasoning, lineage-specific genetic variants identified in the North

American and European columbines should be functionally correlated with flower development and

ecological specialization. Alternatively, independent adaptive radiations of the North American and

European lineages are mainly caused by different ecological opportunities in respective geographic

regions. Phenotypic diversity of the North American and European species is potentially determined by

natural selection acting on similar genetic architectures. In our study, a considerable of the candidate

genes and their enrichment pathways share between the two lineages and Asian species, suggesting the

possibility that similar genetic architectures contributed to the adaptive radiations of the North

American and European columbine species. Nevertheless, we find that majority of the functionally
important pathways show complementary enrichment pattern in the inter-lineage comparisons, clearly
indicating that lineage-specific differences play crucial roles in the adaptive radiations of the North
American and European species. More importantly, a part of the identified lineage-specific genetic
differences (e.g., CCVs and d_N/d_S) are associated with adaptive divergence of the three lineages. For
example, four tandem duplicates of the AGAMOUS MADS-box gene exhibit high genetic divergence
among the North American, European and Asian lineages, particularly that three copies harbor multiple
lineage-specific non-synonymous substitutions. The MADS-box AGAMOUS gene plays critical roles in the
determination of reproductive floral oranges (e.g., carpel and stamen), floral meristem determinacy and
flowering transition (Kramer *et al.*, 2004; Tapia-Lopez *et al.*, 2008; Dreni & Kater, 2014). It indicates that
the four AGAMOUS genes are possibly associated with the floral development and reproductive
isolation of the columbine species. Taken together, our findings indicate that both the lineage-specific
and shared genetic architectures are the intrinsic factor interacts with extrinsic trigger to determine the
adaptive radiations of the North American and European species.
406
Evolutionary potential of cytosine methylation in the adaptation of *Aquilegia* species
The role of epigenetic modification in the long-term evolutionary process has long been debated
(Bossdorf *et al.*, 2008; Diez *et al.*, 2014; Verhoeven *et al.*, 2016). It has been noted that epigenetic
variations are usually under the genetic control and vary rapidly as a result of environmental induction
and stochastic epimutation (Richards *et al.*, 2010; Richards *et al.*, 2017). Nevertheless, it has been
recently recognized that some epigenetic variants can persist over generations and be highly correlated
with adaptive phenotypes (Verhoeven *et al.*, 2016). As illustrated in Arabidopsis, changes in cytosine
methylation can produce meiotically stable epialleles, which could eventually lead to phenotypic
diversity in the absence of genetic variations (Becker *et al.*, 2011; Schmitz *et al.*, 2011; Cortijo *et al.*,
2014). In this study, we aimed to assess whether the epigenetic modification also contributed the
adaptive speciation of the *Aquilegia* species. Consistent with the genomic features detailed above,
clearly high divergence of cytosine methylation was observed among the Asian, European and North
American lineages, particularly that differential cytosine methylation is not only found among the seven
chromosomes but also evident in the gene body and CG island regions. These observations indicate that
cytosine methylation may have contributed to the adaptability of *Aquilegia* species. Between the *A.*
japonica and *A. oxysepala*, for example, candidate genes identified at the genetic level are functionally
related to cell production, stress tolerance and basic cellular activities. At the cytosine methylation level,

the identified DMGs show significant enrichment in categories related to diverse important phenotypic

traits, such as photosynthesis, plant growth and development. Given that adaptive speciation of the two

species is mainly caused by ecological specialization (Li *et al.*, 2014; Li *et al.*, 2019), we then propose that

epigenetic modification is also a complementary mechanism facilitated phenotypic divergence between

the two species. A similar phenomenon was also observed from the inter-lineage comparison where

differences in cytosine methylation are associated with adaptive phenotypes such as the plant growth

and stress tolerance. These findings indicate that both the genetic and epigenetic architectures interact

to respond to diverse extrinsic conditions.

We noted that some DMGs and enriched pathways are commonly found from the genetic and

epigenetic comparisons, especially for those that related to plant growth and stress tolerance. We then

asked whether the epigenetic mechanism works independently or largely under the control of genetic

variability. Intuitively, genetic variations are important drivers of epigenetic variability. Many studies

based on human and mouse have shown genetic variations can not only directly control CG methylation

patterns (Drong *et al.*, 2013; Smith *et al.*, 2014), but also manipulate long-range CG methylation at

specific sites to further influence phenotypes, where CG methylation serves as a mediator (Fisher *et al.*,

2018; Li *et al.*, 2018). However, precedent studies on explaining, in a time-span of evolutionary history,

whether CG methylation functions independent of genetic variations are few to our knowledge. By

analyzing the associations between genetic and epigenetic variability, we find that while the CG-loss

variation can directly lead to the depletion in CG methylation level, there still exist a considerable of

DMRs not attributable to *cis*-variation. Though potential non-linear additivity and possible existence of

distal driver mutations beyond the 1kb window of DMRs may introduce biases, our findings at least

suggest that the epigenetic variability also possesses autonomy to some extent. Furthermore, since

gene body CG methylation in plants generally stabilizes gene expression and is positively correlated with

gene expression level (Takuno & Gaut, 2013; Bewick *et al.*, 2016; Bewick & Schmitz, 2017; Zilberman,

2017), differential methylation in our study is indicative of the amount of eventual gene products. Based

on these attributes together with the significant association between differential methylation (DMGs)

and positive selection (d_N/d_S), we propose that the intrinsic epigenetic variability and the extrinsic

selection pressures are intertwined, where differential methylation could assist selection pressures.

Conclusions

Elucidating how the intrinsic and extrinsic mechanisms promoted species diversification is critical to

understanding the evolution and persistence of biodiversity. Here, we took the advantage of the model

system *Aquilegia* to address genetic and epigenetic bases underlying the rapid species radiation. Our
results demonstrated that intrinsic factors related to specific genetic and epigenetic architectures
conferred adaptability to the *Aquilegia* species in response to diverse external conditions, and that the
epigenetic modification is a complementary mechanism that assists the selection pressure to shape the
phenotypic diversity. Our study provides a genome-wide view of how the high adaptability of *Aquilegia*
species have evolved in the North American, European and Asian columbine species.

Author Contributions

464   L.F.L. conceived this project. T.L. and M.R.L. developed statistical analysis pipeline. T.L., M.R.L., N.D.,
465   Z.H.W., L.Z.L., and X.G. carried out experiments and analyzed the data. T.L., M.R.L., N.D., Z.H.W., L.Z.L.,
466   X.G., and L.F.L. interpreted the data and participated in discussion and wrote the manuscript.

Acknowledgments

469   We thank Aköz Gökçe for her constructive comments on our manuscript. This work was financially
470   supported by National Natural Science Foundation of China (31670382), Shanghai Pujiang Program
471   (19PJ1401500), Start-up funding at Fudan University (JIH1322105) and the Department of Science and
472   Technology of Jilin Province (20190201299JC).

References

Aguirre-Portoles C, Bird AW, Hyman A, Canamero M, de Castro IP, Malunnbres M. 2012. Tpx2 controls
spindle integrity, genome stability, and tumor development. *Cancer Research* **72**(6): 1518-1528.
Alcántara JM, Bastida JM, Rey PJ. 2010. Linking divergent selection on vegetative traits to
environmental variation and phenotypic diversification in the Iberian columbines (*Aquilegia*).
Journal of Evolutionary Biology **23**(6): 1218-1233.
Alexa A, Rahnenfuhrer J, Lengauer T. 2006. Improved scoring of functional groups from gene expression
data by decorrelating GO graph structure. *Bioinformatics* **22**(13): 1600-1607.
Alexander DH, Novembre JK. 2009. Fast model-based estimation of ancestry in unrelated individuals.
Genome Research **19**(9): 1655-1664.
Bastida JM, Herrera CM. 2010. Extended phylogeny of *Aquilegia*: the biogeographical and ecological
patterns of two simultaneous but contrasting radiations. *Plant Systematics and Evolution* **284**(3-
4): 171-185.
Dardou P, Mariette J, Escudie F, Djemiel C, Klopp C. 2014. Jvenn: an interactive venn diagram viewer.
BMC Bioinformatics **51**(1):293.
Becker C, Hagmann J, Jonas M, Daniel K, Oliver S, Karsten B, Detlef W. 2011. Spontaneous epigenetic
variation in the *Arabidopsis thaliana* methylome. *Nature* **480**(7376): 245.
Berner D, Salzburger W. 2015. The genomics of organismal diversification illuminated by adaptive
radiations. *Trends in Genetics* **31**(9): 491-499.

Bewick AJ, Ji L, Niederhuth CE, Willing EM, Hofmeister BT, Shi X, Wang L, Lu Z, Rohr NA, Hartwig B.
2016. On the origin and evolutionary consequences of gene body DNA methylation. *Proceedings*
of the National Academy of Sciences of the United States of America **113**(32): 9111-9116.
Bewick AJ, Schmitz RJ. 2017. Gene body DNA methylation in plants. *Current Opinion in Plant Biology* **36**:
103-110.
Bossdorf O, Richards CL, Massimo P. 2008. Epigenetics for ecologists. *Ecology Letters* **11**(2): 106-115.
Brawand D, Wagner CE, Li YI, Malinsky M, Keller I, Fan SH, Simakov O, Ng AY, Lim ZW, Bezault E, et al.
2014. The genomic substrate for adaptive radiation in African cichlid fish. *Nature* **513**(7518):
375-381.
Carlquist, Motley TJ. 2003. Tarweeds and Silverswords, Evolution of the Madiinae (*Asteraceae*).
Economic Botany **58**(1): 123-124.
Cingolani P, Platts A, Le LW, Coon M, Nguyen T, Luan W, Land SJ, Lu X, Ruden DM. 2012. A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly* **6**(2): 80-
92.
Cortijo S, Wardenaar R, Colome-Tatche M, Gilly A, Etcheverry M, Labadie K, Caillieux E, Hospital F,
Aury JM, Wincker P, et al. 2014. Mapping the epigenetic basis of complex traits. *Science*
343(6175): 1145-1148.
Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT,
Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* **27**(15): 2156-2158.
Diez CM, Roessler K, Gaut BS. 2014. Epigenetics and plant genome evolution. *Current Opinion in Plant*
Biology **18**(1): 1-8.
Dreni L, Kater MM. 2014. MADS reloaded: evolution of the AGAMOUS subfamily genes. *New Phytologist*
201(3): 717-732.
Drong AW, George N, Hedman AK, Eshwar M, Elin G, Small KS, So-Youn S, Bell JT, Fredrik K, Nicole S.
2013. The presence of methylation quantitative trait loci indicates a direct genetic influence on
the level of DNA methylation in adipose tissue. *Plos One* **8**(2): e55923.
Filiault DL, Ballerini ES, Mandakova T, Akoz G, Derieg NJ, Schmutz J, Jenkins J, Grimwood J, Shu S,
Hayes RD, et al. 2018. The *Aquilegia* genome provides insight into adaptive radiation and
reveals an extraordinarily polymorphic chromosome with a unique history. *Elife* **7**.
Fior S, Mingai L, Bengt O, Roberto V, Hodges SA, Lino O, Claudio V. 2013. Spatiotemporal
reconstruction of the *Aquilegia* rapid radiation through next-generation sequencing of rapidly
evolving cpDNA regions. *New Phytologist* **198**(2): 579-592.
Fisher VA, Wang L, Deng X, Sarnowski C, Cupples LA, Liu CT. 2018. Do changes in DNA methylation
mediate or interact with SNP variation? A pharmacoepigenetic analysis. *BMC Genetics* **19**(Suppl
1): 70.
Ford AG, Rüber L, Newton J, Dasmahapatra KK, Balarin JD, Bruun K, Day JJ. 2016. Niche divergence
facilitated by fine-scale ecological partitioning in a recent cichlid fish adaptive radiation.
Evolution **70**(12): 2718-2735.
Fukui K, Nakagawa N, Kitamura Y, Nishida Y, Masui R, Kuramitsu S. 2008. Crystal structure of MutS2
endonuclease domain and the mechanism of homologous recombination suppression. *Journal*
of Biological Chemistry **283**(48): 33417-33427.
Garrido JL, Alcántara JM, Rey PJ, Medrano M, Guitián J, Castellanos MC, Bastida JM, Jaime R, Herrera
CM. 2017. Geographic genetic structure of Iberian columbines (gen. *Aquilegia*). *Plant*
Systematics and Evolution **303**(9): 1145-1160.
Givnish TJ. 1997. Adaptive radiation and molecular systematics: aims and conceptual issues. In: Givnish
TJ, Systma KJ, eds. . *Molecular Evolution and Adaptive Radiation.*: 1-54.
Givnish TJ. 2015. Adaptive radiation versus 'radiation' and 'explosive diversification': why conceptual
distinctions are fundamental to understanding evolution. *New Phytologist* **207**(2): 297-303.

Irisarri I, Singh P, Koblmüller S, Torresdowdall J, Henning F, Franchini P, Fischer C, Lemmon AR,
Lemmon EM, Thallinger GG. 2018. Phylogenomics uncovers early hybridization and adaptive
loci shaping the radiation of Lake Tanganyika cichlid fishes. *Nature Communications* **9**(1): 3159-
3171.
Kramer EM, Hodges SA. 2010. *Aquilegia* as a model system for the evolution and ecology of petals.
Philosophical Transactions Biological Sciences **365**(1539): 477-490.
Kramer EM, Jaramillo MA, Di Stilio VS. 2004. Patterns of gene duplication and functional evolution
during the diversification of the *AGAMOUS* subfamily of MADS box genes in angiosperms.
Genetics **166**(2): 1011-1023.
Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. *Bioinformatics* **27**(11): 1571-1572.
Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for
bigger datasets. *Molecular Biology and Evolution* **33**(7): 1870.
Lamichhaney S, Jonas B, Markus S?Llman A, Khurram M, Manfred G, Alvaro MB, Marta P, Carl-Johan
R, Chao W, Neda Z. 2015. Evolution of Darwin's finches and their beaks revealed by genome
sequencing. *Nature* **518**(7539): 371-375.
Li H, Durbin R. 2009. *Fast and accurate short read alignment with Burrows-Wheeler transform*.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Proc GPD.
2009. The sequence alignment/map format and SAMtools. *Bioinformatics* **25**(16): 2078-2079.
Li L, Wang C, Lu T, Lin S, Hu YQ. 2018. Indirect effect inference and application to GAW20 data. *BMC*
Genetics **19**(Suppl 1): 67.
Li LF, Pang D, Liao QL, Xiao HX. 2011. Genomic and EST microsatellite markers for *Aquilegia* flabellata
and cross-amplification in *A. oxysepala* (Ranunculaceae). *American Journal of Botany* **98**(8): 213-
215.
Li LF, Wang HY, Pang D, Liu Y, Liu B, Xiao HX. 2014. Phenotypic and genetic evidence for ecological
speciation of *Aquilegia japonica* and A. *oxysepala*. *New Phytologist* **204**(4): 1028-1040.
Li MR, Wang HY, Ding N, Lu T, Huang YC, Xiao HX, Liu B, Li LF. 2019. Rapid divergence followed by
adaptation to contrasting ecological niches of two closely related columbine species *Aquilegia*
japonica and A. *oxysepala*. *Genome Bioligy and Evolution* **11**(3): 919-930.
Losos JB. 2010. Adaptive radiation, ecological opportunity, and evolutionary determinism. American
Society of Naturalists E. O. Wilson award address. *American Naturalist* **175**(6): 623-639.
Losos JB, Ricklefs RE. 2009. Adaptation and diversification on islands. *Nature* **457**(7231): 830-836.
Machado-Schiaffino G, Henning F, Meyer A. 2014. Species-specific differences in adaptive phenotypic
plasticity in an ecologically relevant trophic trait: hypertrophic lips in Midas cichlid fishes.
Evolution **68**(7): 2086-2091.
Mckenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,
Gabriel S, Daly M. 2010. The Genome Analysis Toolkit: A map reduce framework for analyzing
next-generation DNA sequencing data. *Genome Research* **20**(9): 1297-1303.
Munz. 1946. Aquilegia: the cultivated and wild columbines. *Gentes Herbarum* **7**: 1-150.
O''Sullivan DE, Johnson KC, Skinner L, Koestler DC, Christensen BC. 2016. Epigenetic and genetic
burden measures are associated with tumor characteristics in invasive breast carcinoma.
Epigenetics **11**(5): 344-353.
Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, David R. 2006. Principal components
analysis corrects for stratification in genome-wide association studies.
Rice P, Longden I, Bleasby A. 2000. EMBOSS: The European molecular biology open software suite.
Trends in Genetics **16**(6): 276-277.

Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, Colome-Tatche M, Durka W, Engelhardt J,
Gaspar B, Gogol-Doring A, et al. 2017. Ecological plant epigenetics: Evidence from model and
non-model species, and the way forward. *Ecology Letters* **20**(12): 1576-1590.
Richards CL, Oliver B, Verhoeven KJF. 2010. Understanding natural epigenetic variation. *New*
Phytologist **187**(3): 562-564.
Schluter D. 1996. Ecological causes of adaptive radiation. *American Naturalist* **148**: S40-S64.
Schluter D. 2000. The Ecology of Adaptive Radiation. *Heredity* **86**(6): 749-750.
Schmitz RJ, Schultz MD, Lewsey MG, O'Malley RC, Urich MA, Ondrej L, Schork NJ, Ecker JR. 2011.
Transgenerational epigenetic instability is a source of novel methylation variants. *Science*
334(6054): 369-373.
Sharma B, Yant L, Hodges SA, Kramer EM. 2014. Understanding the development and evolution of
novel floral form in *Aquilegia*. *Current Opinion in Plant Biology* **17**(1): 22-27.
Simpson EH. 1949. Measurement of Diversity. *Nature* **163**(4148): 688-688.
Simpson GG, Olson EC. 1953. The major features of evolution. *Quarterly Review of Biology* **8**(1): 87-88.
Smith AK, Kilaru V, Kocak M, Almli LM, Mercer KB, Ressler KJ, Tylavsky FA, Conneely KN. 2014.
Methylation quantitative trait loci (meQTLs) are consistently detected across ancestry,
developmental stage, and tissue type. *Bmc Genomics* **15**(1): 145.
Takuno S, Gaut BS. 2013. Gene body methylation is conserved between plant orthologs and is of
evolutionary consequence. *Proceedings of the National Academy of Sciences of the United*
States of America **110**(5): 1797-1802.
Tang LL, Yu Q, Sun JF, Huang SQ. 2007. Floral traits and isolation of three sympatric *Aquilegia* species in
the Qinling Mountains, China. *Plant Systematics and Evolution* **267**(1/4): 121-128.
Tapia-Lopez R, Garcia-Ponce B, Dubrovsky JG, Garay-Arroyo A, Perez-Ruiz RV, Kim SH, Acevedo F,
Pelaz S, Alvarez-Buylla ER. 2008. An *AGAMOUS*-related MADS-box gene, XAL1 (*AGL12*),
regulates root meristem cell proliferation and flowering transition in *Arabidopsis*. *Plant*
Physiology **146**(3): 1182-1192.
Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. *Brief Bioinformatics* **14**(2): 178-192.
Verhoeven KJ, Vonholdt BM, Sork VL. 2016. Epigenetics in ecology and evolution: what we know
and what we need to know. *Molecular Ecology* **25**(8): 1631-1638.
Wagner CE, Harmon LJ, Seehausen O. 2012. Ecological opportunity and sexual selection together
predict adaptive radiation. *Nature* **487**(7407): 366-U124.
Whittall JB, Hodges SA. 2007. Pollinator shifts drive increasingly long nectar spurs in columbine flowers.
Nature **447**(7145): 706-709.
Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*
24(8): 1586-1591.
Zhang H, Dawe RK. 2011. Mechanisms of plant spindle formation. *Chromosome Research* **19**(3): 335-
344.
Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. 2012. A high-performance computing
toolset for relatedness and principal component analysis of SNP data. *Bioinformatics* **28**(24):
3326-3328.
Zilberman D. 2017. An evolutionary case for functional gene body methylation in plants and animals.
Genome Biology **18**(1): 87.
630
631

Figure legend

Figure 1. **Phylogenetic relationship and population structure of the ten worldwide *Aquilegia* species**.

(a) Phylogenetic tree of the 36 accessions constructed by neighbor joining algorithm based on whole-

genome SNPs. (b) PCA reveals genetic similarity within each of the three lineages and genetic disparity

between lineages based on 15,988 LD-pruned SNPs. Curves for each lineage denote 99% probability

ellipses estimated from distribution of the first two principal components. (c) Population admixture of

the 36 *Aquilegia* accessions. Most accessions exhibit almost pure ancestry composition matching the

presupposed evolutionary history of the corresponding species.

Figure 2. **Functional enrichment of genes harboring highly impactful CCVs and DMGs.** CCV-containing

genes specific to either of the two lineages/species being compared were merged to construct a target

gene set. Ratio denotes proportion of CCV-containing genes or DMGs in the corresponding gene set of

interest. Absence of dot indicates no significant enrichment.

Figure 3. **Patterns of cytosine methylation for the ten worldwide *Aquilegia* species.** (a) Genome-wide

CG methylation level distributions of 36 accessions. (b) MDI illustrates chromosome-level CG

methylation similarity. *Aquilegia viridiflora* was used as the reference. (c) CG methylation profiling in

genic region across the four *Aquilegia* groups. Each row represents one genic region starting at 5kb

upstream of its TSS and terminating at 5kb downstream of its TES, sorted by mean methylation level of

all analyzed CG sites. Length of gene body was normalized. (d) CG methylation profiling in and around

CG island.

Figure 4. **Association between CG-loss variations and epigenetic variability**. (a) Top 3,000 most variable

CG sites containing CG-loss variations. (b) top 3,000 most variable non-variant CG sites across 36

accessions show clade-specific methylation patterns. (c) Linear regression demonstrates that CG-loss

variations explain a large proportion of CG methylation variation. Composition of each category is

summarized in (d) with regard to whether each CG site contains a CG-loss variation. Epigenetic

variability was determined by standard deviation in methylation β value across all 36 accessions. CG

sites with top 10,000, 10,001-50,000 and 50,001-150,000 largest standard deviation were ordinally

labelled as possessing "very high", "high" and "moderate" variability respectively. The rest CG sites were

labelled as possessing "low" variability. (e) Association test shows most DMRs are independent of *cis*-

acting SNPs.

Supplementary file

Figure S1. Per-chromosome phylogenetic trees reconstructed using neighbor-joining method.
Polymorphisms detected on each chromosome were retrieved separately to infer the phylogeny.

Figure S2. Distribution of nucleotide diversity (π) at the whole-genome level and the per-chromosome
level. Nucleotide diversity was estimated for each lineage pooling corresponding species, as well as for
A. japonica and *A. oxysepala*.

Figure S3. Distribution of inter-lineage and inter-specific genetic divergence (Fst) at the whole-genome
level and the per-chromosome level.

Figure S4. (a) Overlapping of HGDRs in three inter-lineage comparisons. 148 regions with 5% highest
genetic divergence were considered HGDRs in each comparison. 73 regions were shared by two
comparisons and 23 regions showed high genetic divergence in all inter-lineage comparisons. (b)
Overlapping of LDNRs in three lineages. 148 regions with 5% lowest nucleotide diversity were
considered LDNRs in each lineage. 29 regions had low genetic variability in two lineages and 57 regions
had low genetic variability in all three lineages.

Figure S5. Overlapping of genes harboring CCVs (a), or indicating strong positive (b) and purifying (c)
selection at inter-lineage/species comparisons. Genes having a high d_N/d_S (among the highest 5% or a
5% highest d_N when $d_S = 0$) were considered to be under strong positive selection. Genes having a low
d_N/d_S (among the lowest 5% or a 5% highest d_S when $d_N = 0$) were considered to be under strong
purifying selection.

Figure S6. Functional enrichment of genes indicating strong positive selection or purifying selection.
Functional enrichment analysis was performed for each inter-lineage/species comparison.

Figure S7. PCA illustrates three distinct clusters corresponding to the three lineages. Asian species
further demonstrated higher inter-specific divergence than the American and the European species. PCA
was performed based on 588,659 loci with sufficiently high sequencing quality.

Figure S8. Illustration of differential methylation in two photosynthesis genes. CG methylation pattern
of two genes, Aqcoe7G230600 photosystem I *PsaA*/*PsaB* (a) and Aqcoe7G231300 *CemA* (b) in *A.*
japonica and *A. oxysepala* throughout the gene body region. Red bars indicate methylation level (0-100)
at CG sites. Genomic coordinates on the chromosome 7 are annotated.

Figure S9. Overlapping of DMGs identified in inter-lineage/species comparisons. A considerable
proportion (51.3%) of these DMGs were shared by two or more inter-lineage/species comparisons.

Figure S10. Venn analyses of the candidate genes under relaxed selection (CCV-carrying), positive
selection (high d_N/d_S value), purifying selection (low d_N/d_S value) and differential methylated (DMG).

Each subpanel indicates the comparison between the *A. japonica* and *A. oxysepala* (a), *A. japonica* and

North American (b), *A. japonica* and European (c), *A. oxysepala* and North American (d), *A. oxysepala*

and European (e), North American and European (f).

Table 1. Information of the high-impact conservative clade-specific variants (CCVs) in the cell
reproduction related genes.

Gene	Variant-carrying	Reference	Chromosome	Position	Reference allele	Variant	Annotation	Gene function
Aqcoe1G273400	Asian	American	Chr1	18994915	GAA	GAAA	frameshift	DNA mismatch repair protein *MutS2*
Aqcoe2G151500	European	American	Chr2	15305837	A	G	splicing	PIF1-like helicase
	European	American		15307442	A	C	stop gain	
	European	American		15309865	AATATATAT	AATATATATAT	frameshift	
	European	Asian		15307442	A	C	stop gain	
	European	Asian		15309865	AATATATAT	AATATATATAT	frameshift	
	A. oxysepala	*A. japonica*		15305837	A	G	splicing	
	A. oxysepala	*A. japonica*		15309267	AT	A	frameshift	
Aqcoe2G177700	European	American	Chr2	21794397	TATGCACCAAAGGTATCACGATGC	TATGC	frameshift	PIF1-like helicase
	European	American		21794979	TT	TTGT	frameshift	
	European	Asian		21794397	TATGCACCAAAGGTATCACGATGC	TATGC	frameshift	
	A. oxysepala	*A. japonica*		21795089	CA	C	frameshift	
Aqcoe6G208600	European	American	Chr6	15364081	A	ATCTCTTCG	frameshift	PIF1-like helicase
	European	Asian		15364081	A	ATCTCTTCG	frameshift	
	A. japonica	*A. oxysepala*		15364330	TAA	TA	frameshift	
Aqcoe6G253800	European	American	Chr6	22789898	C	T	stop gain	DNA helicase
	European	American		22790012	G	A	splicing	
	European	Asian		22789898	C	T	stop gain	
	A. japonica	*A. oxysepala*		22790012	G	A	splicing	
Aqcoe2G276600	Asian	American	Chr2	33314422	AGGGGG	AGGGGGG	frameshift	DNA mismatch repair protein *Msh6*
Aqcoe6G160300	*A. japonica*	*A. oxysepala*	Chr6	9414625	G	A	stop gain	*TPX2*
Aqcoe7G062500	*A. oxysepala*	*A. japonica*	Chr7	3789055	G	A	stop gain	cell cycle regulated microtubule associated protein

703
704
705

706   **Table 2. Correlation test between the differential methylation and d_N/d_S value for the overall genes.**

Type of selection	Differential methylation	Jap-Oxy[*]	Jap-Ame	Jap-Eur	Oxy-Ame	Oxy-Eur	Ame-Eur
Positive selection	DMG	7.2%	7.3%	11.9%	6.7%	8.4%	8.9%
	non-DMG	4.4%	5.0%	5.6%	4.7%	5.4%	5.4%
	p value	0.11	7.3e-02	3.9e-05	6.7e-02	1.8e-02	2.8e-04
Purifying selection	DMG	3.1%	1.8%	2.4%	2.3%	2.0%	1.9%
	non-DMG	4.3%	4.3%	4.7%	4.9%	5.1%	4.0%
	p value	0.53	3.2e-02	9.1e-02	1.3e-02	8.4e-03	1.0e-02

708   *, Jap, *A. japonica*; Oxy, *A. oxysepala*; Ame, American species; Eur, European species. Each percentage
709   represents the proportion of genes belonging to either DMGs or non-DMGs compared between the two
710   corresponding clades that are under corresponding or higher strength of positive selection. The *p* values
711   were obtained from Chi-square tests.

Whole genome   **Chr1**   **Chr2**   **Chr3**

Chr4   **Chr5**   **Chr6**   **Chr7**

American vs. European
American vs. Asian
European vs. Asian
A. japonica vs. *A. oxysepala*

Genetic divergence