Systematic profiling of full-length immunoglobulin and T-cell receptor

repertoire diversity in rhesus macaque through long read transcriptome

sequencing

Running title: Long read transcriptome sequencing of rhesus Ig and TCR

Hayden N. Brochu[*,†], Elizabeth Tseng[‡], Elise Smith[§], Matthew J. Thomas[§,¶], Aiden

Jones[*,‖], Kayleigh Diveley[*,‖], Lynn Law[§,¶], Scott G. Hansen[#], Louis J. Picker[#], Michael

Gale, Jr.[§,¶,**], and Xinxia Peng[*,†,††]

*Department of Molecular Biomedical Sciences, North Carolina State University College

of Veterinary Medicine, Raleigh, NC 27607, USA, †Bioinformatics Graduate Program,

North Carolina State University, Raleigh, NC 27695, USA, ‡Pacific Biosciences, Menlo

Park, CA 94025, USA, §Department of Immunology, University of Washington, Seattle,

WA, USA, ¶Center for Innate Immunity and Immune Diseases, University of Washington,

WA, USA, ‖Genetics Graduate Program, North Carolina State University, Raleigh, NC

27695, #Vaccine and Gene Therapy Institute, Oregon Health & Science University,

Beaverton, OR 97006, USA, **Washington National Primate Research Center, University

of Washington, Seattle, WA, USA, ††Bioinformatics Research Center, North Carolina State

University, Raleigh, NC 27695, USA

Address correspondence to Dr. Xinxia Peng, 919-515-4481, xpeng5@ncsu.edu

Abstract

The diversity of immunoglobulin (Ig) and T-cell receptor (TCR) repertoires is a focal point of immunological studies. Rhesus macaques are key for modeling human immune responses, placing critical importance on the accurate annotation and quantification of their Ig and TCR repertoires. However, due to incomplete reference resources, the coverage and accuracy of the traditional targeted amplification strategies for profiling rhesus Ig and TCR repertoires are largely unknown. Here, using long read sequencing, we sequenced four Indian-origin rhesus macaque tissues and obtained high quality, full-length sequences for over 6,000 unique Ig and TCR transcripts, without the need for sequence assembly. We constructed the first complete reference set for the constant regions of all known isotypes and chain types of rhesus Ig and TCR repertoires. We show that sequence diversity exists across the entire variable regions of rhesus Ig and TCR transcripts. Consequently, existing strategies using targeted amplification of rearranged variable regions comprised of V(D)J gene segments miss a significant fraction (27% to 53% and 42% to 49%) of rhesus Ig/TCR diversity. To overcome these limitations, we designed new rhesus-specific assays that remove the need for primers conventionally targeting variable regions and allow single cell-level Ig and TCR repertoire analysis. Our improved approach will enable future studies to fully capture rhesus Ig and TCR repertoire diversity and is applicable for improving annotations in any model organism.

Introduction

Rhesus macaque (*Macaca mulatta*) is one of the most commonly used and best-studied non-human primate (NHP) animal models. NHPs are key to studying human biology and human diseases due to their close phylogenetic relationship and similar physiology to humans [1-4]. They are frequently used for vaccine development [5] and to model infection with human pathogens, such as Mycobacterium tuberculosis [6, 7], HIV [8, 9], influenza A virus [10], and Zika virus [11], among others [12, 13]. Developing complete and accurate NHP genomic resources, especially for the immune system, is imperative for efficient translational interpretation [14, 15].

Critical for mounting adaptive immune responses, immunoglobulin (Ig) and T-cell receptor (TCR) repertoires house an enormous amount of diversity responsible for recognizing a near limitless array of antigens presented through environmental exposures. Ig and TCR have two domains: a constant region and a variable region. The repertoire diversity is mostly present in the variable region domain, which is comprised of a Variable (V), Joining (J), and in some cases a Diversity (D) gene segment (Fig. 1). Duplicated copies of these gene segments cluster into several large loci in the genome. As a result, correct assembly of their respective loci with highly repetitive sequences is a major technical challenge, one that has yet to be fully resolved in rhesus macaque despite significant improvements [16] and the recent release of a new genome assembly, rheMac10 (GCA_003339765.3). As demonstrated by a recent vaccine-related study in rhesus macaque [17], the complexity of these regions requires much longer reads. Standard databases (e.g. the international ImMunoGeneTics information system or IMGT) for these diverse sequences also remain fairly meager relative to their human counterparts, although

there are new tools developed to address these gaps [18]. Since the design of available

rhesus-specific assays for profiling Ig and TCR diversity has heavily relied on these limited

rhesus reference resources, the coverage and the accuracy of these assays still require

unbiased assessment and potentially improved approaches might be necessary. We begin

by briefly summarizing the complexity of these immune repertoires in general and how

this complexity has constrained the development of resources and assays for rhesus

macaque.

In individual B and T cells, different gene segments of both variable and constant

regions are combined at the DNA level to encode distinctive functional Ig and TCR genes

through a process known as somatic recombination (Fig. 1) (reviewed in [19-21]). Somatic

recombination accounts for the majority of the diversity within the variable region domain,

with the number of unique Ig and TCR variable region domains estimated to be on the

order of 10^{13} and 10^{18}, respectively [22]. To further increase the repertoire diversity, within

each cell a fully assembled Ig molecule is composed of a unique pairing of a heavy chain

(IGH) and a light chain: either kappa (IGK) or lambda (IGL) (Fig. 1). Similarly, a TCR is

composed of the pairing of an alpha (TCRA) and beta (TCRB) chain or, less commonly, a

delta (TCRD) and gamma (TCRG) chain (Fig. 1). The process of somatic hypermutation

(SHM) further expands the diversity of Ig molecules, where mutations are introduced to

the variable region domain of recombined DNA at rates roughly one million times higher

than background rates observed in other genes [23].

The constant region domain determines various chain types of Ig and TCR [22, 24].

Within each chain type, these variable gene segments are classified into families (e.g. Ig

heavy chain variable gene family 4 or IGHV4). Each gene family includes one or more

unique gene segments (e.g. IGHV4-1) that share high degrees of similarity. In B cells, the

process by which the variable region domain is juxtaposed from the originally utilized

heavy (H) chain constant region gene (usually IGHM) to a different constant region domain

is referred to as class-switch recombination (CSR), a process essential to B lymphocyte

differentiation. In addition to conferring the non-antigen binding functions of antibody

molecules, the heavy chain constant region of secreted Ig differs from Ig bound to the

surface of B cells (the B cell receptor for antigen) based on alternative splicing at the 3'

end.

96   The genetic diversity of the Ig and TCR loci present a unique challenge for accurate

measurement. Traditionally, these immune repertoires are targeted for amplification either

by multiplex polymerase chain reaction (MPCR) [25], RNA capture [26], or 5' rapid

amplification of cDNA ends (5' RACE) [27]. For rhesus macaque, many Ig repertoire

sequencing efforts use a MPCR approach [28, 29], while some employ 5' RACE [30].

Typically, such PCR-based approaches are designed for individually sorted B or T cells

[28] and facilitate cloning efforts [31]. A more recent rhesus-specific MPCR design aimed

to expand coverage of the Ig repertoire [32]. There were also attempts to improve rhesus

V and J germline gene annotation using 5' RACE sequencing (RACE-seq) [18, 33]. The

recently developed IgDiscover tool [18] now makes it possible to leverage germline

databases of related species to improve those of model organisms, for example using

human germline databases to study Indian- and Chinese-origin rhesus macaques. Authors

from the same lab also developed a strategy for targeting the 5' UTR of V genes, often

conserved among these gene clusters, thus reducing the number of primers needed to target

the variable region in human [34]. However, rhesus MPCR amplification systems are

inherently biased to the V and J gene segments they target because the primers are designed

based on the consensus of a limited number of reference sequences [35].

RACE-seq has the advantage of only targeting the constant region, where primer

design is significantly easier given the low sequence variability. 5' RACE in addition to

MPCR have also been modified to incorporate unique molecular identifiers (UMIs) for

repertoire sequencing [36, 37], mitigating issues arising from PCR bias [38]. However,

such protocols are optimized for applications in human and mouse [39] and have not yet

been applied for Ig/TCR analysis in rhesus macaques. Moreover, even with MPCR or 5'

RACE it is still difficult to capture complete Ig mRNA transcripts with the commonly

available sequencing instruments like the 2 x 250 bp HiSeq system and the 2 x 300 bp

MiSeq system, in part due to the large length variability of recombined molecules [40] and

the potential for non-canonical recombination events [41]. Library preparation techniques

have recently been improved to support this capability but only in human [34]. Despite

these technical advances, it is still unclear how well these strategies perform in rhesus

macaque, where the relatively limited resources preclude similar benchmarking efforts

previously done for human Ig repertoires [42] and TCR repertoires [43].

The technical robustness and unique advantages of scRNA-seq [44, 45] have

revolutionized the analysis of human immune systems [46-50], resulting in an increasing

number of studies for single cell-based Ig and TCR repertoire sequencing in human [51-

54]. Single cell protocols target repertoire sequences similarly to RACE-seq by only

targeting the constant regions of sequences and by incorporating UMIs. Interrogation of

Ig/TCR repertoire and B/T cell transcriptomes in the same single cells has provided novel

insights in human, such as evaluating T cell clones within tumors [51], modeling of epitope

specificities [52, 53], and assessment of host-microbiome immune homeostasis [54].

However, to the best of our knowledge, for NHP models including rhesus macaque,

equivalent assays for single cell Ig and TCR repertoire analysis are still not available.

In this study, we sought to address the technical limitations of existing immune

repertoire sequencing protocols in rhesus macaque. To circumvent the highly difficult task

of correctly assembling Ig/TCR transcripts from short sequencing reads, we performed

long read transcriptome sequencing of rhesus tissue samples. We constructed the first

complete reference set for the constant regions of all known isotypes and chain types of

rhesus Ig and TCR repertoires from high-quality, full-length Ig and TCR transcript

sequences. Our resource provides the multiple rhesus Ig and TCR constant regions still

missing in the current IMGT database, avoiding complications due to incomplete

assemblies of their respective genomic loci. Using this unbiased collection of full-length

Ig and TCR transcript sequences, we were able to examine the diversities across the entire

variable regions of rhesus Ig and TCR transcripts and the coverage of available assays for

profiling rhesus Ig and TCR repertoires. Further, we designed new rhesus-specific assays

that allow much broader rhesus Ig and TCR repertoire analysis both at the single cell level

and using 5' RACE. These results furthermore demonstrate that immunoprofiling

resources and assays can be developed for other organisms similarly without complete

genomic assemblies of immune loci.

Materials and Methods

Tissue sample preparation and transcriptome sequencing

To construct cDNA libraries for PacBio SMRT sequencing, we used previously isolated Indian-origin rhesus macaque RNA from four tissue types: lymph node (LN), peripheral blood mononuclear cells (PBMC), whole blood (WB), and rectal biopsy (RB). To maximize the coverage of transcript diversity, we pooled RNA from multiple Indian-origin rhesus macaques. The pooled WB RNA was from six healthy, uninfected macaques, while the pooled LN, PBMC, and RB RNA were from 18 macaques infected with SIVmac251. The LN RNA came from a combination of peripheral and proximal mesenteric lymph nodes. Using a Clontech SMARTer kit, cDNA was produced from each pooled RNA sample followed by PCR amplification. The amplified cDNA samples were size selected using BluePippin (Sage Science) into four size fractions: 1-2 kb, 2-3 kb, 3-6 kb, >6 kb. Libraries were prepared using the SMRTbell template prep kit 1.0 and sequenced on the PacBio RS II platform with P6-C4 chemistry and 4-hour movie times at the University of Washington PacBio Sequencing Services site.

Data processing

Raw PacBio sequencing data was first run through the CCS protocol in SMRT Analysis 2.3.0 to generate Circular Consensus Sequence (CCS) reads. Each CCS read is the circular consensus sequence of a single molecule. CCS reads were classified as non-full-length and full-length, where the latter has all of the following detected: polyA tail, 5' cDNA primer, and 3' cDNA primer. full-length and non-full-length CCS sequences were aligned to immunoglobulin (Ig) and T cell receptor (TCR) databases using IGBLAST v1.8.0 [55] and

IMGT annotation [56]. Separate searches were carried out using either human or rhesus

macaque germline V, D, and J sequences where available (both for Ig, only human for

TCR). Full-length Ig hits were kept for downstream analyses when they were deemed

functional by IgBLAST output, meaning they lacked premature stop codons and were in-

frame. When Ig hits had both rhesus and human database hits, the rhesus annotation was

used in all subsequent analyses.

Generation of consensus constant region sequences

To validate the chain types reported by IgBLAST and gather additional information about

rhesus Ig/TCR isotypes and their subclasses, we further processed full-length Ig and TCR

constant region sequences as below. Using the J region reported by IgBLAST, we extracted

the constant region sequence for each full-length sequence. Next, we clustered extracted

constant region sequences using CD-HIT [57] with the following parameters: -c 0.97 -G 0

-aL   0.95   -AL   100   -aS   0.99   -AS   30

(https://github.com/Magdoll/cDNA_Cupcake/wiki/Tutorial:-Collapse-redundant-

isoforms-without-genome). Ig and TCR sequences were clustered separately. We only kept

clusters with at least three constituent sequences for downstream consensus sequence

analysis. For those clusters with at least ten sequences we also performed a second round

of more stringent clustering using a 99% identity cutoff (-c 0.99 with CD-HIT) to separate

quality sequences from those with multiple indels and to identify potential allotypes. From

the second round of more stringent clustering, consensus sequences were generated using

only the sequences from the largest remaining clusters, with the goal of removing low

quality sequences that formed many small clusters. When this second round of clustering

resulted in multiple large clusters of comparable size, a consensus sequence was generated

for each cluster. We used the cons tool from the EMBOSS v6.6.0 suite [58] to generate

consensus sequences, identifying and removing any remaining insertions at predominantly

gapped locations. Protein sequences were derived from the generated consensus sequences

using the EMBOSS transeq tool [58], where the frame with the longest protein sequence

was kept. We verified their identities via global alignments of the coding regions of cDNA

sequences to a database of IMGT human constant region nucleotide coding regions [56]

using the usearch_global tool in the USEARCH suite [59], considering only the best

alignment. We aligned consensus sequences belonging to the same chain type (and isotype

if applicable) using Clustal Omega [60], and removed any remaining redundancies in the

consensus sequences after manual visualization and curation in UGENE [61].

Next, we used these databases to systematically classify putative Ig and TCR

sequences using all CCS reads from the complete Iso-Seq dataset as input. CCS reads were

separately globally aligned to the custom databases of consensus Ig and TCR constant

region sequences using the usearch_global tool [59] with a 90% identity threshold to

ascertain chain type and isotype/subclass where appropriate. Sequences without large gaps

(>10bp) in their alignment were kept in the final assignment of Ig and TCR sequences, as

it was unclear if those sequences with large gaps were rare alternatively spliced transcripts

or the result of sequencing errors. We then compared the CCS reads confirmed as Ig/TCR

by our custom databases with the CCS reads identified by the initial IgBLAST search.

Based on these comparisons, the final set of Ig transcripts were filtered to only include

CCS reads that were confirmed by the custom database and also IgBLAST hits deemed

functional by rhesus IMGT annotation (Supplementary Fig. 1B). Further, the final set of

TCR transcripts only required CCS reads to be confirmed by the custom database

(Supplementary Fig. 1A), as it was unclear if IMGT functional annotation for human would

be informative for rhesus-derived transcripts. We quantified the rates of insertions and

deletions in the constant regions of confirmed Ig and TCR sequences using the CIGAR

strings in the global alignments of CCS reads against this custom constant reference

database by usearch_global [59]. Non Ig/TCR CCS reads of this complete Iso-Seq dataset

are being analyzed in a separate study.

Analysis of V-J usage and variable region diversity

The frequencies of V-J combinations were measured using the IMGT annotation reported

by IGBLAST [55, 56] for all final Ig transcript sequences ascertained in the above analysis.

Chi-square test for independence based on V-J combination frequencies was performed in

R [62] and by requiring that at least 50% of elements in each row and column have at least

10 counts. The diversity of the variable region sequences was determined by first removing

the constant region sequence determined by its alignment to the corresponding consensus

constant region sequences. Separate multiple sequence alignments of the most common V-

J regions were performed using Clustal Omega [60] and the consensus profiles were

visualized in R with ggplot2 [62, 63].

In silico **PCR analysis**

Common rhesus-specific MPCR assays for amplifying IGH, IGK, and IGL sequences [28,

29, 32] and TCRA and TCRB sequences [64, 65] were tested via *in silico* PCR analysis

using the UCSC isPcr tool [66]. For those primer sets with a nested design [28, 29], only

the inner primers were used for the analysis. The standard inner and outer reverse primers

used for the 10x B cell and T cell V(D)J assays were also assessed via *in silico* PCR analysis

[66] where a dummy adapter sequence was prepended to the 5' end of sequences to enable

analysis with the isPcr tool and only test reverse primer efficiencies. The efficiencies of

rhesus-specific MPCR assays [28, 29, 32] in amplifying the variable region (forward

primer) and constant region (reverse primer) were similarly tested by either appending and

prepending a dummy adapter sequence, respectively. In all uses of the isPcr tool, default

parameters were used, requiring a perfect match for the first 15 nucleotides on the 3' end

of the primer.

Design of rhesus-specific primers and PCR validation

Ig and TCR constant region inner and outer reverse primers were designed using NCBI

primerBLAST [67] with target Tms consistent with current 10x V(D)J assay forward

primers. For those targeting multiple consensus sequences, the search space was

constrained to consensus regions. To validate these 10x optimized reverse primers,

primerBLAST [67] was used to design standard PCR assays each with a forward primer

compatible with its respective 10x optimized reverse primer and target reference sequence.

This was only done for custom primers in the V(D)J assays, while primers adopted from

Sundling et al [28] were assumed to be efficacious. 100 nmole HPLC purified DNA

oligonucleotides were ordered from IDT. Reverse transcription was performed using

Qiagen QuantiTect Reverse Transcription Kit (Cat. No. 205313, Lot 157037104) following

the manufacturer's protocol with 1 ug rhesus macaque lymph node tissue RNA. The

resulting cDNA was then amplified using Takara Titanium TaqPCR kit (Cat. No. 639210,

Lot   1805465A)   following   the   manufacturer's   protocol

(https://www.takarabio.com/assets/documents/User%20Manual/PT3304-2.pdf),   altering

the recommended cycling conditions with 30 cycles and an annealing temperature and time

of 50°C and 1 minute. 400 ng of each PCR reaction was then run on a 10% TBE

polyacrylamide gel alongside Thermo Scientific 100 bp GeneRuler to confirm product size

from primer pairs.

Results

Generation of a complete reference collection for rhesus macaque Ig and TCR constant regions

Using PacBio transcriptome sequencing (the Iso-Seq method), we obtained over 2.8 million Circular Consensus Sequence (CCS) reads from four different rhesus macaque tissues (Supplementary Table 1). Each CCS read is the circular consensus sequence of a single transcript molecule [68]. About 33% of these CCS reads were full-length, i.e. contained the 5' cDNA primer, 3' cDNA primer, and polyA tail (Supplementary Table 1). Using the available IMGT annotation of the variable regions of Ig (human and rhesus) and TCR (human) germline sequences [56] and IgBLAST [55], we recovered 13,118 Ig and 2,534 TCR putative transcript sequences, respectively accounting for 1.4% and 0.28% of the total number of full-length CCS reads (Supplementary Table 1). Only functional Ig transcripts (i.e. those in-frame and without premature stop codons according to IMGT germline annotation) were used in downstream analyses (5,666 / 13,118 or 43% of full-length Ig transcripts). We kept all full-length TCR transcripts at this step, as it was unclear if their functionalities could be correctly determined with the use of human germline TCR reference sequences.

We first sought to characterize the constant regions of these full-length Ig and TCR transcript sequences. Since constant regions are significantly less variable, this would allow us to compile a complete reference set and accurately classify the different chain types and isotypes. Using the V(D)J annotation provided by IMGT [56], we extracted constant regions from the full-length Ig and TCR transcript sequences. We then clustered and curated these sequences to generate complete cDNA and coding sequences (Materials and

Methods). Complete cDNA sequences represent the entire constant region consensus (i.e.

including the 3' UTR), while the coding regions were generated at both the cDNA and

protein level. We obtained the complete consensus sequences for the following chain types

and isotypes with multiple references noted in parentheses: IGHA (5), IGHD, IGHE,

IGHG1, IGHG2, IGHG3, IGHG4, IGHM (2), IGLC, IGKC, TCRAC, TCRBC1, TCRBC2,

TCRDC, TCRGC1, and TCRGC2. We thus recovered complete reference sequences for

all known secreted IGH isotypes, and additionally two IGHA and one IGHM membrane-

bound reference sequences. We numbered membrane-bound rhesus Ig reference sequences

to be consistent with their corresponding secreted forms.

We identified three unique IGHA consensus sequences: two in both secreted and

bound form and one in only secreted form. Much of the variation among these sequences

was confined to the hinge region, which has previously been reported to be highly

heterogeneous among rhesus macaques [69]. In fact, we recovered three of the eight unique

hinge regions previously identified by [70] (data not shown). We numbered these unique

IGHA sequences based on the relative abundances we later determined within our samples

(i.e. IGHA*01 was the most abundant), following conventional naming for allelic variants.

We also identified all known subclasses of IGHG, TCRBC, and TCRGC. We

aligned the four IGHG cDNA coding sequences we recovered to the available rhesus IGHG

gDNA in IMGT to determine their subclasses. Each cDNA coding sequence reference

corresponded to one of the four IGHG subclasses with high sequence identities (98.6% to

99.9%). To properly classify the TCRBC subclasses, we aligned the two TCRBC protein

sequences to the two reference ORFs available in IMGT, yielding perfect matches (data

not shown). We leveraged the available human ORFs in IMGT to determine subclasses of

TCRGC, as reference sequences were not available for rhesus macaque. Using a multiple

sequence alignment, we assigned orthologous subclass naming to the rhesus references

based on the human reference with the highest structural similarity (data not shown).

Interestingly, 39% (979 of 2,534) of putative TCR transcript sequences identified

by the initial IgBLAST search were in fact Ig transcripts, evidenced by recapitulation of

several Ig consensus sequences among the set of TCR clusters and by successful alignment

of such sequences to the set of Ig reference sequences (Supplementary Fig. 1A). While this

did not prevent our complete recovery of TCR consensus sequences, it suggests that

commonly used human reference based IgBLAST searches could be inaccurate for rhesus

TCR repertoire analysis.

Classification of rhesus Ig and TCR transcript sequences
331

Next, we used these consensus sequences of rhesus constant regions to classify the full-

length rhesus Ig and TCR transcript sequences. When we aligned the constant regions of

full-length functional Ig transcripts to this newly constructed database of rhesus Ig cDNA

consensus sequences, we obtained successful assignments for 96% (5,415 / 5,666) of these

full-length Ig transcript sequences. Those transcript sequences that failed to align tended

to be other molecules from the immunoglobulin superfamily that contain V-set domains

(i.e. a variable region), representing false positives from our initial IgBLAST screen (data

not shown). Among those aligned, a small fraction (1%) of sequences had large structural

differences, indicating infrequent alternative splicing events (data not shown). To simplify

the downstream analysis, we removed these transcripts from further processing, yielding a

final set of 5,384 full-length Ig transcript sequences (Table 1). Among four tissue samples

sequenced here, the rectal biopsy and lymph node had the largest number of unique Ig

transcript sequences overall. Secreted IGHA and IGHG transcripts had the highest relative

abundance and were most prevalent in the rectal biopsy and lymph node samples. IGK and

IGL sequences were significantly less abundant than IGH isotypes in general, but they were

still detected across each tissue.

Similarly, we identified rhesus TCR transcript sequences by aligning raw full-

length CCS reads directly to this custom collection of rhesus TCR cDNA consensus

sequences. This precluded any bias against rhesus TCR transcript sequences that contained

V, D, and J genes with low similarity to those in the human IMGT reference. In total, we

assigned 741 full-length CCS reads as rhesus TCR transcript sequences (Table 2), 50 of

which (7%) we did not originally recover using the human IMGT reference database

(Supplementary Fig. 1A). This suggests there was a significant divergence between human

and rhesus TCR germline V(D)J genes, indicating that a species-specific germline database

is necessary for complete repertoire recovery. We detected the fewest TCR transcript

sequences in the rectal biopsy sample, while the whole blood, PBMC, and lymph node

samples each had larger recoveries of all TCR isotypes (Table 2). TCRA was the most

abundant in each tissue, representing 61% of all sequences recovered (455 / 741). TCRG2

had the second highest abundance with its strongest representation in PBMCs (Table 2).

We then assessed the sequence quality of these Ig and TCR transcripts, by

quantifying insertion and deletion events within the alignments of their constant region

sequences and the corresponding reference cDNA sequences. We elected not to evaluate

the rate of substitutions, as they are difficult to discern from allelic variation given our

sequencing depth and are generally less common than insertions and deletions in CCS reads

[33]. The rate of insertions in Ig constant region sequences was very low, with a mean of

0.11% and median of 0.06%. The mean and median deletion rates were slightly lower at

0.04% and 0%, respectively. Error rates within the TCR constant regions were comparable

to those of Ig constant regions, with mean insertion and deletion rates of 0.15% and 0.07%

and median rates of 0.07% and 0%, respectively. These low error rates reflected the high

quality of the final set of full-length rhesus Ig and TCR transcript sequences we obtained

in this study.

We found the recovery of TCR transcripts to be significantly more accurate and

marginally more sensitive when using our custom TCR constant region database for

identification instead of the traditional IgBLAST approach with human germline

annotation (Supplementary Fig. 1A). We thus elected to align CCS reads directly to our

custom Ig constant region database to make a similar comparison. We recovered 11,451 Ig

transcripts using this strategy; of these, 5,384 (47%) were functional IgBLAST hits (see

Table 1), 5,858 (51%) were non-functional IgBLAST hits, and 209 (2%) were not detected

by IgBLAST (Supplementary Fig. 1B). B cells that contain non-functional receptor

sequences are known to apoptose early in their development; therefore, the proportion of

non-functional IgBLAST hits observed here may be a reflection of the B cell population

captured with these full-length transcripts. Many of the full-length Ig transcripts that eluded

IgBLAST detection had enough sequence upstream of the constant region to harbor a

complete variable region (127 or 1% overall) (Supplementary Fig. 1B), indicating that

some variable region genes may be significantly diverged from those annotated in IMGT.

Usage of known V, D, and J genes in Ig transcripts indicates broad coverage of rhesus Ig repertoire

To assess our overall coverage of the rhesus Ig repertoire, we examined the usage of individual V, D, and J genes among these full-length rhesus Ig sequences based on the alignment of their variable regions to rhesus Ig germline annotations in IMGT [56]. We detected all known gene families of rhesus IGHV, IGHD, and IGHJ genes among the four tissue samples (Fig. 2). The majority of IGHV and IGHJ genes detected were from the IGHV4 and IGHJ4 families, respectively, while there was broad coverage of IGHD gene families. Among light chain gene families, IGKV1 and 2 as well as IGLV1, 2, and 3 were the most frequent. IGKJ families were relatively less skewed in frequency, though IGLJ1 was in slightly higher frequency than other IGLJ families. The only known gene families we did not detect in these tissue samples were IGKV5, IGLJ4, and IGLJ5, likely due to the overall lower abundance of light chain Ig sequences we observed (Table 1). Interestingly, relative frequencies of these different gene families were highly consistent across the four tissues analyzed here (Supplementary Fig. 2), despite drastically different sampling depth (Fig. 2). Only light chain gene families within PBMCs and whole blood deviated from this trend. Since we had significantly less detection of IGK and IGL sequences in these tissues (Table 1), their observed deviations require further investigation.

Given the sufficient depth of coverage observed for IGH V, D, and J gene families, we calculated the combination frequencies of these V and J genes. Consistent with V and J gene frequencies observed in Fig. 2, the majority of V-J combination events contained IGHV4-2 and/or IGHJ4 (Fig. 3). Using the most abundant V and J genes, we tested the independence of V-J recombination events. We observed a significant nonrandom

distribution of V-J recombination frequencies (chi square = 77.6, df = 15, p = 1.9e-10). Furthermore, we discovered a strong positive correlation between the V-J recombination rates in the lymph node and rectal biopsy samples (Pearson correlation = 0.86). Since these two samples had highly different compositions of IGH isotypes (Table 1), we also reasoned that there might be no discernable association between the variable regions and constant regions in general.

Sequence diversity existing across the entire variable region shows MPCR amplification of rhesus immune repertoires is inherently biased

The collection of unbiased full-length Ig and TCR transcript sequences also provided a unique opportunity for exploring the sequence diversity across the entire variable region, the target of current repertoire amplification efforts. We assessed the efficiencies of three commonly used rhesus-specific Ig MPCR strategies [28, 29, 32], by examining the sites targeted by their primers in the context of their respective Ig V genes. We selected the three most abundant IGHV genes represented among the unique Ig transcripts: IGHV4-2 (1,383 sequences), IGHV1-1 (163 sequences), and IGHV3-9 (128 sequences), and generated consensus profiles for each. We discovered that the rhesus primers designed to target these genes all locate in regions rich in sequence diversity (Fig. 4). For example, Sundling et al [28] targeted well-conserved regions of IGHV4-1 and IGHV1-1 (Fig. 4A-B), yet appeared to target a region in IGHV3-9 that had a low percentage of sequences sharing the same consensus nucleotides (Fig. 4C). The low consensus observed in all three profiles at the 5' end of the cDNA sequence (left side of profiles) was likely a result of the lack of guaranteed

5' capture among CCS reads. However, it was confined to regions upstream of the primer

target sites (Fig. 4A-C) and thus did not affect our assessment of available primers.

To illustrate the limitation that the observed diversity could reduce the overall

coverage of these MPCR approaches, we evaluated these primer sets via *in silico* PCR

analysis with the full-length Ig transcript sequences as the source of potential template

transcripts (Fig. 5). Similarly, we also leveraged known rhesus-specific MPCR strategies

for TCR [64, 65] in this analysis (Fig. 5). We first tested only the forward (V gene) primers

from these Ig and TCR primer sets [28, 29, 32, 64, 65] and found that IGH, IGK, and IGL

amplification rates did not exceed 87% and were as low as 45% (Fig. 5A). Amplification

rates for TCRA and TCRB were similar, ranging from 53% to 73% (Fig. 5A). The Ig primer

set designed earliest by Sundling et al in 2012 [28] had the highest recovery of IGH

sequences (76%), while the most recent Ig primer set designed by Rosenfeld et al in 2019

[32] had the highest recovery of IGK (87%) and IGL (71%). The TCRB specific primers

designed by Li et al [65] recovered 73% of the TCRB sequences, while Chen et al [64]

recovered 53% of TCRA sequences.

As expected, when we assessed both forward and reverse primers together, the low

percentages of sequences amplified largely remained the same (Fig. 5B). Notably, the

percentages for Rosenfeld et al [32] and the percentage for Li et al [65] were even lower,

as their reverse primers targeted J gene segments instead of the constant region. Indeed,

when we tested the constant region primers alone (Fig. 5C), we found that Sundling et al

[28], Wiehe et al [29], and Chen et al [64] had near perfect amplification rates, which was

expected.

456 **Design of rhesus-specific B- and T-cell V(D)J assays for single cell analysis**

457 Both our consensus profiling (Fig. 4) and *in silico* PCR analysis (Fig. 5A-C) showed that

458 primers targeting variable regions are the source of bias in targeted amplification of rhesus

459 Ig and TCR repertoires. Other methods, particularly RACE-seq and those using scRNA-

460 seq, avoid this problem by only using primers targeting the constant region. Since there are

461 no known single cell repertoire analysis assays for rhesus macaque, we first checked the

462 efficiency of human B and T cell V(D)J assays developed by 10x Genomics [66] against

463 our collection of Ig and TCR transcript sequences. We found that neither assay was able to

464 fully capture this set of rhesus Ig and TCR transcript sequences (Fig. 5D). The human inner

465 primers successfully amplified IGHD (100%), IGHE (83%), IGHG1 (99%), IGHG2

466 (98%), IGHG3 (97%), IGHG4 (100%), TCRA (94%), TCRB1 (100%), and TCRB2 (93%).

467 Meanwhile, the outer primers only amplified IGHG1 (99%), IGHG2 (97%), IGHG3 (97%),

468 IGHG4 (100%) and IGL (99%). Notably, TCRD, TCRG1, and TCRG2 were not included

469 in this analysis, as 10x Genomics assays do not cover these transcripts.

470 We further assessed sequence homology between human and rhesus reference

471 sequences for the constant regions. For each Ig and TCR isotype, we aligned our reference

472 cDNA coding sequences for rhesus macaque with those available for human and evaluated

473 the compatibility of their respective 10x human primers where relevant. For example, we

474 aligned human IGHA1 and IGHA2 to the five rhesus IGHA consensus sequences (3

475 secreted, 2 membrane-bound) and examined the regions targeted by 10x human primers

476 (Fig. 6A). Overall, the two human cDNA coding sequences had high similarity to the

477 rhesus consensus sequences, averaging 89.7% identity. However, multiple nucleotide

478 differences existed in the regions of rhesus reference sequences targeted by the 10x human

outer and inner primers (Fig. 6A-C). These primer inconsistencies were not specific to

IGHA, as all other human primers with the exception of the IGHG outer primer had

mismatches with at least one of its corresponding rhesus reference sequences (data not

shown). These results clearly indicated rhesus-specific primers are necessary to perform

comparable single cell analysis for rhesus macaques.

To properly design rhesus-specific Ig and TCR constant region primers that are

compatible with respective 10x human V(D)J assay designs, we first selected existing

primers that were free of mismatches, which included the 10x human IGHG outer primer

and rhesus primers designed by Sundling et al [28] (Supplementary data). Notably, some

primers that successfully *in silico* PCR amplified Ig/TCR transcripts contained one or more

mismatches thus requiring updated designs (Materials and Methods). For isotypes that

lacked an inner and/or outer primer, we designed new primers with melting temperatures

(Tms) compatible with that of the 10x forward primer (Supplementary data). Similar to the

Ig primers, 10x human TCR primers were also largely incompatible with the rhesus

consensus sequences. We found that only the inner primer for TCRB had a perfect match

and thus designed all other primers for rhesus macaque (Supplementary data). We also

included primers that target TCRD and TCRG in the final set of our rhesus-specific T-cell

V(D)J primers, which are not included in the human 10x assays [66].

To ensure these newly designed primers can amplify rhesus transcripts

experimentally, for each of them we designed a corresponding forward primer that was

upstream but still within the constant region (Supplementary data). Then we tested all these

custom inner and outer primers by pairing with their forward primers via PCR (Materials

and Methods). We obtained the desired product size for each assay (Supplementary Fig.

3), confirming that these rhesus specific primers newly designed for single cell based

V(D)J assays targeted rhesus Ig/TCR transcripts as expected. In some assays, we also

observed additional bands, but the desired bands tended to be visibly the most prominent

(7 of 8 outer primers and 2 of 5 inner primers tested). Many of those additional bands were

larger than the desired product size (Supplementary Fig. 3), suggesting some forward

primers might have shared certain sequence similarities with the highly diverse upstream

variable regions. The actual single cell assays use nested template switch PCR; therefore,

these forward primers are not needed and such off-target amplification would not be a

concern. Together, these results indicate that the primers we designed will be useful for

single cell assays and compatible with the 10x Genomics platform.

Discussion

In this study, we profiled Ig and TCR repertoires using long read transcript sequencing of tissues collected from Indian-origin rhesus macaques, one of the most widely used NHP models. Since we obtained high-quality, full-length sequences for individual Ig and TCR transcripts, no sequence assembly was necessary. This unbiased profiling analysis allowed us to compile the first complete reference set for the constant regions of all expected rhesus Ig and TCR isotypes and chain types based on homology with human, including three newly identified reference sequences. While most of publicly available rhesus Ig and TCR constant region sequences were computationally predicted from genomic sequences and/or partial sequences, our reference sequences were from directly sequenced transcripts and full length. We were able to define the coding regions for these constant regions, yielding full coding sequences as well as cDNA sequences containing 3' UTRs.

An immediate benefit of having this complete constant region reference is improved accuracy of rhesus Ig/TCR transcripts identification. For example, we found that about 39% of rhesus TCR transcripts that we initially identified by the commonly used combination of IgBLAST and IMGT databases were actually Ig transcripts, based on the alignment of their constant region portion against these reference sequences. This miss-assignment could be due to several reasons. First, the current IMGT database has a limited number of TCR variable region germline sequences for rhesus macaque. Second, the alignment of rhesus Ig/TCR transcripts to the human variable region reference is not sufficiently reliable. The incomplete 5' end of some rhesus Ig/TCR transcript sequences was also a contributing factor. In the future, we expect that our reference sequences will improve identification of novel germline variable genes extracted from recombined Ig and

TCR sequences in rhesus macaque, as the number of false positives will be substantially

abrogated.

These full-length Ig/TCR transcript sequences offered insights into the rhesus

immune system. For the Ig constant regions, we identified both cell membrane-bound and

secreted forms that we differentiated by distinct 3' end splicing. This demonstrates our

ability to detect the alternative splicing events among these multi-exonic regions. As the

genomic assembly and annotation of these constant region loci improves, long read

sequencing may enable discovery of similar alternative exon usage as described for human

Ig heavy chain constant region genes [71]. In addition, we recovered three different

allotypes for IGHA with significant variation in the hinge region. When this hinge region

is longer, the avidity for antigen interactions is increased at the expense of increased

susceptibility to pathogen-derived proteases known to target the hinge region [72]. We

suspect that our ability to identify hinge region variation was driven by the high

heterogeneity of the alpha constant region [69] as well as its highest frequency among the

Ig isotypes we recovered. Given the interspecies variability of IGHG between macaques

and humans [73] and the heterogeneity of rhesus IGHC genes in general [16], it is

imperative that future rhesus Ig gene analyses also consider this allotypic diversity to

improve the translatability of NHP models.

Since we did not use any targeted experimental approaches to obtain these full-

length Ig/TCR transcripts, we were able to examine the diversities of rhesus Ig/TCR

variable regions in an unbiased manner. We found that large diversities existed across the

entire variable regions of rhesus Ig and TCR transcripts. This is especially relevant, since

available assays for profiling rhesus Ig/TCR repertoires have relied on the consensus

primers derived from a limited number of available rhesus variable region sequences. Previous benchmarking analyses for targeted repertoire sequencing in human used either spiked-in samples [43] or samples of unknown composition [42], motivating us to leverage our collected sequences for a truly unbiased assessment of conventional rhesus-specific MPCR designs for Ig [28, 29, 32] and TCR [64, 65]. MPCR targets specific V gene segments (forward primer) and either specific J gene segments or constant regions (reverse primer) to generate amplicon libraries. Not surprisingly, we found that all of these assays failed to cover a significant percentage of these rhesus Ig/TCR transcripts. We do not contend that this method be avoided altogether moving forward, but additional improvements will be necessary. For example, databases of variable region gene segments can be improved using the IgDiscover tool [18]. Targeting the more conserved 5' UTR of V genes may reduce the amount of primers needed in MPCR designs and yield amplicons covering the entire variable region but still need to be designed for rhesus macaque [34]. New algorithms such as the recently described version of the tool for Ig Genotype Elucidation via Rep-Seq (TIgGER) for identifying human Ig [74] can also improve germline allele assignment if adapted for rhesus macaque.

Alternatively, scRNA-seq and RACE-seq only use primers that target the constant regions of these Ig/TCR transcripts. In particular, scRNA-seq combines repertoire analysis and transcriptome profiling in single B and T cells [51-54], and it can identify chain pairs within single cells [75, 76]. Our analysis showed that human-specific constant region primers from commercially available assays were not compatible for use in rhesus macaque. For these reasons, we elected to design the first single cell based B- and T-cell V(D)J assays for rhesus macaque. We leveraged the complete set of rhesus constant region

reference sequences compiled here to ensure a broad coverage of rhesus Ig/TCR diversity

and experimentally validated the newly designed rhesus-specific primers. Our assays are

fully compatible with the standard protocols from 10x Genomics, currently a popular

platform for single cell analysis. These primers can also be used or modified for RACE-

seq applications, considering similar library preparation techniques.

More broadly, this study also demonstrated an accessible strategy for developing

immune repertoire resources and assays for less studied organisms. Traditionally, the

development of Ig/TCR resources would start with collecting large numbers of species-

specific transcript and germline sequences. Since Ig/TCR loci are among the most complex

regions in the genome due to their duplicated and polymorphic nature, proper sequencing

and assembly of these genomic regions requires sophisticated efforts [77]. Consequently,

complete assemblies are only available for selected model organisms such as human and

mouse; meanwhile, these loci still do not have complete assemblies for rhesus macaque

despite recent efforts [16]. The PacBio Iso-Seq approach we used here sequences full-

length Ig/TCR transcripts directly and in a high-throughput manner, bypassing transcript

assembly issues. The performance of this strategy could be further improved as well. For

example, compared to the whole tissue sample used here, use of isolated B and T cells may

significantly improve the recovery of these repertoire sequences. The custom

computational solution we developed and describe here can be modified for other species

by replacing constant region references from a related species to identify putative species-

specific Ig/TCR transcript sequences. However, additional information like Ig/TCR

germline allele assignment, the genomic order of individual genes, and copy number

variations will still require complete genomic sequencing and assembly.

In this study, we systematically profiled the diversity of rhesus Ig and TCR genes

using full-length transcriptome sequencing. We performed benchmark analysis of common

MPCR based targeted sequencing strategies for rhesus macaque and demonstrated that

available MPCR assays do not cover the rhesus repertoire diversity adequately. Our

construction of a complete reference set for rhesus macaque Ig/TCR constant regions

enabled the design of the first rhesus-specific single cell based V(D)J assays, addressing

many of the current technical limitations of rhesus macaque repertoire analysis.

Furthermore, our approach circumvents the computational challenges concomitant with the

genomic assembly of these complex immune genes by combining long read sequencing

and new computational solutions. This study thus represents a new direction for developing

nascent immune resources.

616   Footnotes

617   Grant support

This project has been funded in part with funds from the National Institute of Allergy and

Infectious Diseases, National Institutes of Health, Department of Health and Human

Services, R21AI120713 (XP). This project has been funded in part with funds from the

National Institute of Allergy and Infectious Diseases, National Institutes of Health,

Department of Health and Human Services, under Contract No. HHSN272201300010C

and HHSN272201800008C (MG) and the National Institutes of Health, Office of the

Director P51OD010425 (MG).

626   Abbreviations

CCS, circular consensus sequence; NHP, non-human primate; Ig, immunoglobulin; IGH, Ig heavy chain; IGHA, IGH alpha; IGHD, IGH delta; IGHE, IGH epsilon; IGHG, IGH gamma; IGHM, IGH mu; IGK, Ig kappa chain; IGL, Ig lambda chain; TCR, T cell receptor; TCRA, TCR alpha; TCRB, TCR beta; TCRD, TCR delta; TCRG, TCR gamma; IMGT, the international ImMunoGeneTics information system; MPCR, multiplex PCR; RACE, rapid amplification of cDNA ends.

References

1. Williams, K., A. Lackner, and J. Mallard, *Non-human primate models of SIV infection and CNS neuropathology*. Current Opinion in Virology, 2016. **19**: p. 92-98.

2. Zhao, H., Y.H. Jiang, and Y.Q. Zhang, *Modeling autism in non-human primates: Opportunities and challenges*. Autism Res, 2018. **11**(5): p. 686-694.

3. Flynn, J.L., et al., *Immunology studies in non-human primate models of tuberculosis*. Immunol Rev, 2015. **264**(1): p. 60-73.

4. Reyes, L.F., et al., *A Non-Human Primate Model of Severe Pneumococcal Pneumonia*. PLOS ONE, 2016. **11**(11): p. e0166092.

5. Estes, J.D., S.W. Wong, and J.M. Brenchley, *Nonhuman primate models of human viral infections*. Nature reviews. Immunology, 2018. **18**(6): p. 390-404.

6. Hansen, S.G., et al., *Prevention of tuberculosis in rhesus macaques by a cytomegalovirus-based vaccine*. Nature Medicine, 2018. **24**: p. 130.

7. Carpenter, S.M. and S.M. Behar, *A new vaccine for tuberculosis in rhesus macaques*. Nature Medicine, 2018. **24**: p. 124.

8.   Hansen, S.G., et al., *Profound early control of highly pathogenic SIV by an*

effector memory T-cell vaccine. Nature, 2011. **473**: p. 523.

9.   Hansen, S.G., et al., *Immune clearance of highly pathogenic SIV infection.*

Nature, 2013. **502**: p. 100.

10.   Carroll, T.D., et al., *Efficacy of Influenza Vaccination of Elderly Rhesus*

Macaques Is Dramatically Improved by Addition of a Cationic Lipid/DNA

Adjuvant. The Journal of Infectious Diseases, 2013. **209**(1): p. 24-33.

11.   Adams Waldorf, K.M., et al., *Fetal brain lesions after subcutaneous inoculation*

of Zika virus in a pregnant nonhuman primate. Nat Med, 2016. **22**(11): p. 1256-

1259.

12.   Clement, K.H., et al., *Vaccination of Rhesus Macaques with the Anthrax Vaccine*

Adsorbed Vaccine Produces a Serum Antibody Response That Effectively

Neutralizes Receptor-Bound Protective Antigen In Vitro. Clinical and Vaccine

Immunology, 2010. **17**(11): p. 1753-1762.

13.   Awasthi, S., et al., *An HSV-2 Trivalent Vaccine Is Immunogenic in Rhesus*

Macaques and Highly Efficacious in Guinea Pigs. PLoS Pathog, 2017. **13**(1): p.

e1006141.

14.   Zimin, A.V., et al., *A new rhesus macaque assembly and annotation for next-*

generation sequencing analyses. Biol Direct, 2014. **9**(1): p. 20.

15.   Harding, J.D., *Nonhuman Primates and Translational Research: Progress,*

Opportunities, and Challenges. ILAR Journal, 2017. **58**(2): p. 141-150.

16.   Ramesh, A., et al., *Structure and Diversity of the Rhesus Macaque*

672   *Immunoglobulin Loci through Multiple De Novo Genome Assemblies.* Front

673   Immunol, 2017. **8**: p. 1407.

674   17.   Cirelli, K.M., et al., *Slow Delivery Immunization Enhances HIV Neutralizing*

675   *Antibody and Germinal Center Responses via Modulation of Immunodominance.*

676   Cell, 2019. **177**(5): p. 1153-1171.e28.

677   18.   Corcoran, M.M., et al., *Production of individualized V gene databases reveals*

678   *high levels of immunoglobulin genetic diversity.* Nature Communications, 2016.

679   **7**: p. 13642.

680   19.   Nishana, M. and S.C. Raghavan, *Role of recombination activating genes in the*

681   *generation of antigen receptor diversity and beyond.* Immunology, 2012. **137**(4):

682   p. 271-281.

683   20.   Little, A.J., et al., *Chapter 2 - The Mechanism of V(D)J Recombination*, in

684   *Molecular Biology of B Cells (Second Edition)*, F.W. Alt, et al., Editors. 2015,

685   Academic Press: London. p. 13-34.

686   21.   Hesslein, D.G.T. and D.G. Schatz, *Factors and Forces Controlling V(D)J*

687   *Recombination*, in *Advances in Immunology*, F.J. Dixon, Editor. 2001, Academic

688   Press. p. 169-232.

689   22.   Murphy, K. and C. Weaver, *Janeway's immunobiology*. 2017.

690   23.   Lucas, J.S., et al., *Chapter 1 - The Structure and Regulation of the*

691   *Immunoglobulin Loci*, in *Molecular Biology of B Cells (Second Edition)*, F.W.

692   Alt, et al., Editors. 2015, Academic Press: London. p. 1-11.

24. Manis, J.P., M. Tian, and F.W. Alt, *Mechanism and control of class-switch recombination.* Trends in Immunology, 2002. **23**(1): p. 31-39.

25. Markoulatos, P., N. Siafakas, and M. Moncany, *Multiplex polymerase chain reaction: a practical approach.* J Clin Lab Anal, 2002. **16**(1): p. 47-51.

26. Hodges, E., et al., *Genome-wide in situ exon capture for selective resequencing.* Nat Genet, 2007. **39**(12): p. 1522-7.

27. Yeku, O. and M.A. Frohman, *Rapid Amplification of cDNA Ends (RACE)*, in *RNA: Methods and Protocols*, H. Nielsen, Editor. 2011, Humana Press: Totowa, NJ. p. 107-122.

28. Sundling, C., et al., *Isolation of antibody V(D)J sequences from single cell sorted rhesus macaque B cells.* J Immunol Methods, 2012. **386**(1-2): p. 85-93.

29. Wiehe, K., et al., *Antibody light-chain-restricted recognition of the site of immune pressure in the RV144 HIV-1 vaccine trial is phylogenetically conserved.* Immunity, 2014. **41**(6): p. 909-18.

30. Fu, L., et al., *A comprehensive profiling of T- and B-lymphocyte receptor repertoires from a Chinese-origin rhesus macaque by high-throughput sequencing.* PLOS ONE, 2017. **12**(8): p. e0182733.

31. Sundling, C., et al., *Single-cell and deep sequencing of IgG-switched macaque B cells reveal a diverse Ig repertoire following immunization.* Journal of immunology (Baltimore, Md. : 1950), 2014. **192**(8): p. 3637-3644.

32. Rosenfeld, R., et al., *A primer set for comprehensive amplification of V-genes from rhesus macaque origin based on repertoire sequencing.* J Immunol Methods, 2019. **465**: p. 67-71.

33.   Zhang, W., et al., *Identification of Variable and Joining Germline Genes and Alleles for Rhesus Macaque from B Cell Receptor Repertoires.* The Journal of Immunology, 2019. **202**(5): p. 1612-1622.

34.   Vázquez Bernat, N., et al., *High-Quality Library Preparation for NGS-Based Immunoglobulin Germline Gene Inference and Repertoire Expression Analysis.* Frontiers in Immunology, 2019. **10**(660).

35.   Benichou, J., et al., *Rep-Seq: uncovering the immunological repertoire through next-generation sequencing.* Immunology, 2012. **135**(3): p. 183-191.

36.   Mamedov, I.Z., et al., *Preparing unbiased T-cell receptor and antibody cDNA libraries for the deep next generation sequencing profiling.* Front Immunol, 2013. **4**: p. 456.

37.   He, L., et al., *Toward a more accurate view of human B-cell repertoire by next-generation sequencing, unbiased repertoire capture and single-molecule barcoding.* Sci Rep, 2014. **4**: p. 6778.

38.   Fu, G.K., et al., *Molecular indexing enables quantitative targeted RNA sequencing and reveals poor efficiencies in standard library preparations.* Proc Natl Acad Sci U S A, 2014. **111**(5): p. 1891-6.

39.   Turchaninova, M.A., et al., *High-quality full-length immunoglobulin profiling with unique molecular barcoding.* Nat Protoc, 2016. **11**(9): p. 1599-616.

40.   Rock, E.P., et al., *CDR3 length in antigen-specific immune receptors.* The Journal of experimental medicine, 1994. **179**(1): p. 323-328.

41.   Safonova, Y. and P.A. Pevzner, *De novo Inference of Diversity Genes and*
738   *Analysis of Non-canonical V(DD)J Recombination in Immunoglobulins.* Frontiers
739   in Immunology, 2019. **10**(987).

740   42.   Bashford-Rogers, R.J., et al., *Capturing needles in haystacks: a comparison of B-*
741   *cell receptor sequencing methods.* BMC Immunology, 2014. **15**(1): p. 29.

742   43.   Liu, X., et al., *Systematic Comparative Evaluation of Methods for Investigating*
743   *the TCRβ Repertoire.* PLOS ONE, 2016. **11**(3): p. e0152464.

744   44.   Macosko, Evan Z., et al., *Highly Parallel Genome-wide Expression Profiling of*
745   *Individual Cells Using Nanoliter Droplets.* Cell, 2015. **161**(5): p. 1202-1214.

746   45.   Zheng, G.X.Y., et al., *Massively parallel digital transcriptional profiling of single*
747   *cells.* Nature Communications, 2017. **8**: p. 14049.

748   46.   Chattopadhyay, P.K., et al., *Single-cell technologies for monitoring immune*
749   *systems.* Nature Immunology, 2014. **15**: p. 128.

750   47.   Proserpio, V. and B. Mahata, *Single-cell technologies to study the immune system.*
751   Immunology, 2016. **147**(2): p. 133-140.

752   48.   Papalexi, E. and R. Satija, *Single-cell RNA sequencing to explore immune cell*
753   *heterogeneity.* Nature Reviews Immunology, 2017. **18**: p. 35.

754   49.   Stubbington, M.J.T., et al., *Single-cell transcriptomics to explore the immune*
755   *system in health and disease.* Science, 2017. **358**(6359): p. 58-63.

756   50.   Chen, H., F. Ye, and G. Guo, *Revolutionizing immunology with single-cell RNA*
757   *sequencing.* Cellular & Molecular Immunology, 2019. **16**(3): p. 242-249.

758   51.   Neal, J.T., et al., *Organoid Modeling of the Tumor Immune Microenvironment.*
759   Cell, 2018. **175**(7): p. 1972-1988.e16.

52.   Glanville, J., et al., *Identifying specificity groups in the T cell receptor repertoire.*

Nature, 2017. **547**: p. 94.

53.   Dash, P., et al., *Quantifiable predictive features define epitope-specific T cell*

receptor repertoires. Nature, 2017. **547**: p. 89.

54.   Ichinohe, T., et al., *Next-Generation Immune Repertoire Sequencing as a Clue to*

Elucidate the Landscape of Immune Modulation by Host–Gut Microbiome

Interactions. Frontiers in Immunology, 2018. **9**(668).

55.   Ye, J., et al., *IgBLAST: an immunoglobulin variable domain sequence analysis*

tool. Nucleic Acids Res, 2013. **41**(Web Server issue): p. W34-40.

56.   Lefranc, M.-P., et al., *IMGT, the international ImMunoGeneTics database.*

Nucleic Acids Research, 1999. **27**(1): p. 209-212.

57.   Fu, L., et al., *CD-HIT: accelerated for clustering the next-generation sequencing*

data. Bioinformatics, 2012. **28**(23): p. 3150-2.

58.   Rice, P., I. Longden, and A. Bleasby, *EMBOSS: the European Molecular Biology*

Open Software Suite. Trends Genet, 2000. **16**(6): p. 276-7.

59.   Edgar, R.C., *Search and clustering orders of magnitude faster than BLAST.*

Bioinformatics, 2010. **26**(19): p. 2460-1.

60.   Chojnacki, S., et al., *Programmatic access to bioinformatics tools from EMBL-*

EBI update: 2017. Nucleic Acids Res, 2017. **45**(W1): p. W550-w553.

61.   Okonechnikov, K., et al., *Unipro UGENE: a unified bioinformatics toolkit.*

Bioinformatics, 2012. **28**(8): p. 1166-1167.

62.   R Development Core Team, *R: A language and environment for statistical*

computing. 2010, R Foundation for Statistical Computing: Vienna, Austria.

63.   Wickham, H., *ggplot2: Elegant Graphics for Data Analysis*. 2016: Springer-

Verlag New York.

64.   Chen, Z.W., et al., *Conserved T-cell receptor repertoire in simian*

immunodeficiency virus-infected rhesus monkeys. J Immunol, 1993. **151**(4): p.

2177-87.

65.   Li, Z., et al., *Comprehensive analysis of the T-cell receptor beta chain gene in*

rhesus monkey by high throughput sequencing. Scientific Reports, 2015. **5**: p.

10092.

66.   Kuhn, R.M., D. Haussler, and W.J. Kent, *The UCSC genome browser and*

associated tools. Brief Bioinform, 2013. **14**(2): p. 144-61.

67.   Ye, J., et al., *Primer-BLAST: a tool to design target-specific primers for*

polymerase chain reaction. BMC Bioinformatics, 2012. **13**: p. 134.

68.   Eid, J., et al., *Real-time DNA sequencing from single polymerase molecules*.

Science, 2009. **323**(5910): p. 133-8.

69.   Scinicariello, F. and R. Attanasio, *Intraspecies heterogeneity of immunoglobulin*

alpha-chain constant region genes in rhesus macaques. Immunology, 2001.

103(4): p. 441-448.

70.   Rogers, K.A., et al., *Nonhuman Primate IgA: Genetic Heterogeneity and*

Interactions with CD89. The Journal of Immunology, 2008. **180**(7): p. 4816-4824.

71.   Vollmers, C., et al., *Novel Exons and Splice Variants in the Human Antibody*

Heavy Chain Identified by Single Cell and Single Molecule Sequencing. PLOS

ONE, 2015. **10**(1): p. e0117050.

72.   Woof, J.M. and M.A. Kerr, *IgA function--variations on a theme.* Immunology,

806   2004. **113**(2): p. 175-177.

807   73.   Crowley, A.R. and M.E. Ackerman, *Mind the Gap: How Interspecies Variability*

808   *in IgG and Its Receptors May Complicate Comparisons of Human and Non-*

809   *human Primate Effector Function.* Frontiers in Immunology, 2019. **10**(697).

810   74.   Gadala-Maria, D., et al., *Identification of Subject-Specific Immunoglobulin Alleles*

811   *From Expressed Repertoire Sequencing Data.* Frontiers in Immunology, 2019.

812   **10**(129).

813   75.   Grigaityte, K., et al., *Single-cell sequencing reveals αβ chain pairing shapes the T*

814   *cell repertoire.* bioRxiv, 2017: p. 213462.

815   76.   Carter, J.A., et al., *T-cell receptor αβ chain pairing is associated with CD4+ and*

816   *CD8+ lineage specification.* bioRxiv, 2018: p. 293852.

817   77.   Watson, C.T., et al., *Complete haplotype sequence of the human immunoglobulin*

818   *heavy-chain variable, diversity, and joining genes and characterization of allelic*

819   *and copy-number variation.* American journal of human genetics, 2013. **92**(4): p.

820   530-546.

Figure Legends

Fig 1. Overview of how immunoglobulin (Ig) and T cell receptors (TCR) form, beginning

with germline DNA. Within a single cell (B or T), two chains are generated separately

through either VJ or VDJ rearrangement (somatic recombination) at the DNA level. These

molecules are then transcribed and spliced to form mRNA molecules that include a

constant region (C) domain. The constant region is what determines the specific chain type

shown in the table beneath the graphic. For example, in a B cell, Ig consists of a heavy

chain and a light chain. Depending on its constant region, the heavy chain is further

classified as alpha (IGHA), delta (IGHD), epsilon (IGHE), gamma (IGHG), or mu

(IGHM), while the light chain is either kappa (IGK) or lambda (IGL). The sources of

diversity are analogous across B and T cells, with the exception of the former undergoing

somatic hypermutation, a process that introduces random mutations in the variable (VJ or

VDJ) region.

Fig 2. Stacked bar plot of absolute transcript abundance of V, D, and J genes for each Ig

chain type, where colors of the bar indicate the tissue source of the gene. Variable gene

abundance was computed using full-length Ig transcripts reported in Table 1. Each

individual plot shows the various known gene families of a particular gene segment type

(e.g. VH), where gene families are clusters of highly similar gene segments within that

gene segment type.

Fig 3. Dot plot of various IGHV and IGHJ combinations. The sizes of dots are proportional

to the relative frequencies among the full-length Ig transcript sequences. The gene

segments are conventionally named according to gene family (e.g. IGHV1) followed by

the specific gene segment number. Gene segments simply have the family gene name (e.g.

IGHJ1) when there is only one known gene segment in the family.

Fig 4. Consensus profiles of the most abundant IGHV genes detected. Panels **A-C**

respectively show the percentage of nucleotides at each position that match the consensus

(most abundant) nucleotide for IGHV4-2, IGHV1-1, and IGHV3-9. The IGHV4-2

consensus was constructed from 1,383 sequences, while IGHV1-1 was from 163 sequences

and IGHV3-9 was from 128 sequences. The regions targeted by primers from (S)undling

et al [28], (W)iehe et al [29], and (R)osenfeld et al [32] respectively have either a yellow,

red, or green bar above.

Fig 5. *In silico* PCR analysis of rhesus-specific primers from (S)undling et al [28], (W)iehe

et al [29], (R)osenfeld et al [32], (C)hen et al [64], and (L)i et al [65] as well as human 10x

B and T cell V(D)J reverse primers [66]. Only the inner primers were tested for S, W, while

both inner and outer 10x primers were tested. The primers from C target both TCRA and

TCRB, while those from L only target TCRB, as indicated by the bars shown in **A-C**. The

percentage of transcripts amplified *in silico* are shown as bars for different primer sets and

types of transcripts. **A)** Amplification where only primers targeting the V gene segment

within the variable region from S, W, R, C, and L are tested with the total number of

primers (after enumerating primers with degenerate nucleotides) shown above the bars. **B)**

Amplification using forward (V gene) and reverse (J gene segment or constant region)

primers from S, W, R, C, and L. **C)** Amplification where only primers targeting the constant

region from S, W, and C are tested with the total number of primers (after enumerating

primers with degenerate nucleotides) shown above the bars. R and L are not shown here as

their reverse primers target J gene segments. **D)** Analysis of 10x B and T cell V(D)J inner

and outer reverse primers. The different groups of transcripts (Ig and TCR) are noted below

the bar plot.

Fig 6. Comparison of human and rhesus IGHA reference sequences and illustration of

human 10x primer target regions. A black H and red R, respectively highlight human and

rhesus sequences on the left side of each panel. **A)** Multiple sequence alignment of putative

human IGHA1 (J00220) and IGHA2 (J00221) CDSs with rhesus consensus cDNA coding

sequences recovered (IGHA*01, IGHA*02, and IGHA*03) in either secreted (s) or

membrane-bound (b) form. The target sites of human 10x primers are denoted by red

boxes. **B-C)** Zoom in of the inner and outer primer target sites, where mismatches between

the primer and rhesus consensus sequence(s) are denoted by red boxes.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

	Rectal Biopsy	Whole Blood	PBMC	Lymph Node	Total
IGHA*01	1411 (8)	67 (0)	21 (2)	224 (0)	1723 (10)
IGHA*02	705 (4)	7 (0)	21 (1)	275 (6)	1008 (11)
IGHA*03	101	0	5	9	115
IGHD	0	12	4	3	19
IGHE	1	1	0	4	6
IGHG1	78	21	21	1030	1150
IGHG2	9	9	4	311	333
IGHG3	0	1	0	30	31
IGHG4	1	0	0	13	14
IGHM	129 (3)	20 (21)	18 (14)	169 (8)	336 (46)
IGK	97	30	12	155	294
IGL	100	30	18	140	288
Total	2632 (15)	198 (21)	124 (17)	2363 (14)	5317 (67)

Table 1. Number of unique full-length rhesus IG transcript sequences identified in each tissue. Counts for secreted heavy chain Ig sequences (IGH A, D, E, G1-4, and M) are shown with membrane-bound counts in parenthesis where applicable. IGHA is further classified by allotype, conventionally denoted by an asterisk followed by a number.

	Rectal Biopsy	Whole Blood	PBMC	Lymph Node	Total
TCRA	14	71	103	267	455
TCRB1	1	9	11	26	47
TCRB2	0	13	16	44	73
TCRD	1	1	27	10	39
TCRG1	2	3	6	5	16
TCRG2	3	11	75	22	111
Total	21	108	238	374	741

Table 2. Number of unique full-length rhesus TCR transcript sequences identified in each tissue.