Title

A native prokaryotic voltage-dependent calcium channel with a novel selectivity filter

sequence

Authors

Takushi Shimomura[1,3], Yoshiki Yonekawa[2], Hitoshi Nagura[1], Michihiro Tateyama[3],

Yoshinori Fujiyoshi[1‡,4] and Katsumasa Irie[1,2].

[1]Cellular and Structural Physiology Institute (CeSPI), and [2]Graduate School of

Pharmaceutical Sciences, Nagoya University, Furo-cho, Chikusa, Nagoya 464-8601,

Japan.

[3]Division of Biophysics and Neurobiology, National Institute for Physiological Sciences,

Okazaki, Aichi, 444-8585, Japan

[4]CeSPIA Inc., 2-1-1, Otemachi, Chiyoda, Tokyo, 100-0004, Japan

‡present address: Advanced Research Institute, Tokyo Medical and Dental University,

1-5-45 Yushima, Bunkyo-ku, Tokyo 113 - 8510, Japan

Address correspondence to: Katsumasa Irie, Cellular and Structural Physiology Institute

(CeSPI), Nagoya University, Furo-cho, Chikusa, Nagoya 464-8601, Japan Tel.:

+81-52-747-6838; Fax: +81-52-747-6795; E-mail: kirie@cespi.nagoya-u.ac.jp

Abstract

Voltage-dependent Ca^{2+} channels (Cavs) are indispensable for coupling action potentials with Ca^{2+} signaling in living organisms. The structure of Cavs is similar to that of voltage-dependent Na^+ channels (Navs). It is known that prokaryotic Navs can obtain Ca^{2+} selectivity by negative charge mutations of the selectivity filter, but native prokaryotic Cavs had not yet been identified.

Here, we report the first identification of a native prokaryotic Cav, CavMr, and its relative, NavPp. Although CavMr contains a smaller number of negatively charged residues in the selectivity filter than artificial prokaryotic Cavs, CavMr exhibits high Ca^{2+} selectivity. In contrast, NavPp, which has similar filter sequence to artificial Cavs, mainly allows Na^+ to permeate. Interestingly, a NavPp mutant whose selectivity filter was replaced with that of CavMr exhibits high Ca^{2+} selectivity. Mutational analyses revealed that the glycine residue of the CavMr selectivity filter is a determinant for Ca^{2+} selectivity. This glycine residue is well conserved among subdomains I and III of eukaryotic Cavs.

These findings provide new insight into the Ca^{2+} selectivity mechanism conserved from prokaryotes to eukaryotes.

Introduction

Voltage-dependent Ca^{2+} channels (Cavs), which couple the membrane voltage with Ca^{2+} signaling, regulate some important physiological functions, such as neurotransmission and muscle contraction (Hille, 2001). The channel subunits of mammalian Cavs as well as mammalian voltage-dependent Na^+ channels (Navs) have 24 transmembrane helices (24TM) (Catterall, 2000), and comprise 4 homologous subdomains with 6 transmembrane helices that correspond to one subunit of homo-tetrameric channels, such as Kvs and prokaryotic Navs (BacNavs). Comparison of the sequences between Navs and Cavs indicate that Navs derived from Cavs. Their two pairs of subdomains, domains I and III, and domains II and IV, are evolutionarily close to each other (Rahman et al., 2014; Strong et al., 1993). Therefore, the 24TM-type of Cavs are thought to have evolved from the single-domain type of Cavs with two times of domain duplications. Although prokaryotes are expected to have such ancestor-like channels, native prokaryotic Cavs have not yet been identified. The lack of ancestor-like prokaryotic Cavs is a missing link in the evolution of voltage-dependent cation channels.

In contrast to the lack of prokaryotic Cavs, many BacNavs have been characterized (Irie et al., 2010; Ito et al., 2004; Koishi et al., 2004; Lee et al., 2012; Nagura et al., 2010; Ren et al., 2001; Shimomura et al., 2016, 2011). The simple structure of BacNavs has helped to elucidate the molecular mechanisms of Nav (Irie et al., 2018, 2012; Tsai et al., 2013; Yue et al., 2002). In addition to this point, BacNavs has been used as the genetic tool for manipulating the neuronal excitability in vivo (Bando et al., 2014; Kamiya et al., 2019; Lin et al., 2010). The introduction of a several negatively charged amino acids into the selectivity filter of BacNavs leads to the acquisition of Ca^{2+} selectivity (Tang et al., 2013; Yue et al., 2002). Such a mutant of NavAb (a BacNav from *Arcobacter butzleri*) showed high Ca^{2+} selectivity, and the structural basis of Ca^{2+} selectivity has been discussed on the basis of its crystal structures (Tang et al., 2016, 2013). The selectivity filter sequences with a large number of aspartates in CavAb, which was made by mutations of NavAb, are quite different from those of the original mammalian Cavs.

Here, we newly characterized two BacNav homologues, CavMr from *Meiothermus ruber* and NavPp from *Plesiocystis pacifica*. These two channels are evolutionarily distant from the previously reported canonical BacNavs. We confirmed that CavMr has clear Ca^{2+} selectivity, and NavPp has Na^+ selectivity with Ca^{2+}-dependent inhibition. The discovery of these channels suggests the possible importance of voltage-regulated Ca^{2+} signaling in prokaryotes and may be a new genetic tool for controlling Ca^{2+}

signaling. Furthermore, mutational analyses indicate that the glycine residue of the
CavMr selectivity filter is important for Ca^{2+} selectivity. The glycine residue is also well
conserved in the selectivity filter of the subdomain I and III of mammalian Cavs. On the
basis of these observations, we propose that CavMr is an ancestral-type of native
prokaryotic Cav with a Ca^{2+} selectivity mechanism different from that in artificial
CavAb. CavMr and NavPp are expected to advance our understanding of Ca^{2+}
recognition and the evolution of voltage-dependent cation channels.
8
Results
Identification of two prokaryotic channels with Ca^{2+} permeability and inhibition.
We searched for the primary sequences of prokaryotic Cavs in the GenBank[TM]
database. In mammalian and prokaryotic Navs and Cavs, a larger number of negative
charges in the filter increases Ca^{2+} selectivity (Heinemann et al., 1992; Tang et al., 2014;
Yue et al., 2002). Several BLAST search rounds using the pore regions (S5-S6) of
NaChBac (or NavBh; a BacNav from *Bacillus halodurans*) as templates revealed a
series of candidate prokaryotic Cavs (Fig.1a) with a selectivity filter sequence similar to
the "TLESW" motif, but more negatively charged: ZP_04038264 from *Meiothermus*
ruber, ZP_01909854 from *Plesiocystis pacifica*, YP 003896792_from *Halomonas*
elongata, and YP_003073405 from *Teredinibacter turnerae* (*SI Appendix*, Fig.S1a).
These channels belong to a different branch of the phylogenic tree than that of canonical
BacNavs (Fig.1a) and have some negatively charged residues in their selectivity filter,
similar to CavAb (Fig.1b). In the case of the expression of prokaryotic channels, insect
cells are better than mammalian cells to generate large current amplitudes (Irie et al.,
2018). We therefore transfected Sf9 cells with these genes and measured the resulting
whole-cell currents. Although the cells transfected with genes from *H. elongata* and *T.*
turnerae failed to show detectable currents, those from *M. ruber* and *P. pacifica* showed
currents in response to a depolarizing stimulus from a -140 mV holding potential
(Fig.1c and d). To estimate the Ca^{2+} permeability, we measured their current-voltage
relationships. The *M. ruber* channel had clearly larger currents in the high-Ca^{2+} solution
than in the high-Na^+ solution, and very positive reversal potential was observed in a
high-Ca^{2+} bath solution (Fig.1e). In contrast, the currents derived from the *P. pacifica*
channel increased with increases in the bath Na^+ concentration and significantly
decreased when the Na^+ solution was replaced with a high Ca^{2+} solution. The reversal
potential fit well to the Na^+ equilibrium potential in the high-Na^+ solution (Fig.1f).
These current-voltage relationships suggest that the *M. ruber* channel has a preference
for Ca^{2+} and the *P. pacifica* channel has a preference for Na^+. Therefore, the two newly

identified channels from *M. ruber* and *P. pacifica* are abbreviated as CavMr and NavPp

respectively, based on their ion selectivity and species name.

3
Figure 1. Sequence analysis and the representative current recordings of the
novel BacNav homologues
a). Phylogenetic tree of the BacNav homologues with their GenBank[TM] accession
numbers. The ClustalW program was used to align the multiple protein sequences of
the BacNav homologues. The phylogenetic tree was generated using "PROTDIST",
one of the PHYLIP package (Phylogeny Inference Package:
http://evolution.genetics.washington.edu/phylip.html). The branch lengths are
proportional to the sequence divergence, with the scale bar corresponding to 0.1
substitution per amino acid position. Four homologues colored that are not included in
canonical BacNavs were cloned and expressed to check the current activity. Those of
two which are underlined in red and shown as bold generated the detectable currents.
b). Schematic secondary structure and selectivity filter sequence of BacNavs and
human Cavs. Cylinder indicates α-helix. The selectivity filter sequences are indicated
by alphabetical characters. Negatively charged residues are colored by red. Glycine
residues in the position 4 are colored by cyan. The straight lines indicate the other part
of pore domain. The selectivity filter sequence of hCav1.1 (UniProt ID: Q13698),
hCav2.1 (O00555) and hCav3.1 (O43497), were used. c and d). Representative current
traces to obtain the current-voltage relationships of CavMr (c) and NavPp (d) in Sf9
cells. The straight lines indicating the zero-current level in the representative current
traces. Currents were generated under the bath solutions containing high Na$^+$ (top) and
high Ca^{2+} (middle), by a series of step-pulses shown in bottom. e and f).
Current-voltage relationships of CavMr (e) and NavPp (f) measured under the
different bath solutions (filled black; 150 mM NaCl, open black; 75 mM NaCl and 75
mM NMDG-HCl, open red; 75 mM NaCl and 50 mM CaCl$_2$, filled red; 50 mM CaCl$_2$
and 75 mM NMDG-HCl). Currents of CavMr and NavPp were normalized to that by
0mV depolarization stimuli under 75 mM NaCl and 50 mM CaCl$_2$ bath solution and
150 mM NaCl bath solution, respectively.

To clearly compare the positions of the residues in the selectivity filter in each channel,

we renumbered the seven residues comprising the selectivity filter in the following

description. For example, the seven residues of the CavMr selectivity filter are

183-TLEGWVD-189, and thus Thr183 and Asp189 were renumbered to Thr1 and Asp7

(Fig.1b). Notably, the amino acid sequence of the selectivity filter in CavMr is similar to

the conserved features of domains I/III in mammalian Cavs, a glycine at position 4 and

a polar or negatively charged residue at position 7 (Fig.1b), which are not observed in

the BacNav family. In addition, its sequence is quite similar to that of the human Cav

subdomain I, or even the same as Cav3.1 and 3.2 (Fig.1b).

In the following experiments, to accurately evaluate the reversal potential for the ion

selectivity analysis, we introduced a single mutation that resulted in large and

long-lasting channel currents. T220A and G229A mutations in NaChBac led to slower

inactivation, indicating suppression of the transition to the inactivated state (Irie et al.,

2010; Shimomura et al., 2016). We introduced a T232A mutation to NavPp and a

G240A mutation to CavMr, corresponding to the NaChBac mutations of T220A and

G229A, respectively. These mutants stably showed measurable currents, even after they

were administrated multiple depolarizing stimuli (*SI Appendix*, Fig.S1b-e).

Fig. S1. NavPp T232A and CavMr G240A mutants suppress inactivation
a). Alignment of the deduced amino acid sequences of P1 helix to P2 helix domain of novel cloned homologues with well characterized BacNavs. b) and c). Whole-cell current in CavMr wild type (WT; b) and a G240A mutant (c) when the pulse of -20 mV was given for 500 ms and 1sec, respectively. d) and e). Whole-cell recording of NavPp wild type (WT; d) and NavPp T232A mutant (e) when the pulse of -10 mV was given for 250 ms and 500ms, respectively.

3   **<u>CavMr has high Ca^{2+} selectivity over Na^{+}.</u>**

Figure 2. Cation selectivity of CavMr

4
Figure 2. Cation selectivity of CavMr
a and b). Recordings of the reversal potential of CavMr currents using the ramp
protocol. Currents were generated by the step pulse of -20 mV from -140 mV holding
potential, followed by the ramp pulses with different voltage values (shown at the
bottom of panels a and b). The values of the reversal potential recorded with three
different ramp pulses were averaged. Currents were measured under the bath solution
containing 4 mM (a) and 10 and 20 mM (b) CaCl$_2$ and the pipette solution with 150
mM NaCl. c). The plot of the reversal potential to the bath [Ca^{2+}]$_{out}$. Each value was
obtained using the protocol shown in a and b. The relationship was fitted by a line
with the slope of 41.07±2.64 mV per decade ($n = 4$). d). Representative current traces
to obtain the reversal potential under the condition of 100 mM [Sr^{2+}]$_{out}$ and 150 mM
[Na^{+}]$_{in}$. Currents were generated by the protocol shown in the lower part. e).
Representative current traces to investigate the P_{Cs}/P_{Ca} and P_K/P_{Ca}, the pipette
solutions contained 150 mM [Cs^{+}]$_{in}$ for P_{Cs}/P_{Ca} and 150 mM [K^{+}]$_{in}$ for P_K/P_{Ca}, while
the bath solution contained 10 mM [Ca^{2+}]$_{out}$ in both cases. f). The relative permeability
of Ca^{2+} or Sr^{2+} to Na^{+} in CavMr, calculated from the reversal potential that was
obtained in a, b, and d. g). The relative permeability of each monovalent cation to Ca^{2+}
in CavMr, derived from the data shown in e.

We accurately quantified the selectivity of CavMr for Na^{+} and Ca^{2+} (P_{Ca}/P_{Na}) by

measuring the reversal potential under bi-ionic conditions, in which the Ca^{2+}

concentration in the bath solution was changed to 1.5, 4, 10, 20, and 40 mM while the

intracellular Na^{+} concentration was held constant at 150 mM(Fig.2a and b). The plot of

the reversal potentials as a function of [Ca^{2+}] had a slope of 41.07±2.64 mV /decade (n

= 4), a value close to that for Ca^{2+} (Fig. 2c), and indicated that CavMr had a P_{Ca}/P_{Na} of

218 ± 38 (Fig.2a and b, Table1). This high P_{Ca}/P_{Na} value is comparable to that of CavAb.

Among several species of cations, including Sr^{2+}, K^{+}, and Cs^{+} (Fig.2d and e), Ca^{2+} had

the highest permeability relative to Na^+ (Fig.2f and g, Table1). On the basis of these

results, CavMr was confirmed to be a native prokaryotic Cav with high Ca^{2+} selectivity.

We also investigated whether CavMr shows the typical anomalous mole fraction effect

(Almers and McCleskey, 1984) and the non-monotonic mole fraction effect observed in

NaChBac (Finol-Urdaneta et al., 2014). CavMr did not allow Na^+ permeation under

Ca^{2+}-free (0 mM $CaCl_2$ and 1 mM EGTA) conditions (*SI Appendix*, Fig. S2a and b).

Also, different from the recording of NaChBac currents in a solution containing Na^+ and

K^+, CavMr had an apparently monotonic current increase depending on the Ca^{2+} mole

fraction to Na^+ (*SI Appendix*, Fig. S2c and d).

Fig. S2. The characterization of the selectivity filter of CavMr

a). No observation of anomalous mole fraction effects in CavMr. CavMr currents

were recorded in the bath solution containing the following concentration of Na^+ and

Ca^{2+}; [Na^+]:[Ca^{2+}] is 0:90, 133.7:0.9, 135:0 (mM), respectively. The 0 mM Ca^{2+}

solution also contains 1 mM EGTA. b). The plot of the normalized current amplitude

of CavMr obtained from a. c). Representative current traces of CavMr under different

mole fraction of Ca^{2+}. f_{Ca} indicates [Ca^{2+}]$_{out}$ / ([Ca^{2+}]$_{out}$ + [Na^+]$_{out}$). d). The plot of the

normalized current amplitude to each mole fraction, as measured in c). e). For the

evaluation of the relative permeability of Ca^{2+} and Sr^{2+} to Na^+ of CavMr-D7M, Ca

solution (100mM $CaCl_2$, 10mM HEPES (pH 7.4 adjusted with $Ca(OH)_2$) and 10 mM

glucose) and Sr solution (100 mM $SrCl_2$ 10 mM HEPES (pH 7.4 adjusted by $Sr(OH)_2$)

and 10 mM glucose) were used as bath solution, respectively. High Na pipette solution

(115 mM NaF, 35 mM NaCl, 10 mM EGTA, and 10 mM HEPES (pH 7.4 adjusted by

NaOH)) was used. Currents were generated by the step pulse of -20 mV from -140 mV

holding potential, followed by the ramp pulses with different voltage values. The time

courses of the membrane potentials were shown at the bottom of each current traces. f)

The relative permeability of divalent cation to Na^+ in the CavMr-D7M, whose position

6 residue of the selectivity filter was neutralized by the corresponding residue of
NavAb.

Studies of an artificial Cav, CavAb, revealed that Ca^{2+} selectivity depends on a large

number of aspartates in the filter sequence (Tang et al., 2013). The high Ca^{2+} selectivity

in CavMr was unexpected because the filter sequence contained only one aspartate

residue (Fig.1b). Furthermore, CavMr-D7M, which has only one negatively charged

residue in the selectivity filter "TLEGWVM", still had high Ca^{2+} selectivity comparable

to that of wild-type CavMr ($P_{Ca}/P_{Na} = 144 \pm 12$; *SI Appendix*, Fig. S2e and f and Table1).

These findings indicate that CavMr and artificial CavAb have different Ca^{2+} selective

mechanisms.

	P_{Ca}/P_{Na}			P_{Sr}/P_{Na}			P_{K}/P_{Na}			P_{Cs}/P_{Na}		
CavMr G240A	218	±	38	40.6	±	3.4	0.0036±0.00072 [a]			0.0021±0.00042 [b]		
	(*n*=	20	)	(*n*=	6	)	(*n*=	4	)	(*n*=	4	)
Pp	13.8	±	2.0	24.5	±	0.3	0.95	±	0.04	0.57	±	0.05
	(*n*=	7	)	(*n*=	5	)	(*n*=	4	)	(*n*=	3	)
G4D	7.73	±	2.24	18.6	±	6.1	1.20	±	0.28	0.87	±	0.21
	(*n*=	11	)	(*n*=	4	)	(*n*=	4	)	(*n*=	4	)
G4S	11.9	±	1.5	4.23	±	0.27	1.54	±	0.12	2.02	±	0.48
	(*n*=	5	)	(*n*=	5	)	(*n*=	5	)	(*n*=	3	)
V6T	40.1	±	9.7	13.3	±	2.5	0.69	±	0.26	0.54	±	0.60
	(*n*=	5	)	(*n*=	5	)	(*n*=	3	)	(*n*=	3	)
D7M	144	±	12	20.7	±	2.7	N.D.			N.D.		
	(*n*=	5	)	(*n*=	5	)						
NavPp T232A	0.308±0.028			0.38±0.027			0.16±0.026			0.0052±0.0006		
	(*n*=	10	)	(*n*=	9	)	(*n*=	9	)	(*n*=	7	)
Mr	215	±	33	86.3	±	12.2	0.0045±0.00072 [a]			0.0135±0.0039 [b]		
	(*n*=	7	)	(*n*=	4	)	(*n*=	4	)	(*n*=	9	)
D4G	41.4	±	6.7	8.85	±	0.95	<<	0.01		<<	0.01	
	(*n*=	10	)	(*n*=	4	)	(*n*=	3	)	(*n*=	4	)
D4S	1.49	±	0.18	0.47	±	0.08	0.24	±	0.01	0.11	±	0.02
	(*n*=	6	)	(*n*=	6	)	(*n*=	4	)	(*n*=	3	)
T6V	1.72	±	0.19	33.9	±	5.0	0.99	±	0.03	0.84	±	0.02
	(*n*=	10	)	(*n*=	8	)	(*n*=	4	)	(*n*=	4	)

11 **Table1. Relative permeability of CavMr and NavPp**
12 [a] Because of high Ca^{2+} selectivity, P_{K}/P_{Ca} were indicated
13 [b] Because of high Ca^{2+} selectivity, P_{Cs}/P_{Ca} were indicated

NavPp is permeable to Na⁺ and is blocked by extracellular Ca²⁺.

The current-voltage relationships of NavPp revealed a preference for Na^+ (Fig.1d and
f). Interestingly, NavPp, although having one more aspartate in the selectivity filter than
CavMr, exhibited larger currents in Na^+ solutions than in Ca^{2+} solutions (Fig.1b and d).
Recordings in bath solution containing both Na^+ and Ca^{2+} demonstrated that increasing
the extracellular Ca^{2+} decreased the current in NavPp and led to a positive shift in the
voltage dependence, indicating that a higher concentration of Ca^{2+} inhibited NavPp
(Fig.1f and Fig. 3a and *SI Appendix*, Fig. S3a). With NavAb, increasing the Ca^{2+}
concentration with a constant Na^+ concentration in the bath solution led to a small
increase in the current amplitude, probably due in part to Ca^{2+} permeability (Fig. 3a and
SI Appendix, Fig. S3b). We also investigated the dependence of the direction of current
flow on Ca^{2+} inhibition by comparing pipette solutions containing 10 mM or 150 mM
Na^+, in which the current flowed in an inward or outward direction, respectively, even
under the same -10‑mV depolarizing stimulus (Fig. 3a and *SI Appendix*, Fig. S3a and
c). The results demonstrated that the inhibitory effects of Ca^{2+} on NavPp were
independent of the current direction.

**Figure 3. Na⁺ permeability and the extracellular Ca²⁺-dependent inhibition in
NavPp**

a) $[Ca^{2+}]_{out}$ dependent inhibitory effects in NavPp. Currents were normalized to those
under 1.5 mM $[Ca^{2+}]_{out}$(filled square; NavAb current in the 30 mM $[Na^+]_{out}$ and 10 mM
$[Na^+]_{in}$ (representative data are shown in Fig. S3b), open triangle down; NavPp inward
current in the 30 mM $[Na^+]_{out}$ and 10 mM $[Na^+]_{in}$ (Fig. S3a), open triangle up; NavPp
outward current in the 30 mM $[Na^+]_{out}$ and 150 mM $[Na^+]_{in}$ (Fig. S3c), filled triangle
up; NavPp outward current in the 0 mM $[Na^+]_{out}$ and 150 mM $[Na^+]_{in}$. b).
Representative currents to obtain the reversal potential in NavPp for P_{Ca}/P_{Na}. Currents
were generated by the ramp protocol shown in bottom. $[Ca^{2+}]_{out}$ was varied from 10 to
60 mM with the fixed $[Na^+]$ both in the bath (60mM) and pipette (150mM). c).
Representative currents to obtain the reversal potential in NavPp for P_{Sr}/P_{Na}. Currents
were generated by the ramp protocol shown in bottom. $[Sr^{2+}]_{out}$ was 40 mM with the
fixed $[Na^+]$ both in the bath (40mM) and pipette (150mM). d). The relative
permeability of different cation species to Na^+ in NavPp, calculated from the reversal
potential that was obtained in b, c and Fig.S3d and e.

We then compared the relative permeability of various cations with that of Na^+ in

NavPp. In bi-ionic conditions with high concentration of Ca^{2+} in the bath and high Na^+
in the pipette, the outward current was completely blocked by Ca^{2+} (Fig. 3a), and the
inward current was hardly observed. The reversal potential was obtained under an
extracellular solution containing Na^+ ions, however, despite a partial Ca^{2+} or
Sr^{2+}-induced block (Fig. 3b and c). The selectivity of NavPp was higher for Na^+ than for
Ca^{2+}, Sr^{2+}, K^+, and Cs^+ (Fig. 3d and *SI Appendix*, Fig. S3d and e). The P_{Ca}/P_{Na} was
0.308 ± 0.028 in a bath solution containing both Ca^{2+} and Na^+, suggesting that a larger
fraction of Ca^{2+} is allowed to permeate with outside Na^+ ions through NavPp than
through canonical BacNavs. Similar to Ca^{2+}, Sr^{2+} also blocked the NavPp current, but
may also permeate the channel along with Na^+ ions (Fig. 3c). These findings
demonstrate a unique feature of NavPp, a low affinity Ca^{2+} block, which is not reported
in canonical BacNavs.

13
Fig. S3. The characterization of the selectivity filter of NavPp
a) and b) NavPp and NavAb currents generated by the step pulses from -140 mV
holding potential to -10 mV under the various $[Ca^{2+}]_{out}$ ranging from 1.5 to 80 mM.
Inward currents are observed in the 30 mM $[Na^+]_{out}$ and 10 mM $[Na^+]_{in}$. c) NavPp
currents generated by the same pulse protocol under various $[Ca^{2+}]_{out}$. Outward
currents are observed in the 30 mM $[Na^+]_{out}$ and 150 mM $[Na^+]_{in}$. d and e) For the
evaluation of the relative permeability of K^+ and Cs^+ to Na^+ of NavPp, K solution (150
mM KCl, 2mM $CaCl_2$ 10 mM HEPES (pH 7.4 adjusted by KOH) and 10 mM glucose)
and low Na solution (50 mM NaCl, 100mM NMDG-HCl, 2mM $CaCl_2$ 10 mM HEPES
(pH 7.4 adjusted by NaOH) and 10 mM glucose) were used as bath solution,
respectively. High Na pipette solution (115 mM NaF, 35 mM NaCl, 10 mM EGTA,
and 10 mM HEPES (pH 7.4 adjusted by NaOH)) and Cs pipette solution (115 mM CsF,
35 mM CsCl, 10 mM EGTA, and 10 mM HEPES (pH 7.4 adjusted by CsOH)) were
used for K^+ and Cs^+ selectivity, respectively. Currents were generated by the step pulse

of -20 mV from -140 mV holding potential, followed by the ramp pulses with different
voltage values. The time courses of the membrane potentials were shown at the
bottom of each current traces. f) The relative permeability of each cation species to
Na$^+$ in the single-point mutants of NavPp. The selectivity filter of NavPp was changed
to the Ca^{2+}-selective canonical-BacNavs mutants (T6S; TLEDWSD and T6A;
TLEDWAD).

Interestingly, the filter sequence of NavPp, "TLEDWTD", has three
negatively-charged residues, similar to the filter sequences of the artificial
Ca^{2+}-selective BacNav mutant ("TLEDWSD" mutant of NavAb and "TLEDWAD"
mutant of NaChBac) (Tang et al., 2013; Yue et al., 2002). NavPp does not show Ca^{2+}
permeability, however, but rather a Ca^{2+} block. We also investigated NavPp mutants
with the same filter sequences as the artificial Cavs. NavPp-T6S "TLEDWSD"
exhibited Ca^{2+}-blocked currents similar to wild-type NavPp (*SI Appendix*, Fig. S3f).
Further, NavPp-T6A "TLEDWAD" showed no inward current in bath solutions
containing divalent cations, suggesting that the Ca^{2+}-induced block was enforced.
Therefore, both of the selectivity filter sequences providing Ca^{2+} selectivity to canonical
BacNavs failed to generate Ca^{2+}-permeable NavPp, indicating that the cation permeable
mechanism of NavPp differs from that of canonical BacNavs, as well as that of CavMr.
19

20 **Swapping the filter regions between CavMr and NavPp revealed the importance of**
21 **the glycine residue at position 4 for Ca^{2+} selective permeation.**
22

Figure 4. The cation selectivity of the swapping mutant channels in their selectivity filter
between CavMr and NavPp

a) Amino acid sequences of the selectivity filter in the swapped mutants, CavMr-Pp, Nav-Pp,
NavAb_Mr, and CavMr_Ab. The selectivity filter sequences of CavMr, and NavPp, and NavAb
are indicated by alphabetical characters with cyan, red, and gray shade, respectively. Negatively
charged residues are colored by red. Glycine residues are colored by cyan. The straight lines of
cyan, red, and black indicates the other part of pore domain of CavMr, NavPp, and NavAb,
respectively. b) Pore domains of crystal structure of NavAb (PDB code:5YUA). The selectivity
filter, which corresponds to the sequences shown in a, was indicated in red. c) The relative
permeability of divalent cations to Na^+ (left) and that of monovalent cations to Ca^{2+} (right) in
NavPp-Mr. d) The relative permeability of different cation species to Na^+ in CavMr-Pp.

To search for the determinants of Ca^{2+} selectivity in CavMr, we investigated a series of
mutants in which the filter regions were swapped between CavMr and NavPp (Fig.4a
and b). The mutants with filter sequences swapped between NavPp and CavMr
exhibited channel activity (*SI Appendix*, Fig. S4). A NavPp mutant whose selectivity
filter was replaced with that of CavMr, named NavPp-Mr, exhibited much higher Ca^{2+}
selectivity ($P_{Ca}/P_{Na} = 215 \pm 33$) as well as high Sr^{2+} selectivity, comparable to that of
CavMr (Fig. 4c). In addition, NavPp-Mr excluded Cs^+ similar to CavMr, but weakly
allowed K^+ permeation in contrast to CavMr. On the other hand, a CavMr mutant whose
selectivity filter was replaced with that of NavPp (CavMr-Pp) almost lost its Ca^{2+}
selectivity ($P_{Ca}/P_{Na} = 13.8 \pm 2.0$), and was less able to discriminate Cs^+ and K^+ from
Na^+ (Fig. 4d). That is, CavMr-Pp was a more non-selective channel than the wild-type
CavMr, rather than a Na^+-selective channel. Namely, the Ca^{2+} selectivity (from NavPp
to CavMr) was almost transferable, but the Na^+ selectivity was not. We also investigated
the full swapping of the filter sequences between CavMr and NavAb (Fig. 4a), but
neither swapped mutant of CavMr and NavAb had detectable currents. This finding
suggested that CavMr and NavAb achieve cation selectivity by different structural
backbones and mechanisms.

Fig. S4. Representative current traces of the ramp pulse of selectivity filter-swapped mutants, CavMr_Pp and NavPp_Mr
 For the evaluation of the relative permeability of Ca^{2+}, Sr^{2+}, K^+ and Cs^+ to Na^+, Ca solution (100mM $CaCl_2$, 10mM HEPES (pH 7.4 adjusted with $Ca(OH)_2$) and 10 mM glucose), Sr solution (100 mM $SrCl_2$, 10 mM HEPES (pH 7.4 adjusted by $Sr(OH)_2$) and 10 mM glucose), K solution (150 mM KCl, 2mM $CaCl_2$, 10 mM HEPES (pH 7.4 adjusted by KOH) and 10 mM glucose) and Cs solution (150 mM CsCl, 2mM $CaCl_2$, 10 mM HEPES (pH 7.4 adjusted by CsOH) and 10 mM glucose) were used as bath solution, respectively. High Na pipette solution (115 mM NaF, 35 mM NaCl, 10 mM EGTA and

10 mM HEPES (pH 7.4 adjusted by NaOH)) was used. In the case of NavPp-Mr, the relative permeability of K^+ and Cs^+ to Ca^{2+} were evaluated in behalf of the relative permeability to Na^+ because of high Ca selectivity of NavPp-Mr. For the evaluation, high K pipette solution (115 mM KF, 35 mM KCl, 10 mM EGTA and 10 mM HEPES (pH 7.4 adjusted by KOH)) and high Cs pipette solution (115 mM CsF, 35 mM CsCl, 10 mM EGTA and 10 mM HEPES (pH 7.4 adjusted by CsOH)) were used. As bath solution, Ca solution (100mM $CaCl_2$, 10mM HEPES (pH 7.4 adjusted with $Ca(OH)_2$) and 10 mM glucose was used. Currents were generated by the step pulse of -20 mV from -140 mV holding potential, followed by the ramp pulses with different voltage values. The time courses of the membrane potentials were shown at the bottom of each current traces.

Positions 4 and/or 6 of the filter sequences are thought to be important for Ca^{2+}-selective permeation in NavPp-Mr and CavMr, because only these two positions were mutated in the swapping experiments. We investigated which of the mutations in positions 4 and 6 had greater effects on the loss of and acquisition of Ca^{2+} selectivity in CavMr and NavPp, respectively. In CavMr, both of two single mutants, CavMr-G4D and CavMr-V6T, decreased Ca^{2+} selectivity and allowed K^+ and Cs^+ permeation (Fig. 5a, *SI Appendix*, Fig.S5). Especially, the mutational effect was greater in CavMr-G4D, whose P_{Ca}/P_{Na} was less than 10 (7.73 ± 2.24). CavMr-G4S, in which Gly4 was replaced with the Ser4 of NavAb, also exhibited lower Ca^{2+} selectivity (P_{Ca}/P_{Na} = 11.9 ± 1.5) and was also K^+ and Cs^+ permeable, indicating that even a minor substitution by serine is not tolerable and does not allow for the selection of specific cations (Fig. 5b, Table1, and *SI Appendix*, Fig. S6). In the case of NavPp, NavPp-D4G acquired Ca^{2+} selectivity over Na^+, and also showed a greater exclusion to K^+ and Cs^+ than wild-type NavPp (Fig. 5c and *SI Appendix*, Fig.S5). In contrast, NavPp-T6V failed to acquire the high Ca^{2+} selectivity (P_{Ca}/P_{Na} = 1.72 ± 1.09) and also allowed K^+ and Cs^+ permeation, while it had relatively high Sr^{2+} selectivity. These results indicate that, in both CavMr and NavPp, a glycine residue at position 4 is a key determinant for Ca^{2+} selectivity. It is noteworthy that the glycine is a conserved residue at position 4 of subdomains I and III in all subtypes of mammalian Cavs (Fig.1b).

Figure 5. The single point mutations losing and obtaining Ca^{2+} selectivity of CavMr and NavPp, respectively.

a) The relative permeability of each cation species to Na^+ in the single-point mutants of CavMr. The selectivity filter of CavMr was changed to the corresponding residues of NavPp at position 4 (G4D) and position 6 (V6T), respectively b) The relative permeability of each cation species to Na^+ in the G4S mutants of CavMr, whose position 4 residue of the selectivity filter was mutated to the corresponding residue of canonical BacNavs. c) The relative permeability of each cation species to Na^+ in the single-point mutants of NavPp. The selectivity filter of NavPp was changed to the corresponding residues of CavMr at position 4 (D4G) and position 6 (T6V), respectively.

Fig. S5. Representative current traces of the ramp pulse of single point-swapped mutants between CavMr

To evaluate the relative permeability of Ca^{2+}, Sr^{2+}, K^+ and Cs^+ to Na^+, Ca solution, Sr solution, K solution and Cs solution were used as the bath solution, respectively. High Na pipette solution was used as the pipette solution. The solution contents were described in material method and Fig. S4. The time courses of the membrane potentials were shown at the bottom of each current traces.

1
Fig. S6. Representative current traces of the ramp pulse of single point-swapped
mutants of NavPp
To evaluate the relative permeability of Ca^{2+}, Sr^{2+}, K^+ and Cs^+ to Na^+, Ca solution, Sr
solution, K solution and Cs solution were used as the bath solution, respectively. High
Na pipette solution was used as the pipette solution. The solution contents were
described in material method and Fig. S4. The time course of the membrane potentials
was shown at the bottom of each current traces.
9

DISCUSSION

A native prokaryotic voltage-dependent Ca^{2+} channel has a unique Ca^{2+} selective

mechanism.

In this study, we newly characterized two prokaryotic voltage-dependent cation

channels, CavMr and NavPp. CavMr is the first native prokaryotic Cavs reported

despite its BacNav-like "TxExW" motif, and NavPp could be inhibited by high

concentrations of extracellular Ca^{2+}. The P_{Ca}/P_{Na} of CavMr was more than 200 (Fig. 2e

and Table 1), comparable to that of CavAb, an artificial Ca^{2+} channel. Anomalous mole

fraction effects were not observed in CavMr (*SI Appendix*, Fig. S2a and b), suggesting

that CavMr has a very high affinity for Ca^{2+}. In addition to providing new insights about

general Ca^{2+}-selective mechanisms, CavMr has the potential to be a new genetic tool for

upregulating calcium signaling, as BacNavs are useful genetic tools for increasing

action potential firing in mice (Bando et al., 2014; Kamiya et al., 2019; Lin et al., 2010).

Phylogenetic analysis demonstrated that CavMr and NavPp are similar to each other,

but distant from canonical BacNavs (Fig. 1a). The high Ca^{2+} selectivity of CavMr was

transferable to NavPp. Intriguingly, two pairs of mutants with the same selectivity filter

(CavMr-G4D and NavPp-T6V, CavMr-V6T and NavPp-D4G) showed a very similar

tendency with regard to both the order and extent of cation selectivity (Fig. 5a and c).

Therefore, the basic overall architecture of the NavPp selectivity filter could be similar

to that of CavMr. On the other hand, the Ca^{2+} selectivity mechanism of CavMr

completely differs from that of CavAb. Structural comparison of NavAb and CavAb

showed that the aspartate mutations did not alter the main chain trace, and simply

introduced the negative charges around the ion pathway to increase Ca^{2+} permeability

(Fig. 7a and b) (Tang et al., 2013). In contrast, in the case of CavMr, two non-charged

residues (Gly4 and Val6) are required for the high Ca^{2+} selectivity (Fig. 4c, 5a), but

Asp7 is not necessary (*SI Appendix*, Fig. S2f). A no-charge mutation at position 7,

CavMr-D7M "TLEGWVM", is an outstanding example demonstrating that high Ca^{2+}

selectivity can be achieved in the absence of any aspartates in its filter region (*SI*

Appendix, Fig. S2f). Furthermore, the introduction of a negative charge into the

selectivity filter (G4D mutation) had the opposite effect on the Ca^{2+} selectivity of

CavMr compared with NavAb and NaChBac (Tang et al., 2014; Yue et al., 2002).

Moreover, the decreased selectivity in G4S also indicates that the glycine at position 4 is

indispensable for Ca^{2+} selectivity in CavMr (Fig. 5b). The flexibility and/or small size

of the glycine at position 4 in CavMr might be critical. These findings are inconsistent

with the notion derived from the Ca^{2+}-selective mutants of NavAb and NaChBac, and

therefore the native structure of the selectivity filter and the molecular mechanism of

ion selectivity of CavMr are thought to differ from those of CavAb. While the structure of CavMr is not yet available, we are able to speculate on the structure of the selectivity filter of CavMr on the basis of the structure of human Cav1.1 subdomains I and III (Wu et al., 2016) (Fig. 6c), whose selectivity filter sequences are very similar to that of CavMr. In the selectivity filter of Cav1.1 subdomains I and III, the side chain of the residue at position 7 is shifted outward. The position-4 glycine residue widens the entrance of the selectivity filter, which would facilitate the entry of hydrated cations into the ion pore and might increase Ca^{2+} selectivity.

Cavs in prokaryotes and the species-specific tuning of homo-tetrameric channels.

Prokaryotes have a number of putative Ca^{2+} binding proteins, such as EF-hand proteins, P-type Ca^{2+} pumps, and Ca^{2+} transporters (Domínguez et al., 2015). The intracellular Ca^{2+} concentration is kept low and changes in response to mechanical and chemical stimuli (Dominguez, 2004). These features imply that prokaryotic Ca^{2+} signaling is similar to that of eukaryotes. The strong ability of CavMr to exclude Na^+ and K^+ along with Ca^{2+} permeation suggests that its primary physiological role is Ca^{2+} intake in response to a voltage change (Fig. 2f and g). In some bacteria, the direction of flagellar rotation and chemotaxis changes depending on the internal Ca^{2+} concentration (Ordal, 1977; Tisa et al., 1993; Tisa and Adler, 1995). *M. ruber* was isolated from hot springs, and therefore a sufficient amount of Ca^{2+} is likely to exist in its native environment (Loginova et al., 1984). CavMr activation by a voltage change, which could vary depending on the environmental ionic conditions, might lead to any response to adapt to the new environment, such as flagellar rotation. These characteristics indicate the existence of signal coupling between the membrane voltage and Ca^{2+}, even in the early stages of life, which might be the origin of the corresponding functions in eukaryotes, such as muscle contraction.

NavPp permeates more Na^+ than Ca^{2+}, but its selectivity is modest (Fig. 3f and Table 1). Notably, *P. pacifica* is a marine myxobacterium that requires NaCl for its growth (Iizuka et al., 2003). As mentioned above, the basic architecture of the CavMr/NavPp group is thought to produce a preference for Ca^{2+}. *P. pacifica* might modify this channel architecture to acquire a Na^+ intake pathway, which would likely result in the remaining feature of low-affinity Ca^{2+} inhibition in NavPp. This flexible usage of homo-tetrameric channels to allow different cations to permeate is also reported in another bacterium, *Bacillus alkalophilus* (DeCaen et al., 2014). NsvBa from *B. alkalophilus* is a non-selective channel whose selectivity filter is changed from "TLESWAS", a typical Na^+-selective sequence in alkaliphilic bacillus, to "TLDSWGS", possibly to adapt to its

ionic environment. Recently, an early eukaryote, diatom, was found to have another
homo-tetrameric channel with no selectivity that has an important role in electrical
signaling in this species (Helliwell et al., 2019). These findings suggest that the cation
selectivity of the homo-tetrameric channel family can be flexibly tuned to realize the
required roles specific to its original species. We also found four homologues that have
filter sequences similar to CavMr "Tx(E/D)GWx(D/E)" in the NCBI database
(WP_009945599.1, WP_012983075.1, WP_024079824.1, and XP_002186055.1).
XP_002186055.1 belongs to diatoms, which implies the wide use of monomeric Cavs in
various organisms – not only prokaryotes, but also eukaryotes.

Insights into Ca^{2+} selectivity and the evolution of mammalian Cavs.

12
Figure 6. Comparison between mammalian and prokaryotic Cav.
a and b). Structures of the selectivity filter in CavAb (PDB code: 4MVZ) and NavAb
(PDB code: 5YUA). c and d) Structure of the rabbit Cav1.1 selectivity filter (PDB
code: 5GJV). The subdomains I and III (c), and II and IV (d) were separately shown.
The carbon atoms of negatively charged residues were indicated in pink. Dashed green
circle indicates the wide entrance of the selectivity filter.

Aspartate residues are generally observed in the Ca^{2+} permeation pathway in ion
channels, as well as many Ca^{2+} binding proteins (Halling et al., 2016; Yan et al., 2015;
Zalk et al., 2015). Actually, NavAb and NaChBac were successfully transformed to
Ca^{2+}-selective channels with the aspartate-introduced filter sequences "TLDDW(S/A)D"

(Tang et al., 2014; Yue et al., 2002). But, our results elucidate that this strategy is not the only method for achieving high Ca^{2+} selectivity. Human Cavs subdomains possess, at most, two aspartate residues in the selectivity filters in other part than position 3. In addition, the negatively charged residue at position 3, which is thought to be the most critical for cation selectivity in both Navs and Cavs, is not aspartate, but glutamate in most of the human Cavs subdomains (Yu and Catterall, 2004). CavAb has 12 aspartates in the selectivity filter of a channel tetramer, while there are 4 aspartates in CavMr. The net negative charge is 5~7 in mammalian Cavs, 8 in CavMr, and 12 in CavAb. These findings indicate that Ca^{2+} selectivity can be achieved with even fewer negative charges than CavAb and close to mammalian Cav, probably with the contribution of a special backbone structure around the selectivity filters.

It is noteworthy that the selectivity filter sequence of CavMr is very similar to those of human Cav subdomains I and III, both of which possess a glycine at position 4 (Fig. 6c). Especially, the Cav3.1 and 3.2 subdomains I have the same sequence as CavMr. On the other hand, the sequences of subdomains II and IV are relatively similar to that of CavAb (Fig. 6d). These sequence similarities of the glycine residue at position 4 are also found in CatSper, the sperm calcium permeable channel (Darszon et al., 2011). The channel region of CatSper is formed by four different subunits (CatSper1-4). The selectivity filters of CatSper 3 "TVDGWTD" and CatSper 4 "TQDGWVD" are similar to that of CavMr and share a glycine residue at position 4 as well as subdomains I and III of 24TM Cavs. These similarities indicate the generality of the CavMr-like Ca^{2+}-selectivity mechanism. Further investigation of the detailed structure of CavMr may help to elucidate the principles and origin underlying Ca^{2+} selectivity.

Material and methods

Cloning of BacNav homologues and site-directed mutagenesis

The NaChBac amino acid sequence (NP_242367) was used as the query for a BLASTP search against the Microbial Genomic database at NCBI. The identified primary sequence data were obtained from Entrez at NCBI (*Meiothermus ruber* as ZP_04038264, *Plesiocystis pacifica* as ZP_01909854, *Halomonas elongata* as YP_003896792 and *Teredinibacter turnerae* as YP_003073405). These DNAs were synthesized by Genscript Inc. and subcloned into the pCI vector (Promega) using the EcoRI and SalI sites and the pBiEX vector (Novagen) using NcoI and BamHI sites, respectively. Site-directed mutagenesis was achieved by polymerase chain reaction (PCR) of the full-length plasmid containing the Nav gene using PrimeSTAR® MAX DNA Polymerase (Takara Bio.). All clones were confirmed by DNA sequencing.

Electrophysiological analysis using mammalian cells

For the recordings related to mole fraction effects (Fig. S2a-d), currents were recorded from Chinese hamster ovary (CHO) -K1 cells expressing channels. The recordings were performed as described previously (Tateyama and Kubo, 2018). Cells were transfected with channel DNAs using the LipofectAMINE 2000 (Invitrogen) and plated onto cover slips. Currents were recorded 24-36 h after transfection. Current recording by the whole cell patch clamp technique was performed using Axopatch 200B amplifiers, Digidata1332A, and pClamp 9 software (Molecular Devices). The pipette solution contained 130 mM KCl, 5 mM Na_2-ATP, 3 mM EGTA, 0.1 mM $CaCl_2$, 4 mM $MgCl_2$ and 10 mM HEPES (pH 7.2 adjusted with KOH). The bath solution contained 135 mM NaCl, 4 mM KCl, 1 mM $CaCl_2$, 5 mM $MgCl_2$ and 10 mM HEPES (pH 7.4 adjusted with NaOH). For the measurement of mole fraction effects, the bath solutions containing different ratio of NaCl / $CaCl_2$ (135/0, 108/18, 81/36, 54/54, 27/82 and 0/90 mM) were used. The Ca^{2+}-free solution was achieved with the solution containing 135 mM NaCl, 1 mM EGTA and 0 mM $CaCl_2$.

Electrophysiological measurement in insect cells

The recordings other than those for mole fraction effects were performed using SF-9 cells. SF-9 cells were grown in Sf-900™ III medium (Gibco) complemented with 0.5% 100× Antibiotic-Antimycotic (Gibco) at 27°C. Cells were transfected with target channel-cloned pBiEX vectors and enhanced green fluorescent protein (EGFP)-cloned pBiEX vectors using Fugene HD transfection reagent (Promega). The channel-cloned

vector (2 μg) was mixed with 0.5 μg of the EGFP-cloned vector in 100μL of the culture

medium. Next, 3 μL Fugene HD reagent was added and the mixture was incubated for

10 min before the transfection mixture was gently dropped onto cultured cells. After

incubation for 16-48 h, the cells were used for electrophysiological measurements. In

the measurement of I-V relation curves, the pipette solution contains 75 mM NaF, 40

mM CsF, 35 mM CsCl, 10 mM EGTA, and 10 mM HEPES (pH 7.4 adjusted by CsOH)

For evaluation of ion selectivity, high Na pipette solution (115 mM NaF, 35 mM NaCl,

10 mM EGTA, and 10 mM HEPES (pH 7.4 adjusted by NaOH)) was used. For the

evaluation of Ca, Sr, K and Cs selectivity, Ca solution (100 mM $CaCl_2$, 10 mM HEPES

(pH 7.4 adjusted by $Ca(OH)_2$), and 10 mM glucose), Sr solution (100 mM $SrCl_2$, 10

mM HEPES (pH 7.4 adjusted by $Sr(OH)_2$), and 10 mM glucose), K solution (150 mM

KCl, 2mM $CaCl_2$, 10 mM HEPES (pH 7.4 adjusted by KOH), and 10 mM glucose), and

Cs solution (150 mM CsCl, 2mM $CaCl_2$, 10 mM HEPES (pH 7.4 adjusted by CsOH),

and 10 mM glucose) were used as the bath solution, respectively. E_{rev} of high Ca^{2+}

selective channels were measured under three external solutions containing 144mM

NMDG-Cl and 4mM $CaCl_2$, 135mM NMDG-Cl and 10mM $CaCl_2$, 120mM NMDG-Cl

and 20mM $CaCl_2$ (10mM HEPES pH 7.4 adjusted with HCl). And E_{rev} of high Ca^{2+}

selective channels for the calculation of P_K/P_{Ca} and P_{Cs}/P_{Ca} were measured under

external solutions containing 135mM NMDG-Cl and 10mM $CaCl_2$ (10mM HEPES pH

7.4 adjusted with HCl) with high K pipette solution (115 mM KF, 35 mM KCl, 10 mM

EGTA, and 10 mM HEPES (pH 7.4 adjusted by KOH)) and high Cs pipette solution

(115 mM CsF, 35 mM CsCl, 10 mM EGTA, and 10 mM HEPES (pH 7.4 adjusted by

CsOH)), respectively. E_{rev} of NavPp for the calculation of P_{Ca}/P_{Na} or P_{Sr}/P_{Na} were

measured under external solution containing 50mM NMDG-Cl, 40mM NaCl, 40mM

$CaCl_2$ or $SrCl_2$ and 10mM HEPES pH 7.4 adjusted with NaOH. E_{rev} of NavPp for the

calculation of P_{Cs}/P_{Na} was measured under high Cs pipette solution and external

solution containing 110mM NMDG-Cl, 40mM NaCl, 3mM $CaCl_2$ and 10mM HEPES

pH 7.4 adjusted with NaOH.

As the pipette solution for measurement of the Ca block effect in NavPp, low Na

pipette solution (140 mM CsF, 10 mM NaCl, 10 mM EGTA, and 10 mM HEPES (pH

7.4 adjusted by CsOH)) and high Na pipette solution were used for inward and outward

current measurement, respectively. As a bath solution, Ca blocking solution (30 mM

NaCl, 120 mM NMDG-Cl, 1.5 mM $CaCl_2$, 10 mM HEPES (pH 7.4 adjusted by NaOH)

and 10 mM glucose) was used for the 1.5 mM Ca blocking condition. In 10mM Ca

blocking condition, a bath solution contains 30 mM NaCl, 105 mM NMDG-Cl, 10 mM

$CaCl_2$, 10 mM HEPES (pH 7.4 adjusted by NaOH) and 10 mM glucose. And, in each

Ca blocking conditions, 15 mM NMDG-Cl was replaced per 10 mM $CaCl_2$. The bath

solution was changed using the Dynaflow® Resolve system. All experiments were

conducted at $25 \pm 2°C$. All results are presented as mean $\pm$ standard error.

Calculation of ion selectivity by the GHK equation

To determine the ion selectivity of each channel, the intracellular solution and

extracellular solution were arbitrarily set and the reversal potential at each concentration

was measured by giving the ramp pulse of membrane potential. The applied ramp pulse

was set to include the reversal potential. In addition, a depolarization stimulus of 2-10

ms was inserted to check whether the behavior of the cell changed for each

measurement. As a result, P_{Ca}/P_{Na} was calculated by substituting the obtained reversal

potential (E_{rev}) into the expression derived from the GHK equation (Frazier et al., 2000);

$$P_{Ca}/P_{Na} = \frac{-([Na]_i - [Na]_o e^{-E_{rev}F/RT})(1 - e^{-2E_{rev}F/RT})}{4([Ca]_i - [Ca]_o e^{-2E_{rev}F/RT})(1 - e^{-E_{rev}F/RT})}$$

where F is Faraday's constant, R is the Gas constant, and T is 298.1[K]. The same

expression was used for Sr^{2+}. The Sr^{2+} selectivity (P_{Sr}/P_{Na}) was measured in the same

way.

Na^+ selectivity against monovalent cations (P_M/P_{Na}) was calculated by substituting the

obtained reversal potential and P_{Ca}/P_{Na} into the expression derived from the GHK

equation (Lopin et al., 2012):

$$P_M/P_{Na} = [\frac{-4([Ca]_i - [Ca]_o e^{-2E_{rev}F/RT})(1 - e^{-E_{rev}F/RT})}{([Na]_i - [Na]_o e^{-E_{rev}F/RT})(1 - e^{-2E_{rev}F/RT})} \cdot (P_{Ca}/P_{Na})$$
$$- 1][\frac{([Na]_i - [Na]_o e^{-E_{rev}F/RT})}{([M]_i - [M]_o e^{-E_{rev}F/RT})}]$$

References

Almers W, McCleskey EW. 1984. Non-selective conductance in calcium channels of

frog muscle: calcium selectivity in a single-file pore. *J Physiol* **353**:585–608.

doi:10.1113/jphysiol.1984.sp015352

Bando Y, Irie K, Shimomura T, Umeshima H, Kushida Y, Kengaku M, Fujiyoshi Y,

Hirano T, Tagawa Y. 2014. Control of Spontaneous Ca2+ Transients Is Critical for

Neuronal Maturation in the Developing Neocortex. *Cereb Cortex*.

doi:10.1093/cercor/bhu180

Catterall W. 2000. From ionic currents to molecular mechanisms: the structure and

function of voltage-gated sodium channels. *Neuron* **26**:13–25.

Darszon A, Nishigaki T, Beltran C, Treviño CL. 2011. Calcium Channels in the

Development, Maturation, and Function of Spermatozoa. *Physiol Rev*

91:1305–1355. doi:10.1152/physrev.00028.2010

DeCaen PG, Takahashi Y, Krulwich T a, Ito M, Clapham DE. 2014. Ionic selectivity

and thermal adaptations within the voltage-gated sodium channel family of

alkaliphilic Bacillus. *Elife* **3**:1–15. doi:10.7554/eLife.04387

Dominguez DC. 2004. Calcium signalling in bacteria. *Mol Microbiol* **54**:291–297.

doi:10.1111/j.1365-2958.2004.04276.x

Domínguez DC, Guragain M, Patrauchan M. 2015. Calcium binding proteins and

calcium signaling in prokaryotes. *Cell Calcium* **57**:151–165.

doi:10.1016/j.ceca.2014.12.006

Finol-Urdaneta RK, Wang Y, Al-Sabi A, Zhao C, Noskov SY, French RJ. 2014.

Sodium channel selectivity and conduction: prokaryotes have devised their own

molecular strategy. *J Gen Physiol* **143**:157–71. doi:10.1085/jgp.201311037

Frazier CJ, George EG, Jones SW. 2000. Apparent change in ion selectivity caused by

changes in intracellular K(+) during whole-cell recording. *Biophys J* **78**:1872–80.

doi:10.1016/S0006-3495(00)76736-1

Halling DB, Liebeskind BJ, Hall AW, Aldrich RW. 2016. Conserved properties of

individual Ca2+-binding sites in calmodulin. *Proc Natl Acad Sci*

113:E1216–E1225. doi:10.1073/pnas.1600385113

Heinemann SH, Terlau H, Stühmer W, Imoto K, Numa S. 1992. Calcium channel

characteristics conferred on the sodium channel by single mutations. *Nature*

356:441–443.

Helliwell KE, Chrachri A, Koester JA, Wharam S, Verret F, Taylor AR, Wheeler GL,

Brownlee C. 2019. Alternative Mechanisms for Fast Na + /Ca 2+ Signaling in

Eukaryotes via a Novel Class of Single-Domain Voltage-Gated Channels. *Curr*

Biol **29**:1503-1511.e6. doi:10.1016/j.cub.2019.03.041

Hille B. 2001. Ion Channels of Excitable Membranes, Third Edtition (Sunderland, MA:

Sinauer Associates Inc).

Iizuka T, Jojima Y, Fudou R, Hiraishi A, Ahn JW, Yamanaka S. 2003. Plesiocystis

pacifica gen. nov., sp. nov., a marine myxobacterium that contains dihydrogenated

menaquinone, isolated from the Pacific coasts of Japan. *Int J Syst Evol Microbiol*

53:189–195. doi:10.1099/ijs.0.02418-0

Irie K, Haga Y, Shimomura T, Fujiyoshi Y. 2018. Optimized expression and

purification of NavAb provide the structural insight into the voltage dependence.

FEBS Lett **592**:274–283. doi:10.1002/1873-3468.12955

Irie K, Kitagawa K, Nagura H, Imai T, Shimomura T, Fujiyoshi Y. 2010. Comparative

study of the gating motif and C-type inactivation in prokaryotic voltage-gated sodium channels. *J Biol Chem* **285**:3685–94. doi:10.1074/jbc.M109.057455

Irie K, Shimomura T, Fujiyoshi Y. 2012. The C-terminal helical bundle of the tetrameric prokaryotic sodium channel accelerates the inactivation rate. *Nat Commun* **3**:793. doi:10.1038/ncomms1797

Ito M, Xu H, Guffanti AA, Wei Y, Zvi L, Clapham DE, Krulwich TA. 2004. The voltage-gated Na+ channel NaVBP has a role in motility, chemotaxis, and pH homeostasis of an alkaliphilic Bacillus. *Proc Natl Acad Sci* **101**:10566–10571. doi:10.1073/pnas.0402692101

Kamiya A, Hayama Y, Kato S, Shimomura A, Shimomura T, Irie K, Kaneko R, Yanagawa Y, Kobayashi K, Ochiya T. 2019. Genetic manipulation of autonomic nerve fiber innervation and activity and its effect on breast cancer progression. *Nat Neurosci*. doi:10.1038/s41593-019-0430-3

Koishi R, Xu H, Ren D, Navarro B. 2004. A superfamily of voltage-gated sodium channels in bacteria. *J Biol Chem* **279**:9532–9538. doi:10.1074/jbc.M313100200

Lee S, Goodchild SJ, Ahern CA. 2012. Molecular and functional determinants of local anesthetic inhibition of NaChBac. *Channels (Austin)* **6**:403–406. doi:10.4161/chan.21807

Lin CW, Sim S, Ainsworth A, Okada M, Kelsch W, Lois C. 2010. Genetically Increased Cell-Intrinsic Excitability Enhances Neuronal Integration into Adult Brain Circuits. *Neuron* **65**:32–39. doi:10.1016/j.neuron.2009.12.001

Loginova LG, Egorova LA, Golovacheva RS, Seregina LM. 1984. Thermus ruber sp. nov., nom. rev. *Int J Syst Bacteriol* **34**:498–499. doi:10.1099/00207713-34-4-498

Lopin K V., Thevenod F, Page JC, Jones SW. 2012. Cd2+ Block and Permeation of CaV3.1 (α1G) T-Type Calcium Channels: Candidate Mechanism for Cd2+ Influx. *Mol Pharmacol* **82**:1183–1193. doi:10.1124/mol.112.080176

Nagura H, Irie K, Imai T, Shimomura T, Hige T, Fujiyoshi Y. 2010. Evidence for lateral mobility of voltage sensors in prokaryotic voltage-gated sodium channels. *Biochem Biophys Res Commun* **399**:341–6. doi:10.1016/j.bbrc.2010.07.070

Ordal GW. 1977. Calcium ion regulates chemotactic behaviour in bacteria. *Nature* **270**:66–67. doi:10.1038/270066a0

Rahman T, Cai X, Brailoiu GC, Abood ME, Brailoiu E, Patel S. 2014. Two-pore channels provide insight into the evolution of voltage-gated Ca2+ and Na+ channels. *Sci Signal* **7**:ra109–ra109. doi:10.1126/scisignal.2005450

Ren D, Navarro B, Xu H, Yue L, Shi Q, Clapham DE. 2001. A prokaryotic voltage-gated sodium channel. *Science* **294**:2372–5. doi:10.1126/science.1065635

Shimomura T, Irie K, Fujiyoshi Y. 2016. Molecular determinants of prokaryotic voltage-gated sodium channels for recognition of local anesthetics. *FEBS J* **283**:2881–2895. doi:10.1111/febs.13776

Shimomura T, Irie K, Nagura H, Imai T, Fujiyoshi Y. 2011. Arrangement and mobility of the voltage sensor domain in prokaryotic voltage-gated sodium channels. *J Biol Chem* **286**:7409–7417. doi:10.1074/jbc.M110.186510

Strong M, Chandy KG, Gutman GA. 1993. Molecular evolution of voltage-sensitive ion channel genes: on the origins of electrical excitability. *Mol Biol Evol* **10**:221–42. doi:10.1093/oxfordjournals.molbev.a039986

Tang L, Gamal El-Din TM, Payandeh J, Martinez GQ, Heard TM, Scheuer T, Zheng N, Catterall W a. 2014. Structural basis for Ca2+ selectivity of a voltage-gated calcium channel. *Nature* **505**:56–61. doi:10.1038/nature12775

Tang L, Gamal El-Din TM, Payandeh J, Martinez GQ, Heard TM, Scheuer T, Zheng N, Catterall WA. 2013. Structural basis for Ca2+ selectivity of a voltage-gated calcium channel. *Nature* **505**:56–61. doi:10.1038/nature12775

Tang L, Gamal El-Din TM, Swanson TM, Pryde DC, Scheuer T, Zheng N, Catterall WA. 2016. Structural basis for inhibition of a voltage-gated Ca2+ channel by Ca2+ antagonist drugs. *Nature* **537**:117–121. doi:10.1038/nature19102

Tateyama M, Kubo Y. 2018. Gi/o-coupled muscarinic receptors co-localize with GIRK channel for efficient channel activation. *PLoS One* **13**:1–18. doi:10.1371/journal.pone.0204447

Tisa LS, Adler J. 1995. Cytoplasmic free-Ca2+ level rises with repellents and falls with attractants in Escherichia coli chemotaxis. *Proc Natl Acad Sci U S A* **92**:10777–81. doi:10.1073/pnas.92.23.10777

Tisa LS, Olivera BM, Adler J. 1993. Inhibition of Escherichia coli chemotaxis by omega-conotoxin, a calcium ion channel blocker. *J Bacteriol* **175**:1235–1238. doi:10.1128/jb.175.5.1235-1238.1993

Tsai CJ, Tani K, Irie K, Hiroaki Y, Shimomura T, McMillan DG, Cook GM, Schertler GFX, Yoshinori F, Li XD. 2013. Two alternative conformations of a voltage-gated sodium channel. *J Mol Biol* **425**:4074–4088. doi:10.1016/j.jmb.2013.06.036

Wu J, Yan Z, Li Z, Qian X, Lu S, Dong M, Zhou Q, Yan N. 2016. Structure of the voltage-gated calcium channel Cav1.1 at 3.6 Å resolution. *Nature* **537**:191–196. doi:10.1038/nature19321

Yan Z, Bai XC, Yan C, Wu J, Li Z, Xie T, Peng W, Yin CC, Li X, Scheres SHW, Shi Y, Yan N. 2015. Structure of the rabbit ryanodine receptor RyR1 at near-atomic resolution. *Nature* **517**:50–55. doi:10.1038/nature14063

Yu FH, Catterall W a. 2004. The VGL-chanome: a protein superfamily specialized for

electrical signaling and ionic homeostasis. *Sci STKE* **2004**:re15.

doi:10.1126/stke.2532004re15

Yue L, Navarro B, Ren D, Ramos A, Clapham DE. 2002. The cation selectivity filter of

the bacterial sodium channel, NaChBac. *J Gen Physiol* **120**:845.

doi:10.1085/jgp.20028699

Zalk R, Clarke OB, Georges A Des, Grassucci RA, Reiken S, Mancia F, Hendrickson

WA, Frank J, Marks AR. 2015. Structure of a mammalian ryanodine receptor.

Nature **517**:44–49. doi:10.1038/nature13950

Acknowledgements

We very appreciate to Dr. Yoshihiro Kubo for the critical suggestion about the structure of our manuscript. This work was supported by Grants-in-Aid for Scientific Research (S), a Grant-in-Aid for Young Scientists (B), the Japan Agency for Medical Research and Development and the Toyoaki Scholarship Foundation.

Author contribution

T.S. and K.I. conducted the experiments; T.S. searched for homologues; Y.Y. and K.I. performed the electrophysiological experiments of insect cells; M.T. performed the electrophysiological experiments of mammalian cells; T.S., Y.Y., H.N. and K.I. optimized the measurement conditions; T.S., Y.Y., H.N., Y.F. and K.I. contributed to the study design and wrote the paper.

Figure legends.

Table1. Relative permeability of CavMr and NavPp

Figure 1. Sequence analysis and the representative current recordings of the novel

BacNav homologues

a). Phylogenetic tree of the BacNav homologues with their GenBank[TM] accession

numbers. The ClustalW program was used to align the multiple protein sequences of the

BacNav homologues. The phylogenetic tree was generated using "PROTDIST", one of

the PHYLIP package (Phylogeny Inference Package:

http://evolution.genetics.washington.edu/phylip.html). The branch lengths are

proportional to the sequence divergence, with the scale bar corresponding to 0.1

substitution per amino acid position. Four homologues colored that are not included in

canonical BacNavs were cloned and expressed to check the current activity. Those of

two which are underlined in red and shown as bold generated the detectable currents. b).

Schematic secondary structure and selectivity filter sequence of BacNavs and human

Cavs. Cylinder indicates α-helix. The selectivity filter sequences are indicated by

alphabetical characters. Negatively charged residues are colored by red. Glycine

residues in the position 4 are colored by cyan. The straight lines indicate the other part

of pore domain. The selectivity filter sequence of hCav1.1 (UniProt ID: Q13698),

hCav2.1 (O00555) and hCav3.1 (O43497), were used. c and d). Representative current

traces to obtain the current-voltage relationships of CavMr (c) and NavPp (d) in Sf9

cells. The straight lines indicating the zero-current level in the representative current

traces. Currents were generated under the bath solutions containing high Na^+ (top) and

high Ca^{2+} (middle), by a series of step-pulses shown in bottom. e and f). Current-voltage

relationships of CavMr (e) and NavPp (f) measured under the different bath solutions

(filled black; 150 mM NaCl, open black; 75 mM NaCl and 75 mM NMDG-HCl, open

red; 75 mM NaCl and 50 mM $CaCl_2$, filled red; 50 mM $CaCl_2$ and 75 mM NMDG-HCl).

Currents of CavMr and NavPp were normalized to that by 0mV depolarization stimuli

under 75 mM NaCl and 50 mM $CaCl_2$ bath solution and 150 mM NaCl bath solution,

respectively.

Figure 2. Cation selectivity of CavMr

a and b). Recordings of the reversal potential of CavMr currents using the ramp protocol.

Currents were generated by the step pulse of -20 mV from -140 mV holding potential,

followed by the ramp pulses with different voltage values (shown at the bottom of

panels a and b). The values of the reversal potential recorded with three different ramp pulses were averaged. Currents were measured under the bath solution containing 4 mM (a) and 10 and 20 mM (b) $CaCl_2$ and the pipette solution with 150 mM NaCl. c). The plot of the reversal potential to the bath $[Ca^{2+}]_{out}$. Each value was obtained using the protocol shown in a and b. The relationship was fitted by a line with the slope of 41.07 ± 2.64 mV per decade ($n = 4$). d). Representative current traces to obtain the reversal potential under the condition of 100 mM $[Sr^{2+}]_{out}$ and 150 mM $[Na^+]_{in.}$ Currents were generated by the protocol shown in the lower part. e). Representative current traces to investigate the P_{Cs}/P_{Ca} and P_K/P_{Ca}, the pipette solutions contained 150 mM $[Cs^+]_{in}$ for P_{Cs}/P_{Ca} and 150 mM $[K^+]_{in}$ for P_K/P_{Ca}, while the bath solution contained 10 mM $[Ca^{2+}]_{out}$ in both cases. f). The relative permeability of Ca^{2+} or Sr^{2+} to Na^+ in CavMr, calculated from the reversal potential that was obtained in a, b, and d. g). The relative permeability of each monovalent cation to Ca^{2+} in CavMr, derived from the data shown in e.

Figure 3. Na^+ permeability and the extracellular Ca^{2+}-dependent inhibition in NavPp

a) $[Ca^{2+}]_{out}$ dependent inhibitory effects in NavPp. Currents were normalized to those under 1.5 mM $[Ca^{2+}]_{out}$(filled square; NavAb current in the 30 mM $[Na^+]_{out}$ and 10 mM $[Na^+]_{in}$ (representative data are shown in Fig. S3b), open triangle down; NavPp inward current in the 30 mM $[Na^+]_{out}$ and 10 mM $[Na^+]_{in}$ (Fig. S3a), open triangle up; NavPp outward current in the 30 mM $[Na^+]_{out}$ and 150 mM $[Na^+]_{in}$ (Fig. S3c), filled triangle up; NavPp outward current in the 0 mM $[Na^+]_{out}$ and 150 mM $[Na^+]_{in}$. b). Representative currents to obtain the reversal potential in NavPp for P_{Ca}/P_{Na}. Currents were generated by the ramp protocol shown in bottom. $[Ca^{2+}]_{out}$ was varied from 10 to 60 mM with the fixed $[Na^+]$ both in the bath (60mM) and pipette (150mM). c). Representative currents to obtain the reversal potential in NavPp for P_{Sr}/P_{Na}. Currents were generated by the ramp protocol shown in bottom. $[Sr^{2+}]_{out}$ was 40 mM with the fixed $[Na^+]$ both in the bath (40mM) and pipette (150mM). d). The relative permeability of different cation species to Na^+ in NavPp, calculated from the reversal potential that was obtained in b, c and Fig.S3d and e.

Figure 4. The cation selectivity of the swapping mutant channels in their selectivity filter between CavMr and NavPp

a) Amino acid sequences of the selectivity filter in the swapped mutants, CavMr-Pp, Nav-Pp, NavAb_Mr, and CavMr_Ab. The selectivity filter sequences of CavMr, and

NavPp, and NavAb are indicated by alphabetical characters with cyan, red, and gray shade, respectively. Negatively charged residues are colored by red. Glycine residues are colored by cyan. The straight lines of cyan, red, and black indicates the other part of pore domain of CavMr, NavPp, and NavAb, respectively. b) Pore domains of crystal structure of NavAb (PDB code:5YUA). The selectivity filter, which corresponds to the sequences shown in a, was indicated in red. c) The relative permeability of divalent cations to Na^+ (left) and that of monovalent cations to Ca^{2+} (right) in NavPp-Mr. d) The relative permeability of different cation species to Na^+ in CavMr-Pp.

Figure 5. The single point mutations losing and obtaining Ca^{2+} selectivity of CavMr and NavPp, respectively.

a) The relative permeability of each cation species to Na^+ in the single-point mutants of CavMr. The selectivity filter of CavMr was changed to the corresponding residues of NavPp at position 4 (G4D) and position 6 (V6T), respectively b) The relative permeability of each cation species to Na^+ in the G4S mutants of CavMr, whose position 4 residue of the selectivity filter was mutated to the corresponding residue of canonical BacNavs. c) The relative permeability of each cation species to Na^+ in the single-point mutants of NavPp. The selectivity filter of NavPp was changed to the corresponding residues of CavMr at position 4 (D4G) and position 6 (T6V), respectively.

Figure 6. Comparison between mammalian and prokaryotic Cav.

a and b). Structures of the selectivity filter in CavAb (PDB code: 4MVZ) and NavAb (PDB code: 5YUA). c and d) Structure of the rabbit Cav1.1 selectivity filter (PDB code: 5GJV). The subdomains I and III (c), and II and IV (d) were separately shown. The carbon atoms of negatively charged residues were indicated in pink. Dashed green circle indicates the wide entrance of the selectivity filter.