DISTANCES WALKED BY BEACH USERS AND PROTECTING SHOREBIRD

HABITAT ZONES FROM DISTURBANCE

STEPHEN TOTTERMAN[1]

[1] Empire Vale, NSW 2478, AUSTRALIA.

Email: stephen@totterman.net.au

RUNNING TITLE: Distances walked by beach users

7 ABSTRACT

Human population growth along Australia's coast is increasing development and recreation pressures on
beaches and shorebirds. This study observed human recreation on 18 beaches on the far north coast of
New South Wales in February and March 2019. The far north coast supports the largest numbers of beach-
resident Australian Pied Oystercatchers *Haematopus longirostris* in the State. The most frequent activities
observed were walking (29%), dog walking (21%) and swimming (16%). Walkers covered greater
distances compared to other beach users. For beach walkers and dog walkers combined, the mean along
shore distance walked from an access point was 809 m and the 95th percentile was 1990 m. Noting that
human recreation disturbance is a major conservation threat to beach nesting birds and that pro-
environmental behaviour is uncommon among beach users, large separation distances between beach
access points and shorebird habitat zones are recommended to reduce human intrusions into those zones.
This spatial zoning and passive human exclusion strategy can be applied to long and less-developed
beaches.

20 INTRODUCTION

Growing human populations and demand for coastal lifestyles away from the major cities (the "sea
change" phenomenon) are strong demographic forces that are increasing development and recreation
pressures on Australia's coastal areas (Priest *et al.* 2002). BirdLife Australia (2018) have stated that the
greatest conservation threat to beach nesting birds is disturbance from people visiting the beach.

Birds typically respond to a perceived threat, such as a potential predator or a human, by moving
away (Blumstein 2006). Frequent human recreation disturbance can result in displacement of shorebirds
from their preferred habitat (Schultz & Stock 1993, Liley & Sutherland 2007), increased energy
expenditure in flight responses (West *et al.* 2002), reduced energy intake through disrupted foraging
(Burger 1994, West *et al.* 2002) and exposure of eggs or dependent young to diminished parental care
(Lord *et al.* 2001, Verhulst *et al.* 2001, Harrison 2009). Disturbance has the potential to result in declines
in bird populations if displacement becomes permanent and there is little surplus habitat to move into, if
survival is reduced or if breeding success is lowered (Gill *et al.* 2001). However, it is difficult to
extrapolate observed individual or local effects to population impacts (Davidson & Rothwell 1993, Hill *et*
al. 1997, Gill *et al.* 2001).

Human recreation disturbance rates on beaches often outnumber natural disturbances, even on
beaches with low visitation rates (Weston & Elgar 2005, Harrison 2009). Habitat and life history attributes
make beach nesting birds particularly sensitive to human recreation disturbance. Ocean beaches are
narrow, linear habitats where birds can find it difficult to maintain a safe distance from people
(Pienkowski 1993, Harrison 2009). Territorial behaviour and high site fidelity of oystercatchers (Taylor *et*
al. 2014), for example, means that breeding pairs are unlikely to relocate to alternative sites, if they exist.
Australian pied oystercatchers are large shorebirds (length 42–50 cm; Marchant & Higgins 1990) and
large birds are generally more "flighty" than are smaller species (Blumstein 2006, Weston *et al.* 2012a).
Beach nesting bird populations are often concentrated at a number of sites and local changes in breeding
populations at those sites are population impacts.

Manfredo and Dayer (2004) noted that the thoughts and actions of humans ultimately determine
the course and resolution (if any) of human-wildlife conflicts. Practical research into human recreation
disturbance of wildlife must then include studies of human attitudes and behaviour (*e.g.* Antos *et al.* 2007,
Maguire *et al.* 2011, Guiness *et al.* 2020, Schneider *et al.* 2020). However, far more studies into human
disturbance of birds have simply measured flight initiation distances (FIDs), *i.e.* the distance at which an
individual bird or flock moves away from a stimulus (typically a human walking towards the focal bird or
flock) (see Weston *et al.* 2012a for a summary). A common application for FID data is for selecting
"buffer distances" between humans and wildlife. However, Hill *et al.* (1997) cautioned that there are
potentially many experimental variables that affect FIDs (species identity, individuals or flocks, breeders
or non-breeders, the habitat, activity of the birds prior to disturbance, stimulus applied, starting distance,
etc.) and that the results of any one study are often limited to specific experimental conditions. There is a
risk that end users may generalise and misuse FIDs when developing management plans for human-
wildlife conflicts.

Codes of conduct and enforceable regulations are commonly applied with the aim to reduce
human-wildlife conflict. However, evidence is growing that "pro-environmental" behaviour is uncommon
in coastal areas and that compliance with regulations is low (Dowling & Weston 1999, Antos *et al.* 2007,
Weston *et al.* 2014, Maguire *et al.* 2019, Schneider *et al.* 2020). Glover *et al.* (2011) surveyed 295 visitors
and residents at Western Port Bay, Victoria and reported that while 86% of respondents supported
protection of shorebird habitat, only 39% were supportive of limitations on access for walkers to avoid
disturbance to wildlife. Given that it is difficult and costly to manage human behaviour, the application of
FID data for shorebirds on beaches may be limited to designing human exclosures around individual nest
sites or colonies and shorebird roosts (*e.g.* Lafferty *et al.* 2006, Weston *et al.* 2012b).

This study recommends, for longer and less-developed beaches, spatial zoning to restrict
disturbance activity to certain areas of the beach (Hill *et al.* 1997). Large separation distances between
access points and shorebird habitat zones (*i.e.* greater than the distances people typically walk) can reduce
intrusion rates into those zones and disturbance to shorebirds (Dowling & Weston 1999, Antos *et al.* 2007,
Liley & Sutherland 2007).

Selecting an appropriate separation distance depends on visitation rates to the beach. For the
simplest case of one habitat zone and one access point, the intrusion rate (R) into the habitat zone (*e.g.*
persons per hour*)* is:

$$R = V \times [1 - p(s)]$$

where V is the visitation rate, s is the separation distance between the access point and the start of the
habitat zone and $p(s)$ is the proportion of beach users walking to the start of the zone (Figure 1).

Given a fixed intrusion rate (*e.g.* potential disturbance events per hour), the separation distance
must be increased as the visitation rate V increases. Recognising strong among person/group variability in
distances walked and anticipating future growth in human populations and beach recreation, it is
recommended to simply adopt the 95th percentile distance under the assumption that patterns of beach
recreation and distances walked hold into the future. However, as daily visitation rates increase to
hundreds of people or more, other management actions that attempt to promote coexistence between
humans and shorebirds could be investigated (reviewed by Pienkowski 1993).

<INSERT FIGURE 1 HERE?>

This study observed distances walked for beach users on the far north coast of NSW. This region
supports the largest numbers of beach-resident Australian Pied Oystercatchers *Haematopus longirostris* in
the State including the South Ballina Beach, Broadwater Beach (which includes Airforce Beach and Salty
Lagoon), Bombing Range Beach and Yuraygir Key Management Sites currently recognised by the NSW
Department of Planning, Industry and Environment (2019). The Australian Pied Oystercatcher is currently
listed as Endangered in Schedule 1 of the NSW Biodiversity Conservation Act 2016 (NSW Government
2016).

Preceding studies for the Australian Pied Oystercatcher have reported mean flight initiation
distances from 39–129 m (Table 1). This study was motivated by concerns that oystercatchers at South

Ballina are increasingly threatened by "beach house" developments and that misuse of FID data has
resulted in inadequate separation distances between new beach accesses and oystercatcher territories. For
example, the FID results of Harrison (2009, Table 1) could be misused to allow access points as close as
46 m to oystercatcher nests. There are several problematic assumptions underlying this choice, including:
nest locations are fixed and do not change from year to year, a breeding pair needs only 0.7 Ha of space
(*i.e.* a circular area with radius 46 m) and people will not bring domestic dogs onto the beach, *etc*.

<INSERT TABLE 1 HERE>

METHODS

Study sites
Twenty locations (sites) were sampled on 18 beaches between Tweed Heads and Coffs Harbour, NSW,
between 8 February and 24 March 2019 (Figure 2, Table 2). Five oystercatcher breeding beaches were
included. Most beaches were moderate to long (mean length 6.5 km). On shorter beaches, *e.g.* Chinaman's
Beach (length 0.8 km), walking distances are constrained by the beach length.

<INSERT FIGURE 2 HERE>

<INSERT TABLE 2 HERE>

Sites were selected to include a variety of adjoining land uses and dogs permitted and dogs
prohibited zones (Table 2). Thirteen sites had a number of closely spaced access points that were observed
together as one site. The mean between-site distance of 87.7 km was far greater than the mean within-site
distance of 353 m (for sites with multiple access points). Remote and infrequently used sites are not
productive and were not sampled. For example, visitation rates to the picnic area north of Broadwater
Headland (at the southern end of South Ballina Beach) were too low to persevere with sampling.

Observations
Sampling occurred in the morning and on any day of the week. There were no school holidays during this
study. Ten observations of beach users were made at each site. Social groups of beach users (*e.g.* a picnic)
were recorded as one observation, to avoid pseudoreplication. The total sample size was 200 observations
(20 sites × 10 observations per site).

For busy access points, every third person/group entering the beach was observed, counting from
when the observer was within sight of the access point and ready to record new observations. For less
busy beaches or at less busy times, accumulation of observations was accelerated by observing every
consecutive person/group and/or observing multiple access points. Patchs Beach was visited twice to
accumulate ten observations and South Ballina was visited six times. Sampling of other sites was
completed in one visit.

The response variable was the maximum one-way distance a focal person/group walked from an
access point. Distances were measured using a handheld Global Positioning System (GPS) satellite
receiver (Garmin GPSmap 62s) or estimated if less than *c.* 50 m (where GPS imprecision is large relative
to the distance to be measured). Distances were measured in the along shore direction and included zero
distances where the person/group did not move along shore from the access point (*e.g.* walking directly
down shore to the water). Focal persons/groups were observed discreetly from some distance to avoid
disturbing them and potentially affecting their behaviour. Binoculars were used as a visual aid and, on
longer beaches, a bicycle was used (at low tide) to quickly move between observation points.

Predictor variables recorded together with each focal person/group observation were: day, time
the person/group entered the beach (Australian Eastern Standard Time, without daylight saving), number
of adults in the group, number of children in the group, numerically dominant gender of adults in the
group (male, female or balanced), number of accompanying dogs, activity of the person/group, count of
other people within 500 m of the access (excluding the observer), presence/absence of lifeguards on the
beach, nearest tide (high or low), a visual estimate of percent cloud cover, wind speed in five subjective
categories (calm, very light, light, moderate, strong) and wind direction (eight cardinal points: N, NE, E,
SE *etc*). Daily 9 am temperature (°C) and rainfall (mm) observations from the nearest Australian Bureau
of Meteorology weather station (http://www.bom.gov.au/climate/data/index.shtml) were later added to the

data table. Other variables recorded were access type (pedestrian, vehicle, combined) and distance to
parking (m) for each access point and beach length (km) for each site.

Group composition variables (adults, children, gender, *etc.*) were recorded because they may
affect the motivation or capability for walking long distances. Counts of people on the beach were
recorded because new arrivals may have to walk some distance to find their own space. Similarly, the
presence/absence of lifeguards was recorded because people may have to walk some distance to swim in
the patrolled zone. Tide was recorded because the firm intertidal zone sand at low tide can facilitate
walking longer distances. Weather variables were recorded because uncomfortable conditions can deter
walking long distances. Time of day was recorded because it is related to weather conditions and because
some persons would have been walking early before work and had limited time. Weekday was recorded
because some persons may have had more time for recreation on weekends. Distance to parking was
recorded because it could affect the motivation or capability for additional walking along the beach.
Access type was disregarded because there was little variation for this variable (96% pedestrian access
points).

All 10 activities recorded are described in Table 3. Only activities that involved walking onto and
along the beach were recorded. Joggers, runners, cyclists and horse riders were not included in this study
because, even though they often cover longer distances than walkers, those activities were uncommon.
Four-wheel drives (4WDs) were not part of this study and beach driving was permitted on only seven of
the 18 study beaches.

<INSERT TABLE 3 HERE>

Statistical analysis
Conditional inference regression trees (Hothorn *et al.* 2006) were used to discover relationships between
distance walked and predictor variables. Regression trees are a machine learning method that explains
variation in a single variable by repeatedly splitting the data into homogeneous groups using combinations
of predictor variables (De'ath & Fabricius 2000). Machine learning methods can accommodate high-
dimensional data, with many predictor variables (Breiman 2001). Conditional inference trees utilise
statistical tests to avoid variable selection bias and overfitting.

Evaluation of machine learning models concerns the ability of models to predict new or withheld
data rather than goodness of fit for a single model fitted to the all the data. A "useful" model is one that
can make acceptably accurate predictions for new data. Accuracy (square Root Mean Squared Error;
RMSE) of distance walked predictions was evaluated using five-fold cross-validation with ten repeats.
Relative performance of regression tree models was compared to a mean distance walked null model and
is reported as the reduction in cross-validated deviance: $D^2 = (MSE_{null} - MSE_{fitted}) / MSE_{null}$ (Guisan &
Zimmerman 2000).

The statistical analysis was performed in R version 3.5.2 (R Core Team 2018) and using the R
packages *party* version 1.3-1 (Hothorn *et al.* 2018) and *caret* version 6.0-81 (Kuhn *et al.* 2018).
Multiplicity adjusted *P*-values for conditional inference trees were computed using Monte Carlo
permutation tests (with 9999 resamples), the minimum number of observations in a terminal node ('leaf')
was set to 10 (5% of the sample size $n = 200$) and the minimum sum of numbers of observations in a node
in order to be considered for splitting was set to 30 (3×10).

RESULTS
A total of 200 observations of beach users were made from 20 sites on 18 beaches between 8 February and
24 March 2019. Twelve of the 20 sites (60%) were adjacent to residential areas and the other eight
(Hastings Point, Potts Point, Wooyung Corner, South Ballina, Chinamans Beach, Shark Bay, Lake
Arragan and Illaroo) were non-urban sites. Observations were nearly equally distributed between
weekdays (51%) and weekends (49%) and high (43%) and low tides (57%). Most observations were early
in the morning, between 0645–0930h AEST (interquartile range). Adults were present in 98% of
observations and the mean group size was 1.4 adults ($n = 197$; 95% CI = 1.3–1.5) . The dataset was nearly
uniformly distributed among majority female (32%), male (38%) and balanced gender (30%) groups. The
study took place at the end of a dry summer (mean rainfall 2 mm on sampling days). Mean 9 am
temperature was 25 °C, mean cloud cover was 30% and wind speed was most frequently light.

The most frequent activities observed were walking (29%), dog walking (21%) and swimming
(16%). The distribution of distances people walked had a strong positive skew (Figure 3). The mean
distance was 444 m ($n = 200$; 95% CI = 365–523 m) and the 95th percentile distance was 1683 m,
meaning that five per cent of persons/groups walked further than that distance. Walkers covered the
longest distances compared to other beach users (Figure 4). Mean distances walked were similar for
walkers with or without dogs at 748 m ($n = 41$; 95% CI = 605–892 m) and 851 m ($n = 58$; 95% CI = 667–
1035 m) respectively. For both walking activities combined, the mean distance was 809 m ($n = 99$; 95%
CI = 687–930 m) and the 95th percentile distance was 1990 m.

<INSERT FIGURE 3 HERE>

<INSERT FIGURE 4 HERE>

A conditional inference regression tree model with minimum criterion 0.95 (1 – P-value)
indicated that longer distances were associated with beach walking, with or without dogs ($P < 0.001$) and
low wind speed (calm or very light winds; $P = 0.048$) (Figure 5). Other beach users walked shorter
distances with a slight increase in the middle of the week (Wednesday and Thursday; $P = 0.02$), however
there were only 17 observations in that node and that result is not expected to appear in replicate studies.

<INSERT FIGURE 5 HERE>

The regression tree model resulted in a 38% reduction in cross-validated deviance relative to the
mean distance walked null model, however the large 443 m residual RMSE indicates large among
person/group variability in distances walked.

DISCUSSION

This study purposively selected a variety of beaches and sites and, although sampling was somewhat
haphazard, the dataset was fairly well-balanced among urban and non-urban sites, weekdays and
weekends, high and low tides and gender of beach users. It is suggested that the sample was reasonably
representative of morning, mostly local beach users in late summer in northern NSW. It would be easy to
conduct similar studies for other regions and at other times of the year, if necessary.

Walking, with or without dogs, was the most frequent beach recreation activity observed in this
study. Walking has also been reported to be the most frequent activity for other beaches and coastal
wetlands in Australia (Dowling & Weston 1999, Antos *et al.* 2007, Glover *et al.* 2011; Maguire *et al.*
2011) although driving can take over on beaches where 4WDs are permitted (Schlacher *et al.* 2013).

The 95th percentile distance for beach walkers and dogs walkers combined was 1990 m. Applying
this result, it is recommended that beach access points be approximately two km distant from shorebird
habitat to reduce human intrusions into such zones. Of course, the beach should be longer than two km
and less-developed for this separation distance to be implemented. Such beaches are also where larger
numbers of beach nesting birds can be found (*e.g.* the Australian Pied Oystercatcher Key Management
Sites that were noted in the Introduction).

Compared to flight initiation distances, a two km separation distance seems extremely large but is
consistent with the ecology of beach nesting birds and is supported by anecdotal evidence. In particular,
beach nesting Australian pied oystercatchers occupy linear territories that can be hundreds of metres in
length. For example, the maximum recorded oystercatcher breeding density for South Ballina Beach was
18 pairs in 2000 (NSW Department of Planning, Industry and Environment, *unpubl. data*), over 18.7 km
of available habitat, which is equivalent to a mean territory length of 1040 m. Large separation distances
can protect such long territories from human intrusions.

Oystercatcher nest location data from 2001–2017 show a 4.3 km gap in the distribution of
breeding oystercatchers around the village of Patch's Beach (currently with 19 dwellings) on South
Ballina Beach (NSW Department of Planning, Industry and Environment, *unpubl. data*). Harrison (2009)
suggested that this gap was the result of permanent and daily human recreation disturbance which deters
oystercatchers from occupying otherwise suitable habitat close to Patch's Beach. The equivalent 2150 m
disturbance radius (= 0.5×4300) is close to the 1990 m 95th percentile distance from this study.

Finally, it is suggested that human-wildlife coexistence strategies are a "last-resort" option, for
overdeveloped beaches that have already been let down by poor land-use planning and where human

recreation is established and prevalent. While coexistence strategies could deliver positive conservation
outcomes, managing human behaviour is known to be difficult and costly. Moreover, there is a danger that
planners will "cherry pick" elements of coexistence strategies to further the development and recreational
use of beaches.

ACKNOWLEDGEMENTS

REFERENCES

Antos, M.J., G.C. Ehmke, C.L. Tzaros & M.A. Weston 2007. Unauthorised human use of an urban
coastal wetland sanctuary: current and future patterns, *Landscape and Urban Planning* 80:173–
183.
BirdLife Australia 2018. Beach-nesting Birds. BirdLife Australia. Available at:
http://www.birdlife.org.au/projects/beach-nesting-birds (accessed 17 November 2018).
Blumstein, D.T. 2006. Developing an evolutionary ecology of fear: how life history and natural history
traits affect disturbance tolerance in birds. *Animal Behaviour* 71:389–399.
Breiman, L. 2001. Statistical modeling: the two cultures. *Statistical Science* 16:199–231.
Burger, J. 1994. The effect of human disturbance on foraging behaviour and habitat use in Piping Plover
(*Charadrius melodus*). *Estuaries* 17:695–701.
Davidson, N.C. & P.I. Rothwell 1993. Human disturbance to waterfowl on estuaries: conservation and
coastal management implications of current knowledge. *Wader Study Group Bulletin* 68:97-105.
De'ath, G. & K.E. Fabricius 2000. Classification and regression trees: a powerful yet simple technique
for ecological data analysis. *Ecology* 81:3178–3192.
Dowling, B. & M.A. Weston 1999. Managing a breeding population of the Hooded Plover *Thinornis*
rubricollis in a high-use recreational environment. *Bird Conservation International* 9:255–270.
Gill, J.A., K. Norris & W.J. Sutherland 2001. Why behavioural responses may not reflect the
population consequences of human disturbance. *Biological Conservation* 97:265–268.
Glover, H.K., M.A. Weston & G.S. Maguire 2011. Towards ecologically meaningful and socially
acceptable buffers: Response distances of shorebirds in Victoria, Australia, to human disturbance.
Landscape and Urban Planning 103:326–334.
Guinness, S.J, G.S. Maguire, K.K. Miller & M.A. Weston 2020. My dog, my beach! Attitudes towards
dog management on Victorian beaches. *Australasian Journal of Environmental Management*
27:329–342.
Guisan, A. & N.E. Zimmermann 2000. Predictive habitat distribution models in ecology. *Ecological*
Modelling 135:147–186.
Hill, D., D. Hockin, D. Price, G. Tucker, R. Morris & J. Treweek 1997. Bird disturbance: improving
the quality and utility of disturbance research. *Journal of Applied Ecology* 34:275–288.
Harrison, A.E. 2009. The ecology of two vulnerable shorebirds (*Haematopus fuliginosus* and *H.*
longirostris) in sub-tropical northern NSW, Australia: implications for conservation and
management. PhD thesis, University of New England. Armidale, New South Wales, Australia.
Hothorn, T., K. Hornik & A. Zeileis 2006. Unbiased recursive partitioning: a conditional inference
framework. *Journal of Computational and Graphical Statistics* 15:651--674.
Hothorn, T., K. Hornik, C. Strobl & A. Zeileis 2018. party: a laboratory for recursive partytioning. R
package version 1.3-1. Available at: http://party.R-forge.R-project.org
Kuhn, M. 2018. caret: Classification and Regression Training. R package version 6.0-81. Available at:
https://CRAN.R-project.org/package=caret
Lafferty, K.D., D. Goodman & C.P. Sandoval 2006. Restoration of breeding by snowy plovers
following protection from disturbance. *Biodiversity and Conservation* 15:2217–2230.
Liley, D. & W.J. Sutherland 2007. Predicting the population consequences of human disturbance for
ringed plovers *Charadrius hiaticula*: a game theory approach. *Ibis* 149:82–94.
Lord, A., J.R. Waas, J. Innes & M.J. Whittingham 2001. Effects of human approaches to nests of
northern New Zealand dotterel. *Biological Conservation* 98:233–240.

Maguire, G.S., K.K. Miller, M.A. Weston & K. Young 2011. Being beside the seaside: beach use and
preferences among coastal residents of south-eastern Australia. *Ocean and Coastal Management*
54:781–788.

Maguire, G.S., K.K. Miller & M.A. Weston 2019. Only the strictest rules apply: investigating regulation
compliance of beaches to minimize invasive dog impacts on threatened shorebird populations. Ch.
11. *In:* Makowski, C. & C.W. Finkl. (Eds.) Impacts of invasive species on coastal environments.
Coastal Research Library 29. Springer, Cham, Switzerland.

Manfredo, M.J. & A.A. Dayer 2004. Concepts for exploring the social aspects of human-wildlife
conflict in a global context. *Human Dimensions of Wildlife* 9:1–20.

Marchant S. & P.J. Higgins (Eds.) 1993. *Haematopus longirostris* Pied Oystercatcher. Pp. 716–726. *In:*
Handbook of Australian, New Zealand and Antarctic birds. Vol. 2, raptors to lapwings. Oxford
University Press, Melbourne.

NSW Department of Planning, Industry & Environment 2019. Saving our species. Pied Oystercatcher
(*Haematopus longirostris*) Key Management Sites. NSW Government. Available at:
https://www.environment.nsw.gov.au/savingourspeciesapp/project.aspx?ProfileID=10386
(accessed 26 August 2019).

NSW Government 2016. Biodiversity Conservation Act 2016. Available at:
https://www.legislation.nsw.gov.au/~/view/act/2016/63/ (accessed 18 February 2019).

Paton, D.C., M. Ziembicki, P. Owen & C. Heddle 2000. Disturbance distances for water birds and the
management of human recreation with special reference to the Coorong region of South Australia.
University of Adelaide, Adelaide, Australia.

Pienkowski, M.W. 1993. The impact of tourism on coastal breeding waders in western and southern
Europe: an overview. *Wader Study Group Bulletin* 68:92–96.

Priest, B., P. Straw, & M.A. Weston 2002. Shorebird conservation in Australia. Supplement to
Wingspan 12: 1–15.

R Core Team 2018. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. Available at https://www.R-project.org/

Schlacher, T.A., T. Nielsen & M.A. Weston 2013. Human recreation alters behaviour profiles of non-
breeding birds on open-coast sandy shores. *Estuarine, Coastal and Shelf Science* 118:31–42.

Schneider, T.J., G.S. Maguire, G.S., D.A. Whisson & M.A. Weston 2020. Regulations fail to constrain
dog space use in threatened species beach habitats. *Journal of Environmental Planning and*
Management 63: 1022–1036.

Schultz, R. & M. Stock 1993. Kentish Plovers and tourists: competitors on sandy coasts? *Wader Study*
Group Bulletin 68:83–91.

Taylor, I.R.; O.M.G. Newman; P. Park; B. Hansen; C.D.T. Minton & R. Jessop 2014. Conservation
assessment of the Australian Pied Oystercatcher *Haematopus longirostris*. *International Wader*
Studies 20:116–128.

Verhulst, S., K. Oosterbeek K. & B.J. Ens 2001. Experimental evidence for effects of human
disturbance on foraging and parental care in oystercatchers. *Biological Conservation* 101:375–
380.

West, A.D., J.D. Goss-Custard, R.A. Stillman, R.W.G. Caldow, S.E.A. le V. dit Durrell & S.
McGrorty 2002. Predicting the impacts of disturbances on shorebird mortality using a behaviour-
based model. *Biological Conservation* 106:319–328.

Weston, M.A. & Elgar, M.A. 2005. Disturbance to brood-rearing Hooded Plover *Thinornis rubricollis*:
responses and consequences. *Bird Conservation International* 15:193–209.

Weston, M.A., E.M. McLeod, D.T. Blumstein & P.-J. Guay 2012a. A review of flight initiation
distances and their application to managing disturbance to Australian birds. *Emu* 112:269–286.

Weston, M.A., F. Dodge, A. Bunce, D.G. Nimmo & K.K. Miller 2012b. Do temporary beach closures
assist in the conservation of breeding shorebirds on recreational beaches? *Pacific Conservation*
Biology 18:47–55.

Weston, M.A., T.A. Schlacher & D. Lynn 2014. Pro-environmental beach driving is uncommon and
ineffective in reducing disturbance to beach-dwelling birds. *Environmental Management* 53:999–
1004.

Table 1. Mean flight initiation distances (FIDs) for Australian Pied Oystercatchers in preceding studies. For
comparison, the mean distance walked by beach users in this study was 444 m (see Results).

Reference	Habitat	Stimulus	Mean FID (m)	Breeding status
Paton *et al.* 2000	Estuary	Human	83	Non-breeding
Blumstein 2006	Unknown	Human	39 ($n = 23$)	Unknown
Harrison 2009	Beach	Human	46 ($n = 79$)	Breeding
		Human and dog	129 ($n = 20$)	Breeding
Glover *et al.* 2011	Varied	Human	42 ($n = 21$)	Non-breeding

351 **Table 2.** Study sites and beach access points sampled in this study (north to south). Beaches known to have breeding
352 oystercatchers (OC) are recorded as 'Yes'. Access points where dogs are allowed onto the beach are recorded as
353 'Yes'. Major land uses within 200–300 m of each access point are classified into four categories: R = Residential; P
354 = Public Reserve (typically at 'peri-urban' sites); T = Tourism; C = Conservation.

Beach	Length (km)	OC	Site	Access	Dogs	Land use	Co-ordinates (S, E)
Letitia Spit	3.3		Fingal	Caravan park	Yes	T	28° 11' 45.6", 153° 33' 59.9"
				Fingal SLSC	Yes	R	28° 11' 47.3", 153° 34' 01.6"
Bogangar	8.6		Salt	Salt SLSC	Yes	R	28° 16' 38.2", 153° 34' 47.6"
Cudgera	2.8		Hastings Point	Opp. Shell stn.	Yes	P	28° 22' 10.1", 153° 34' 30.8"
Wooyung	16.6		Potts Point	Bridge	Yes	P	28° 23' 32.9", 153° 34' 05.2"
Wooyung	16.6		Wooyung Corner	Opp. c'van park	Yes	T	28° 27' 25.0", 153° 33' 14.8"
				WOO46	Yes	T	28° 27' 26.1", 153° 33' 14.6"
Tallow	7.4		Suffolk Park	Clifford St.		R	28° 41' 26.9", 153° 36' 51.3"
				Caravan park		T	28° 41' 29.9", 153° 36' 50.6"
				Alcorn St. S.		R	28° 41' 31.5", 153° 36' 50.3"
Seven Mile	8.4		Lennox Head	Lake Ainsworth	Yes	P	28° 46' 56.0", 153° 35' 39.3"
Angels	1.7		Angels (East Ballina)	Overpass	Yes	P	28° 50' 57.2", 153° 36' 05.0"
				Old 4WD track	Yes	R	28° 51' 04.8", 153° 36' 01.1"
				Car park	Yes	R	28° 51' 10.8", 153° 35' 57.8"
				Angels Beach	Yes	R	28° 51' 13.5", 153° 35' 56.3"
South Ballina	23.0	Yes	South Ballina (near c'van park)	Pedestrian		T, C[1]	28° 53' 16.1", 153° 34' 21.4"
				4WD		T, C[1]	28° 53' 17.8", 153° 34' 19.1"
			Patchs Beach	Pedestrian	Yes	R	28° 56' 59.0", 153° 30' 56.6"
Airforce	5.8	Yes	Evans Head	SLSC		R	29° 06' 44.8", 153° 26' 04.3"
Chinamans	0.8		Chinamans	Dirrawong Res.		C	29° 07' 49.5", 153° 26' 52.3"
Ten Mile	12.5	Yes	Shark Bay (Bundjalung NP)	4WD		C	29° 21' 31.4", 153° 21' 16.7"
				Picnic area		T, C	29° 21' 36.2", 153° 21' 34.3"
Iluka	2.7		Iluka	Car park	Yes	P	29° 25' 12.9", 153° 21' 48.3"
				Breakwall	Yes	P	29° 25' 18.0", 153° 21' 46.4"
Pippi	1.7		Yamba	Dolphin Park	Yes	R	29° 26' 40.6", 153° 21' 56.2"
				Dolphin Park	Yes	R	29° 26' 42.2", 153° 21' 55.3"
Plumbago	1.9	Yes	Lake Arragan camping area (Yuraygir NP)	Picnic area		T, C	29° 33' 56.7", 153° 20' 14.6"
				Camp		T, C	29° 34' 01.6", 153° 20' 14.5"
				Camp		T, C	29° 34' 05.7", 153° 20' 14.0"
				Red Cliff		T, C	29° 34' 08.7", 153° 20' 16.4"
Illaroo	9.0		Illaroo camping area (Yuraygir NP)	Steps		T, C	29° 45' 22.7", 153° 17' 31.3"
				Barbeque		T, C	29° 45' 25.2", 153° 17' 31.5"
				Gully		T, C	29° 45' 27.9", 153° 17' 31.6"
				Picnic area		T, C	29° 45' 32.1", 153° 17' 34.5"
Wooli	6.7	Yes	Wooli	Scope St.	Yes	R	29° 51' 47.8", 153° 16' 04.1"
				Firth Ln.	Yes	R	29° 52' 17.9", 153° 15' 58.9"
				Water tower	Yes	R	29° 52' 22.4", 153° 15' 58.6"
				Main Beach	Yes	R	29° 52' 32.2", 153° 15' 58.1"
				South Tce.	Yes	P	29° 52' 47.6", 153° 15' 58.0"
Sandy	1.1		Sandy Beach	Car park		R[2]	30° 08' 57.2", 153° 12' 00.2"
				Park		R[2]	30° 09' 04.3", 153° 12' 00.2"
				Park		R[2]	30° 09' 08.6", 153° 12' 00.8"
Sapphire	2.2		Sapphire Beach	Cafe	Yes	T, C[2]	30° 13' 30.8", 153° 09' 20.8"
				Private	Yes	R[2]	30° 13' 37.8", 153° 09' 17.7"
				Caravan park	Yes	T[2]	30° 13' 45.2", 153° 09' 14.3"
				Park	Yes	R[2]	30° 13' 49.0", 153° 09' 13.7"
				Lakeside Dr.	Yes	R[2]	30° 13' 51.1", 153° 09' 13.3"

Acronyms: 4WD = four-wheel drive; NP = National Park; SLSC = Surf Life Saving Club

[1] The northern end of South Ballina Beach adjoins the Richmond River Nature Reserve (Conservation) however the major land use near the caravan park is tourism.

[2] Sandy and Sapphire Beaches adjoin the typically narrow Coffs Coast Regional Park (Conservation) and the major land uses around those areas sampled were Residential and Tourism.

356 **Table 3.** Descriptions of beach recreation activities recorded in this study.

Activity	Description
Walking (WK)	Walking along the beach, usually direct and close to the shoreline. Walkers walk for some distance and then turn back or exit the beach via a different access point.
Dog walking (WD)	Walking along the beach accompanied by one or more dogs, on or off leash.
Swimming (SW)	Does not include wading (to approximately knee-depth).
Surfing (SF)	Mostly on surfboards.
Picnic (PC)	Picnics generally involved multiple persons and a variety of activities. Features of this activity can include beach shelters, food and beverages, recreational equipment and the group stays in one location for an extended period.
Sunbathing (SB)	Self-explanatory.
Looking (LK)	Viewing the beach or ocean, checking the beach, fishing or surf conditions and often where there is seating on the beachfront.
Fishing (FI)	Recreational line fishing.
Playing (PL)	Young children playing in the sand. Unlike picnickers, these groups don't bring so much equipment, don't stay long and adults present are supervising and not engaged in other recreation.
Photography (PH)	Recreational photography.

358
359 **Figure 1.** Beach user intrusions from an access point (arrowed) into a shorebird habitat zone (grey area). The
360 distributions are (from top to bottom) a frequency distribution of distances walked ($f(d)$), a cumulative frequency
361 distribution ($p(d)$) and a distribution of intrusions ($V \times [1 - p(d)]$). V is the number of people using the access point
362 and walking along shore towards the habitat zone. The separation distance is s and R is the number of intrusions into
363 the habitat zone.

364
365 **Figure 2.** Map of the far north coast of New South Wales showing the location of access points (white-filled circles)
366 where beach recreation was studied (see Table 2 for details of beaches and sites). The background image shows
367 human population densities (scale 0–2000 persons.km^{-2}; darker values indicate higher densities) from the Australian
368 Population Grid 2011 (https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.007). Text annotations identify larger
369 coastal towns and cities. The inset map shows the location of the study region within Australia.

370
371   **Figure 3.** Distances walked for all beach users (n =200). Vertical dotted lines indicate the mean and 95th percentile
372   distances (dotted and dashed lines respectively).

373
374   **Figure 4.** Distances walked for each activity: FI = fishing, LK = looking, PC = picnic, PH = photography, PL =
375   playing, SB = sunbathing, SF = surfing, SW = swimming, WD = dog walking, WK = walking. Boxes show the first
376   quartile, median and third quartile, whiskers extend to a maximum of 1.5 times the interquartile range and data
377   outside the whiskers are plotted as individual points (white-filled circles).

378
379 **Figure 5.** Conditional inference regression tree for distances people walked. The left branch indicates that:1) beach
380 walkers (WD = dog walking, WK = without dogs) typically covered longer distances than did other beach users
381 (right branch), and 2) distances walked decreased at higher wind speeds (> 1 = light to moderate; stronger winds
382 were not observed)..