Antimicrobial resistance gene levels are independent of levofloxacin administration but related to

abundance of specific pathobionts in dementia patients

Aislinn D. Rowan-Nash[1], Rafael Araos[2,3,4], Erika M.C. D'Agata[5], & Peter Belenky[1*]

[1] Department of Molecular Microbiology and Immunology, Brown University, Providence RI,

USA

[2] Instituto de Ciencias e Innovación en Medicina (ICIM), Facultad de Medicina Clinica Alemana

Universidad del Desarrollo, Chile

[3] Millenium Nucleus for Collaborative Research on Bacterial Resistance (MICROB-R), Chile

[4] Advanced Center for Chronic Diseases (ACCDiS), Facultad de Medicina Clinica Alemana

Univarsidad del Desarrollo, Santiago Chile.

[5] Infectious Diseases Division, Rhode Island Hospital, Warren Alpert Medical School of Brown

University, Providence, RI, USA

* indicates corresponding author

PB: peter_belenky@brown.edu

ABSTRACT

Background: The issue of antimicrobial resistance continues to grow worldwide, and long-term

care facilities are significant reservoirs of antimicrobial-resistant organisms, in part due to high

frequency of antimicrobial use. Patients with advanced dementia are particularly vulnerable to

multidrug-resistant organism acquisition and antimicrobial overuse, which has negative

consequences for the gut microbiome and can contribute to the selection and propagation of

antimicrobial resistance genes. In this study, we examined the impacts of the fluoroquinolone

antimicrobial levofloxacin on the gut microbiome of advanced dementia patients and the

correlation between abundance of pathogens and antimicrobial resistance genes.

Results: We did not find significant impacts of levofloxacin on the diversity, composition,

function, or resistome of the gut microbiota of this population. In fact, we observed significant

inter- and intra-subject heterogeneity in the composition of the microbiota, suggesting temporal

instability. However, we were able to link the antimicrobial resistance gene burden in a sample

with the relative abundance of several pathobionts – particularly *Escherichia coli*, *Proteus*

mirabilis, and *Enterococcus faecalis*, as well as less-prevalent species including *Providencia*

stuartii and *Staphylococcus haemolyticus*. Furthermore, we used metagenomic assembly and

binning to demonstrate that these species had higher genomic resistance gene levels than common

gut commensals, and we were able to predict antimicrobial resistance gene burden from the relative

abundances of these species in a separate, larger cohort from the same population.

Conclusions: We observe that levofloxacin did not have an impact on the microbiota, potentially

due to high host absorption rate of this antimicrobial and high intra-subject temporal variability.

However, we found that the relative abundances of several pathobionts were correlated with and

were even predictive of the level of antimicrobial resistance genes in corresponding samples, and

that these species carried high levels of resistances genes in their assembled genomes. Given the

high frequency with which these species were found at high levels in this population and the

underlying vulnerability to infection with multidrug resistant organisms of advanced dementia

patients, attention to microbial blooms of these species may be warranted.

Keywords: antimicrobial resistance, pathobiont, dementia, microbiome, metagenome assembly

BACKGROUND:

It is well-recognized that there is a growing threat of antimicrobial-resistant (AMR) bacterial strains that threaten the health and lives of millions worldwide. In the United States alone, the Centers for Disease Control and Prevention estimates that at least 2 million people get an AMR infection each year, and at least 23,000 die as a result[1]. A number of factors have driven the rise in AMR bacteria worldwide, including overprescription of antibiotics in the healthcare setting, over-the-counter access to antibiotics in some countries, and widespread use of antibiotics in animal husbandry for non-veterinary purposes[2-4]. Concerningly, hospitals and other medical institutions are frequent sites of AMR bacteria acquisition, where patients may already be ill or immunocompromised, antimicrobial use is common, and patient-to-patient transmission of AMR isolates can occur via inadequate hygiene or environmental contamination[5-8]. For example, AMR bacteria are highly prevalent in nursing homes, with estimates that over 35% of nursing home residents are colonized with multidrug resistant organisms (MDROs)[9-14]. This is particularly problematic in light of the fact that elderly patients in long-term care facilities may be frequently hospitalized, potentially serving as a mode of bidirectional transport of MDROs between healthcare facilities[15-17]. They are also prone to infections and are frequently treated with antimicrobials[18-20], which has long been associated with acquisition of MDROs and may not always be indicated[12, 17, 20-28].

The problem of MDRO prevalence and inappropriate antimicrobial use is of particular relevance in elderly subjects with advanced dementia, a population which receives extensive antimicrobial treatment which becomes more frequent closer to death, calling its benefit and effectiveness into question[28, 29]. Accordingly, advanced dementia specifically has been shown to be a risk factor of MDRO colonization[13, 30]. To examine this issue, the Study of Pathogen

Resistance and Exposure to Antimicrobials in Dementia (SPREAD) was undertaken from 2009-

2012 in order to analyze MDRO acquisition and appropriateness of antimicrobial prescription in

elderly adults with advanced dementia residing in nursing homes[31]. Supporting the widespread

nature of MDRO carriage in this population, analysis of SPREAD subjects revealed that there were

significant baseline levels and new acquisitions of MDROs, and that there was notable spread of

MDRO strains within and even between nursing home facilities[28, 32].

In addition to potential facilitation of MDRO acquisition or spread, antimicrobial overuse

may also have negative impacts on the diversity, composition, or function of the gut microbiota,

which may already be vulnerable in elderly populations. Healthy younger adults tend to have a

fecal microbiome characterized by relatively high diversity of species and populated primarily by

members of the phyla *Bacteroidetes* and *Firmicutes*, largely obligate anaerobes which exist in

homeostasis with the intestinal epithelium[33-37]. However, it has been found that during

senescence, the gut tends to have higher levels of *Bacteroidetes* and *Proteobacteria* and harbors

higher levels of facultative aerobes and potential pathobionts, including *Enterobacterales* such as

E. coli [36, 38-45]. These changes become more pronounced as aging progresses, and several

studies have indicated that age-related alterations to the gut microbiota are relatively minor in

septuagenarians, but become more pronounced over time and are clear in centenarians and

supercentenarians[39, 43, 46-48]. This is likely due to a number of factors, including the decline

of immune function, onset of age-related diseases (including metabolic disorders), changes to diet

and mobility, and the increased likelihood of medication utilization and/or hospitalization[42, 49].

However, lifestyle of elderly adults has an important impact, as research suggests that community-

resident elderly subjects have a distinct and more diverse microbiome compared with those of their

hospitalized or institutionalized peers, which was suggested to be at least in part due to nutritional

differences[49, 50]. Furthermore, reduced microbiome diversity has been associated with

increased frailty of elderly subjects[49, 51]. Accordingly, given that the microbiomes of

institutionalized elderly patients are perhaps already at risk, understanding the impacts of

antimicrobial use and MDRO acquisition on this population is of importance.

We analyzed the gut microbiomes of eleven subjects from SPREAD to examine the impact

of antimicrobial use on the gut microbiota composition, function, and antimicrobial resistance gene

(ARG) profile of elderly dementia patients. These subjects were chosen as they were the largest

cohort who had received a single antimicrobial (levofloxacin) during the collection period, and we

anticipated that this intervention could have an impact on the already-vulnerable microbiota of this

elderly, institutionalized cohort. Levofloxacin is an antimicrobial of the fluoroquinolone class with

high oral bioavailability[52-54] which has been found to reduce levels of Gram-negative aerobic

bacteria – including *Proteobacteria* and particularly *Enterobacterales* – in the fecal microbiota[55-

61], although fluoroquinolone resistance among this taxon has been spreading[62-69]. A

maximum of five rectal swab samples, collected every three months, were taken from each subject,

and both 16S rRNA amplicon and shotgun metagenomics sequencing were performed. We

analyzed alpha and beta diversity, taxonomic composition, functional potential, and antimicrobial

resistance gene profiles before and after administration of levofloxacin. Interestingly, we did not

find a specific impact of levofloxacin on any of these measures. However, we did find an

association between blooms of particular enteric species and ARG burden, including in samples

where MDRO were not detected by culture, suggesting that certain pathobionts carrying high ARG

burdens may frequently colonize this population and that metagenomics may allow detection of

resistant bacteria not flagged by culture-based methods.

RESULTS

Overview of Subjects

Elderly patients in long-term care facilities, and particularly patients with advanced

dementia, are frequently exposed to antimicrobials and are at high-risk of acquisition and carriage

of MDRO[9-13, 18-20, 27-30, 32]. We were interested in whether levofloxacin, one of the most

commonly prescribed antimicrobials, had impacts on the gut microbiota and resistome of these

subjects. We analyzed up to five rectal swabs, taken every three months over the course of a year,

from eleven subjects in the SPREAD cohort[31], using both 16S rRNA and shotgun metagenomics

sequencing (Figure 1A). During their participation in the study, these subjects had received only

levofloxacin (average course of eight days), which has previously been shown to decrease the

proportion of the *Enterobacterales* order of *Proteobacteria*[55-61]. Of the eleven subjects, all but

Subject I were female and all but Subject G were white. They ranged in age from 72 to 101, and

six members of the cohort did not survive for the full year of the study (Additional Table 1). All

but two subjects (C and G) resided in different nursing homes. Overall, there were 38 samples for

metagenomics sequencing (Additional Table 2). Culture-based methods indicated that four of the

eleven subjects acquired a MDRO during the study: Subject A acquired methicillin-resistant *S.*

aureus (MRSA) at the 12-month timepoint, Subject B acquired multidrug-resistant *E. coli* at the

3-month timepoint, and Subjects C and D both acquired multidrug-resistant *P. mirabilis* at the 3-

month timepoint (Additional Table 1).

Alpha and Beta Diversity Metrics

We first used the metagenomic sequencing data to compare the alpha diversity, or the

diversity within each sample, of samples collected before and after levofloxacin administration.

According to Shannon's Diversity Index, which incorporates both richness and evenness of

samples, there was no significant difference between the pre- and post-levofloxacin samples

(Figure 1B). Furthermore, the alpha diversity was variable over time even within the same subject,

and there was no clear trend of recovery in alpha diversity after antibiotic cessation. This suggests

a degree of temporal instability, in which the richness and/or evenness of the samples varies

changes over time.

We then examined beta diversity, or the diversity between samples. We utilized the Bray-

Curtis Dissimilarity metric, which considers the identity and abundance of taxa shared between

samples. Plotting this metric in a principal coordinate analysis (PCoA) revealed no apparent

pattern of clustering based on either subject or sample collection point relative to levofloxacin, and

in fact, samples from the same subject were often located quite distantly from one another (Figure

1C). We then compared the within-subjects dissimilarity of sequential samples within a subject

when both were pre-levofloxacin, both were post-levofloxacin, or one sample was pre- and one

was post-levofloxacin; there was no significant difference between any of the groups (Figure 1D),

suggesting that levofloxacin was not associated with community disruption. Furthermore, while

within-subject dissimilarity was lower than between-subjects dissimilarity, the effect size was low

(0.7013 vs. 0.7712, respectively; Figure 1E).

Taxonomic Composition

We utilized Kraken2 in conjunction the with Bayesian Reestimation of Abundance with

KrakEN2 (Bracken2) pipeline to assign taxonomy to our metagenomic sequencing samples[70,

71]. Corresponding to the high between-subjects beta-diversity, the taxonomic composition of the

gut microbiome varied significantly between subjects. As is typical for the human gut microbiome,

most bacteria belonged to the five major phyla of *Firmicutes*, *Bacteroidetes*, *Proteobacteria*,

Actinobacteria, and *Verrucomicrobia*. However, consistent with the high within-subjects beta

diversity, the dominant phyla varied greatly even between samples from the same subject

(Additional Figure 1); for example, the most abundant phylum in Subject E was *Bacteroidetes* at

two timepoints, *Proteobacteria* at two timepoints, and *Firmicutes* at one (Additional Figure 1F).

Overall, the most abundant phylum was *Actinobacteria* in three samples, *Bacteroidetes* in

seventeen samples, *Firmicutes* in seven samples, and *Proteobacteria* in eleven samples

(Additional Figure 1A-L); averaging across all samples, *Bacteroidetes* was highest at 34.2%,

followed by *Proteobacteria* (26.9%), *Firmicutes* (23.3%), and *Actinobacteria* (11.2%) (Additional

Figure 1A). Qualitatively, many of the samples from this population represent highly divergent

and dysbiotic microbiomes compared with what is typically seen with younger subjects, in which

Proteobacteria in particular make up a much smaller proportion of the microbiome than in these

elderly dementia subjects[33].

The genus- and species-level taxonomic composition was also variable. Blooms of

potential   pathogens[72],   including   *Campylobacter   ureolyticus*[73]*,   Corynebacterium*

urealyticum[74], *Enterococcus faecalis*[75, 76], *Escherichia coli* [77, 78], *Oligella urethralis*[79-

82], *Proteus mirabilis*[83, 84], *Providencia stuartii*[85, 86], *Pseudomonas aeruginosa*[87, 88],

Staphylococcus aureus[89-91], and *Staphylococcus haemolyticus*[92-94], were fairly common,

both before and after levofloxacin administration (Figure 2A, Additional Figure 2). Across

subjects, even baseline samples varied in composition, as expected from beta-diversity analysis.

Averaging across all samples, the single most-abundant species was *E. coli*, further supporting the

qualitatively dysbiotic nature of the gut microbiome of this cohort (Figure 2A). Despite the high

proportion of members of *Enterobacterales* in this cohort, Linear Discriminant Analysis Effect

Size (LEfSe) analysis[95] did not reveal biomarkers for pre- or post-levofloxacin samples at the

phylum, genus, or species level. Full data on taxonomic composition at the phylum and species

levels can be found in Additional Data 1.

As we had access to full 16S rRNA and shotgun metagenomics data for our samples, we

compared their taxonomic identifications at the genus level. The two methods of analysis were

generally consistent, and blooms of prominent genera (including *Escherichia, Proteus,*

Enterococcus, Providencia, Staphylococcus, and *Bacteroides*) were generally detected by both

analysis pipelines (Additional Figure 3A). Metagenomics analysis was unsurprisingly able to

detect more distinct genera, and of the genera that were called by both pipelines, LEfSe analysis

revealed biases in both methods. For example, metagenomics analysis by Kraken2 and Bracken2

detected higher levels of *Bacteroides*, while 16S rRNA analysis with Quantitative Insights Into

Microbial Ecology 2 (QIIME2)[96] detected higher levels of *Ruminiclostridium*. Full data on

taxonomic abundances at the genus level can be found in Additional Data 1 for metagenomics and

Additional Data 2 for 16S rRNA.

Functional Potential

We used the Human Microbiome Project Unified Metabolic Analysis Network 2

(HUMAnN2) pipeline[97] to analyze the genetic content of the metagenomic samples. We utilized

LEfSe to compare community function at the Kyoto Encyclopedia of Genes and Genomes (KEGG)

ortholog, Gene Ontology (GO) term, and MetaCyc pathway levels. As in the taxonomic analysis,

there were no significant biomarkers of either pre-or post-levofloxacin administration samples.

However, while the taxonomic profile of the samples varied greatly, the functional capacity of the

samples was fairly consistent across samples (Additional Figure 4). Full data on functional

potential can be found in Additional Data 3.

Antimicrobial Resistance Gene Profile

We used the DeepARG machine-learning program[98] to detect resistance genes in the

metagenomic samples. Across all samples, the most abundant class of ARG was "multidrug",

followed by "macrolide-lincosamide-streptogramin" (MLS), and "tetracycline". The most

common specific gene detected was the multidrug resistance *rpoB2* variant of the RNA polymerase

beta subunit, followed by the MLS resistance gene *macB* and a multidrug ABC transporter (Figure

2B). LEfSe analysis revealed no ARG biomarkers of either pre- or post-levofloxacin samples. Full

data on ARG composition can be found in Additional Data 4.

However, we were able to detect changes in specific ARG classes and genes that

corresponded with the detection of antimicrobial-resistant organisms in two subjects. Subject A

acquired MRSA at the 12-month timepoint, and a bloom of this species to 25.0% could be detected

in the metagenomic taxonomic data (Figure 3A, Additional Figure 2B). While the overall level of

ARGs did not notably increase at this timepoint, there was a clear expansion in beta-lactam

resistance genes (Figures 3B, Additional Figure 5B), including the *mecA*/*mecR1*/*mecI* operon,

which regulates expression of the low-affinity penicillin-binding protein *mecA* (PBP-2A)[99-102]

(Figure 3C). This operon is characteristic of MRSA strains[99-102], supporting the culture-based

classification of this *S. aureus* isolate as MRSA.

Similarly, Subject B acquired multidrug-resistant *E. coli* (resistant to the beta-lactams

ampicillin/sulbactam, cefazolin, ceftazidime, and ceftriaxone and to the fluoroquinolone

ciprofloxacin) at the three-month timepoint, and the proportion of this species expanded to 47.3%

of the population in the corresponding sample (Figure 3D, Additional Figure 2C). Accordingly,

this sample showed a notable increase in the relative abundance of ARGs, which was in large part

driven by an increase in a number of multidrug resistance genes (Figure 3E); there was also a clear

increase in several beta-lactam resistance genes, including the low-affinity penicillin-binding

protein genes *PBP-1A, PBP-1B,* and *penA* (*PBP2*) as well as class C beta-lactamase genes[103-

108], and several fluoroquinolone resistance genes, including the transporters *patA* and *mdtK* [109-

113] (Figure 3F-G).

Despite the acquisition of multidrug-resistant *P. mirabilis* at the three-month timepoint in

Subjects C and D, there was no corresponding increase in ARGs. ARG levels stayed approximately

the same in Subject C (0.372% at baseline and 0.384% at three months) and decreased in Subject

D from 0.482% at baseline to 0.364% at the three-month timepoint (Figures 2B, Additional Figure

5D-E). However, this corresponds to the taxonomic data; levels of *P. mirabilis* were low and stable

in Subject C (0.55% at baseline and 0.61% three months later), and while *P. mirabilis* made up

13.8% of the population at baseline in Subject D, it underwent a reduction to 2.3% at the three-

month timepoint (Figures 2A, Additional Figure 2D-E). Taken together, this data indicates that

our metagenomics pipeline can detect blooms of AMR pathogens and that the corresponding

change in ARG levels aligns with culture-based detection of MDROs. At the same time,

metagenomic analysis of some samples found blooms of pathogens and ARGs that were not

associated with culture-based MDRO detection.

Attribution of ARG Density to Specific Species

While total ARG density within samples did not vary by levofloxacin administration, there

was significant variability between samples. In fact, most samples had similar baseline levels of

ARGs of 0.3% to 0.4% of the total reads, while only a few samples rose above this value to between

0.6% and 0.8%. Close inspection of the taxonomic composition of the samples revealed that

samples with higher levels of ARGs tended to have blooms of one or more of the *Proteobacteria*

species *E. coli* and *P. mirabilis* and the *Firmicutes* species *E. faecalis*, strains of which are common

pathobionts[114-120] (Figure 4A). Confirming this association, correlation analysis between ARG

levels and the sum of the relative abundances of these three species showed a strong and significant

positive correlation (r = 0.791, R^2 = 0.6254, p <0.0001, Pearson's correlation; Figure 4B). This

suggests that in samples with higher-than-baseline ARG levels, ARG abundance is being driven

by high relative abundance of these three species.

However, there were two notable exceptions: Samples E9 and H6 had high levels of ARGs

without corresponding blooms of these three species. However, *P. stuartii* bloomed to 41.9%

relative abundance in Sample E9 and *S. haemolyticus* bloomed to 36.9% in Sample H6 (Figure

2A, Additional Figure 2F&I). Both species have long been associated with AMR phenotypes[93,

94, 121-127] and were not found at high levels in other samples, but could explain the higher ARG

abundance in these samples (Figure 4A). Supporting this possibility, an examination of the ARGs

in Sample H6 showed a distinct profile relative to other samples, with high levels of staphylococcal

resistance genes including fluoroquinolone resistance gene *norB* and macrolide-streptogramin

resistance gene *msrA* [128-131](Figures 2B, 3I-L). Accordingly, addition of *P. stuartii* and *S.*

haemolyticus abundances to the analysis resulted in a stronger correlation (r = 0.933, R^2 = 0.8706,

p < 0.0001, Pearson's correlation; Figure 4C).

To more rigorously examine the relationship between the species of interest and ARG

levels, we performed metagenomic assembly and binning to compare the levels of ARGs in these

organisms to levels in other common and abundant species, including likely commensals and

potential pathogens (Figure 4D). Specifically, we analyzed bins that passed various quality

controls (Additional Table 3) and corresponded to species identified by Kraken2/Bracken2 to

make up greater than 0.1% of their source samples (Additional Table 4).

As anticipated, we found that the levels of ARGs in bins from *E. coli* and *P. mirabilis* were

consistently high compared to other species analyzed. In fact, *E. coli* had the highest average ARG

density of any species analyzed, while *P. mirabilis* was the fifth-highest. Notably, the ARG

composition of the bins of these species from samples in which MDROs were detected (B3, C3,

and D3) did not appear to be different from those of other samples (Additional Figure 6A-B),

although it is possible that some resistance genes were carried on plasmids that were not assembled

into genomes. *P. stuartii* had the second-highest average ARG density, reflecting the expansion of

ARGs detected in sample E9, where this species bloomed to 41.9% of the population. The third

and fourth positions were taken by the single bins constructed for *Klebsiella oxytoca* and

Morganella morganii, other *Proteobacteria* with pathogenic potential[132-134]. *P. aeruginosa*

bins rounded out the top six, with similar levels to the other top species. However, as *K. oxytoca*

and *M. morganii* were never present at greater than 3% and *P. aeruginosa* bloomed in only two

samples, they did not significantly contribute to overall ARG density in the cohort. Importantly,

high ARG density was not a universal feature of *Proteobacteria*, or even of pathogenic

Proteobacteria; bins constructed for the *Campylobacter* species *C. hominis* and *C. ureolyticus* had

universally low ARG levels. Additionally, while we could not construct a high-quality bin for *O.*

urethralis, the low ARG densities in the samples in which this species bloomed (C0 and C3)

suggests that it also has low genomic ARG content. This suggests that high ARG density among

the *Proteobacteria* analyzed was restricted to the *Gammaproteobacteria* class, primarily of the

order *Enterobacterales* but also including *Pseudomonadales*.

293   We were only able to construct two good-quality bins for *E. faecalis*, which varied in their

ARG levels, particularly on the basis of bacitracin resistance. On average, while the two bins did

not have ARG levels as high as the *Proteobacteria* of interest, they did rank among the highest of

the *Firmicutes* bins tested. We were also able to create a single bin for *S. haemolyticus* from

Sample H6 in which it made up 36.9% of the population. This bin had an ARG density higher than

the average for any other non-*Proteobacteria* species, supporting its role in the high ARG levels

found in the corresponding sample. As expected from the analysis of the total ARG population of

that sample (Figure 3G), the staphylococcal resistance genes *norB* and *msrA* were found in this

bin. We were also able to create two bins for *S. aureus*, including from sample A12 where MRSA

was detected. The A12 bin contained the characteristic MRSA gene *mecA* while the H6 bin did

not, suggesting that the *S. aureus* strain found in H6 was not MRSA (Figure S6C). In general, bins

from the phyla *Actinobacteria* (including *Bifidobacterium* and *Corynebacterium* species) and

Bacteroidetes (including *Bacteroides* and *Parabacteroides* species) had low ARG levels. Full data

on the ARGs and classes found in species-level bins can be found in Additional Data 4.

Prediction of ARG Density from Species Abundances

Our initial analysis only considered the eleven subjects for whom we had longitudinal

metagenomics data due to their receiving levofloxacin. We also had access to a larger dataset:

shotgun metagenomics had been performed on a further 67 samples for a related study. In this

case, the data was not longitudinal and encompassed an array of antibiotic treatment conditions

across 67 subjects, providing a diverse set of taxonomic and ARG data on which to test whether

the relationship between *E. coli*, *P. mirabilis*, and *E. faecalis* and ARG density held true. As an

initial test, we performed the same correlation analyses between species of interest and ARG levels

as on the levofloxacin dataset, finding that both the simple and complex models showed significant

correlation ($r = 0.7179$, $r^2 = 0.5154$, $p < 0.0001$ and $r = 0.7627$, $r2 = 0.5817$, p<0.0001, respectively;

Pearson's correlation; Figure 5A-B). This provided initial support for the trend being present in

the wider dataset.

We then created a multiple linear regression model to predict ARG density using the

relative abundances (RA) of the three main species of interest in the initial levofloxacin dataset,

with the following equation: (ARG density) = 0.003482 + 0.006221(*E. coli* RA) + 0.006248(*P.*

mirabilis RA) + 0.006920(*E. faecalis* RA) (Figure 5B). We then used this equation to predict the

ARG density in the larger metagenomics dataset and found that it was able to accurately predict

the true ARG level of those samples, with predicted and actual values correlating significantly (r

= 0.7139, r^2 = 0.5096, p<0.0001; Pearson's correlation; Figure 5C). As before, there were a few

notable outliers with higher ARG levels than predicted by the model; those three samples

contained high levels of *P. stuartii*, *P. aeruginosa*, or *Klebsiella pneumoniae* This maps well to

the fact that we observed high levels of ARGs in bins constructed from *P. stuartii*, *P. aeruginosa*,

and the related species *K. oxytoca* (Figure 4D).

We also created a multiple linear regression model that incorporated the relative

abundances of *P. stuartii* and *S. haemolyticus*, which caused outliers from the original species-

ARG correlation: (ARG density) = 0.003253 + 0.006715(*E. coli* RA) + 0.006748(*P. mirabilis* RA)

+ 0.003461(*E. faecalis* RA) + 0.01123(*S. haemolyticus* RA) + 0.007569(*P. stuartii* RA) (Figure

5E). As before, we tested this equation against the larger dataset, and found that it slightly

increased the accuracy of the predictions; specifically, it eliminated the outlier which had high *P.*

stuartii levels and slightly improved the correlation between predicted and actual ARG levels (r =

0.7753, r^2 = 0.6012, p<0.0001; Pearson's correlation; Figure 5F). However, the simpler model is

more broadly applicable, as blooms of *P. stuartii* and *S. haemolyticus* are relatively uncommon.

Similarly, while *Klebsiella* spp. and *P. aeruginosa* may also contribute to high ARG density in

samples, they do not bloom as commonly in this cohort as the core predictive species of *E. coli*, *P.*

mirabilis, and *E. faecalis*.

These results indicate that in this population, levels of only a few key species could predict

the majority of ARG abundance beyond background levels. Both the core predictive species (*E.*

coli, *P. mirabilis*, *E. faecalis*) and others that are associated with high ARG levels in samples (*P.*

stuartii, *S. haemolyticus*, *P. aeruginosa*, *Klebsiella* spp.) are pathogens and/or pathobionts.

Monitoring levels of these species may be helpful in elderly, institutionalized populations, as these

patients may be vulnerable to developing or transmitting AMR infections from high-level carriage

of these species.

DISCUSSION

Overall, we found that the microbial composition of the gut microbiome of elderly patients

with advanced dementia was quite variable, both between subjects and over time within the same

subject. Even in the absence of antimicrobial treatment, there was notable fluctuation in the

abundance of a number of species, including pathobionts such as *E. coli*, *P. mirabilis*, and *E.*

faecalis. When comparing the taxonomic composition, functional potential, and resistome of pre-

and post-levofloxacin samples, we did not find any significant differences. One potential reason

for this finding is that oral levofloxacin is well-absorbed by the host, with greater than 99%

bioavailability[52, 53, 135-137], and therefore may not be directly available to the luminal

microbiota of the lower gastrointestinal tract at high levels. Furthermore, other studies have

suggested that levofloxacin has a relatively minor impact on the gut microbiome, primarily

reducing levels of *Enterobacterales*[55-61], and it may be less-associated with *Clostridiodes*

difficile-associated diarrhea outbreaks than other antimicrobials, including other

fluoroquinolones[138].

Additionally, in this cohort, levofloxacin was typically administered at least two weeks

prior to collected timepoints, potentially allowing sufficient time for the microbiome to recover

from or shift away from its immediately post-antibiotic state. Furthermore, the impacts of

levofloxacin on the gut microbiome may be dependent upon the initial state upon administration.

If the microbiome is initially relatively diverse and healthy, antibiotic administration may be

disruptive and allow blooms of atypical dominant species such as members of *Proteobacteria*;

such an occurrence might be observed in Subject F, where a diverse *Bacteroides*-dominated

microbiome was overtaken by several *Enterobacterales* after levofloxacin treatment (Additional

Figure 2G). Alternatively, if the microbiome is initially dominated by one or more pathogens,

antimicrobial administration may correct such blooms and allow for the restoration of a diverse

community, as might have occurred in Subject E as a *P. stuartii* bloom was eliminated (Additional

Figure 2F).

Finally, since the pre-existing temporal instability of this community was high,

levofloxacin-related changes may not be detectable through the noise of this cohort's general

microbiome instability. In contrast to our observations, studies in adults have generally found that

the within-subjects dissimilarity is much lower than between-subjects dissimilarity, in line with

the fact that the gut microbiome tends to be relatively stable within the same subject over time –

including in an elderly cohort[33, 34, 41, 139, 140]. This suggests that the gut microbiomes of the

subjects in this study were less stable than that of other cohorts, potentially suggesting that this

institutionalized population with advanced functional impairment is more prone to infections or

has weaker immune systems than young healthy adults or even community-resident elderly adults.

Interestingly, despite the taxonomic variability, the functional composition of the cohort was

relatively similar across samples and subjects. This is in line with previous studies of the human

gut microbiome, which suggest that variable taxa can fill the same functional niches, resulting in

a more similar functional composition across individuals despite inter-individual differences in the

taxonomic composition[33, 35, 141].

As all of the subjects had been given an antibiotic, we were particularly interested in the antibiotic resistance profile of the subjects before and after levofloxacin administration. However, as observed in the taxonomic and functional data, there was no apparent association of any ARG genes or classes with either pre- or post-levofloxacin status. This may be due to the fact that levofloxacin did not have any specific impacts on the resistome of this cohort, or due to the factors that may have concealed any impacts of levofloxacin, as discussed above. However, we were particularly intrigued by the finding that ARG density in a particular sample could be linked to the abundance of a few key species. *E. coli*, *P. mirabilis*, and *E. faecalis* are all pathobionts that are often found at low levels in a healthy microbiome, but bloomed frequently at various timepoints across a majority of our subjects. All three species can cause severe illness, have been previously observed to colonize nursing home residents, and include well-known multidrug-resistant strains[11-13, 28, 30, 32, 68, 75, 114-120, 142]. In fact, three of the subjects (B, C, and D) are known to have acquired multidrug-resistant strains of *E. coli* and *P. mirabilis* during the study. However, we observed an association between these three species and ARG levels across the entire sample set (Figure 4B), and the ARG composition of the bins of *E. coli* and *P. mirabilis* from samples where MDROs were detected were not distinct from their other bins (Additional Figure 6A-B). This suggests that metagenomic sequencing may allow the identification of antimicrobial-resistant organisms that escape detection via culture-based techniques, although it is also possible that the multidrug-resistant isolates contained ARG-carrying plasmids that were not captured by our assembly and binning strategy.

A major implication of this finding is that metagenomic analysis could be a particularly useful tool to track antimicrobial resistance in institutions like nursing homes and hospitals, particularly with the capability to construct contigs and bins that allow examination of specific

genomes. In this case, it has allowed us to connect the high levels of ARGs in certain samples with

correspondingly high levels of specific pathobionts, which had high proportions of ARGs within

their genomes even in samples where MDROs were not detected. In a vulnerable population

already prone to infections and carriage of MDROs, metagenomics could be a useful surveillance

tool to assess the prevalence or transmission of ARGs in long-term care facilities.

Importantly, all of the subjects in our study were institutionalized in nursing homes, and

there exists significant potential for transfer of bacteria between patients. As all but two of our

subjects (C and G) lived in different homes, we could not directly examine this possibility

ourselves, but it is possible that the high abundance of pathobionts and/or ARGs in our cohort is

related to the spread of isolates within nursing homes. This also raises the possibility that we would

not find a similar association between pathobionts and ARG levels in a healthy or community-

based elderly cohort, who might be less likely to harbor or transmit such high levels of these

bacteria. However, if an association between particular "sentinel" species and ARGs holds true in

other elderly institutionalized populations, qPCR detection of the loads of these such pathobionts

may allow for prediction of resistant bacterial outbreaks before they occur.

In addition to the increased potential for spread of resistant strains through institutions,

there are some other potential explanations for the association between ARGs and these particular

species. In particular, all of the species that we found to be associated with ARG density are

potential human pathogens, can be grown *in vitro*, and have been previously associated with AMR

phenotypes. ARGs, as well as mobile genetic elements carrying them, from these species may be

better-studied than those from organisms less likely to pose a threat to human health, including gut

commensals. If ARGs from these organisms are well-represented in databases, it could potentially

bias analyses based on these databases toward detecting pathogen- over commensal-associated

ARGs. However, there has been significant work done on the resistome of the human commensal

microbiome, including functional metagenomics to detect new ARGs. These have found that

commensal anaerobes may serve as significant reservoirs of ARGs, and may in some cases

contribute to the transfer of resistance to pathobionts[143-150]. Commensal carriage of

antimicrobial resistance genes may correspond to the baseline level of 0.3-0.4% ARGs observed

in samples without pathobiont dominance.

Some limitations to the findings of this study must be acknowledged. First, as for all

database-based methodologies, we are limited by accuracy and completeness of those databases.

While the human gut microbiome is fairly well-characterized, there may be so-called microbial

dark matter that is not well-represented in the taxonomic database used for species identification.

We also used a database composed of bacterial and archaeal genomes, excluding consideration of

bacteriophage and microbial eukaryotes from our analyses. As mentioned, database representation

is particularly relevant for our ARG analysis, as the genes in this database may be skewed towards

easily-culturable and pathogenic source species, and our analysis may have missed ARGs found

in commensal or unculturable gut species. Additionally, critics have noted that some genes found

in ARG databases used have unclear links to resistance phenotypes, and may perform regulatory,

efflux, or other functions not always related to antimicrobial resistance[143, 151].

Second, we were limited by the original SPREAD population, in which few subjects

received only a single antimicrobial during the course of the study; this makes it difficult to say

whether the temporal variability we observed was widespread in the cohort, although the fact that

there were frequently high pathobiont levels observed in the larger metagenomics dataset we used

to test our multiple linear regression suggests that this may be the case. Third, in this study we

worked with rectal swabs, which are similar but not identical to fecal samples and may be

susceptible to cross-contamination from urinary pathogens or skin flora, particularly in incontinent

advanced dementia patients[152-155]. Fourth, metagenomic assembly has limitations. It cannot

create bins of all species found in a given sample, genome reconstruction is based on the isolates

present in the database used, and analysis of assembled genomes may exclude consideration of

plasmids – which are often sources of ARGs. Finally, as we analyzed metagenomic data, we cannot

comment on the actual antimicrobial resistance phenotypes of the communities or individual

bacteria that we studied.

CONCLUSIONS:

The gut microbiome was highly variable both between and within subjects, with frequent

blooms and reductions of bacterial species both before and after levofloxacin treatment. We did

not find that levofloxacin had a consistent impact on specific taxa or functions, levels of

antimicrobial resistance genes, or overall microbiome diversity in these subjects. This may suggest

that administration of levofloxacin to this population does not have a detectable impact on their

microbiomes. However, while we could not link levofloxacin to antimicrobial resistance gene

levels, there were a number of samples that had higher relative abundances of these genes. In our

original metagenomics dataset, we were able to identify that levels of these genes could be linked

to blooms of specific bacterial species, including *E. coli*, *P. mirabilis*, and *E. faecalis*. We were

able to build a model to predict total ARG levels in a sample from the relative abundance of these

species, and confirm the validity of this model in a larger metagenomics dataset from the rest of

the SPREAD study, including subjects taking a range of antibiotics. Furthermore, use of

metagenomic assembly and binning allowed us to confirm that our species of interest carry greater

ARG densities than other abundant members of the microbiome, even in subjects where MDROs

were not detected by culturing.

This demonstrates that there is a significant amount of information that can be obtained

from metagenomic assembly and binning. With sufficient depth, powerful computational tools

allow whole genomes to be assembled from short-read metagenomic sequencing, which permits

interrogation of the likely features of species of interest in complex microbial communities. In our

case, we were able to confirm the association between pathobiont blooms and ARG levels in the

gut, showing that the genomes of pathobionts contained a greater proportion of ARGs than gut

commensals such as *Bacteroides* and *Bifidobacterium* species. This suggests that while the

commensal microbiota are known to serve as reservoirs of antimicrobial resistance, in this cohort

blooms of pathobionts may serve as the driver of ARG levels in the gut microbiome. Given how

frequently these blooms occurred, special attention should be paid to these species in dementia

patients in long-term care facilities, a vulnerable group which is often immunocompromised,

frequently administered medication including antimicrobials, and may carry MDRO at relatively

high levels.

METHODS

Sample Collection and Preparation

Subjects:

Eleven subjects were chosen from the SPREAD cohort based on the following inclusion

criteria: at least two consecutive rectal swabs were collected from the subject during the study,

subjects had received a single oral course of levofloxacin during the study (average course of 8

days), and subjects received no other antimicrobials during the study or in the 3 months prior to

the first sample collection. Of the 11 subjects, 10 were female and 10 were white, while ages

ranged from 72 to 101. Five subjects lived through the entire sample collection period, while the

other six passed away at some point prior to the final collection; between this attrition, one sample

that was not collected, and three samples that were not well-sequenced, we had a total of 38 usable

metagenomic samples (Figure 1A; Additional Tables S1-2). All samples were collected under

SPREAD, which was approved by the Institutional Review Board at Hebrew Life[31].

Sample Collection:

Samples were collected by insertion of sterile double-tipped swabs (Starswab II; Starplex

Scientific Inc., Ontario, Canada) into the anus of the subject. The first swab was used to identify

multidrug-resistant organisms (including methicillin-resistant *S. aureus*, vancomycin-resistant

enterococci, and multidrug-resistant Gram-negative organisms such as *E. coli*, *P. mirabilis*, *P.*

aeruginosa, or *P. stuartii*) via culturing techniques as described previously[156]. The second swab

was frozen in 20% glycerol at -80°C for DNA extraction and sequencing.

Sample Processing:

Frozen rectal swabs were thawed and placed into 96-well plates, before extraction using

the PowerSoil DNA Isolation Kit (MOBIO, West Carlsbad, CA) according to the manufacturer's

instructions. DNA concentrations were measured using a Nanodrop 1000 (Thermo Scientific,

Waltham, MA) and extracted DNA was stored at -20°C until further use.

16S rRNA Amplicon Sequencing

Sequencing:

The V4 hypervariable region of the 16S rRNA gene was amplified according to Earth

Microbiome Project protocols. Amplification was performed using Illumina-adapted universal 16S

primers 515F and 806R under the following conditions: 3 minutes at 94°C, 45 cycles of [45

seconds at 94°C, 60 seconds at 50°C, 90 seconds at 72°C], 10 minutes at 72°C. All reactions were

prepared using 5 PRIME polymerase 1X HotMasterMix (5PRIME, Gaithersburg, MD) and run in

triplicate to alleviate primer bias. Triplicates were pooled before cleaning with a PCR Purification

Kit (Qiagen). These products were quantified using the Qubit dsDNA High Sensitivity Assay Kit

(Invitrogen, Eugene, OR) and samples were pooled in equimolar amounts. Sequencing was

performed using the Illumina MiSeq platform located at the New York University Langone

Medical Center Genome Technology Core. Sequences can be found under the BioProject

accession number PRJNA573963 (http://www.ncbi.nlm.nih.gov/bioproject/573963).

Data Processing:

Data processing was performed using the QIIME2 (v 2019.1) pipeline[96]. The Divisive

Amplicon Denoising Algorithm 2 (DADA2) method was used to quality-filter sequences and

categorize amplicon sequence variants (ASVs)[157], and the SILVA (release 132) 99% identity

V4 classifier was used to assign taxonomy to ASVs[158]. See Additional File 1 for more

information. Taxonomic relative abundances were exported at the genus level for further analysis.

Output data can be found in Additional Data 2.

Shotgun Sequencing

Sample Preparation and Sequencing:

Extracted DNA (2 ng DNA in 50 uL buffer) was sheared to 450bp using a Covaris LE220

system. Library preparation was performed using a Biomek FXP Automated Liquid Handling

Workstation (Beckman Coulter) with the KAPA HyperPrep Kit (Roche), with 12 cycles of PCR.

Final libraries were normalized and pooled, with 20 samples per poor. Each pool was run on 2

lanes of an Illumina HiSeq 4000 using the paired-end 2x150bp protocol. Library preparation and

sequencing was performed at the New York University Langone Medical Center Genome

Technology Core. Sequences can be found under the BioProject accession number PRJNA573963

(http://www.ncbi.nlm.nih.gov/bioproject/573963) for the levofloxacin dataset and under the

BioProject accession number PRJNA531921 (https://www.ncbi.nlm.nih.gov/bioproject/531921)

for the test dataset.

Data Processing:

Raw shotgun sequencing reads were processed using Kneaddata (v0.6.1) to remove

contaminating human sequences from the dataset[159]. Briefly, the *kneaddata* function was used

with the Bowtie2 *Homo sapiens* database (v0.1)[160] to remove contaminating host reads from

the sequencing files. See Additional File 1 for more information.

Taxonomic Classification:

Kraken2, a taxonomic classifier that maps shotgun sequencing k-mers to genomic

databases, was used to assign taxonomy to kneaddata-processed shotgun sequencing reads[70].

Briefly, the *kraken2-build* function was used to create a custom database containing the "bacteria"

and "archaea" from NCBI libraries, and the *kraken2* function was used to run the kneaddata-

filtered shotgun sequencing reads against this database and assign taxonomy. While Kraken2 does

not estimate species abundances, Bracken2 (Bayesian Reestimation of Abundance with KrakEN)

uses the taxonomy assigned by Kraken2 to estimate the number of reads per sample that originate

from individual species[71]. The Kraken2 database was used to create a Bracken-compatible

database using the *bracken-build* function, and the Kraken2 report files for each sample were run

against the Bracken database using the *bracken* function for the phylum, genus, and species levels.

Phylum- and species-level relative abundance outputs were formatted for biomarker discovery

using LEfSe. The *kraken-biom* function was used to convert the Bracken report files into a biom

file for import into R. Output data can be found in Additional Data 1. Relative abundance plots

were generated in GraphPad Prism v8.

Taxonomic Diversity Analysis:

Alpha and beta diversity analyses were performed using the phyloseq (v1.27.2)[161, 162]

and vegan (v2.5-4)[163] packages in R (v3.4.3). Briefly, the biom file was imported into a

phyloseq object. The phyloseq *estimate_richness* function was used to obtain Shannon's Diversity

Index values for all samples, while the vegan *phyloseq::distance* and *ordinate* functions were used

to generate a Bray-Curtis matrix and PCoA values. See Additional File 1 for more information.

Data was exported as csv files for formatting, and plotting was performed in GraphPad Prism v8.

Gene and Pathways Analysis:

The Human Microbiome Project Unified Metabolic Analysis Network 2 (HUMAnN2)

pipeline was used to profile the presence and abundance of genetic pathways in our samples[97].

Briefly, the *humann2* function was used with the kneaddata-filtered metagenomic sequences to

estimate genes and MetaCyc pathways present in the samples based on the UniRef90 database,

files were joined using the *humann2_join_tables* function and the full tables were de-leveled using

the *humann2_split_stratified_table* function. The unstratified gene-level abundances were

converted to both GO terms and KEGG orthologs using the *humann2_regroup_table* function, and

the *humann2_renorm_table* function was used to normalize the MetaCyc pathway, GO term, and

KEGG ortholog tables by computing relative abundance. These relative abundance tables were

formatted for biomarker discovery with LEfSe. Additionally, the, and LEfSe was also used to

analyze pre- and post-treated samples using both outputs. See Additional File 1 for more

information. Output data can be found in Additional Data 3. Plots were generated in Graphpad

Prism 8.

<u>Resistome Analysis:</u>

The ARG content of the samples was analyzed using DeepARG-SS, a deep learning model

that can predict ARGs from short-read metagenomic data[98]. We first analyzed the data using the

deeparg function with the *-reads* flag. The mapped ARGs output was then imported into R, where

it was processed to obtain tables of the ARGs detected per sample at both the specific gene and

antibiotic class levels. The ARGs detected were normalized to the number of reads per sample.

Additionally, after metagenomic assembly and binning was performed (see below),

individual bins were analyzed using DeepARG-LS, a deep learning model optimized to predict

ARGS from gene-level input. The *DNA_features* output from selected bins was analyzed using the

deeparg function with the *-genes* flag to analyze whether the levels or identity of ARGs could be

linked to specific species of interest. The ARGs detected were normalized to the number of features

per bin. All output data can be found in Additional Data 4. Plots were generated in GraphPad Prism

8.

<u>Metagenomic Assembly and Binning:</u>

To further examine the ARGs present in the samples, kneaddata-filtered reads were

uploaded to the web-based Pathosystems Resource Integration Center (PATRIC)[164]. Reads

were assembled into contigs using the *auto* option of the Assembly service, which provides both

raw output contigs from specific assembly algorithms and contigs of the "best" assembly as judged

by the in-house PATRIC script ARAST. We ran the assembly using two different inputs: reads

that had been processed by *kneaddata* as pairs, which has the advantage of utilizing mate-pairing

information for longer total reads, and reads that had been processed by *kneaddata* after pairs were

concatenated into a single file, which has the advantage of keeping reads whose mates failed

trimming.

Both the raw SPAdes assembly algorithm contigs[165] and the best assembly contigs were

then processed using the Metagenomics Binning service, which assigns contigs to specific

organisms and annotates the bin's genome. Quality measures were used to define bins as either

"good", "acceptable", or "bad" according to the criteria in Additional Table 3, and only "good" or

"acceptable" bins were used moving forward. When more than one binning strategy (paired

assembly or single assembly, SPAdes contigs or best contigs) called a particular bin as "good" or

"acceptable", quality measures from the four strategies were compared and the highest-quality bin

for a given species of interest was chosen for ARG analysis. Finally, only bins of species present

at 0.1% or greater relative abundance in the corresponding sample were selected for further

analysis. A list of bins used, their source, and quality measures can be found in Additional Table

4.

<u>Taxonomic Biomarker Analysis:</u>

LEfSe was used to identify potential biomarkers distinguishing levofloxacin-treated

samples[95]. In all cases (taxonomic abundances, MetaCyc pathways, KEGG orthologs, GO

terms, ARGs), data was formatted into csv files and uploaded to the Galaxy webserver. LEfSe was

run under default parameters for biomarker detection, comparing either all pre-levofloxacin to all

post-levofloxacin or immediately pre-levofloxacin to immediately post-levofloxacin. LEfSe was

also used to compare genus-level taxonomic abundance outputs from Kraken2/Bracken2 and

QIIME2, again under default parameters.

<u>DECLARATIONS</u>

<u>Ethics Approval and Consent to Participate:</u> Approval for the SPREAD study was obtained from

the Institutional Review Board committee at Hebrew SeniorLife.

<u>Consent for Publication:</u> Not applicable

<u>Competing Interests:</u> The authors declare no competing interests.

<u>Funding:</u> This study was funded by: the National Institute of General Medical Sciences

institutional development award P20GM121344 for the COBRE Center for Antimicrobial

Resistance and Therapeutic Discovery at Brown University (PB), the Centers for Disease Control

and Prevention award 200-2016-91939 (EMCD), the National Institutes of Allergy and Infectious

Diseases award K24AI119158 (EMCD), the National Institute of Aging award R01AG032982

(EMCD), and the Millennium Nucleus for Collaborative Research on Bacterial Resistance

(MICROB-R) supported by the Millennium Scientific Initiative of the Ministry of Economy,

Development and Tourism (Chile) (RA).

<u>Author's Contributions:</u> ADR conceptualized the project, performed analysis, generated figures,

and wrote the manuscript. RA collected the data and contributed to manuscript preparation. EMCD

collected the data and contributed to manuscript preparation. PB conceptualized the project and

wrote the manuscript. All authors read and approved the final manuscript.

<u>Acknowledgements:</u> The authors acknowledge the laboratory of Dr. Martin Blaser and the New

York University Langone Medical Center Genome Technology Core, which prepared

metagenomic libraries and performed all 16S rRNA and shotgun metagenomics sequencing for

this study.

<u>ADDITIONAL DATA FILES</u>

Additional Data 1: Taxonomic Classifications from Shotgun Metagenomics Kraken2/Bracken2

This file includes the relative abundances of the taxonomic classifications at the phylum, genus,

and species level for both the initial levofloxacin-treated dataset (tabs 1, 2, and 3) and the second,

larger test dataset (tabs 4, 5, and 6). Additional Table 5 links the sample names used for the test

dataset in this study with their identifiers in BioProject PRJNA531921.

Additional Data 2: Taxonomic Classifications from 16S rRNA Sequencing QIIME2

This file includes the relative abundances of the taxonomic classifications at the phylum (tab 1)

and genus (tab 2) level for the initial levofloxacin-treated dataset.

Additional Data 3: Metagenomic Classifications from HUMANn2

This file includes the relative abundances of the MetaCyc pathway (tab 1), KEGG ortholog (tab

2), and GO term (tab 3) outputs for the initial levofloxacin-treated dataset.

Additional Data 4: Antimicrobial Resistance Gene profiles from DeepARG

This file includes the relative abundances of antimicrobial class and specific resistance genes at

the phylum, genus, and species level for the initial levofloxacin-treated dataset (tabs 1 and 2), the

second, larger test dataset (tabs 3 and 4), and the bins generated by PATRIC (tabs 5 and 6).

Additional Table 5 links the sample names used for the test dataset in this study with their

identifiers in BioProject PRJNA531921.

Additional File 1: Code Used for Analysis

This file includes the analysis code used for QIIME2, Kraken2 and Bracken, Phyloseq,

HUMANn2, and DeepARG.

List of Abbreviations in Manuscript and Figures

A:A   Aminoglycoside:Aminocoumarin

AG   Aminoglycoside

AMR   Antimicrobial Resistant

ARG   Antimicrobial Resistance Gene

BC   Bacitracin

BL   Beta-Lactam

Bracken2   Bayesian Reestimation of Abundance with KrakEN2

DP   Diaminopyrimidine

FFM   Fosfomycin

FMM   Fosmidomycin

FQ   Fluoroquinolone

GP   Glycopeptide

GO   Gene Ontology

HUMANn2   Human Microbiome Project Unified Metabolic Analysis Network 2

709	KEGG	Kyoto Encyclopedia of Genes and Genomes
710	LEfSe	Linear Discriminant Analysis Effect Size
711	MD	Multidrug
712	MDRO	Multidrug-Resistant Organism
713	MLS	Macrolide-Lincosamide-Streptogramin
714	MP	Mupirocin
715	MRSA	Methicillin-resistant *Staphylococcus aureus*
716	NM	Nitromidazole
717	OZ	Oxazolidinone
718	PATRIC	Pathosystems Resource Integration Center
719	PC	Phenicol
720	PMT	Pleuromutilin
721	PMX	Polymyxin
722	PT	Peptide
723	QIIME2	Quantitative Insights Into Microbial Ecology 2
724	TC	Tetracycline
725	SPREAD	Study of Pathogen Resistance and Exposure to Antimicrobials in Dementia
726	UC	Unclassified
727		

REFERENCES

1. Centers for Disease Control and Prevention: **Antibiotic Resistance Threats in the United States, 2013**. US Department of Health and Human Services. Atlanta, GA, U.S.A.; 2013.

2. Aslam B, Wang W, Arshad MI, Khurshid M, Muzammil S, Rasool MH, Nisar MA, Alvi RF, Aslam MA, Qamar MU *et al*: **Antibiotic resistance: a rundown of a global crisis**. *Infect Drug Resist* 2018, **11**:1645-1658.

3. Michael CA, Dominey-Howes D, Labbate M: **The antimicrobial resistance crisis: causes, consequences, and management**. *Front Public Health* 2014, **2**:145.

4. Ventola CL: **The antibiotic resistance crisis: part 1: causes and threats**. *P T* 2015, **40**(4):277-283.

5. Paterson DL: **The role of antimicrobial management programs in optimizing antibiotic prescribing within hospitals**. *Clin Infect Dis* 2006, **42 Suppl 2**:S90-95.

6. Struelens MJ: **The epidemiology of antimicrobial resistance in hospital acquired infections: problems and possible solutions**. *BMJ* 1998, **317**(7159):652-654.

7. Mulvey MR, Simor AE: **Antimicrobial resistance in hospitals: how concerned should we be?** *CMAJ* 2009, **180**(4):408-415.

8. Cookson B: **Clinical significance of emergence of bacterial antimicrobial resistance in the hospital environment**. *J Appl Microbiol* 2005, **99**(5):989-996.

9. Cassone M, Mody L: **Colonization with Multi-Drug Resistant Organisms in Nursing Homes: Scope, Importance, and Management**. *Curr Geriatr Rep* 2015, **4**(1):87-95.

10. Aliyu S, Smaldone A, Larson E: **Prevalence of multidrug-resistant gram-negative bacteria among nursing home residents: A systematic review and meta-analysis**. *Am J Infect Control* 2017, **45**(5):512-518.

11. O'Fallon E, Gautam S, D'Agata EM: **Colonization with multidrug-resistant gram-negative bacteria: prolonged duration and frequent cocolonization**. *Clin Infect Dis* 2009, **48**(10):1375-1381.

12. O'Fallon E, Kandel R, Schreiber R, D'Agata EM: **Acquisition of multidrug-resistant gram-negative bacteria: incidence and risk factors within a long-term care population**. *Infect Control Hosp Epidemiol* 2010, **31**(11):1148-1153.

13. Pop-Vicas A, Mitchell SL, Kandel R, Schreiber R, D'Agata EM: **Multidrug-resistant gram-negative bacteria in a long-term care facility: prevalence and risk factors**. *J Am Geriatr Soc* 2008, **56**(7):1276-1280.

14. Trick WE, Weinstein RA, DeMarais PL, Kuehnert MJ, Tomaska W, Nathan C, Rice TW, McAllister SK, Carson LA, Jarvis WR: **Colonization of skilled-care facility residents with antimicrobial-resistant pathogens**. *J Am Geriatr Soc* 2001, **49**(3):270-276.

15. van den Dool C, Haenen A, Leenstra T, Wallinga J: **The Role of Nursing Homes in the Spread of Antimicrobial Resistance Over the Healthcare Network**. *Infect Control Hosp Epidemiol* 2016, **37**(7):761-767.

16. Trick WE, Kuehnert MJ, Quirk SB, Arduino MJ, Aguero SM, Carson LA, Hill BC, Banerjee SN, Jarvis WR: **Regional dissemination of vancomycin-resistant enterococci resulting from interfacility transfer of colonized patients**. *J Infect Dis* 1999, **180**(2):391-396.

17. Morrill HJ, Caffrey AR, Jump RL, Dosa D, LaPlante KL: **Antimicrobial Stewardship in Long-Term Care Facilities: A Call to Action**. *J Am Med Dir Assoc* 2016, **17**(2):183 e181-116.

18.   Bonomo RA: **Multiple antibiotic-resistant bacteria in long-term-care facilities: An**

emerging problem in the practice of infectious diseases. *Clin Infect Dis* 2000,

31(6):1414-1422.

19.   Daneman N, Gruneir A, Newman A, Fischer HD, Bronskill SE, Rochon PA, Anderson

GM, Bell CM: **Antibiotic use in long-term care facilities**. *J Antimicrob Chemother* 2011,

66(12):2856-2863.

20.   van Buul L.W. vdSJT, Veenhuizen R.B., Achterberg W.P., Schellevis F.G., Essink

R.T.G.M., van Benthem B.H.B., Natsch S., Hertogh C.M.P.M.: **Antibiotic Use and**

Resistance in Long Term Care Facilities. *Journal of the American Medical Directors*

Association 2012, **12**(6):568.e561-568.e513.

21.   McGowan JE, Jr.: **Antimicrobial resistance in hospital organisms and its relation to**

antibiotic use. *Rev Infect Dis* 1983, **5**(6):1033-1048.

22.   Nicolle LE, Bentley DW, Garibaldi R, Neuhaus EG, Smith PW: **Antimicrobial use in**

long-term-care facilities. SHEA Long-Term-Care Committee. *Infect Control Hosp*

Epidemiol 2000, **21**(8):537-545.

23.   Jones SR, Parker DF, Liebow ES, Kimbrough RC, 3rd, Frear RS: **Appropriateness of**

antibiotic therapy in long-term care facilities. *Am J Med* 1987, **83**(3):499-502.

24.   Dyar OJ, Pagani L, Pulcini C: **Strategies and challenges of antimicrobial stewardship**

in long-term care facilities. *Clin Microbiol Infect* 2015, **21**(1):10-19.

25.   Peron EP, Hirsch AA, Jury LA, Jump RL, Donskey CJ: **Another setting for stewardship:**

high rate of unnecessary antimicrobial use in a veterans affairs long-term care

facility. *J Am Geriatr Soc* 2013, **61**(2):289-290.

26.   Loeb M: **Antibiotic use in long-term-care facilities: many unanswered questions**. *Infect*
Control Hosp Epidemiol 2000, **21**(10):680-683.

27.   D'Agata E, Loeb MB, Mitchell SL: **Challenges in assessing nursing home residents with**
advanced dementia for suspected urinary tract infections. *J Am Geriatr Soc* 2013,
61(1):62-66.

28.   Mitchell SL, Shaffer ML, Loeb MB, Givens JL, Habtemariam D, Kiely DK, D'Agata E:
Infection management and multidrug-resistant organisms in nursing home residents
with advanced dementia. *JAMA Intern Med* 2014, **174**(10):1660-1667.

29.   D'Agata E, Mitchell SL: **Patterns of antimicrobial use among nursing home residents**
with advanced dementia. *Arch Intern Med* 2008, **168**(4):357-362.

30.   Snyder GM, O'Fallon E, D'Agata EM: **Co-colonization with multiple different species**
of multidrug-resistant gram-negative bacteria. *Am J Infect Control* 2011, **39**(6):506-
510.

31.   Mitchell SL, Shaffer ML, Kiely DK, Givens JL, D'Agata E: **The study of pathogen**
resistance and antimicrobial use in dementia: study design and methodology. *Arch*
Gerontol Geriatr 2013, **56**(1):16-22.

32.   D'Agata EM, Habtemariam D, Mitchell S: **Multidrug-Resistant Gram-Negative**
Bacteria: Inter- and Intradissemination Among Nursing Homes of Residents With
Advanced Dementia. *Infect Control Hosp Epidemiol* 2015, **36**(8):930-935.

33.   Human Microbiome Project C: **Structure, function and diversity of the healthy human**
microbiome. *Nature* 2012, **486**(7402):207-214.

34.   Lloyd-Price J, Abu-Ali G, Huttenhower C: **The healthy human microbiome**. *Genome*
Med 2016, **8**(1):51.

35.   Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N,

819   Levenez F, Yamada T *et al*: **A human gut microbial gene catalogue established by**

820   **metagenomic sequencing**. *Nature* 2010, **464**(7285):59-65.

821   36.   O'Toole P.W. CMJ: **Gut microbiota: Changes throughout the lifespan from infancy to**

822   **elderly**. *International Dairy Journal* 2010, **20**(4):281-291.

823   37.   Rowan-Nash AD, Korry BJ, Mylonakis E, Belenky P: **Cross-Domain and Viral**

824   **Interactions in the Microbiome**. *Microbiol Mol Biol Rev* 2019, **83**(1).

825   38.   Nagpal R, Mainali R, Ahmadi S, Wang S, Singh R, Kavanagh K, Kitzman DW,

826   Kushugulova A, Marotta F, Yadav H: **Gut microbiome and aging: Physiological and**

827   **mechanistic insights**. *Nutr Healthy Aging* 2018, **4**(4):267-285.

828   39.   Odamaki T, Kato K, Sugahara H, Hashikura N, Takahashi S, Xiao JZ, Abe F, Osawa R:

829   **Age-related changes in gut microbiota composition from newborn to centenarian: a**

830   **cross-sectional study**. *BMC Microbiol* 2016, **16**:90.

831   40.   Wang F, Yu T, Huang G, Cai D, Liang X, Su H, Zhu Z, Li D, Yang Y, Shen P *et al*: **Gut**

832   **Microbiota Community and Its Assembly Associated with Age and Diet in Chinese**

833   **Centenarians**. *J Microbiol Biotechnol* 2015, **25**(8):1195-1204.

834   41.   Claesson MJ, Cusack S, O'Sullivan O, Greene-Diniz R, de Weerd H, Flannery E, Marchesi

835   JR, Falush D, Dinan T, Fitzgerald G *et al*: **Composition, variability, and temporal**

836   **stability of the intestinal microbiota of the elderly**. *Proc Natl Acad Sci U S A* 2011, **108**

837   **Suppl 1**:4586-4591.

838   42.   Salazar N, Valdes-Varela L, Gonzalez S, Gueimonde M, de Los Reyes-Gavilan CG:

839   **Nutrition and the gut microbiome in the elderly**. *Gut Microbes* 2017, **8**(2):82-97.

840   43.   Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turroni S, Consolandi C,

841   Quercia S, Scurti M, Monti D *et al*: **Gut Microbiota and Extreme Longevity**. *Curr Biol*

842   2016, **26**(11):1480-1485.

843   44.   Alfa MJ, Strang D, Tappia PS, Graham M, Van Domselaar G, Forbes JD, Laminman V,

844   Olson N, DeGagne P, Bray D *et al*: **A randomized trial to determine the impact of a**

845   **digestion resistant starch composition on the gut microbiome in older and mid-age**

846   **adults**. *Clin Nutr* 2018, **37**(3):797-807.

847   45.   Park SH, Kim KA, Ahn YT, Jeong JJ, Huh CS, Kim DH: **Comparative analysis of gut**

848   **microbiota in elderly people of urbanized towns and longevity villages**. *BMC Microbiol*

849   2015, **15**:49.

850   46.   Biagi E, Nylund L, Candela M, Ostan R, Bucci L, Pini E, Nikkila J, Monti D, Satokari R,

851   Franceschi C *et al*: **Through ageing, and beyond: gut microbiota and inflammatory**

852   **status in seniors and centenarians**. *PLoS One* 2010, **5**(5):e10667.

853   47.   Rampelli S, Candela M, Turroni S, Biagi E, Collino S, Franceschi C, O'Toole PW, Brigidi

854   P: **Functional metagenomic profiling of intestinal microbiome in extreme ageing**.

855   *Aging (Albany NY)* 2013, **5**(12):902-912.

856   48.   Santoro A, Ostan R, Candela M, Biagi E, Brigidi P, Capri M, Franceschi C: **Gut**

857   **microbiota changes in the extreme decades of human life: a focus on centenarians**.

858   *Cell Mol Life Sci* 2018, **75**(1):129-148.

859   49.   Claesson MJ, Jeffery IB, Conde S, Power SE, O'Connor EM, Cusack S, Harris HM,

860   Coakley M, Lakshminarayanan B, O'Sullivan O *et al*: **Gut microbiota composition**

861   **correlates with diet and health in the elderly**. *Nature* 2012, **488**(7410):178-184.

50.   Ticinesi A, Milani C, Lauretani F, Nouvenne A, Mancabelli L, Lugli GA, Turroni F,

Duranti S, Mangifesta M, Viappiani A *et al*: **Gut microbiota composition is associated**

with polypharmacy in elderly hospitalized patients. *Sci Rep* 2017, **7**(1):11102.

51.   van Tongeren SP, Slaets JP, Harmsen HJ, Welling GW: **Fecal microbiota composition**

and frailty. *Appl Environ Microbiol* 2005, **71**(10):6438-6442.

52.   Fish DN, Chow AT: **The clinical pharmacokinetics of levofloxacin**. *Clin Pharmacokinet*

1997, **32**(2):101-119.

53.   Chien SC, Rogge MC, Gisclon LG, Curtin C, Wong F, Natarajan J, Williams RR, Fowler

CL, Cheung WK, Chow AT: **Pharmacokinetic profile of levofloxacin following once-**

daily 500-milligram oral or intravenous doses. *Antimicrob Agents Chemother* 1997,

41(10):2256-2260.

54.   Anderson VR, Perry CM: **Levofloxacin : a review of its use as a high-dose, short-course**

treatment for bacterial infection. *Drugs* 2008, **68**(4):535-565.

55.   Inagaki Y, Nakaya R, Chida T, Hashimoto S: **The effect of levofloxacin, an optically-**

active isomer of ofloxacin, on fecal microflora in human volunteers. *Jpn J Antibiot*

1992, **45**(3):241-252.

56.   Edlund C, Nord CE: **Effect of quinolones on intestinal ecology**. *Drugs* 1999, **58 Suppl**

2:65-70.

57.   Edlund C, Nord CE: **Effect on the human normal microflora of oral antibiotics for**

treatment of urinary tract infections. *J Antimicrob Chemother* 2000, **46 Suppl A**:41-48.

58.   Edlund C, Sjostedt S, Nord CE: **Comparative effects of levofloxacin and ofloxacin on**

the normal oral and intestinal microflora. *Scand J Infect Dis* 1997, **29**(4):383-386.

59.   Bhalodi AA, van Engelen TSR, Virk HS, Wiersinga WJ: **Impact of antimicrobial therapy**
on the gut microbiome. *J Antimicrob Chemother* 2019, **74**(Supplement_1):i6-i15.

60.   Ziegler M, Han JH, Landsburg D, Pegues D, Reesey E, Gilmar C, Gorman T, Bink K,
Moore A, Kelly BJ *et al*: **Impact of Levofloxacin for the Prophylaxis of Bloodstream**
Infection on the Gut Microbiome in Patients With Hematologic Malignancy. *Open*
Forum Infect Dis 2019, **6**(7):ofz252.

61.   Sullivan A, Edlund C, Nord CE: **Effect of antimicrobial agents on the ecological balance**
of human microflora. *Lancet Infect Dis* 2001, **1**(2):101-114.

62.   de Lastours V, Chau F, Roy C, Larroque B, Fantin B: **Emergence of quinolone resistance**
in the microbiota of hospitalized patients treated or not with a fluoroquinolone. *J*
Antimicrob Chemother 2014, **69**(12):3393-3400.

63.   Lautenbach E, Strom BL, Nachamkin I, Bilker WB, Marr AM, Larosa LA, Fishman NO:
Longitudinal trends in fluoroquinolone resistance among Enterobacteriaceae isolates
from inpatients and outpatients, 1989-2000: differences in the emergence and
epidemiology of resistance across organisms. *Clin Infect Dis* 2004, **38**(5):655-662.

64.   Dalhoff A: **Global fluoroquinolone resistance epidemiology and implictions for**
clinical use. *Interdiscip Perspect Infect Dis* 2012, **2012**:976273.

65.   Spellberg B, Doi Y: **The Rise of Fluoroquinolone-Resistant Escherichia coli in the**
Community: Scarier Than We Thought. *J Infect Dis* 2015, **212**(12):1853-1855.

66.   Acar JF, Goldstein FW: **Trends in bacterial resistance to fluoroquinolones**. *Clin Infect*
Dis 1997, **24 Suppl 1**:S67-73.

67.   Nordmann P, Poirel L: **Emergence of plasmid-mediated resistance to quinolones in**
Enterobacteriaceae. *J Antimicrob Chemother* 2005, **56**(3):463-469.

68.   Paterson DL: **Resistance in gram-negative bacteria: enterobacteriaceae**. *Am J Med*
908   2006, **119**(6 Suppl 1):S20-28; discussion S62-70.

909   69.   Ruppe E, Woerther PL, Barbier F: **Mechanisms of antimicrobial resistance in Gram-**
910   **negative bacilli**. *Ann Intensive Care* 2015, **5**(1):61.

911   70.   Wood DE, Salzberg SL: **Kraken: ultrafast metagenomic sequence classification using**
912   **exact alignments**. *Genome Biol* 2014, **15**(3):R46.

913   71.   Lu JB, F.P.; Thielen P.; Salzberg, S.L.: **Bracken: estimating species abundance in**
914   **metagenomics data**. *PeerJ Computer Science* 2017:3:e104.

915   72.   Weiner LM, Webb AK, Limbago B, Dudeck MA, Patel J, Kallen AJ, Edwards JR, Sievert
916   DM: **Antimicrobial-Resistant Pathogens Associated With Healthcare-Associated**
917   **Infections: Summary of Data Reported to the National Healthcare Safety Network at**
918   **the Centers for Disease Control and Prevention, 2011-2014**. *Infect Control Hosp*
919   *Epidemiol* 2016, **37**(11):1288-1301.

920   73.   O'Donovan D, Corcoran GD, Lucey B, Sleator RD: **Campylobacter ureolyticus: a**
921   **portrait of the pathogen**. *Virulence* 2014, **5**(4):498-506.

922   74.   Salem N, Salem L, Saber S, Ismail G, Bluth MH: **Corynebacterium urealyticum: a**
923   **comprehensive review of an understated organism**. *Infect Drug Resist* 2015, **8**:129-145.

924   75.   Agudelo Higuita NI, Huycke MM: **Enterococcal Disease, Epidemiology, and**
925   **Implications for Treatment**. In: *Enterococci: From Commensals to Leading Causes of*
926   *Drug Resistant Infection.* Edited by Gilmore MS, Clewell DB, Ike Y, Shankar N. Boston;
927   2014.

928   76.   Fiore E, Van Tyne D, Gilmore MS: **Pathogenicity of Enterococci**. *Microbiol Spectr* 2019,
929   **7**(4).

77.   Conway T, Cohen PS: **Commensal and Pathogenic Escherichia coli Metabolism in the**
Gut. *Microbiol Spectr* 2015, **3**(3).

78.   Woodward SE, Krekhno Z, Finlay BB: **Here, there, and everywhere: How pathogenic**
E. coli sense and respond to gastrointestinal biogeography. *Cell Microbiol*
2019:e13107.

79.   Baqi M, Mazzulli T: **Oligella infections: Case report and review of the literature**. *Can*
J Infect Dis 1996, **7**(6):377-379.

80.   Graham DR, Band JD, Thornsberry C, Hollis DG, Weaver RE: **Infections caused by**
Moraxella, Moraxella urethralis, Moraxella-like groups M-5 and M-6, and Kingella
kingae in the United States, 1953-1980. *Rev Infect Dis* 1990, **12**(3):423-431.

81.   Pugliese A, Pacris B, Schoch PE, Cunha BA: **Oligella urethralis urosepsis**. *Clin Infect*
Dis 1993, **17**(6):1069-1070.

82.   Wilmer AW, P.; Press, N.; Leung, V.; Romney, M.; Champagne, S.: **Oligella urethralis**
as a Cause of Urosepsis. *Clinical Microbiology Newsletter* 2013, **35**(10):84-85.

83.   Schaffer JN, Pearson MM: **Proteus mirabilis and Urinary Tract Infections**. *Microbiol*
Spectr 2015, **3**(5).

84.   Chen CY, Chen YH, Lu PL, Lin WR, Chen TC, Lin CY: **Proteus mirabilis urinary tract**
infection and bacteremia: risk factors, clinical presentation, and outcomes. *J*
Microbiol Immunol Infect 2012, **45**(3):228-236.

85.   Wie SH: **Clinical significance of Providencia bacteremia or bacteriuria**. *Korean J*
Intern Med 2015, **30**(2):167-169.

86. Kurmasheva N, Vorobiev V, Sharipova M, Efremova T, Mardanova A: **The Potential Virulence Factors of Providencia stuartii: Motility, Adherence, and Invasion**. *Biomed Res Int* 2018, **2018**:3589135.

87. Bassetti M, Vena A, Croxatto A, Righi E, Guery B: **How to manage Pseudomonas aeruginosa infections**. *Drugs Context* 2018, **7**:212527.

88. Mittal R, Aggarwal S, Sharma S, Chhibber S, Harjai K: **Urinary tract infections caused by Pseudomonas aeruginosa: a minireview**. *J Infect Public Health* 2009, **2**(3):101-111.

89. Gordon RJ, Lowy FD: **Pathogenesis of methicillin-resistant Staphylococcus aureus infection**. *Clin Infect Dis* 2008, **46 Suppl 5**:S350-359.

90. Lowy FD: **Staphylococcus aureus infections**. *N Engl J Med* 1998, **339**(8):520-532.

91. Naimi TS, LeDell KH, Como-Sabetti K, Borchardt SM, Boxrud DJ, Etienne J, Johnson SK, Vandenesch F, Fridkin S, O'Boyle C *et al*: **Comparison of community- and health care-associated methicillin-resistant Staphylococcus aureus infection**. *JAMA* 2003, **290**(22):2976-2984.

92. Becker K, Heilmann C, Peters G: **Coagulase-negative staphylococci**. *Clin Microbiol Rev* 2014, **27**(4):870-926.

93. Czekaj T, Ciszewski M, Szewczyk EM: **Staphylococcus haemolyticus - an emerging threat in the twilight of the antibiotics age**. *Microbiology* 2015, **161**(11):2061-2068.

94. Froggatt JW, Johnston JL, Galetto DW, Archer GL: **Antimicrobial resistance in nosocomial isolates of Staphylococcus haemolyticus**. *Antimicrob Agents Chemother* 1989, **33**(4):460-466.

95. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C: **Metagenomic biomarker discovery and explanation**. *Genome Biol* 2011, **12**(6):R60.

96.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer

975   N, Pena AG, Goodrich JK, Gordon JI *et al*: **QIIME allows analysis of high-throughput**

976   **community sequencing data**. *Nat Methods* 2010, **7**(5):335-336.

977   97.   Franzosa EA, McIver LJ, Rahnavard G, Thompson LR, Schirmer M, Weingart G, Lipson

978   KS, Knight R, Caporaso JG, Segata N *et al*: **Species-level functional profiling of**

979   **metagenomes and metatranscriptomes**. *Nat Methods* 2018, **15**(11):962-968.

980   98.   Arango-Argoty G, Garner E, Pruden A, Heath LS, Vikesland P, Zhang L: **DeepARG: a**

981   **deep learning approach for predicting antibiotic resistance genes from metagenomic**

982   **data**. *Microbiome* 2018, **6**(1):23.

983   99.   Hiramatsu K: **Molecular evolution of MRSA**. *Microbiol Immunol* 1995, **39**(8):531-543.

984   100.   Hiramatsu K, Cui L, Kuroda M, Ito T: **The emergence and evolution of methicillin-**

985   **resistant Staphylococcus aureus**. *Trends Microbiol* 2001, **9**(10):486-493.

986   101.   Tsubakishita S, Kuwahara-Arai K, Sasaki T, Hiramatsu K: **Origin and molecular**

987   **evolution of the determinant of methicillin resistance in staphylococci**. *Antimicrob*

988   *Agents Chemother* 2010, **54**(10):4352-4359.

989   102.   Enright MC, Robinson DA, Randle G, Feil EJ, Grundmann H, Spratt BG: **The**

990   **evolutionary history of methicillin-resistant Staphylococcus aureus (MRSA)**. *Proc*

991   *Natl Acad Sci U S A* 2002, **99**(11):7687-7692.

992   103.   Smith AM, Klugman KP: **Alterations in PBP 1A essential-for high-level penicillin**

993   **resistance in Streptococcus pneumoniae**. *Antimicrob Agents Chemother* 1998,

994   **42**(6):1329-1333.

995   104.   Zapun A, Contreras-Martel C, Vernet T: **Penicillin-binding proteins and beta-lactam**

996   **resistance**. *FEMS Microbiol Rev* 2008, **32**(2):361-385.

105.   Brannigan JA, Tirodimos IA, Zhang QY, Dowson CG, Spratt BG: **Insertion of an extra**
amino acid is the main cause of the low affinity of penicillin-binding protein 2 in
penicillin-resistant strains of Neisseria gonorrhoeae. *Mol Microbiol* 1990, **4**(6):913-
919.
106.   Dowson CG, Jephcott AE, Gough KR, Spratt BG: **Penicillin-binding protein 2 genes of**
non-beta-lactamase-producing, penicillin-resistant strains of Neisseria gonorrhoeae.
Mol Microbiol 1989, **3**(1):35-41.
107.   Thulin S, Olcen P, Fredlund H, Unemo M: **Total variation in the penA gene of Neisseria**
meningitidis: correlation between susceptibility to beta-lactam antibiotics and penA
gene heterogeneity. *Antimicrob Agents Chemother* 2006, **50**(10):3317-3324.
108.   Beceiro AB, G.: **Class C Beta-Lactamases: An Increasing Problem Worldwide**.
Reviews in Medical Microbiology 2004, **15**(4):141-152.
109.   Garvey MI, Baylay AJ, Wong RL, Piddock LJ: **Overexpression of patA and patB, which**
encode ABC transporters, is associated with fluoroquinolone resistance in clinical
isolates of Streptococcus pneumoniae. *Antimicrob Agents Chemother* 2011, **55**(1):190-
196.
110.   El Garch F, Lismond A, Piddock LJ, Courvalin P, Tulkens PM, Van Bambeke F:
Fluoroquinolones induce the expression of patA and patB, which encode ABC efflux
pumps in Streptococcus pneumoniae. *J Antimicrob Chemother* 2010, **65**(10):2076-2082.
111.   Hernando-Amado S, Blanco P, Alcalde-Rico M, Corona F, Reales-Calderon JA, Sanchez
MB, Martinez JL: **Multidrug efflux pumps as main players in intrinsic and acquired**
resistance to antimicrobials. *Drug Resist Updat* 2016, **28**:13-27.

112.   Sato T, Yokota S, Uchida I, Okubo T, Usui M, Kusumoto M, Akiba M, Fujii N, Tamura

1020   Y: **Fluoroquinolone resistance mechanisms in an Escherichia coli isolate, HUE1,**

1021   **without quinolone resistance-determining region mutations**. *Front Microbiol* 2013,

1022   **4**:125.

1023   113.   Li XZ, Plesiat P, Nikaido H: **The challenge of efflux-mediated antibiotic resistance in**

1024   **Gram-negative bacteria**. *Clin Microbiol Rev* 2015, **28**(2):337-418.

1025   114.   Chow J, Tang H, Mazmanian SK: **Pathobionts of the gastrointestinal microbiota and**

1026   **inflammatory disease**. *Curr Opin Immunol* 2011, **23**(4):473-480.

1027   115.   Dzutsev A, Trinchieri G: **Proteus mirabilis: The Enemy Within**. *Immunity* 2015,

1028   **42**(4):602-604.

1029   116.   Butto LF, Schaubeck M, Haller D: **Mechanisms of Microbe-Host Interaction in Crohn's**

1030   **Disease: Dysbiosis vs. Pathobiont Selection**. *Front Immunol* 2015, **6**:555.

1031   117.   Mirsepasi-Lauridsen HCV, B.A.; Krogfelt, K.A.; Petersen, A.M.: **Escherichia coli**

1032   **pathobionts associated with inflammatory bowel disease**. *Clinical Microbiology*

1033   *Reviews* 2019:32:e00060-00018.

1034   118.   Yang HM-L, H.C.; Struve, C.; Allaire, J.M.; Bosman, E.; Sivignon, A.; Vogl, W.; Ma, C.;

1035   Reid, G.; Li, X.; Petersen, A.M.; Jacobson, K.; Gouin, S.; Barnich, N.; Yu, H.; Krogfelt,

1036   K.A.; Vallance, B.A.: **Ulcerative Colitis-associated E. coli Pathobionts Potentiate**

1037   **Colitis in Susceptible Hosts**. *The Journal of Immunology* 2019, **202**(1):192.193.

1038   119.   Hamilton AL, Kamm MA, Ng SC, Morrison M: **Proteus spp. as Putative**

1039   **Gastrointestinal Pathogens**. *Clin Microbiol Rev* 2018, **31**(3).

120. Archambaud C, Derre-Bobillot A, Lapaque N, Rigottier-Gois L, Serror P: **Intestinal**

translocation of enterococci requires a threshold level of enterococcal overgrowth in

the lumen. *Sci Rep* 2019, **9**(1):8926.

121. Overturf GD, Wilkins J, Ressler R: **Emergence of resistance of Providencia stuartii to**

multiple antibiotics: speciation and biochemical characterization of Providencia. *J*

Infect Dis 1974, **129**(3):353-357.

122. McHale PJ, Keane CT, Dougan G: **Antibiotic resistance in Providencia stuartii isolated**

in hospitals. *J Clin Microbiol* 1981, **13**(6):1099-1104.

123. Hawkey PM: **Providencia stuartii: a review of a multiply antibiotic-resistant**

bacterium. *J Antimicrob Chemother* 1984, **13**(3):209-226.

124. Warren JW: **Providencia stuartii: a common cause of antibiotic-resistant bacteriuria**

in patients with long-term indwelling catheters. *Rev Infect Dis* 1986, **8**(1):61-67.

125. Oikonomou O, Liakopoulos A, Phee LM, Betts J, Mevius D, Wareham DW: **Providencia**

stuartii Isolates from Greece: Co-Carriage of Cephalosporin (blaSHV-5, blaVEB-1),

Carbapenem (blaVIM-1), and Aminoglycoside (rmtB) Resistance Determinants by a

Multidrug-Resistant Outbreak Clone. *Microb Drug Resist* 2016, **22**(5):379-386.

126. Barros EM, Ceotto H, Bastos MC, Dos Santos KR, Giambiagi-Demarval M:

Staphylococcus haemolyticus as an important hospital pathogen and carrier of

methicillin resistance genes. *J Clin Microbiol* 2012, **50**(1):166-168.

127. Maleki A, Ghafourian S, Taherikalani M, Soroush S: **Alarming and Threatening Signals**

from Health Centers About Multi Drug Resistance Staphylococcus haemolyticus.

Infect Disord Drug Targets 2019, **19**(2):118-127.

128.   Costa SS, Viveiros M, Amaral L, Couto I: **Multidrug Efflux Pumps in Staphylococcus**

1063   **aureus: an Update**. *Open Microbiol J* 2013, **7**:59-71.

1064   129.   Truong-Bolduc QC, Dunman PM, Strahilevitz J, Projan SJ, Hooper DC: **MgrA is a**

1065   **multiple regulator of two new efflux pumps in Staphylococcus aureus**. *J Bacteriol*

1066   2005, **187**(7):2395-2405.

1067   130.   Lina G, Quaglia A, Reverdy ME, Leclercq R, Vandenesch F, Etienne J: **Distribution of**

1068   **genes encoding resistance to macrolides, lincosamides, and streptogramins among**

1069   **staphylococci**. *Antimicrob Agents Chemother* 1999, **43**(5):1062-1066.

1070   131.   Ross JI, Eady EA, Cove JH, Cunliffe WJ, Baumberg S, Wootton JC: **Inducible**

1071   **erythromycin resistance in staphylococci is encoded by a member of the ATP-binding**

1072   **transport super-gene family**. *Mol Microbiol* 1990, **4**(7):1207-1214.

1073   132.   Singh L, Cariappa MP, Kaur M: **Klebsiella oxytoca: An emerging pathogen?** *Med J*

1074   *Armed Forces India* 2016, **72**(Suppl 1):S59-S61.

1075   133.   Moradigaravand D, Martin V, Peacock SJ, Parkhill J: **Population structure of multidrug**

1076   **resistant Klebsiella oxytoca within hospitals across the UK and Ireland identifies**

1077   **sharing of virulence and resistance genes with K. pneumoniae**. *Genome Biol Evol* 2017.

1078   134.   Liu H, Zhu J, Hu Q, Rao X: **Morganella morganii, a non-negligent opportunistic**

1079   **pathogen**. *Int J Infect Dis* 2016, **50**:10-17.

1080   135.   Noel GJ: **A Review of Levofloxacin for the Treatment of Bacterial Infections**. *Clinical*

1081   *Medicine Insights: Therapeutics* 2009, **1**.

1082   136.   Bush LM, Chaparro-Rojas F, Okeh V, Etienne J: **Cumulative clinical experience from**

1083   **over a decade of use of levofloxacin in urinary tract infections: critical appraisal and**

1084   **role in therapy**. *Infect Drug Resist* 2011, **4**:177-189.

137. Croom KF, Goa KL: **Levofloxacin: a review of its use in the treatment of bacterial**

infections in the United States. *Drugs* 2003, **63**(24):2769-2802.

138. Deshpande A, Pant C, Jain A, Fraser TG, Rolston DD: **Do fluoroquinolones predispose**

patients to Clostridium difficile associated disease? A review of the evidence. *Curr*

Med Res Opin 2008, **24**(2):329-333.

139. Mehta RS, Abu-Ali GS, Drew DA, Lloyd-Price J, Subramanian A, Lochhead P, Joshi AD,

Ivey KL, Khalili H, Brown GT *et al*: **Stability of the human faecal microbiome in a**

cohort of adult men. *Nat Microbiol* 2018, **3**(3):347-355.

140. Liskiewicz P, Pelka-Wysiecka J, Kaczmarczyk M, Loniewski I, Wronski M, Baba-Kubis

A, Skonieczna-Zydecka K, Marlicz W, Misiak B, Samochowiec J: **Fecal Microbiota**

Analysis in Patients Going through a Depressive Episode during Treatment in a

Psychiatric Hospital Setting. *J Clin Med* 2019, **8**(2).

141. Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, Sogin ML,

Jones WJ, Roe BA, Affourtit JP *et al*: **A core gut microbiome in obese and lean twins**.

Nature 2009, **457**(7228):480-484.

142. Davies EV, James CE, Williams D, O'Brien S, Fothergill JL, Haldenby S, Paterson S,

Winstanley C, Brockhurst MA: **Temperate phages both mediate and drive adaptive**

evolution in pathogen biofilms. *Proc Natl Acad Sci U S A* 2016, **113**(29):8266-8271.

143. van Schaik W: **The human gut resistome**. *Philos Trans R Soc Lond B Biol Sci* 2015,

370(1670):20140087.

144. Francino MP: **Antibiotics and the Human Gut Microbiome: Dysbioses and**

Accumulation of Resistances. *Front Microbiol* 2015, **6**:1543.

145.   Penders J, Stobberingh EE, Savelkoul PH, Wolffs PF: **The human microbiome as a**

1108   **reservoir of antimicrobial resistance**. *Front Microbiol* 2013, **4**:87.

1109   146.   Hu Y, Yang X, Qin J, Lu N, Cheng G, Wu N, Pan Y, Li J, Zhu L, Wang X *et al*:

1110   **Metagenome-wide analysis of antibiotic resistance genes in a large cohort of human**

1111   **gut microbiota**. *Nat Commun* 2013, **4**:2151.

1112   147.   Jackson RW, Vinatzer B, Arnold DL, Dorus S, Murillo J: **The influence of the accessory**

1113   **genome on bacterial pathogen evolution**. *Mob Genet Elements* 2011, **1**(1):55-65.

1114   148.   Scott KP: **The role of conjugative transposons in spreading antibiotic resistance**

1115   **between bacteria that inhabit the gastrointestinal tract**. *Cell Mol Life Sci* 2002,

1116   **59**(12):2071-2082.

1117   149.   Salyers AA, Gupta A, Wang Y: **Human intestinal bacteria as reservoirs for antibiotic**

1118   **resistance genes**. *Trends Microbiol* 2004, **12**(9):412-416.

1119   150.   Kazimierczak KA, Scott KP: **Antibiotics and resistance genes: influencing the**

1120   **microbial ecosystem in the gut**. *Adv Appl Microbiol* 2007, **62**:269-292.

1121   151.   Martinez JL, Coque TM, Baquero F: **What is a resistance gene? Ranking risk in**

1122   **resistomes**. *Nat Rev Microbiol* 2015, **13**(2):116-123.

1123   152.   Budding AE, Grasman ME, Eck A, Bogaards JA, Vandenbroucke-Grauls CM, van

1124   Bodegraven AA, Savelkoul PH: **Rectal swabs for analysis of the intestinal microbiota**.

1125   *PLoS One* 2014, **9**(7):e101344.

1126   153.   Bassis CM, Moore NM, Lolans K, Seekatz AM, Weinstein RA, Young VB, Hayden MK,

1127   Program CDCPE: **Comparison of stool versus rectal swab samples and storage**

1128   **conditions on bacterial community profiles**. *BMC Microbiol* 2017, **17**(1):78.

154. Biehl LM, Garzetti D, Farowski F, Ring D, Koeppel MB, Rohde H, Schafhausen P, Stecher

B, Vehreschild M: **Usability of rectal swabs for microbiome sampling in a cohort study**

of hematological and oncological patients. *PLoS One* 2019, **14**(4):e0215428.

155. Fair K, Dunlap DG, Fitch A, Bogdanovich T, Methe B, Morris A, McVerry BJ, Kitsios

GD: **Rectal Swabs from Critically Ill Patients Provide Discordant Representations of**

the Gut Microbiome Compared to Stool Samples. *mSphere* 2019, **4**(4).

156. Snyder GM, D'Agata EM: **Diagnostic accuracy of surveillance cultures to detect**

gastrointestinal colonization with multidrug-resistant gram-negative bacteria. *Am J*

Infect Control 2012, **40**(5):474-476.

157. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP: **DADA2: High-**

resolution sample inference from Illumina amplicon data. *Nat Methods* 2016,

13(7):581-583.

158. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO: **The**

SILVA ribosomal RNA gene database project: improved data processing and web-

based tools. *Nucleic Acids Res* 2013, **41**(Database issue):D590-596.

159. McIver LJ, Abu-Ali G, Franzosa EA, Schwager R, Morgan XC, Waldron L, Segata N,

Huttenhower C: **bioBakery: a meta'omic analysis environment**. *Bioinformatics* 2018,

34(7):1235-1237.

160. Langmead B, Trapnell C, Pop M, Salzberg SL: **Ultrafast and memory-efficient**

alignment of short DNA sequences to the human genome. *Genome Biol* 2009,

10(3):R25.

161. McMurdie PJ, Holmes S: **phyloseq: an R package for reproducible interactive analysis**

and graphics of microbiome census data. *PLoS One* 2013, **8**(4):e61217.

162.   McMurdie PJ, Holmes S: **Phyloseq: a bioconductor package for handling and analysis**

1153   **of high-throughput phylogenetic sequence data**. *Pac Symp Biocomput* 2012:235-246.

1154   163.   Dixon P: **VEGAN, a package of R functions for community ecology**. *J Veg Sci* 2009,

1155   **14**:927-930.

1156   164.   Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz

1157   T, Gabbard JL *et al*: **Improvements to PATRIC, the all-bacterial Bioinformatics**

1158   **Database and Analysis Resource Center**. *Nucleic Acids Res* 2017, **45**(D1):D535-D542.

1159   165.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA: **metaSPAdes: a new versatile**

1160   **metagenomic assembler**. *Genome Res* 2017, **27**(5):824-834.

1161

Figure 1: Subject Overview and Diversity Metrics

(A) Metagenomics sequencing was performed on longitudinal samples from eleven subjects from SPREAD who had received a single course of levofloxacin during their participation in the study. Points represent collection of samples, at intervals of approximately 3 months, relative to administration of levofloxacin. (B) Shannon diversity over time of all subjects. The dashed line indicates administration of levofloxacin. p = 0.175 for immediately pre-levofloxacin vs. immediately post-levofloxacin samples and p = 0.1006 for all pre-levofloxacin vs. all post-levofloxacin samples; Mann-Whitney test. (C) PCoA analysis of Bray-Curtis Dissimilarity. Solid arrows connect immediately pre- with immediately post-levofloxacin samples, dashed arrows connect other sequential samples, and dotted arrows connect samples where an intermediate sample is missing. (D) Within-subjects Bray-Curtis Dissimilarity of sequential samples. p = 0.6248 between pre-levofloxacin samples, post-levofloxacin samples, or pre-post levofloxacin samples; ANOVA). (E) Overall within-subjects, T0 between-subjects, and overall between-subjects Bray-Curtis Dissimilarity. p = 0.0262 for overall within-subjects vs. T0 between-subjects and p = 0.0175 for overall within-subjects vs. overall between-subjects; t-test with Welch's correction.

Figure 2: Relative Abundances of Species and Antimicrobial Resistance Genes
(A) Relative abundance of the most-abundant species across all samples, with all other species grouped in the "other" category. Species are grouped by genus and phylum, and are ranked within those levels by average relative abundance across all samples. Broad color categories distinguish phylum (*Proteobacteria* are red, *Bacteroidetes* are blue, *Firmicutes* are green, and *Actinobacteria* are purple), while different species of the same genus are given the same specific background color. Red lines indicate levofloxacin administration; dashed lines indicate usage between consecutive timepoints, while dotted lines indicate usage where the immediately post-levofloxacin sample is missing (B) Relative abundance of the most-abundant antimicrobial resistance genes (ARGs) across all samples. Specific ARGs are grouped by the class of antimicrobials they provide resistance to. Broad color categories distinguish class (Multidrug RGs are blue, MLS RGs are red, etc.), while related gene categories (ex: the *mec* operon or *mex* efflux proteins) are given the same specific background color. All ARGs were normalized to the total number of reads.

Figure 3: Antimicrobial Resistance Gene Profiles Reflect Taxonomic Observations
(A) Relative abundance of species in Subject A, showing a bloom in *S. aureus* at T12. (B) Relative abundance of ARG classes in Subject A, showing an expansion in beta-lactam resistance genes at T12. (C) Relative abundance of beta-lactam resistance genes in Subject A, showing increases in the *mecA/mecI/mecRI* operon at T12. (D) Relative abundance of species in Subject B, showing a bloom in *E. coli* at T3. (E) Relative abundance of ARG classes in Subject B, showing an expansion in multidrug, beta-lactam, and fluoroquinolone resistance genes at T3. (F) Relative abundance of multidrug resistance genes in Subject B, showing increases in various ARGs at T3. (G) Relative abundance of fluoroquinolone resistance genes in Subject B, showing increases in genes including *patA* and *mdtK* at T3. (H) Relative abundance of beta-lactam resistance genes in Subject B, showing increases in genes including penicillin-binding proteins and class C beta-lactamase at T3. (I) Relative abundance of species in Subject H, showing a bloom in *S. haemolyticus* at T6. (J) Relative abundance of ARG classes in Subject H, showing increases in MLS and fluoroquinolone resistance genes. (K) Relative abundance of MLS resistance genes in Subject H, showing an increase in staphylococcal resistance gene *msrA* and others at T6. (L) Relative abundance of fluoroquinolone resistance genes in Subject H, showing an increase in staphylococcal resistance gene *norB* and others at T6.

Figure 4: Relationship of ARG Levels to the Relative Abundance of Specific Pathobionts
(A) Correspondence between the relative abundances of key species of interest (*E. coli*, *P. mirabilis*, *E. faecalis*, *P. stuartii*, and *S. haemolyticus*) and total ARG density in each sample. (B) Correlation between the sum of the relative abundances of *E. coli*, *P. mirabilis*, and *E. faecalis* and the total ARG density in each sample (r = 0.791, R^2 = 0.6254, p <0.0001; Pearson's correlation). (C) Correlation between the sum of the relative abundances of *E. coli*, *P. mirabilis*, *E. faecalis*, *P. stuartii*, and *S. haemolyticus* and the total ARG density in each sample (r = 0.933, R^2 = 0.8706, p < 0.0001; Pearson's correlation). (D) Average ARG density in bins of species across all samples in which we were able to construct a bin for that species. Specific genes are grouped and colored by their ARG class, and bins are grouped by phylum and ranked by their total average ARG density within that phylum.

Figure 5: Prediction of ARG Density From Relative Abundance of Specific Pathobionts
(A) Correlation between the sum of the relative abundances of *E. coli*, *P. mirabilis*, and *E. faecalis* and the total ARG density in each sample in the test dataset (r = 0.7139, r^2 = 0.5096, p<0.0001; Pearson's correlation). (B) Correlation between the sum of the relative abundances of *E. coli*, *P. mirabilis*, *E. faecalis*, *P. stuartii*, and *S. haemolyticus* and the total ARG density in each sample in the test dataset (r = 0.7753, r^2 = 0.6012, p<0.0001; Pearson's correlation). (C) Multiple linear regression of relative abundances of *E. coli*, *P. mirabilis*, and *E. faecalis* to ARG density in samples in the levofloxacin dataset (38 samples). (D) Correlation between the predicted ARG density and actual ARG density in the test dataset (67 samples) based on the relative abundances of *E. coli*, *P. mirabilis*, and *E. faecalis*. (E) Multiple linear regression of relative abundances of *E. coli*, *P. mirabilis*, *E. faecalis*, *P. stuartii*, and *S. haemolyticus* to ARG density in samples in the levofloxacin dataset (38 samples). (F) Correlation between the predicted ARG density and actual ARG density in the test dataset (67 samples) based on the relative abundances of *E. coli*, *P. mirabilis*, *E. faecalis*, *P. stuartii*, and *S. haemolyticus*.