Evaluating the informativeness of deep learning annotations for human complex diseases

Kushal K. Dey[1], Bryce Van de Geijn [1], Samuel Sungil Kim [1,3], Farhad Hormozdiari[1], David R. Kelley[4] & Alkes L. Price [1,2,5]

1 Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, MA
2 Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA
3 Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Cambridge, MA
4 Calico Labs. South San Francisco, CA
5 Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA

Abstract

Deep learning models have shown great promise in predicting genome-wide regulatory effects from DNA sequence, but their informativeness for human complex diseases and traits is not fully understood. Here, we evaluate the disease informativeness of two types of deep learning annotations: (1) variant-level annotations (based on the reference allele), assessing whether they are more informative for complex disease than the underlying experimental data used to train the predictive models; and (2) allelic-effect annotations (absolute value of the predicted difference between reference and variant alleles), which have been a major focus of recent work. In each case, we primarily consider annotations constructed using two previously trained deep learning models, DeepSEA and Basenji. We apply stratified LD score regression (S-LDSC) to 41 independent diseases and complex traits (average N=320K) to evaluate each annotation's informativeness for disease heritability conditional on a broad set of coding, conserved, regulatory and LD-related annotations from the baseline-LD model and other sources; as a secondary metric, we also evaluate the accuracy of models that incorporate deep learning annotations in predicting disease-associated or fine-mapped SNPs. We aggregated annotations across all tissues (resp. blood cell types or brain tissues) in meta-analyses across all 41 traits (resp. 11 blood-related traits or 8 brain-related traits). Variant-level annotations, despite being highly enriched for disease heritability, produced no conditionally significant results in meta-analyses across all 41 traits or 11 blood-related traits, but brain-specific DeepSEA-H3K4me3 and Basenji-H3K27ac annotations were conditionally significant in meta-analyses across 8 brain-related traits; a sequence motif analysis suggests that these annotations could be capturing unique information about nucleosome occupancy. Allelic-effect annotations were also highly enriched for disease heritability, and produced conditionally significant results for Basenji-H3K4me3 in meta-analyses across all 41 traits and brain-specific Basenji-H3K4me3 in meta-analyses across 8 brain-related traits. We conclude that deep learning models are informative for disease, but their informativeness cannot be inferred from metrics based on their accuracy in predicting regulatory annotations.

Introduction

Disease risk variants identified by genome-wide association studies (GWAS) lie predominantly in non-coding regions of the genome[1,2,3,4,5,6,7], motivating broad efforts to generate genome-wide maps of regulatory marks across tissues and cell types[8,9,10,11]. Recently, deep learning models trained using these genome-wide maps have shown considerable promise in predicting regulatory marks directly from DNA sequence[12,13,14,15,16,17,18]. In particular, these studies showed that variant-level deep learning annotations (based on the reference allele) attained high accuracy in predicting the underlying chromatin marks[13,14,15,16], and that models incorporating allelic-effect deep learning annotations (absolute value of the predicted difference between reference and variant alleles) attained high accuracy in predicting disease-associated SNPs[13,14,15,16]; additional applications of deep learning models, including analyses of signed allelic-effect annotations, are discussed in the Discussion section. However, it is unclear whether deep learning annotations at commonly varying SNPs contain unique information for complex disease that is not present in other annotations.

Here, we evaluate the informativeness for complex disease of both variant-level and allelic-effect annotations at commonly varying SNPs constructed using two deep learning models previously trained on tissue-specific regulatory features (DeepSEA[13,15] and Basenji[16]); we also train a new deep learning model to prioritize SNPs within non-tissue-specific features (BiClassCNN). We apply stratified LD score regression[5,19] (S-LDSC) to 41 independent diseases and complex traits (average N=320K) to evaluate each annotation's informativeness for disease heritability conditional on a broad set of coding, conserved, regulatory and LD-related annotations from the baseline-LD model[19] and other sources (imputed Roadmap and ChromHMM annotations[11,20,21,22]); as a secondary metric, we also evaluate the accuracy of models that incorporate deep learning annotations in predicting disease-associated or fine-mapped SNPs[23,24]. We aggregate DeepSEA and Basenji annotations across all tissues in meta-analyses across all 41 traits, across blood cell types in meta-analyses across 11 blood-related traits, and across brain tissues in meta-analyses across 8 brain-related traits. Our analyses of variant-level annotations assess the specific value of deep learning models in denoising chromatin data (ref.[25]; see Discussion), analogous to methods such as ChromImpute[20] and ChromHMM[21,22] that do not make use of DNA sequence; we thus condition our variant-level analysis on ChromImpute and ChromHMM annotations. Our analyses of allelic-effect annotations further condition on informative variant-level annotations, specifically assessing whether predicted allelic effects contain additional information about disease beyond the information in variant-level annotations.

Results

Overview of methods

We define a genomic annotation as an assignment of a numeric value to each SNP; annotations can be either binary or continuous-valued (Methods). Our focus is on continuous-valued annotations (with values between 0 and 1) trained by deep learning models to predict biological function from DNA sequence. Annotation values are defined for each SNP with minor allele count ≥ 5 in a 1000 Genomes Project European reference panel[26], as in our previous work[5]. We have publicly released all new annotations analyzed

in this study (see URLs).

In our analysis of variant-level deep learning annotations across 41 traits, we analyzed 25 non-tissue-specific deep learning annotations: 8 DeepSEA annotations[13,15] previously trained to predict 4 tissue-specific chromatin marks (DNase, H3K27ac, H3K4me1, H3K4me3) known to be associated with active promoter and enhancer regions across 127 Roadmap tissues[11,27], aggregated using the average (Avg) or maximum (Max) across tissues; 8 analogous Basenji annotations[16], quantile-matched with DeepSEA annotations to lie between 0 and 1; and 9 BiClassCNN annotations that we trained to prioritize SNPs within non-tissue-specific features (Table 1 and Methods). To assess whether the deep learning annotations provided unique information for disease, we conservatively included a broad set of coding, conserved, regulatory and LD-related annotations in our analyses: 86 annotations from the baseline-LD (v2.1) model[19], which has been shown to effectively model LD-dependent architectures[28]; 8 Roadmap annotations[11] (analogous to DeepSEA and Basenji annotations), imputed using ChromImpute[20]; and 40 ChromHMM annotations[21,22] based on 20 ChromHMM states across 127 Roadmap tissues[11] (Table S1). In our analysis of allelic-effect (Δ) annotations, we analyzed 16 non-tissue-specific deep learning annotations (8 DeepSEAΔ and 8 BasenjiΔ), again aggregated using the average (Avg) or maximum (Max) across tissues (Table 1 and Methods). When comparing pairs of annotations that differed only in their aggregation strategy (Avg/Max), chromatin mark (DNase/H3K27ac/H3K4me1/H3K4me3), model (DeepSEA/Basenji) or type (variant-level/allelic-effect), respectively, we observed large correlations across aggregation strategies (average $r = 0.71$), chromatin marks (average $r = 0.58$), models (average $r = 0.54$) and types (average $r = 0.48$) (Figure S1).

In our analysis of 11 blood-related traits (resp. 8 brain-related traits), for both variant-level and allelic-effect annotations, we analyzed 8 DeepSEA annotations and 8 Basenji annotations that were aggregated across 27 blood cell types (resp. 13 brain tissues), instead of all 127 tissues. Details of other annotations included in these analyses are provided below (see Variant-level annotations for blood and brain traits subsection).

We assessed the informativeness of these annotations for disease heritability using stratified LD score regression (S-LDSC) with the baseline-LD model[5,19]. We considered two metrics, enrichment and standardized effect size ($\tau^\star$). Enrichment is defined as the proportion of heritability explained by SNPs in an annotation divided by the proportion of SNPs in the annotation[5], and generalizes to continuous-valued annotations with values between 0 and 1[29]. Standardized effect size ($\tau^\star$) is defined as the proportionate change in per-SNP heritability associated with a 1 standard deviation increase in the value of the annotation, conditional on other annotations included in the model[19]; unlike enrichment, $\tau^\star$ quantifies effects that are unique to the focal annotation. In our "marginal" analyses, we estimated $\tau^\star$ for each focal annotation conditional on annotations from the baseline-LD model. In our "joint" analyses, we merged baseline-LD model annotations with focal annotations that were marginally significant after Bonferroni correction and performed forward stepwise elimination to iteratively remove focal annotations that had conditionally non-significant $\tau^\star$ values after Bonferroni correction, as in ref.[19]. All analyses of allelic-effect annotations were further conditioned on jointly significant annotations from the variant-level analysis. Distinct from evaluating deep learning annotations using S-LDSC, we also evaluated the accuracy of models that incorporate deep learning annotations in predicting disease-associated or fine-mapped SNPs[23,24] (Methods).

Variant-level annotations for 41 diseases and traits

We evaluated the informativeness of 8 DeepSEA, 8 Basenji and 9 BiClassCNN annotations for disease heritability by applying S-LDSC with the baseline-LD model[5,19] to summary association statistics for 41 independent diseases and complex traits (average N=320K); for 6 traits we analyzed two different data sets, leading to a total of 47 data sets analyzed (Table S2). We meta-analyzed results across these 47 data sets, which were chosen to be independent[29]. The 41 traits include 27 UK Biobank traits[30] for which summary association statistics are publicly available (see URLs).

We constructed 8 variant-level DeepSEA predictive annotations by applying previously trained DeepSEA models[15] (see URLs) for each of 4 tissue-specific chromatin marks (DNase, H3K27ac, H3K4me1, H3K4me3) across 127 Roadmap tissues[11] to 1kb of human reference sequence around each SNP; for each chromatin mark, we aggregated variant-level DeepSEA annotations across the 127 tissues using either the average (Avg) or maximum (Max) across tissues (Table 1 and Methods). The DeepSEA model was highly predictive of the corresponding tissue-specific chromatin marks, with AUROC values reported by ref.[15] ranging from $0.77 - 0.97$ (Table S3). In our marginal analysis of disease heritability using S-LDSC, all DeepSEA annotations were highly enriched for disease heritability (average enrichment 2.2x), but only one DeepSEA annotation (DeepSEA-H3K4me3-Max) attained a Bonferroni-significant standardized effect size ($\tau^\star$) conditional on the baseline-LD model (Figure 1 and Table S4). This implies that high AUROC values for variant-level annotations do not necessarily translate into conditional informativeness for human disease.

We constructed 8 variant-level Basenji annotations by applying previously trained Basenji models[16] (see URLs) and aggregating across tissues in analogous fashion (Table 1 and Methods); Basenji uses a Poisson predictive deep learning model framework, unlike the binary classification approach of DeepSEA, and analyzes 130kb of human reference sequence around each SNP using dilated convolutional layers. The constituent tissue-specific Basenji annotations do not lie between 0 and 1, and ref.[16] did not report AUROC values; we transformed these annotations to lie between 0 and 1 via quantile matching with corresponding DeepSEA annotations, to ensure a fair comparison of the two approaches (Methods). In our marginal analysis of disease heritability using S-LDSC, all Basenji annotations were highly enriched for disease heritability (average enrichment 2.2x), but no Basenji annotation attained a Bonferroni-significant $\tau^\star$ conditional on the baseline-LD model (Table S4).

We constructed 9 BiClassCNN annotations (Table 1 and Methods); unlike variant-level DeepSEA and Basenji annotations, the BiClassCNN model was only applied to non-tissue-specific annotations (e.g. coding, promoter, or TSS), and was used to prioritize SNPs within these annotations by restricting BiClassCNN predictions to the underlying annotations; prior to this restriction, the BiClassCNN model was highly predictive of the underlying annotations, with AUROC values ranging from $0.67 - 0.84$ (Table S3). In our marginal analysis of disease heritability using S-LDSC, all BiClassCNN annotations were highly enriched for disease heritability (average enrichment 2.9x) except for BiClassCNN-Repressed (which was depleted, 0.68x), and 4 BiClassCNN annotations attained a Bonferroni-significant $\tau^\star$ conditional on the baseline-LD model (Figure 1 and Table S5). 3 of these 4 annotations (BiClassCNN-Coding, BiClassCNN-Repressed, and BiClassCNN-TSS) produced independent signals (Table S6).

To ensure a conservative assessment of which annotations provide unique information for disease, we included a broader set of annotations in our analyses. First, we included 8

Roadmap annotations[11] (analogous to DeepSEA and Basenji annotations) imputed using ChromImpute[20], and 40 ChromHMM annotations[21,22] based on 20 ChromHMM states across 127 Roadmap tissues[11], again aggregated using the average (Avg) or maximum (Max) across tissues (Table S1 and Methods). Of these, 4 Roadmap annotations and 17 ChromHMM annotations attained a Bonferroni-significant $\tau^\star$ conditional on the baseline-LD model (Figure S2, Figure S3 and Table S7). Second, motivated by the BiClassCNN results (see above), we investigated additional annotations related to coding, repressed and TSS regions. We determined that BiClassCNN-TSS was highly correlated with CpG-island, local CpG-content ($\pm$1kb) and local GC-content ($\pm$1kb) annotations (Figure S4). We thus incorporated CpG-island, local CpG-content and local GC-content annotations, as well as these annotations restricted to coding, repressed and TSS regions. We also considered a gene-level annotation ($\pm$5kb) based on probability of loss-of-function intolerance (pLI)[31], the pLI annotation restricted to coding and TSS regions, and PolyPhen[32,33] and SIFT[34] coding annotations ($12 + 5 = 17$ additional annotations; Table S8 and Methods). 11 of these 17 annotations attained a Bonferroni-significant $\tau^\star$ conditional on the baseline-LD model (Table S8).

We jointly analyzed all 38 annotations that were Bonferroni-significant in our marginal analyses (Figure 1, Figure S2, Figure S3, Table S4, Table S5, Table S7, Table S8) by performing forward stepwise elimination to iteratively remove annotations that had conditionally non-significant τ^* values after Bonferroni correction (based on a total of 106 (variant-level + allelic-effect) non-tissue-specific annotations tested in marginal analyses). Of these, 0 DeepSEA, 0 BiClassCNN, 1 Roadmap, 1 ChromHMM, 3 local GC-content and 2 pLI annotations were jointly significant in the resulting joint model (Figure 1, Figure S5 and Table S9); these annotations were included in all of the conditional analyses below. We determined that the surviving Roadmap-H3K4me1-Avg annotation was sufficient to eliminate the DeepSEA-H3K4me3-Max annotation that was significant in our marginal analysis (Table S10), and that the local GC-content annotations were sufficient to eliminate the 4 BiClassCNN annotations that were significant in our marginal analysis (Table S11).

As an alternative to conditional analysis using S-LDSC, we analyzed various sets of annotations by training a gradient boosting model to classify $12,296$ SNPs from the NIH GWAS catalog[23] and assessing the AUROC, as in ref.[13,16] (Methods); although this is not a formal conditional analysis, comparing the AUROC achieved by different sets of annotations can provide an indication of which annotations provide unique information for disease. Results are reported in Table S12. We reached three main conclusions. First, the aggregated variant-level DeepSEA and Basenji annotations were only moderately informative for disease (AUROC $= 0.582$ and 0.612, respectively, consistent with moderate enrichments (DeepSEA: 1.49x, Basenji: 1.49x) of these annotations for NIH GWAS SNPs; Table S13). Second, including tissue-specific variant-level DeepSEA and Basenji annotations for all 127 tissues had little impact on the results (AUROC $= 0.591$ and 0.620, respectively). Third, the disease informativeness of the baseline-LD model (AUROC $= 0.758$) was not substantially impacted by adding the aggregated variant-level DeepSEA and Basenji annotations (AUROC $= 0.759$ and 0.766, respectively); results were similar for the non-tissue-specific joint model (baseline-LD model $+$ 7 annotations). These findings were consistent with our S-LDSC analyses.

We conclude that variant-level DeepSEA and Basenji annotations that were aggregated across tissues were highly enriched for heritability but not conditionally informative across the 41 traits, and that marginally significantly informative BiClassCNN annotations were explained away by local GC-content.

Variant-level annotations for blood and brain traits

We next evaluated the informativeness of blood-specific variant-level annotations for blood-related diseases and traits. We analyzed 11 independent blood-related traits (6 autoimmune diseases and 5 blood cell traits; ref.[29] and Table S2) by running S-LDSC conditioned on the non-tissue-specific variant-level joint model (baseline-LD model + 7 annotations; Figure S5).

We analyzed 8 blood-specific variant-level DeepSEA and 8 blood-specific variant-level Basenji annotations (Table S14), representing the blood-specific analogues of the non-tissue-specific variant-level DeepSEA and Basenji annotations from Table 1; in each case we computed the Average (Avg) or Maximum (Max) of cell-type-specific DeepSEA and Basenji variant-level annotations across blood cell types. In our marginal S-LDSC analysis, both blood-specific Basenji annotations and blood-specific DeepSEA annotations were very highly enriched (average enrichment 4.15x and 3.75x, respectively). However, only 0 blood-specific DeepSEA annotations and 4 blood-specific Basenji annotations attained a Bonferroni-significant standardized effect size ($\tau^\star$) conditional on the non-tissue-specific variant-level joint model (Figure 2 and Table S14).

We also analyzed 8 blood-specific Roadmap and 40 blood-specific ChromHMM annotations (Table S15), representing the blood-specific analogues of the non-tissue-specific Roadmap and ChromHMM annotations from Table S1. In our marginal S-LDSC analysis, all 8 blood-specific Roadmap annotations and 23 blood-specific ChromHMM annotations attained a Bonferroni-significant $\tau^\star$ (Figure S6, Figure S7 and Table S15).

We jointly analyzed the 35 blood-specific annotations that were Bonferroni-significant in marginal analyses (4 Basenji, 8 Roadmap, 23 ChromHMM; Figure 2, Figure S6 and Figure S7) by performing forward stepwise elimination to iteratively remove annotations that had conditionally non-significant $\tau^\star$ values after Bonferroni correction (based on the 80 (variant-level + allelic effect) blood-specific annotations tested in marginal analyses). Of these, 0 Basenji, 2 Roadmap and 4 ChromHMM annotations were jointly significant in the resulting joint model (Figure 2, Figure S8 and Table S16). We determined that the 2 surviving blood-specific Roadmap annotations were sufficient to eliminate 3 of the 4 blood-specific Basenji annotations that were significant in marginal analyses (Table S17). We note that several of the blood-specific Roadmap and ChromHMM annotations had $\tau^\star > 0.5$ in the blood-specific joint model (Figure S8) (annotations with $\tau^\star > 0.5$ are unusual, and considered to be important[35]), consistent with the well-documented importance of tissue-specific annotations, particularly for blood-related traits[5,11]. We also analyzed various sets of blood-specific annotations by training a gradient boosting model to classify 8,741 fine-mapped autoimmune disease SNPs[24] (relevant to blood-specific annotations only) and assessing the AUROC (analogous to Table S12). Results are reported in Table S18. We reached three main conclusions. First, the aggregated blood-specific variant-level DeepSEA and Basenji annotations were informative for disease, with Basenji being more informative (AUROC = 0.621 and 0.664, respectively, consistent with moderate enrichments (DeepSEA: 1.75x, Basenji: 1.97x) of these annotations for the fine-mapped SNPs; Table S19). Second, including cell-type-specific variant-level DeepSEA and Basenji annotations for all 27 blood cell types had little impact on the results (AUROC = 0.631 and 0.671, respectively). Third, the disease informativeness of the non-tissue-specific joint model (AUROC = 0.845) was not substantially impacted by adding aggregated blood-specific DeepSEA and Basenji variant-level annotations (AUROC = 0.849 and 0.853, respectively); results were similar for the the blood-specific joint model (non-tissue-specific joint model + 6 blood annotations). These findings were

consistent with our S-LDSC analyses.

We next evaluated the informativeness of brain-specific variant-level annotations for brain-related diseases and traits, analogous to blood. We analyzed 8 independent brain-related traits (ref.[29] and Table S2) by running S-LDSC conditioned on the non-tissue-specific variant-level joint model (baseline-LD model + 7 annotations; Figure S5).

We analyzed 8 brain-specific variant-level DeepSEA and 8 brain-specific variant-level Basenji annotations (Table S20), representing the brain-specific analogues of the non-tissue-specific variant-level DeepSEA and Basenji annotations from Table 1; in each case we computed the Average (Avg) or Maximum (Max) of tissue-type-specific DeepSEA and Basenji variant-level annotations across brain tissues. In our marginal S-LDSC analysis, both brain-specific DeepSEA and Basenji annotations were highly enriched (average enrichment 2.71x and 2.73x, respectively; Table S20). Furthermore, all 8 brain-specific DeepSEA annotations and all 8 blood-specific Basenji annotations attained a Bonferroni-significant standardized effect size ($\tau^\star$) conditional on the non-tissue-specific variant-level joint model (Figure 3 and Table S20).

We also analyzed 8 brain-specific Roadmap and 40 brain-specific ChromHMM annotations (Table S21), representing the brain-specific analogues of the non-tissue-specific Roadmap and ChromHMM annotations from Table S1. In our marginal S-LDSC analysis, all 8 brain-specific Roadmap annotations and 13 brain-specific ChromHMM annotations attained a Bonferroni-significant τ^* (Figure S9, Figure S10 and Table S21).

We jointly analyzed the 37 brain-specific annotations that were Bonferroni-significant in marginal analyses (8 DeepSEA, 8 Basenji, 8 Roadmap, 13 ChromHMM; Figure 3, Figure S9 and Figure S10) by performing forward stepwise elimination to iteratively remove annotations that had conditionally non-significant $\tau^\star$ values after Bonferroni correction (based on the 80 (variant-level + allelic-effect) brain-specific annotations tested in marginal analyses). Of these, 1 DeepSEA, 1 Basenji, 1 Roadmap and 3 ChromHMM annotations were jointly significant in the resulting joint model (Figure 3, Figure S11 and Table S22). We determined that neither the 1 surviving brain-specific Roadmap annotation nor the 3 surviving brain-specific ChromHMM annotations were sufficient to eliminate any of the 8 DeepSEA and 8 Basenji brain annotations that were significant in marginal analyses (Table S23 and Table S24). We note that none of the brain-specific DeepSEA, Basenji, Roadmap and ChromHMM annotations had $\tau^\star > 0.5$ in the brain-specific joint model (Figure 3 and Figure S11). We did not consider secondary analyses of fine-mapped SNPs for brain-related traits, due to the lack of a suitable resource analogous to ref.[24].

We investigated the k-mer composition of regions proximal to the 1 DeepSEA and 1 Basenji brain-specific variant-level annotations that were jointly significant (Figure 3). For each of these 2 continuous-valued annotations and each of all 682 possible k-mers with $1 \leq k \leq 5$ (merged with their reverse complements), we assessed the weighted k-mer enrichment in 1kb regions around each SNP in the annotation (Methods). Many CpG-related k-mers ($k \geq 3$) attained Bonferroni-significant enrichments, with the largest and most significant enrichments attained by CGCGC (8.8x and P=9.6e-55 for DeepSEA-H3K4me3-brain-Max, 2.8x and P=6.4e-70 for Basenji-H3K27ac-brain-Max; the latter is a larger annotation, such that smaller enrichments are expected) and CGGCG (8.7x and P=2.4e-53 for DeepSEA-H3K4me3-brain-Max, 2.8x and P=3.3e-72 for Basenji-H3K27ac-brain-Max) (Table S25). We hypothesize that these deep learning annotations could be capturing unique information about nucleosome occupancy, which is known to correlate

with the CGCG motif[36,37]; however, our analysis does not formally implicate nucleosome occupancy, and other explanations are also possible (although shorter motifs reflecting GC-content did not produce significant enrichments).

We conclude that blood-specific variant-level DeepSEA and Basenji annotations that were aggregated across blood cell types were very highly enriched for heritability but not uniquely informative for blood-related traits. On the other hand, brain-specific variant-level DeepSEA and Basenji annotations that were aggregated across brain tissues were uniquely informative for brain-related traits. This may be because brain tissues are not as well-represented in Roadmap data as blood cell types, leaving more room for new information to be retrieved by deep learning models.

Allelic-effect annotations for 41 diseases and traits

Distinct from variant-level annotations, we evaluated the informativeness of allelic-effect annotations (absolute value of the predicted difference between reference and variant alleles), which have been a major focus of recent work[13,14,15,16]. We evaluated the informativeness of 8 non-tissue-specific DeepSEA and 8 non-tissue-specific Basenji allelic-effect annotations (Table 1) for disease heritability by applying S-LDSC to the 41 traits (Table S2). We did not consider BiClassCNN allelic-effect annotations, whose interpretation for annotations such as coding, repressed and TSS is unclear. Analyses of allelic-effect annotations were conditioned on the non-tissue-specific variant-level joint model (baseline-LD model + 7 annotations; Figure S5); this distinguishes allelic effects from regional sequence context. For ease of comparison, allelic-effect Basenji annotations were quantile-matched with corresponding allelic-effect DeepSEA annotations, analogous to analyses of variant-level annotations.

In our marginal analysis of disease heritability using S-LDSC, allelic-effect Basenji annotations were more enriched for disease heritability (2.40x) than allelic-effect DeepSEA annotations (1.91x) (Table S26). However, only 0 DeepSEA annotations and 1 Basenji annotation, BasenjiΔ-H3K4me3-Max, attained a Bonferroni-significant standardized effect size ($\tau^\star$) conditional on the non-tissue-specific variant-level joint model (Figure 4 and Table S26); results were similar when conditioned on just the baseline-LD model (Table S27). Despite the high correlation between variant-level and allelic-effect annotations ($r = 0.48$; Figure S1), the corresponding variant-level annotation (Basenji-H3K4me3-Max) did not produce significant signal (Figure 4 and Table S28), consistent with our previous variant-level analysis (Figure 1). We note that since BasenjiΔ-H3K4me3-Max was the only marginally significant annotation in the non-tissue-specific allelic-effect analysis, the final non-tissue-specific joint model is defined by the addition of this annotation to non-tissue-specific variant-level joint model. A k-mer enrichment analysis (analogous to Table S25) indicated that BasenjiΔ-H3K4me3-Max was enriched for the k-mers CGCGC (4.1x and P=3.5e-10) and CGGCG (4.1x and P=3.6e-10), once again suggesting a potential connection to nucleosome occupancy (Table S29).

We also analyzed various sets of allelic-effect annotations by training a gradient boosting model to classify $12,296$ SNPs from the NIH GWAS catalog[23] and assessing the AUROC (analogous to Table S12), as in ref.[13,16] (Methods). Results are reported in Table S30. We reached three main conclusions. First, the aggregated allelic-effect DeepSEA and Basenji annotations were informative for disease (AUROC = 0.584 and 0.592, respectively, consistent with enrichments of these annotations (DeepSEA: 1.50x, Basenji: 1.75x) for NIH GWAS SNPs; Table S31). Second, including tissue-specific

allelic-effect DeepSEA and Basenji annotations for all 127 tissues slightly improved the results (AUROC = 0.602 and 0.611, respectively; lower than AUROC = 0.657 and 0.666 reported in ref.[16] because our analysis was restricted to chromatin marks and did not consider transcription factor binding site (TFBS) or cap analysis of gene expression (CAGE) data). Third, the disease informativeness of the non-tissue-specific variant-level joint model (AUROC = 0.762) was not substantially impacted by adding the aggregated allelic-effect DeepSEA and Basenji annotations (AUROC = 0.766 and 0.769, respectively). These findings were consistent with our S-LDSC analyses; in particular, the slightly higher AUROC for Basenji and DeepSEA allelic-effect annotations (across all analyses) was consistent with our S-LDSC results showing higher enrichments and a conditionally significant signal for Basenji annotations.

We conclude that allelic-effect DeepSEA and Basenji annotations that were aggregated across tissues were enriched for heritability across the 41 traits (with higher enrichments for Basenji), and that one Basenji allelic-effect annotation was conditionally informative.

Allelic-effect annotations for blood and brain traits

We next evaluated the informativeness of 8 blood-specific DeepSEA and 8 blood-specific Basenji allelic-effect annotations across 11 blood-related traits (Table S2) by running S-LDSC conditional on the blood-specific variant-level joint model (Figure S8) and BasenjiΔ-H3K4me3-Max (the 1 significant non-tissue-specific allelic-effect annotation; Figure 4 and Table S26).

In our marginal analysis of disease heritability using S-LDSC, blood-specific allelic-effect Basenji annotations were much more enriched for disease heritability (4.57x) than blood-specific allelic-effect DeepSEA annotations (2.20x), despite similar annotation sizes. However, none of the blood-specific allelic-effect annotations attained a Bonferroni-significant standardized effect size ($\tau^\star$) when conditioned on the blood-specific variant-level joint model and BasenjiΔ-H3K4me3-Max (Table S32). (In contrast, when we analyzed the blood-specific allelic-effect annotations conditional only on the non-tissue-specific final joint model, 0 DeepSEA annotations and 6 Basenji annotations attained a Bonferroni-significant $\tau^\star$ (Table S33); when these annotations were analyzed conditional only on the blood-specific variant-level joint model, 0 DeepSEA annotations and 1 Basenji annotation attained a Bonferroni-significant $\tau^\star$ (Table S34)).

We also analyzed various sets of blood-specific allelic-effect annotations by training a gradient boosting model to classify 8,741 fine-mapped autoimmune disease SNPs[24] (relevant to blood-specific annotations only) and assessing the AUROC (analogous to Table S18). Results are reported in Table S35. We reached three main conclusions. First, the aggregated blood-specific allelic-effect DeepSEA and Basenji annotations were informative for disease, with Basenji being more informative (AUROC = 0.613 and 0.672, respectively, consistent with moderate enrichments (DeepSEA: 1.71x, Basenji: 2.37x) of these annotations for the fine-mapped SNPs; Table S36). Second, including cell-type-specific allelic-effect DeepSEA and Basenji annotations for all 27 blood cell types slightly improved the results (AUROC = 0.633 and 0.684, respectively). Third, the disease informativeness of the blood-specific variant-level joint model plus BasenjiΔ-H3K4me3-Max (AUROC = 0.848) was not substantially impacted by adding the aggregated blood-specific DeepSEA and Basenji allelic-effect annotations (AUROC = 0.847 and 0.851, respectively). These findings were consistent with our S-LDSC analysis.

Finally, we evaluated the informativeness of 8 brain-specific DeepSEA and 8 brain-

specific Basenji allelic-effect annotations across 8 brain-related traits, by running S-LDSC conditional on the brain-specific variant-level joint model and BasenjiΔ-H3K4me3-Max. In our marginal S-LDSC analysis, brain-specific allelic-effect Basenji annotations were more enriched for disease heritability (2.53x) than brain-specific allelic-effect DeepSEA annotations (1.94x), despite similar annotation sizes. Two brain-specific allelic-effect Basenji annotations (BasenjiΔ-H3K4me3-brain-Max and BasenjiΔ-H3K4me3-brain-Avg) attained a Bonferroni-significant standardized effect size ($\tau^\star$) when conditioned on the brain-specific variant-level joint model and BasenjiΔ-H3K4me3-Max (Figure 5 and Table S37). (In contrast, when we analyzed the brain-specific allelic-effect annotations conditional only on the non-tissue-specific final joint model, 7 DeepSEA annotations and 7 Basenji annotations attained a Bonferroni-significant $\tau^\star$ (Table S38); when these annotations were analyzed conditional only on the brain-specific variant-level joint model, 0 DeepSEA annotations and 3 Basenji annotations attained a Bonferroni-significant $\tau^\star$ (Table S39)). Despite the high correlation between variant-level and allelic-effect annotations ($r = 0.48$; Figure S1), the corresponding variant-level annotations (Basenji-H3K4me3-brain-Max and Basenji-H3K4me3-brain-Avg) did not produce significant signal (Figure 5 and Table S40), consistent with our previous variant-level analysis (Figure 3).

We jointly analyzed the two annotations, BasenjiΔ-H3K4me3-brain-Max and BasenjiΔ-H3K4me3-brain-Avg, that were Bonferroni-significant in marginal analyses (Figure 5) by performing forward stepwise elimination to iteratively remove annotations that had conditionally non-significant $\tau^\star$ values after Bonferroni correction (based on the 80 variant-level and allelic-effect brain-specific annotations tested in marginal analyses). Of these, only BasenjiΔ-H3K4me3-brain-Max was jointly significant in the resulting brain-specific final joint model, with $\tau^\star$ very close to 0.5 (Figure 5, Table S37 and Table S41). A k-mer enrichment analysis (analogous to Table S25) indicated that BasenjiΔ-H3K4me3-brain-Max was enriched for the k-mers CGCGC (6.2x and p=1.1e-25) and CGGCG (6.1x and p=4.9e-25), once again suggesting a potential connection to nucleosome occupancy (Table S29). We did not consider secondary analyses of fine-mapped SNPs for brain-related traits, due to the lack of a suitable resource analogous to ref.[24].

We conclude that blood-specific allelic-effect annotations were very highly enriched for heritability but not uniquely informative for blood-related traits, whereas one brain-specific allelic-effect annotation was uniquely informative for brain-related traits, similar to our variant-level analyses. Blood-specific and brain-specific allelic-effect Basenji annotations generally outperformed DeepSEA annotations, yielding higher enrichments and the sole conditionally significant annotation, similar to our non-tissue-specific allelic-effect analyses.

Discussion

We have evaluated the informativeness for disease of variant-level and allelic-effect annotations constructed using two previously trained deep learning models, DeepSEA[13,15] and Basenji[16]. We evaluated each annotation's informativeness using S-LDSC[5,19]; as a secondary metric, we also evaluated the accuracy of gradient boosting models incorporating deep learning annotations in predicting disease-associated or fine-mapped SNPs[23,24], as in previous work[13,16]. In non-tissue-specific analyses, we identified one allelic-effect Basenji annotation that was uniquely informative for 41 diseases and complex traits. In blood-specific analyses, we identified no deep learning annotations that were uniquely informative for 11 blood-related traits. In brain-specific analyses, we identified

brain-specific variant-level DeepSEA and Basenji annotations, as well as a brain-specific allelic-effect Basenji annotation, that were uniquely informative for 8 brain-related traits. Sequence motif analyses suggest that the uniquely informative annotations could be capturing unique information about nucleosome occupancy, although other explanations are also possible.

Our results imply that the informativeness of deep learning annotations for disease cannot be inferred from metrics such as AUROC that evaluate their accuracy in predicting underlying regulatory annotations derived from experimental assays. Instead, deep learning annotations must be evaluated using methods that specifically evaluate their informativeness for disease, conditional on a broad set of other functional annotations; the S-LDSC method that we applied here is one such method, and the accuracy of gradient boosting models incorporating both deep learning annotations and other functional annotations can also be a useful metric. We emphasize the importance of conditioning on a broad set of functional annotations, in order to assess whether deep learning models leveraging DNA sequence provide unique (as opposed to redundant) information; previous work has robustly linked deep learning annotations to disease[12,13,14,15,16], but those analyses did not condition on a broad set of other functional annotations.

Our work has several limitations, representing important directions for future research. First, our analyses of deep learning annotations using S-LDSC are inherently focused on common variants, but deep learning models have also shown promise in prioritizing rare pathogenic variants[15,38,39]. The value of deep learning models for prioritizing rare pathogenic variants has been questioned in a recent analysis focusing on Human Gene Mutation Database (HGMD) variants[40], meriting further investigation. Second, our analyses of allelic-effect annotations are restricted to unsigned analyses, but signed analyses have also proven valuable in linking deep learning annotations to molecular traits and complex disease[16,41,42]; however, genome-wide signed relationships are unlikely to hold for the regulatory marks (DNase and histone marks) that we focus on here, which do not correspond to specific genes or pathways. Third, we focused here on deep learning models trained to predict specific regulatory marks, but deep learning models have also been used to predict a broader set of regulatory features, including gene expression levels and cryptic splicing[15,16,39], that may be informative for complex disease. We have also not considered the application of deep learning models to TFBS, CAGE and ATAC-seq data[16,42], which is a promising future research direction. Fourth, we focused here on deep learning models trained using human data, but models trained using data from other species may also be informative for human disease[43,42]. Fifth, the forward stepwise elimination procedure that we use to identify jointly significant annotations[19] is a heuristic procedure whose choice of prioritized annotations may be close to arbitrary in the case of highly correlated annotations; nonetheless, our framework does impose rigorous criteria for conditional informativeness. Finally, beyond deep learning models, it is of high interest to evaluate other machine learning methods for predicting regulatory effects[44,45,46,47,48].

URLs

- Annotations analyzed in this study:
  https://data.broadinstitute.org/alkesgroup/LDSCORE/DeepLearning/

- 1000 Genomes Project Phase 3 data:
  ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502

- UK Biobank summary statistics:
  `https://data.broadinstitute.org/alkesgroup/UKBB/`

- Baseline-LD annotations:
  `https://data.broadinstitute.org/alkesgroup/LDSCORE/`

- BOLT-LMM software:
  `https://data.broadinstitute.org/alkesgroup/BOLT-LMM`

- DeepSEA pre-trained model:
  `https://github.com/FunctionLab/ExPecto`

- Basenji pre-trained model:
  `https://github.com/calico/basenji`

Acknowledgments

We thank Huwenbo Shi and Steven Gazal for helpful discussions. This research was funded by NIH grants U01 HG009379, R01 MH101244, and R01 MH107649. This research was conducted using the UK Biobank Resource under application 16549.

Methods

Genomic annotations and the baseline-LD model

We define a functional annotation as an assignment of a numeric value to each SNP; annotations can be either binary or continuous-valued (Methods). Our focus is on continuous-valued annotations (with values between 0 and 1) trained by deep learning models to predict biological function from DNA sequenceWe define a genomic annotation as an assignment of a numeric value to each SNP in a predefined reference panel (e.g., 1000 Genomes Project[26]; see URLs). Binary annotations can have value 0 or 1 only. Continuous-valued annotations can have any real value; our focus is on continuous-valued annotations with values between 0 and 1. Annotations that correspond to known or predicted function are referred to as functional annotations. The baseline-LD model (v.2.1) contains 86 functional annotations (see URLs). These annotations include binary coding, conserved, and regulatory annotations (e.g., promoter, enhancer, histone marks, TFBS) and continuous-valued linkage disequilibrium (LD)-related annotations.

DeepSEA and Basenji annotations

Tissue-specific deep learning annotations were derived using two pre-trained Convolutional Neural Net (CNN) models: DeepSEA[13,15] (architecture from ref.[15]) and Basenji[16] (see URLs). DeepSEA is a classification based model trained on binary peak call data from $2,002$ cell-type specific TFBS, histone mark and chromatin accessibility annotations from the ENCODE[21] and Roadmap Epigenomics[11] projects. Basenji is a Poisson prediction based model trained on original count data from $4,229$ cell-type specific histone mark, chromatin accessibility and FANTOM5 CAGE[49,50] annotations. Additionally, Basenji uses dilated convolutional layers that allow scanning much larger contiguous sequence around a variant ($\approx$ 130kb) compared to DeepSEA (1kb). We restricted our analyses to DNase-I Hypersensitivity Sites (DHS) and 3 histone marks (H3K27ac, H3K4me1 and H3K4me3) that are known to be associated with active enhancers and promoters[51].

For each SNP with minor allele count ≥ 5 in 1000 Genomes, we applied the pre-trained DeepSEA and Basenji models to the surrounding DNA sequence (based on the reference allele) to compute the predicted probability of a tissue-specific chromatin mark (DNase, H3K27ac, H3K4me1, H3K4me3) to generate the corresponding variant-level annotation. To generate the corresponding allelic-effect annotation, we compute the predicted difference in probability between the reference and the alternate alleles. The Basenji annotations were quantile-matched to corresponding DeepSEA annotations to ensure a fair comparison of the two approaches. We aggregated these probabilistic annotations across all 127 Roadmap tissues by taking either the average (Avg) or maximum (Max) to generate non-tissue specific annotations, yielding 8 DeepSEA annotations and 8 Basenji annotations. Similarly, we aggregated over 27 blood cell types (resp. 13 brain tissues) to generate blood (resp. brain) specific annotations for each chromatin mark.

BiClassCNN annotations

We used trained a new deep learning model, BiClassCNN, to prioritize SNPs within non-tissue-specific annotations. BiClassCNN analyzes 1kb of human reference sequence around each SNP (analogous to DeepSEA). The positive training set for BiClassCNN consists of 1kb of reference sequence around SNPs that are known to have the functionality of interest (e.g. coding); we included all such sequences in the positive training set. The negative training set consists of 1kb of reference sequence around SNPs that are 1kb away from all SNPs with the functionality of interest; we included a subset of such sequences in the negative training set, so as to match the overall size, GC content and repeat element content of the positive set (as in ref.[44,52]). We used a shallow Convolutional Neural Net architecture for training (see Figure S12).

We ran two training models, one for the even chromosomes and one for odd chromosomes, and used the trained model on even (resp. odd) chromosomes to assign a predicted probability of functionality (e.g. coding), based on sequence context, to each SNP on odd (resp. even) chromosomes. Unlike DeepSEA and Basenji, BiClassCNN annotations were restricted to regions of known functionality (e.g. coding) by setting annotation values to 0 outside those regions; thus, BiClassCNN prioritizes SNPs within regions of known functionality (e.g. coding). (BiClassCNN annotations that were not restricted in this fashion were far less informative for disease.)

We restricted S-LDSC analyses of BiClassCNN annotations to annotations for which the BiClassCNN AUROC value was at least 0.6 (Table 1 and Table S3). This eliminated three annotations (Intron, H3K27ac and UTR-3'), leaving a total of 12 BiClassCNN annotations.

Other annotations

We also considered:

- (Table S7) 8 Roadmap annotations[11] (analogous to DeepSEA and Basenji annotations) imputed using ChromImpute[20].

- (Table S7) 40 ChromHMM annotations[21,22] based on 20 ChromHMM states across 127 Roadmap tissues[11], again aggregated using the average (Avg) or maximum (Max) across tissues.

- (Table S8) 12 annotations consisting of CpG-island, local CpG-content and local GC-content annotations, as well as these annotations restricted to coding, repressed and TSS regions (for which BiClassCNN produced conditionally significant signals). The CpG-island annotation was retrieved from the UCSC genome browser[53]. Local CpG-content and local GC-content denote the proportion of CpG and $G + C$ dinuclotides in ± 1kb regions around each variant of the genome, computed using the hg19 reference genome fasta file. By definition, the LocalGCcontent annotation is of larger size than the LocalCpGcontent annotation.

- (Table S8) 3 annotations consisting of a pLI annotation, as well as this annotation restricted to coding and TSS regions. The pLI annotation was defined by annotating each SNP in a 5kb window around a gene with the pLI score of that gene[31]. We did not consider the pLI annotation restricted to repressed regions because unlike TSS and coding, repressed regions are not directly linked to a gene.

- (Table S8) 2 coding annotations, SIFT[34] and Polyphen[32,33], which have been analyzed in previous work[54,55].

Stratified LD score regression

Stratified LD score regression (S-LDSC) is a method that assesses the contribution of a genomic annotation to disease and complex trait heritability[5,19]. S-LDSC assumes that the per-SNP heritability or variance of effect size (of standardized genotype on trait) of each SNP is equal to the linear contribution of each annotation

$$var\left(\beta_j\right) := \sum_c a_{cj}\tau_c, \tag{1}$$

where a_{cj} is the value of annotation c for SNP j, where a_{cj} may be binary (0/1), continuous or probabilistic, and τ_c is the contribution of annotation c to per-SNP heritability conditioned on other annotations. S-LDSC estimates the τ_c for each annotation using the following equation

$$E\left[\chi_j^2\right] = N\sum_c l(j,c)\tau_c + 1, \tag{2}$$

where $l(j,c) = \sum_k a_{ck}r_{jk}^2$ is the *stratified LD score* of SNP j with respect to annotation c and r_{jk} is the genotypic correlation between SNPs j and k computed using data from 1000 Genomes Project[26] (see URLs); N is the GWAS sample size.

We assess the informativeness of an annotation c using two metrics. The first metric is enrichment (E), defined as follows (for binary and probabilistic annotations only):

$$E = \frac{\frac{h_g^2(c)}{h_g^2}}{\frac{\sum_j a_{cj}}{M}}, \tag{3}$$

where $h_g^2(c)$ is the heritability explained by the SNPs in annotation c, weighted by the annotation values.

The second metric is standardized effect size ($\tau^\star$) defined as follows (for binary, probabilistic, and continuous-valued annotations):

$$\tau_c^\star = \frac{\tau_c sd_c}{\frac{h_g^2}{M}}, \tag{4}$$

where sd_c is the standard error of annotation c, h_g^2 the total SNP heritability and M is the total number of SNPs on which this heritability is computed (equal to $5,961,159$ in our analyses). $\tau_c^\star$ represents the proportionate change in per-SNP heritability associated to a 1 standard deviation increase in the value of the annotation.

Weighted k-mer enrichment Analysis

We performed weighted k-mer enrichment analyses of the four deep learning annotations (DeepSEA-H3K4me3-Brain-Max, Basenji-H3K27ac-brain-Max, BasenjiΔ-H3K4me3-Max and BasenjiΔ-H3K4me3-brain-Max) that were conditionally informative for disease heritability (Figure 3, Figure4, Figure5), for all 682 possible k-mers with $1 \leq k \leq 5$ (merged with their reverse complements). Results of these analyses are reported in Table S25 and Table S29.

For each k-mer i, we computed k-mer counts $\kappa_s^{(i)}$ in the 1kb regions around each SNP s in the genome.

For each deep learning annotation D, for each k-mer i, we computed the weighted average $\mathcal{W}_D^{(i)}$ of k-mer counts $\kappa^{(i)}$, weighted by values of the probabilistic annotation:

$$\mathcal{W}_D^{(i)} := \sum_s D_s \kappa_s^{(i)}. \tag{5}$$

We compared $\mathcal{W}_D^{(i)}$ with $\mathcal{W}_{D^{\text{null}}}^{(i)}$, where D^{null} is defined as the probabilistic annotation with all valuesuniformly equal to $\bar{D}$, the average value (annotation size) of annotation D.

We computed the *weighted k-mer enrichment* of annotation D with respect to k-mer i as

$$WKE_D^{(i)} := \mathcal{W}_D^{(i)} / \mathcal{W}_{D^{\text{null}}}^{(i)} \tag{6}$$

We assessed the statistical significance of the weighted k-mer enrichment via a permutation test in which we randomly permuted the values of the deep learning annotation D across SNPs and compared $WKE_D^{(i)}$ to values of $WKE_{D^{\text{perm}}}^{(i)}$ for each permuted annotation D^{perm}. We computed p-values by fitting a Gaussian distribution to the values of $WKE_{D^{\text{perm}}}^{(i)}$ across 10,000 such permutations.

Classification of disease-associated or fine-mapped SNPs

As an alternative to conditional analysis using S-LDSC, we evaluated the efficacy of various sets of annotations for classifying 12,296 disease-associated SNPs from the NIH GWAS catalog[23] (as in ref.[13,16]) or 8,741 fine-mapped autoimmune disease SNPs[24] against the same number of control SNPs, matched for minor allele frequency. We used XGBoost, a machine learning technique based on gradient tree boosting[56,57]. To optimize classification performance, we selected XGBoost parameter settings to minimize overfitting, as in ref.[58].

Tables

Table 1. List of non-tissue-specific deep learning annotations analyzed. We list the 25 variant-level deep learning annotations (8 DeepSEA, 8 Basenji, 9 BiClass-CNN) and their annotation sizes (average annotation value across SNPs), and 16 allelic-effect deep learning annotations (8 DeepSEAΔ, 8 BasenjiΔ) and their annotation sizes.

Variant-level annotations	Size (%)
DeepSEA-DNase-Avg	1.6
DeepSEA-DNase-Max	13.2
DeepSEA-H3K27ac-Avg	3.2
DeepSEA-H3K27ac-Max	13.7
DeepSEA-H3K4me1-Avg	5.5
DeepSEA-H3K4me1-Max	23.2
DeepSEA-H3K4me3-Avg	1.7
DeepSEA-H3K4me3-Max	8.8
Basenji-DNase-Avg	1.6
Basenji-DNase-Max	13.1
Basenji-H3K27ac-Avg	3.2
Basenji-H3K27ac-Max	13.7
Basenji-H3K4me1-Avg	5.5
Basenji-H3K4me1-Max	23.2
Basenji-H3K4me3-Avg	1.7
Basenji-H3K4me3-Max	8.8
BiClassCNN-Coding	1.0
BiClassCNN-Enhancer	2.6
BiClassCNN-Promoter	2.7
BiClassCNN-Repressed	36.0
BiClassCNN-SuperEnhancer	6.7
BiClassCNN-TFBS	7.9
BiClassCNN-TSS	0.8
BiClassCNN-UTR-5'	0.4
BiClassCNN-WeakEnhancer	1.1
Allelic-effect annotations	Size (%)
DeepSEAΔ-DNase-Avg	0.3
DeepSEAΔ-DNase-Max	2.0
DeepSEAΔ-H3K27ac-Avg	0.2
DeepSEAΔ-H3K27ac-Max	0.9
DeepSEAΔ-H3K4me1-Avg	0.3
DeepSEAΔ-H3K4me1-Max	1.7
DeepSEAΔ-H3K4me3-Avg	0.1
DeepSEAΔ-H3K4me3-Max	0.7
BasenjiΔ-DNase-Avg	0.3
BasenjiΔ-DNase-Max	2.1
BasenjiΔ-H3K27ac-Avg	0.2
BasenjiΔ-H3K27ac-Max	0.9
BasenjiΔ-H3K4me1-Avg	0.3
BasenjiΔ-H3K4me1-Max	1.7
BasenjiΔ-H3K4me3-Avg	0.1
BasenjiΔ-H3K4me3-Max	0.7

Figures

Figure 1. Disease informativeness of non-tissue-specific variant-level deep learning annotations: (A) Heritability enrichment, conditional on the baseline-LD model. Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditional on either the baseline-LD model (marginal analyses: left column, white) or the baseline-LD model plus 1 Roadmap, 1 ChromHMM, 3 LocalGCcontent and 2 pLI annotations (non-tissue-specific variant-level joint model: right column, shaded). Results are meta-analyzed across 41 traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 106 (variant-level + allelic-effect) non-tissue-specific annotations tested ($P < 0.05/106$). None of these annotations were significant after conditioning on the non-tissue-specific variant-level joint model (right column). Error bars denote 95% confidence intervals. Numerical results are reported in Table S4, Table S5 and Table S9.

Figure 2. Disease informativeness of blood-specific variant-level deep learning annotations: (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model. Horizontal line denotes no enrichment. (B) Standardized effect size $\tau^\star$ conditioned on either the non-tissue-specific variant-level joint model (marginal analysis: left column, white) or the non-tissue-specific variant-level joint model plus 2 Roadmap and 4 ChromHMM blood-specific annotations (blood-specific variant-level joint model: right column, dark shading). Results are meta-analyzed across 11 blood-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). None of these annotations were significant after conditioning on the blood-specific variant-level joint model. Error bars denote 95% confidence intervals. Numerical results are reported in Table S14 and Table S16.

Figure 3. Disease informativeness of brain-specific variant-level deep learning annotations: (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model. Horizontal line denotes no enrichment. (B) Standardized effect size $\tau^\star$ conditioned on either the non-tissue-specific variant-level joint model (marginal analysis: left column, white) or the non-tissue-specific variant-level joint model plus 1 DeepSEA, 1 Basenji, 1 Roadmap and 3 ChromHMM brain-specific annotations (brain-specific variant-level joint model: right column, dark shading). Results are meta-analyzed across 8 brain-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) brain-specific annotations tested ($P < 0.05/80$). For brain-specific variant-level joint model, ** denotes $P < 0.05/80$. Error bars denote 95% confidence intervals. Numerical results are reported in Table S20 and Table S22.

Figure 4. Disease informativeness of non-tissue-specific allelic-effect deep learning annotations: (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model. Horizontal line denotes no enrichment. (B) Standardized effect size $\tau^\star$ conditioned on either the non-tissue-specific variant-level joint model (marginal analysis: left column, white) or the variant-level joint model plus 1 non-tissue-specific allelic-effect Basenji annotation (BasenjiΔ-H3K4me3-Max) (non-tissue-specific final joint model: right column, dark shading. Results are meta-analyzed across 41 traits. Results are displayed only for the allelic-effect annotation (BasenjiΔ-H3K4me3-Max) with significant $\tau^\star$ in marginal analyses after correcting for 106 (variant-level + allelic-effect) non-tissue-specific annotations tested ($P < 0.05/106$), along with the corresponding variant-level annotation; the correlation between the two annotations is 0.43. For non-tissue-specific final joint model (right column), ** denotes $P < 0.05/106$. Error bars denote 95% confidence intervals. Numerical results are reported in Table S26 and Table S28.

Figure 5. Disease informativeness of brain-specific allelic-effect deep learning annotations: (A) Heritability enrichment, conditioned on the brain-specific variant-level joint model and the 1 significant non-tissue-specific allelic-effect annotation (BasenjiΔ-H3K4me3-Max). Horizontal line denotes no enrichment. (B) Standardized effect size $\tau^\star$ conditioned on either the brain-specific variant-level joint model and BasenjiΔ-H3K4me3-Max (marginal analysis: left column, white) or the same model plus 1 brain-specific allelic-effect annotation (BasenjiΔ-H3K4me3-brain-Max) (brain-specific final joint model: right column, dark shading). Results are meta-analyzed across 8 brain-related traits. Results are displayed only for the 2 allelic-effect annotations with significant τ^* in marginal analyses after correcting for 80 (variant-level + allelic-effect) brain-specific annotations tested ($P < 0.05/80$), along with the corresponding variant-level annotations; the correlation between the two allelic-effect annotations is 0.78, and the average correlation between the two pairs of variant-level (Basenji) and allelic-effect (BasenjiΔ) annotations is 0.44. For brain-specific final joint model (right column), ** denotes $P < 0.05/80$. Error bars denote 95% confidence intervals. Numerical results are reported in Table S37, Table S40 and Table S41.

Table S1. List of non-tissue-specific Roadmap and ChromHMM annotations analyzed. We list the 8 Roadmap and 40 ChromHMM annotations that we analyzed, together with the size of the annotation. These annotations are probabilistic, thus the size of the annotation is defined as the average annotation across all reference SNPs.

Annotation	Size (in %)
Roadmap-DNase-Avg	2.2%
Roadmap-H3K27ac-Avg	3.1%
Roadmap-H3K4me1-Avg	4.4%
Roadmap-H3K4me3-Avg	1.7%
Roadmap-DNase-Max	13.4%
Roadmap-H3K27ac-Max	21.4%
Roadmap-H3K4me1-Max	25.4%
Roadmap-H3K4me3-Max	6.6%
ChromHMM-DNase-Avg	0.7%
ChromHMM-Active_Enhancer_1-Avg	0.3%
ChromHMM-Active_Enhancer_2-Avg	0.4%
ChromHMM-Enhancer_acetylation-Avg	0.3%
ChromHMM-Active_Enhancer_Flanking-Avg	0.5%
ChromHMM-Weak_Enhancer_1-Avg	0.3%
ChromHMM-Weak_Enhancer_2-Avg	1.2%
ChromHMM-Heterochromatin-Avg	1.4%
ChromHMM-Promoter_Bivalent-Avg	0.3%
ChromHMM-Promoter_Downstream_1-Avg	0.4%
ChromHMM-Promoter_Downstream_2-Avg	0.2%
ChromHMM-Promoter_Poised-Avg	0.2%
ChromHMM-Promoter_Upstream-Avg	0.4%
ChromHMM-Quiescent-Avg	78.4%
ChromHMM-Repressed_Polycomb-Avg	1.7%
ChromHMM-TSS-Avg	0.2%
ChromHMM-Transcription_Enhancer_3'-Avg	0.2%
ChromHMM-Transcription_Enhancer_5'-Avg	0.4%
ChromHMM-Transcription_Weak_Enhancer-Avg	0.5%
ChromHMM-Transcriptional_Regulator-Avg	0.3%
ChromHMM-DNase-Max	9.2%
ChromHMM-Active_Enhancer_1-Max	7.1%
ChromHMM-Active_Enhancer_2-Max	9.9%
ChromHMM-Enhancer_acetylation-Max	11.3%
ChromHMM-Active_Enhancer_Flanking-Max	11.1%
ChromHMM-Weak_Enhancer_1-Max	7.7%
ChromHMM-Weak_Enhancer_2-Max	19.9%
ChromHMM-Heterochromatin-Max	7.1%
ChromHMM-Promoter_Bivalent-Max	1.9%
ChromHMM-Promoter_Downstream_1-Max	1.4%
ChromHMM-Promoter_Downstream_2-Max	1.5%
ChromHMM-Promoter_Poised-Max	6.7%
ChromHMM-Promoter_Upstream-Max	2.5%
ChromHMM-Quiescent-Max	91.0%
ChromHMM-Repressed_Polycomb-Max	9.9%
ChromHMM-TSS-Max	0.9%
ChromHMM-Transcription_Enhancer_3'-Max	3.1%
ChromHMM-Transcription_Enhancer_5'-Max	4.8%
ChromHMM-Transcription_Weak_Enhancer-Max	4.7%
ChromHMM-Transcriptional_Regulator-Max	3.3%

Table S2. List of 41 diseases and complex traits analyzed. We first list the 11 blood-related traits (5 blood cell traits and 6 autoimmune diseases), followed by 10 brain traits (with 2 traits analyzed by two different datasets). Overall, for 6 traits we analyzed two different data sets, leading to a total of 47 data sets.

Trait	Source	N
Platelet Count	UKBiobank[30]	444382
Red Blood Cell Count	UKBiobank[30]	445174
Red Blood Cell Distribution Width	UKBiobank[30]	442700
Eosinophil Count	UKBiobank[30]	439938
White Blood Cell Count	UKBiobank[30]	444502
Auto Immune Traits (Sure)	UKBiobank[30]	459324
Crohn's Disease	Jostins et al., 2012 Nature[59]	20883
Rheumatoid Arthritis	Okada et al., 2014 Nature[60]	37681
Ulcerative Colitis	Jostins et al., 2012 Nature[59]	27432
Lupus	Bentham et al., 2015[61]	14267
Celiac	Dubois et al., 2010[62]	15283
Age at Menarche	UKBiobank[30]	242278
BMI	Speliotes et al., 2010 Nat Genet[63]	122033
BMI	UKBiobank[30]	457824
Depressive symptoms	Okbay et al., 2016 Nat Genet[64]	161460
Neuroticism	UKBiobank[30]	372066
Schizophrenia	SCZ Working Group of the PGC, 2014 Nature[4]	70100
Years of Education	Okbay et al., 2016 Nature[64]	328917
Ever Smoked	TAG Consortium, 2010 Nat Genet[65]	74035
Smoking Status	UKBiobank[30]	457683
Bipolar Disorder	PGC Bipolar Disorder Group[66]	16731
Age at Menopause	UKBiobank[30]	143025
Age first birth	Barban et al., 2016 Nat Genet[67]	222037
Anorexia	Boraska et al., 2014 Mol Psych[68]	32143
Autism Spectrum	PGC Cross-Disorder Group, 2013 Lancet[69]	10263
College Education	UKBiobank[30]	454813
Coronary Artery Disease	Schunkert et al., 2011 Nat Genet[70]	77210
Dermatologic Diseases	UKBiobank[30]	459324
Eczema	UKBiobank[30]	458699
FEV1-FVC Ratio	UKBiobank[30]	371949
Forced Vital Capacity (FVC)	UKBiobank[30]	371949
Hair Color	UKBiobank[30]	452720
HDL	Teslovich et al., 2010 Nature[71]	97749
Heel T Score	UKBiobank[30]	445921
Height	Lango Allen et al., 2010 Nature[72]	131547
Height	UKBiobank[30]	458303
High Cholesterol	UKBiobank[30]	459324
Hypothyroidism	UKBiobank[30]	459324
LDL	Teslovich et al., 2010 Nature[71]	93354
Morning Person	UKBiobank[30]	410520
Number children ever born	Barban et al., 2016 Nat Genet[67]	318863
Respiratory and Ear-nose-throat Diseases	UKBiobank[30]	459324
Skin Color	UKBiobank[30]	453609
Sunburn Occasion	UKBiobank[30]	344229
Tanning	UKBiobank[30]	449984
Type 2 Diabetes	Morris et al., 2012 Nat Genet[73]	60786
Type 2 Diabetes	UKBiobank[30]	459324

Table S3. AUROC values of deep learning annotations. AUROC (Area under the ROC curve) for DeepSEA (reported by ref.[15]) and BiClassCNN annotations. For the DeepSEA model, the Minimum-Maximum (Average) AUROC across all tissues is reported. The constituent tissue-specific Basenji annotations do not lie on a probabilistic scale, and ref.[16] did not report AUROC values

Annotation	Area under ROC curve (AUROC)
DeepSEA-DNase	All: 0.83-0.95 (0.91)
	brain: 0.84-0.92 (0.88)
	blood: 0.86-0.94 (0.89)
DeepSEA-H3K27ac	All: 0.78-0.88 (0.84)
	brain: 0.80-0.85 (0.82)
	blood: 0.78-0.87 (0.83)
DeepSEA-H3K4me1	All: 0.77-0.86 (0.81)
	brain: 0.79-0.83 (0.80)
	blood: 0.78-0.83 (0.81)
DeepSEA-H3K4me3	All: 0.88-0.97 (0.92)
	brain: 0.89-0.95 (0.91)
	blood: 0.88-0.95 (0.92)
BiClassCNN-Coding	0.79
BiClassCNN-DNase	0.69
BiClassCNN-Enhancer	0.72
BiClassCNN-H3K4me1	0.70
BiClassCNN-H3K4me3	0.73
BiClassCNN-Promoter	0.76
BiClassCNN-Repressed	0.70
BiClassCNN-SuperEnhancer	0.67
BiClassCNN-TFBS	0.75
BiClassCNN-TSS	0.84
BiClassCNN-UTR-5'	0.75
BiClassCNN-WeakEnhancer	0.74

Table S4. S-LDSC results for marginal analysis of non-tissue-specific variant-level DeepSEA and Basenji annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 non-tissue-specific variant-level DeepSEA and 8 non-tissue-specific variant-level Basenji annotations, when conditioned on 86 baseline-LD annotations. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Basenji-DNase-Avg (1.7%)	-0.078	0.07	0.26	2.4	0.2	4.9e-09
Basenji-DNase-Max (13.6%)	-0.095	0.065	0.14	1.8	0.079	1.7e-16
Basenji-H3K27ac-Avg (3.4%)	0.14	0.059	0.016	2.5	0.14	2.7e-16
Basenji-H3K27ac-Max (14.2%)	-0.026	0.031	0.41	1.8	0.074	9.4e-16
Basenji-H3K4me1-Avg (5.6%)	0.08	0.042	0.055	2	0.076	2.7e-17
Basenji-H3K4me1-Max (23.7%)	-0.057	0.027	0.032	1.6	0.046	1.4e-17
Basenji-H3K4me3-Avg (2.0%)	0.17	0.07	0.014	3.7	0.26	1.4e-16
Basenji-H3K4me3-Max (9.4%)	0.079	0.036	0.029	2	0.088	5.3e-15
DeepSEA-DNase-Avg (1.7%)	-0.12	0.11	0.27	2.3	0.2	3e-07
DeepSEA-DNase-Max (13.6%)	-0.17	0.078	0.028	1.8	0.075	1.1e-15
DeepSEA-H3K27ac-Avg (3.4%)	0.19	0.071	0.0089	2.4	0.12	1.1e-17
DeepSEA-H3K27ac-Max (14.2%)	0.0092	0.06	0.88	1.8	0.066	2.6e-17
DeepSEA-H3K4me1-Avg (5.6%)	0.015	0.059	0.8	1.9	0.059	2.2e-17
DeepSEA-H3K4me1-Max (23.7%)	-0.13	0.06	0.027	1.6	0.041	8.8e-18
DeepSEA-H3K4me3-Avg (2.0%)	0.27	0.093	0.0033	4	0.25	7.1e-19
DeepSEA-H3K4me3-Max (9.4%)	0.25	0.06	4.5e-05	2.2	0.087	2.2e-17

Table S5. S-LDSC results for marginal analysis of non-tissue-specific Bi-ClassCNN annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 12 non-tissue-specific BiClassCNN annotations. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
BiClassCNN-Coding (1.3%)	0.93	0.16	2.5e-09	4.9	0.34	1e-12
BiClassCNN-DNase (9.1%)	-0.02	0.04	0.59	2.1	0.18	7.9e-06
BiClassCNN-Enhancer (2.8%)	0.31	0.18	0.09	3.4	0.25	2.3e-17
BiClassCNN-H3K4me1 (29.6%)	-0.05	0.07	0.50	1.8	0.073	4.3e-18
BiClassCNN-H3K4me3 (7.4%)	0.42	0.13	0.0019	2.9	0.18	8.8e-16
BiClassCNN-Promoter (2.7%)	0.19	0.05	0.00012	2.5	0.16	8.4e-17
BiClassCNN-Repressed (31.8%)	0.24	0.05	2.6e-06	0.68	0.048	1.2e-05
BiClassCNN-SuperEnhancer (8.0%)	-0.03	0.04	0.42	2.2	0.11	1.8e-15
BiClassCNN-TFBS (8.2%)	0.36	0.13	0.0041	2.9	0.21	2.3e-14
BiClassCNN-TSS (1.0%)	0.8	0.13	3.1e-10	6.9	0.61	1e-19
BiClassCNN-UTR-5' (0.5%)	0.017	0.12	0.89	2.9	0.49	1.7e-06
BiClassCNN-WeakEnhancer (1.3%)	-0.15	0.13	0.26	2.3	0.24	1.9e-14

Table S6. . BiClassCNN-Coding, BiClassCNN-Repressed and BiClassCNN-TSS produced independent signals.. We report standardized effect sizes ($\tau^\star$) and enrichment (E) of the 4 significant BiClassCNN annotations from Figure 1, when modeled jointly and passed through the forward stepwise elimination, all conditioned on the baseline-LD model. The three annotations that remain are BiClassCNN-Coding, BiClassCNN-Repressed and BiClassCNN-TSS. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
BiClassCNN-Coding (1.3%)	0.73	0.17	2.2e-05	4.9	0.34	1.4e-15
BiClassCNN-Repressed (31.8%)	0.19	0.051	0.00018	0.68	0.048	2.9e-07
BiClassCNN-TSS (1.0%)	0.68	0.13	9.6e-08	6.8	0.6	0.00023

Table S7. S-LDSC results for marginal analysis of non-tissue-specific Roadmap and ChromHMM annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 non-tissue-specific Roadmap and 40 non-tissue-specific ChromHMM annotations. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Roadmap-DNase-Avg (2.2%)	0.39	0.11	0.00034	5.3	0.46	9.7e-13
Roadmap-DNase-Max (13.4%)	0.18	0.075	0.02	3	0.23	3.9e-12
Roadmap-H3K27ac-Avg (3.1%)	0.37	0.084	1.1e-05	4.2	0.27	9.4e-17
Roadmap-H3K27ac-Max (21.4%)	0.16	0.079	0.046	2.5	0.13	6.4e-17
Roadmap-H3K4me1-Avg (4.4%)	0.47	0.077	9.1e-10	3.6	0.19	1.9e-17
Roadmap-H3K4me1-Max (25.4%)	0.094	0.099	0.34	2.2	0.1	8.1e-19
Roadmap-H3K4me3-Avg (1.7%)	0.43	0.088	7.2e-07	5.8	0.41	2.7e-18
Roadmap-H3K4me3-Max (6.6%)	-0.12	0.065	0.075	3.3	0.23	5.7e-16
ChromHMM-DNase-Avg (0.7%)	-0.12	0.049	0.012	1.4	0.39	0.4
ChromHMM-DNase-Max (9.2%)	-0.19	0.061	0.0021	1.5	0.24	0.71
ChromHMM-Active-Enhancer-1-Avg (0.3%)	-0.15	0.03	3.2e-07	2.7	0.3	1.2e-08
ChromHMM-Active-Enhancer-1-Max (7.1%)	-0.17	0.044	8.1e-05	2.3	0.22	0.0025
ChromHMM-Active_Enhancer_2-Avg (0.4%)	-0.073	0.052	0.16	2.9	0.34	0.0011
ChromHMM-Active_Enhancer_2-Max (9.9%)	-0.13	0.057	0.027	2.2	0.2	0.011
ChromHMM-Enhancer_acetylation-Avg (0.3%)	-0.19	0.039	8.5e-07	1.4	0.25	0.79
ChromHMM-Enhancer_acetylation-Max (11.3%)	-0.19	0.029	1.8e-10	1.5	0.13	0.0066
ChromHMM-Active_Enhancer_Flanking-Avg (0.5%)	-0.12	0.035	0.00085	2.3	0.22	0.0092
ChromHMM-Active_Enhancer_Flanking-Max (11.1%)	-0.15	0.046	0.00078	1.9	0.15	0.11
ChromHMM-Weak_Enhancer_1-Avg (0.3%)	-0.14	0.035	7.1e-05	1.8	0.31	9.9e-08
ChromHMM-Weak_Enhancer_1-Max (7.7%)	-0.21	0.034	8.8e-10	1.8	0.19	5.1e-06
ChromHMM-Weak_Enhancer_2-Avg (1.2%)	-0.052	0.036	0.15	2.1	0.14	1.5e-10
ChromHMM-Weak_Enhancer_2-Max (19.9%)	-0.12	0.052	0.02	1.7	0.093	5.8e-11
ChromHMM-Heterochromatin-Avg (1.4%)	-0.074	0.02	0.00027	0.3	0.058	2e-18
ChromHMM-Heterochromatin-Max (7.1%)	-0.055	0.018	0.0022	0.39	0.038	3.5e-21
ChromHMM-Promoter_Bivalent-Avg (0.3%)	0.1	0.042	0.017	6.1	0.48	8.2e-14
ChromHMM-Promoter_Bivalent-Max (1.9%)	0.055	0.041	0.18	4	0.27	2.1e-14
ChromHMM-Promoter_Downstream_1-Avg (0.4%)	0.18	0.062	0.0029	7	0.89	7.8e-12
ChromHMM-Promoter_Downstream_1-Max (1.4%)	0.24	0.063	0.0002	6.5	0.6	8.9e-16
ChromHMM-Promoter_Downstream_2-Avg (0.2%)	-0.014	0.045	0.76	2.5	0.67	0.1
ChromHMM-Promoter_Downstream_2-Max (1.5%)	0.095	0.033	0.004	3.6	0.34	2.2e-14
ChromHMM-Promoter_Poised-Avg (0.2%)	-0.14	0.046	0.0018	1.3	0.25	0.074
ChromHMM-Promoter_Poised-Max (6.8%)	-0.23	0.037	1.1e-09	1.2	0.098	0.0011

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se(E)	p(E)
ChromHMM-Promoter_Upstream-Avg (0.4%)	0.04	0.051	0.43	5.7	0.45	5.3e-16
ChromHMM-Promoter_Upstream-Max (2.5%)	-0.13	0.045	0.0052	3.8	0.34	7.5e-14
ChromHMM-Quiescent-Avg (78.4%)	-0.061	0.018	0.00074	0.73	0.02	3e-18
ChromHMM-Quiescent-Max (91.0%)	-0.086	0.014	1.4e-09	0.86	0.01	6.6e-20
ChromHMM-Repressed_Polycomb-Avg (1.7%)	-0.052	0.016	0.0011	1.5	0.065	1.5e-16
ChromHMM-Repressed_Polycomb-Max (9.9%)	-0.072	0.016	6.5e-06	1.3	0.037	2.9e-20
ChromHMM-TSS-Avg (0.2%)	-0.0011	0.075	0.99	5.1	1	5.1e-05
ChromHMM-TSS-Max (0.9%)	-0.058	0.073	0.42	4.5	0.49	1.7e-10
ChromHMM-Transcription_Enhancer_3'-Avg (0.2%)	0.078	0.027	0.0038	3.4	0.29	1.7e-16
ChromHMM-Transcription_Enhancer_3'-Max (3.1%)	0.088	0.035	0.013	2.8	0.21	6.7e-18
ChromHMM-Transcription Enhancer-5'-Avg (0.4%)	0.12	0.024	9.7e-07	3.6	0.23	5.4e-18
ChromHMM-Transcription_Enhancer_5'-Max (4.8%)	0.16	0.022	2.2e-13	2.9	0.15	6.5e-19
ChromHMM-Transcription_Weak_Enhancer-Avg (0.5%)	0.12	0.024	1.3e-06	2.8	0.16	7.3e-19
ChromHMM-Transcription_Weak_Enhancer-Max (4.7%)	0.16	0.022	3e-12	2.6	0.12	1.7e-21
ChromHMM-Transcriptional_Regulator-Avg (0.3%)	0.27	0.039	5.2e-12	6.8	0.51	4.3e-17
ChromHMM-Transcriptional_Regulator-Max (3.3%)	0.22	0.037	4.2e-09	4	0.26	1.8e-18

Table S8. S-LDSC results for marginal analysis of CpG-island, local CpG-content, local GC-content, pLI, SIFT and Polyphen annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of the weighted Coding, TSS, Repressed and Gene $\pm 5KB$ annotations, where the weights may be driven by the underlying sequence context such as CpG-island, LocalCpG-content ($\pm 1kb$), LocalGC-content ($\pm 1kb$) etc as well as pLI, pLI-Coding, pLI-TSS, PolyPhen and SIFT. All analyses are conditioned on baseline-LD annotations. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
CpGisland (0.8%)	0.2	0.051	6.1e-05	6.7	0.61	5e-14
CpGisland-Coding (0.2%)	0.094	0.049	0.056	7.7	1.2	1.6e-11
CpGisland-Repressed (0.02%)	-0.074	0.023	0.001	3.9	1.4	0.19
CpGisland-TSS (0.4%)	0.23	0.074	0.0019	8.8	0.98	2.2e-11
LocalCpGcontent (5.5%)	-0.19	0.052	0.00025	1.1	0.013	1.5e-15
LocalCpGcontent-Coding (0.2%)	0.44	0.075	4.9e-09	6	0.45	7.6e-16
LocalCpGcontent-Repressed (2.2%)	-0.34	0.058	4.9e-09	0.59	0.055	3.7e-10
LocalCpGcontent-TSS (0.2%)	0.66	0.12	3.8e-08	7.1	0.59	1.1e-17
LocalGCcontent (41.0%)	-0.15	0.03	5.7e-07	1.1	0.0062	1.1e-11
LocalGCcontent-Coding (0.8%)	1.1	0.17	1.9e-10	5	0.35	2e-16
LocalGCcontent-Repressed (18.0%)	-0.68	0.11	5.3e-10	0.63	0.05	1.1e-08
LocalGCcontent-TSS (1.0%)	1.8	0.28	1.7e-10	6.4	0.56	5.7e-19
pLI (14.0%)	0.095	0.011	3.9e-18	1.5	0.034	9.5e-17
pLI-Coding (0.4%)	0.62	0.057	2.5e-27	13	0.78	5.6e-17
pLI-TSS (0.3%)	0.52	0.039	3.5e-41	14	1.1	6.4e-20
Polyphen (0.1%)	0.23	0.081	0.0046	13	2.2	1.8e-10
SIFT (0.1%)	-0.067	0.065	0.3	4.6	1.5	4e-06

Table S9. S-LDSC results for non-tissue-specific variant-level joint model.
Joint S-LDSC model fit of all non-tissue-specific annotations as well as LocalGC-content, LocalCpG-content and CpG-island weighted annotations of TSS, Repressed and Coding regions, and pLI, pLI-TSS, pLI-Coding, SIFT and Polyphen. Only 7 annotations were found to be significant conditioned on all other annotations and the baseline-LD model. Results are meta-analyzed across 41 traits. We also highlight in red annotations from Figure 1 that were not significant under the joint model

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
ChromHMM-Active-Enhancer-1-Avg (0.3%)	-0.16	0.03	6.8e-08	2.6	0.29	1.8e-10
Roadmap-H3K4me1-Avg (4.4%)	0.46	0.079	5e-09	3.6	0.19	6.5e-21
pLI-Coding (0.4%)	0.45	0.054	4e-17	11	0.71	3e-16
pLI-TSS (0.3%)	0.35	0.039	1.4e-19	12	1	8.2e-21
LocalGCcontent (41.0%)	-0.16	0.03	1.1e-07	1.1	0.0066	4.2e-17
LocalGCcontent-TSS (1.0%)	1.2	0.28	1.6e-05	5.8	0.52	5.8e-26
LocalGCcontent-Coding (0.8%)	0.91	0.18	7.8e-07	5.2	0.34	6.7e-20
DeepSEA-H3K4me3-Max (9.4%)	0.13	0.06	0.029	2.2	0.077	6.8e-19
BiClassCNN-Coding (1.3%)	0.43	0.28	0.13	5.1	0.34	6.5e-19
BiClassCNN-Promoter (2.0%)	0.14	0.048	0.0036	2.7	0.17	2.2e-20
BiClassCNN-Repressed (31.8%)	0.1	0.043	0.02	0.66	0.046	1.1e-08
BiClassCNN-TSS (1.0%)	0.19	0.25	0.44	6.2	0.55	3.9e-18

Table S10. The Roadmap-H3K4me1-Avg annotation was sufficient to eliminate the DeepSEA-H3K4me3-Max annotation that was significant in marginal analysis. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of the significant non-tissue-specific variant-level DeepSEA and Roadmap annotations from marginal analysis, when modeled conditioned on each other and the 86 baseline-LD annotations, and passed through Forward Stepwise Elimination. The results for the penultimate step of the elimination process is shown. DeepSEA-H3K4me3-Max is not significant after correcting for all tested hypothesis (0.05/106) and gets eliminated next. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Roadmap-H3K4me1-Avg (4.4%)	0.455	0.08	1e-08	3.63	0.19	3e-19
DeepSEA-H3K4me3-Max (9.4%)	0.182	0.06	0.004	2.14	0.08	1e-16

Table S11. The Local GC-content annotations were sufficient to eliminate the 4 BiClassCNN annotations that were significant in our marginal analysis. Meta-analyzed $\tau^\star$ and Enrichment E of the Joint S-LDSC model fit of the 4 BiClassCNN annotations from Figure 1 conditioned on the 86 baseline-LD annotations and 3 LocalGCcontent annotations from Table S9. Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
BiClassCNN-Coding (1.3%)	0.69	0.3	0.023	5	0.35	1e-16
BiClassCNN-Promoter (2.7%)	0.15	0.052	0.0039	2.7	0.17	3.1e-20
BiClassCNN-Repressed (31.8%)	0.1	0.045	0.022	0.69	0.047	1.9e-07
BiClassCNN-TSS (1.0%)	0.13	0.28	0.65	6.6	0.58	3e-19

Table S12. AUROC of various sets of annotations incorporating non-tissue-specific variant-level deep learning annotations in predicting $12,296$ **NIH GWAS SNPs**. We report the AUROC for a gradient boosting model trained using each respective set of annotations, either without including baseline-LD model annotations; including baseline-LD model annotations; or including annotations from the non-tissue-specific variant-level joint model (baseline-LD model + 7 annotations; baseline-LD+7).

Feature	GWAS SNPs
DeepSEA-Avg/Max	0.582
Basenji-Avg/Max	0.612
DeepSEA-All	0.591
Basenji-All	0.620
baseline-LD	0.758
baseline-LD + DeepSEA-Avg/Max	0.759
baseline-LD + Basenji-Avg/Max	0.766
baseline-LD + DeepSEA-All	0.765
baseline-LD + Basenji-All	0.770
baseline-LD+7	0.762
baseline-LD+7 + DeepSEA-Avg/Max	0.764
baseline-LD+7 + Basenji-Avg/Max	0.765
baseline-LD+7 + DeepSEA-All	0.769
baseline-LD+7 + Basenji-All	0.772

Table S13. Enrichment of NIH GWAS SNPs for non-tissue-specific variant-level annotations. We report the enrichment in deep learning annotations (along with Jackknife standard error) at top 12, 296 known disease-associated SNPs from NIH GWAS catalog[23].

Feature	GWAS SNPs
Basenji-DNase-Avg	1.61 (0.010)
Basenji-DNase-Max	1.35 (0.005)
Basenji-H3K27ac-Avg	1.62 (0.009)
Basenji-H3K27ac-Max	1.42 (0.005)
Basenji-H3K4me1-Avg	1.48 (0.005)
Basenji-H3K4me1-Max	1.31 (0.003)
Basenji-H3K4me3-Avg	1.81 (0.016)
Basenji-H3K4me3-Max	1.30 (0.006)
DeepSEA-DNase-Avg	1.58 (0.009)
DeepSEA-DNase-Max	1.39 (0.004)
DeepSEA-H3K27ac-Avg	1.58 (0.008)
DeepSEA-H3K27ac-Max	1.39 (0.004)
DeepSEA-H3K4me1-Avg	1.46 (0.001)
DeepSEA-H3K4me1-Max	1.31 (0.003)
DeepSEA-H3K4me3-Avg	1.81 (0.013)
DeepSEA-H3K4me3-Max	1.43 (0.005)

Table S14. S-LDSC results for marginal analysis of blood-specific variant-level DeepSEA and Basenji annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 blood-specific variant-level DeepSEA and 8 blood-specific variant-level Basenji annotations, when conditioned on non-tissue-specific variant-level joint model (baseline-LD and annotations from Figure S5). Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Basenji-DNase-blood-Avg (1.9%)	0.06	0.11	0.6	5	0.37	1.2e-12
Basenji-DNase-blood-Max (5.4%)	0.22	0.1	0.028	3.5	0.2	8.8e-06
Basenji-H3K27ac-blood-Avg (2.5%)	0.59	0.13	4.2e-06	4.6	0.25	3.2e-05
Basenji-H3K27ac-blood-Max (8.5%)	0.38	0.13	0.0045	3.2	0.13	4.5e-05
Basenji-H3K4me1-blood-Avg (6.2%)	0.77	0.13	4.4e-09	3.2	0.15	3.7e-05
Basenji-H3K4me1-blood-Max (13.7%)	0.58	0.13	9.8e-06	2.6	0.095	6.4e-05
Basenji-H3K4me3-blood-Avg (2.0%)	0.34	0.13	0.011	6.9	0.41	5.8e-05
Basenji-H3K4me3-blood-Max (5.5%)	0.34	0.098	0.00065	4.2	0.18	6.7e-05
DeepSEA-DNase-blood-Avg (1.9%)	-0.12	0.13	0.35	4.9	0.36	6e-05
DeepSEA-DNAse-blood-Max (5.4%)	0.055	0.11	0.6	3.4	0.18	6.9e-05
DeepSEA-H3K27ac-blood-Avg (2.5%)	0.12	0.11	0.29	4	0.21	9.1e-05
DeepSEA-H3K27ac-blood-Max (8.5%)	0.14	0.11	0.2	2.8	0.1	9e-05
DeepSEA-H3K4me1-blood-Avg (6.2%)	0.14	0.086	0.093	2.8	0.11	0.0001
DeepSEA-H3K4me1-blood-Max (13.7%)	0.14	0.073	0.048	2.4	0.074	0.0001
DeepSEA-H3K4me3-blood-Avg (2.0%)	-0.25	0.13	0.052	5.8	0.42	4.1e-05
DeepSEA-H3K4me3-blood-Max (5.5%)	0.064	0.096	0.5	3.9	0.19	8.3e-05

Table S15. S-LDSC results for marginal analysis of blood-specific Roadmap and ChromHMM annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 blood-specific Roadmap and 40 blood-specific ChromHMM annotations, when conditioned on non-tissue-specific variant-level joint model (baseline-LD and annotations from Figure S5). Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Roadmap-DNase-blood-Avg (2.1%)	1.8	0.29	1.6e-09	14	1.2	7.1e-06
Roadmap-DNase-blood-Max (6.2%)	1.4	0.22	9.8e-10	8.7	0.68	1e-05
Roadmap-H3K27ac-blood-Avg (3.2%)	1.5	0.32	2e-06	9.5	0.77	4.9e-06
Roadmap-H3K27ac-blood-Max (10.2%)	1.3	0.21	1.7e-10	6.4	0.43	9.1e-06
Roadmap-H3K4me1-blood-Avg (4.9%)	1.5	0.33	7.3e-06	7.1	0.54	6.4e-06
Roadmap-H3K4me1-blood-Max (11.2%)	1.4	0.24	5.7e-09	5.8	0.42	2.8e-06
Roadmap-H3K4me3-blood-Avg (1.8%)	1.3	0.3	1.5e-05	13	1.1	1e-05
Roadmap-H3K4me3-blood-Max (3.6%)	1.4	0.23	3.6e-09	11	0.86	2.9e-06
ChromHMM-DNase-blood-Avg (0.4%)	-0.017	0.11	0.88	2.9	0.91	0.0015
ChromHMM-DNase-blood-Max (2.2%)	0.37	0.099	0.00019	5.2	0.59	3e-07
ChromHMM-Active_Enhancer_1-blood-Avg (0.3%)	1	0.25	3.4e-05	12	2.2	2.1e-05
ChromHMM-Active_Enhancer_1-blood-Max (2.6%)	1	0.17	7.7e-09	9.3	1.1	1.7e-07
ChromHMM-Active_Enhancer_2-blood-Avg (0.4%)	1.1	0.26	2.1e-05	12	1.8	6.4e-07
ChromHMM-Active_Enhancer_2-blood-Max (3.5%)	1.2	0.22	3.3e-08	9	1.1	3.3e-08
ChromHMM-Enhancer_acetylation-blood-Avg (0.2%)	0.24	0.13	0.059	5	0.9	4.9e-06
ChromHMM-Enhancer_acetylation-blood-Max (2.8%)	0.54	0.14	8.3e-05	5.6	0.72	9.5e-06
ChromHMM-Active_Enhancer_Flanking-blood-Avg (0.4%)	0.96	0.24	4.6e-05	10	1.6	1.9e-06
ChromHMM-Active_Enhancer_Flanking-blood-Max (3.3%)	1.1	0.22	1.1e-06	8.2	1.1	2.3e-07
ChromHMM-Weak_Enhancer_1-blood-Avg (0.3%)	0.43	0.11	5.3e-05	8.8	1	7.4e-08
ChromHMM-Weak_Enhancer_1-blood-Max (2.2%)	0.78	0.14	1.4e-08	8.6	0.89	9.7e-06
ChromHMM-Weak_Enhancer_2-blood-Avg (1.1%)	0.81	0.19	1.3e-05	6.6	0.88	9e-07
ChromHMM-Weak_Enhancer_2-blood-Max (6.5%)	1	0.19	4.1e-08	5.8	0.62	1.6e-07
ChromHMM-Heterochromatin-blood-Avg (1.2%)	-0.13	0.031	3.2e-05	0.12	0.16	0.00025
ChromHMM-Heterochromatin-blood-Max (3.2%)	-0.097	0.031	0.002	0.01	0.13	0.00057
ChromHMM-Promoter_Bivalent-blood-Avg (0.3%)	-0.34	0.12	0.0053	2.1	1.5	0.08
ChromHMM-Promoter_Bivalent-blood-Max (0.8%)	-0.078	0.13	0.54	4.7	0.96	1.3e-05
ChromHMM-Promoter_Downstream_1-blood-Avg (0.4%)	0.66	0.18	0.00034	19	2.1	1.4e-05

Table S15. *(continued)*

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se(E)	p(E)
ChromHMM-Promoter_Downstream_1-blood-Max (0.9%)	1.1	0.18	3.2e-10	19	1.5	8.7e-06
ChromHMM-Promoter_Downstream_2-blood-Avg (0.1%)	0.51	0.13	0.0001	14	2.2	7.7e-08
ChromHMM-Promoter_Downstream_2-blood-Max (0.7%)	0.65	0.11	1.4e-08	12	1.5	3.8e-11
ChromHMM-Promoter_Poised-blood-Avg (0.2%)	-0.12	0.12	0.32	2.8	1.4	0.0029
ChromHMM-Promoter_Poised-blood-Max (1.6%)	0.065	0.075	0.38	3.6	0.53	2.5e-05
ChromHMM-Promoter_Upstream-blood-Avg (0.5%)	-0.15	0.12	0.22	8.7	1.2	6.9e-08
ChromHMM-Promoter_Upstream-blood-Max (1.3%)	0.59	0.13	1e-05	12	0.86	2.1e-05
ChromHMM-Quiescent-blood-Avg (80.0%)	-0.21	0.082	0.012	0.52	0.022	4.9e-05
ChromHMM-Quiescent-blood-Max (86.0%)	-0.092	0.049	0.06	0.64	0.017	6e-05
ChromHMM-Repressed_Polycomb-blood-Avg (1.4%)	-0.17	0.047	0.00037	1.1	0.36	0.089
ChromHMM-Repressed_Polycomb-blood-Max (0.04%)	-0.12	0.04	0.0038	1.3	0.22	0.0036
ChromHMM-TSS-blood-Avg (0.1%)	-0.35	0.15	0.022	8.9	3.9	0.0041
ChromHMM-TSS-blood-Max (0.3%)	-0.34	0.12	0.0051	7.5	2	0.00049
ChromHMM-Transcription_Enhancer_3'-blood-Avg (0.2%)	0.29	0.11	0.0067	7.7	1.4	1.4e-06
ChromHMM-Transcription_Enhancer_3'-blood-Max (1.1%)	0.45	0.11	2.9e-05	7.6	0.95	4.2e-06
ChromHMM-Transcription_Enhancer_5'-blood-Avg (0.5%)	0.28	0.11	0.0085	6.2	0.96	2.6e-10
ChromHMM-Transcription_Enhancer_5'-blood-Max (2.5%)	0.45	0.11	5.4e-05	5.8	0.68	1.4e-06
ChromHMM-Transcription_Weak_Enhancer-blood-Avg (0.6%)	-0.035	0.1	0.72	3.1	0.87	6.9e-07
ChromHMM-Transcription_Weak_Enhancer-blood-Max (2.6%)	0.19	0.09	0.035	3.9	0.64	6.8e-10
ChromHMM-Transcriptional_Regulator-blood-Avg (0.4%)	0.76	0.11	2.8e-12	14	1.3	1.3e-10
ChromHMM-Transcriptional_Regulator-blood-Max (1.7%)	0.74	0.16	2.6e-06	9.6	1.1	2.4e-11

Table S16. S-LDSC results for blood-specific variant-level joint model.
Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of the significant blood-specific variant-level DeepSEA, Basenji, Roadmap and ChromHMM annotations from Table S14 and Table S15, conditioned on each other and the non-tissue-specific variant-level joint model (baseline-LD and annotations from Figure S5). Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
ChromHMM-Active_Enhancer_2-blood-Max (3.5%)	0.73	0.18	6.9e-05	7.5	0.89	1.7e-06
ChromHMM-Promoter_Downstream_1-blood-Max (1.1%)	1	0.21	1.4e-06	15	1.5	3.5e-05
ChromHMM-Transcriptional_Regulator-blood-Avg (0.4%)	0.84	0.13	1.6e-11	12	1.1	7e-09
ChromHMM-Repressed_Polycomb-blood-Avg (1.4%)	-0.11	0.032	5.8e-04	1.5	0.35	0.0026
Roadmap-H3K4me1-blood-Max (11.2%)	0.68	0.18	0.00023	5.4	0.39	6.9e-06
Roadmap-H3K4me3-blood-Avg (1.8%)	-0.77	0.19	7.4e-05	9.4	0.71	3.8e-05

Table S17. The 2 surviving blood-specific Roadmap annotations were sufficient to eliminate 3 of the 4 blood-specific variant-level Basenji annotations that were significant in marginal analysis. Joint S-LDSC model fit of the blood-specific variant-level Basenji annotations from Table S14 conditioned on the non-tissue-specific variant-level joint model plus the two Roadmap blood annotations from Table S16. Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Basenji-H3K27ac-blood-Avg (2.5%)	0.16	0.12	0.17	4.4	0.23	2e-05
Basenji-H3K4me1-blood-Avg (6.2%)	0.34	0.094	0.00031	3.1	0.12	2.5e-05
Basenji-H3K4me1-blood-Max (13.7%)	0.18	0.11	0.096	2.5	0.086	3.9e-05
Basenji-H3K4me3-blood-Max (5.5%)	-0.052	0.1	0.61	4.1	0.19	3.1e-05

Table S18. AUROC of various sets of annotations incorporating blood-specific variant-level deep learning annotations in predicting 8,741 fine-mapped autoimmune diseease SNPs. We report the AUROC for a gradient boosting model trained using each respective set of annotations, either without including baseline-LD model annotations; including baseline-LD model annotations; including annotations from the non-tissue-specific variant-level joint model (baseline-LD model + 7 annotations; baseline-LD+7); or including annotations from the blood-specific joint model (non-tissue-specific joint variant-level joint model + 6 blood annotations; baseline-LD+7+6Blood).

Feature	finemapped SNPs
DeepSEA-blood-Avg/Max	0.621
Basenji-blood-Avg/Max	0.664
DeepSEA-blood-All	0.631
Basenji-blood-All	0.671
baseline-LD	0.841
baseline-LD + DeepSEA-blood-Avg/Max	0.846
baseline-LD + Basenji-blood-Avg/Max	0.852
baseline-LD + DeepSEA-blood-All	0.849
baseline-LD + Basenji-blood-All	0.854
baseline-LD+7	0.845
baseline-LD+7 + DeepSEA-blood-Avg/Max	0.849
baseline-LD+7 + Basenji-blood-Avg/Max	0.853
baseline-LD+7 + DeepSEA-blood-All	0.848
baseline-LD+7 + Basenji-blood-All	0.853
baseline-LD+7+6Blood	0.851
baseline-LD+7+6Blood + DeepSEA-blood-Avg/Max	0.853
baseline-LD+7+6Blood + Basenji-blood-Avg/Max	0.854
baseline-LD+7+6Blood + DeepSEA-blood-All	0.853
baseline-LD+7+6Blood + Basenji-blood-All	0.853

Table S19. Enrichment of fine-mapped autoimmune disease SNPs for blood-specific variant-level annotations. We report the enrichment in blood-specific variant-level deep learning annotations (along with Jackknife standard error) for 8741 fine-mapped SNPs in immune-related traits[24].

Feature	finemapped SNPs
Basenji-DNase-blood-Avg	2.02 (0.03)
Basenji-DNase-blood-Max	1.80 (0.03)
Basenji-H3K27ac-blood-Avg	2.14 (0.02)
Basenji-H3K27ac-blood-Max	1.85 (0.02)
Basenji-H3K4me1-blood-Avg	2.04 (0.03)
Basenji-H3K4me1-blood-Max	1.81 (0.02)
Basenji-H3K4me3-blood-Avg	2.17 (0.02)
Basenji-H3K4me3-blood-Max	1.95 (0.03)
DeepSEA-DNase-blood-Avg	1.88 (0.03)
DeepSEA-DNase-blood-Max	1.68 (0.02)
DeepSEA-H3K27ac-blood-Avg	1.81 (0.02)
DeepSEA-H3K27ac-blood-Max	1.67 (0.02)
DeepSEA-H3K4me1-blood-Avg	1.70 (0.01)
DeepSEA-H3K4me1-blood-Max	1.55 (0.02)
DeepSEA-H3K4me3-blood-Avg	1.95 (0.02)
DeepSEA-H3K4me3-blood-Max	1.76 (0.02)

Table S20. S-LDSC results for marginal analysis of brain-specific variant-level DeepSEA and Basenji annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 brain-specific variant-level DeepSEA and 8 brain-specific variant-level Basenji annotations corresponding to 4 chromatin marks, DNase-1 Hypersensitivity Sites (DHS), H3K27ac, H3K4me1 and H3K4me3, when conditioned on the non-tissue-specific variant-level joint model (baseline-LD and annotations from Figure S5). Results are meta-analyzed across 8 brain-related traits..

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEA-DNase-brain-Avg (2.9%)	0.5	0.081	8.6e-10	2.9	0.23	5e-05
DeepSEA-DNase-brain-Max (4.5%)	0.41	0.067	1.7e-09	2.5	0.19	3.4e-05
DeepSEA-H3K27ac-brain-Avg (3.2%)	0.6	0.066	5.4e-20	2.3	0.072	1.1e-08
DeepSEA-H3K27ac-brain-Max (7.6%)	0.55	0.062	1.5e-18	2.1	0.076	8.8e-09
DeepSEA-H3K4me1-brain-Avg (4.7%)	0.37	0.06	5.8e-10	2.1	0.06	3.7e-11
DeepSEA-H3K4me1-brain-Max (10.5%)	0.36	0.053	5.3e-12	1.9	0.07	2.6e-11
DeepSEA-H3K4me3-brain-Avg (1.9%)	0.77	0.095	4.6e-16	4.4	0.26	1.7e-08
DeepSEA-H3K4me3-brain-Max (3.4%)	0.77	0.085	1e-19	3.8	0.19	2.2e-08
Basenji-DNase-brain-Avg (2.9%)	0.47	0.078	1.3e-09	2.7	0.16	8.8e-08
Basenji-DNase-brain-Max (4.5%)	0.49	0.076	9.3e-11	2.8	0.18	1.2e-07
Basenji-H3K27ac-brain-Avg (3.2%)	0.34	0.049	3.1e-12	2.3	0.079	1.6e-08
Basenji-H3K27ac-brain-Max (7.6%)	0.4	0.058	3.6e-12	2.3	0.12	1.7e-06
Basenji-H3K4me1-brain-Avg (4.7%)	0.37	0.059	4.1e-10	2.1	0.059	7.6e-11
Basenji-H3K4me1-brain-Max (10.5%)	0.33	0.09	0.00022	2	0.062	2.6e-10
Basenji-H3K4me3-brain-Avg (1.9%)	0.55	0.074	1.1e-13	3.6	0.18	5.2e-07
Basenji-H3K4me3-brain-Max (3.4%)	0.77	0.089	8.2e-18	4	0.3	8.8e-06

Table S21. S-LDSC for marginal analysis of brain-specific Roadmap and ChromHMM annotations. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 brain-specific Roadmap and 40 brain-specific ChromHMM annotations, when conditioned on the non-tissue-specific variant-level joint model annotations (baseline-LD and annotations from Figure S5). Results are meta-analyzed across 8 brain-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Roadmap-DNase-brain-Avg (2.4%)	0.55	0.084	5.5e-11	4.9	0.41	5.4e-06
Roadmap-DNase-brain-Max (5.1%)	0.39	0.079	7.1e-07	3.5	0.21	2.4e-06
Roadmap-H3K27ac-brain-Avg (3.2%)	0.34	0.09	0.00019	3.5	0.15	1.1e-10
Roadmap-H3K27ac-brain-Max (8.2%)	0.37	0.095	0.00012	2.9	0.11	7.6e-11
Roadmap-H3K4me1-brain-Avg (4.4%)	0.48	0.12	5e-05	3.2	0.12	4.4e-11
Roadmap-H3K4me1-brain-Max (11.2%)	0.46	0.12	0.00016	2.7	0.09	3.6e-11
Roadmap-H3K4me3-brain-Avg (1.5%)	0.86	0.15	7.7e-09	6.8	0.37	1.1e-12
Roadmap-H3K4me3-brain-Max (2.9%)	0.72	0.12	1.8e-09	5.7	0.28	4e-12
ChromHMM-DNase-brain-Avg (0.8%)	-0.12	0.096	0.21	0.86	0.8	0.45
ChromHMM-DNase-brain-Max (3.0%)	0.062	0.072	0.39	1.8	0.5	0.0031
ChromHMM-Active-Enhancer-1-brain-Avg (0.4%)	0.24	0.086	0.0061	3.6	0.5	0.06
ChromHMM-Active-Enhancer-1-brain-Max (1.5%)	0.32	0.1	0.002	3.5	0.45	0.056
ChromHMM-Active_Enhancer_2-brain-Avg (0.4%)	0.39	0.083	3.4e-06	4.8	0.48	0.00043
ChromHMM-Active_Enhancer_2-brain-Max (2.2%)	0.36	0.082	8e-06	4	0.3	1e-05
ChromHMM-Enhancer_acetylation-brain-Avg (0.3%)	0.13	0.054	0.02	2.3	0.29	0.077
ChromHMM-Enhancer_acetylation-brain-Max (2.2%)	0.19	0.068	0.0047	2.5	0.29	0.016
ChromHMM-Active_Enhancer_Flanking-brain-Avg (0.8%)	0.13	0.057	0.024	2.5	0.21	0.0031
ChromHMM-Active_Enhancer_Flanking-brain-Max (3.1%)	0.16	0.062	0.011	2.3	0.17	0.002

Table S21. *(continued)*

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se(E)	p(E)
ChromHMM-Weak_Enhancer_1-brain-Avg (0.4%)	0.19	0.073	0.0087	3.9	0.56	0.03
ChromHMM-Weak_Enhancer_1-brain-Max (1.7%)	0.29	0.091	0.0012	4.1	0.51	0.014
ChromHMM-Weak_Enhancer_2-brain-Avg (1.3%)	0.25	0.074	0.00065	2.8	0.27	3.5e-10
ChromHMM-Weak_Enhancer_2-brain-Max (5.6%)	0.27	0.085	0.0015	2.5	0.23	3.6e-10
ChromHMM-Heterochromatin-brain-Avg (1.2%)	-0.00042	0.021	0.98	0.37	0.12	0.016
ChromHMM-Heterochromatin-brain-Max (2.3%)	0.0051	0.021	0.81	0.42	0.1	0.0058
ChromHMM-Promoter_Bivalent-brain-Avg (0.2%)	0.15	0.06	0.01	7.4	0.97	0.00052
ChromHMM-Promoter_Bivalent-brain-Max (0.6%)	0.29	0.058	3e-07	7.2	0.57	2.7e-05
ChromHMM-Promoter_Downstream_1-brain-Avg (0.4%)	0.29	0.072	7e-05	7.7	0.9	1.5e-05
ChromHMM-Promoter_Downstream_1-brain-Max (0.6%)	0.47	0.077	1.2e-09	9.1	0.83	1.9e-08
ChromHMM-Promoter_Downstream_2-brain-Avg (0.2%)	0.12	0.069	0.075	4.4	1.2	0.095
ChromHMM-Promoter_Downstream_2-brain-Max (0.4%)	0.22	0.068	0.0011	5.5	0.98	0.00026
ChromHMM-Promoter_Poised-brain-Avg (0.2%)	0.23	0.083	0.0064	4.3	0.81	0.059
ChromHMM-Promoter_Poised-brain-Max (1.1%)	0.24	0.1	0.018	3.7	0.75	0.11
ChromHMMM-Promoter_Upstream-brain-Avg (0.5%)	0.46	0.11	2.7e-05	9.6	0.82	1e-10
ChromHMM-Promoter_Upstream-brain-Max (1.0%)	0.5	0.11	5.9e-06	8.1	0.6	1.3e-11

Table S21. *(continued)*

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se(E)	p(E)
ChromHMM-Quiescent-brain-Avg (77.6%)	-0.22	0.054	5.1e-05	0.81	0.013	8.1e-07
ChromHMM-Quiescent-brain-Max (86.0%)	-0.088	0.029	0.0025	0.87	0.0098	3.4e-06
ChromHMM-Repressed_Polycomb-brain-Avg (1.4%)	-0.069	0.025	0.005	1	0.16	0.31
ChromHMM-Repressed_Polycomb-brain-Max (4.3%)	-0.046	0.024	0.054	1.2	0.094	0.012
ChromHMM-TSS-brain-Avg (0.2%)	0.32	0.092	0.00048	7.9	1.6	0.00061
ChromHMM-TSS-brain-Max (0.4%)	0.44	0.097	5e-06	9	1.3	0.0017
ChromHMM-Transcription_Enhancer_3'-brain-Avg (0.3%)	0.12	0.038	0.0027	3.6	0.39	0.039
ChromHMM-Transcription_Enhancer_3'-brain-Max (1.0%)	0.092	0.039	0.017	2.9	0.31	0.0022
ChromHMM-Transcription_Enhancer_5'-brain-Avg (0.5%)	0.027	0.032	0.4	2.8	0.24	1.7e-08
ChromHMM-Transcription_Enhancer_5'-brain-Max (1.6%)	0.083	0.034	0.016	2.8	0.2	5.4e-10
ChromHMM-Transcription_Weak_Enhancer-brain-Avg (0.6%)	0.21	0.049	2.8e-05	3.3	0.33	0.00026
ChromHMM-Transcription_Weak_Enhancer-brain-Max (1.8%)	0.22	0.05	1.7e-05	3	0.25	2.2e-11
ChromHMM-Transcriptional_Regulator-brain-Avg (0.3%)	0.11	0.045	0.019	5	0.52	1.8e-06
ChromHMM-Transcriptional_Regulator-brain-Max (0.9%)	0.056	0.047	0.24	3.7	0.35	3.3e-06

Table S22. S-LDSC results for brain-specific variant-level joint model.
Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of the significant brain-specific
variant-level DeepSEA, Basenji, Roadmap and ChromHMM annotations from Table
S20 and Table S21, conditioned on each other and the non-tissue-specific variant-level
joint model annotations (baseline-LD and annotations from Figure S5). Results are
meta-analyzed across 8 brain-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Basenji-H3K27ac-brain-Max (7.7%)	0.27	0.059	5.4e-06	2.3	0.077	1.3e-08
ChromHMM-Promoter_Bivalent-brain-Max (0.6%)	0.21	0.056	0.00014	6.9	0.56	3.3e-06
ChromHMM-Transcriptional-Weak-Enhancer-brain-Max (1.8%)	0.26	0.06	1.2e-05	3	0.25	4.8e-16
ChromHMM-Weak_Enhancer_2-brain-Avg (1.3%)	0.27	0.078	0.00067	2.7	0.25	3.4e-12
DeepSEA-H3K4me3-brain-Max (3.4%)	0.35	0.09	0.0001	3.6	0.2	5.8e-08
Roadmap-H3K4me3-brain-Max (2.9%)	0.4	0.091	1.3e-05	5.2	0.28	1.5e-10

Table S23. The surviving brain-specific Roadmap annotations do not eliminate any brain specific variant-level DeepSEA and brain-specific variant-level Basenji annotations that were significant in marginal analysis. Standardized Effect sizes ($\tau^{\star}$) and Enrichment (E) of brain-specific DeepSEA and brain-specific Basenji annotations, conditioned on the non-tissue-specific variant-level joint model annotations and 1 significant Roadmap brain annotation (Roadmap-H3K4me3-brain-Max) that is significant in the joint model in Table S22. Results are meta-analyzed across 8 brain-related traits.

Annotation	$\tau^{\star}$	se($\tau^{\star}$)	p($\tau^{\star}$)	E	se (E)	p(E)
DeepSEA-DNase-brain-Avg (2.9%)	0.31	0.081	0.00011	2.8	0.24	3.8e-08
DeepSEA-DNase-brain-Max (4.5%)	0.25	0.069	0.00022	2.5	0.2	2.3e-08
DeepSEA-H3K27ac-brain-Avg (3.2%)	0.47	0.068	2.4e-12	2.4	0.072	9e-12
DeepSEA-H3K27ac-brain-Max (7.7%)	0.43	0.064	1.7e-11	2.2	0.072	5.6e-12
DeepSEA-H3K4me1-brain-Avg (4.7%)	0.3	0.054	1.6e-08	2.1	0.057	4.1e-13
DeepSEA-H3K4me1-brain-Max (10.5%)	0.29	0.053	4.9e-08	1.9	0.068	5e-13
DeepSEA-H3K4me3-brain-Avg (1.9%)	0.5	0.092	5.7e-08	4.4	0.26	1.3e-11
DeepSEA-H3K4me3-brain-Max (3.4%)	0.54	0.087	6.4e-10	3.8	0.19	6.7e-12
Basenji-DNase-brain-Avg (2.9%)	0.34	0.064	8.7e-08	2.8	0.17	1.4e-10
Basenji-DNase-brain-Max (4.5%)	0.36	0.068	1.4e-07	2.7	0.15	1.5e-10
Basenji-H3K27ac-brain-Avg (3.2%)	0.25	0.055	4e-06	2.4	0.11	1.3e-10
Basenji-H3K27ac-brain-Max (7.7%)	0.34	0.056	1.1e-09	2.3	0.073	1.3e-10
Basenji-H3K4me1-brain-Avg (4.7%)	0.3	0.053	2.2e-08	2.1	0.059	7e-12
Basenji-H3K4me1-brain-Max (10.5%)	0.29	0.087	0.00077	2.1	0.061	1.6e-12
Basenji-H3K4me3-brain-Avg (1.9%)	0.31	0.081	0.00014	4.1	0.28	4.8e-11
Basenji-H3K4me3-brain-Max (3.4%)	0.61	0.08	2.3e-14	3.8	0.17	3.5e-11

Table S24. The surviving brain-specific ChromHMM annotations do not eliminate any brain specific variant-level DeepSEA and brain-specific variant-level Basenji annotations that were significant in marginal analysis. Standardized Effect sizes ($\tau^{\star}$) and Enrichment (E) of brain-specific variant-level DeepSEA and brain-specific variant-level Basenji annotations, conditioned on the non-tissue-specific variant-level joint model annotations and 3 significant ChromHMM brain annotations that are significant in the joint model in Table S22. Results are meta-analyzed across 8 brain-related traits.

Annotation	$\tau^{\star}$	se($\tau^{\star}$)	p($\tau^{\star}$)	E	se (E)	p(E)
DeepSEA-DNase-brain-Avg (2.9%)	0.44	0.079	2e-08	2.8	0.23	3.5e-06
DeepSEA-DNase-brain-Max (4.5%)	0.36	0.067	5.9e-08	2.5	0.2	1.3e-06
DeepSEA-H3K27ac-brain-Avg (3.2%)	0.55	0.065	2.6e-17	2.2	0.085	7.1e-08
DeepSEA-H3K27ac-brain-Max (7.6%)	0.5	0.062	5.3e-16	2.1	0.086	5.4e-08
DeepSEA-H3K4me1-brain-Avg (4.7%)	0.28	0.053	9.9e-08	1.9	0.07	4.4e-08
DeepSEA-H3K4me1-brain-Max (10.5%)	0.28	0.052	5.5e-08	1.8	0.08	2.3e-08
DeepSEA-H3K4me3-brain-Avg (1.9%)	0.73	0.094	8.4e-15	4.2	0.26	5.4e-08
DeepSEA-H3K4me3-brain-Max (3.4%)	0.72	0.086	3.9e-17	3.6	0.2	6.3e-08
Basenji-DNase-brain-Avg (2.9%)	0.4	0.071	1.7e-08	2.7	0.18	7.2e-08
Basenji-DNase-brain-Max (4.5%)	0.42	0.073	5.4e-09	2.7	0.16	5.1e-08
Basenji-H3K27ac-brain-Avg (3.2%)	0.30	0.05	1.6e-09	2.2	0.13	1.1e-07
Basenji-H3K27ac-brain-Max (7.6%)	0.36	0.057	2.3e-10	2.2	0.084	5.7e-08
Basenji-H3K4me1-brain-Avg (4.7%)	0.31	0.057	6.1e-08	2	0.072	6.9e-08
Basenji-H3K4me1-brain-Max (10.5%)	0.30	0.089	0.00061	2	0.054	4e-09
Basenji-H3K4me3-brain-Avg (1.9%)	0.52	0.073	1e-12	3.8	0.31	2.4e-07
Basenji-H3K4me3-brain-Max (3.4%)	0.68	0.096	1.5e-12	3.5	0.2	1.4e-07

Table S25. Weighted k-mer enrichments for significant variant level brain-specific variant-level DeepSEA and Basenji annotations. We report the weighted k-mer enrichment and enrichment p-value for the top significant enriched k-mers ($1 \leq k \leq 5$) for H3K4me3-DeepSEA-brain-Max and H3K27ac-Basenji-brain-Max which were jointly significant in the brain-specific variant-level joint model. The Bonferonni corrcection threshold is $0.05/(4*682)$ where 682 is the number of k-mers analyzed for the 2 significant variant-level annotations analyzed here and the 2 significant allelic-effect annotations analyzed in Table S29.

H3K4me3-DeepSEA-brain-Max		
kmer	enrichment	pvalue
CGCGC	8.8	9.6e-55
CGGCG	8.7	2.4e-53
CGCCG	8.5	1.2e-50
CGCG	5.9	5.1e-23
CGCGG	5.7	2.1e-21
CCGCG	5.7	5.0e-21
CGCGA	5.0	7.5e-16
AGCGC	4.8	1.0e-14
CCCGC	4.4	3.9e-12
CGCCC	4.3	1.1e-11
CCGCC	4.1	2.1e-10
ACGCG	4.1	2.2e-10
CCGGC	4.0	1.2e-09
CGGC	4.0	1.4e-09
CGGCC	4.0	1.4e-09
CGCGT	3.9	2.4e-09
CCGC	3.8	6.3e-09
CGCC	3.8	7.4e-09
CGGGC	3.8	9.7e-09
CGAGC	3.7	2.6e-08
CGCAG	3.6	6.2e-08
CGC	3.6	9.3e-08
CTGCG	3.6	1.5e-07
CGCTC	3.4	6.2e-07
CTCGC	3.4	1.1e-06

H3K27ac-Basenji-brain-Max		
kmer	enrichment	pvalue
CGGCG	2.8	3.3e-72
CGCGC	2.8	6.4e-70
CGCCG	2.7	1.1e-66
CGCG	2.1	6.7e-27
CGCGG	2	7.4e-22
CCGCG	1.9	2.2e-21
CGCGA	1.9	9.8e-20
AGCGC	1.9	1.2e-18
ACGCG	1.7	2.1e-12
CCCGC	1.7	8.9e-12
CGCCC	1.7	2.3e-11
CGCGT	1.7	3.1e-11
CGAGC	1.6	1.1e-10
CGGC	1.6	2.8e-10
CGCAG	1.6	3.8e-10
CTGCG	1.6	4.4e-10
CCGCC	1.6	1.5e-09
CCGGC	1.6	4.3e-09
CGGCC	1.6	4.7e-09
CCGC	1.6	1.5e-08
CGCC	1.6	1.7e-08
CGC	1.5	2.3e-08
CCGCT	1.5	3.5e-08
CGCTC	1.5	3.4e-08
AGCGG	1.5	4.7e-08

Table S26. S-LDSC results for marginal analysis of non-tissue-specific allelic-effect DeepSEA and Basenji annotations conditioned on non-tissue-specific variant-level joint model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 non-tissue-specific DeepSEAΔ and 8 non-tissue-specific BasenjiΔ annotations, when conditioned on the non-tissue-specific variant-level joint model (Figure S5). Results are meta-analyzed across 41 traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-Avg (0.3%)	0.07	0.07	0.32	2.3	0.21	3.3e-07
DeepSEAΔ-DNase-Max (2.0%)	-0.034	0.066	0.61	1.6	0.1	1.5e-05
DeepSEAΔ-H3K27ac-Avg (0.2%)	0.056	0.075	0.45	2	0.12	9.7e-10
DeepSEAΔ-H3K27ac-Max (0.9%)	0.02	0.064	0.75	1.7	0.095	3.3e-10
DeepSEAΔ-H3K4me1-Avg (0.3%)	0.014	0.057	0.81	1.7	0.066	1.9e-12
DeepSEAΔ-H3K4me1-Max (1.7%)	-0.051	0.06	0.4	1.4	0.061	2.7e-07
DeepSEAΔ-H3K4me3-Avg (0.1%)	0.078	0.07	0.27	2.6	0.16	0.00042
DeepSEAΔ-H3K4me3-Max (0.7%)	0.069	0.071	0.33	2	0.087	2.6e-13
BasenjiΔ-DNase-Avg (0.3%)	-0.001	0.13	0.99	2.7	0.36	0.00038
BasenjiΔ-DNase-Max (2.0%)	0.14	0.075	0.071	2.3	0.17	1.1e-10
BasenjiΔ-H3K27ac-Avg (0.3%)	0.19	0.11	0.075	2.4	0.18	6.3e-11
BasenjiΔ-H3K27ac-Max (0.9%)	0.078	0.036	0.032	2.1	0.11	1.3e-18
BasenjiΔ-H3K4me1-Avg (0.3%)	0.14	0.063	0.027	2	0.1	5.6e-16
BasenjiΔ-H3K4me1-Max (1.7%)	0.13	0.05	0.0093	1.9	0.091	4.2e-16
BasenjiΔ-H3K4me3-Avg (0.1%)	0.16	0.071	0.023	3.2	0.19	6.8e-14
BasenjiΔ-H3K4me3-Max (0.7%)	0.32	0.053	7.7e-10	2.6	0.13	3.3e-18

Table S27. "S-LDSC results for marginal analysis of non-tissue-specific allelic-effect DeepSEA and Basenji annotations conditioned on baseline-LD model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 non-tissue-specific DeepSEAΔ and 8 non-tissue-specific BasenjiΔ annotations, when conditioned on the baseline-LD model. Results are meta-analyzed across 41 independent traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-Avg (0.3%)	-0.059	0.08	0.46	2	0.22	2.9e-05
DeepSEAΔ-DNase-Max (2.0%)	-0.24	0.079	0.0019	1.3	0.11	7.7e-05
DeepSEAΔ-H3K27ac-Avg (0.2%)	0.046	0.073	0.52	2	0.12	6e-13
DeepSEAΔ-H3K27ac-Max (0.9%)	-0.05	0.068	0.47	1.7	0.093	2.7e-11
DeepSEAΔ-H3K4me1-Avg (0.3%)	-0.054	0.061	0.38	1.6	0.062	3.6e-14
DeepSEAΔ-H3K4me1-Max (1.7%)	-0.21	0.068	0.002	1.3	0.059	4.2e-06
DeepSEAΔ-H3K4me3-Avg (0.1%)	0.19	0.069	0.006	2.9	0.18	1.7e-11
DeepSEAΔ-H3K4me3-Max (0.7%)	0.074	0.076	0.34	2.1	0.095	2.1e-16
BasenjiΔ-DNase-Avg (0.3%)	-0.037	0.12	0.77	2.6	0.36	3.9e-05
BasenjiΔ-DNase-Max (2.0%)	0.032	0.07	0.65	2.2	0.17	1.3e-06
BasenjiΔ-H3K27ac-Avg (0.2%)	0.21	0.1	0.042	2.4	0.18	7e-08
BasenjiΔ-H3K27ac-Max (0.9%)	0.011	0.036	0.76	2.1	0.11	5.2e-14
BasenjiΔ-H3K4me1-Avg (0.3%)	0.089	0.056	0.11	1.9	0.097	1.6e-
BasenjiΔ-H3K4me1-Max (1.7%)	0.034	0.046	0.45	1.9	0.089	1.6e-12
BasenjiΔ-H3K4me3-Avg (0.1%)	0.22	0.075	0.0028	3.2	0.21	4.6e-13
BasenjiΔ-H3K4me3-Max (0.7%)	0.33	0.058	1.1e-08	2.6	0.14	6.3e-16

Table S28. S-LDSC results for marginal analysis of non-tissue-specific variant-level Basenji annotations conditioned on the non-tissue-specific variant-level joint model and the non-tissue-specific final joint model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 non-tissue-specific Basenji variant level annotations, when conditioned on (Top) non-tissue-specific variant-level joint model (FigureS5) and (Bottom) non-tissue-specific final joint model comprising of the non-tissue-specific variant-level joint model plus 1 significant allelic-effect annotation (BasenjiΔ-H34me3-Max) from Figure 4. Results are meta-analyzed across 41 independent traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Conditional on non-tissue-specific variant-level joint model						
Basenji-DNase-Avg (1.6%)	-0.085	0.064	0.18	2.5	0.19	1.4e-13
Basenji-DNase-Max (13.2%)	-0.052	0.06	0.38	1.8	0.075	2.3e-19
Basenji-H3K27ac-Avg (3.2%)	0.047	0.055	0.39	2.5	0.14	1.2e-17
Basenji-H3K27ac-Max (13.7%)	-0.018	0.03	0.55	1.8	0.073	1e-17
Basenji-H3K4me1-Avg (5.5%)	0.041	0.04	0.31	2	0.075	1.6e-18
Basenji-H3K4me1-Max (23.2%)	-0.046	0.025	0.067	1.6	0.046	1.6e-19
Basenji-H3K4me3-Avg (1.7%)	0.017	0.058	0.77	3.7	0.24	5.1e-17
Basenji-H3K4me3-Max (8.8%)	0.029	0.033	0.38	2	0.083	1.6e-17
Conditional on non-tissue-specific final joint model						
Basenji-DNase-Avg (1.6%)	-0.11	0.065	0.08	1.8	0.074	1.4e-38
Basenji-DNase-Max (13.2%)	-0.11	0.065	0.099	2.5	0.19	8.6e-32
Basenji-H3K27ac-Avg (3.2%)	0.0051	0.055	0.93	2.5	0.13	1.3e-33
Basenji-H3K27ac-Max (13.7%)	-0.075	0.03	0.012	1.8	0.072	1.5e-31
Basenji-H3K4me1-Avg (5.5%)	-0.011	0.04	0.79	1.9	0.073	7.9e-32
Basenji-H3K4me1-Max (23.2%)	-0.11	0.036	0.76	1.6	0.046	3.5e-33
Basenji-H3K4me3-Avg (1.7%)	-0.031	0.061	0.61	3.5	0.23	5.3e-34
Basenji-H3K4me3-Max (8.8%)	-0.026	0.036	0.47	1.9	0.081	3.4e-32

Table S29. Weighted k-mer enrichments for significant allelic-effect Basenji annotations. We report the weighted k-mer enrichment and enrichment p-value for the top significantly enriched k-mers ($1 \le k \le 5$) for the two allelic effect annotations found significant: BasenjiΔ-H3K4me3-Max and BasenjiΔ-H3K4me3-brain-Max annotations. The Bonferonni corrcection threshold is $0.05/(4*682)$ where 682 is the number of k-mers analyzed for the 2 significant variant-level annotations analyzed in Table S25 and the 2 significant allelic-effect annotations analyzed here.

BasenjiΔ-H3K4me3-Max		
kmer	enrichment	pvalue
CGGCG	4.1	3.6e-10
CGCGC	4.1	3.5e-10
CGCCG	4	7.4e-10
BasenjiΔ-H3K4me3-brain-Max		
kmer	enrichment	pvalue
CGCGC	6.2	1.1e-25
CGGCG	6.1	4.9e-25
CGCCG	6	2.8e-24
CGCG	4.3	1.8e-11
CGCGG	4.2	4.9e-11
CCGCG	4.2	9.8e-11
CGCGA	3.7	5.4e-08
AGCGC	3.5	2.8e-07
CCCGC	3.4	8.9e-07
CGCCC	3.3	2.2e-06
ACGCG	3.2	7.7e-06

Table S30. AUROC of various sets of annotations incorporating non-tissue-specific allelic-effect deep learning annotations in predicting $12,296$ **NIH GWAS SNPs**. We report the AUROC for a gradient boosting model trained using each respective set of annotations, either without including baseline-LD model annotations; including baseline-LD model annotations; or including annotations from the non-tissue-specific variant-level joint model (baseline-LD model + 7 annotations; baseline-LD+7).

Feature	GWAS SNPs
DeepSEAΔ-Avg/Max	0.584
BasenjiΔ-Avg/Max	0.592
DeepSEAΔ-All	0.602
BasenjiΔ-All	0.611
baseline-LD	0.758
baseline-LD + DeepSEAΔ-Avg/Max	0.767
baseline-LD + BasenjiΔ-Avg/Max	0.767
baseline-LD + DeepSEAΔ-All	0.770
baseline-LD + BasenjiΔ-All	0.772
baseline-LD+7	0.762
baseline-LD+7 + DeepSEAΔ-Avg/Max	0.766
baseline-LD+7 + BasenjiΔ-Avg/Max	0.769
baseline-LD+7 + DeepSEAΔ-All	0.771
baseline-LD+7 + BasenjiΔ-All	0.773

Table S31. Enrichment of NIH GWAS SNPs for non-tissue-specific allelic-effect annotations. We report the enrichment in deep learning annotations (along with Jackknife standard error) at top 12, 296 known disease-associated SNPs from NIH GWAS catalog[23].

Feature	GWAS SNPs
BasenjiΔ-DNase-Avg	1.93 (0.020)
BasenjiΔ-DNase-Max	1.70 (0.012)
BasenjiΔ-H3K27ac-Avg	1.76 (0.01)
BasenjiΔ-H3K27ac-Max	1.63 (0.009)
BasenjiΔ-H3K4me1-Avg	1.54 (0.008)
BasenjiΔ-H3K4me1-Max	1.53 (0.008)
BasenjiΔ-H3K4me3-Avg	2.08 (0.022)
BasenjiΔ-H3K4me3-Max	1.79 (0.011)
DeepSEAΔ-DNase-Avg	1.64 (0.01)
DeepSEAΔ-DNase-Max	1.41 (0.007)
DeepSEAΔ-H3K27ac-Avg	1.57 (0.009)
DeepSEAΔ-H3K27ac-Max	1.46 (0.008)
DeepSEAΔ-H3K4me1-Avg	1.43 (0.007)
DeepSEAΔ-H3K4me1-Max	1.33 (0.006)
DeepSEAΔ-H3K4me3-Avg	1.67 (0.013)
DeepSEAΔ-H3K4me3-Max	1.50 (0.007)

Table S32. S-LDSC results for marginal analysis of blood-specific allelic-effect deep learning annotations against blood-specific variant-level joint model and 1 significant non-tissue-specific BasenjiΔ annotation. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 blood-specific DeepSEAΔ and 8 blood-specific BasenjiΔ annotations, when conditioned on the blood-specific joint model annotations and 1 significant allelic-effect annotation (BasenjiΔ-H3K4me3-Max) from Table S26. Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-blood-Avg (0.2%)	-0.4	0.12	0.00074	2.3	0.38	1.8e-05
DeepSEAΔ-DNase-blood-Max (0.6%)	-0.35	0.11	0.0011	2	0.26	5.1e-11
DeepSEAΔ-H3K27ac-blood-Avg (0.2%)	-0.23	0.11	0.028	2.6	0.23	1.6e-08
DeepSEAΔ-H3K27ac-blood-Max (0.5%)	-0.34	0.11	0.002	2.1	0.16	1.8e-08
DeepSEAΔ-H3K4me1-blood-Avg (0.3%)	-0.35	0.13	0.0035	1.9	0.16	1.3e-08
DeepSEAΔ-H3K4me1-blood-Max (0.8%)	-0.34	0.12	0.0023	1.7	0.14	6.4e-07
DeepSEAΔ-H3K4me3-blood-Avg (0.1%)	-0.3	0.11	0.0063	2.7	0.35	4.3e-08
DeepSEAΔ-H3K4me3-blood-Max (0.4%)	-0.38	0.14	0.0056	2.2	0.24	7.9e-10
BasenjiΔ-DNase-blood-Avg (0.2%)	-0.31	0.2	0.12	5	0.41	9.5e-11
BasenjiΔ-DNase-blood-Max (0.6%)	-0.012	0.22	0.96	4.7	0.28	4e-08
BasenjiΔ-H3K27ac-blood-Avg (0.2%)	0.34	0.17	0.046	4.8	0.29	1.5e-08
BasenjiΔ-H3K27ac-blood-Max (0.5%)	0.23	0.17	0.17	4	0.19	2.9e-08
BasenjiΔ-H3K4me1-blood-Avg (0.3%)	0.033	0.1	0.75	3.6	0.2	4.9e-09
BasenjiΔ-H3K4me1-blood-Max (0.8%)	-0.006	0.15	0.97	3.3	0.17	1.3e-08
BasenjiΔ-H3K4me3-blood-Avg (0.1%)	0.57	0.33	0.083	6.5	0.55	2.6e-09
BasenjiΔ-H3K4me3-blood-Max (0.4%)	0.39	0.3	0.21	4.7	0.3	7.8e-09

Table S33. S-LDSC results for marginal analysis of blood-specific alllelic-effect deep learning annotations conditioned on the non-tissue-specific final joint model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 blood-specific DeepSEAΔ and 8 blood-specific BasenjiΔ annotations, when conditioned on the non-tissue-specific final joint model. Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-blood-Avg (0.2%)	-0.37	0.14	0.0064	2.9	0.52	1.8e-06
DeepSEAΔ-DNase-blood-Max (0.6%)	-0.49	0.12	2.4e-05	2	0.28	1.6e-06
DeepSEAΔ-H3K27ac-blood-Avg (0.2%)	0.024	0.11	0.83	3.3	0.29	1.2e-18
DeepSEAΔ-H3K27ac-blood-Max (0.5%)	-0.23	0.11	0.038	2.4	0.2	1.2e-16
DeepSEAΔ-H3K4me1-blood-Avg (0.3%)	-0.33	0.099	0.0011	2	0.18	2e-10
DeepSEAΔ-H3K4me1-blood-Max (0.8%)	-0.47	0.098	1.4e-06	1.7	0.15	4.8e-08
DeepSEAΔ-H3K4me3-blood-Avg (0.1%)	-0.071	0.12	0.56	3.8	0.41	1.4e-08
DeepSEAΔ-H3K4me3-blood-Max (0.4%)	-0.23	0.13	0.078	2.7	0.28	5.7e-09
BasenjiΔ-DNase-blood-Avg (0.2%)	0.56	0.2	0.0049	7.2	0.57	1.6e-09
BasenjiΔ-DNase-blood-Max (0.6%)	0.69	0.21	0.00081	5.8	0.37	5.1e-09
BasenjiΔ-H3K27ac-blood-Avg (0.2%)	1.8	0.26	1.5e-12	6.2	0.38	6.6e-09
BasenjiΔ-H3K27ac-blood-Max (0.5%)	1.5	0.22	1.1e-11	4.8	0.25	1.5e-08
BasenjiΔ-H3K4me1-blood-Avg (0.3%)	1.2	0.2	7.7e-09	4.6	0.28	2.1e-09
BasenjiΔ-H3K4me1-blood-Max (0.8%)	1	0.19	8.7e-08	4	0.22	3.4e-09
BasenjiΔ-H3K4me3-blood-Avg (0.1%)	1.9	0.36	1.3e-07	8.6	0.68	3e-09
BasenjiΔ-H3K4me3-blood-Max (0.4%)	2.1	0.36	7.4e-09	5.7	0.34	1.3e-08

Table S34. S-LDSC results for marginal analysis of blood-specific allelic-effect deep learning annotations against blood-specific variant-level joint model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 blood-specific DeepSEAΔ and 8 blood-specific BasenjiΔ annotations, when conditioned on the blood-specific variant-level joint model annotations. Results are meta-analyzed across 11 blood-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-blood-Avg (0.2%)	-0.24	0.14	0.089	2.4	0.38	0.00027
DeepSEAΔ-DNase-blood-Max (0.6%)	-0.18	0.1	0.075	2	0.26	5e-06
DeepSEAΔ-H3K27ac-blood-Avg (0.2%)	-0.073	0.1	0.48	2.7	0.23	9.2e-09
DeepSEAΔ-H3K27ac-blood-Max (0.5%)	-0.22	0.12	0.066	2.1	0.16	1.1e-07
DeepSEAΔ-H3K4me1-blood-Avg (0.3%)	-0.21	0.089	0.021	1.9	0.16	2.6e-08
DeepSEAΔ-H3K4me1-blood-Max (0.8%)	-0.22	0.1	0.036	1.7	0.14	7.7e-06
DeepSEAΔ-H3K4me3-blood-Avg (0.1%)	-0.12	0.11	0.27	2.9	0.36	1.7e-06
DeepSEAΔ-H3K4me3-blood-Max (0.4%)	-0.23	0.15	0.11	2.3	0.25	1.8e-07
BasenjiΔ-DNase-blood-Avg (0.2%)	0.039	0.17	0.81	5.1	0.41	3.7e-11
BasenjiΔ-DNase-blood-Max (0.6%)	0.22	0.18	0.22	4.7	0.29	7.1e-08
BasenjiΔ-H3K27ac-blood-Avg (0.2%)	0.45	0.12	0.00025	4.8	0.31	2.3e-07
BasenjiΔ-H3K27ac-blood-Max (0.5%)	0.36	0.11	0.0017	4	0.21	2.1e-07
BasenjiΔ-H3K4me1-blood-Avg (0.3%)	0.22	0.096	0.021	3.6	0.21	4.5e-08
BasenjiΔ-H3K4me1-blood-Max (0.8%)	0.18	0.11	0.11	3.3	0.18	1.1e-07
BasenjiΔ-H3K4me3-blood-Avg (0.1%)	0.67	0.2	0.00067	6.5	0.52	5.6e-09
BasenjiΔ-H3K4me3-blood-Max (0.4%)	0.47	0.15	0.0018	4.8	0.32	1.2e-07

Table S35. AUROC of various sets of annotations incorporating blood-specific allelic-effect deep learning annotations in predicting 8,741 fine-mapped autoimmune diseease SNPs. We report the AUROC for a gradient boosting model trained using each respective set of annotations, either without including baseline-LD model annotations; including baseline-LD model annotations; or including annotations from the blood-specific variant-level joint model plus BasenjiΔ-H3K4me3-Max (baseline-LD+7+6blood+1).

Feature	Farh et al
DeepSEAΔ-blood-Avg/Max	0.613
BasenjiΔ-blood-Avg/Max	0.672
DeepSEAΔ-blood-All	0.633
BasenjiΔ-blood-All	0.684
baseline-LD	0.841
baseline-LD + DeepSEAΔ-blood-Avg/Max	0.847
baseline-LD + BasenjiΔ-blood-Avg/Max	0.852
baseline-LD + DeepSEAΔ-blood-All	0.848
baseline-LD + BasenjiΔ-blood-All	0.851
baseline-LD+7+6blood+1	0.848
baseline-LD+7+6blood+1 + DeepSEAΔ-blood-Avg/Max	0.847
baseline-LD+7+6blood+1 + BasenjiΔ-blood-Avg/Max	0.851
baseline-LD+7+6blood+1 + DeepSEAΔ-blood-All	0.849
baseline-LD+7+6blood+1 + BasenjiΔ-blood-All	0.851

Table S36. Enrichment of fine-mapped autoimmune disease SNPs for blood-specific allelic-effect annotations. We report the enrichment in blood-specific deep learning allelic-effect annotations (along with Jackknife standard error) for 8741 fine-mapped SNPs in immune-related traits[24].

Feature	Farh et al
BasenjiΔ-DNase-blood-Avg	2.59 (0.05)
BasenjiΔ-DNase-blood-Max	2.41 (0.04)
BasenjiΔ-H3K27ac-blood-Avg	2.44 (0.03)
BasenjiΔ-H3K27ac-blood-Max	2.29 (0.03)
BasenjiΔ-H3K4me1-blood-Avg	2.21 (0.03)
BasenjiΔ-H3K4me1-blood-Max	2.09 (0.02)
BasenjiΔ-H3K4me3-blood-Avg	2.52 (0.03)
BasenjiΔ-H3K4me3-blood-Max	2.40 (0.03)
DeepSEAΔ-DNase-blood-Avg	1.87 (0.03)
DeepSEAΔ-DNase-blood-Max	1.62 (0.02)
DeepSEAΔ-H3K27ac-blood-Avg	1.83 (0.03)
DeepSEAΔ-H3K27ac-blood-Max	1.60 (0.01)
DeepSEAΔ-H3K4me1-blood-Avg	1.58 (0.01)
DeepSEAΔ-H3K4me1-blood-Max	1.45 (0.01)
DeepSEAΔ-H3K4me3-blood-Avg	2.01 (0.04)
DeepSEAΔ-H3K4me3-blood-Max	1.66 (0.02)

Table S37. S-LDSC results for marginal analysis of brain-specific allelic-effect deep learning annotations against brain-specific variant-level joint model and 1 significant non-tissue-specific BasenjiΔ annotation. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 brain-specific DeepSEAΔ and 8 brain-specific BasenjiΔ annotations, when conditioned on brain-specific variant-level joint model and 1 significant non-tissue-specific annotation from Table S26). Results are meta-analyzed across 8 brain-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-brain-Avg (0.4%)	-0.027	0.075	0.72	1.9	0.25	6.1e-05
DeepSEAΔ-DNase-brain-Max (0.7%)	-0.011	0.074	0.88	1.8	0.26	2.8e-05
DeepSEAΔ-H3K27ac-brain-Avg (0.2%)	0.029	0.071	0.68	1.8	0.1	6e-07
DeepSEAΔ-H3K27ac-brain-Max (0.4%)	-0.021	0.072	0.77	1.7	0.11	5.6e-07
DeepSEAΔ-H3K4me1-brain-Avg (0.3%)	0.0093	0.065	0.89	1.7	0.086	3.4e-07
DeepSEAΔ-H3K4me1-brain-Max (0.6%)	-0.013	0.069	0.85	1.6	0.1	1.6e-07
DeepSEAΔ-H3K4me3-brain-Avg (0.1%)	-0.001	0.08	0.99	2.5	0.22	3.6e-05
DeepSEAΔ-H3K4me3-brain-Max (0.2%)	-0.055	0.088	0.54	2.5	0.19	6.5e-06
BasenjiΔ-DNase-brain-Avg (0.4%)	0.25	0.083	0.0024	2.9	0.16	1.5e-07
BasenjiΔ-DNase-brain-Max (0.7%)	0.12	0.077	0.12	2.6	0.18	5.9e-06
BasenjiΔ-H3K27ac-brain-Avg (0.2%)	0.11	0.05	0.023	2.1	0.12	4.4e-07
BasenjiΔ-H3K27ac-brain-Max (0.4%)	0.1	0.055	0.071	2	0.15	2.2e-07
BasenjiΔ-H3K4me1-brain-Avg (0.3%)	0.11	0.058	0.058	1.9	0.077	1e-06
BasenjiΔ-H3K4me1-brain-Max (0.6%)	0.083	0.06	0.17	1.9	0.085	1.1e-07
BasenjiΔ-H3K4me3-brain-Avg (0.1%)	0.34	0.097	0.00047	3.5	0.18	2.5e-08
BasenjiΔ-H3K4me3-brain-Max (0.2%)	0.46	0.095	1.4e-06	3.4	0.19	3.2e-08

Table S38. S-LDSC results for marginal analysis of brain-specific allelic-effect deep learning annotations conditioned on the non-tissue-specific final joint model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 brain-specific DeepSEAΔ and 8 brain-specific BasenjiΔ annotations, when conditioned on the non-tissue-specific final joint model. Results are meta-analyzed across 8 brain-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-brain-Avg (0.4%)	0.25	0.073	0.00053	2.4	0.17	1.4e-05
DeepSEAΔ-DNase-brain-Max (0.7%)	0.21	0.072	0.0034	2.2	0.19	3.9e-08
DeepSEAΔ-H3K27ac-brain-Avg (0.2%)	0.3	0.061	9.4e-07	2	0.094	4.8e-09
DeepSEAΔ-H3K27ac-brain-Max (0.4%)	0.28	0.064	1.1e-05	2	0.1	5.2e-07
DeepSEAΔ-H3K4me1-brain-Avg (0.3%)	0.27	0.056	1e-06	1.9	0.08	7e-10
DeepSEAΔ-H3K4me1-brain-Max (0.6%)	0.27	0.062	2e-05	1.9	0.097	1.7e-08
DeepSEAΔ-H3K4me3-brain-Avg (0.1%)	0.44	0.071	4.5e-10	3.4	0.22	7.8e-05
DeepSEAΔ-H3K4me3-brain-Max (0.2%)	0.5	0.072	2.6e-12	3.3	0.19	3.3e-06
BasenjiΔ-DNase-brain-Avg (0.4%)	0.56	0.097	9.1e-09	3.1	0.16	3.5e-09
BasenjiΔ-DNase-brain-Max (0.7%)	0.46	0.073	3.5e-10	2.9	0.16	9.2e-08
BasenjiΔ-H3K27ac-brain-Avg (0.2%)	0.32	0.046	5.4e-12	2.1	0.096	7.8e-08
BasenjiΔ-H3K27ac-brain-Max (0.4%)	0.25	0.048	1.6e-07	2.1	0.13	4.5e-08
BasenjiΔ-H3K4me1-brain-Avg (0.3%)	0.34	0.069	6.3e-07	2.1	0.072	4.8e-09
BasenjiΔ-H3K4me1-brain-Max (0.6%)	0.23	0.084	0.0072	2	0.08	7.6e-11
BasenjiΔ-H3K4me3-brain-Avg (0.1%)	0.79	0.11	2.6e-13	3.7	0.18	1.1e-08
BasenjiΔ-H3K4me3-brain-Max (0.2%)	0.91	0.079	4e-30	3.4	0.18	2.3e-08

Table S39. S-LDSC results for marginal analysis of brain-specific allelic-effect deep learning annotations conditioned on the brain-specific variant-level joint model. Standardized Effect sizes ($\tau^\star$) and Enrichment (E) of 8 brain-specific DeepSEAΔ and 8 brain-specific BasenjiΔ annotations, when conditioned on the brain-specific variant-level joint model. Results are meta-analyzed across 8 brain-related traits.

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
DeepSEAΔ-DNase-brain-Avg (0.4%)	0.0038	0.076	0.96	1.9	0.25	1.7e-06
DeepSEAΔ-DNase-brain-Max (0.7%)	0.022	0.075	0.77	1.8	0.26	1.9e-06
DeepSEAΔ-H3K27ac-brain-Avg (0.2%)	0.065	0.072	0.36	1.8	0.1	1.8e-09
DeepSEAΔ-H3K27ac-brain-Max (0.4%)	0.022	0.073	0.76	1.7	0.11	1.2e-09
DeepSEAΔ-H3K4me1-brain-Avg (0.3%)	0.039	0.065	0.55	1.7	0.087	1.5e-09
DeepSEAΔ-H3K4me1-brain-Max (0.6%)	0.022	0.069	0.76	1.6	0.1	7.6e-11
DeepSEAΔ-H3K4me3-brain-Avg (0.1%)	0.025	0.08	0.75	2.6	0.22	8.5e-07
DeepSEAΔ-H3K4me3-brain-Max (0.2%)	-0.021	0.089	0.81	2.5	0.19	1.3e-08
BasenjiΔ-DNase-brain-Avg (0.4%)	0.29	0.081	0.00041	2.8	0.16	1e-08
BasenjiΔ-DNase-brain-Max (0.7%)	0.16	0.076	0.035	2.5	0.18	3.7e-07
BasenjiΔ-H3K27ac-brain-Avg (0.2%)	0.15	0.048	0.0026	2	0.12	1.1e-08
BasenjiΔ-H3K27ac-brain-Max (0.4%)	0.13	0.048	0.0054	2	0.16	2e-0
BasenjiΔ-H3K4me1-brain-Avg (0.3%)	0.14	0.056	0.014	1.9	0.076	3.8e-08
BasenjiΔ-H3K4me1-brain-Max (0.6%)	0.12	0.058	0.041	1.9	0.083	2.6e-09
BasenjiΔ-H3K4me3-brain-Avg (0.1%)	0.36	0.08	7.7e-06	3.6	0.18	5.5e-10
BasenjiΔ-H3K4me3-brain-Max (0.2%)	0.41	0.074	4.1e-08	3.4	0.19	6.3e-10

Table S40. S-LDSC results for marginal analysis of brain-specific Basenji variant-level annotations and allelic-effect annotations conditioned on either the brain-specific variant-level joint model and BasenjiΔ-H3K4me3-Max or the brain-specific final joint model. We report the standardized effect sizes (τ^*) and enrichment (E) of Basenji-H3K4me3-brain-Avg, Basenji-H3K4me3-brain-Max and BasenjiΔ-H3K4me3-brain-Avg, when conditioned on either the brain-specific variant-level joint model and BasenjiΔ-H3K4me3-Max (top panel) or the brain-specific final joint model (bottom panel). Results are meta-analyzed across 8 brain-related traits.

Conditional on brain specific variant-level joint model and BasenjiΔ-H3K4me3-Max						
Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Basenji-H3K4me3-brain-Avg (1.9%)	-0.21	0.097	0.02	3.9	0.29	4.3e-08
Basenji-H3K4me3-brain-Max (3.4%)	0.16	0.11	0.16	3.6	0.19	2.4e-08
BasenjiΔ-H3K4me3-brain-Avg (0.1%)	0.34	0.097	0.00047	3.5	0.18	2.5e-08
Conditional on brain-specific final joint model						
Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
Basenji-H3K4me3-brain-Avg (1.9%)	-0.21	0.098	0.02	3.6	0.3	3e-10
Basenji-H3K4me3-brain-Max (3.4%)	0.049	0.11	0.65	3.4	0.2	1.6e-10
BasenjiΔ-H3K4me3-brain-Avg (0.1%)	-0.62	0.33	0.061	3.3	0.18	1.3e-08

Table S41. S-LDSC results for brain-specific final joint model. The results are conditional on the brain-specific variant-level joint model and 1 significant non-tissue-specific allelic-effect annotation. Results are meta-analyzed across 8 brain-related traits

Annotation	$\tau^\star$	se($\tau^\star$)	p($\tau^\star$)	E	se (E)	p(E)
BasenjiΔ-H3K4me3-brain-Avg (0.1%)	-0.62	0.33	0.06	3.3	0.18	1.3e-08
BasenjiΔ-H3K4me3-brain-Max (0.2%)	0.98	0.30	9e-04	3.3	0.18	2.7e-08

Figure S1. Correlations between deep learning annotations. We report correlations between each pair of variant-level (DeepSEA and Basenji) and/or allelic-effect (DeepSEAΔ and BasenjiΔ) annotations. When comparing pairs of annotations that differed only in their aggregation strategy (Avg/Max), chromatin mark (DNase/H3K27ac/H3K4me1/H3K4me3), model (DeepSEA/Basenji) or type (variant-level/allelic-effect), respectively, we observed large correlations across aggregation strategies (average $r = 0.71$), chromatin marks (average $r = 0.58$), models (average $r = 0.54$) and types (average $r = 0.48$).

Figure S2. Disease informativeness of non-tissue-specific Roadmap annotations in marginal analysis. (A) Heritability enrichment, conditioned on the baseline-LD model. Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on the baseline-LD model. Results are meta-analyzed across 41 traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 106 (variant-level + allelic-effect) non-tissue-specific annotations tested ($P < 0.05/106$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S7.

Figure S3. Disease informativeness of non-tissue-specific ChromHMM annotations in marginal analysis. (A) Heritability enrichment, conditioned on the baseline-LD model. Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on the baseline-LD model. Results are meta-analyzed across 41 traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 106 (variant-level + allelic-effect) non-tissue-specific annotations tested ($P < 0.05/106$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S7.

Figure S4. BiClassCNN-TSS is highly correlated with CpG-island, local CpG-content($\pm$1kb) and local GC-content($\pm$1kb) annotations. (A) Violin plot of the BiClassCNN TSS annotation for variants that are TSS and fall in a CpG island and variants that are TSS but do not fall in a CpG island. (B) Scatter plot of the BiClassCNN TSS annotation with the local GC content (1KB) around the variant for all TSS variants. (C) Scatter plot of the BiClassCNN TSS annotation with the local CpG content (1KB) around the variant for all TSS variants. Here, by TSS variants, we imply variants that are annotated to fall in a TSS region as predicted by Segway[74,75]. The BiClassCNN TSS annotations showed strong correlation with all three features - CpG-island, Local GC and Local CpG content at the TSS variants with the correlation with the Local GC content at TSS variants being the strongest.

Figure S5. Disease informativeness of non-tissue-specific Roadmap, ChromHMM, LocalGCcontent and pLI annotations in non-tissue-specific variant-level joint model. (A) Heritability enrichment, conditioned on the baseline-LD model. Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on each other as well as the baseline-LD model. Results are meta-analyzed across 41 traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 106 (variant-level + allelic-effect) non-tissue-specific annotations tested ($P < 0.05/106$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S9.

Figure S6. Disease informativeness of blood-specific Roadmap annotations in marginal analysis conditional on the non-tissue-specific variant-level joint model. (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model (original baseline-LD annotations and the annotations from Figure S5). Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on the non-tissue-specific variant-level joint model. Results are meta-analyzed across 11 blood-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S15.

Figure S7. Disease informativeness of blood-specific ChromHMM annotations in marginal analysis conditional on the non-tissue-specific variant-level joint model. (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model (original baseline-LD annotations and the annotations from Figure S5). Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$)conditioned on the non-tissue-specific variant-level joint model. Results are meta-analyzed across 11 blood-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S15.

Figure S8. Disease informativeness of blood-specific Roadmap and ChromHMM annotations in blood-specific final joint model. (A) Heritability enrichment, conditioned on each other and non-tissue-specific variant-level joint model (original baseline-LD annotations and the annotations from Figure S5). Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on each other and the non-tissue-specific variant-level joint model. Results are meta-analyzed across 11 blood-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). Error bars denote 95% confidence intervals. Numerical results are reported in Table Table S16.

Figure S9. Disease informativeness of brain-specific Roadmap annotations in marginal analysis conditional on the non-tissue-specific variant-level joint model. (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model (original baseline-LD annotations and the annotations from Figure S5). Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on the non-tissue-specific variant-level joint model. Results are meta-analyzed across 8 brain-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S21.

Figure S10. Disease informativeness of brain-specific ChromHMM annotations in marginal analysis conditional on the non-tissue-specific variant-level joint model. (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model (original baseline-LD annotations and the annotations from Figure S5). Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^\star$) conditioned on the non-tissue-specific variant-level joint model. Results are meta-analyzed across 8 brain-related traits. Results are displayed only for annotations with significant $\tau^\star$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). Error bars denote 95% confidence intervals. Numerical results are reported in Table S21.

Figure S11. Disease informativeness of brain-specific Roadmap and ChromHMM annotations in brain-specific final joint model. (A) Heritability enrichment, conditioned on the non-tissue-specific variant-level joint model (original baseline-LD annotations and the annotations from Figure S5), and all brain-specific variant-level DeepSEA and Basenji, and Roadmap and ChromHMM annotations. Horizontal line denotes no enrichment. (B) Standardized effect size ($\tau^{\star}$) conditioned on the non-tissue-specific variant-level joint model and all brain-specific variant-level DeepSEA and Basenji, and Roadmap and ChromHMM annotations. Results are meta-analyzed across 8 brain-related traits. Results are displayed only for annotations with significant $\tau^{\star}$ in marginal analyses after correcting for 80 (variant-level + allelic-effect) blood-specific annotations tested ($P < 0.05/80$). Error bars denote 95% confidence intervals. Numerical results are reported in Table Table S22.

Figure S12. The Neural net architecture for BiClassCNN. The architecture of the Covolutional Neural Net (CNN) model used for BiClassCNN training. It comprioses of two layers - 1 layer being Convolutional+MaxPool with 5 filters and 1 Fully Connected hidden layer with 10 nodes leading to the final output layer with 2 nodes distinguishing between the positive and negative sets. We chose the convolutional filter to be of size 4 and stride length 1 and the MaxPool filter to pool every 4 bases at a time. RELU activation was used.

References

1. M.T. Maurano et al. Systematic localization of common disease-associated variation in regulatory dna. *Science*, 337(6099):pp.1190–1195, 2012.

2. G. Trynka et al. Chromatin marks identify critical cell types for fine mapping complex trait variants. *Nature genetics*, 45(2):p.124, 2013.

3. J.K. Pickrell. Joint analysis of functional genomic data and genome-wide association studies of 18 human traits. *The American Journal of Human Genetics*, 94(4):pp.559–573, 2014.

4. S. Ripke et al. Biological insights from 108 schizophrenia-associated genetic loci. *Nature*, 511(7510):421, 2014.

5. H.K. Finucane, B. Bulik-Sullivan, A. Gusev, G. Trynka, Y. Reshef, P.R. Loh, V. Anttila, H. Xu, C. Zang, K. Farh, and S. Ripke. Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nature genetics*, 47:1228, 2015.

6. A.L. Price, C.C. Spencer, and P. Donnelly. Progress and promise in understanding the genetic basis of common diseases. *Proceedings of the Royal Society B: Biological Sciences*, 282(1821):p.20151684, 2015.

7. P.M. Visscher et al. 10 years of GWAS discovery: biology, function, and translation. *The American Journal of Human Genetics*, 101(1):5–22, 2017.

8. J. Ernst et al. Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature*, 473(7345):p.43, 2011.

9. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature*, 489:57, 2012.

10. R. Andersson et al. An atlas of active enhancers across human cell types and tissues. *Nature*, 507(7493):p.455, 2014.

11. A. Kundaje et al. Integrative analysis of 111 reference human epigenomes. *Nature*, 518:317, 2015.

12. B. Alipanahi, A. Delong, M.T. Weirauch, and B.J. Frey. Predicting the sequence specificities of DNA-and RNA-binding proteins by deep learning. *Nature biotechnology*, 33:831–838, 2015.

13. J. Zhou and O.G. Troyanskaya. Predicting effects of noncoding variants with deep learning–based sequence model. *Nature methods*, 12:931, 2015.

14. D.R. Kelley, J. Snoek, and J.L. Rinn. Basset: learning the regulatory code of the accessible genome with deep convolutional neural networks. *Genome research*, 26:990–999, 2016.

15. J. Zhou, C.L. Theesfeld, K. Yao, K.M. Chen, A.K. Wong, and O.G. Troyanskaya. Deep learning sequence-based ab initio prediction of variant effects on expression and disease risk. *Nature genetics*, 50:1171, 2018.

16. D.R. Kelley, Y.A. Reshef, M. Bileschi, D. Belanger, C.Y. McLean, and J. Snoek. Sequential regulatory activity prediction across chromosomes with convolutional neural networks. *Genome research*, 28:739–750, 2018.

17. J. Zou et al. A primer on deep learning in genomics. *Nature genetics*, 51:12–18, 2019.

18. G. Eraslan et al. Deep learning: new computational modelling techniques for genomics. *Nature Reviews Genetics*, page p.1, 2019.

19. S. Gazal et al. Linkage disequilibrium–dependent architecture of human complex traits shows action of negative selection. *Nat. Genet*, 49 (10):1421, 2017.

20. J. Ernst and M. Kellis. Large-scale imputation of epigenomic datasets for systematic annotation of diverse human tissues. *Nature biotechnology*, 33:364, 2015.

21. J. Ernst and M. Kellis. ChromHMM: automating chromatin-state discovery and characterization. *Nature methods*, 9:215, 2012.

22. J. Ernst and M. Kellis. Chromatin-state discovery and genome annotation with ChromHMM. *Nature protocols*, 12:2478, 2017.

23. J. MacArthur et al. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic acids research*, 45(D1):pp.D896–D901, 2017.

24. K.K.H. Farh et al. Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature*, 518:p.337, 2015.

25. P.W. Koh, E. Pierson, and A. Kundaje. Denoising genome-wide histone ChIP-seq with convolutional neural networks. *Bioinformatics*, 33:i225–i233, 2017.

26. 1000 Genomes Project Consortium. A global reference for human genetic variation. *Molecular cell*, 526(7571):p.68, 2015.

27. D. Shlyueva, G. Stampfel, and A. Stark. Transcriptional enhancers: from properties to genome-wide predictions. *Nature Reviews Genetics*, 15(4):pp.272–286, 2014.

28. S. Gazal, C. Marquez-Luna, H.K. Finucane, and A.L. Price. Reconciling s-ldsc and ldak models and functional enrichment estimates. *Nat. Genet*, page p.256412, 2019.

29. F. Hormozdiari et al. Leveraging molecular quantitative trait loci to understand the genetic architecture of diseases and complex traits. *Nature genetics*, 50(7):p.1041, 2018.

30. C. Bycroft et al. The uk biobank resource with deep phenotyping and genomic data. *Nature*, 562(7726):p.203, 2018.

31. M. Lek, K.J. Karczewski, E.V. Minikel, K.E. Samocha, E. Banks, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature*, 536:p.285, 2016.

32. I.A. Adzhubei, S. Schmidt, L. Peshkin, V.E. Ramensky, A. Gerasimova, et al. A method and server for predicting damaging missense mutations. *Nature methods*, 7:p.248, 2010.

33. I. Adzhubei, D.M. Jordan, and S.R. Sunyaev. Predicting functional effect of human missense mutations using PolyPhen-2. *Current protocols in human genetics*, 76:p.7–20, 2013.

34. P. Kumar, S. Henikoff, and P.C. Ng. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nature protocols*, 4:p.1073, 2009.

35. F.I. Hormozdiari et al. Functional disease architectures reveal unique biological role of transposable elements. *Nature Communications*, 10(1):4054, 2019.

36. A. Weiner et al. High-resolution nucleosome mapping reveals transcription-dependent promoter packaging. *Genome research*, 20(1):pp.90–100, 2010.

37. A. Mahpour et al. A methyl-sensitive element induces bidirectional transcription in tata-less cpg island-associated promoters. *PloS one*, 13(10):p.e0205608, 2018.

38. J. Zhou et al. Whole-genome deep-learning analysis identifies contribution of noncoding mutations to autism risk. *Nature genetics*, 51(6):p.973, 2019.

39. K. Jaganathan et al. Predicting splicing from primary sequence with deep learning. *Cell*, 176(3):pp.535–548, 2019.

40. L. Liu et al. Biological relevance of computationally predicted pathogenicity of noncoding variants. *Nature Communications*, 10:p.330, 2019.

41. Y.A. Reshef, H.K. Finucane, D.R. Kelley, A. Gusev, D. Kotliar, J.C. Ulirsch, et al. Detecting genome-wide directional effects of transcription factor binding on polygenic disease risk. *Nature genetics*, 50:1483, 2018.

42. D.R. Kelley. Cross-species regulatory sequence activity prediction. *BioRxiv*, page p.660563, 2019.

43. H. Yoshida et al. The cis-Regulatory Atlas of the Mouse Immune System. *Cell*, 176:pp.897–912, 2019.

44. M. Ghandi, D. Lee, M. Mohammad-Noori, and M.A. Beer. Enhanced regulatory sequence prediction using gapped k-mer features. *PLoS computational biology*, 10:p.e1003711, 2014.

45. J.W. Whitaker, Z. Chen, and W. Wang. Predicting the human epigenome from DNA motifs. *Nature methods*, 12:265–272, 2015.

46. D. Lee, D.U. Gorkin, M. Baker, B.J. Strober, A.L. Asoni, A.S. McCallion, and M.A. Beer. A method to predict the impact of regulatory variants from DNA sequence. *Nature genetics*, 47:955, 2015.

47. D. Smedley et al. A whole-genome analysis framework for effective identification of pathogenic regulatory variants in Mendelian disease. *The American Journal of Human Genetics*, 99(3):595–606, 2016.

48. A.C. Wells et al. Identification of essential regulatory elements in the human genome. *biorxiv*, page p.444562, 2018.

49. M. Lizio et al. Gateways to the fantom5 promoter level mammalian expression atlas. *Genome biology*, 16 (1):p.22, 2015.

50. M. Lizio et al. Update of the fantom web resource: high resolution transcriptome of diverse cell types in mammals. *Nucleic acids research*, 45 (D1):p.D737, 2017.

51. B. van de Geijn et al. Annotations capturing cell-type-specific tf binding explain a large fraction of disease heritability. *bioRxiv*, page p.474684, 2018.

52. M. Ghandi, M. Mohammad-Noori, N. Gharegani, D. Lee, L. Garraway, and M.A. Beer. gkmSVM: an R package for gapped-kmer SVM. *Bioinformatics*, 32:2205–2207, 2016.

53. D. Karolchik, A.S. Hinrichs, T.S. Furey, K.M. Roskin, C.W. Sugnet, et al. The UCSC Table Browser data retrieval tool. *Nucleic acids research*, 32:pp.D493–D496, 2004.

54. P. Rentzsch et al. CADD: predicting the deleteriousness of variants throughout the human genome. *Nucleic acids research*, 47(D1):pp.D886–D894, 2018.

55. S. Gazal et al. Functional architecture of low-frequency variants highlights strength of negative selection across coding and non-coding annotations. *Nat. Genet*, 50:1600–1607, 2018.

56. J.H. Friedman. Greedy function approximation: a gradient boosting machine. *Annals of statistics*, pages pp.1189–1232, 2001.

57. T. Chen and C. Guestrin. Xgboost: A scalable tree boosting system. *In Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining*, ACM:785–794, 2016.

58. B. Caron, Y. Luo, and A. Rausell. NCBoost classifies pathogenic non-coding variants in Mendelian diseases through supervised learning on purifying selection signals in humans. *Genome biology*, 20:32, 2019.

59. L. Jostins et al. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature*, 491:119–124, 2012.

60. Y. Okada et al. Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature*, 506:376–381, 2014.

61. J. Bentham et al. Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus. *Nature genetics*, 47(12):p.1457, 2015.

62. P.C. Dubois et al. Multiple common variants for celiac disease influencing immune gene expression. *Nature genetics*, 42(4):p.295, 2010.

63. E.K. Speliotes et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature genetics*, 42:937–948, 2010.

64. A. Okbay et al. Genome-wide association study identifies 74 loci associated with educational attainment. *Nature*, 533:539, 2016.

65. Tobacco and . Genetics Consortium. Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nature genetics*, 42:441–447, 2010.

66. Psychiatric GWAS Consortium Bipolar Disorder Working Group. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near odz4. *Nature genetics*, 43:977–983, 2011.

67. N. Barban et al. Genome-wide analysis identifies 12 loci influencing human reproductive behavior. *Nature genetics*, 48(12):p.1462, 2016.

68. V. Boraska et al. A genome-wide association study of anorexia nervosa. *Molecular psychiatry*, 19(10):p.1085, 2014.

69. Cross-Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *The Lancet*, 381(9875):p.1371–1379, 2013.

70. H. Schunkert et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nature genetics*, 43:333–338, 2011.

71. T.M. Teslovich et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*, 466:707–713, 2010.

72. A. Lango et al. Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature*, 467:832–838, 2010.

73. A.P. Morris et al. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nature genetics*, 44:981–990, 2012.

74. M.M. Hoffman, J. Ernst, S.P. Wilder, A. Kundaje, R.S. Harris, and M. Libbrecht. A method to predict the impact of regulatory variants from DNA sequence. *Nucleic acids research*, 41:827–841, 2012.

75. M.M. Hoffman, O.J. Buske, J. Wang, Z. Weng, J.A. Bilmes, and W.S. Noble. Unsupervised pattern discovery in human chromatin structure through genomic segmentation. *Nature methods*, 9:473, 2012.