TITLE

ISG15/USP18/STAT2 is a molecular hub regulating autocrine IFN I-
mediated control of Dengue and Zika virus replication

Constanza Eleonora Espada[1], Edroaldo Lummertz da Rocha[1], Adara Aurea
dos Santos[1], Zamira Guerra Soares[1], Greicy Malaquias[1], Daniel Oliveira
Patrício[1], Edgar Gonzalez Kozlova[1], Paula Fernandes dos Santos[1], Juliano
Bordignon[2], Thomas J. Sanford[3], Teodoro Fajardo[3], Trevor R. Sweeney[3],
André Báfica[1], Daniel Santos Mansur[1*]

[1]Laboratório de Imunobiologia, Departamento de Microbiologia, Imunologia e
Parasitologia, Centro de Ciências Biológicas, Universidade Federal de Santa
Catarina, Santa Catarina CEP 88040-900, Brazil.

[2]Laboratório de Virologia Molecular, Instituto Carlos Chagas, ICC/Fiocruz-PR,
Curitiba, Paraná, CEP 81350-010, Brazil.

[3]Division of Virology, Department of Pathology, University of Cambridge,
Addenbrooke's Hospital, Hills Road, Cambridge, CB2 0QQ, UK.

[*]Correspondence: daniel.mansur@ufsc.br

SUMMARY

The establishment of a virus infection is the result of the pathogen's ability to
replicate in a hostile environment generated by the host's immune system.
Here, we found that ISG15 restricts Dengue and Zika viruses' replication
through the stabilization of its binding partner USP18. ISG15 expression was
necessary to control DV replication driven by both autocrine and paracrine
type one interferon (IFN-I) signaling. Moreover, USP18 competes with NS5-
mediated STAT2 degradation, a major mechanism for establishment of
flavivirus infection. Strikingly, reconstitution of USP18 in ISG15-deficient cells
was sufficient to restore the cells' immune response and restrict virus growth,

suggesting that the IFNAR-mediated ISG15 activity is also antiviral. Our
results add a novel layer of complexity in the virus/host interaction interface
and suggest that NS5 has a narrow window of opportunity to degrade STAT2,
therefore suppressing host's IFN-I mediated response and promoting virus
replication.
40
41 **KEYWORDS**
42 Dengue virus, Zika virus, ISG15, USP18, type one interferon, ISGylation,
43 antiviral response, immune evasion, NS5, STAT2
44
45 **INTRODUCTION**
46
Cells detect infection by recognizing molecular patterns derived from
pathogen's constituents (PAMPs) or cell damage (DAMPs). During viral
infection, nucleic acid is a major signal that triggers the innate immune
response, inducing a type one interferon (IFN-I)-mediated antiviral state (1, 2).
IFN-I binds to its cognate receptor and activates the JAK/STAT pathway,
leading to expression of hundreds of interferon stimulated genes (ISGs) that
make the intracellular environment hostile to viral replication in infected and
proximal cells (3). The evolutionary arms race between viruses and its hosts
led to evolution of immune evasion mechanisms that are crucial for successful
viral replication. Considering IFN-I's importance in viral infection control, many
immunomodulatory proteins target this signaling pathway (4).
Several flaviviruses, such as dengue virus (DV), Zika virus (ZIKV) and yellow
fever virus (YFV), have emerged and re-emerged over recent years and are
the leading cause of human arbovirus infection (5, 6). DV alone infects nearly
400 million people every year (7) with extensive health and economic burden
(8). One possibility for effective flavivirus emergence is their ability to
counteract the human immune system.
The compact flavivirus genome encodes seven non-structural proteins that
are responsible for viral replication and immune evasion. Six of these proteins
are not secreted implying that intracellular pathways are central targets for
evasion (9–13). For instance, DV non-structural protein 5 (NS5), which is the
viral RNA-dependent-RNA-polymerase (RdRp) and a methyltransferase,

mediates STAT2 degradation by facilitating its interaction with UBR4, leading
to its ubiquitination and subsequent proteasomal targeting (14). This evasion
pathway is functional in humans but not mice due differences in the amino
acid sequence of mouse STAT2 (15).
ISG15 is an intracellular and secreted ubiquitin-like protein that has three
described functions. Extracellular ISG15 acts as a cytokine, leading to the
expression of IFNγ and IL-10 in diverse immune cells (16–18). It has been
suggested that humans lacking ISG15 have severe mycobacterial disease
due to deficiency in IFNγ production by NK cells (19). Moreover, ISG15 mRNA
is highly expressed in active tuberculosis and strongly correlates with disease
severity (17, 20).
ISG15 is conjugated to other proteins through a three-step ubiquitination-like
process (21) in which the main ligase for ISGylation is the HECT domain and
RCC1-like domain-containing protein 5 (HERC5) (22, 23). ISGylated proteins
are affected in several different ways, such as increased or reduced stability
and activity. ISG15 can also be conjugated to viral proteins, impacting their
function, and therefore belong in the plethora of ISGs with a direct antiviral
function (3, 24).
The third and more recently described role of ISG15 is its IFN-I modulatory
function. Non-conjugated ISG15 binds and stabilizes the ISG USP18, a
protease that negatively regulates IFN-I signaling and also serve as a
deISGylating protein (25–27). Indeed, individuals lacking ISG15 are prone to
severe interferonopathies due to decreased USP18 function and increased
IFN-I signaling. Interestingly, ISG15-deficient patients do not have enhanced
susceptibility to virus suggesting ISG15 is not necessary to control ubiquitous
viral infections in vivo (28). Despite cells deficient for ISG15 having less
USP18 and sustained IFN-I signaling in humans (28, 29), in mice ISG15 itself
blocks replication of human viruses such as Influenza and HSV-1(30).
Although this evidence places ISG15 as an important molecule inhibiting IFN-
I-mediated actions, it also suggests that ISG15 mediates host cell intrinsic
mechanisms triggered by viruses. However, how ISG15 bridges these
apparently two paradoxical phenomena is unclear. Specifically, it is possible

that ISG15 directly regulates proteins explored by viruses during early
intracellular infection events.
Here, we observed that, unlike several other ISGs, ISG15 is highly expressed
in infected cells containing the DV genome. Furthermore, independently of its
ISGylation function, ISG15 restricts flavivirus replication primarily in the
infected cell by stabilizing USP18, which in turn competes with viral NS5 for
binding to STAT2. Our results suggest that flaviviruses exploit an ISG15-
mediated IFN-I regulatory mechanism to evade innate immunity and enable
replication.
110
111
112 **RESULTS**
113
114 *ISG15 is expressed in DV-infected cells*
115
The cell is a fundamental unit for viral infection control and developments in
single-cell sequencing technology have enabled examination of host-
pathogen interactions in great detail. Recently, Zanini and colleagues
generated single-cell RNA sequencing of two independent data sets of DV-
infected human cells (PBMCs and the HuH7 hepatoma cell line) (31, 32). We
re-analyzed available single-cell transcriptomic data dividing cells into three
categories: uninfected, infected and bystander. Here we define uninfected
cells as the ones derived from healthy donors, bystander as cells that have
been exposed to the virus or were derived from an infected patient but did not
have the viral RNA detected and infected cells as the ones where the viral
genome was found. We used t-Distributed Stochastic Neighbor Embedding
(tSNE) analysis to visualize cell-to-cell relationships in space of reduced
dimensionality. As reported previously (31, 32), global cellular mRNA
expression profiling was not sufficient to separate infected or bystander from
uninfected cells, suggesting a high variability of gene expression in these
samples (Figure 1A and S1A). As IFN-I are key elements in controlling
infection, we filtered the results of the differential gene expression analysis
using the gene ontology (GO) term for "type one interferon". The Venn
diagram in figure 1B shows that from the 394 differentially expressed (DE)

genes in peripheral blood mononuclear cells (PBMCs) of patients infected with DV or healthy donors, 37 were ISGs (IFN-I GO). ISG15, UBE2L6, HERC5 and USP18, members of the ISGylation pathway, were differentially expressed during DV infection (Figure 1C in bold). Interestingly, in contrast to other single cell experiments using Influenza virus as a model where all the members of the ISGylation pathway seem to be expressed at similar levels in both infected and bystander cells (33), HERC5 was the only member of the ISGylation pathway with a higher expression level in bystander cells (Figure 1C and D). In the data set derived from the Huh7 cell line, ISG15 was the only canonical antiviral protein expressed in DV genome-containing cells (Figure S1C and D). This is in agreement with previous reports that the Huh7 cell line does not produce IFN-I upon viral infection (34–36) and could explain the high number of infected cells (Figure S1A) in comparison with the number of infected PMBCs (Figure 1A). In PBMCs, NK, monocytes and B cells were the infected cells with higher ISG15 expression and similar to the data from Zanini and colleagues (32) B cells and monocytes being proportionately the most infected cells (Figure 1E). These results show that in contrast to most ISGs, ISG15 and other components of the ISGylation pathway are enriched in cells where the DV genome was present.

ISG15 restricts DV and ZIKV replication

The enrichment of ISG15-related genes at a single-cell level led us to investigate how ISG15 might impact flavivirus replication in a human cell. We used an A549 cell line lacking ISG15, previously generated using CRISPR/Cas9 in our lab (17). A549 cells were chosen due to their ability to support flavivirus replication and most importantly to produce and respond to IFN-I (37–40).

ISG15-deficient cells were more susceptible to DV infection as shown by an increased plaque size (Figure 2A and B) and number of infected cells per plaque (Figure 2C). In addition, we determined the kinetics of DV replication and dissemination by using a low multiplicity of infection (MOI), to allow for viral spread through secondary infection events. Percentages of infected cells over time (Figure 2D), relative DV Pre-Membrane RNA quantification in the

supernatant (Figure 2E) and infectious particle formation (Figure 2F) were

also increased in the absence of ISG15. DV infection at 4^o C for 2 hours

followed by relative intracellular DV genome quantification indicated no

differences in viral RNA between WT and knockout cells up to 48 hours post-

infection (Figure 2G), suggesting that ISG15 plays a role in the DV life cycle at

a stage following viral entry. Importantly, reconstitution of ISG15 expression in

knockout cells led to phenotypic reversion (Figure 2H and I), confirming that

the effects observed in our experiments were indeed caused by the lack of

ISG15.

Finally, lack of ISG15 expression led to an increase in plaque size in cells

infected with a phylogenetically related virus, such as ZIKV (Figure 2J and K)

but not with HSV-1 (Figure S2A and B) or VSV (Figure S2C and D). These

results are in line with Speer and colleagues' data where cells isolated from

humans deficient for ISG15 do not have enhanced susceptibility to HSV-1 or

VSV (29). Altogether these results suggest a specific role for ISG15 in the

regulation of flavivirus replication and dissemination.

ISGylation deficiency does not affect DV spread

ISG15 is an IFN-I-inducible ubiquitin-like molecule and can be conjugated to

target proteins by HERC5, an ISG15 ligase also induced by IFN (23, 41). Of

note, ISGylation of host or viral proteins was reported to inhibit replication of

several viruses such as influenza (IAV) (42), human cytomegalovirus (HCMV)

(43) and human respiratory syncytial virus (RSV) (44). Moreover, DV proteins

were also shown to be ISGylated (45). To determine whether ISGylation could

be involved in DV restriction, we generated HERC5-deficient A549 cells using

CRISPR/Cas9 (Figure S3A) and accordingly, A549 HERC5 null cells were not

able to perform ISGylation after IFN-I treatment (Figure 3A). In contrast to

ISG15 null cells, we did not observe differences between WT and HERC5-

deficient cells when we evaluated both plaque area (Figure 3B) and number

of infected cells per plaque following infection with DV (Figure 3C). These

results suggest that ISGylation is not sufficient to inhibit DV replication.

ISG15 is necessary for autocrine IFNAR1-mediated control of DV replication

ISG15 and its binding partner USP18 are crucial for IFN-I pathway down-regulation, which is pivotal for infection control and immune-regulation in humans (28, 29, 46). The A549 ISG15-KO cell line exhibited a lower expression of USP18 and sustained ISG expression, as exemplified by IFIT3, after IFN-I stimulation (Figure S4), reproducing the phenotype observed in cells isolated from humans lacking ISG15 (29).

While our single-cell RNA-seq analysis showed that both ISG15 and USP18 are upregulated in PMBCs containing DV RNA, there was an asynchrony of ISGs being expressed in infected versus bystander cells, with a greater spectrum present in the latter (Figure 1C). This is expected as flavivirus' control of IFN-I signaling occurs mainly via its intracellular non-structural proteins (47, 48) and few ISGs can be directly induced by IRF3 (49–51).

Of interest, STAT2 mRNA is expressed at higher levels in DV RNA-containing cells when compared to bystanders and uninfected cells (Figure 1C). Hence, we hypothesized that ISG15 interference with DV replication is associated with autocrine IFN-I signaling in infected cells. Confirming the results from PMBCs and HuH7 cells, ISG15 mRNA was induced in A549 cells during DV infection (Figure 4A). As expected, this expression was abolished in knockout cells. Therefore, we measured the activation of the IFN-I pathway in WT and ISG15-KO cells infected with DV. Despite ISG15-knockout cells having full machinery to control viral infection as well as hyper-responsiveness to exogenous IFNα (Figure S5, Figure 4E and F) (28, 29), they did not respond properly to DV infection. Infected ISG15-deficient cells had less STAT1 phosphorylation and IFIT1 expression when compared to WT cells and as expected had no detectable USP18 (Figure 4B). Infected knockout cells, despite inducing IFNβ mRNA at higher levels than the WT, had less IFIT1 mRNA (Figures 4C and D) indicating that DV infection impairs the response at both mRNA and protein levels downstream of IFN-I induction.

Taken together, our results suggest that ISG15 inhibits DV infection by
regulating early events of the IFN-I receptor activation. To test this, we
constructed an ISG15/IFNAR1 double knockout cell line based on the ISG15-
KO background (Figure S3B). Cells were infected with DV and plaque area
and number of infected cells per plaque was quantified. In parallel, we
performed this experiment in IFNAR1 knockout cells, previously generated by
our group (52, 53). The same phenotype was observed in all three cell lines,
with no additive or synergistic effects in the double knockout (Figures 4G and
H), which places intracellular ISG15 downstream of IFNAR1 signaling in the
control of DV infection.
245
246   *ISG15 counteracts DV IFN-I evasion*
247
Flaviviruses are known to counteract IFN-I signaling by inducing the
degradation of STAT2, a key protein in the interferon signal transduction
pathway (11, 14, 54). As our previous results suggest that ISG15's role during
DV infection is downstream of IFNAR engagement (Figure 4), we evaluated
DV-mediated STAT2 degradation in the absence of ISG15. In agreement with
others (55), STAT2 degradation in the A549 cell line occurs rapidly after
infection. Despite showing sustained activation of IFN-I signaling (Figure S4),
ISG15-KO cells showed pronounced STAT2 degradation when infected with
DV (Figure 5A and E-upper panel). As infected IFN-secreting cells are able to
induce an antiviral state in neighboring-bystander cells by inducing the
expression of ISGs (56), we then evaluated in which cell population, infected
and/or bystander, ISG15 impacted DV infection. DV plaques were co-stained
for flavivirus E protein and IFIT3 and confocal microscopy was performed.
Both infected and bystander cells were able to respond to infection, producing
IFIT3 (Figure 5D, top panel). However, IFIT3 expression in the plaque context
was largely impaired and restricted to infected cells in the absence of ISG15
(Figures 5B, C and D). This data suggests that restriction of DV replication
and dissemination is achieved by both autocrine and paracrine IFN-I response
amplification and is dependent on ISG15 expression. To further investigate
this, we sorted DV envelope protein positive and negative cells (Figure S5A
and B), therefore enabling us to evaluate the impact of infection in virus-

containing and bystander cells, respectively. Strikingly, DV-positive cells had
a lower expression of STAT2, IFIT3 and USP18 when compared to bystander
cells; a phenotype that was markedly enhanced in ISG15-deficient cells
(Figure 5E, *bottom panel*). These results confirm DV inhibits IFNAR signaling
and suggest ISG15 function is targeted for successful viral infection.

USP18 expression overcomes ISG15 deficiency

STAT2 degradation mediated by all DV serotypes and ZIKV is largely
dependent on NS5 (11, 14), the virus RdRp and methyltransferase. NS5
were shown to bind to the N-terminus of human STAT2 (11, 15). Interestingly,
USP18 interacts with STAT2 via its coiled-coil and DNA binding domains. This
causes negative regulation of IFNAR signaling by displacing JAK1 from chain
2 of the receptor (25). As shown here (Figures 4B, 5E and S4) and elsewhere
(28, 29), the absence of ISG15 results in USP18's destabilization. We
therefore assessed whether NS5 is also present in the STAT2/USP18
complex. In HEK293 cells lacking ISG15 (Figure S3C, D and E) and primed
with IFNα for 18 hours, both endogenous STAT2 and overexpressed USP18,
immuno-precipitated with ZIKV's NS5 (Figure 6A, lane 2). This interaction was
enhanced when a protease-deficient high-expressing USP18 mutant (C64A)
was used (Figure 6A, lane 3). We then hypothesized that the interaction of
USP18 with STAT2 competes with NS5 binding, which in turn could result in a
lower efficiency of virus evasion mechanism. To test this, we performed an
endogenous STAT2 pull-down in ISG15-deficient HEK293 cells transfected
with NS5, USP18 or a combination of both, followed by treatment with IFNα
overnight. In the absence of NS5, USP18 interaction with STAT2 was
markedly enhanced (Figure 6B, lanes 2 and 3), suggesting that both proteins
indeed compete for the same region of the protein.
Thus, we evaluated whether the presence of USP18 could restore STAT2
expression in ISG15 knockout cells during DV infection. Overexpression of
USP18 led to an increase of STAT2 levels similar to ones seen in wild type
cells (Figure 6C). Importantly, USP18 reconstitution in ISG15-deficient cells
was also able to reduce virus replication (Figure 6D), suggesting that

reestablishment of the IFN-I signaling was sufficient to recover the WT
phenotype in ISG15 knockout cells.
Together, the results presented here reveal that human ISG15 restricts DV
and ZIKV replication via its ability to stabilize USP18 and regulate the type 1
IFN signaling pathway.
307
308 **DISCUSSION**
309
A successful infection is dependent on the virus replication machinery and its
ability to evade host immunity. One of the first lines of defense a virus has to
overcome is a plethora of antiviral genes induced by IFN-I (57).
ISG15 is induced early during infection (33, 58) and has been shown to have
a viral restriction role in several infection models. Most of those have been
reported to be a consequence of ISGylation (42–44, 59), even though this
process was suggested to be both inefficient and unspecific (59). Here we
show that during the IFN-I response elicited throughout DV infection,
ISGylation is redundant for antiviral immunity; rather, ISG15's ability to
stabilize USP18 prevents NS5-mediated STAT2 degradation, thus leading to
a more effective interferon response that culminates in DV and ZIKV
restriction.
Many of ISG15's functions have been shown to be immunomodulatory. For
instance, ISGylation stabilizes IRF3 by occluding its ubiquitylation sites (60),
negatively regulates RIG-I (61) and activates PKR (62). Secreted ISG15
works as a cytokine (16, 63), leading to the production of IFNγ and IL-10 in
human cells, crucial to the control of pathogens such as *Mycobacterium*
tuberculosis (17, 19). Moreover, free intracellular ISG15 is essential for
USP18 stability (Figures, 4B, 5E, S4 and (28, 29) and its absence leads to
severe interferonopathy in humans (28). Also there are reports of ISG15
having an antiviral function independent of Ube1L during Chikungunya virus
infection in mice (28, 64), where free ISG15 contributes to infection control by
blunting potentially pathologic levels of cytokine effectors. Considering this
range of functions, it is expected that different pathogens might interact with
this pathway in different ways, according to its co-evolutionary history.

Of note, NK, NKT and monocytes were the PBMC populations with higher
upregulation of ISG15 mRNA in single-cell gene expression studies. These
have been shown to be the major producers and/or targets of free
extracellular ISG15 in other contexts (17–19, 63). The influence of
extracellular ISG15 during viral infections should be further explored in the
future.

Interestingly, humans lacking ISG15 do not have increased susceptibility to
common viral infections, such as influenza and HSV-1. The explanation for
that, as suggested elsewhere, might lie on the sustained IFN-I response of
their cells creating a hostile environment for virus growth (28, 29). Here, we
further expand these studies and reveal that ISG15 restrict DV and ZIKV
replication by stabilizing USP18 and therefore competing for a niche exploited
by such viruses. We also demonstrate that USP18, STAT2 and NS5 co-
immunoprecipitate, suggesting a very narrow window of opportunity that NS5
has to degrade STAT2. As USP18/ ISG15 interaction seems to result in the
down-regulation of IFN-I signaling in humans but not mice (29), it is tempting
to speculate that NS5 interaction with STAT2, a major flavivirus immune
evasion mechanism and also restricted to humans (15), was shaped by the
USP18/ISG15 interaction.

Our results suggest an unexpected mechanism by which ISG15 can exert an
antiviral function distinct from the debilitating effects of its conjugation to viral
proteins. The key role of IFN-I in viral infections might lead to the perception
of ISGs having a necessarily direct antiviral function, a paradigm that is
recently being reassessed, with a range of ISGs being implicated in infection-
independent functions (65, 66). Here we provide mechanistic insight of the
arms race between ISG15, USP18 and NS5 that suggests that protein/protein
dynamics adjacent to IFNAR are a key determinant for the outcome of
flavivirus infection.

ACKNOWLEDGMENTS

This work was supported by Comissão de Aperfeiçoamento de Pessoal de
Nível   Superior   (CAPES)   Computational   Biology   programme
(23038.010048/2013-27), Conselho Nacional de Desenvolvimento Científico e
Tecnológico (CNPq) (311406/2017-3 and 407609/2018-0) and the Academy

of Medical Sciences/U.K. (NAF004/1005) to D.S.M. Howard Hughes Medical
Institute – Early Career Scientist to A.B. (55007412), Wellcome Trust and
Royal Society Sir Henry Dale Fellowship (202471/Z/16/Z) to T.R.S. Wellcome
Trust PhD Studentship (105389/Z/14/Z) to T.J.S. CAPES scholarship
(88887.195782/2018-00) to E.L.R. CNPq scholarship (205096/2018-2) to J.B.
We thank Dr. Sandra Pellegrini and Dr. Carsten Münk for kindly providing
human USP18 plasmids. We thank Dr. Brian Ferguson for critical reading of
the manuscript.
377

AUTHOR CONTRIBUTIONS
Conceptualization, C.E.E., A.B. and D.S.M.; Methodology, C.E.E., E.L.R and
D.S.M.; Software, E.L.R; Formal Analysis, C.E.E., E.L.R and E.G.K.,
Investigation, C.E.E., A.A.S., D.O.P, G.M. and JB; Resources, C.E.E., A.A.S,
P.F.S., Z.G, T.J.S., T.F. and T.R.S. Data Curation, E.L.R. and E.G.K., Writing-
Original Draft, C.E.E. and D.S.M., Writing-Review and editing, C.E.E, E.L.R,
D.O.P., E.G.K., P.F.S., J.B., T.J.S., T.F., T.R.S, A.B., and D.S.M.,
Visualization, C.E.E., E.L.R. and D.S.M., Supervision, D.S.M., Project
Administration and Funding Acquisition, D.S.M.
387

388 **DECLARATION OF INTERESTS**
389 The authors declare no competing interests.
390

FIGURE LEGENDS
Figure 1. ISG15 is expressed in DV-infected cells
(A) tSNE was used to visualize the single-cell global transcriptome data. Blue
dots represent uninfected cells, derived from healthy donors. Beige dots
represent bystander cells and red dots represent infected cells, both derived
from patients infected with DV.
(B) Differential expression of ISGs in PBMCs of patients infected with DV. The
Gene Ontology term "type one interferon" was used to filter the results from
the single-cell RNA sequencing.
(C) Single cell ISG expression variability in uninfected, bystander and
infected. ISGylation related genes are underlined. Blue bar represents
uninfected cells, derived from healthy donors. Beige bar represents bystander

cells and red bar represents infected cells, both derived from patients infected

with DV.

(D) Violin plot representing the expression of ISGylation family members in

uninfected [U], bystander [B] and infected [I] PMBC.

(E) Expression of ISG15 in DV infected PMBCs. Size is proportional to the

percentage of infected cells in each cell type. Color intensity represents

ISG15 average expression.

Figure 2. ISG15 restricts DV and ZIKV replication

(A) A549 WT and ISG15 KO were infected with 20 DV PFUs. At 36 hpi cells

were fixed, permeabilized and stained for the flavivirus E protein using 4G2

antibody. Displayed images were acquired with a Leica DMI6000 B

microscope.

(B and C) DV relative plaque area (B) and the number of infected cells per

plaque (C) quantified by ImageJ software and analyzed using Welch's t test.

Error bars represent mean ± SD. Results are representative of three

independent experiments.

(D, E and F) Multiple-step DV growth curve in A549 cells. Shown is the

percentage of cells infected as measured by E protein staining (4G2+) (D),

extracellular DV mRNA (PrM) relative expression by RT-qPCR (E) and

titration by focus forming assay (FFA) (F). Statistical analyses were conducted

using unpaired t tests. Error bars represent mean ± SD. Results are

representative of three independent experiments.

(G) Changes in genomic DV mRNA relative expression over time following

binding of DV to A549 WT and ISG15 KO cells at 4°C and analyzed using

unpaired t test. Error bars represent mean ± SD. Results are representative of

two independent experiments.

(H and I) Complementation of A549 ISG15 KO cells with ectopically

expressed ISG15. Shown is the percentage of cells infected 36 hpi as

measured by E protein staining (4G2+) (H) and DV mRNA relative expression

by RT-qPCR (I). One-way ANOVA was used to analyze these experiments.

EV: empty vector. Error bars represent mean ± SD. Results are

representative of two independent experiments.

(J and I) A549 WT and ISG15 KO were infected with 20 ZIKV PFUs. At 36 hpi
cells were fixed, permeabilized and stained for the flavivirus E protein. ZIKV
relative plaque area (J) and number of infected cells per plaque (I), quantified
by ImageJ software and analyzed using unpaired t test with Welch's
correction. Images were acquired with an Olympus IX83 inverted microscope.
Error bars represent mean ± SD. Results are representative of three
independent experiments.
Statistical analyses were performed using Prism 8 (GraphPad Software). p
values *<0.05; **<0.01; ***<0.001.
445

Figure 3. ISGylation is not sufficient to restrict DV spread
(A) ISGylation profile of A549 WT and HERC5 KO cells by Western blot. Cells
were primed with IFNα2b (100 IU/ml) for 24 h and cell lysates were analyzed
with an ISG15 antibody. (*) indicates antibody unspecific band.
(B and C) A549 cells were infected with 20 DV PFUs. At 36 hpi cells were
fixed, permeabilized and stained for the flavivirus E protein. DV relative
plaque area (B) and the number of infected cells per plaque (C) quantified by
ImageJ software and analyzed by one-way ANOVA. Images were acquired
with an Olympus IX83 inverted microscope.
Error bars represent mean ± SD. Results are representative of three or more
independent experiments. Statistical analyses were performed using Prism 8
(GraphPad Software). p values ****<0.0001.
458

Figure 4. ISG15 is necessary for autocrine IFNAR1-mediated control of
DV replication
(A-D) A549 cells were infected with 20 DV PFU. At 36 hpi, cells were
harvested and the indicated mRNA transcripts were quantified by RT-qPCR
(A, C and D). Cell lysates were analyzed by immunoblotting (B). Error bars
represent mean ± SD. Results are representative of two independent
experiments. Data was analyzed by unpaired t test.
(E and F) A549 ISG15 KO cells were primed with IFNα2b (100 IU/ml) for 12 h,
washed three times with DPBS and allowed to rest 12 h before infection with
DV or ZIKV at an MOI of 0.1. Shown is the percentage of cells infected at 36
hpi, as measured by E protein staining (4G2+). Error bars represent mean ±

SD. Results are representative of three independent experiments. Data was

analyzed using unpaired t test.

(G and H) A549 cells were infected with 20 DV PFUs. At 36 hpi cells were

fixed, permeabilized and stained for the flavivirus E protein. DV relative

plaque area (E) and number of infected cells per plaque (F) quantified by

ImageJ software and analyzed by one-way ANOVA. Images were acquired

with an Olympus IX83 inverted microscope.

Error bars represent mean ± SD. Results are representative of three

independent experiments. Statistical analyses were performed using Prism 8

(GraphPad Software). p values *<0.05; ***<0.001.

Figure 5. ISG15 counteracts DV IFN-I evasion.

(A) A549 WT and ISG15 KO were infected with 20 DV PFU. Cells were

harvested at the indicated time points after infection (hpi) and cell lysates

were analyzed by Western blot using STAT2 and Actin antibodies.

(B, C and D) A549 WT and ISG15 KO immunofluorescence assay (IFA) 36

hpi for cellular IFIT3 and flavivirus E protein expression (D). Percentage of

IFIT3 (B) and DV (C) positive cells per plaque were quantified by ImageJ

software and analyzed using unpaired t test with Welch's correction when

appropriate. Displayed images were acquired with a Leica DMI6000 B

microscope.

(E) A549 cells were infected with DV at MOI 0.01. At 36 hpi, cells were fixed,

permeabilized and stained for flavivirus E protein. Cell lysates were analyzed

by Western blot with the indicated antibodies before (upper panel) and after

(bottom panel) cells were sorted by fluorescence-activated cell sorting (FACS)

based on E protein expression.

U: uninfected. B: bystander. I: infected.

Error bars represent mean ± SD. Results are representative of two

independent experiments. Statistical analyses were performed using Prism 8

(GraphPad Software). p values ***<0.001.

Figure 6. USP18 expression overcomes ISG15 deficiency

(A) Flag-tag immunoprecipitation (IP) assay and Western blot analysis of

HEK293 ISG15 KO cells transfected with ZIKV NS5-FLAG, human USP18

WT, human USP18 C64A mutant or the empty vector (pcDNA3.1) plasmids, followed by IFNα2b (100 IU/ml) priming for 18 h. WCL, whole cell lysate. Results are representative of three independent experiments.

(B) STAT2 IP assay and Western blot analysis of HEK293 ISG15 KO cells transfected with the indicated plasmids, followed by IFNα2b (100 IU/ml) priming for 18 h. Results are representative of three independent experiments.

(C and D) Complementation with USP18 in A549 ISG15 KO cells. Cells were stably transfected with human USP18 or the empty vector and infected with DV at an MOI of 0.01. At 36 hpi, cells were harvested and cell lysates were analyzed by Western blot with the corresponding antibodies (C). Percentage of cells infected as measured by E protein staining (D). Error bars represent mean ± SD. Results are representative of two independent experiments. Statistical analyses were conducted using Mann-Whitney's test in Prism 8 (GraphPad Software. p value **<0.01 (D). EV: empty vector.

STAR METHODS

KEY RESOURCES TABLE

LEAD CONTACT AND MATERIAL AVAILABILITY
Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Daniel Santos Mansur (daniel.mansur@ufsc.br)

EXPERIMENTAL MODELS AND SUBJECT DETAILS

Cells and viruses
Mammalian cell lines were maintained at 37 °C under the conditions of a humidified atmosphere and 5% CO2. The human alveolar adenocarcinoma-derived A549 cells, human embryonic kidney HEK293 and the African green monkey kidney-derived Vero cells were cultured in Dulbecco's modified Eagle's medium F-12 (DMEM F12) (Gibco) supplemented with 5% fetal bovine serum (FBS) (Gibco) and streptomycin/penicillin (100 U/ml) (Gibco).

The *Aedes albopictus* mosquito-derived cell line C6/36 was maintained at 28°C in a BOD in Leibovitz's L-15 medium (L-15) (Gibco) supplemented with 10% FBS (Gibco), 0.26% tryptose phosphate broth (Sigma) and 50 µg/ml gentamicin (Sigma). A549 ISG15 KO and IFNAR1-KO have been described elsewhere (17, 52, 53). All cells were negative for mycoplasma.

Dengue virus serotype 4 (DENV-4 TVP/360 – GenBank accession number: KU513442) and Zika virus (ZV BR 2015/15261 - GenBank accession number: MF073358) stocks were propagated in C6/36 cells and titrated in Vero cells. Vesicular stomatitis virus-green fluorescent protein (VSVeGFP) (Indiana strain, Marques-JT, Plos Pathogens 2013) and Herpes simplex virus-1-green fluorescent protein (HSV-1eGFP) (SC16) viruses were propagated and titrated in Vero cells. HSV-1eGFP was a kind gift of Professor Stacey Efstathiou.

METHOD DETAIL

Single-cell RNA sequencing analysis

Processed, publicly available single-cell RNA-seq data are available through the GEO accession numbers GSE116672 and GSE110496. We downloaded processed single-cell data and metadata from the supplementary information from the respective publications (31, 32).

Then, we used CellRouter to analyze these datasets. To perform the tSNE analysis using single-cell data generated by Zanini 2018 (32), we set the parameters num.pcs=10, seed=1 and max_iter=1000 in the computeTSNE function. Next, we identified genes preferentially expressed in Uninfected, Bystander and Infected cells using a cutoff for the log2 fold change of 0.25. We used a customized script to obtain all genes containing the keywords "type I interferon" in the Gene Ontology Biological Processes (package versions: org.Hs.eg.db_3.10.0, GO.db_3.10.0). Next, we took the overlap of type I interferon genes with the genes preferentially in each condition reported above. The remaining analyses were focused on these genes.

To perform the tSNE analysis using the single-cell data generated by Zanini 2018 (31), we set the parameters num.pcs=20, seed=1 and max_iter=1000 in the computeTSNE function. We used a strategy similar to the one described

above to identify genes differentially expressed in each condition but used a
cutoff of 0.15 for this dataset. The parameter num.pcs was determined using
the elbow approach, as described in the CellRouter tutorial at
https://github.com/edroaldo/cellrouter.

CRISPR/Cas9-mediated gene editing
A549 WT and ISG15 KO cells were co-transfected with two Herc5 or Ifnar1-
targeting gRNA CRISPR/Cas9-GFP plasmids, respectively. HEK293 WT cells
were transfected with three Isg15-targeting gRNA CRISPR/Cas9-GFP
plasmids (Table S1) (Horizon Cambridge, UK). After 72 h, cells were sorted
by FACS (FACSMelody, BD) and single-cell derived clones were initially
screened by PCR genotyping. Additionally, both HERC5 and ISG15/IFNAR1
(dKO) clones were functionally tested by assessing their ISGylation profile
and expression of ISGs, respectively, after IFNα priming. Briefly, A549 WT
and HERC5 and dKO clones were primed with IFNα2b (100 IU/ml) (PBL
Assay Science) for 24 h and the expression of ISG15-conjugates and IFIT3
was analyzed by Western blot. HEK293 cells were primed with IFNα2b (1000
IU/ml) for 8 h, total RNA was isolated and Isg15 mRNA expression was
assessed by RT-qPCR.

Viral infection
A549 cells were seeded one day prior to infection in appropriate multi-well
plates. For plaque assay, a viral inoculum containing 20 plaque forming units
(PFU) of the corresponding virus was added to each well, and virus
adsorption was performed in DMEM supplemented with
penicillin/streptomycin (100 U/ml) for 90 min at 37°C. Cells were washed with
PBS to remove un-adsorbed virus, and maintained in DMEM 1.5%
carboxymethylcellulose sodium (CMC) (Sigma). Alternatively, cells were
infected with DV at the indicated MOI, as described above, and maintained in
DMEM supplemented with 1% FBS.

Titration
DV titration was performed by focus forming assay (FFA) in C6/36 cells.
Briefly, C6/36 cells were seeded (1×10^5 cells/well in 24 well plate) and after

overnight incubation were infected with a 10-fold serial dilution of virus
samples (cell culture supernatants) in L-15 with 0.26% tryptose and 25 µg/ml
of gentamicin. After 90 minutes, the inoculum was removed and a CMC
overlay media (L-15 media with 5% FCS, 0.26% tryptose, 25 µg/ml of
gentamicin and 1.6% of CMC) was added and the plates were incubated for 7
days at 28°C. After incubation, cells were washed, fixed with 3%
paraformaldehyde (PFA) (Sigma-Aldrich) and permeabilized with 0.5% triton
X100 (Sigma-Aldrich). After washing, cells were immunostained with mouse
monoclonal anti flavivirus E protein antibody 4G2 (ATCC® HB-112™, dilution
1:100), followed by goat anti-mouse immunoglobulin conjugated to alkaline
phosphatase (Promega S3721, dilution 1:7500). Focuses of infection were
revealed using NBT/BCIP reagent (Promega), following the manufacturer's
instructions and the virus titer calculated as follow: media of focus
number/inoculum volume x dilution. The results are expressed as $FFU_{C6/36}$/ml
(Gould et al., 1985).
VSVeGFP and HSV-1eGFP titrations were performed by plaque assay.
Briefly, Vero cells were seeded in 24-well plates 24 h prior to infection. Cell
monolayers were washed with PBS and inoculated with 0.3 ml of serial 10-
fold dilutions of the virus in duplicates. After 90 min adsorption at 37°C, each
well was re-suspended in DMEM 1.5% CMC. At 48 hpi, cells were fixed with
3% PFA (Sigma) for 30 min, washed 3 times with PBS and stained with 1%
crystal violet (Sigma) for 30 min at room temperature. Virus yield was
calculated and expressed as plaque forming units per ml (PFU/ml).
629
Immunofluorescence
At 24 hpi and 18 hpi, respectively, HSVeGFP and VSVeGFP infected cells
were fixed with 3% PFA for 20 min at room temperature, permeabilized with
0.5% Triton X100 in PBS for 4 min and stained with DAPI counterstain
(Molecular Probes). At 36 hpi, DV and ZIKV infected cells were fixed,
permeabilized and stained with mouse monoclonal 4G2 antibody (10 µg/ml
dilution 1:100), followed by incubation with Alexa Fluor 488 rabbit anti-mouse
IgG (H+L, Life Technologies, dilution 1:500), and DAPI counterstain.
Images were acquired with an Olympus IX83 inverted microscope. Briefly,

virus plaques, determined by eGFP or flavivirus E protein expression, were
delimited; images were converted to 16-bit and processed to be analyzed with
the ImageJ Software Cell Counter Plugin (W. S. Rasband, ImageJ, US
National Institutes of Health, Bethesda, MD; http://rsb.info.nih.gov/ij/, 1997–
2006).

For confocal analysis, A549 cells grown on glass coverslips were mock
infected or infected with 10 DV PFU. At 36 hpi, cells were fixed and
permeabilized. Following washes with PBS, cells were stained with mouse
monoclonal anti-E protein (4G2, dilution 1:100) and rabbit polyclonal anti-
IFIT3 (Proteintech, 15201-1-AP, dilution 1:200) for 1 h at room temperature.
The cells were washed with PBS and stained with secondary antibodies Alexa
Fluor 488 goat anti-mouse IgG (H+L, Life Technologies, dilution 1:500) and
Alexa 568 goat anti-rabbit IgG (H+L, Life Technologies, dilution 1:500) and
DAPI counterstain (Molecular Probes). Cells were washed and coverslips
mounted using Prolong antifade reagent (Invitrogen). Z-stack and max
intensity projection images were generated with a Leica DMI6000 B confocal
microscope and Leica Application Suite X software for image analysis (Leica
Microsystems).

Flow cytometry and cell sorting
Cells were fixed with 2% PFA in PBS, washed twice with PBS and
permeabilized with 0.5% saponin in 1% BSA in PBS. Anti-flavivirus E protein
mAb 4G2 was conjugated to Alexa Fluor 488 5-SDP (Life Technologies)
following the manufacturer's instructions. Cells were incubated with FITC-
conjugated 4G2 (dilution 1:1000) in permeabilization buffer for 40 minutes at
room temperature, washed once and resuspended in FACS buffer. The cell
suspensions were analyzed by flow cytometry on a FACSVerse instrument
(BD Biosciences) and analyzed using FlowJo V10 software (BD). Cell sorting
experiments were performed on a FACSMelody cell sorter (BD Biosciences).

RT-qPCR
vRNA was isolated by using QIAamp viral RNA Mini Kit (Qiagen). Intracellular
total RNA was isolated with TRIZOL (Thermo Life) following manufacturer's

instructions. A total of 1 µg was reverse transcribed using the High capacity

cDNA reverse transcription kit (Applied Biosystems). Quantitative PCRs

(qPCRs) were performed with GoTaq ® qPCR Master mix (Promega) using

StepOne Plus real time PCR system (Applied Biosystems). VSVeGFP and

18S mRNA were used as control housekeeping genes. Amounts of DV or ISG

mRNA were calculated by using the ΔΔCT method. Primers specific the

mRNA analyzed are listed in Table S1.

Western blot

Human antibodies used for immunoblot were as follows: mouse mAb to β-

actin (Abcam, ab6276, dilution 1:4000), rabbit mAb to IFIT1 (Abcam,

ab137632, dilution 1:1000), mouse mAb to pSTAT1(Y701) [M135] (Abcam,

ab29045, dilution 1:1000), mouse mAb to STAT1 (Abcam, ab3987, dilution

1:1000), rabbit pAb to IFIT3 (ProteinTech, 15201-1-AP, dilution 1:1000), rabbit

mAb to USP18 [D4E7] (Cell Signaling Technologies, 4813, dilution 1:1000),

rabbit mAb to STAT2 [D9J7L] (Cell Signaling Technologies, 72604, dilution

1:1000), mouse mAb to ISG15 (R&D System, MAB4845, dilution 1:1000),

home-made mouse mAb anti-GFP (67, 68) (dilution 1:1000) and mouse mAb

to FLAG tag (Sigma, F3165, dilution 1:500).

Cells were treated as indicated, wash two times with ice-cold PBS and then

lysed in RIPA buffer [50 mM tris-HCl (pH 7.5), 150 mM NaCl, 1% Triton X100,

0.5% sodium deoxycholate, 0.1% SDS, 5 mM EDTA] supplemented with 1x

protease inhibitors (Mini Protease Inhibitor Tablets, Roche). Total protein

concentration was determined by BSA Protein Assay Kit (Thermo Life). For

western blotting, 20 µg of total protein were prepared in dithiothreitol-

containing Laemmli sample buffer, separated and transferred to a

nitrocellulose blotting membrane (GE Healthcare Amersham). After transfer,

the membrane was blocked with 5% nonfat dry milk in TBS 0.1% Tween 20

(TBST) for 1 h at room temperature. Membrane was incubated with primary

Abs diluted in 2% BSA in TBST at 4°C with gentle shaking overnight.

Membrane was washed three times with TBST and then incubated with the

appropriate secondary HRP-linked antibody for 1 h at room temperature.

Membranes were washed and covered with ECL developing solution

(Pierce™ ECL WB substrate, Thermo Fisher Scientific).

Plasmids and transfections

Herc5 and Ifnar1 sgRNA/Cas9/GFP plasmids were provided by Horizon (Cambridge, UK). sgRNA sequences are described in Supplementary Table 1. Expression plasmid for ISG15 was described elsewhere (17). Pmax^{TM} GFP expression vector was acquired from Lonza (cat numb #D-00061). Expression plasmid for ZIKV NS5 was generated by amplifying the NS5 coding sequence (amino acids 2521-3423 in the polyprotein) from a previously described plasmid-based ZIKV reverse genetic system (69) using primers containing an N-terminal FLAG tag and inserted into pcDNA3.1. Expression plasmids for human USP18 WT and USP18 C64A mutant were kindly provided by Dr Carsten Münk (70). Transfections were performed with FuGene6 (Promega), following the manufacturer's instructions. Stably transfected cells were selected with geneticin (500 µg/ml) (Sigma).

Immunoprecipitation

For Stat2 co-immunoprecipitation assays, cells were lysed in RIPA buffer composed of 20 mM Tris-HCl pH 7.4, 150 mM NaCl,10 mM CaCl2, 0.1% v/v Triton X-100, 10% v/v glycerol and complete protease inhibitor cocktail (Mini Protease Inhibitor Tablets, Roche). The supernatant was separated by centrifugation at 12.000 g at 4 °C for 10 min and incubated with STAT2 antibody (2 µg) (Santa Cruz Biotechnology, 514193) for 3 h at 4°C with gentle shaking. Complexes were precipitated with protein A/G Plus-agarose (Santa Cruz Biotechnology), washed with TBS and resuspended in SDS sample buffer. Immunoprecipitates were subjected to SDS-PAGE and western blotting, as described above.

FLAG-tagged proteins were immunoprecipitated with anti-FLAG M2-agarose (Sigma), following the manufacturer's instructions.

All assays were performed three times and representative blots are presented.

QUANTIFICATION AND STATISTICAL ANALYSIS

Details concerning the statistical analysis methods are provided in each figure legend. Briefly, all data were analyzed using GraphPad Prism 8 software and

740 were shown as mean and the standard deviation (SD). Statistical significance
741 was determined by Welch's t test or one-way ANOVA for plaque size
742 analyses, unpaired t test for virus multiple-step growth curve, cellular mRNA
743 quantification and percentage of cells infected. Statistical significance is
744 indicated by ∗, $p < 0.05$; ∗∗, $p < 0.01$; ∗∗∗, $p < 0.001$.
745

SUPPLEMENTAL INFORMATION

Figure S1
(A) tSNE was used to visualize the single-cell global transcriptome data. Blue
dots represent uninfected cells, derived from healthy donors. Beige dots
represent bystander cells and red dots represent infected cells, both derived
from patients infected with DV.
(B) Differential expression of ISGs in HuH7 cells infected with DV. The Gene
Ontology term "type one interferon" was used to filter the results from the
single-cell RNA sequencing.
(C) Single cell ISG expression variability in uninfected, bystander and
infected. ISGylation related genes are underlined. Blue bar represents
uninfected cells. Beige bar represents bystander cells and red bar represents
infected cells, both derived from cells exposed to DV.
(D) Violin plot representing the expression of ISGylation family members in
uninfected [U], bystander [B] and infected [I] HuH7 cell line.
762

Figure S2. ISG15 does not restrict HSV-1 and VSV spread
(A and B) A549 cells were infected with 20 HSV-1eGFP PFU. At 24 hpi, cells
were fixed and stained with DAPI counterstain. HSV-1eGFP relative plaque
area (A) and number of cells per plaque (B). Images were acquired with an
Olympus IX83 inverted microscope and quantified by ImageJ software.
(C and D) A549 cells were infected with 20 VSVeGFP PFU. At 18 hpi, cells
were fixed and stained with DAPI counterstain. VSVeGFP relative plaque
area (C) and number of cells per plaque (D). Images were acquired with an
Olympus IX83 inverted microscope and quantified by ImageJ software.

Error bars represent mean ± SD. Results are representative of three or more

independent experiments. Statistical analyses were conducted using Mann-

Whitney's test in Prism 8 (GraphPad Software).

Figure S3. Characterization of knockout cell lines

(A) Agarose gel electrophoresis of A549 WT and HERC5 KO PCR products

using primers surrounding CRISPR/Cas9 ISG15 sgRNA guides editing region.

PCR product size: WT: 310 bp; HERC5 KO: ~250 bp. (*) indicates PCR

unspecific band. WT: A549 WT cell line; KO: A549 HERC5 KO; B: blank

(B) Clonal isolates of A549 cells were immunoblotted for IFIT3 after 24 h

treatment with IFNα2b (100 IU/ml) (+, control IFIT3 -proficient cells; 1–9,

different clonal isolates after limiting dilution).

(C) Agarose gel electrophoresis of HEK293 WT and ISG15 KO PCR products

using primers surrounding CRISPR/Cas9 ISG15 sgRNA guides editing region.

PCR product size: WT: 1016bp; ISG15 KO: ~180bp. (*) indicates PCR

unspecific band. WT: HEK293 WT cell line; KO: HEK293 ISG15 KO; B: blank

(D) HEK293 WT and ISG15 KO cells were stimulated with IFNα2b (1000

IU/ml) for 8 h. Cells were harvested and total RNA was isolated. Isg15 mRNA

was analyzed by RT-qPCR.

(E) HEK293 ISG15KO PCR product was cloned into pGEM vector and

sequenced by Sanger method. Nucleotide sequence was aligned with Isg15

reference sequence retrieved from GenBank (NM_005101) and translated

into primary amino acid sequence. (.) indicates the same sequence; (-) gap;

(*) stop codon.

Figure S4. A549 ISG15 KO cell line exhibits sustained ISG expression

(A) A549 cells were primed with IFNα2b (100 IU/ml) for 12 h, washed three

times with DPBS and allowed to rest. Cells were harvested at the indicated

time point and cell lysates were analyzed by Western blot with the

corresponding antibodies.

Results are representative of two independent experiments.

Figure S5. Sorting of infected cells

(A and B) A549 cells were infected with DV at MOI 0.01. Representative FACS profile and DV mRNA qPCR of WT (A) and ISG15-KO (B) cells sorted by flavivirus E protein expression (4G2, FITC-A axis).
B: bystander. I: infected

REFERENCES

1. D. S. Mansur, G. L. Smith, B. J. Ferguson, Intracellular sensing of viral DNA by the innate immune system. *Microbes Infect.* **16**, 1002–1012 (2014).

2. A. Gebhardt, B. T. Laudenbach, A. Pichlmair, Discrimination of Self and Non-Self Ribonucleic Acids. *J. Interferon Cytokine Res.* **37**, 184–197 (2017).

3. J. W. Schoggins, Interferon-Stimulated Genes: What Do They All Do? *Annu Rev Virol* (2019) https:/doi.org/10.1146/annurev-virology-092818-015756.

4. A. García-Sastre, Ten Strategies of Interferon Evasion by Viruses. *Cell Host Microbe* **22**, 176–184 (2017).

5. T. C. Pierson, M. S. Diamond, The emergence of Zika virus and its new clinical syndromes. *Nature* **560**, 573–581 (2018).

6. A. D. T. Barrett, The reemergence of yellow fever. *Science* **361**, 847–848 (2018).

7. S. Bhatt, *et al.*, The global distribution and burden of dengue. *Nature* **496**, 504–507 (2013).

8. D. S. Shepard, E. A. Undurraga, Y. A. Halasa, J. D. Stanaway, The global economic burden of dengue: a systematic analysis. *Lancet Infect. Dis.* **16**, 935–941 (2016).

9. L. Miorin, A. M. Maestre, A. Fernandez-Sesma, A. García-Sastre, Antagonism of type I interferon by flaviviruses. *Biochem. Biophys. Res. Commun.* **492**, 587–596 (2017).

10. A. E. Ngono, S. Shresta, Immune Response to Dengue and Zika. *Annu. Rev. Immunol.* **36**, 279–308 (2018).

11. A. Grant, *et al.*, Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. *Cell Host Microbe* **19**, 882–890 (2016).

12. M. Laurent-Rolle, *et al.*, The interferon signaling antagonist function of yellow fever virus NS5 protein is activated by type I interferon. *Cell Host Microbe* **16**, 314–327 (2014).

13. S. Chen, Z. Wu, M. Wang, A. Cheng, Innate Immune Evasion Mediated by Flaviviridae Non-Structural Proteins. *Viruses* **9** (2017).

14. J. Morrison, *et al.*, Dengue virus co-opts UBR4 to degrade STAT2 and antagonize type I interferon signaling. *PLoS Pathog.* **9**, e1003265 (2013).

15. J. Ashour, *et al.*, Mouse STAT2 restricts early dengue virus replication. *Cell Host Microbe* **8**, 410–421 (2010).

16.   P. F. Dos Santos, D. S. Mansur, Beyond ISGlylation: Functions of Free
Intracellular and Extracellular ISG15. *J. Interferon Cytokine Res.* **37**, 246–253
(2017).

17.   P. F. Dos Santos, *et al.*, ISG15-Induced IL-10 Is a Novel Anti-Inflammatory
Myeloid Axis Disrupted during Active Tuberculosis. *J. Immunol.* **200**, 1434–1442
(2018).

18.   C. D. Swaim, A. F. Scott, L. A. Canadeo, J. M. Huibregtse, Extracellular ISG15
Signals Cytokine Secretion through the LFA-1 Integrin Receptor. *Mol. Cell* **68**,
581–590.e5 (2017).

19.   D. Bogunovic, *et al.*, Mycobacterial disease and impaired IFN-γ immunity in
humans with inherited ISG15 deficiency. *Science* **337**, 1684–1688 (2012).

20.   M. Delgobo, *et al.*, An evolutionary recent IFN-IL-6-CEBP axis is linked to
monocyte expansion and tuberculosis severity in humans
https:/doi.org/10.1101/514943.

21.   B. Skaug, Z. J. Chen, Emerging role of ISG15 in antiviral immunity. *Cell* **143**,
187–190 (2010).

22.   A. Dastur, S. Beaudenon, M. Kelley, R. M. Krug, J. M. Huibregtse, Herc5, an
interferon-induced HECT E3 enzyme, is required for conjugation of ISG15 in
human cells. *J. Biol. Chem.* **281**, 4334–4338 (2006).

23.   J. J. Y. Wong, Y. F. Pung, N. S.-K. Sze, K.-C. Chin, HERC5 is an IFN-induced
HECT-type E3 protein ligase that mediates type I IFN-induced ISGylation of
protein targets. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 10735–10740 (2006).

24.   Y.-C. Perng, D. J. Lenschow, ISG15 in antiviral immunity and beyond. *Nat. Rev.*
Microbiol. **16**, 423–439 (2018).

25.   K.-I. Arimoto, *et al.*, STAT2 is an essential adaptor in USP18-mediated
suppression of type I interferon signaling. *Nat. Struct. Mol. Biol.* **24**, 279–289
(2017).

26.   M. P. Malakhov, O. A. Malakhova, K. I. Kim, K. J. Ritchie, D.-E. Zhang, UBP43
(USP18) specifically removes ISG15 from conjugated proteins. *J. Biol. Chem.*
277, 9976–9981 (2002).

27.   O. Malakhova, M. Malakhov, C. Hetherington, D.-E. Zhang, Lipopolysaccharide
activates the expression of ISG15-specific protease UBP43 via interferon
regulatory factor 3. *J. Biol. Chem.* **277**, 14703–14711 (2002).

28.   X. Zhang, *et al.*, Human intracellular ISG15 prevents interferon-α/β over-
amplification and auto-inflammation. *Nature* **517**, 89–93 (2015).

29.   S. D. Speer, *et al.*, ISG15 deficiency and increased viral resistance in humans
but not mice. *Nat. Commun.* **7**, 11496 (2016).

30.   D. J. Lenschow, *et al.*, IFN-stimulated gene 15 functions as a critical antiviral
molecule against influenza, herpes, and Sindbis viruses. *Proc. Natl. Acad. Sci.*
U. S. A. **104**, 1371–1376 (2007).

31.   F. Zanini, S.-Y. Pu, E. Bekerman, S. Einav, S. R. Quake, Single-cell
transcriptional dynamics of flavivirus infection. *Elife* **7** (2018).

32. F. Zanini, *et al.*, Virus-inclusive single-cell RNA sequencing reveals the
molecular signature of progression to severe dengue. *Proc. Natl. Acad. Sci. U.*
S. A. **115**, E12363–E12369 (2018).

33. I. Ramos, *et al.*, Innate immune response to influenza virus at single-cell
resolution in human epithelial cells revealed paracrine induction of interferon
lambda 1. *J. Virol.* (2019) https://doi.org/10.1128/JVI.00559-19.

34. P. Keskinen, *et al.*, Impaired antiviral response in human hepatoma cells.
Virology **263**, 364–375 (1999).

35. B. Schmid, *et al.*, Live Cell Analysis and Mathematical Modeling Identify
Determinants of Attenuation of Dengue Virus 2'-O-Methylation Mutant. *PLoS*
Pathog. **11**, e1005345 (2015).

36. H. Roth, *et al.*, Flavivirus Infection Uncouples Translation Suppression from
Cellular Stress Responses. *MBio* **8** (2017).

37. K. Wang, *et al.*, Interferon-stimulated TRIM69 interrupts dengue virus replication
by ubiquitinating viral nonstructural protein 3. *PLoS Pathog.* **14**, e1007287
(2018).

38. J. N. Whelan, Y. Li, R. H. Silverman, S. R. Weiss, Zika Virus Production Is
Resistant to RNase L Antiviral Activity. *J. Virol.* **93** (2019).

39. R. C. Gullberg, *et al.*, Stearoly-CoA desaturase 1 differentiates early and
advanced dengue virus infections and determines virus particle infectivity. *PLoS*
Pathog. **14**, e1007261 (2018).

40. J. Zhang, *et al.*, Flaviviruses Exploit the Lipid Droplet Protein AUP1 to Trigger
Lipophagy and Drive Virus Production. *Cell Host Microbe* **23**, 819–831.e5
(2018).

41. C. Zhao, C. Denison, J. M. Huibregtse, S. Gygi, R. M. Krug, Human ISG15
conjugation targets both IFN-induced and constitutively expressed proteins
functioning in diverse cellular pathways. *Proc. Natl. Acad. Sci. U. S. A.* **102**,
10200–10205 (2005).

42. C. Zhao, T.-Y. Hsiang, R.-L. Kuo, R. M. Krug, ISG15 conjugation system targets
the viral NS1 protein in influenza A virus-infected cells. *Proc. Natl. Acad. Sci. U.*
S. A. **107**, 2253–2258 (2010).

43. Y. J. Kim, *et al.*, Consecutive Inhibition of ISG15 Expression and ISGylation by
Cytomegalovirus Regulators. *PLoS Pathog.* **12**, e1005850 (2016).

44. R. González-Sanz, *et al.*, ISG15 Is Upregulated in Respiratory Syncytial Virus
Infection and Reduces Virus Growth through Protein ISGylation. *J. Virol.* **90**,
3428–3438 (2016).

45. T. Hishiki, *et al.*, Interferon-mediated ISG15 conjugation restricts dengue virus 2
replication. *Biochem. Biophys. Res. Commun.* **448**, 95–100 (2014).

46. V. François-Newton, *et al.*, USP18-based negative feedback control is induced
by type I and type III interferons and specifically inactivates interferon α
response. *PLoS One* **6**, e22200 (2011).

47. N. A. Dalrymple, V. Cimica, E. R. Mackow, Dengue Virus NS Proteins Inhibit

RIG-I/MAVS Signaling by Blocking TBK1/IRF3 Phosphorylation: Dengue Virus
Serotype 1 NS4A Is a Unique Interferon-Regulating Virulence Determinant.
MBio **6**, e00553–15 (2015).

48. J. L. Muñoz-Jordán, *et al.*, Inhibition of alpha/beta interferon signaling by the
NS4B protein of flaviviruses. *J. Virol.* **79**, 8004–8013 (2005).

49. C. Daly, N. C. Reich, Double-stranded RNA activates novel factors that bind to
the interferon-stimulated response element. *Mol. Cell. Biol.* **13**, 3756–3764
(1993).

50. C. Daly, N. C. Reich, Characterization of specific DNA-binding factors activated
by double-stranded RNA as positive regulators of interferon alpha/beta-
stimulated genes. *J. Biol. Chem.* **270**, 23739–23746 (1995).

51. B. K. Weaver, K. P. Kumar, N. C. Reich, Interferon regulatory factor 3 and
CREB-binding protein/p300 are subunits of double-stranded RNA-activated
transcription factor DRAF1. *Mol. Cell. Biol.* **18**, 1359–1368 (1998).

52. E. S. Guimarães, *et al.*, Cyclic Dinucleotides Trigger STING-Dependent
Unfolded Protein Response That Favors Bacterial Replication. *J. Immunol.* **202**,
2671–2681 (2019).

53. K. H. Antunes, *et al.*, Microbiota-derived acetate protects against respiratory
syncytial virus infection through a GPR43-type 1 interferon response. *Nat.*
Commun. **10**, 3273 (2019).

54. J. Ashour, M. Laurent-Rolle, P.-Y. Shi, A. García-Sastre, NS5 of dengue virus
mediates STAT2 binding and degradation. *J. Virol.* **83**, 5408–5418 (2009).

55. M. Jones, *et al.*, Dengue virus inhibits alpha interferon signaling by reducing
STAT2 expression. *J. Virol.* **79**, 5414–5420 (2005).

56. U. Rand, *et al.*, Multi-layered stochasticity and paracrine signal propagation
shape the type-I interferon response. *Mol. Syst. Biol.* **8**, 584 (2012).

57. W. M. Schneider, M. D. Chevillotte, C. M. Rice, Interferon-stimulated genes: a
complex web of host defenses. *Annu. Rev. Immunol.* **32**, 513–545 (2014).

58. K. R. Loeb, A. L. Haas, The interferon-inducible 15-kDa ubiquitin homolog
conjugates to intracellular proteins. *J. Biol. Chem.* **267**, 7806–7813 (1992).

59. L. A. Durfee, N. Lyon, K. Seo, J. M. Huibregtse, The ISG15 conjugation system
broadly targets newly synthesized proteins: implications for the antiviral function
of ISG15. *Mol. Cell* **38**, 722–732 (2010).

60. H.-X. Shi, *et al.*, Positive regulation of interferon regulatory factor 3 activation by
Herc5 via ISG15 modification. *Mol. Cell. Biol.* **30**, 2424–2436 (2010).

61. M.-J. Kim, S.-Y. Hwang, T. Imaizumi, J.-Y. Yoo, Negative feedback regulation of
RIG-I-mediated antiviral signaling by interferon-induced ISG15 conjugation. *J.*
Virol. **82**, 1474–1483 (2008).

62. F. Okumura, *et al.*, Activation of double-stranded RNA-activated protein kinase
(PKR) by interferon-stimulated gene 15 (ISG15) modification down-regulates
protein translation. *J. Biol. Chem.* **288**, 2839–2847 (2013).

63.   J. D'Cunha, E. Knight Jr, A. L. Haas, R. L. Truitt, E. C. Borden,
Immunoregulatory properties of ISG15, an interferon-induced cytokine. *Proc.*
Natl. Acad. Sci. U. S. A. **93**, 211–215 (1996).

64.   S. W. Werneke, *et al.*, ISG15 is critical in the control of Chikungunya virus
infection independent of UbE1L mediated conjugation. *PLoS Pathog.* **7**,
e1002322 (2011).

65.   F. Vuillier, Z. Li, P.-H. Commere, L. T. Dynesen, S. Pellegrini, USP18 and ISG15
coordinately impact on SKP2 and cell cycle progression. *Sci. Rep.* **9**, 4066
(2019).

66.   S. Dabo, *et al.*, Inhibition of the inflammatory response to stress by targeting
interaction between PKR and its cellular activator PACT. *Sci. Rep.* **7**, 16129
(2017).

67.   C. Zanluca, G. A. C. A. Mazzarotto, J. Bordignon, C. N. Duarte Dos Santos,
Development, characterization and application of monoclonal antibodies against
Brazilian Dengue virus isolates. *PLoS One* **9**, e110620 (2014).

68.   G. F. Silveira, *et al.*, Human T Lymphocytes Are Permissive for Dengue Virus
Replication. *J. Virol.* **92** (2018).

69.   M. Mutso, *et al.*, Reverse genetic system, genetically stable reporter viruses and
packaged subgenomic replicon based on a Brazilian Zika virus isolate. *J. Gen.*
Virol. **98**, 2712–2724 (2017).

70.   E. Osei Kuffour, *et al.*, USP18 (UBP43) Abrogates p21-Mediated Inhibition of
HIV-1. *J. Virol.* **92** (2018).

Figure 1-Espada *et al.*

Figure 2-Espada *et al.*

Figure 3-Espada *et al.*

Figure 4-Espada *et al.*

Figure 5-Espada *et al.*

Figure 6-Espada *et al.*