TITLE

ISG15/USP18/STAT2 is a molecular hub regulating autocrine IFN I-mediated

control of Dengue and Zika virus replication

Constanza Eleonora Espada[1], Edroaldo Lummertz da Rocha[1], Taissa Ricciardi-

Jorge[1], Adara Aurea dos Santos[1], Zamira Guerra Soares[1], Greicy Brisa Malaquias

Dias[1], Daniel de Oliveira Patricio[1], Edgar Gonzalez-Kozlova[1], Paula Fernandes dos

Santos[1], Juliano Bordignon[2#], Thomas J. Sanford[3], Teodoro Fajardo[3], Trevor R.

Sweeney[3], André Báfica[1], Daniel Santos Mansur[1*]

[1]Laboratório de Imunobiologia, Departamento de Microbiologia, Imunologia e

Parasitologia, Centro de Ciências Biológicas, Universidade Federal de Santa

Catarina, Santa Catarina CEP 88040-900, Brazil.

[2]Laboratório de Virologia Molecular, Instituto Carlos Chagas, ICC/Fiocruz-PR,

Curitiba, Paraná, CEP 81350-010, Brazil.

[3]Division of Virology, Department of Pathology, University of Cambridge,

Addenbrooke's Hospital, Hills Road, Cambridge, CB2 0QQ, UK.

#Current address: Instituto Carlos Chagas, ICC/Fiocruz-PR, Curitiba, Paraná, CEP

81350-010, Brazil.

[*]Correspondence: daniel.mansur@ufsc.br

SUMMARY

The establishment of a virus infection is the result of the pathogen's ability to replicate in a hostile environment generated by the host's immune system. Here, we found that ISG15 restricts Dengue and Zika viruses' replication through the stabilization of its binding partner USP18. ISG15 expression was necessary to control DV replication driven by both autocrine and paracrine type one interferon (IFN-I) signaling. Moreover, USP18 competes with NS5-mediated STAT2 degradation, a major mechanism for establishment of flavivirus infection. Strikingly, reconstitution of USP18 in ISG15-deficient cells was sufficient to restore the cells' immune response and restrict virus growth, suggesting that the IFNAR-mediated ISG15 activity is also antiviral. Our results reveal a novel layer of complexity in the virus/host interaction interface and suggest that NS5 has a narrow window of opportunity to degrade STAT2, thereby suppressing host's IFN-I mediated response and promoting virus replication.

IMPORTANCE

Disease is an emergent property that results from a microorganism's ability to replicate in a given host and the latter's immune response. Here we describe how the immunoregulatory function of ISG15 (an interferon stimulated gene) affects Dengue and Zika virus replication by occupying a niche used by the virus non-structural protein 5 (NS5) to evade host's immunity. In the absence of ISG15, NS5 efficiently degrades a main signaling hub of innate immunity, Signal transducer and activator of transcription 2 (STAT2), leading to cell immune suppression and

consequently virus growth. This sheds light into how flaviviruses intimately interact

with the host immune system and could lead to a host-based therapy target.

KEYWORDS

Dengue virus, Zika virus, ISG15, USP18, type one interferon, ISGylation, antiviral

response, immune evasion, NS5, STAT2

INTRODUCTION

Cells detect infection by recognizing molecular patterns derived from pathogen's

constituents (PAMPs) or cell damage (DAMPs). During viral infection, nucleic acid is

a major signal that triggers the innate immune response, inducing a type one

interferon (IFN-I)-mediated antiviral state (1, 2). IFN-I binds to its cognate receptor

and activates the JAK/STAT pathway, leading to expression of hundreds of

interferon stimulated genes (ISGs) that make the intracellular environment hostile to

viral replication in infected and proximal cells (3). The evolutionary arms race

between viruses and its hosts led to evolution of immune evasion mechanisms that

are crucial for successful viral replication. Considering IFN-I's importance in infection

control, many viral immunomodulatory proteins target this signaling pathway (4).

Several flaviviruses, such as dengue virus (DV), Zika virus (ZIKV) and yellow fever

virus (YFV), have emerged and re-emerged over recent years and are the leading

cause of human arbovirus infection (5, 6). DV alone infects nearly 400 million people

every year (7) with extensive health and economic burden (8). A requirement for

effective flavivirus emergence is the ability to counteract the human immune system.

The compact flavivirus genome encodes seven non-structural proteins that are

responsible for viral replication and immune evasion. Six of these proteins are not

secreted implying that intracellular pathways are central targets for evasion (9–13).

For instance, DV non-structural protein 5 (NS5), which is the viral RNA-dependent-

RNA-polymerase (RdRp) and a methyltransferase, mediates STAT2 degradation by

facilitating its interaction with UBR4, leading to its ubiquitination and subsequent

proteasomal targeting (14). This evasion pathway is functional in humans but not

mice due to differences in the amino acid sequence of human and murine STAT2

(15).

ISG15 is an intracellular and secreted ubiquitin-like protein that has three described

functions. Extracellular ISG15 acts as a cytokine, leading to the expression of IFNγ

and IL-10 in diverse immune cells (16–18). It has been suggested that humans

lacking ISG15 have severe mycobacterial disease due to deficiency in IFNγ

production by NK cells (19). Moreover, ISG15 mRNA is highly expressed in active

tuberculosis and strongly correlates with disease severity (17, 20).

ISG15 is conjugated to other proteins through a three-step ubiquitination-like process

(21) in which the main ligase for ISGylation is the HECT domain and RCC1-like

domain-containing protein 5 (HERC5) (22, 23). ISGylated proteins are affected in

several different ways, such as increased or reduced stability and activity. ISG15 can

also be conjugated to viral proteins, impacting their function, and therefore belong in

the plethora of ISGs with a direct antiviral function (3, 24).

The third and more recently described role of ISG15 is its IFN-I modulatory function.

Non-conjugated ISG15 binds and stabilizes the ISG USP18, a protease that

negatively regulates IFN-I signaling and also serves as a deISGylation protein (25–

27). Correspondingly, individuals lacking ISG15 are prone to severe

interferonopathies due to decreased USP18 function and increased IFN-I signaling.

Interestingly, ISG15-deficient patients do not have enhanced susceptibility to viruses

suggesting ISG15 is not necessary to control ubiquitous viral infections in vivo (28).

In contrast to the indirect role of ISG15 in negative regulation of IFN-I signaling

through the USP18 axis in humans (28, 29), murine ISG15 itself blocks replication of

human viruses such as Influenza and HSV-1 (30). These findings implicate ISG15 as

an important molecule inhibiting IFN-I-mediated actions but also suggest that ISG15

may mediate host cell intrinsic mechanisms triggered by viruses. However, how

ISG15 bridges these apparently two paradoxical phenomena is unclear. Specifically,

it is possible that ISG15 directly regulates proteins exploited by viruses during early

intracellular infection events.

Here, we observed that, unlike several other ISGs, ISG15 is highly expressed in

infected cells containing the DV genome. Furthermore, independently of its

ISGylation function, ISG15 restricts flavivirus replication primarily in the infected cell

by stabilizing USP18, which in turn competes with viral NS5 for binding to STAT2.

Our results suggest that flaviviruses exploit an ISG15-mediated IFN-I regulatory

mechanism to evade innate immunity and enable replication.

RESULTS

ISG15 is expressed in DV-infected cells

The cell is a fundamental unit for viral infection control and developments in single-

cell sequencing technology have enabled examination of host-pathogen interactions

in great detail. Recently, Zanini and colleagues generated single-cell RNA

sequencing data from human cells (PBMCs and the HuH7 hepatoma cell line)

independently infected with DV (31, 32). We re-analyzed available single-cell

transcriptomic data dividing cells into three categories: uninfected, infected and

bystander. Here we define uninfected cells as those derived from healthy donors,

bystander cells as those derived from an infected patient or have been exposed to

the virus but did not have the viral RNA detected and infected cells as those in which

viral genome was detected. We used t-Distributed Stochastic Neighbor Embedding

(tSNE) analysis to visualize cell-to-cell relationships in space of reduced

dimensionality. As reported previously (31, 32), global cellular mRNA expression

profiling was not sufficient to separate infected or bystander from uninfected cells,

suggesting a high variability of gene expression in these samples (Figure 1A). As

IFN-I are key elements in controlling infection, we filtered the results of the

differential gene expression analysis using the gene ontology (GO) term for "type

one interferon". The Venn diagram in figure 1B shows that from the 394 differentially

expressed (DE) genes in peripheral blood mononuclear cells (PBMCs) of patients

infected with DV or healthy donors, 37 were ISGs (IFN-I GO). ISG15, UBE2L6,

HERC5 and USP18, members of the ISGylation pathway, were differentially

expressed during DV infection (Figure 1C underlined). Interestingly, in contrast to

other single cell experiments using Influenza virus as a model where all the

members of the ISGylation pathway seem to be expressed at similar levels in both

infected and bystander cells (33), HERC5 was the only member of the ISGylation

pathway with a higher expression level in bystander cells (Figure 1C and D). In the

data set derived from the HuH7 cell line, ISG15 was the only canonical antiviral

protein expressed in DV genome-containing cells (data not shown). This is in

agreement with previous reports that the HuH7 cell line does not produce IFN-I upon

viral infection (34–36) and could explain the high number of infected cells (data not

shown) in comparison with the number of infected PMBCs (Figure 1A). From

PBMCs, NK, monocytes and B cells were the infected cells with higher ISG15

expression and similar to the data from Zanini and colleagues (32) B cells and

monocytes being proportionately the most infected cells (Figure 1E). These results

show that in contrast to most ISGs, ISG15 and other components of the ISGylation

pathway are enriched in cells where the DV genome was present.

ISG15 restricts DV and ZIKV replication

The enrichment of ISG15-related genes at a single-cell level led us to investigate

how ISG15 might impact flavivirus replication in a human cell. We used an A549 cell

line lacking ISG15, previously generated using CRISPR/Cas9 (17). A549 cells were

chosen due to their ability to support flavivirus replication and most importantly to

produce and respond to IFN-I (37–40).

ISG15-deficient cells were more susceptible to DV infection as shown by an

increased foci size (Figure 2A and B) and number of infected cells per foci (Figure

2C). In addition, we determined the kinetics of DV replication and dissemination by

using a low multiplicity of infection (MOI), to allow for viral spread through secondary

infection events. Percentages of infected cells over time (Figure 2D), relative DV

Pre-Membrane RNA quantification in the supernatant (Figure 2E) and infectious

particle formation (Figure 2F) were also increased in the absence of ISG15. DV

infection at 4 ^{o}C for 2 hours followed by relative intracellular DV genome

quantification indicated no differences in viral RNA between WT and knockout cells

up to 48 hours post-infection (Figure 2G), indicating that ISG15 plays a role in the

DV life cycle at a stage after viral entry. Importantly, reconstitution of ISG15

expression in knockout cells led to phenotypic reversion (Figure 2H and I),

confirming that the effects observed in our experiments were caused by depletion of

ISG15.

Finally, lack of ISG15 expression led to an increase in foci size in cells infected with

a phylogenetically related virus, such as ZIKV (Figure 2J and K) but not with HSV-1

or VSV (data not shown). These results are in line with Speer and colleagues' data

where cells isolated from humans deficient for ISG15 do not have enhanced

susceptibility to HSV-1 or VSV (29). Altogether these results suggest a specific role

for ISG15 in the regulation of flavivirus replication and dissemination.

ISGylation deficiency does not affect DV spread

ISG15 is an IFN-I-inducible ubiquitin-like molecule and can be conjugated to target

proteins by HERC5, an ISG15 ligase also induced by IFN (23, 41). Of note,

ISGylation of host or viral proteins was reported to inhibit replication of several

viruses such as influenza (IAV) (42), human cytomegalovirus (HCMV) (43) and

human respiratory syncytial virus (RSV) (44). Moreover, DV proteins were also

shown to be ISGylated (45). To determine whether ISGylation could be involved in

DV restriction, we generated HERC5-deficient A549 cells using CRISPR/Cas9 (data

not shown) and accordingly, A549 HERC5 null cells were not able to perform

ISGylation after IFN-I treatment (Figure 3A). In contrast to ISG15 null cells, we did

not observe differences between WT and HERC5-deficient cells when we evaluated

both foci area (Figure 3B) and number of infected cells per foci following infection

with DV (Figure 3C). Therefore, ISGylation is not sufficient to inhibit DV replication.

ISG15 is necessary for autocrine IFNAR1-mediated control of DV replication

ISG15 and its binding partner USP18 are crucial for IFN-I pathway down-regulation,

which is pivotal for infection control and immune-regulation in humans (28, 29, 46).

The A549 ISG15-KO cell line exhibited a lower expression of USP18 and sustained

ISG expression, as exemplified by IFIT3, after IFN-I stimulation (data not shown),

reproducing the phenotype observed in cells isolated from humans lacking ISG15

(29).

While our single-cell RNA-seq analysis showed that both ISG15 and USP18 are

upregulated in PBMCs containing DV RNA, there was an asynchrony of ISGs being

expressed in infected versus bystander cells, with a greater spectrum present in the

latter (Figure 1C). This is expected as flavivirus' control of IFN-I signaling occurs

mainly via its intracellular non-structural proteins (47, 48) and few ISGs can be

directly induced by IRF3 (49–51).

Of interest, STAT2 mRNA is expressed at higher levels in DV RNA-containing cells

when compared to bystanders and uninfected cells (Figure 1C and 4A). Hence, we

hypothesized that ISG15 interference with DV replication is associated with autocrine

IFN-I signaling in infected cells. Confirming the results from PBMCs and HuH7 cells,

ISG15 mRNA was induced in A549 cells during DV infection (Figure 4C). As

expected, this expression was abolished in knockout cells. Therefore, we measured

the activation of the IFN-I pathway in WT and ISG15-KO cells infected with DV.

Despite ISG15-knockout cells having full machinery to control viral infection as well

as hyper-responsiveness to exogenous IFNα (data not shown) (28, 29), they did not

respond robustly to DV infection. Infected ISG15-deficient cells showed reduced

STAT1 phosphorylation and STAT2 and IFIT1 expression when compared to WT

cells. As expected, A549 lacking ISG15 had no detectable USP18 (Figure 4B).

Infected knockout cells induced IFNβ mRNA at higher levels than the WT but had

less IFIT1 mRNA (Figures 4C) indicating that DV infection impairs the response at

both mRNA and protein levels downstream of IFN-I induction.

Taken together, our results suggest that ISG15 inhibits DV infection by regulating

early events of the IFN-I receptor activation. To test this, we constructed an

ISG15/IFNAR1 double knockout cell line based on the ISG15-KO background

(Figure 4E). Cells were infected with DV and foci area and number of infected cells

per foci was quantified. In parallel, we performed this experiment in IFNAR1

knockout cells, previously generated by our group (52, 53). The same phenotype

was observed in all three cell lines, with no additive or synergistic effects in the

double knockout (Figures 4F and G), which places intracellular ISG15 downstream

of IFNAR1 signaling in the control of DV infection.

ISG15 counteracts DV IFN-I evasion

Flaviviruses are known to counteract IFN-I signaling by inducing the degradation of

STAT2, a key protein in the interferon signal transduction pathway (11, 14, 54). As

our previous results suggest that ISG15's role during DV infection is downstream of

IFNAR engagement (Figure 4), we evaluated DV-mediated STAT2 degradation in

the absence of ISG15. In agreement with others (55), STAT2 degradation in the

A549 cell line occurs rapidly after infection. Despite showing sustained activation of

IFN-I signaling (data not shown), ISG15-KO cells showed pronounced STAT2

degradation when infected with DV (Figure 5A and E-upper panel). As infected IFN-

secreting cells are able to induce an antiviral state in neighboring-bystander cells by

inducing the expression of ISGs (56), we then examined in which cell population,

infected and/or bystander, ISG15 impacted DV infection. DV foci were co-stained for

flavivirus E protein and IFIT3, as an ISG marker, and confocal microscopy was

performed. Both infected and bystander cells were able to respond to infection,

producing IFIT3 (Figure 5D, top panel). However, IFIT3 expression in the foci context

was largely impaired and restricted to infected cells in the absence of ISG15 (Figures

5B, C and D). This data suggests that restriction of DV replication and dissemination

is achieved by both autocrine and paracrine IFN-I response amplification and is

dependent on ISG15 expression. To further investigate this, we sorted DV envelope

protein positive and negative cells (data not shown), therefore enabling us to

evaluate the impact of infection in virus-containing and bystander cells, respectively.

Strikingly, DV-positive cells had a lower expression of STAT2, IFIT3 and USP18

when compared to bystander cells; a phenotype that was markedly enhanced in

ISG15-deficient cells (Figure 5E, *bottom panel*). While also confirming DV inhibition

of IFNAR signaling, our results further suggest that ISG15 function is targeted for

successful viral infection.

USP18 expression overcomes ISG15 deficiency

STAT2 degradation mediated by all DV serotypes and ZIKV is largely dependent on

NS5 (11, 14), the virus RdRp and methyltransferase. NS5 was shown to bind to the

N-terminus of human STAT2 (11, 15). Interestingly, USP18 interacts with STAT2 via

its coiled-coil and DNA binding domains. This causes negative regulation of IFNAR

signaling by displacing JAK1 from chain 2 of the receptor (25). As shown here

(Figures 4B, 5E) and elsewhere (28, 29), the absence of ISG15 results in USP18's

destabilization. We therefore examined if NS5, STAT2 and USP18 are part of the

same complex. In HEK293 cells lacking ISG15 (data not shown) and primed with

IFNα for 18 hours, both endogenous STAT2 and overexpressed USP18, immuno-

precipitated with ZIKV's NS5 (Figure 6A, lane 2). This interaction was enhanced

when a protease-deficient high-expressing USP18 mutant (C64A) was used (Figure

6A, lane 3). We then hypothesized that the interaction of USP18 with STAT2

competes with NS5 binding, which in turn may result in decreased efficiency of virus

dependent STAT2 degradation. To test this, V5-tagged STAT2 was expressed in

HEK293 alone (Figure 6B, lane 1) or with FLAG-NS5 (lanes 2-5) in the absence

(lane 2) or presence (lanes 3-5) of increasing amounts of HA-tagged USP18,

followed by a treatment with IFNα for 18 hrs. STAT2 was immunoprecipitated with

V5 antibody and co-precipitation of NS5 and USP18 was assessed by immunoblot.

As expected, STAT2 interacted with NS5, while the amount of NS5 co-precipitated

with STAT2 decreased with increasing expression of USP18, suggesting that both

proteins indeed compete for the same binding region of STAT2.

Thus, we evaluated whether the presence of USP18 could restore STAT2

expression in ISG15 knockout cells during DV infection. Overexpression of USP18

led to an increase of STAT2 protein level similar to that in wild type cells (Figure 6C).

Importantly, USP18 reconstitution in ISG15-deficient cells was also able to reduce

virus replication (Figure 6D), suggesting that reestablishment of the IFN-I signaling

was sufficient to recover the WT phenotype in ISG15 knockout cells.

Together, the results presented here reveal that human ISG15 indirectly restricts DV

and ZIKV replication via its ability to stabilize USP18 and regulate the type 1 IFN

signaling pathway.

DISCUSSION

A successful infection is dependent on the virus replication machinery and its ability

to evade host immunity. One of the first lines of defense a virus has to overcome is a

plethora of antiviral genes induced by IFN-I (57).

ISG15 is induced early during infection (33, 58) and has been shown to have a viral

restriction role in several infection models. Most of those have been reported to be a

consequence of ISGylation (42–44, 59), even though this process was suggested to

be both inefficient and unspecific (59). Here we show that during the IFN-I response

elicited throughout DV infection, direct viral protein ISGylation is redundant for

antiviral immunity; rather, ISG15's ability to stabilize USP18 prevents NS5-mediated

STAT2 degradation, thus leading to a more effective interferon response that

culminates in DV and ZIKV restriction.

Many of ISG15's functions have been shown to be immunomodulatory. For instance,

ISGylation stabilizes IRF3 by occluding its ubiquitylation sites (60), negatively

regulates RIG-I (61) and activates PKR (62). Secreted ISG15 functions as a cytokine

(16, 63), leading to the production of IFNγ and IL-10 in human cells, crucial to the

control of pathogens such as *Mycobacterium tuberculosis* (17, 19). Moreover, free

intracellular ISG15 is essential for USP18 stability (Figures 4B, 5E, S4 and (28, 29)

and its absence leads to severe interferonopathy in humans (28). Also, there are

reports of ISG15 having an antiviral function independent of Ube1L during Chikungunya virus infection in mice (28, 64), where free ISG15 contributes to infection control by blunting potentially pathologic levels of cytokine effectors. Considering this range of functions, it is expected that different pathogens might interact with this pathway in different ways, according to its co-evolutionary history.

Of note, NK, NKT and monocytes were the PBMC populations with higher upregulation of ISG15 mRNA in single-cell gene expression studies. These have been shown to be the major producers and/or targets of free extracellular ISG15 in other contexts (17–19, 63). The influence of extracellular ISG15 during viral infections should be further explored in the future.

Interestingly, humans lacking ISG15 do not have increased susceptibility to common viral infections, such as influenza and HSV-1. The explanation for this, as suggested elsewhere, may lie in the sustained IFN-I response of their cells creating a hostile environment for virus growth (28, 29). Here, we reveal that ISG15 restricts DV and ZIKV replication indirectly by stabilizing USP18 and thereby disrupting NS5-STAT2 interaction: ISG15 promotes competition for a niche exploited by such viruses. We also demonstrate that USP18, STAT2 and NS5 co-immunoprecipitate, suggesting a very narrow window of opportunity that NS5 has to degrade STAT2. As USP18/ ISG15 interaction is reported to down-regulate IFN-I signaling in humans but not mice (29), it is tempting to speculate that NS5 interaction with STAT2, a major flavivirus immune evasion mechanism and also restricted to humans (15), was shaped by the USP18/ISG15 interaction.

Our results suggest an unexpected mechanism by which ISG15 can exert an antiviral function distinct from the debilitating effects of its conjugation to viral proteins. The key role of IFN-I in viral infections might lead to the perception of ISGs

having a necessarily direct antiviral function, a paradigm that is recently being reassessed, with a range of ISGs being implicated in infection-independent functions (65, 66). Here we provide mechanistic insight of the arms race between ISG15, USP18 and NS5 which suggests that protein/protein dynamics adjacent to IFNAR are a key determinant for the outcome of flavivirus infection.

ACKNOWLEDGMENTS

This work was supported by Comissão de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) Computational Biology programme (23038.010048/2013-27), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (311406/2017-3, 307889/2020-3 and 407609/2018-0), CNPq INCTV and the Academy of Medical Sciences/U.K. (NAF004/1005) to D.S.M. Howard Hughes Medical Institute – Early Career Scientist to A.B. (55007412), Wellcome Trust and Royal Society Sir Henry Dale Fellowship (202471/Z/16/Z) to T.R.S. Wellcome Trust PhD Studentship (105389/Z/14/Z) to T.J.S. CAPES scholarship (88887.195782/2018-00) to E.L.R. CNPq scholarship (205096/2018-2) to J.B. DSM and AB are CNPQ-PQ Scholars. We thank Dr. Sandra Pellegrini and Dr. Carsten Münk for kindly providing human USP18 plasmids. We thank Dr. Brian Ferguson for critical reading of the manuscript.

AUTHOR CONTRIBUTIONS

Conceptualization, C.E.E., A.B. and D.S.M.; Methodology, C.E.E., E.L.R and D.S.M.; Software, E.L.R; Formal Analysis, C.E.E., E.L.R and E.G.K., Investigation, C.E.E., A.A.S., D.O.P, G.M. and JB; Resources, C.E.E., A.A.S, P.F.S., Z.G, T.J.S., T.F. and T.R.S. Data Curation, E.L.R. and E.G.K., Writing-Original Draft, C.E.E. and D.S.M., Writing-Review and editing, C.E.E, E.L.R, D.O.P., E.G.K., P.F.S., J.B., T.J.S., T.F.,

T.R.S, A.B., and D.S.M., Visualization, C.E.E., E.L.R. and D.S.M., Supervision,

D.S.M., Project Administration and Funding Acquisition, D.S.M.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. ISG15 is expressed in DV-infected PBMC

(A) tSNE was used to visualize the single-cell global transcriptome data. Blue dots

represent uninfected cells, derived from healthy donors. Beige dots represent

bystander cells and red dots represent infected cells, both derived from patients

infected with DV.

(B) Differential expression of ISGs in PBMCs of patients infected with DV. The Gene

Ontology term "type one interferon" was used to filter the results from the single-cell

RNA sequencing.

(C) Single cell ISG expression variability in uninfected, bystander and infected.

ISGylation related genes are underlined. Blue bar represents uninfected cells,

derived from healthy donors. Beige bar represents bystander cells and red bar

represents infected cells, both derived from patients infected with DV.

(D) Violin plot representing the expression of ISGylation family members in

uninfected [U], bystander [B] and infected [I] PMBC.

(E) Expression of ISG15 in DV infected PMBCs. Size is proportional to the

percentage of infected cells in each cell type. Color intensity represents ISG15

average expression.

Figure 2. ISG15 restricts DV and ZIKV replication

(A) A549 WT and ISG15 KO were infected with 20 DV PFUs. At 36 hpi cells were

fixed, permeabilized and stained for the flavivirus E protein using 4G2 antibody.

Displayed images were acquired with a Leica DMI6000 B microscope.

(B and C) DV relative foci area (B) and the number of infected cells per foci (C)

quantified by ImageJ software and analyzed using Welch's t test. Error bars

represent mean ± SD. Results are representative of three independent experiments.

(D, E and F) Multiple-step DV growth curve in A549 cells. Cells were infected at an

MOI of 0.01 and harvested at multiple time points. Shown is the percentage of cells

infected as measured by E protein staining (4G2+) (D), extracellular viral RNA

relative expression by RT-qPCR (E) and titration by focus forming assay (FFA) (F).

Statistical analyses were conducted using unpaired t tests. Error bars represent

mean ± SD. Results are representative of three independent experiments.

(G) Changes in viral RNA relative expression over time following binding of DV to

A549 WT and ISG15 KO cells at 4°C and analyzed using unpaired t test. Error bars

represent mean ± SD. Results are representative of two independent experiments.

(H and I) Complementation of A549 ISG15 KO cells with ectopically expressed

ISG15. Cells were infected at an MOI of 0.01 and at 36 hpi cells were fixed and

processed for measurement by flow cytometry. Shown is the percentage of infected

cells as measured by E protein staining (4G2+) (H) and viral RNA relative expression

by RT-qPCR (I). One-way ANOVA was used to analyze these experiments. EV:

empty vector. Error bars represent mean ± SD. Results are representative of two

independent experiments.

(J and I) A549 WT and ISG15 KO were infected with 20 ZIKV PFUs. At 36 hpi cells

were fixed, permeabilized and stained for the flavivirus E protein. ZIKV relative foci

area (J) and number of infected cells per foci (I), quantified by ImageJ software and

analyzed using unpaired t test with Welch's correction. Images were acquired with

an Olympus IX83 inverted microscope. Error bars represent mean ± SD. Results are

representative of three independent experiments.

Statistical analyses were performed using Prism 8 (GraphPad Software). p values

*<0.05; **<0.01; ***<0.001.

Figure 3. ISGylation is not sufficient to restrict DV spread

(A) ISGylation profile of A549 WT and HERC5 KO cells by Western blot (WB). Cells

were primed with IFNα2b (100 IU/ml) for 24 h and cell lysates were analyzed with an

ISG15 antibody. (*) indicates antibody unspecific band.

(B and C) A549 cells were infected with 20 DV PFUs. At 36 hpi cells were fixed,

permeabilized and stained for the flavivirus E protein. DV relative foci area (B) and

the number of infected cells per foci (C) quantified by ImageJ software and analyzed

by one-way ANOVA. Images were acquired with an Olympus IX83 inverted

microscope.

Error bars represent mean ± SD. Results are representative of three or more

independent experiments. Statistical analyses were performed using Prism 8

(GraphPad Software). p values ****<0.0001.

Figure 4. **ISG15 is necessary for autocrine** IFNAR**1-mediated control of DV**

replication

(A) Violin plot representing the expression of STAT1 and STAT2 in uninfected [U],

bystander [B] and infected [I] PBMC.

(B and C) A549 cells were infected with 20 DV PFU. At 36 hpi, cells were harvested,

cell lysates were analyzed by WB (B) and the indicated mRNA transcripts were

quantified by RT-qPCR (C).   Error bars represent mean ± SD. Results are

representative of two independent experiments. Data was analyzed by unpaired t

test.

(D) A549 ISG15 KO cells were primed with IFNα2b (100 IU/ml) for 12 h, washed

three times with DPBS and allowed to rest 12 h before infection with DV or ZIKV at

an MOI of 0.1. Shown is the percentage of cells infected at 36 hpi, as measured by E

protein staining (4G2+). Error bars represent mean ± SD. Results are representative

of three independent experiments. Data was analyzed using unpaired t test.

(E) Clonal isolates of A549 cells were immunoblotted for IFIT3 after 24 h treatment

with IFNα2b (100 IU/ml) (+, control IFIT3 -proficient cells; 1–9, different clonal

isolates after limiting dilution). dKO refers to an ISG15/IFNAR1 double knockout.

(F and G) A549 cells were infected with 20 DV PFUs. At 36 hpi cells were fixed,

permeabilized and stained for the flavivirus E protein. DV relative foci area (F) and

number of infected cells per foci (G) quantified by ImageJ software and analyzed by

one-way ANOVA. Images were acquired with an Olympus IX83 inverted microscope.

Error bars represent mean ± SD. Results are representative of three independent

experiments. Statistical analyses were performed using Prism 8 (GraphPad

Software). p values *<0.05; ***<0.001.

Figure 5. ISG15 counteracts DV IFN-I evasion.

(A) A549 WT and ISG15 KO were infected with 20 DV PFU. Cells were harvested at

the indicated time points after infection (hpi) and cell lysates were analyzed by WB

using STAT2 and Actin antibodies.

(B, C and D) A549 WT and ISG15 KO immunofluorescence assay (IFA) 36 hpi for

cellular IFIT3 and flavivirus E protein expression (D). Percentage of IFIT3 (B) and DV

(C) positive cells per foci were quantified by ImageJ software and analyzed using

unpaired t test with Welch's correction when appropriate. Displayed images were

acquired with a Leica DMI6000 B microscope.

(E) A549 cells were infected with DV at MOI 0.01. At 36 hpi, cells were fixed,

permeabilized and stained for flavivirus E protein. Cell lysates were analyzed by WB

with the indicated antibodies before (upper panel) and after (bottom panel) cells were

sorted by fluorescence-activated cell sorting (FACS) based on E protein expression.

U: uninfected. B: bystander. I: infected.

Error bars represent mean ± SD. Results are representative of two independent

experiments. Statistical analyses were performed using Prism 8 (GraphPad

Software). p values ***<0.001.

Figure 6. USP18 expression overcomes ISG15 deficiency

(A) Flag-tag immunoprecipitation (IP) assay and WB analysis of HEK293 ISG15 KO

cells transfected with ZIKV NS5-FLAG, HA-tagged human USP18 WT or human

USP18 C64A mutant or the empty vector (pcDNA3.1) plasmids, followed by IFNα2b

(100 IU/ml) priming for 18 h. WCL, whole cell lysate. Results are representative of

three independent experiments.

(B) V5-tagged STAT2 IP assay and WB analysis of HEK293 cells transfected with

the indicated plasmids, followed by IFNAα2b (100 IU/ml) priming for 18 h. Results

are representative of three independent experiments.

(C and D) Complementation with USP18 in A549 ISG15 KO cells. Cells were stably

transfected with human USP18 or the empty vector and infected with DV at an MOI

of 0.01. At 36 hpi, cells were harvested and cell lysates were analyzed by WB with

the corresponding antibodies (C). Percentage of cells infected as measured by E

protein staining (D). Error bars represent mean ± SD. Results are representative of

two independent experiments. Statistical analyses were conducted using Mann-

Whitney's test in Prism 8 (GraphPad Software. p value **<0.01 (D). EV: empty

vector.

LEAD CONTACT AND MATERIAL AVAILABILITY

Further information and requests for resources and reagents should be directed to

and will be fulfilled by the Lead Contact, Daniel Santos Mansur

(daniel.mansur@ufsc.br)

MATERIAL AND METHODS

Cells and viruses

Mammalian cell lines were maintained at 37°C under the conditions of a humidified

atmosphere and 5% CO_2. The human alveolar adenocarcinoma-derived A549 cells,

human embryonic kidney HEK293 and the African green monkey kidney-derived

Vero cells were cultured in Dulbecco's modified Eagle's medium F-12 (DMEM F12)

(Gibco) supplemented with 5% fetal bovine serum (FBS) (Gibco) and

streptomycin/penicillin (100 U/ml) (Gibco). The *Aedes albopictus* mosquito-derived

cell line C6/36 was maintained at 28°C in a BOD in Leibovitz's L-15 medium (L-15)

(Gibco) supplemented with 10% FBS (Gibco), 0.26% tryptose phosphate broth

(Sigma) and 50 µg/ml gentamicin (Sigma). A549 ISG15 KO and IFNAR1 KO have

been described elsewhere (17, 52, 53). All cells were negative for mycoplasma.

Dengue virus serotype 4 (DENV-4 TVP/360 – GenBank accession number:

KU513442) and Zika virus (ZV BR 2015/15261 - GenBank accession number:

MF073358) stocks were propagated in C6/36 cells and titrated in Vero cells.

Vesicular stomatitis virus-green fluorescent protein (VSVeGFP) (Indiana strain,

Marques-JT, Plos Pathogens 2013) and Herpes simplex virus-1-green fluorescent

protein (HSV-1eGFP) (SC16) viruses were propagated and titrated in Vero cells.

HSV-1eGFP was a kind gift of Professor Stacey Efstathiou.

Single-cell RNA sequencing analysis

Processed, publicly available single-cell RNA-seq data are available through the

GEO accession numbers GSE116672 and GSE110496. We downloaded processed

single-cell data and metadata from the supplementary information from the

respective publications (31, 32).

Then, we used CellRouter to analyze these datasets. To perform the tSNE analysis

using single-cell data generated by Zanini 2018 (32), we set the parameters

num.pcs=10, seed=1 and max_iter=1000 in the computeTSNE function. Next, we

identified genes preferentially expressed in Uninfected, Bystander and Infected cells

using a cutoff for the log2 fold change of 0.25. We used a custom script to obtain all

genes containing the keywords "type I interferon" in the Gene Ontology Biological

Processes (package versions: org.Hs.eg.db_3.10.0, GO.db_3.10.0). Next, we took

the overlap of type I interferon genes with the genes preferentially expressed in each

condition reported above. The remaining analyses were focused on these genes.

To perform the tSNE analysis using the single-cell data generated by Zanini 2018

(31), we set the parameters num.pcs=20, seed=1 and max_iter=1000 in the

computeTSNE function. We used a strategy similar to the one described above to

identify genes differentially expressed in each condition but used a cutoff of 0.15 for

this dataset. The parameter num.pcs was determined using the elbow approach, as

described in the CellRouter tutorial at https://github.com/edroaldo/cellrouter.

CRISPR/Cas9-mediated gene editing

A549 WT and ISG15 KO cells were co-transfected with two Herc5 or Ifnar1-targeting

gRNA CRISPR/Cas9-GFP plasmids, respectively. HEK293 WT cells were

transfected with three Isg15-targeting gRNA CRISPR/Cas9-GFP plasmids (Table

S1) (Horizon Cambridge, UK). After 72 h, cells were sorted by FACS (FACSMelody,

BD) and single-cell derived clones were initially screened by PCR genotyping.

Additionally, both HERC5 and ISG15/IFNAR1 (dKO) clones were functionally tested

by assessing their ISGylation profile and expression of ISGs, respectively, after IFNα

priming. Briefly, A549 WT and HERC5 and dKO clones were primed with IFNα2b

(100 IU/ml) (PBL Assay Science) for 24 h and the expression of ISG15-conjugates

and IFIT3 was analyzed by Western blot. HEK293 cells were primed with IFNα2b

(1000 IU/ml) for 8 h, total RNA was isolated and Isg15 mRNA expression was

assessed by RT-qPCR.

Viral infection

A549 cells were seeded one day prior to infection in appropriate multi-well plates.

For foci assay, a viral inoculum containing 20 foci forming units (PFU) of the

corresponding virus was added to each well, and virus adsorption was performed in

DMEM supplemented with penicillin/streptomycin (100 U/ml) for 90 min at 37°C.

Cells were washed with PBS to remove un-adsorbed virus, and maintained in DMEM

1.5% carboxymethylcellulose sodium (CMC) (Sigma). Alternatively, cells were

infected with DV at the indicated MOI, as described above, and maintained in DMEM

supplemented with 1% FBS.

Titration

DV titration was performed by focus forming assay (FFA) in C6/36 cells. Briefly,

C6/36 cells were seeded (1×10^5 cells/well in 24 well plate) and after overnight

incubation were infected with a 10-fold serial dilution of virus samples (cell culture

supernatants) in L-15 with 0.26% tryptose and 25 µg/ml of gentamicin. After 90

minutes, the inoculum was removed and a CMC overlay media (L-15 media with 5%

FCS, 0.26% tryptose, 25 µg/ml of gentamicin and 1.6% of CMC) was added and the

plates were incubated for 7 days at 28°C. After incubation, cells were washed, fixed

with 3% paraformaldehyde (PFA) (Sigma-Aldrich) and permeabilized with 0.5% triton

X100 (Sigma-Aldrich). After washing, cells were immunostained with mouse

monoclonal anti flavivirus E protein antibody 4G2 (ATCC® HB-112™, dilution 1:100),

followed by goat anti-mouse immunoglobulin conjugated to alkaline phosphatase

(Promega S3721, dilution 1:7500). Focuses of infection were revealed using

NBT/BCIP reagent (Promega), following the manufacturer's instructions and the virus

titer calculated as follow: media of focus number/inoculum volume x dilution. The

results are expressed as $FFU_{C6/36}$/ml (Gould et al., 1985).

VSVeGFP and HSV-1eGFP titrations were performed by foci assay. Briefly, Vero

cells were seeded in 24-well plates 24 h prior to infection. Cell monolayers were

washed with PBS and inoculated with 0.3 ml of serial 10-fold dilutions of the virus in

duplicates. After 90 min adsorption at 37°C, each well was re-suspended in DMEM

1.5% CMC. At 48 hpi, cells were fixed with 3% PFA (Sigma) for 30 min, washed 3

times with PBS and stained with 1% crystal violet (Sigma) for 30 min at room

temperature. Virus yield was calculated and expressed as foci forming units per ml

(PFU/ml).

Immunofluorescence

At 24 hpi and 18 hpi, respectively, HSVeGFP and VSVeGFP infected cells were

fixed with 3% PFA for 20 min at room temperature, permeabilized with 0.5% Triton

X100 in PBS for 4 min and stained with DAPI counterstain (Molecular Probes). At 36

hpi, DV and ZIKV infected cells were fixed, permeabilized and stained with mouse

monoclonal 4G2 antibody (10 µg/ml dilution 1:100), followed by incubation with Alexa

Fluor 488 rabbit anti-mouse IgG (H□+□L, Life Technologies, dilution 1:500), and

DAPI counterstain. Images were acquired with an Olympus IX83 inverted

microscope.　Briefly, virus foci, determined by eGFP or flavivirus E protein

expression, were delimited; images were converted to 16-bit and processed to be

analyzed with the ImageJ Software Cell Counter Plugin (W. S. Rasband, ImageJ, US

National Institutes of Health, Bethesda, MD; http://rsb.info.nih.gov/ij/, 1997–2006).

For confocal analysis, A549 cells grown on glass coverslips were mock infected or

infected with 10 DV PFU. At 36 hpi, cells were fixed and permeabilized. Following

washes with PBS, cells were stained with mouse monoclonal anti-E protein (4G2,

dilution 1:100) and rabbit polyclonal anti-IFIT3 (Proteintech, 15201-1-AP, dilution

1:200) for 1 h at room temperature. The cells were washed with PBS and stained

with secondary antibodies Alexa Fluor 488 goat anti-mouse IgG (H□+□L, Life

Technologies, dilution 1:500) and Alexa 568 goat anti-rabbit IgG (H+L, Life

Technologies, dilution 1:500) and DAPI counterstain (Molecular Probes). Cells were

washed and coverslips mounted using Prolong antifade reagent (Invitrogen). Z-stack

and max intensity projection images were generated with a Leica DMI6000 B

confocal microscope and Leica Application Suite X software for image analysis

(Leica Microsystems).

Flow cytometry and cell sorting

Cells were fixed with 2% PFA in PBS, washed twice with PBS and permeabilized

with 0.5% saponin in 1% BSA in PBS. Anti-flavivirus E protein mAb 4G2 was

conjugated to Alexa Fluor 488 5-SDP (Life Technologies) following the

manufacturer's instructions. Cells were incubated with FITC-conjugated 4G2 (dilution

1:1000) in permeabilization buffer for 40 minutes at room temperature, washed once

and resuspended in FACS buffer. The cell suspensions were analyzed by flow

cytometry on a FACSVerse instrument (BD Biosciences) and analyzed using FlowJo

V10 software (BD). Cell sorting experiments were performed on a FACSMelody cell

sorter (BD Biosciences).

RT-qPCR

vRNA was isolated by using QIAamp viral RNA Mini Kit (Qiagen). Intracellular total

RNA was isolated with TRIZOL (Thermo Life) following manufacturer's instructions.

A total of 1 µg was reverse transcribed using the High-capacity cDNA reverse

transcription kit (Applied Biosystems). Quantitative PCRs (qPCRs) were performed

with GoTaq ® qPCR Master mix (Promega) following the standard cycling conditions

suggested by the manufacturer in a StepOnePlus real time PCR system (Applied

Biosystems). VSVeGFP and 18S mRNA were used as control housekeeping genes.

Amounts of DV or ISG mRNA were calculated by using the ΔΔCT method. Primers

specific to the mRNA analyzed are listed in Table S1.

Western blot

Human antibodies used for immunoblot were as follows: mouse mAb to β-actin

(Abcam, ab6276, dilution 1:4000), rabbit mAb to IFIT1 (Abcam, ab137632, dilution

1:1000), mouse mAb to pSTAT1(Y701) [M135] (Abcam, ab29045, dilution 1:1000),

mouse mAb to STAT1 (Abcam, ab3987, dilution 1:1000), rabbit pAb to IFIT3

(ProteinTech, 15201-1-AP, dilution 1:1000), rabbit mAb to USP18 [D4E7] (Cell

Signaling Technologies, 4813, dilution 1:1000), rabbit mAb to STAT2 [D9J7L] (Cell

Signaling Technologies, 72604, dilution 1:1000), mouse mAb to ISG15 (R&D

System, MAB4845, dilution 1:1000), home-made mouse mAb anti-GFP (67, 68)

(dilution 1:1000), mouse mAb to FLAG tag (Sigma, F3165, dilution 1:500), rabbit

mAb to HA (Sigma, H6908, dilution 1:2000) and rabbit mAb to V5 (CST, D3H8Q,

dilution 1:1000)

Cells were treated as indicated, wash two times with ice-cold PBS and then lysed in

RIPA buffer [50 mM tris-HCl (pH 7.5), 150 mM NaCl, 1% Triton X100, 0.5% sodium

deoxycholate, 0.1% SDS, 5 mM EDTA] supplemented with 1x protease inhibitors

(Mini Protease Inhibitor Tablets, Roche). Total protein concentration was determined

by BSA Protein Assay Kit (Thermo Life). For western blotting, 20 µg of total protein

were prepared in dithiothreitol-containing Laemmli sample buffer, separated and

transferred to a nitrocellulose blotting membrane (GE Healthcare Amersham). After

transfer, the membrane was blocked with 5% nonfat dry milk in TBS 0.1% Tween 20

(TBST) for 1 h at room temperature. Membrane was incubated with primary Abs

diluted in 2% BSA in TBST at 4°C with gentle shaking overnight. Membrane was

washed three times with TBST and then incubated with the appropriate secondary

HRP-linked antibody for 1 h at room temperature. Membranes were washed and

covered with ECL developing solution (Pierce^{TM} ECL WB substrate, Thermo Fisher

Scientific).

Plasmids and transfections

Herc5 and Ifnar1 sgRNA/Cas9/GFP plasmids were provided by Horizon (Cambridge,

UK). sgRNA sequences are described in Supplementary Table 1. Expression

plasmid for ISG15 was described elsewhere (17). pMax^{TM} GFP expression vector

was acquired from Lonza (cat numb #D-00061). Expression plasmid for ZIKV NS5

was generated by amplifying the NS5 coding sequence (amino acids 2521-3423 in

the polyprotein) from a previously described plasmid-based ZIKV reverse genetic

system (69) using primers containing an N-terminal FLAG tag and inserted into

pcDNA3.1. V5-STAT2-expressing plasmid was generated by inserting the STAT2

coding sequence into pcDNA5/FRT/TO/V5 (Addgene #9081) vector. Expression

plasmids for human USP18 WT and USP18 C64A mutants were kindly provided by

Dr Carsten Münk (70). Transfections were performed with FuGene6 (Promega),

following the manufacturer's instructions. Stably transfected cells were selected with

geneticin (500 µg/ml) (Sigma).

Immunoprecipitation

For STAT2 co-immunoprecipitation assays, cells were lysed in RIPA buffer

composed of 20 mM Tris-HCl pH 7.4, 150 mM NaCl,10 mM CaCl2, 0.1% v/v Triton

X-100, 10% v/v glycerol and complete protease inhibitor cocktail (Mini Protease

Inhibitor Tablets, Roche). The supernatant was separated by centrifugation at 12.000

g at 4 °C for 10 min and incubated with STAT2 antibody (2 µg) (Santa Cruz

Biotechnology, 514193) for 3 h at 4°C with gentle shaking. Complexes were

precipitated with protein A/G Plus-agarose (Santa Cruz Biotechnology), washed with

TBS and resuspended in SDS sample buffer. Immunoprecipitates were subjected to

SDS-PAGE and western blotting, as described above.

FLAG-tagged proteins were immunoprecipitated with anti-FLAG M2-agarose

(Sigma), following the manufacturer's instructions.

All assays were performed three times and representative blots are presented.

QUANTIFICATION AND STATISTICAL ANALYSIS

Details concerning the statistical analysis methods are provided in each figure

legend. Briefly, all data were analyzed using GraphPad Prism 8 software and were

shown as mean and the standard deviation (SD). Statistical significance was

determined by Welch's t test or one-way ANOVA for foci size analyses, unpaired t

test for virus multiple-step growth curve, cellular mRNA quantification and

percentage of cells infected. Statistical significance is indicated by ∗, $p < 0.05$; ∗∗, $p <$

0.01; ∗∗∗, $p < 0.001$.

REFERENCES

1. Mansur DS, Smith GL, Ferguson BJ. 2014. Intracellular sensing of viral DNA by

the innate immune system. Microbes Infect 16:1002–1012.

2. Gebhardt A, Laudenbach BT, Pichlmair A. 2017. Discrimination of Self and Non-

Self Ribonucleic Acids. J Interferon Cytokine Res 37:184–197.

3. Schoggins JW. 2019. Interferon-Stimulated Genes: What Do They All Do? Annu

Rev Virol https://doi.org/10.1146/annurev-virology-092818-015756.

4. García-Sastre A. 2017. Ten Strategies of Interferon Evasion by Viruses. Cell

Host Microbe 22:176–184.

5. Pierson TC, Diamond MS. 2018. The emergence of Zika virus and its new

clinical syndromes. Nature 560:573–581.

6. Barrett ADT. 2018. The reemergence of yellow fever. Science.

7. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM,

Brownstein JS, Hoen AG, Sankoh O, Myers MF, George DB, Jaenisch T, Wint

GRW, Simmons CP, Scott TW, Farrar JJ, Hay SI. 2013. The global distribution

and burden of dengue. Nature 496:504–507.

8. Shepard DS, Undurraga EA, Halasa YA, Stanaway JD. 2016. The global

economic burden of dengue: a systematic analysis. Lancet Infect Dis 16:935–

742  941.

743 9. Miorin L, Maestre AM, Fernandez-Sesma A, García-Sastre A. 2017. Antagonism

744  of type I interferon by flaviviruses. Biochem Biophys Res Commun 492:587–596.

745 10. Ngono AE, Shresta S. 2018. Immune Response to Dengue and Zika. Annu Rev

746  Immunol 36:279–308.

747 11. Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M,

748  Schwarz MC, Sánchez-Seco MP, Evans MJ, Best SM, García-Sastre A. 2016.

749  Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host

750  Microbe 19:882–890.

751 12. Laurent-Rolle M, Morrison J, Rajsbaum R, Macleod JML, Pisanelli G, Pham A,

752  Ayllon J, Miorin L, Martinez C, tenOever BR, García-Sastre A. 2014. The

753  interferon signaling antagonist function of yellow fever virus NS5 protein is

754  activated by type I interferon. Cell Host Microbe 16:314–327.

755 13. Chen S, Wu Z, Wang M, Cheng A. 2017. Innate Immune Evasion Mediated by

756  Flaviviridae Non-Structural Proteins. Viruses 9.

757 14. Morrison J, Laurent-Rolle M, Maestre AM, Rajsbaum R, Pisanelli G, Simon V,

758  Mulder LCF, Fernandez-Sesma A, García-Sastre A. 2013. Dengue virus co-opts

759  UBR4 to degrade STAT2 and antagonize type I interferon signaling. PLoS

760  Pathog 9:e1003265.

761 15. Ashour J, Morrison J, Laurent-Rolle M, Belicha-Villanueva A, Plumlee CR,

762  Bernal-Rubio D, Williams KL, Harris E, Fernandez-Sesma A, Schindler C,

763  García-Sastre A. 2010. Mouse STAT2 restricts early dengue virus replication.

Cell Host Microbe 8:410–421.

16. Dos Santos PF, Mansur DS. 2017. Beyond ISGlylation: Functions of Free

Intracellular and Extracellular ISG15. J Interferon Cytokine Res 37:246–253.

17. Dos Santos PF, Van Weyenbergh J, Delgobo M, Oliveira Patricio D de,

Ferguson BJ, Guabiraba R, Dierckx T, Menezes SM, Báfica A, Mansur DS.

2018. ISG15-Induced IL-10 Is a Novel Anti-Inflammatory Myeloid Axis Disrupted

during Active Tuberculosis. J Immunol 200:1434–1442.

18. Swaim CD, Scott AF, Canadeo LA, Huibregtse JM. 2017. Extracellular ISG15

Signals Cytokine Secretion through the LFA-1 Integrin Receptor. Mol Cell

68:581–590.e5.

19. Bogunovic D, Byun M, Durfee LA, Abhyankar A, Sanal O, Mansouri D, Salem S,

Radovanovic I, Grant AV, Adimi P, Mansouri N, Okada S, Bryant VL, Kong X-F,

Kreins A, Velez MM, Boisson B, Khalilzadeh S, Ozcelik U, Darazam IA,

Schoggins JW, Rice CM, Al-Muhsen S, Behr M, Vogt G, Puel A, Bustamante J,

Gros P, Huibregtse JM, Abel L, Boisson-Dupuis S, Casanova J-L. 2012.

Mycobacterial disease and impaired IFN-γ immunity in humans with inherited

ISG15 deficiency. Science 337:1684–1688.

20. Delgobo M, Daniel A G, Kozlova E, Rocha EL, Rodrigues-Luiz GF, Mascarin L,

Dias G, Patrício DO, Dierckx T, Bicca MA, Breton G, de Menezes YKT, Starick

MR, Rovaris D, Del Moral J, Mansur DS, Van Weyenbergh J, Báfica A. An

evolutionary recent IFN-IL-6-CEBP axis is linked to monocyte expansion and

tuberculosis severity in humans https://doi.org/10.1101/514943.

21. Skaug B, Chen ZJ. 2010. Emerging role of ISG15 in antiviral immunity. Cell

143:187–190.

22. Dastur A, Beaudenon S, Kelley M, Krug RM, Huibregtse JM. 2006. Herc5, an

interferon-induced HECT E3 enzyme, is required for conjugation of ISG15 in

human cells. J Biol Chem 281:4334–4338.

23. Wong JJY, Pung YF, Sze NS-K, Chin K-C. 2006. HERC5 is an IFN-induced

HECT-type E3 protein ligase that mediates type I IFN-induced ISGylation of

protein targets. Proc Natl Acad Sci U S A 103:10735–10740.

24. Perng Y-C, Lenschow DJ. 2018. ISG15 in antiviral immunity and beyond. Nat

Rev Microbiol 16:423–439.

25. Arimoto K-I, Löchte S, Stoner SA, Burkart C, Zhang Y, Miyauchi S, Wilmes S,

Fan J-B, Heinisch JJ, Li Z, Yan M, Pellegrini S, Colland F, Piehler J, Zhang D-E.

2017. STAT2 is an essential adaptor in USP18-mediated suppression of type I

interferon signaling. Nat Struct Mol Biol 24:279–289.

26. Malakhov MP, Malakhova OA, Kim KI, Ritchie KJ, Zhang D-E. 2002. UBP43

(USP18) specifically removes ISG15 from conjugated proteins. J Biol Chem

277:9976–9981.

27. Malakhova O, Malakhov M, Hetherington C, Zhang D-E. 2002.

Lipopolysaccharide activates the expression of ISG15-specific protease UBP43

via interferon regulatory factor 3. J Biol Chem 277:14703–14711.

28. Zhang X, Bogunovic D, Payelle-Brogard B, Francois-Newton V, Speer SD, Yuan

C, Volpi S, Li Z, Sanal O, Mansouri D, Tezcan I, Rice GI, Chen C, Mansouri N,

Mahdaviani SA, Itan Y, Boisson B, Okada S, Zeng L, Wang X, Jiang H, Liu W,

Han T, Liu D, Ma T, Wang B, Liu M, Liu J-Y, Wang QK, Yalnizoglu D,

Radoshevich L, Uzé G, Gros P, Rozenberg F, Zhang S-Y, Jouanguy E,

Bustamante J, García-Sastre A, Abel L, Lebon P, Notarangelo LD, Crow YJ,

Boisson-Dupuis S, Casanova J-L, Pellegrini S. 2015. Human intracellular ISG15

prevents interferon-α/β over-amplification and auto-inflammation. Nature

517:89–93.

29. Speer SD, Li Z, Buta S, Payelle-Brogard B, Qian L, Vigant F, Rubino E, Gardner

TJ, Wedeking T, Hermann M, Duehr J, Sanal O, Tezcan I, Mansouri N, Tabarsi

P, Mansouri D, Francois-Newton V, Daussy CF, Rodriguez MR, Lenschow DJ,

Freiberg AN, Tortorella D, Piehler J, Lee B, García-Sastre A, Pellegrini S,

Bogunovic D. 2016. ISG15 deficiency and increased viral resistance in humans

but not mice. Nat Commun 7:11496.

30. Lenschow DJ, Lai C, Frias-Staheli N, Giannakopoulos NV, Lutz A, Wolff T, Osiak

A, Levine B, Schmidt RE, García-Sastre A, Leib DA, Pekosz A, Knobeloch K-P,

Horak I, Virgin HW 4th. 2007. IFN-stimulated gene 15 functions as a critical

antiviral molecule against influenza, herpes, and Sindbis viruses. Proc Natl Acad

Sci U S A 104:1371–1376.

31. Zanini F, Pu S-Y, Bekerman E, Einav S, Quake SR. 2018. Single-cell

transcriptional dynamics of flavivirus infection. Elife 7.

32. Zanini F, Robinson ML, Croote D, Sahoo MK, Sanz AM, Ortiz-Lasso E, Albornoz

LL, Rosso F, Montoya JG, Goo L, Pinsky BA, Quake SR, Einav S. 2018. Virus-

inclusive single-cell RNA sequencing reveals the molecular signature of

progression to severe dengue. Proc Natl Acad Sci U S A 115:E12363–E12369.

33. Ramos I, Smith G, Ruf-Zamojski F, Martinez-Romero C, Fribourg M, Carbajal

EA, Hartmann BM, Nair VD, Marjanovic N, Monteagudo PL, DeJesus VA,

Mutetwa T, Zamojski M, Tan GS, Jayaprakash C, Zaslavsky E, Albrecht RA,

Sealfon SC, García-Sastre A, Fernandez-Sesma A. 2019. Innate immune

response to influenza virus at single-cell resolution in human epithelial cells

revealed paracrine induction of interferon lambda 1. J Virol

https://doi.org/10.1128/JVI.00559-19.

34. Keskinen P, Nyqvist M, Sareneva T, Pirhonen J, Melén K, Julkunen I. 1999.

Impaired antiviral response in human hepatoma cells. Virology 263:364–375.

35. Schmid B, Rinas M, Ruggieri A, Acosta EG, Bartenschlager M, Reuter A, Fischl

W, Harder N, Bergeest J-P, Flossdorf M, Rohr K, Höfer T, Bartenschlager R.

2015. Live Cell Analysis and Mathematical Modeling Identify Determinants of

Attenuation of Dengue Virus 2'-O-Methylation Mutant. PLoS Pathog

11:e1005345.

36. Roth H, Magg V, Uch F, Mutz P, Klein P, Haneke K, Lohmann V, Bartenschlager

R, Fackler OT, Locker N, Stoecklin G, Ruggieri A. 2017. Flavivirus Infection

Uncouples Translation Suppression from Cellular Stress Responses. MBio 8.

37. Wang K, Zou C, Wang X, Huang C, Feng T, Pan W, Wu Q, Wang P, Dai J.

2018. Interferon-stimulated TRIM69 interrupts dengue virus replication by

ubiquitinating viral nonstructural protein 3. PLoS Pathog 14:e1007287.

38. Whelan JN, Li Y, Silverman RH, Weiss SR. 2019. Zika Virus Production Is

Resistant to RNase L Antiviral Activity. J Virol 93.

39. Gullberg RC, Steel JJ, Pujari V, Rovnak J, Crick DC, Perera R. 2018. Stearoly-

CoA desaturase 1 differentiates early and advanced dengue virus infections and

determines virus particle infectivity. PLoS Pathog 14:e1007261.

40. Zhang J, Lan Y, Li MY, Lamers MM, Fusade-Boyer M, Klemm E, Thiele C,

Ashour J, Sanyal S. 2018. Flaviviruses Exploit the Lipid Droplet Protein AUP1 to

Trigger Lipophagy and Drive Virus Production. Cell Host Microbe 23:819–

831.e5.

41. Zhao C, Denison C, Huibregtse JM, Gygi S, Krug RM. 2005. Human ISG15

conjugation targets both IFN-induced and constitutively expressed proteins

functioning in diverse cellular pathways. Proc Natl Acad Sci U S A 102:10200–

10205.

42. Zhao C, Hsiang T-Y, Kuo R-L, Krug RM. 2010. ISG15 conjugation system

targets the viral NS1 protein in influenza A virus-infected cells. Proc Natl Acad

Sci U S A 107:2253–2258.

43. Kim YJ, Kim ET, Kim Y-E, Lee MK, Kwon KM, Kim KI, Stamminger T, Ahn J-H.

2016. Consecutive Inhibition of ISG15 Expression and ISGylation by

Cytomegalovirus Regulators. PLoS Pathog 12:e1005850.

44. González-Sanz R, Mata M, Bermejo-Martín J, Álvarez A, Cortijo J, Melero JA,

Martínez I. 2016. ISG15 Is Upregulated in Respiratory Syncytial Virus Infection

and Reduces Virus Growth through Protein ISGylation. J Virol 90:3428–3438.

45. Hishiki T, Han Q 'en, Arimoto K-I, Shimotohno K, Igarashi T, Vasudevan SG,

Suzuki Y, Yamamoto N. 2014. Interferon-mediated ISG15 conjugation restricts

dengue virus 2 replication. Biochem Biophys Res Commun 448:95–100.

46. François-Newton V, Magno de Freitas Almeida G, Payelle-Brogard B, Monneron

D, Pichard-Garcia L, Piehler J, Pellegrini S, Uzé G. 2011. USP18-based

negative feedback control is induced by type I and type III interferons and

specifically inactivates interferon α response. PLoS One 6:e22200.

47. Dalrymple NA, Cimica V, Mackow ER. 2015. Dengue Virus NS Proteins Inhibit

RIG-I/MAVS Signaling by Blocking TBK1/IRF3 Phosphorylation: Dengue Virus

Serotype 1 NS4A Is a Unique Interferon-Regulating Virulence Determinant. MBio

6:e00553–15.

48. Muñoz-Jordán JL, Laurent-Rolle M, Ashour J, Martínez-Sobrido L, Ashok M,

Lipkin WI, García-Sastre A. 2005. Inhibition of alpha/beta interferon signaling by

the NS4B protein of flaviviruses. J Virol 79:8004–8013.

49. Daly C, Reich NC. 1993. Double-stranded RNA activates novel factors that bind

to the interferon-stimulated response element. Mol Cell Biol 13:3756–3764.

50. Daly C, Reich NC. 1995. Characterization of specific DNA-binding factors

activated by double-stranded RNA as positive regulators of interferon

alpha/beta-stimulated genes. J Biol Chem 270:23739–23746.

51. Weaver BK, Kumar KP, Reich NC. 1998. Interferon regulatory factor 3 and

CREB-binding protein/p300 are subunits of double-stranded RNA-activated

transcription factor DRAF1. Mol Cell Biol 18:1359–1368.

52. Guimarães ES, Gomes MTR, Campos PC, Mansur DS, Dos Santos AA, Harms

J, Splitter G, Smith JA, Barber GN, Oliveira SC. 2019. Cyclic Dinucleotides

Trigger STING-Dependent Unfolded Protein Response That Favors Bacterial

Replication. J Immunol 202:2671–2681.

53. Antunes KH, Fachi JL, de Paula R, da Silva EF, Pral LP, Dos Santos AÁ, Dias

GBM, Vargas JE, Puga R, Mayer FQ, Maito F, Zárate-Bladés CR, Ajami NJ,

Sant'Ana MR, Candreva T, Rodrigues HG, Schmiele M, Silva Clerici MTP,

Proença-Modena JL, Vieira AT, Mackay CR, Mansur D, Caballero MT, Marzec J,

Li J, Wang X, Bell D, Polack FP, Kleeberger SR, Stein RT, Vinolo MAR, de

Souza APD. 2019. Microbiota-derived acetate protects against respiratory

syncytial virus infection through a GPR43-type 1 interferon response. Nat

Commun 10:3273.

54. Ashour J, Laurent-Rolle M, Shi P-Y, García-Sastre A. 2009. NS5 of dengue virus

mediates STAT2 binding and degradation. J Virol 83:5408–5418.

55. Jones M, Davidson A, Hibbert L, Gruenwald P, Schlaak J, Ball S, Foster GR,

Jacobs M. 2005. Dengue virus inhibits alpha interferon signaling by reducing

STAT2 expression. J Virol 79:5414–5420.

56. Rand U, Rinas M, Schwerk J, Nöhren G, Linnes M, Kröger A, Flossdorf M, Kály-

Kullai K, Hauser H, Höfer T, Köster M. 2012. Multi-layered stochasticity and

paracrine signal propagation shape the type-I interferon response. Mol Syst Biol

8:584.

57. Schneider WM, Chevillotte MD, Rice CM. 2014. Interferon-stimulated genes: a

complex web of host defenses. Annu Rev Immunol 32:513–545.

58. Loeb KR, Haas AL. 1992. The interferon-inducible 15-kDa ubiquitin homolog

conjugates to intracellular proteins. J Biol Chem 267:7806–7813.

59. Durfee LA, Lyon N, Seo K, Huibregtse JM. 2010. The ISG15 conjugation system

broadly targets newly synthesized proteins: implications for the antiviral function

of ISG15. Mol Cell 38:722–732.

60. Shi H-X, Yang K, Liu X, Liu X-Y, Wei B, Shan Y-F, Zhu L-H, Wang C. 2010.

Positive regulation of interferon regulatory factor 3 activation by Herc5 via ISG15

modification. Mol Cell Biol 30:2424–2436.

61. Kim M-J, Hwang S-Y, Imaizumi T, Yoo J-Y. 2008. Negative feedback regulation

of RIG-I-mediated antiviral signaling by interferon-induced ISG15 conjugation. J

Virol 82:1474–1483.

62. Okumura F, Okumura AJ, Uematsu K, Hatakeyama S, Zhang D-E, Kamura T.

2013. Activation of double-stranded RNA-activated protein kinase (PKR) by

interferon-stimulated gene 15 (ISG15) modification down-regulates protein

translation. J Biol Chem 288:2839–2847.

63. D'Cunha J, Knight E Jr, Haas AL, Truitt RL, Borden EC. 1996. Immunoregulatory

properties of ISG15, an interferon-induced cytokine. Proc Natl Acad Sci U S A

93:211–215.

64. Werneke SW, Schilte C, Rohatgi A, Monte KJ, Michault A, Arenzana-Seisdedos

F, Vanlandingham DL, Higgs S, Fontanet A, Albert ML, Lenschow DJ. 2011.

ISG15 is critical in the control of Chikungunya virus infection independent of

UbE1L mediated conjugation. PLoS Pathog 7:e1002322.

65. Vuillier F, Li Z, Commere P-H, Dynesen LT, Pellegrini S. 2019. USP18 and

ISG15 coordinately impact on SKP2 and cell cycle progression. Sci Rep 9:4066.

66. Dabo S, Maillard P, Collados Rodriguez M, Hansen MD, Mazouz S, Bigot D-J,

Tible M, Janvier G, Helynck O, Cassonnet P, Jacob Y, Bellalou J, Gatignol A,

Patel RC, Hugon J, Munier-Lehmann H, Meurs EF. 2017. Inhibition of the

inflammatory response to stress by targeting interaction between PKR and its

cellular activator PACT. Sci Rep 7:16129.

67. Zanluca C, Mazzarotto GACA, Bordignon J, Duarte Dos Santos CN. 2014.

Development, characterization and application of monoclonal antibodies against

Brazilian Dengue virus isolates. PLoS One 9:e110620.

68. Silveira GF, Wowk PF, Cataneo AHD, Dos Santos PF, Delgobo M, Stimamiglio

MA, Lo Sarzi M, Thomazelli APFS, Conchon-Costa I, Pavanelli WR, Antonelli

LRV, Báfica A, Mansur DS, Dos Santos CND, Bordignon J. 2018. Human T

Lymphocytes Are Permissive for Dengue Virus Replication. J Virol 92.

69. Mutso M, Saul S, Rausalu K, Susova O, Žusinaite E, Mahalingam S, Merits A.

2017. Reverse genetic system, genetically stable reporter viruses and packaged

subgenomic replicon based on a Brazilian Zika virus isolate. J Gen Virol

98:2712–2724.

70. Osei Kuffour E, Schott K, Jaguva Vasudevan AA, Holler J, Schulz WA, Lang PA,

Lang KS, Kim B, Häussinger D, König R, Münk C. 2018. USP18 (UBP43)

Abrogates p21-Mediated Inhibition of HIV-1. J Virol 92.

Figure 1-Espada *et al.*

Figure 3-Espada *et al.*

Figure 4-Espada *et al.*

Figure 5-Espada *et al.*

Figure 6-Espada *et al.*